Effects of prey turnover on poison frog toxins: using landscape ecology to assess how biotic

interactions affect species phenotypes

Ivan Prates[1*], Andrea Paz[2], Jason L. Brown[3], Ana C. Carnaval[2]

[1]Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian

Institution, Washington, DC, USA.

[2]Department of Biology, City College of New York, and Graduate Center, City University of

New York, New York, NY, USA.

[3]Zoology Department, Southern Illinois University, Carbondale, IL, USA.

[*]Correspondence author. E-mail: ivanprates@gmail.com

Running Title: Drivers of toxin turnover in poison frogs.

Keywords: Dendrobatidae, *Oophaga pumilio*, ant, alkaloid, chemical ecology, eco-evolutionary

dynamics, Generalized Dissimilarity Modeling, Multiple Matrix Regression.

Abstract

Ecological studies of species pairs demonstrated that biotic interactions promote

phenotypic change and eco-evolutionary feedbacks. However, we have a limited understanding

of how phenotypes respond to interactions with multiple taxa. We investigate how interactions

with a network of prey species contribute to spatially structured variation in the skin toxins of the

Neotropical poison frog *Oophaga pumilio*. Specifically, we assess how beta-diversity of

alkaloid-bearing arthropod prey assemblages (68 ant species) and evolutionary divergence

among populations (from a neutral genetic marker) contribute to frog poison dissimilarity (toxin

profiles composed of 230 different lipophilic alkaloids sampled from 934 frogs at 46 sites). We

show that ant assemblage turnover predicts alkaloid turnover and unique toxin combinations

across the range of *O. pumilio*. By contrast, evolutionary relatedness is barely correlated with

toxin variation. We discuss how the analytical framework proposed here can be extended to

other multi-trophic systems, coevolutionary mosaics, microbial assemblages, and ecosystem

services.

Introduction

Phenotypic variation within species, an essential component of evolutionary theory, has received increased attention by ecologists (Bolnick et al., 2011; Vildenes and Langangen, 2015). This interest has been chiefly motivated by evidence showing that phenotypic change, both adaptive and plastic, can happen within contemporary time scales and thus has consequences for ecological processes (Shoener, 2011; Hendry, 2015). Changes in trait frequencies can affect survival and reproduction and ultimately determine population density and persistence of a given species. In turn, these demographic changes may influence community-level and ecosystem functions such as nutrient cycling, decomposition, and primary productivity (Miner et al., 2005; Pelletier et al., 2009; Post and Palkovack, 2009). This interplay between evolutionary and ecological processes, or 'eco-evolutionary dynamics', has brought phenotypes to the center of ecological research (Hendry, 2015).

Several studies focusing on interacting species pairs have demonstrated that population-level phenotypic change can originate from biotic interactions, leading to geographic trait variation within species. For instance, different densities of Killifish predators in streams lead to distinct morphological and life history traits in their Trinidadian guppy prey (Endler, 1995). In western North America, levels of tetrodotoxin resistance in garter snake predators can match toxicity levels in local populations of tetrodotoxin-defended newt prey (Brodie et al., 2002). By focusing on the associations between two species, these studies have provided crucial insights into how interactions can lead to trait divergence and potentially shift the evolutionary trajectory of natural populations (Post and Palkovack, 2009; Hendry, 2015). The resulting trait diversity can have broad ecological consequences, altering the role of species in the ecosystem at a local scale (Palkovacs et al., 2009).

However, we still have a limited understanding of how interactions among multiple co-

distributed organisms contribute to complex phenotypes, particularly when the set of interacting

species and their phenotypes vary geographically. An example of a complex phenotype that is

shaped by a network of interactions with many species is the chemical defense system of poison

frogs (Dendrobatidae). In this clade of Neotropical amphibians, species can exhibit dozens to

hundreds of distinct lipophilic alkaloid toxins in their skin, and aposematic color patterns

advertise their distastefulness (Saporito et al., 2012; Santos et al., 2016). Poison composition

varies spatially within species such that populations closer in geography tend to have alkaloid

profiles more similar to one another than to populations farther away (Mebs et al., 2008; Saporito

et al., 2006, 2007a; Stuckert et al., 2014). This geographic variation in toxin profiles has been

attributed to local differences in prey availability, because poison frogs obtain their defensive

alkaloids from dietary arthropods (Daly et al., 2000; Saporito et al., 2004, 2007b, 2009; Jones et

al., 2012). For instance, specific toxins in the skin of a frog may match those of the arthropods

sampled from its gut (McGugan et al., 2016). However, individual alkaloids may be locally

present in frog skins but absent from the arthropods they consume, and vice-versa; thus, the

extent to which frog chemical traits reflect arthropods assemblages remains unclear (Daly et al.,

2000, 2002; Saporito et al., 2007b; Jones et al., 2012). Population differences may also stem

from an effect of shared evolutionary history, because alkaloid sequestration may be partially

under genetic control in poison frogs (Daly et al., 2003, 2005, 2009). These drivers of toxin

turnover can have broad consequences for the ecology of poison frogs, because alkaloids protect

these amphibians from numerous predators (Darst and Cummings, 2006; Gray et al., 2010;

Weldon et al., 2013; Murray et al., 2016), ectoparasites (Weldon et al., 2006), and pathogenic

microorganisms (Macfoy et al., 2005).

To address the question of how interactions among multiple organisms contribute to

spatially structured phenotypes, we investigate how prey assemblage turnover and evolutionary

divergence among populations predict the rich spectrum of toxins secreted by poison frogs. We

focus on the well-studied toxin profiles of the strawberry poison frog, *Oophaga pumilio*, which

exhibits over 230 distinct alkaloids over its Central American range (Daly et al., 1987, 2002;

Saporito et al., 2006, 2007a). First, we develop correlative models that approximate the spatial

distribution of toxic ants, a crucial source of alkaloids for poison frogs (Saporito et al., 2012;

Santos et al., 2016). Then, we apply Generalized Dissimilarity Modeling (GDM) (Ferrier et al.,

2007) to assess the contribution of projected ant assemblage turnover to chemical trait

dissimilarity among sites that have been screened for amphibian alkaloids, thus treating these

toxins as a "community of traits". To examine the effects of evolutionary relatedness, we

implement a Multiple Matrix Regression approach (MMRR) (Wang, 2013) that incorporates not

only prey turnover but also genetic divergences between *O. pumilio* populations as inferred from

a neutral genetic marker.

Material and Methods

Estimating frog poison composition dissimilarity

As poison composition data of *Oophaga pumilio*, we used lipophilic alkaloid profiles

derived from gas chromatography coupled with mass spectrometry of 934 frog skins sampled in

53 Central American sites (Daly et al., 1987, 2002; Saporito et al., 2006, 2007a), as compiled by

Saporito et al. (2007a). We georeferenced sampled sites based on maps and localities presented

by Saporito et al. (2007a) and other studies of *O. pumilio* (Saporito et al., 2006; Wang and

Shaffer, 2008; Brown et al., 2010; Hauswaldt et al., 2011; Gehara et al., 2013). Because

haphazard sampling may exaggerate poison composition variation in this dataset, and to

maximize alkaloid sampling effort, we combined data from different expeditions to each site,

pooling data from individuals. As such, we do not focus on potential short-term individual

fluctuations in toxin composition (Saporito et al., 2006), but instead on variation tied to spatial

gradients. To match the finest resolution available for the data grid predictors (see below), we

combined alkaloid data within a 1 km^2 grid cell. The final dataset included 230 alkaloids from 21

structural classes in 46 grid cell sites (Fig. 1) (alkaloid and locality data to be presented in the

Supplementary Information 1; presentation of raw data pending manuscript acceptance. See

Supplementary Information 2 for decisions on alkaloid identity). To estimate matrices of alkaloid

composition dissimilarity (pairwise Sorensen's distances) across frog populations, as well as

geographic distances between sites, we used the *fossil* package (Vavrek, 2011) in R v. 3.3.3 (R

Development Core Team, 2018).

Estimating arthropod prey assemblage dissimilarity

To approximate the spatial turnover of prey available to *Oophaga pumilio*, we focused on

alkaloid-bearing ants that occur in (but are not necessarily restricted to) Costa Rica, Nicaragua

and Panama, where that frog occurs. Ants are *O. pumilio*'s primary prey type, corresponding to

more than half of the ingested volume (Donnelly, 1991; Caldwell, 1996; Darst et al., 2004). This

frog also eats a large proportion of mites; however, limited occurrence data and taxonomic

knowledge for mite species (e.g., McGugan et al., 2016) precluded their inclusion in our spatial

analyses. Following a comprehensive literature search of alkaloid occurrence in ant taxa (Ritter

et al., 1973; Wheeler, 1981; Jones et al., 1982a,b, 1988, 1996, 1999, 2007, 2012; Daly et al.,

1994, 2000, 2005; Schroder et al., 1996; Spande et al., 1999; Leclercq et al., 2000; Saporito et

al., 2004, 2009; Clark et al., 2005; Fox et al., 2012; Chen et al., 2013; Adams et al., 2015;

Touchard et al., 2016), we estimated ant composition dissimilarity based on species that belong

to genera known to harbor alkaloids, as follows: *Acromyrmex, Anochetus, Aphaenogaster*, *Atta,*

Brachymyrmex, *Megalomyrmex*, *Monomorium*, *Nylanderia*, *Solenopsis*, and *Tetramorium* (see

Supplementary Information 2 for decisions on alkaloid presence in ant taxa). Georeferenced

records were compiled from the Ant Web database as per June 2017 using the *antweb* R package

(AntWeb, 2017). The search was restricted to the continental Americas between latitudes 40ºN

and 40ºS. We retained only those 68 ant species that had a minimum of five unique occurrence

records after spatial rarefaction (see below); the final dataset included a total of 1,417 unique

records. Ant locality data to be presented in the Supplementary Information 3 (presentation of

raw data pending manuscript acceptance).

Because available ant records rarely matched the exact locations where *O. pumilio*

alkaloids were characterized, we modeled the distribution of ant species at the landscape level to

allow an estimation of prey composition at sites with empirical frog poison data. We created a

species distribution model (SDM) for each ant species using MaxEnt (Phillips et al., 2006) based

on 19 bioclimatic variables from the Worldclim v. 1.4 database (Hijmans et al., 2005; available

at www.worldclim.org). Before modeling, we used the *spThin* R package (Aiello-Lammens,

2015) to rarefy ant records and ensure a minimum distance of 5 km between points, thus

reducing environmental bias from spatial auto-correlation (Veloz, 2009; Boria et al., 2014). To

reduce model over-fitting, we created a minimum-convex polygon defined by a 100 km radius

around each species occurrence points, restricting background point selection by the modeling

algorithm (Phillips et al., 2009; Anderson and Raza, 2010).

To properly parameterize the individual ant SDMs (Shcheglovitova and Anderson, 2013;

Boria et al., 2014), we chose the best combination of feature class (shape of the function

describing species occurrence vs. environmental predictors) and regularization multiplier (how

closely a model fits known occurrence records) using the *ENMeval* R package (Muscarella et al.,

2014). For species with 20 or more records, we evaluated models using k-fold cross-validation,

which segregates training and testing points in different random bins (in this case, k = 5). For

species with less than 20 records, we evaluated model fit using jackknife, a particular case of k-

fold cross validation where the number of bins (k) is equal to the total number of points. We

evaluated model fit under five combinations of feature classes, as follows: (1) Linear, (2) Linear

and Quadratic, (3) Hinge, (4) Linear, Quadratic and Hinge, and (5) Linear, Quadratic, Hinge,

Product, and Threshold. As regularization multipliers, we tested values ranging from 0.5 to 5 in

0.5 increments (Shcheglovitova and Anderson, 2013; Brown, 2014). For each model, the best

parameter combination was selected using AICc. Based on this best combination, we generated a

final SDM for each ant species (parameters used in all SDMs are presented in the Supplementary

Information 4).

To estimate a matrix of prey assemblage turnover based on ant distributions, we

converted SDMs to binary maps using the 10^{th} percentile presence threshold. We then extracted

presences for each ant species at sites sampled for frog alkaloids using the *raster* R package

(Hijmans et al., 2015). Matrices of estimated ant composition dissimilarity (pairwise Sorensen's

distances) were calculated with *fossil* in R. To visualize ant species turnover across the range of

O. pumilio, we reduced the final dissimilarity matrices to three ordination axes by applying

multidimensional scaling using the *cmdscale* function in R. Each axis was then assigned a

separate RGB color (red, green, or blue) as per Brown et al. (2014). A distribution layer for *O.*

pumilio was obtained from the IUCN database (available at www.iucn.org).

Modeling alkaloid composition turnover as a function of ant species turnover

To test the hypothesis that toxin composition in poison frogs vary geographically as a

function of the composition of prey species, we modeled the turnover of *O. pumilio* alkaloids

using a Generalized Dissimilarity Modeling (GDM) approach (Ferrier et al., 2007). GDM is an

extension of matrix regression developed to model species composition turnover across regions

as a function of environmental predictors. Once a GDM model is fitted to available biological

data (estimated from species presences at sampled sites), the compositional dissimilarity across

unsampled areas can be estimated based on environmental predictors (available for both sampled

and unsampled sites). We implemented GDM following the steps of Rosauer et al. (2014) using

the *GDM* R package (Manion et al., 2018).

We used the compiled database of alkaloids per site as the base of our GDMs. As

predictor variables, we initially included geographic distance and all 68 individual models of ant

species distributions at a 1 km^2 resolution. To select the combination of predictors that contribute

the most to GDM models while avoiding redundant variables, we implemented a stepwise

backward elimination process (Williams et al., 2012), as follows: first, we built a model with all

predictor variables; then, those variables that explained less than the arbitrary amount of 0.1% of

the data deviance were removed iteratively, until only those variables contributing more than

0.1% were left.

To visualize estimated alkaloid turnover on geographic space from GDM outputs, we

applied multidimensional scaling on the resulting dissimilarity matrices, following the procedure

outlined above for the ant SDMs.

Estimating frog population genetic divergence

To evaluate associations between alkaloid composition and shared evolutionary history

between frog populations, we used the *cytochrome B* gene dataset of Hauswaldt et al. (2011),

who sampled 197 *O. pumilio* individuals from 25 Central American localities. Because most

sites sampled for genetic data are geographically close to sites sampled for alkaloids (Saporito et

al., 2007a; Hauswaldt et al., 2011), we paired up alkaloid and genetic data based on Voronoi

diagrams. A Voronoi diagram is a polygon whose boundaries encompass the area that is closest

to a reference point relative to all other points of any other polygon (Aurenhammer, 1991).

Specifically, we estimated polygons using the sites sampled for genetic data as references points

and paired them with the sites for toxin data contained within each resulting Voronoi diagram.

To estimate Voronoi diagrams, we used ArcGIS 10.3 (ESRI, Redlands). To calculate a matrix of

average uncorrected pairwise genetic distances among localities, we used Mega 7 (Kumar et al.,

2016). Six genetically sampled sites that had no corresponding alkaloid data were excluded from

the analyses.

Testing associations between toxin composition, prey assemblage dissimilarity, and population

genetic divergence

To test the effect of population evolutionary divergence on poison composition of *O.*

pumilio, we implemented a Multiple Matrix Regression with Randomization (MMRR) approach

in R (Wang, 2013). To allow comparisons between genetic divergence and prey composition, we

also included a matrix of ant assemblage dissimilarity as a predictor variable in MMRR models.

As response variables, we focused on two distinct alkaloid datasets, namely individual alkaloids

(n = 230) and alkaloid structural classes (n = 21). Additionally, to account for variation in poison

composition resulting from ingestion of arthropod sources other than ants (i.e., mites, beetles,

millipedes; Saporito et al., 2009), we performed a second set of analyses restricted to alkaloids (n

= 125) belonging to alkaloid classes (n = 9) reported to occur in ant taxa. To assess the statistical

significance of MMRR models and predictor variables, 10,000 permutations were used.

Matrices of alkaloid composition dissimilarity, estimated ant assemblage dissimilarity,

genetic distances between *O. pumilio* populations, and geographic distances between sites, as

well as Supplementary Information and R scripts used in all analyses, are available online

through GitHub (https://github.com/ivanprates/2019_gh_pumilio).

Results

Toxin diversity

Compilation of toxin profiles revealed large geographic variation in the number of

alkaloids that compose the poison of *Oophaga pumilio*. The richness of individual alkaloids

across sites varied between 7 and 48, while the richness of alkaloid structural classes varied

between 3 and 17. Alkaloid composition turnover among sites was high; pairwise Sorensen

distances ranged from 0.33 to 1 for individual alkaloids, and from 0.08 to 0.86 for alkaloid

structural classes (Sorensen distances vary from 0 to 1, with 0 representing identical

compositions and 1 representing no composition overlap across sites). A matrix regression

framework using MMRR indicated that geographic distances affected the turnover of both

individual alkaloids ($p < 0.0001$; $R^2 = 0.16$; $R = 0.4$) and alkaloid structural classes ($p = 0.003$;

$R^2 = 0.09$; $R = 0.29$).

Ant composition turnover

Distribution modeling of ant species suggested variation of ant richness over the

landscape, with a concentration of species in the central portion of the distribution of *O. pumilio*

in Costa Rica and southern Nicaragua (Fig. 2). The estimated number of alkaloid-bearing ant

species at sites sampled for toxin profiles ranged between 24 and 52. However, ant richness did

not have a significant effect on the number of individual alkaloids (linear regression; $p > 0.22$)

(Fig. 3a) or alkaloid structural classes ($p > 0.07$).

Estimated ant assemblage turnover was low within the distribution of *O. pumilio* (Fig. 2).

Pairwise ant Sorensen distances ranged from 0 to 0.47 at sites sampled for frog alkaloids,

reflecting the broad ranges inferred for several alkaloid-bearing ant species. MMRR analyses

suggested a significant effect of geographic distances between sites on species turnover of

alkaloid-bearing ants ($p < 0.0001$; $R^2 = 0.62$; $R = 0.78$).

Projection of ant beta-diversity on geographic space suggested that prey assemblages are

similar throughout the southern range of *O. pumilio* in Panama and Costa Rica, with a transition

in the northern part of the range in inland Nicaragua (Fig. 2). Inner mid-elevations are expected

to harbor ant assemblages that are distinct from those in the coastal lowlands.

Generalized Dissimilarity Modeling

Dissimilarity modeling supports the idea that ant assemblage turnover affects the spatial

variation of poison composition in *O. pumilio*. A GDM model explained 22.9% of the alkaloid

composition turnover when the model included geographic distances among sampled sites. A

model that did not incorporate geographic distances explained 20% of alkaloid profile

dissimilarity. After eliminating those ant distribution models that contributed very little to the

GDM ($< 0.1\%$) using a stepwise backward elimination procedure, only nine out of 68 species

were retained in the final model, as follows: *Acromyrmex coronatus*, *Anochetus orchidicola*,

Brachymyrmex coactus, *Brachymyrmex pictus*, *Monomorium ebeninum*, *Solenopsis azteca*,

Solenopsis bicolor, *Solenopsis pollux*, and an undescribed *Solenopsis* species (sp. "jtl001"). A

GDM model not including geographic distances explained 20% of the observed alkaloid beta-

diversity also for this reduced ant dataset.

Projection of GDM outputs onto geographic space indicates areas expected to have more

similar amphibian alkaloid profiles based on ant turnover (Fig. 4). The results suggest latitudinal

variation in poison composition, with a gradual transition from the southern part of the

distribution of *O. pumilio* (in Panama) through the central and northern parts of the range (in

Costa Rica and Nicaragua, respectively) (pink to purple to blue in Fig. 4). Another major

transition in poison composition was estimated across the eastern and western parts of the range

of *O. pumilio* in Nicaragua (blue to green in Fig. 4).

Multiple Matrix Regressions

In agreement with the GDM results, MMRR analyses supported the hypothesis that

spatial turnover of *O. pumilio* toxin profiles is affected by prey composition dissimilarity, in spite

of the limited variation of ant assemblages. The estimated turnover of alkaloid-bearing ant

assemblages had a significant effect on individual alkaloid composition dissimilarity ($p <$

0.0001; $R^2 = 0.13$) (Fig. 3b), a result that changed little when considering only those alkaloids

from classes known to occur in ants ($p < 0.0001$; $R^2 = 0.11$). Similarly, estimated alkaloid-

bearing ant assemblages had a significant yet weak effect on the dissimilarity of alkaloid

structural classes among sites ($p = 0.002$; $R^2 = 0.05$), a result that held when considering only

alkaloids classes known to occur in ants ($p = 0.007$; $R^2 = 0.03$).

Similar to the effect of ant composition, genetic distances between populations of *O.*

pumilio had a weak yet significant effect on frog alkaloid composition dissimilarity ($p < 0.0001$;

$R^2 = 0.09$) (Fig. 3c). A MMRR model incorporating both frog population genetic distances and

ant composition dissimilarity explained around 15% of alkaloid composition variation among

populations of *O. pumilio* ($p_{ants} < 0.001$; $p_{gen} = 0.05$).

Discussion

Effects of prey turnover on trait variation, and its ecological consequences

Based on estimates of the distribution of alkaloid-bearing ants, we found associations

between prey composition gradients and spatial turnover in the defensive chemical traits of the

strawberry poison frog, *Oophaga pumilio*. Species distribution modeling supported that the pool

of alkaloid sources varies in space. Moreover, GDM and MMRR analyses inferred that alkaloid

beta diversity in *O. pumilio* covaries with the composition of ant assemblages. These results are

consistent with observations that distinct toxins are restricted to specific arthropod taxa (Saporito

et al., 2007a, 2012); prey species with limited distributions may contribute to unique defensive

phenotypes in different parts of the range of poison frog species. Accordingly, GDM predicted

unique alkaloid combinations in the northern range of *O. pumilio* (Nicaragua) relative to central

and southern areas (Costa Rica and Panama) (Fig. 4); that northern region remains poorly known

in terms of chemical diversity (Mebs et al., 2008), and future surveys may reveal novel toxin

combinations in the poison frogs that occur therein. These results are consistent with the view

that biotic interactions play a role in phenotypic variation within species and across space and

influence functional diversity in ecological communities (Miner et al., 2005; Pelletier et al.,

2009; Post and Palkovack, 2009).

Our analyses also indicate that different prey species may have disparate contributions to

observed chemical trait variation in poison frogs. After eliminating those ants that contributed

very little to the GDM (< 0.1%), only nine out of 68 species were retained in the final model.

This result may imply that a few prey items contribute disproportionately to the uniqueness of

chemical defenses among populations of *O. pumilio*. Alternatively, it may point to redundancy in

the alkaloids provided by prey species to poison frogs. Specifically, it is possible that some of the

ant species contributed less to alkaloid beta-diversity when in the presence of a co-distributed

and potentially functionally equivalent species, therefore being removed during the GDM's

stepwise backward elimination procedure.

These results have potential implications for the ecology and evolution of the interacting

species. For instance, frogs may favor and seek prey types that provide unique chemicals or

chemical combinations, increasing survival rates from encounters with their predators. This idea

is supported by evidence that alkaloid quantity, type, and richness result in differences in the

perceived palatability of poison frogs to predators (Murray et al., 2017; Bolton et al., 2017).

Moreover, because alkaloids vary from mildly unpalatable to lethally toxic (Daly et al., 2005;

Santos et al., 2016), the composition of amphibian poisons may affect the survival and behavior

of their predators following an attack (Darst and Cummings, 2006). From the perspective of

arthropod prey, predator feeding preferences and foraging behavior may affect population

densities and dynamics, potentially in a predator density-dependent fashion (Bolnick et al., 2011;

Pelletier et al., 2009). Finally, functional redundancy among prey types may confer resilience to

the defensive phenotypes of poison frogs, ensuring continued protection from predators despite

fluctuations in prey availability. Future investigations of these topics will advance our

understanding of how the phenotypic outcomes of species interactions affect ecological

processes.

Evolutionary relatedness and phenotypic similarity

In addition to the contribution of prey assemblages, the results support that phenotypic

similarity between populations also varies with evolutionary relatedness. MMRR analyses

indicated that population genetic divergence has a significant effect on alkaloid beta diversity in

O. pumilio, an effect that may reflect physiological or behavioral differences. For instance,

feeding experiments support that poison frog species differ in their capacity for lipophilic

alkaloid sequestration and that at least a few species can perform metabolic modification of

ingested alkaloids (Daly et al., 2003, 2005, 2009). Additionally, an effect of evolutionary

divergence on toxin profiles may stem from population differences in foraging strategies or

behavioral prey choice (Daly et al., 2000; Saporito et al., 2007a). Importantly, however, MMRR

suggested that genetic divergence accounts for less variation in the defensive phenotypes of *O.*

pumilio than alkaloid-bearing prey.

We employed genetic distances from a neutral marker as a proxy for shared evolutionary

history, and do not imply that this particular locus contributes to variation in frog physiology. An

assessment of how genetic variants influence chemical trait diversity in poison frogs will rely on

sampling of functional genomic variation, for which the development of genomic resources (e.g.,

Rogers et al., 2018) will be essential. Nevertheless, our approach can be extended to other

systems where information on both neutral and functional genetic variation is available. These

systems include, for instance, predators where nucleotide substitutions in ion channel genes

confer resistance to toxic prey at a local scale (Feldman et al., 2010), and species where variation

in major histocompatibility complex loci drives resistance to pathogens locally (Savage and

Zamudio, 2016).

Other sources of phenotypic variation

Although we show that prey community composition explains part of the spatial variation

in frog poisons, this proportion is relatively small. This result may stem from challenges in

quantifying turnover of prey species that act as alkaloid sources. Due to restricted taxonomic and

distribution information, we estimated species distribution models only for a set of well-sampled

ant species and were unable to include other critical sources of dietary alkaloids, particularly

mites (Saporito et al., 2012; McGugan et al., 2016). Although not being able to incorporate a

substantial fraction of the prey assemblages expected to influence poison composition in *O.*

pumilio, the GDM was able to explain about 23% of poison variation in this species. This

proportion is likely to increase with the inclusion of additional species of ants and other alkaloid-

bearing prey taxa, such as mites, beetles, and millipedes (Daly et al., 2000; Dumbacher et al.,

2004; Saporito et al., 2003).

Limitations in our knowledge of arthropod chemical diversity may also have affected the

analyses. Contrary to our expectations, we found a slight decrease in the explanatory value of

models that incorporated only alkaloids from structural classes currently known to occur in ants,

a result that may reflect an underestimation of ant chemical diversity. For instance, mites and

beetles are thought to be the source of tricyclics to Neotropical poison frogs, but the discovery of

these alkaloids in African Myrmicinae ants suggests that they may also occur in species from

other regions (Schroder et al., 1996). As studies keep describing naturally occurring alkaloids,

we are far from a complete picture of chemical trait diversity in both arthropods and amphibians

(Saporito et al., 2012).

This study demonstrates that incorporating species interactions can provide new insights

into the drivers of phenotypic diversity even when other potential sources of variation are not

fully understood. It is possible that frog poison composition responds not only to prey

availability but also to alkaloid profiles in prey, since toxins may vary within arthropod species

(Daly et al., 2002; Saporito et al., 2004, 2007b; Dall'Aglio-Holvorcem et al., 2009; Fox et al.,

2012). There is evidence that arthropods synthesize their alkaloids endogenously (Leclercq et al.,

1996; Camarano et al., 2012; Haulotte et al., 2012), but they might also sequester toxins from

plants, fungi, and symbiotic microorganisms (Saporito et al., 2012; Santos et al., 2016).

Accounting for these other causes of variation will be a challenging task to chemical ecologists.

An alternative to bypass knowledge gaps on the taxonomy, distribution, and physiology of these

potential toxin sources may be to focus on their environmental correlates. A possible extension

of our approach is to develop models that incorporate abiotic predictors such as climate and soil

variation, similarly to investigations of community composition turnover at the level of species

(Ferrier et al., 2007; Zamborlini-Saiter et al., 2016).

Eco-evolutionary feedbacks

Associations between prey assemblage composition and predator phenotype provide

opportunities for eco-evolutionary feedbacks, the bidirectional effects between ecological and

evolutionary processes (Post and Palkovacs, 2009). In the case of poison frogs, there is evidence

that toxicity correlates to the conspicuousness of skin color patterns (Maan and Cummings, 2011),

which therefore act as aposematic signals for predators (Hegna et al., 2013). However, coloration

patterns also mediate assortative mating in poison frogs (Maan and Cummings, 2009; Crothers and

Cummings, 2013). When toxicity decreases, the intensity of selection for aposematism also

decreases, and mate choice may become a more important driver of color pattern evolution than

predation (Summers et al., 1999). Divergence due to sexual selection may happen quickly when

effective population sizes are small and population gene flow is limited, which is the case in *O.*

pumilio (Gehara et al., 2013). By affecting chemical defenses, it may be that spatially structured prey

assemblages have contributed to the vast diversity of color patterns seen in poison frogs. On the other

hand, our GDM approach inferred similar alkaloid profiles between populations of *O. pumilio*

that have distinct coloration patterns (Fig. 4). However, it may be challenging to predict frog

toxicity from chemical profiles (Daly et al., 2005), and future studies of this topic will advance

our understanding of the evolutionary consequences of poison composition variation.

Concluding remarks

Integrative approaches have demonstrated how strong associations between species can

lead to tight covariation among species traits, with an iconic example being that of predator and

prey species that engage in "coevolutionary arms races" (Thompson, 2005). However, it may be

harder to assess the outcomes of weaker interactions among multiple co-distributed organisms

(Anderson, 2017). The chemical defense system of poison frogs varies as a function of prey

assemblages composed of tens of arthropod species, each of them providing single or a few toxin

molecules (Daly et al., 2005). Not surprisingly, it has been hard to predict the combinations of

traits emerging from these interactions (Santos et al., 2016). The landscape ecology framework

presented here approaches this problem by incorporating data on biotic interactions throughout

the range of a focal species, a strategy that has also improved correlative models of species

distributions (Lewis et al., 2017; Sanín and Anderson, 2018). Our framework can be extended to

a range of systems that have similar structures, including other multi-trophic interactions (Van

der Putten et al., 2001; Del-Claro, 2004; Scherber et al., 2010), geographic coevolutionary

mosaics (Greene and McDiarmid, 1981; Mallet et al., 1995; Symula et al., 2001), microbial

assemblages (Zomorrodi and Maranas, 2012; Landesman et al., 2014), and ecosystem services

(Moorhead and Sinsabaugh, 2006; Allison, 2012; Gossner et al., 2016). Integrative approaches to

the problem of how spatially structured biotic interactions contribute to phenotypic diversity will

continue to advance our understanding of the interplay between ecological and evolutionary

processes.

Acknowledgments

We are grateful to entomologists David Lohman, Corrie Moreau, and Israel del Toro for

suggestions and advice about ant diversity, as well as its ecological drivers, in the initial stages of

this project. Suggestions by Rayna C. Bell, Kevin P. Mulder, Michael L. Yuan, Edward A.

Myers, Ryan K. Schott, Maria Strangas, Danielle Rivera, Catherine Graham, Robert Anderson,

and the New York Species Distribution Modeling Group greatly improved this manuscript. This

work was co-funded by FAPESP (BIOTA 2011/50146-6 and 2013/50297–0), NSF (DEB

1343578), and NASA through the Dimensions of Biodiversity Program. IP acknowledges

additional funding from a Smithsonian Peter Buck Postdoctoral Fellowship.

References

Adams, R.M., Jones, T.H., Jeter, A.W., Henrik, H., Schultz, T.R. and Nash, D.R., 2012. A

comparative study of exocrine gland chemistry in *Trachymyrmex* and *Sericomyrmex*

fungus-growing ants. Biochemical Systematics and Ecology, 40, pp. 91-97.

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B. and Anderson, R.P., 2015.

spThin: an R package for spatial thinning of species occurrence records for use in

ecological niche models. Ecography, 38(5), pp. 541-545.

Allison, S.D., 2012. A trait-based approach for modelling microbial litter decomposition.

Ecology Letters, 15(9), pp. 1058-1070.

Anderson, R.P., 2017. When and how should biotic interactions be considered in models of

species niches and distributions?. Journal of Biogeography, 44(1), pp. 8-17.

Anderson, R.P. and Raza, A., 2010. The effect of the extent of the study region on GIS models of

species geographic distributions and estimates of niche evolution: preliminary tests with

montane rodents (genus *Nephelomys*) in Venezuela. Journal of Biogeography, 37(7), pp.

1378-1393.

AntWeb. Available from http://www.antweb.org. Accessed June 2017.

Aurenhammer, F., 1991. Voronoi diagrams—a survey of a fundamental geometric data structure.

ACM Computing Surveys (CSUR), 23(3), pp. 345-405.

Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., Rudolf,

V.H., Schreiber, S.J., Urban, M.C. and Vasseur, D.A., 2011. Why intraspecific trait

variation matters in community ecology. Trends in Ecology & Evolution, 26(4), pp. 183-

192.

Bolton, S.K., Dickerson, K. and Saporito, R.A., 2017. Variable alkaloid defenses in the

Dendrobatid poison frog *Oophaga pumilio* are perceived as differences in palatability to

arthropods. Journal of Chemical Ecology, 43(3), pp. 273-289.

Boria, R.A., Olson, L.E., Goodman, S.M. and Anderson, R.P., 2014. Spatial filtering to reduce

sampling bias can improve the performance of ecological niche models. Ecological

Modelling, 275, pp. 73-77.

Brodie Jr, E.D., Ridenhour, B.J. and Brodie III, E.D., 2002. The evolutionary response of

predators to dangerous prey: hotspots and coldspots in the geographic mosaic of

coevolution between garter snakes and newts. Evolution, 56(10), pp. 2067-2082.

Brown, J.L., Maan, M.E., Cummings, M.E. and Summers, K., 2010. Evidence for selection on

coloration in a Panamanian poison frog: a coalescent-based approach. Journal of

Biogeography, 37(5), pp. 891-901.

Brown, J.L., 2014. SDM toolbox: a python-based GIS toolkit for landscape genetic,

biogeographic and species distribution model analyses. Methods in Ecology and

Evolution, 5(7), pp. 694-700.

Caldwell, J.P., 1996. The evolution of myrmecophagy and its correlates in poison frogs (Family

Dendrobatidae). Journal of Zoology, 240(1), pp. 75-101.

Camarano, S., González, A. and Rossini, C., 2012. Origin of *Epilachnapaenulata* defensive

alkaloids: incorporation of [1-13C]-sodium acetate and [methyl-2H3]-stearic acid.

Journal of Insect Physiology, 58(1), pp. 110-115.

Chen, J., Rashid, T., Feng, G., Zhao, L. and Oi, D., 2013. Defensive chemicals of tawny crazy

487   ants, *Nylanderia fulva* (Hymenoptera: Formicidae) and their toxicity to red imported fire

488   ants, *Solenopsis invicta* (Hymenoptera: Formicidae). Toxicon, 76, pp. 160-166.

489   Clark, V.C., Raxworthy, C.J., Rakotomalala, V., Sierwald, P. and Fisher, B.L., 2005. Convergent

490   evolution of chemical defense in poison frogs and arthropod prey between Madagascar

491   and the Neotropics. Proceedings of the National Academy of Sciences, 102(33), pp.

492   11617-11622.

493   Crothers, L.R. and Cummings, M.E., 2013. Warning signal brightness variation: sexual selection

494   may work under the radar of natural selection in populations of a polytypic poison frog.

495   The American Naturalist, 181(5), pp. E116-E124.

496   Dall'Aglio-Holvorcem, C.G., Benson, W.W., Gilbert, L.E., Trager, J.C. and Trigo, J.R., 2009.

497   Chemical tools to distinguish the fire ant species *Solenopsis invicta* and *S. saevissima*

498   (Formicidae: Myrmicinae) in Southeast Brazil. Biochemical Systematics and Ecology,

499   37(4), pp. 442-451.

500   Daly, J.W., Myers, C.W. and Whittaker, N., 1987. Further classification of skin alkaloids from

501   neotropical poison frogs (Dendrobatidae), with a general survey of toxic/noxious

502   substances in the amphibia. Toxicon, 25(10), pp. 1023-1095.

503   Daly, J.W., Secunda, S.I., Garraffo, H.M., Spande, T.F., Wisnieski, A. and Cover Jr, J.F., 1994.

504   An uptake system for dietary alkaloids in poison frogs (Dendrobatidae). Toxicon, 32(6),

505   pp. 657-663.

506   Daly, J.W., Garraffo, H.M., Jain, P., Spande, T.F., Snelling, R.R., Jaramillo, C. and Rand, A.S.,

507   2000. Arthropod–frog connection: decahydroquinoline and pyrrolizidine alkaloids

common to microsympatric myrmicine ants and dendrobatid frogs. Journal of Chemical

Ecology, 26(1), pp. 73-85.

Daly, J.W., Kaneko, T., Wilham, J., Garraffo, H.M., Spande, T.F., Espinosa, A. and Donnelly,

M.A., 2002. Bioactive alkaloids of frog skin: combinatorial bioprospecting reveals that

pumiliotoxins have an arthropod source. Proceedings of the National Academy of

Sciences, 99(22), pp. 13996-14001.

Daly, J.W., Garraffo, H.M., Spande, T.F., Clark, V.C., Ma, J., Ziffer, H. and Cover, J.F., 2003.

Evidence for an enantioselective pumiliotoxin 7-hydroxylase in dendrobatid poison frogs

of the genus *Dendrobates*. Proceedings of the National Academy of Sciences, 100(19),

pp. 11092-11097.

Daly, J.W., Spande, T.F. and Garraffo, H.M., 2005. Alkaloids from amphibian skin: a tabulation

of over eight-hundred compounds. Journal of Natural Products, 68(10), pp. 1556-1575.

Daly, J.W., Ware, N., Saporito, R.A., Spande, T.F. and Garraffo, H.M., 2009. N-

methyldecahydroquinolines: an unexpected class of alkaloids from Amazonian poison

frogs (Dendrobatidae). Journal of Natural Products, 72(6), pp. 1110-1114.

Darst, C.R., Menéndez-Guerrero, P.A., Coloma, L.A. and Cannatella, D.C., 2004. Evolution of

dietary specialization and chemical defense in poison frogs (Dendrobatidae): a

comparative analysis. The American Naturalist, 165(1), pp. 56-69.

Darst, C.R. and Cummings, M.E., 2006. Predator learning favours mimicry of a less-toxic model

in poison frogs. Nature, 440(7081), p. 208.

Del-Claro, K., 2004. Multitrophic relationships, conditional mutualisms, and the study of

interaction biodiversity in tropical savannas. Neotropical Entomology, 33(6), pp. 665-

672.

Donnelly, M.A., 1991. Feeding patterns of the strawberry poison frog, *Dendrobates pumilio*

(Anura: Dendrobatidae). Copeia, pp. 723-730.

Dumbacher, J.P., Wako, A., Derrickson, S.R., Samuelson, A., Spande, T.F. and Daly, J.W.,

2004. Melyrid beetles (Choresine): a putative source for the batrachotoxin alkaloids

found in poison-dart frogs and toxic passerine birds. Proceedings of the National

Academy of Sciences, 101(45), pp. 15857-15860.

Endler, J.A., 1995. Multiple-trait coevolution and environmental gradients in guppies. Trends in

Ecology & Evolution, 10(1), pp. 22-29.

Feldman, C.R., Brodie Jr, E.D., Brodie III, E.D. and Pfrender, M.E., 2010. Genetic architecture

of a feeding adaptation: garter snake (*Thamnophis*) resistance to tetrodotoxin bearing

prey. Proceedings of the Royal Society B: Biological Sciences, 277(1698), pp. 3317-

3325.

Ferrier, S., Manion, G., Elith, J. and Richardson, K., 2007. Using generalized dissimilarity

modelling to analyse and predict patterns of beta diversity in regional biodiversity

assessment. Diversity and Distributions, 13(3), pp. 252-264.

Fox, E.G.P., Pianaro, A., Solis, D.R., Delabie, J.H.C., Vairo, B.C., Machado, E.D.A. and Bueno,

O.C., 2012. Intraspecific and intracolonial variation in the profile of venom alkaloids and

cuticular hydrocarbons of the fire ant *Solenopsis saevissima* Smith (Hymenoptera:

Formicidae). Psyche: A Journal of Entomology, 2012.

Gehara, M., Summers, K. and Brown, J.L., 2013. Population expansion, isolation and selection:

novel insights on the evolution of color diversity in the strawberry poison frog.

Evolutionary Ecology, 27(4), pp. 797-824.

Gossner, M.M., Lewinsohn, T.M., Kahl, T., Grassein, F., Boch, S., Prati, D., Birkhofer, K.,

Renner, S.C., Sikorski, J., Wubet, T. and Arndt, H., 2016. Land-use intensification causes

multitrophic homogenization of grassland communities. Nature, 540(7632), p. 266.

Gray, H.M. and Kaiser, H., 2010. Does alkaloid sequestration protect the green poison frog,

Dendrobates auratus, from predator attacks. Salamandra, 46(4), pp. 235-238.

Greene, H.W. and McDiarmid, R.W., 1981. Coral snake mimicry: does it occur?. Science,

213(4513), pp. 1207-1212.

Haulotte, E., Laurent, P. and Braekman, J.C., 2012. Biosynthesis of defensive coccinellidae

alkaloids: incorporation of fatty acids in adaline, coccinelline, and harmonine. European

Journal of Organic Chemistry, 2012(10), pp. 1907-1912.

Hauswaldt, J.S., Ludewig, A.K., Vences, M. and Pröhl, H., 2011. Widespread co-occurrence of

divergent mitochondrial haplotype lineages in a Central American species of poison frog

(*Oophaga pumilio*). Journal of Biogeography, 38(4), pp. 711-726.

Hendry, A.P., 2015. Key questions on the role of phenotypic plasticity in eco-evolutionary

dynamics. Journal of Heredity, 107(1), pp. 25-41.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. and Jarvis, A., 2005. Very high resolution

interpolated climate surfaces for global land areas. International Journal of Climatology:

A Journal of the Royal Meteorological Society, 25(15), pp. 1965-1978.

Hijmans, R.J., van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J.A., Lamigueiro,

O.P., Bevan, A., Racine, E.B., Shortridge, A. and Hijmans, M.R.J., 2015. Package

'raster'. R package.

Jones, T.H., Blum, M.S. and Fales, H.M., 1982a. Ant venom alkaloids from *Solenopsis* and

Monorium species: recent developments. Tetrahedron, 38(13), pp. 1949-1958.

Jones, T.H., Blum, M.S., Howard, R.W., McDaniel, C.A., Fales, H.M., DuBois, M.B. and

577   Torres, J., 1982b. Venom chemistry of ants in the genus *Monomorium*. Journal of

578   Chemical Ecology, 8(1), pp. 285-300.

579   Jones, T.H., Blum, M.S., Andersen, A.N., Fales, H.M. and Escoubas, P., 1988. Novel 2-ethyl-5-

580   alkylpyrrolidines in the venom of an Australian ant of the genus *Monomorium*. Journal of

581   Chemical Ecology, 14(1), pp. 35-45.

582   Jones, T.H., Torres, J.A., Spande, T.F., Garraffo, H.M., Blum, M.S. and Snelling, R.R., 1996.

583   Chemistry of venom alkaloids in some *Solenopsis* (*Diplorhoptrum*) species from Puerto

584   Rico. Journal of Chemical Ecology, 22(7), pp. 1221-1236.

585   Jones, T.H., Gorman, J.S., Snelling, R.R., Delabie, J.H., Blum, M.S., Garraffo, H.M., Jain, P.,

586   Daly, J.W. and Spande, T.F., 1999. Further alkaloids common to ants and frogs:

587   decahydroquinolines and a quinolizidine. Journal of Chemical Ecology, 25(5), pp. 1179-

588   1193.

589   Jones, T.H., Voegtle, H.L., Miras, H.M., Weatherford, R.G., Spande, T.F., Garraffo, H.M., Daly,

590   J.W., Davidson, D.W. and Snelling, R.R., 2007. Venom chemistry of the ant *Myrmicaria*

591   *melanogaster* from Brunei. Journal of Natural Products, 70(2), pp. 160-168.

592   Jones, T.H., Adams, R.M., Spande, T.F., Garraffo, H.M., Kaneko, T. and Schultz, T.R., 2012.

593   Histrionicotoxin alkaloids finally detected in an ant. Journal of Natural Products, 75(11),

594   pp. 1930-1936.

595   Kumar, S., Stecher, G. and Tamura, K., 2016. MEGA7: molecular evolutionary genetics analysis

596   version 7.0 for bigger datasets. Molecular Biology and Evolution, 33(7), pp. 1870-1874.

Landesman, W.J., Nelson, D.M. and Fitzpatrick, M.C., 2014. Soil properties and tree species

drive ß-diversity of soil bacterial communities. Soil Biology and Biochemistry, 76, pp.

201-209.

Leclercq, S., Braekman, J.C., Daloze, D., Pasteels, J.M. and Van der Meer, R.K., 1996.

Biosynthesis of the solenopsins, venom alkaloids of the fire ants. Naturwissenschaften,

83(5), pp. 222-225.

Leclercq, S., Braekman, J.C., Daloze, D. and Pasteels, J.M., 2000. The defensive chemistry of

ants. In Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of

Organic Natural Products (pp. 115-229). Springer, Vienna.

Lewis, J.S., Farnsworth, M.L., Burdett, C.L., Theobald, D.M., Gray, M. and Miller, R.S., 2017.

Biotic and abiotic factors predicting the global distribution and population density of an

invasive large mammal. Scientific Reports, 7, p. 44152.

Maan, M.E. and Cummings, M.E., 2011. Poison frog colors are honest signals of toxicity,

particularly for bird predators. The American Naturalist, 179(1), pp. E1-E14.

Macfoy, C., Danosus, D., Sandit, R., Jones, T.H., Garraffo, H.M., Spande, T.F. and Daly, J.W.,

2005. Alkaloids of anuran skin: antimicrobial function?. Zeitschrift für Naturforschung

C, 60(11-12), pp. 932-937.

Mallet, J. and Gilbert Jr, L.E., 1995. Why are there so many mimicry rings? Correlations

between habitat, behaviour and mimicry in *Heliconius* butterflies. Biological Journal of

the Linnean Society, 55(2), pp. 159-180.

Manion, G., Lisk, M., Ferrier, S., Nieto-Lugilde, D., Mokany, K. and Fitzpatrick, M.C., 2018.

gdm: generalized dissimilarity modeling. R package ver. 1.3. 11.

McGugan, J.R., Byrd, G.D., Roland, A.B., Caty, S.N., Kabir, N., Tapia, E.E., Trauger, S.A.,

Coloma, L.A. and O'Connell, L.A., 2016. Ant and mite diversity drives toxin variation in

the Little Devil Poison frog. Journal of Chemical Ecology, 42(6), pp. 537-551.

Mebs, D., Pogoda, W., Batista, A., Ponce, M., Köhler, G. and Kauert, G., 2008. Variability of

alkaloid profiles in *Oophaga pumilio* (Amphibia: Anura: Dendrobatidae) from western

Panama and southern Nicaragua. Salamandra, 44(24), p. 247.

Miner, B.G., Sultan, S.E., Morgan, S.G., Padilla, D.K. and Relyea, R.A., 2005. Ecological

consequences of phenotypic plasticity. Trends in Ecology & Evolution, 20(12), pp. 685-

692.

Moorhead, D.L. and Sinsabaugh, R.L., 2006. A theoretical model of litter decay and microbial

interaction. Ecological Monographs, 76(2), pp. 151-174.

Murray, E.M., Bolton, S.K., Berg, T. and Saporito, R.A., 2016. Arthropod predation in a

dendrobatid poison frog: does frog life stage matter?. Zoology, 119(3), pp. 169-174.

Muscarella, R., Galante, P.J., Soley-Guardia, M., Boria, R.A., Kass, J.M., Uriarte, M. and

Anderson, R.P., 2014. ENM eval: an R package for conducting spatially independent

evaluations and estimating optimal model complexity for Maxent ecological niche

models. Methods in Ecology and Evolution, 5(11), pp. 1198-1205.

Palkovacs, E.P., Marshall, M.C., Lamphere, B.A., Lynch, B.R., Weese, D.J., Fraser, D.F.,

Reznick, D.N., Pringle, C.M. and Kinnison, M.T., 2009. Experimental evaluation of

evolution and coevolution as agents of ecosystem change in Trinidadian streams.

Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1523), pp.

1617-1628.

Pelletier, F., Garant, D. and Hendry, A.P., 2009. Eco-evolutionary dynamics. Philosophical

Transactions of the Royal Society B, pp. 1483-1489.

Phillips, S.J., Anderson, R.P. and Schapire, R.E., 2006. Maximum entropy modeling of species

geographic distributions. Ecological Modelling, 190(3-4), pp. 231-259.

Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J. and Ferrier, S.,

2009. Sample selection bias and presence-only distribution models: implications for

background and pseudo-absence data. Ecological Applications, 19(1), pp. 181-197.

Post, D.M. and Palkovacs, E.P., 2009. Eco-evolutionary feedbacks in community and ecosystem

ecology: interactions between the ecological theatre and the evolutionary play.

Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1523), pp.

1629-1640.

Ritter, F.J., Rotgans, I.E.M., Talman, E., Verwiel, P.E.J. and Stein, F., 1973. 5-methyl-3-butyl-

octahydroindolizine, a novel type of pheromone attractive to Pharaoh's ants

(*Monomorium pharaonis* (L.). Experientia, 29(5), pp. 530-531.

Rogers, R.L., Zhou, L., Chu, C., Márquez, R., Corl, A., Linderoth, T., Freeborn, L., MacManes,

M.D., Xiong, Z., Zheng, J. and Guo, C., 2018. Genomic takeover by transposable

elements in the strawberry poison frog. Molecular Biology and Evolution, 35(12), pp.

2913-2927.

Rosauer, D.F., Ferrier, S., Williams, K.J., Manion, G., Keogh, J.S. and Laffan, S.W., 2014.

Phylogenetic generalised dissimilarity modelling: a new approach to analysing and

predicting spatial turnover in the phylogenetic composition of communities. Ecography,

37(1), pp. 21-32.

Sanín, C. and Anderson, R.P., 2018. A framework for simultaneous tests of abiotic, biotic, and

historical drivers of species distributions: empirical tests for North American wood

warblers based on climate and pollen. The American Naturalist, 192(2), pp. E48-E61.

Santos, J.C., Tarvin, R.D. and O'Connell, L.A., 2016. A review of chemical defense in poison

frogs (Dendrobatidae): ecology, pharmacokinetics, and autoresistance. In Chemical

Signals in Vertebrates 13 (pp. 305-337). Springer, Cham.

Saporito, R.A., Donnelly, M.A., Hoffman, R.L., Garraffo, H.M. and Daly, J.W., 2003. A

siphonotid millipede (*Rhinotus*) as the source of spiropyrrolizidine oximes of dendrobatid

frogs. Journal of Chemical Ecology, 29(12), pp. 2781-2786.

Saporito, R.A., Garraffo, H.M., Donnelly, M.A., Edwards, A.L., Longino, J.T. and Daly, J.W.,

2004. Formicine ants: an arthropod source for the pumiliotoxin alkaloids of dendrobatid

poison frogs. Proceedings of the National Academy of Sciences, 101(21), pp. 8045-8050.

Saporito, R.A., Donnelly, M.A., Garraffo, H.M., Spande, T.F. and Daly, J.W., 2006. Geographic

and seasonal variation in alkaloid-based chemical defenses of *Dendrobates pumilio* from

Bocas del Toro, Panama. Journal of Chemical Ecology, 32(4), pp. 795-814.

Saporito, R.A., Donnelly, M.A., Jain, P., Garraffo, H.M., Spande, T.F. and Daly, J.W., 2007a.

Spatial and temporal patterns of alkaloid variation in the poison frog *Oophaga pumilio* in

Costa Rica and Panama over 30 years. Toxicon, 50(6), pp. 757-778.

Saporito, R.A., Donnelly, M.A., Norton, R.A., Garraffo, H.M., Spande, T.F. and Daly, J.W.,

2007b. Oribatid mites as a major dietary source for alkaloids in poison frogs. Proceedings

of the National Academy of Sciences, 104(21), pp. 8885-8890.

Saporito, R.A., Spande, T.F., Garraffo, H.M. and Donnelly, M.A., 2009. Arthropod alkaloids in

poison frogs: a review of the dietary hypothesis. Heterocycles, 79(1), pp. 277-97.

Saporito, R.A., Donnelly, M.A., Spande, T.F. and Garraffo, H.M., 2012. A review of chemical

ecology in poison frogs. Chemoecology, 22(3), pp. 159-168.

Savage, A.E. and Zamudio, K.R., 2016. Adaptive tolerance to a pathogenic fungus drives major

histocompatibility complex evolution in natural amphibian populations. Proceedings of

the Royal Society B: Biological Sciences, 283(1827), p. 20153115.

Scherber, C., Eisenhauer, N., Weisser, W.W., Schmid, B., Voigt, W., Fischer, M., Schulze, E.D.,

Roscher, C., Weigelt, A., Allan, E. and Beßler, H., 2010. Bottom-up effects of plant

diversity on multitrophic interactions in a biodiversity experiment. Nature, 468(7323), p.

553.

Schoener, T.W., 2011. The newest synthesis: understanding the interplay of evolutionary and

ecological dynamics. Science, 331(6016), pp. 426-429.

Schröder, F., Franke, S., Francke, W., Baumann, H., Kaib, M., Pasteels, J.M. and Daloze, D.,

1996. A new family of tricyclic alkaloids from *Myrmicaria* ants. Tetrahedron, 52(43), pp.

13539-13546.

Shcheglovitova, M. and Anderson, R.P., 2013. Estimating optimal complexity for ecological

niche models: a jackknife approach for species with small sample sizes. Ecological

Modelling, 269, pp. 9-17.

Spande, T.F., Jain, P., Garraffo, H.M., Pannell, L.K., Yeh, H.J., Daly, J.W., Fukumoto, S.,

Imamura, K., Tokuyama, T., Torres, J.A. and Snelling, R.R., 1999. Occurrence and

significance of decahydroquinolines from dendrobatid poison frogs and a myrmicine ant:

use of 1H and 13C NMR in their conformational analysis. Journal of Natural Products,

62(1), pp. 5-21.

Stuckert, A.M., Saporito, R.A., Venegas, P.J. and Summers, K., 2014. Alkaloid defenses of co-

mimics in a putative Müllerian mimetic radiation. BMC Evolutionary Biology, 14(1), p.

76.

Summers, K., Symula, R., Clough, M. and Cronin, T., 1999. Visual mate choice in poison frogs.

Proceedings of the Royal Society of London. Series B: Biological Sciences, 266(1434),

pp. 2141-2145.

Symula, R., Schulte, R. and Summers, K., 2001. Molecular phylogenetic evidence for a mimetic

radiation in Peruvian poison frogs supports a Müllerian mimicry hypothesis. Proceedings

of the Royal Society of London. Series B: Biological Sciences, 268(1484), pp. 2415-

2421.

Thompson, J.N., 2005. Coevolution: the geographic mosaic of coevolutionary arms races.

Current Biology, 15(24), pp. R992-R994.

Touchard, A., Aili, S., Fox, E., Escoubas, P., Orivel, J., Nicholson, G. and Dejean, A., 2016. The

biochemical toxin arsenal from ant venoms. Toxins, 8(1), p. 30.

Van der Putten, W.H., Vet, L.E., Harvey, J.A. and Wäckers, F.L., 2001. Linking above-and

belowground multitrophic interactions of plants, herbivores, pathogens, and their

antagonists. Trends in Ecology & Evolution, 16(10), pp. 547-554.

Vavrek, M.J., 2011. Fossil: palaeoecological and palaeogeographical analysis tools.

Palaeontologia Electronica, 14(1), p. 16.

Veloz, S.D., 2009. Spatially autocorrelated sampling falsely inflates measures of accuracy for

presence-only niche models. Journal of Biogeography, 36(12), pp. 2290-2299.

Vindenes, Y. and Langangen, Ø., 2015. Individual heterogeneity in life histories and eco-

evolutionary dynamics. Ecology Letters, 18(5), pp. 417-432.

Wang, I.J. and Shaffer, H.B., 2008. Rapid color evolution in an aposematic species: a

phylogenetic analysis of color variation in the strikingly polymorphic strawberry poison-

dart frog. Evolution: International Journal of Organic Evolution, 62(11), pp. 2742-2759.

Wang, I.J., 2013. Examining the full effects of landscape heterogeneity on spatial genetic

variation: a multiple matrix regression approach for quantifying geographic and

ecological isolation. Evolution, 67(12), pp. 3403-3411.

Weldon, P.J., Kramer, M., Gordon, S., Spande, T.F. and Daly, J.W., 2006. A common

pumiliotoxin from poison frogs exhibits enantioselective toxicity against mosquitoes.

Proceedings of the National Academy of Sciences, 103(47), pp. 17818-17821.

Weldon, P.J., Cardoza, Y.J., Vander Meer, R.K., Hoffmann, W.C., Daly, J.W. and Spande, T.F.,

2013. Contact toxicities of anuran skin alkaloids against the fire ant (*Solenopsis invicta*).

Naturwissenschaften, 100(2), pp. 185-192.

Wheeler, J.W., Olubajo, O., Storm, C.B. and Duffield, R.M., 1981. Anabaseine: venom alkaloid

of Aphaenogaster ants. Science, 211(4486), pp. 1051-1052.

Williams, K.J., Belbin, L., Austin, M.P., Stein, J.L. and Ferrier, S., 2012. Which environmental

variables should I use in my biodiversity model?. International Journal of Geographical

Information Science, 26(11), pp. 2009-2047.

Zamborlini Saiter, F., Brown, J.L., Thomas, W.W., de Oliveira-Filho, A.T. and Carnaval, A.C.,

2016. Environmental correlates of floristic regions and plant turnover in the Atlantic

Forest hotspot. Journal of Biogeography, 43(12), pp. 2322-2331.

Zomorrodi, A.R. and Maranas, C.D., 2012. OptCom: a multi-level optimization framework for

the metabolic modeling and analysis of microbial communities. PLoS Computational

Biology, 8(2), pp. e1002363.

Figure 1. Sites sampled for skin alkaloid profiles of the strawberry poison frog, *Oophaga*

pumilio. Each site corresponds to a 1 km^2 grid cell, matching the resolution of environmental

predictors. Original alkaloid data compiled by Saporito et al. (2017a). The distribution of *O.*

pumilio is indicated in dark grey.

 Figure 2. Estimated species turnover (left) and richness of prey assemblages across the range of

 the poison frog *Oophaga pumilio* based on projected distributions of 68 ant species from

 alkaloid-bearing genera. Inset indicates the natural range of *O. pumilio* (dark grey).

Figure 3. Alkaloid richness in *Oophaga pumilio* as a function of ant assemblage richness across

sites (A); alkaloid dissimilarity in *O. pumilio* as a function of ant assemblage dissimilarity across

sites (B); and alkaloid dissimilarity in *O. pumilio* as a function of population genetic divergence

based on a neutral marker (C). Relationships in B and C are statistically significant; significance

was estimated from a Multiple Matrix Regression with Randomization (MMRR) approach (see

text).

 Figure 4. Estimated composition dissimilarity of alkaloid profiles across the range of *Oophaga*

774 *pumilio* as a function of spatial turnover of alkaloid-bearing ant species, from a Generalized

775 Dissimilarity Modeling approach (GDM). Similar colors on the map indicate similar estimated

776 alkaloid profiles. Pictures indicate dorsal skin coloration patterns in *O. pumilio*. Inset indicates

777 the natural range of *O. pumilio* (dark grey).