EEG power spectra and subcortical pathology in chronic disorders of consciousness

Evan S. Lutkenhoff, Ph.D.[a,b], Anna Nigri, Ph.D [c], Davide Rossi Sebastiano, MD, Ph.D [d], Davide Sattin, PsyD [e], Elisa Visani, M.Sc. [d], Cristina Rosazza, Ph.D. [f], Ludovico D'Incerti, MD [c], Maria Grazia Bruzzone, MD [c], Silvana Franceschetti, MD [d], Matilde Leonardi, MD [e], Stefania Ferraro, Ph.D.[c,*], Martin M. Monti Ph.D.[a,b]

[a]*Department of Psychology, University of California Los Angeles, Los Angeles, CA, USA*
[b]*Brain Injury Research Center (BIRC), Department of Neurosurgery, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA*
[c]*Department of Neuroradiology, Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milan, Italy.*
[d]*Department of Neurophysiology, Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milan, Italy.*
[e]*Neurology, Public Health, Disability Unit and Coma Research Centre, Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milan, Italy.*
[f]*Scientific Direction, Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milan, Italy.*
On the behalf of the Coma Research Center, Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milan, Italy.

Author Manuscript Version

*To whom correspondence should be addressed: Stefania Ferraro, Ph.D., Fondazione IRCCS Neurologico 'Carlo Besta', Via Celoria 11, 20133 Milan, Italy; Ph +39 (02) 23941, Fax +39 (02) 70638217; e-mail: Stefania.Ferraro@istituto-besta.it

Search terms: disorders of consciousness, subcortical pathology, EEG, MRI.

Abstract word count: 249
Manuscript body count: 4,524
References: 57

Abstract

Objective: To determine (i) the association between long-term impairment of consciousness after severe brain injury, spontaneous brain oscillations, and underlying subcortical damage, and (ii) whether such data can be used to aid patient diagnosis, a process known to be susceptible to high error rates.

Methods: Cross-sectional observational sample of 116 patients with a disorder of consciousness secondary to brain injury, collected prospectively at a tertiary center between 2011 and 2013. Multimodal analyses relating clinical measures of impairment, electroencephalographic measures of spontaneous brain activity, and magnetic resonance imaging data of subcortical atrophy were conducted in 2018.

Results: In the final analyzed sample of 61 patients, systematic associations were found between electroencephalographic power spectra and subcortical damage. Specifically, the ratio of beta-to-delta relative power was negatively associated with greater atrophy in regions of the bilateral thalamus and globus pallidus (both left > right) previously shown to be preferentially atrophied in chronic disorders of consciousness. Power spectrum total density was also negatively associated with widespread atrophy in regions of the left globus pallidus, right caudate, and in brainstem. Furthermore, we showed that the combination of demographics, encephalographic, and imaging data in an analytic framework can be employed to aid behavioral diagnosis.

Conclusions: These results ground, for the first time, electroencephalographic presentation detected with routine clinical techniques in the underlying brain pathology of disorders of consciousness and demonstrate how multimodal combination

of clinical, electroencephalographic, and imaging data can be employed in potentially

mitigating the high rates of misdiagnosis typical of this patient cohort.

Keywords: Consciousness, Electroencephalography, Magnetic Resonance Imaging, Severe Brain Injury

Introduction

When patients survive a severe brain injury, but fail to fully recover the two cardinal

elements of consciousness (i.e., awareness and arousal; (Laureys, 2005; Monti, 2012)),

they are said to enter a disorder of consciousness (DOC; (Monti, Laureys, & Owen,

2010)). This label describes a set of conditions including coma, a state of unarousable

unresponsiveness, the vegetative state (VS), a state of arousal in the absence of

(self)awareness, and the minimally conscious state (MCS), a state in which patients

can demonstrate some level of (self)awareness, albeit inconsistently. MCS can be

further subdivided into MCS+, to indicate patients capable of high-level behavioral

responses such as command following and intelligible verbalizations, and MCS-, to

indicate patients only demonstrating low-level behaviors such as visual pursuit and

appropriate contingent behavior (e.g., smiling or crying in response to emotional

stimuli; (Bruno, Vanhaudenhuyse, Thibaut, Moonen, & Laureys, 2011)).

Over the past 20 years, electroencephalography (EEG) and magnetic resonance

imaging (MRI) have been increasingly employed to monitor neurological status

(Brenner, 2005), residual cognitive function (Chennu et al., 2013; Monti et al., 2015;

Schnakers et al., 2008), state of awareness (Crone et al., 2015; Demertzi et al., 2019;

Owen et al., 2006), and potential for recovery (Bagnato et al., 2010; Crone, Bio, Vespa,

Lutkenhoff, & Monti, 2018; Schnakers et al., 2019) in DOC patients. In the context of

bedside EEG, analysis of the magnitude of oscillations at different frequencies (i.e.,

power spectrum analysis) has been shown to be capable of differentiating DOC

patients from patients with severe neurocognitive disorder but no consciousness

impairment (Leon-Carrion, Martin-Rodriguez, Damas-Lopez, Barroso y Martin, &

Dominguez-Morales, 2008), as well as clinical categories of chronic DOC (i.e., VS,

MCS), with depth of impairment correlating with slower, larger amplitude oscillations

(Lechinger et al., 2013; Lehembre et al., 2012). While this technique is sensitive to

different injury etiologies (Fingelkurts, Fingelkurts, Bagnato, Boccagni, & Galardi,

2013), it is blind to the underlying anatomical damage. At a theoretical level, the

mesocircuit theory of recovery of consciousness after brain injury predicts a

relationship between the slowing and amplitude of electrocortical oscillations and the

degree of pathologic changes taking place after trauma, hypoxia, or multifocal

ischemia (Schiff, 2016). While indirect evidence exists in support of this model, with

in vivo and *post-mortem* work demonstrating a relationship between damage to

thalamus, loss of thalamo-cortical structural connectivity, and depth of impairment

(Adams, Graham, & Jennett, 2000; Lutkenhoff et al., 2015; Zheng, Reggente,

Lutkenhoff, Owen, & Monti, 2017), there are no data directly connecting the patterns

of EEG power spectra and subcortical damage in long-term DOC patients, a gap which

is not only problematic for the clinician's interpretation of the observed EEG data, but

also hampers our ability to monitor, through an inexpensive, bedside, repeatable

technique interventions and their effects. In what follows, we address, in a large cohort

of patients with chronic DOC, the heretofore untested relationship between observed

electrocortical rhythms, patterns of subcortical brain atrophy (including thalamus,

brainstem, and basal ganglia), and clinical measures of awareness and arousal.

Methods

Participants

A convenience sample of 116 patients was recruited from a larger database (n=153)

(Nigri et al., 2017; Rosazza et al., 2016; Rossi Sebastiano et al., 2015; Rossi

Sebastiano et al., 2018) of chronic DOC patients with severe brain injury, who

underwent a 1-week assessment at the Coma Research Centre (CRC) of the

Neurological Institute C. Besta in Milan, Italy, between 2011 and 2013. The assessment

included (i) clinical evaluation, including the Coma Recovery Scale Revised (CRS-R;

(Giacino, Kalmar, & Whyte, 2004)), (ii) neurophysiological evaluation, including

resting EEG, and (iii) neuroradiological assessment, including MRI (see Table 1).

Insert Table 1 about here

The acquisition of both resting EEG and structural MRI datasets constituted the

inclusion criteria for the present study. Experienced raters independently assessed each

patient 4 times with the Italian version of the CRS-R (Sacco et al., 2011). The best-

recorded performance was used to classify patients as VS or MCS. As described below,

55 patients were discarded due to the low quality of the MRI data, following previously

established procedures (Lutkenhoff et al., 2015). Specifically, 24 datasets were excluded

because of excessive movement artifacts, 16 datasets were excluded due to software

failure in segmenting subcortical structures, 13 datasets were excluded due to poor

quality in the estimation of normalized brain tissue volume, and 2 subjects were

excluded due to large regions of signal dropout artifacts (e.g., implants) preventing

tissue segmentation (see Figure S1). Importantly, as discussed below, exclusions do

not bias the analyzed sample as compared to the full cohort (see also Table S1). No

patients were excluded on the basis of the EEG data. The final sample included 61 patients

with DOC (median age=53 years, range age = 20-82 years, 36 males), 37 VS and 24 MCS.

Of these, 33% (n = 20) had a traumatic brain injury (TBI) and 67% (n = 41) had non-

traumatic etiology (non-TBI). In particular, among non-TBI patients, 30% (n = 18)

suffered from anoxic brain injury and 37% (n = 23) from haemorrhagic and/or ischemic

brain injury. The median disease duration at the time of the study was 24 months (range =

5-198 months). (See Tables 1 and S1). The local Ethics Committee approved all aspects of

this research and written informed consent was obtained from the legally authorized

representative of the patients prior to their inclusion in the study.

Data acquisition and analysis

EEG data acquisition and processing

Patients underwent polygraphic recordings between 2 pm and 9 am on the following

day (Rossi Sebastiano et al., 2018). Recordings included electrooculography (EOG),

electromyography (EMG) from the sub-mental muscle, a bipolar precordial

electrocardiogram (ECG) derivation and an impedance thoracic pneumogram. EEG

recordings were made with Ag/AgCl surface electrodes (impedances <5kΩ) and

acquired at a sampling rate of 256 Hz using a computerized system (Micromed SpA,

Mogliano Veneto, Treviso, Italy). Raw EEG signals were recorded, against a common

reference electrode to allow off-line data reformatting, using a 19 EEG electrode array

placed according to the 10-20 International System (including frontal (Fp1, Fp2, F3,

F4), central (C3, C4), parietal (P3, P4), and occipital (O1, O2) electrodes). We

preliminarily divided the awake from sleep based on the polygraphic signal. Spectral

EEG analysis was then performed on one epoch of 5-minute, consecutive, artifact-free,

awake EEG signal. To minimize issues deriving from different levels of arousal, we

selected the EEG epochs as close as possible to the beginning of the recording, in

order to prevent signal degradation due to the prolonged recording and to maintain, as

much as possible, the same reproducible characteristics between recordings in different

patients. The selected epochs were filtered (1-70 Hz, 12 db/octave), followed by a 50

Hz notch filter to suppress the noise of the electrical power line, reformatted against

the linked ear-lobe reference. In order to remove blink-artifacts, we applied an ICA-

artifact rejection algorithm. Then, the selected EEG activity was divided into 90 non-

overlapping 2 s segments and analyzed using the fast Fourier transform. Absolute total

power and relative power were evaluated in the delta (1-4 Hz), theta (4-8 Hz), alpha

(8-13 Hz), beta (13-30 Hz) and gamma (>30 Hz) bands, and averaged within each

EEG channel.

MRI data acquisition and processing

Neuroimaging data were obtained with a 3T MRI machine (Achieva, Philips Healthcare

BV, Best, NL; 32-channel coil). The MRI protocol included a high-resolution 3D-TFE

T1-weighted sequence (185 sagittal slices, TR=9.781 ms, TE=4.6 ms, voxel size = 1

mm^3, flip angle = 8°). Analysis of subcortical structures was conducted using a

Figure 1. *Sample structure extraction (left) and 3-dimensional triangle vertex mesh (right). (Abbreviations: A, anterior; BrStem, brain stem; Caud, caudate; GP, globus pallidus; Hipp, hippocampus; L, left; P, posterior; Putm, putamen; R, right; Thal, thalamus. Figure from (Lutkenhoff et al., 2015).)*

technique known as shape or vertex analysis, part of the FMRIB software library

(FSL; FMRIB, Oxford, UK), following a previously established pipeline (Lutkenhoff

et al., 2015; Schnakers et al., 2019). Briefly, MR images were brain-extracted using

optiBET (Lutkenhoff et al., 2014), then the thalamus, caudate, putamen, globus

pallidus, hippocampus, and brainstem were segmented using FSL FIRST (Patenaude,

Smith, Kennedy, & Jenkinson, 2011), for each patient and structure separately, and

reconstructed into 3-dimensional vertex meshes, as depicted in Figure 1. In addition,

the normalized brain volume (NBV), a measure of global brain atrophy including

white and gray matter volume, was calculated for each patient using FSL SIENA

(Smith et al., 2002).

Statistical Analyses

In what follows, we describe four analysis which were employed to assess the

relationship between clinical/behavioral measures of impairment, electrophysiological

recordings, and subcortical measures of atrophy.

EEG analysis

To assess the relationship between EEG spectral power and clinical grouping, we ran a

mixed-model linear analysis with EEG (relative) power as the dependent variable,

laterality (left, right, middle), channel position (Fp, F, C, P, O), and EEG features

(total power, delta, theta, alpha, beta, gamma bands) as the repeated measures,

diagnosis (VS, MCS-, MCS+), etiology (TBI, non-TBI), sex, EEG frequency, age, and

months post-injury as fixed variables and subjects as the random variable. As

described below, the significant interaction between EEG features and diagnosis was

followed up with one mixed-model analysis per each EEG feature (using the same

model, albeit without the EEG feature as repeated variable). Individual mixed-models

were followed up with pairwise post-hoc comparisons between diagnostic groups (i.e.,

VS, MCS-, MCS+), with Šidák correction for multiplicity.

EEG – MRI analysis

In the second analysis, we related EEG spectral features to subcortical shape measures.

However, because of significant correlations among spectral characteristics across

electrodes and frequency bands, spectral data first were entered into a principal

component analysis (PCA), with varimax rotation. The PCA returned 8 components

with eigenvalue greater than 1, cumulatively explaining 91.8% of the total variance. The

first three components presented a similar pattern, each loading negatively, in all

electrodes, on the delta band and positively on beta, alpha, and theta bands,

respectively (henceforth, β/δ ratio component, α/δ ratio component, and θ/δ ratio

component, respectively). The fourth component loaded positively on all gamma

frequency electrodes (henceforth, γ component), while the fifth component loaded

positively on the total power for each electrode (henceforth, total power component).

Finally, the last three components appeared to capture diffuse statistical covariance

between electrodes, although with a preference for loading, respectively, positively on

the delta band and negatively on the alpha band in right hemispheric electrodes

(henceforth, δ/α_{RH} ratio component), loading positively on delta band and negatively

on the alpha band in Fp electrodes (henceforth, δ/α_{Fp} ratio component), and loading

negatively on both the alpha and theta bands in the parietal and occipital electrodes

(henceforth, $\alpha\theta_{(P,O)}^{-}$). (See Figure S2 for a depiction of the strength of each component

for VS, MCS-, and MCS+.) EEG components were then entered, as independent

variables, in a general linear model predicting local shape patterns (e.g., atrophy). Sex,

age, time-post-injury, etiology (i.e., TBI vs non-TBI), were included as covariates,

along with NBV (to ensure that observed tissue displacement reflect local subcortical

shape changes independent of overall brain atrophy (Lutkenhoff et al., 2015;

Schnakers et al., 2019)). Group-level significance was assessed with a non-parametric

permutation test at a level of $p < 0.05$ corrected for multiplicity using family-wise

cluster correction and threshold free cluster enhancement (TFCE) as implemented in

FSL randomize (Smith & Nichols, 2009).

CRS-R – MRI analysis

In this analysis, we related the patients' behavioral presentation, as captured by the

CRS-R subscales, with subcortical atrophy. Because of significant correlations

between the subscales of the CRS-R (i.e., the desired independent variables),

behavioral data were entered into a PCA performed analogously to the one described

above. The analysis returned 3 components with an eigenvalue greater than 1,

cumulatively explaining 69.57% of the variance. The three components loaded on,

respectively, the auditory, visual, and arousal subscales (henceforth, audio-visual-

arousal component), the motor subscale (henceforth, motor component), and the

oromotor and communication subscales (henceforth, oromotor-communication

component). As in the previous analysis, the three components were entered as

independent variables in a general linear model predicting subcortical shape, along with

the same covariates described above. Group-level significance was assessed identically

to the previous analysis.

Predicting DOC level from EEG spectral features

Finally, we employed a binary logistic regression to evaluate the degree to which

global atrophy and EEG measures related to diagnosis (i.e., VS vs. MCS). With a 3-

block model, we attempt to predict diagnosis from a model including demographic

variables only (i.e., age, sex, months post injury, etiology (TBI vs. non-TBI)), a model

including demographics and a measure of global atrophy (including global white

matter and gray matter), and a model including demographic, global atrophy, and EEG

variables (i.e., the 8 EEG components).

Results

EEG results

The mixed-model analysis revealed a significant interaction ($F(10, 1309.055) = 16.599$,

$p < 0.001$) between diagnostic group (i.e., VS, MCS-, MCS+) and EEG features (i.e.,

total power, delta, theta, alpha, beta, gamma frequency bands; see Figure 2 and Table

1), along with a significant main effect of diagnosis ($F(2, 4389.124) = 5.158$, $p =$

0.006), EEG features ($F(5, 1309.055) = 6.368$, $p < 0.001$), and months post-injury (F

$(2, 4220.566) = 5.407$, $p = 0.020$). Follow-up mixed-model analyses (one per EEG

feature) revealed a significant effect of diagnosis on total power ($F(2, 594.374) =$

19.115, $p < 0.001$; with pairwise-comparisons revealing that VS patients show

significantly less total power than both MCS groups), delta ($F(2, 770.006) = 4.620$, p

$= 0.010$; with VS patients having the most relative delta power, significantly more than

MCS- patients), theta ($F(2, 779.789) = 12.268$, $p < 0.001$; with VS patients showing

significantly less relative theta power than both MCS groups), alpha ($F(2, 748.766) =$

13.231, $p < 0.001$; with MCS- patients showing significantly more relative alpha

power than VS and MCS+ patients, and VS patients showing greater relative alpha

power than MCS+ patients), and gamma ($F(2, 746.625) = 5.416, p = 0.005$; with MCS+

patients showing significantly less relative gamma power than MCS- patients). (For

Figure 2. *Summary of EEG data. Total and relative power at each frequency band (to allow displaying on the same axis, values are normalized within each variable). (Error bars represent standard errors.)*

further description of the EEG data in this cohort see (Rossi Sebastiano et al., 2018).)

EEG – MRI analysis

As shown in Figure 3, three of the EEG factors exhibited significant correlations with

local atrophy measurements. Specifically, the β/δ ratio component was negatively

associated with greater atrophy in bilateral thalamus (left: $t = 3.28$, $p = 0.025$, 1,007

significant vertices, (sig. vert.), right: $t = 2.85$, $p = 0.041$, 544 sig. vert.), bilateral

globus pallidus (left: $t = 4.26$, $p = 0.002$, 491 sig. vert.; right: $t = 3.42$, $p = 0.025$, 356

Figure 3. Results for the rest EEG – MRI analysis. (a) Results for the β/δ ratio component; (b) results for the θ/δ component; (c) results for the total power component. Warmer colors indicate regions with greater atrophy; gray indicates no significant effect.

sig. vert.), left caudate ($t = 3.37$, $p = 0.02$, 793 sig. vert.), and right hippocampus ($t =$

3.21, $p = 0.02$, 735 sig. vert.) (see Fig. 3a). The θ/δ component was negatively

associated with increased atrophy in right putamen ($t = 3.72$, $p = 0.037$, 140 sig. vert.)

and right globus pallidus ($t = 3.89$, $p = 0.037$, 30 sig. vert.) (see Fig. 3b). Finally, the

total power component was negatively associated with widespread atrophy in the

brainstem (including the 4th ventricle region; $t = 4.41$, $p = 0.004$, 3704 sig. vert.), as

well as the left globus pallidus ($t = 4.00$, $p = 0.008$, 365 sig. vert.) and right caudate (t

$= 5.48$, $p = 0.018$, 193 sig. vert.) (see Fig. 3c). No significant associations were

detected for the remaining components.

CRS-R – MRI analysis

As shown in Figure 4a,b, two of the CRS-R components exhibited significant

correlations with local atrophy measurements. Specifically, the motor component was

negatively associated with greater atrophy in broad regions of the brainstem ($t = 3.69$,

$p = 0.074$, 4,301 sig. vert.; see Fig. 4a) while the oromotor-communication component

was negatively associated with greater atrophy in regions of the brainstem ($t = 2.74$, p

$= 0.05$, 943 sig. vert.), left putamen ($t = 3.23$, $p = 0.023$, 712 sig. vert.), and right

globus pallidus ($t = 2.38$, $p = 0.049$, 108 sig. vert.).

Figure 4. *Results for the CRS-R – MRI analyses. (a) Results for the CRS-R motor component; (b) results for the CRS-R oromotor-communication component. (See Fig. 3 for color interpretation.)*

Predicting DOC level from EEG spectral features

As shown in Figure 5a and Table 2, diagnosis (i.e., VS vs. MCS) was predicted

significantly better by the model including EEG components, overall brain atrophy

(including both white matter and gray matter), and demographic components (i.e., age,

sex, etiology (TBI vs. non-TBI), and months post injury; $\chi^2(54) = 29.09$, $p < 0.001$,

Nagelkerke pseudo-$R^2_{Nag} = 0.51$), compared to the model with demographics and

brain atrophy ($\chi^2(58) = 16.66$, $p < 0.001$, $R^2_{Nag} = 0.32$), as well as the model with

demographics only ($\chi^2(59) = 8.93$, $p = 0.003$, $R^2_{Nag} = 0.18$). Indeed, the full model

achieved better performance (i.e., area under the curve, $AUC = 0.87$), sensitivity

1   and specificity (sens/spec; 0.79, 0.81, respectively) than both other models ($AUC =$

2   0.78, sens/spec 0.58/0.78 and $AUC = 0.67$, sens/spec 0.29/0.89 for the

3   demographics and atrophy and the demographic only models, respectively).

4   Notably, the contribution of increasingly complex models (i.e., adding brain

5   atrophy and EEG components) is to increase the model's sensitivity to MCS (at the

Figure 5. *(a) ROC curve for the three binary logistic models classifying patient diagnosis (i.e., VS versus MCS) on the basis of demographic components alone (M1), demographic components and overall brain atrophy (M2), and demographic components, overall brain atrophy, and EEG components (M3). (b) Conditional estimate plots for each of the significant variables selected in the full model; top row: Demographic (age, sex) component, brain overall atrophy (i.e., NBV); bottom row: EEG total power component).*

cost of a loss of specificity).

Insert Table 2 about here

Finally, in terms of individual variables, as shown in Figure 5b and Table 2, the

full model selected one demographic component (age/sex; $\beta = 1.44$, $OR = 1.06$, $p =$

0.002), overall brain atrophy ($\beta = 1.27$, $OR = 3.56$, $p < 0.001$), as well as the EEG

total power ($\beta = 0.66$, $OR = 1.94$, $p = 0.017$).

Discussion

We report three main findings addressing the relationship between severity of the

impairment of consciousness, EEG spectral profile, and subcortical atrophy. First, our

results show that spectral profiles recorded with conventional EEG map onto specific

patterns of subcortical brain pathology (as observed with MRI), in line with theoretical

proposals (Schiff, 2010; Schiff, 2016). Indeed, we find that the ratio of fast (i.e., beta) to

slow (i.e., delta) frequencies component is negatively related to atrophy in thalamic regions

well known to be associated with severity of impairment after brain injury, as shown in

post-mortem (Adams et al., 2000) and *in vivo* (Lutkenhoff et al., 2015; Lutkenhoff et al.,

2013) studies. Damage within thalamus, along with functional (Monti et al., 2015)

and/or structural (Zheng et al., 2017) disconnection of thalamo-cortical projections, is

indeed central to current theories of recovery from severe brain injury (Schiff, 2010),

with greater de-afferentiation leading to a state of functional thalamic quiescence and,

through cell-intrinsic mechanisms (Crunelli et al., 2018), delta oscillations (Schiff,

2016; Schiff, Nauvel, & Victor, 2014). Indeed, interventions aimed at up-regulating

thalamic activity have been shown capable of enhancing behavioral responsiveness in

DOC patients (Schnakers & Monti, 2017). The origin of beta frequency oscillations in

the thalamo-cortical circuit remains unclear (Koelman & Lowery, 2019), nonetheless,

there are at least two lines of evidence connecting them to DOC. On the one hand, a

lesser state of thalamic de-afferentiation might result, through circuit-level

interactions, in a thalamic bursting component expected in the beta frequencies (Schiff

et al., 2014). On the other hand, beta oscillations have been associated with the globus

pallidus (and subthalamic nucleus), a region which we find strongly negatively

associated (particularly in its lateral segment), with the beta/delta component, and

which has been previously implicated in electrocortical and behavioral arousal in

animal models (Qiu, Vetrivelan, Fuller, & Lu, 2010) and behavioral arousal in several

pathologies including DOC (Lutkenhoff et al., 2015). Some evidence suggests that

beta oscillations can be induced by increased inhibition of the globus pallidus (Mirzaei

et al., 2017), which, in DOC, would imply a lesser dysfunction within the mesocircuit

(i.e., greater inhibitory input from the striatal medium spiny cells towards the

pallidum; Schiff 2010). Thus, whether of thalamic or pallidal origin, or a superposition

of the two, beta oscillations in DOC might mark a comparatively less disconnected

mesocircuit.

Second, similarly to results obtained in a different (large) cohort of chronic DOC

patients (Lutkenhoff et al., 2015), subscales of the CRS-R mapped onto different

underlying patterns of atrophy, with oromotor/verbal and communication subscales

component loading on left thalamic and bilateral putamen, as well as subregions of

brainstem, and the motor component loading mainly on extensive brainstem atrophy.

While the present results are similar to those reported previously (Lutkenhoff et al.,

2015), there also are important differences. For example, no association was found

between clinical measurement and brainstem atrophy in our previous work, and the

pattern of thalamic atrophy in the two studies is not fully overlapping. It is difficult,

however, to fully evaluate the meaning of these differences since the PCA components

extracted in the two studies loaded differently on each CRS-R subscale. Therefore,

although we do find a consistent set of regions correlating with CRS-R communication

subscale in the two studies, spanning left thalamus and putamen, the details of the

associations between all subscales and subcortical brain pathology remain to be fully

characterized. Third, consistent with prior work (Corchs et al., 2019; Estraneo et al.,

2016; Lechinger et al., 2013), we find that EEG spectral features are relevant to

diagnosing a patient's chronic state of consciousness (i.e., VS versus MCS), a decision

known to be susceptible to a relatively high misdiagnosis rate (Monti & Owen, 2010;

Monti et al., 2015; Schnakers, Giacino, Kalmar, Piret, & Lopez, 2006; Schnakers et

al., 2009). Specifically, EEG components, together with demographic information,

could correctly classify patients across the conscious/unconscious line (as behaviorally

defined) with ∼87% success, leveraging on demographic information (i.e., sex, age),

overall brain atrophy, and EEG features (i.e., total power and θ/δ components). It is

also noteworthy that in our data combination of EEG and MRI data is additive in terms

of enhancing discriminability across groups, strengthening the idea that multimodal

approaches are a desirable way of assaying different – and complementary – aspects of

DOC pathology (Coleman, Bekinschtein, Monti, Owen, & Pickard, 2009; Schiff,

2006) and, more broadly, brain function and health (Kumral et al., 2020; Nentwich et

al., 2020), particularly in the context of deriving predictive models from brain data

(Engemann et al., 2020). Despite the good concordance between behavior-based

diagnosis and the classification based on demographic, brain atrophy, and EEG

variables, the two approaches still disagree over one third of the cases. Seven MCS

patients were classified as being in VS and seven VS patients were classified as being

MCS. The MCS patients classified as VS by our analysis could reflect the fact that

behavioral diagnoses compress very different EEG profiles (for example, patients with

traumatic or anoxic etiology) into the same clinical category. Similarly, the VS

patients classified as MCS by our algorithm could either be a reflection of the variance

in the spectrum of oscillations that are compatible with a state of unconsciousness, or a

genuine misdiagnosis (Edlow et al., 2017; Monti et al., 2015; Owen et al., 2006).

Finally, in evaluating the above results, the reader should be mindful of some

limitations. First, our results are skewed by survivor bias effects; we might thus be

representing a spectrum of impairment which, while severe, excludes the even greater

damage present in patients who do not survive later than a year post injury. The same is

true with respect to the fact that 55 datasets had to be excluded from the collected sample.

While it would have been ideal to be able to retain more of the sample, conventional

quality control limited our ability to analyze the full dataset. Nonetheless, the final

analyzed sample is similar to that of other MR-based work in the field (e.g., (Demertzi et

al., 2015; Monti, Vanhaudenhuyse, et al., 2010; Stender et al., 2014; Vanhaudenhuyse et

al., 2010)). Furthermore, as shown in Table S1, the analyzed sample and the excluded

sample are matched on almost all demographic and clinical variables, suggesting that our

procedure, while affecting statistical power, is not introducing additional bias to the

collected sample. Second, while VS and MCS- subgroups exhibit the expected imbalance

across male and female patients (cf., (Lutkenhoff et al., 2015; Monti, Vanhaudenhuyse, et

al., 2010; Stender et al., 2014)), it should be noted that the MCS+ subgroup is entirely

composed of male patients, thus inviting caution in extrapolating results exclusive to this

subgroup (e.g., differences in relative power distribution). Second, unexpectedly, we

observe that MCS+ patients exhibit lower alpha relative power compared to MCS-

patients. Nonetheless, considering that the two groups are matched in terms of overall

relative power across alpha and beta frequencies (24.4% and 24.2% for MCS- and

MCS+, respectively), it is possible that the decreased relative power in the alpha band

observed in MCS+ is simply the converse of the increased relative power observed in

the beta band (both relative to MCS-). Although caution should be exercised in making

such inferences from a small subset of our data, it is conceivable that MCS- patients

could have a more "rigid" rhythmic EEG less susceptible to variations as compared to

MCS+ patients. Third, due to significant correlations across channels within and across

power bands, in order to perform the regression analyses presented above, we had to

first reduce the independent variables by means of a PCA. While conventional, it does

affect the interpretation of our results in as much as we cannot directly assess whether

the effects we report in mixed component (e.g., the β/δ component) are principally due

to either frequency or to their combination. Nonetheless, each of the three ratio

components correlates strongly with the "raw" ratio of each pair of frequencies

(calculated by taking the ratio of the average relative power across all channels;

specifically, r=0.97, r=0.87, and r=0.85 for β/δ, α/δ, and θ/δ, respectively) as well as

with each numerator and denominator variable (albeit numerically more with the

nominator for β/δ and α/δ components; see Table S2). This suggests that our results

can be reasonably interpreted as reflecting actual ratios in relative power. Third, gamma frequencies are known to often contain residual muscle artifacts. We decided to keep them in the analysis mainly because, even if they do contain artifacts, including this component still contributes to explaining variance in the signal. Had we not included it, any variance across patients due to motion would have *de facto* been subsumed by the unexplained variance term. In this sense, our approach is analogous to the conventional inclusion of motion parameters in functional MRI analysis. Fourth, while we report associations between brain damage in subcortical regions and EEG spectral features, this does not necessarily imply that the pinpointed areas are themselves the generators of specific oscillatory rhythms at rest.

Finally, it should also be pointed out that the present work did not include an additional independent sample to confirm the classification results, thus inviting caution in the extrapolation of the results towards new cohorts. As multi-site efforts increase, larger cohorts of high-quality data will permit full application of such approaches.

In conclusion, this work bridges different levels of analysis of patients surviving severe brain injury, uniting DOC brain pathology (Adams et al., 2000; Lutkenhoff et al., 2015; Lutkenhoff et al., In press), clinical evaluation (Giacino et al., 2004), and power spectral features (Brenner, 2005; Fingelkurts et al., 2013; Schnakers et al., 2019) in a framework that makes use of routine multimodal clinical data and informs both basic science and the diagnostic process. Furthermore, the present data also show that the pattern of association between spectral profile, brain damage, and clinical

variables observed in the acute setting (Schnakers et al., 2019) persist through the

chronic timeframe, in line with the idea that brain injury is best thought of as a long-

term disease as opposed to an "event" (Masel & DeWitt, 2010).

Author Contributions
SF, MMM, ML, MGB conceived the study; DRS, SF, EV, oversaw collection and clinical

interpretation of EEG data; LD, MGB, AN, CR oversaw collection and clinical

interpretation of MRI data; ESL, AN, MMM, carried out the analysis presented in the

paper; ESL, AN, SF, MMM, interpreted the results and drafted the manuscript; all authors

provided critical feedback on the manuscript and revised the manuscript; SF, MMM, MGB

secured funding.

Potential Conflicts of Interest

The Authors declare no conflicts of interest.

Ethical standards

The authors assert that all procedures contributing to this work comply with the ethical

standards of the relevant national and institutional committees on human experimentation

and with the Helsinki Declaration of 1975, as revised in 2008.

Funding sources

This work was supported by the Italian Ministry of Health (research grant RF-2013-

02359287) and by Tiny Blue Dot Foundation.

References

Adams, J. H., Graham, D. I., & Jennett, B. (2000). The neuropathology of the vegetative state after

an   acute   brain   insult.   *Brain,   123   (Pt   7)*,   1327-1338.

https://doi.org/10.1093/brain/123.7.1327

Bagnato, S., Boccagni, C., Prestandrea, C., Sant'Angelo, A., Castiglione, A., & Galardi, G. (2010).

Prognostic value of standard EEG in traumatic and non-traumatic disorders of

consciousness following coma. *Clinical Neurophysiology, 121*(3), 274-280.

https://doi.org/10.1016/j.clinph.2009.11.008

Brenner, R. P. (2005). The interpretation of the EEG in stupor and coma. *Neurologist, 11*(5), 271-

284. https://doi.org/10.1097/01.nrl.0000178756.44055.f6

Bruno, M. A., Vanhaudenhuyse, A., Thibaut, A., Moonen, G., & Laureys, S. (2011). From

unresponsive wakefulness to minimally conscious PLUS and functional locked-in

syndromes: recent advances in our understanding of disorders of consciousness. *Journal of*

Neurology, 258(7), 1373-1384. https://doi.org/10.1007/s00415-011-6114-x

Chennu, S., Finoia, P., Kamau, E., Monti, M. M., Allanson, J., Pickard, J. D., . . . Bekinschtein, T.

A. (2013). Dissociable endogenous and exogenous attention in disorders of consciousness.

Neuroimage Clin, 3, 450-461. https://doi.org/10.1016/j.nicl.2013.10.008

Coleman, M. R., Bekinschtein, T., Monti, M. M., Owen, A. M., & Pickard, J. D. (2009). A

multimodal approach to the assessment of patients with disorders of consciousness.

Progress in Brain Research, 177, 231-248. https://doi.org/10.1016/S0079-6123(09)17716-

6

Corchs, S., Chioma, G., Dondi, R., Gasparini, F., Manzoni, S., Markowska-Kacznar, U., . . .

Morreale, A. (2019). Computational Methods for Resting-State EEG of Patients With

Disorders   of   Consciousness.   *Frontiers   in   Neuroscience,   13*,   807.

https://doi.org/10.3389/fnins.2019.00807

Crone, J. S., Bio, B. J., Vespa, P. M., Lutkenhoff, E. S., & Monti, M. M. (2018). Restoration of

thalamo-cortical connectivity after brain injury: recovery of consciousness, complex

behavior, or passage of time? *Journal of Neuroscience Research, 96*(4), 671-687.

https://doi.org/10.1002/jnr.24115

Crone, J. S., Schurz, M., Holler, Y., Bergmann, J., Monti, M., Schmid, E., . . . Kronbichler, M.

(2015). Impaired consciousness is linked to changes in effective connectivity of the

posterior cingulate cortex within the default mode network. *Neuroimage, 110*, 101-109.

https://doi.org/10.1016/j.neuroimage.2015.01.037

Crunelli, V., Lorincz, M. L., Connelly, W. M., David, F., Hughes, S. W., Lambert, R. C., . . .

Errington, A. C. (2018). Dual function of thalamic low-vigilance state oscillations: rhythm-

regulation and plasticity. *Nature Reviews: Neuroscience, 19*(2), 107-118.

https://doi.org/10.1038/nrn.2017.151

Demertzi, A., Antonopoulos, G., Heine, L., Voss, H. U., Crone, J. S., de Los Angeles, C., . . .

Laureys, S. (2015). Intrinsic functional connectivity differentiates minimally conscious

from unresponsive patients. *Brain, 138*(Pt 9), 2619-2631.

https://doi.org/10.1093/brain/awv169

Demertzi, A., Tagliazucchi, E., Dehaene, S., Deco, G., Barttfeld, P., Raimondo, F., . . . Sitt, J. D.

(2019). Human consciousness is supported by dynamic complex patterns of brain signal

coordination. *Sci Adv, 5*(2), eaat7603. https://doi.org/10.1126/sciadv.aat7603

Edlow, B. L., Chatelle, C., Spencer, C. A., Chu, C. J., Bodien, Y. G., O'Connor, K. L., . . . Wu, O.

(2017). Early detection of consciousness in patients with acute severe traumatic brain

injury. *Brain, 140*(9), 2399-2414. https://doi.org/10.1093/brain/awx176

Engemann, D. A., Kozynets, O., Sabbagh, D., Lemaitre, G., Varoquaux, G., Liem, F., & Gramfort,

A. (2020). Combining magnetoencephalography with magnetic resonance imaging enhances learning of surrogate-biomarkers. *Elife, 9*. https://doi.org/10.7554/eLife.54055

Estraneo, A., Loreto, V., Guarino, I., Boemia, V., Paone, G., Moretta, P., & Trojano, L. (2016). Standard EEG in diagnostic process of prolonged disorders of consciousness. *Clinical Neurophysiology, 127*(6), 2379-2385. https://doi.org/10.1016/j.clinph.2016.03.021

Fingelkurts, A. A., Fingelkurts, A. A., Bagnato, S., Boccagni, C., & Galardi, G. (2013). The value of spontaneous EEG oscillations in distinguishing patients in vegetative and minimally conscious states. *Supplements to Clinical Neurophysiology, 62*, 81-99. https://doi.org/10.1016/b978-0-7020-5307-8.00005-3

Giacino, J. T., Kalmar, K., & Whyte, J. (2004). The JFK Coma Recovery Scale-Revised: measurement characteristics and diagnostic utility. *Archives of Physical Medicine and Rehabilitation, 85*(12), 2020-2029. https://doi.org/10.1016/j.apmr.2004.02.033

Koelman, L. A., & Lowery, M. M. (2019). Beta-Band Resonance and Intrinsic Oscillations in a Biophysically Detailed Model of the Subthalamic Nucleus-Globus Pallidus Network. *Frontiers in Computational Neuroscience, 13*, 77. https://doi.org/10.3389/fncom.2019.00077

Kumral, D., Sansal, F., Cesnaite, E., Mahjoory, K., Al, E., Gaebler, M., . . . Villringer, A. (2020). BOLD and EEG signal variability at rest differently relate to aging in the human brain. *Neuroimage, 207*, 116373. https://doi.org/10.1016/j.neuroimage.2019.116373

Laureys, S. (2005). The neural correlate of (un)awareness: lessons from the vegetative state. *Trends in Cognitive Sciences, 9*(12), 556-559. https://doi.org/10.1016/j.tics.2005.10.010

Lechinger, J., Bothe, K., Pichler, G., Michitsch, G., Donis, J., Klimesch, W., & Schabus, M. (2013). CRS-R score in disorders of consciousness is strongly related to spectral EEG at rest. *Journal of Neurology, 260*(9), 2348-2356. https://doi.org/10.1007/s00415-013-6982-3

Lehembre, R., Bruno, M. A., Vanhaudenhuyse, A., Chatelle, C., Cologan, V., Leclercq, Y., . . .

Noirhomme, Q. (2012). Resting-state EEG study of comatose patients: a connectivity and

frequency analysis to find differences between vegetative and minimally conscious states.

Functional   Neurology,   27(1),   41-47.   Retrieved   from   <u><Go   to</u>

<u>ISI>://WOS:000307137200007</u>

Leon-Carrion, J., Martin-Rodriguez, J. F., Damas-Lopez, J., Barroso y Martin, J. M., &

Dominguez-Morales, M. R. (2008). Brain function in the minimally conscious state: a

quantitative neurophysiological study. *Clinical Neurophysiology, 119*(7), 1506-1514.

<u>https://doi.org/10.1016/j.clinph.2008.03.030</u>

Lutkenhoff, E. S., Chiang, J., Tshibanda, L., Kamau, E., Kirsch, M., Pickard, J. D., . . . Monti, M.

M. (2015). Thalamic and extrathalamic mechanisms of consciousness after severe brain

injury. *Annals of Neurology, 78*(1), 68-76. <u>https://doi.org/10.1002/ana.24423</u>

Lutkenhoff, E. S., McArthur, D. L., Hua, X., Thompson, P. M., Vespa, P. M., & Monti, M. M.

(2013). Thalamic atrophy in antero-medial and dorsal nuclei correlates with six-month

outcome   after   severe   brain   injury.   *Neuroimage   Clin,   3*,   396-404.

<u>https://doi.org/10.1016/j.nicl.2013.09.010</u>

Lutkenhoff, E. S., Rosenberg, M., Chiang, J., Zhang, K., Pickard, J. D., Owen, A. M., & Monti, M.

M. (2014). Optimized brain extraction for pathological brains (optiBET). *PloS One, 9*(12),

e115551. <u>https://doi.org/10.1371/journal.pone.0115551</u>

Lutkenhoff, E. S., Wright, M. J., Shrestha, V., Real, C., McArthur, D. L., Buitrago-Blanco, M., . . .

Monti, M. M. (In press). The subcortical basis of outcome and cognitive impairment in

TBI: A longitudinal cohort study. *Neurology*, 669390.

Masel, B. E., & DeWitt, D. S. (2010). Traumatic brain injury: a disease process, not an event.

Journal of Neurotrauma, 27(8), 1529-1540. <u>https://doi.org/10.1089/neu.2010.1358</u>

Mirzaei, A., Kumar, A., Leventhal, D., Mallet, N., Aertsen, A., Berke, J., & Schmidt, R. (2017).

Sensorimotor Processing in the Basal Ganglia Leads to Transient Beta Oscillations during

Behavior.   *Journal   of   Neuroscience,   37*(46),   11220-11232.

https://doi.org/10.1523/JNEUROSCI.1289-17.2017

Monti, M. M. (2012). Cognition in the vegetative state. *Annual Review of Clinical Psychology, 8*,

431-454. https://doi.org/10.1146/annurev-clinpsy-032511-143050

Monti, M. M., Laureys, S., & Owen, A. M. (2010). The vegetative state. *BMJ, 341*(aug02 1),

c3765. https://doi.org/10.1136/bmj.c3765

Monti, M. M., & Owen, A. M. (2010). Behavior in the Brain Using Functional Neuroimaging to

Assess Residual Cognition and Awareness After Severe Brain Injury. *Journal of

Psychophysiology, 24*(2), 76-82. https://doi.org/10.1027/0269-8803/a000016

Monti, M. M., Rosenberg, M., Finoia, P., Kamau, E., Pickard, J. D., & Owen, A. M. (2015).

Thalamo-frontal connectivity mediates top-down cognitive functions in disorders of

consciousness.   *Neurology,   84*(2),   167-173.

https://doi.org/10.1212/WNL.0000000000001123

Monti, M. M., Vanhaudenhuyse, A., Coleman, M. R., Boly, M., Pickard, J. D., Tshibanda, L., . . .

Laureys, S. (2010). Willful modulation of brain activity in disorders of consciousness. *New

England Journal of Medicine, 362*(7), 579-589. https://doi.org/10.1056/NEJMoa0905370

Nentwich, M., Ai, L., Madsen, J., Telesford, Q. K., Haufe, S., Milham, M. P., & Parra, L. C.

(2020). Functional connectivity of EEG is subject-specific, associated with phenotype, and

different   from   fMRI.   *Neuroimage,   218*,   117001.

https://doi.org/10.1016/j.neuroimage.2020.117001

Nigri, A., Catricala, E., Ferraro, S., Bruzzone, M. G., D'Incerti, L., Sattin, D., . . . members, C. R.

C. C. R. C. (2017). The neural correlates of lexical processing in disorders of

consciousness. *Brain Imaging and Behavior, 11*(5), 1526-1537. https://doi.org/10.1007/s11682-016-9613-7

Owen, A. M., Coleman, M. R., Boly, M., Davis, M. H., Laureys, S., & Pickard, J. D. (2006). Detecting awareness in the vegetative state. *Science, 313*(5792), 1402. https://doi.org/10.1126/science.1130197

Patenaude, B., Smith, S. M., Kennedy, D. N., & Jenkinson, M. (2011). A Bayesian model of shape and appearance for subcortical brain segmentation. *Neuroimage, 56*(3), 907-922. https://doi.org/10.1016/j.neuroimage.2011.02.046

Qiu, M. H., Vetrivelan, R., Fuller, P. M., & Lu, J. (2010). Basal ganglia control of sleep-wake behavior and cortical activation. *European Journal of Neuroscience, 31*(3), 499-507. https://doi.org/10.1111/j.1460-9568.2009.07062.x

Rosazza, C., Andronache, A., Sattin, D., Bruzzone, M. G., Marotta, G., Nigri, A., . . . Coma Research Center, B. I. (2016). Multimodal study of default-mode network integrity in disorders of consciousness. *Annals of Neurology, 79*(5), 841-853. https://doi.org/10.1002/ana.24634

Rossi Sebastiano, D., Panzica, F., Visani, E., Rotondi, F., Scaioli, V., Leonardi, M., . . . Franceschetti, S. (2015). Significance of multiple neurophysiological measures in patients with chronic disorders of consciousness. *Clinical Neurophysiology, 126*(3), 558-564. https://doi.org/10.1016/j.clinph.2014.07.004

Rossi Sebastiano, D., Visani, E., Panzica, F., Sattin, D., Bersano, A., Nigri, A., . . . Franceschetti, S. (2018). Sleep patterns associated with the severity of impairment in a large cohort of patients with chronic disorders of consciousness. *Clinical Neurophysiology, 129*(3), 687-693. https://doi.org/10.1016/j.clinph.2017.12.012

Sacco, S., Altobelli, E., Pistarini, C., Cerone, D., Cazzulani, B., & Carolei, A. (2011). Validation of

the Italian version of the Coma Recovery Scale-Revised (CRS-R). *Brain Injury, 25*(5), 488-495. https://doi.org/10.3109/02699052.2011.558043

Schiff, N. D. (2006). Multimodal neuroimaging approaches to disorders of consciousness. *Journal of Head Trauma Rehabilitation, 21*(5), 388-397. https://doi.org/10.1097/00001199-200609000-00003

Schiff, N. D. (2010). Recovery of consciousness after brain injury: a mesocircuit hypothesis. *Trends in Neurosciences, 33*(1), 1-9. https://doi.org/10.1016/j.tins.2009.11.002

Schiff, N. D. (2016). Mesocircuit mechanisms underlying recovery of consciousness following severe brain injuries: model and predictions. In *Brain Function and Responsiveness in Disorders of Consciousness* (pp. 195-204): Springer.

Schiff, N. D., Nauvel, T., & Victor, J. D. (2014). Large-scale brain dynamics in disorders of consciousness. *Current Opinion in Neurobiology, 25*, 7-14. https://doi.org/10.1016/j.conb.2013.10.007

Schnakers, C., Giacino, J., Kalmar, K., Piret, S., & Lopez, E. (2006). Does the FOUR Score Correctly Diagnose the Vegetative and Minimally Conscious States ? Tomographic Visualization of Cholinesterase. *Annals of Neurology, 60*(6), 2005-2006.

Schnakers, C., Lutkenhoff, E. S., Bio, B. J., McArthur, D. L., Vespa, P. M., & Monti, M. M. (2019). Acute EEG spectra characteristics predict thalamic atrophy after severe TBI. *Journal of Neurology, Neurosurgery and Psychiatry, 90*(5), 617-619. https://doi.org/10.1136/jnnp-2017-317829

Schnakers, C., & Monti, M. M. (2017). Disorders of consciousness after severe brain injury: therapeutic options. *Current Opinion in Neurology, 30*(6), 573-579. https://doi.org/10.1097/WCO.0000000000000495

Schnakers, C., Perrin, F., Schabus, M., Majerus, S., Ledoux, D., Damas, P., . . . Laureys, S. (2008).

Voluntary brain processing in disorders of consciousness. *Neurology, 71*(20), 1614-1620. https://doi.org/10.1212/01.wnl.0000334754.15330.69

Schnakers, C., Vanhaudenhuyse, A., Giacino, J., Ventura, M., Boly, M., Majerus, S., . . . Laureys, S. (2009). Diagnostic accuracy of the vegetative and minimally conscious state: clinical consensus versus standardized neurobehavioral assessment. *BMC Neurology, 9*, 35. https://doi.org/10.1186/1471-2377-9-35

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage, 44*(1), 83-98. https://doi.org/10.1016/j.neuroimage.2008.03.061

Smith, S. M., Zhang, Y., Jenkinson, M., Chen, J., Matthews, P. M., Federico, A., & De Stefano, N. (2002). Accurate, robust, and automated longitudinal and cross-sectional brain change analysis. *Neuroimage, 17*(1), 479-489. https://doi.org/10.1006/nimg.2002.1040

Stender, J., Gosseries, O., Bruno, M. A., Charland-Verville, V., Vanhaudenhuyse, A., Demertzi, A., . . . Laureys, S. (2014). Diagnostic precision of PET imaging and functional MRI in disorders of consciousness: a clinical validation study. *Lancet, 384*(9942), 514-522. https://doi.org/10.1016/S0140-6736(14)60042-8

Vanhaudenhuyse, A., Noirhomme, Q., Tshibanda, L. J., Bruno, M. A., Boveroux, P., Schnakers, C., . . . Boly, M. (2010). Default network connectivity reflects the level of consciousness in non-communicative brain-damaged patients. *Brain, 133*(Pt 1), 161-171. https://doi.org/10.1093/brain/awp313

Zheng, Z. S., Reggente, N., Lutkenhoff, E., Owen, A. M., & Monti, M. M. (2017). Disentangling disorders of consciousness: Insights from diffusion tensor imaging and machine learning. *Human Brain Mapping, 38*(1), 431-443. https://doi.org/10.1002/hbm.23370

Tables

Table 1: Analyzed sample summary statistics of demographic, clinical, and electroencephalographic data per diagnostic group. (Abbreviations: VS, Vegetative State; MCS-Minimally conscious State "minus"; MCS+, Minimally Conscious State "plus"; MPI, months post-injury; T, traumatic; NT, non-traumatic; H, hemorrhagic injury, I, ischemic injury, A, anoxic; Orom/Verb, oromotor/verbal; Communic, communication.) (See Table S1 for a summary of demographic and clinical variables for the excluded datasets and comparison with the analyzed sample.)

	VS	MCS-	MCS+
Age (mean, SD, y)	53 (±14.21)	54 (±18.77)	46 (±16.77)
MPI (mean, SD, mo)	23 (±16.05)	44 (±37.17)	53 (±64.97)
Sex	14F, 23M	11F, 6M	0F, 7M
Etiology	12T, 9H, 1I, 1HI, 14TA	5T, 7H, 1I, 0HI, 4TA	3T, 3H, 1I, 0HI, 0TA

Coma Recovery Scale Revised (CRS-R)			
Total score	6.24 (1.01)	9.24 (1.25)	10.43 (1.90)
Auditory	1.03 (0.50)	1.41 (0.51)	2.00 (1.15)
Visual	0.76 (0.43)	2.59 (0.71)	2.43 (1.40)
Motor	1.92 (0.28)	2.12 (0.33)	2.14 (0.90)
Orom/Verb	1.00 (0.33)	1.24 (0.56)	1.43 (0.98)
Communic	0.00 (0.00)	0.12 (0.33)	0.57 (0.53)
Arousal	1.54 (0.56)	1.76 (0.56)	1.86 (0.90)

Electroencephalography (Total power & frequency relative power)			
Total power (μV^2)	138.22 (140.35)	229.20 (261.81)	178.96 (148.15)
Delta (1–4Hz)	49.46% (13.46%)	41.51% (14.35%)	42.57% (16.97%)
Theta (4–8Hz)	30.21% (11.09%)	34.09% (9.73%)	33.23% (11.58%)
Alpha (8–13Hz)	9.71% (5.64%)	11.89% (6.08%)	8.33% (4.48%)
Beta (13–30Hz)	7.78% (7.09%)	9.59% (8.22%)	13.89% (17.23%)
Gamma (>30Hz)	2.84% (3.00%)	2.91% (3.21%)	1.98% (1.63%)

Table 2: Binary logistic regression results. Top: comparison of the area under the curve, sensitivity, specificity, and precision of the model with demographic components only, the model with demographic components and the global brain atrophy factor, and the model with demographic components, global brain atrophy factor, and EEG components. Bottom: Individual predictors selected for the full model (i.e., DEM+Brain+EEG). (Abbreviations: area under the curve, AUC; sensitivity, sens.; specificity, spec.; precision, prec.)

Model	df	ΔX²	p	R^2	AUC	Sens	Spec	Prec
Intercept	60							
DEM	59	8.93	0.003	0.18	0.67	0.29	0.89	0.64
DEM+Brain	58	16.66	<0.001	0.32	0.78	0.58	0.78	0.64
DEM+Brain+EEG	54	29.09	<0.001	0.51	0.87	0.79	0.81	0.73

							95% CI	
Parameter	b	SE	β	OR	z	p	LB	UB
(Intercept)	-1.92	0.50	-0.79	0.15	-3.86	<0.001	-2.89	-0.95
DEM: Sex-Age	0.05	0.02	1.44	1.06	3.15	0.002	0.02	0.09
Normalized Brain	1.27	0.37	1.27	3.56	3.44	<0.001	0.55	1.99
EEG: Total power	0.66	0.28	0.66	1.94	2.39	0.017	0.12	1.21
EEG: $\delta/\alpha(RH)$	0.63	0.57	0.63	1.88	1.11	0.269	-0.49	1.75
EEG: θ/δ	0.61	0.32	0.61	1.84	1.88	0.060	-0.03	1.25
EEG: $\alpha\theta(P,O)[Neg.]$	0.59	0.36	0.59	1.80	1.65	0.100	-0.11	1.29

Figure Legends

Figure 1. Sample structure extraction (left) and 3-dimensional triangle vertex mesh (right). (Abbreviations: A, anterior; BrStem, brain stem; Caud, caudate; GP, globus pallidus; Hipp, hippocampus; L, left; P, posterior; Putm, putamen; R, right; Thal, thalamus. Figure from (Lutkenhoff et al., 2015).)

Figure 2. Summary of EEG data. Total and relative power at each frequency band (to allow displaying on the same axis, values are normalized within each variable). (Error bars represent standard errors.)

Figure 3. Results for the rest EEG – MRI analysis. (a) Results for the β/δ ratio component; (b) results for the θ/δ component; (c) results for the total power component. Warmer colors indicate regions with greater atrophy; gray indicates no significant effect.

Figure 4. Results for the CRS-R – MRI analyses. (a) Results for the CRS-R motor component; (b) results for the CRS-R oromotor-communication component. (See Fig. 3 for color interpretation.)

Figure 5. (a) ROC curve for the three binary logistic models classifying patient diagnosis (i.e., VS versus MCS) on the basis of demographic components alone (M1), demographic components and overall brain atrophy (M2), and demographic components, overall brain atrophy, and EEG components (M3). (b) Conditional estimate plots for each of the

significant variables selected in the full model; top row: Demographic (age, sex) component, brain overall atrophy (i.e., NBV); bottom row: EEG total power component, EEG theta/delta component).

Supplementary Online Materials for

EEG power spectra and subcortical pathology in chronic disorders of consciousness

Evan S. Lutkenhoff, Ph.D, Anna Nigri, Ph.D, Davide Rossi Sebastiano, MD, Ph.D, Davide Sattin, PsyD, Elisa Visani, M.Sc., Cristina Rosazza, Ph.D., Ludovico D'Incerti, MD, Maria Grazia Bruzzone, MD, Silvana Franceschetti, MD, Matilde Leonardi, MD, Stefania Ferraro, Ph.D, Martin M. Monti Ph.D.

Contents:

Table S1. Comparison of demographic and clinical variables of patients included in the final analyzed sample (n-61) and patients excluded during the MRI processing/QC step (n=55). (Abbreviations: VS, Vegetative State; MCS- Minimally conscious State "minus"; MCS+, Minimally Conscious State "plus"; MPI, months post-injury; T, traumatic; H, hemorrhagic injury, I, ischemic injury, A, anoxic. '†' indicates significant difference across groups prior to correction for multiple comparisons. No significant differences were

	Included (n=61)	Excluded (n=55)
Age (years)	51.04 (15.38)	50.24 (13.25)
MPI (months)†	32.45 (32.81)	51.45 (50.79)
Sex	25F, 36M	22F, 33M
Etiology	20T, 19H, 3I, 1HI, 18A, 0TA	15T, 18H, 0I, 0HI, 21A, 1TA
Diagnosis	37 VS, 17 MCS-, 7 MCS+, 0 E-MCS	36 VS, 11 MCS-, 7 MCS+, 1 E-MCS
Coma Recovery Scale Revised (CRS-R)		
Total score	7.56 (2.05)	8.13 (2.46)
Auditory	1.25 (0.67)	1.45 (0.83)
Visual	1.46 (1.10)	1.62 (1.34)
Motor	2.00 (0.41)	2.00 (0.38)
Oromotor/Verbal	1.11 (0.52)	1.07 (0.38)
Communication	0.10 (0.30)	0.09 (0.29)
Arousal†	1.69 (0.53)	1.89 (0.42)

observed after correction for multiplicity.)

Table S2. Spearman correlation between the three ratio components (β/δ, α/δ, and θ/δ) and the raw ratio of each pair of frequencies (calculated by taking the ratio of the average relative power across all channels), as well as the numerator frequency (β, α, and θ, respectively) and the denominator frequency (δ for all components).

	raw ratio	numerator	denominator
β/δ comp	0.95	0.94	-0.61
α/δ comp	0.87	0.94	-0.78
θ/δ comp	0.85	0.79	-0.80

Figure S1. Examples of excluded dataset (for each category) during the MRI quality control process.

Figure S2. Mean intensity of each EEG component across the three DOC clinical categories. (Error bars show standard error.).