Inoculation with the mycorrhizal fungus *Rhizophagus irregularis* modulates

the relationship between root growth and nutrient content in maize (*Zea mays*

ssp. *mays* L.)

M. Rosario Ramírez-Flores[1], Elohim Bello-Bello[2], Rubén Rellán-Álvarez[2, 3], Ruairidh J. H.

Sawers[2,4*] and Víctor Olalde-Portugal[1*]

[1] Departamento de Biotecnología y Bioquímica, Centro de Investigación y de Estudios

Avanzados (CINVESTAV-IPN), Irapuato, C.P. 36821, Guanajuato, México.

[2] Laboratorio Nacional de Genómica para la Biodiversidad/Unidad de Genómica Avanzada,

Centro de Investigación y de Estudios Avanzados, Instituto Politécnico Nacional (CINVESTAV-

IPN), Irapuato, C.P. 36821, Guanajuato, México.

[3] Department of Molecular and Structural Biochemistry, North Carolina State University, NC,

USA.

[4] Department of Plant Science, The Pennsylvania State University, State College, PA, USA.

* Author for correspondence rjs6686@psu.edu; v_olalde@yahoo.com.mx

ABSTRACT

Plant root systems play an essential role in nutrient and water acquisition. In resource-limited

soils, modification of root system architecture is an important strategy to optimize plant

performance. Most terrestrial plants also form symbiotic associations with arbuscular

mycorrhizal fungi to maximize nutrient uptake. In addition to direct delivery of nutrients,

arbuscular mycorrhizal fungi benefit the plant host by promoting root growth. Here, we aimed to

quantify the impact of arbuscular mycorrhizal symbiosis on root growth and nutrient uptake in

maize. Inoculated plants showed an increase in both biomass and the total content of twenty

quantified elements. In addition, image analysis showed mycorrhizal plants to have denser, more

branched root systems. For most of the quantified elements, the increase in content in

mycorrhizal plants was proportional to root and overall plant growth. However, the increase in

boron, calcium, magnesium, phosphorus, sulfur and strontium was greater than predicted by root

system size alone, indicating fungal delivery to be supplementing root uptake.

KEYWORDS: mycorrhiza, root development, ionome, maize

INTRODUCTION

Plant productivity is typically limited by the availability of nutrients, in both natural and agricultural ecosystems. Under nutrient-poor conditions, plants have the capacity to modulate the architecture and functionality of their root system, potentially increasing nutrient uptake (Lynch, 1995). Beyond a general reallocation of resources from leaf to root, there are nutrient-specific changes in the development of primary (PR) and lateral (LR) roots, building a root system architecture (RSA) that may better optimize resource acquisition (López-Bucio et al., 2003; Osmont et al., 2007; Gruber et al., 2013). Nutrients are heterogeneously distributed in the soil, and plants respond to local concentration, allocating greater root production to regions of higher availability (Campbell et al., 1991; Farley and Fitter, 1999; Grossman and Rice, 2012). In addition, nutrient distribution varies across soil horizons: poorly-mobile nutrients such as phosphorus (P), potassium (K), magnesium (Mg) or calcium (Ca) are typically enriched in topsoil, while more mobile nutrients, such as nitrogen (N), are typically more abundant deeper in the soil (Rubio et al., 2003; Ho et al., 2004; Lynch and Brown, 2008; Postma et al., 2014; Rangarajan et al., 2018). As a consequence, researchers have distinguished RSAs optimized for topsoil versus deeper foraging (Lynch, 2019).

Root development is regulated by the combined action of internal developmental pathways and external environmental stimuli (Malamy and Ryan, 2001), conditioning both plasticity and intra- and inter-specific variation. These pathways are based on the action of plant hormones, signal receptors, transcription factors and secondary messengers, including Ca^{2+}, nitric oxide and reactive oxygen species (Fukaki and Tasaka, 2009; Garay-Arroyo et al., 2012; Jung and McCouch, 2013; Schlicht et al., 2013; Zhang et al., 2015; Shahzad and Amtmann, 2017). Auxin works with cytokinin to regulate LR initiation (Aloni et al., 2006), and with

gibberellin to modulate cell proliferation and elongation (Fu and Harberd, 2003). Strigolactones

impact LR formation and root hair development in a dose-dependent manner (Koltai, 2011), and,

in maize, promote nodal root development (Guan et al., 2012). RSA is largely a product of the

balance between root elongation and branching (Postma et al., 2014). Primary root growth will

continue if the root meristem is active and populations of stem cells within the quiescent center

are maintained. In *Arabidopsis*, the exhaustion of the primary root meristem under low P is

considered the classic example of a plastic response to optimize topsoil foraging (Williamson et

al., 2001; López-Bucio et al., 2002; Mora-Macías et al., 2017). Root branching, through the

production of first- or higher- order LRs, is regulated at the level of the formation of LR

primordia and their subsequent expansion. Low availability of N, P, sulfur (S) or zinc (Zn)

promotes an increase in the density and/or length of LRs, although, as mentioned above, when

the nutrient distribution is patchy, LRs may proliferate in regions of high local nutrient

abundance (Zhang et al., 1999; Kutz et al., 2002; López-Bucio et al., 2002; Bouranis et al., 2008;

Gruber et al., 2013). In cereal crops, the bulk of the adult root system is comprised of shoot-

borne crown (CR) roots and associated LRs (Hochholdinger and Tuberosa, 2009). While greater

branching will increase the surface area for nutrient uptake, it becomes inefficient if placement

of roots in close proximity results in competition for the same nutrients, an effect that will be

greater for more mobile nutrients (Postma et al., 2014).

Significant intra-specific variation has been observed in RSA and plasticity, and has been

linked to superior agronomic performance in crops under specific edaphic conditions. In

common bean, varieties producing shallow basal roots and a large number of adventitious roots

explore the topsoil more efficiently, performing better in low P soils (Rubio et al., 2003; Lynch

and Brown, 2008). In rice, the PHOSPHATE STARVATION TOLERANCE 1 (PSTOl1) kinase,

identified from the traditional variety Kasalath, is associated with enhanced early root growth,

and increases yield under low P conditions (Wissuwa et al., 2005; Gamuyao et al., 2012).

83   In addition to relying on their roots to acquire nutrients, most terrestrial plants also form

mutualistic symbioses with arbuscular mycorrhizal (AM) fungi of the Phylum Glomeromycota

(Parniske, 2008; Smith and Read, 2010). The association with AM fungi is one of the oldest

plant symbioses, and AM symbioses were likely important during plant colonization of the land,

performing essential functions before the evolution of the vascular root system (Schüßler et al.,

2001). Establishment of AM symbiosis requires a complex interchange of signals between plant

and fungus that results in the entry of the fungus into the root and the development of highly

branched arbuscules within the root cortical cells, providing the primary site of nutrient exchange

(Parniske, 2008). AM fungi are obligate symbionts, obtaining carbon from their plant host in

return for providing nutrients and water that are acquired by an extensive network of root-

external hyphae (Bago et al., 2003). The clearest benefit to the plant is enhanced P uptake (Chiu

and Paszkowski, 2019), although AM fungi have been reported to promote uptake of other

elements, including N, Fe, S and Zn (Liu et al., 2000; González-Guerrero et al., 2005;

Govindarajulu et al., 2005; López-Pedrosa and González-Guerrero, 2006; Allen and Shachar-

Hill, 2009). In practice, complex interactions between nutrients influence the outcome of the

symbiosis with respect to any given element (Liu et al., 2000; Gerlach et al., 2015; Ramírez-

Flores et al., 2017). Although AM symbioses are widespread in both natural ecosystems and

cultivated fields, the plant host maintains a degree of control over establishment and the extent of

colonization, rejecting the fungus under high nutrient conditions (Nouri et al., 2014).

AM fungi not only directly deliver nutrients, but also promote growth of the host plant

roots, with secondary effects on nutrient uptake. General improvements in plant health under

AM symbiosis may be correlated with increased root system size (Berta et al., 1995; Tisserant et

al., 1996; Gutjahr et al., 2009; Sawers et al., 2017). AM symbiosis is also associated with

developmental changes, such as an increase in LR production and greater root branching (Berta

et al., 1990; Paszkowski and Boller, 2002; Oláh et al., 2005; Gutjahr et al., 2009). The failure to

form LR primordia in the maize *lateral rootless1* (*lrt1*) mutant can be partially overcome by AM

symbiosis, indicating an influence on plant developmental pathways (Hochholdinger and Feix,

1998; Paszkowski and Boller, 2002). In *Medicago truncatula,* developing fungal spores are

sufficient to trigger LR formation (Oláh et al., 2005; Gutjahr et al., 2009). In rice, AM fungi can

induce LR formation in *pollux*, *ccamk* and *cyclops* mutants, even though symbiosis is not

established, confirming that RSA modification need not be dependent on enhanced host plant

nutrition (Gutjahr et al., 2009).

The impact of AM symbiosis on nutrient uptake can be quantified by comparing the

concentration or content of a given element in the aerial portion of mycorrhizal (M) and non-

colonized (NC) plants (Gerlach et al., 2015; Ramírez-Flores et al., 2017). Such a comparison,

however, does not distinguish direct hyphal nutrient delivery from the secondary consequences

of fungus-induced modification of root architecture or the effects of greater root growth resulting

from a general increase in plant vigor. In the case of P, elegant labelling experiments and

extensive functional studies have clearly demonstrated the direct role of root-external hyphae in

nutrient delivery (Pearson and Jakobsen, 1993; Smith et al., 2003; Chiu and Paszkowski, 2019).

The picture, however, remains less clear with respect to other mineral nutrients.

In this study, we characterized RSA and leaf ionome in young maize plants, grown with

or without inoculation with AM fungi. For each nutrient, we asked whether any increase in

uptake was explainable by changes in the root system alone, or whether there was evidence of

additional hyphal foraging. For an increase resulting from greater root growth, we expected the

relationship between root surface area and uptake to be maintained, albeit that mycorrhizal plants

themselves were larger. If, however, there was a significant effect of hyphal foraging, we

predicted an increase in uptake per unit surface area of root.

MATERIALS AND METHODS

Plant material and growth conditions

Two maize inbred lines (*Zea mays* ssp. *mays* var. B73 and W22) were grown with (M) or

without (NC) inoculation with fungus, in PVC tubes (1 m in height, 15 cm in diameter). A total

of 64 plants (2 genotypes x 2 treatments x 16 replicates) were planted in complete blocks across

four planting dates. Each planting consisted of 16 plants (2 genotypes x 2 treatments x 4

replicates), with a one week interval between planting dates. B73 seed was produced in the

winter of 2015 in Valle de Banderas, Nayarit, México. W22 seed was produced in the summer of

2013 in Aurora, New York, USA. Each tube was filled with 17 L of a sterilized substrate mix

consisting of sand:perlite:silt (4.5:1.5:1, v/v). The substrate Olsen P concentration was 4.8 ppm.

For M plants, we inoculated each tube with ~700 *Rhizophagus irregularis* spores collected from

a commercial liquid inoculant (AGTIV®) and delivered to the middle of the tube at 15 cm depth

before planting. The experiment was conducted under greenhouse conditions, at an average

temperature of 24 ºC and humidity of 48%. Maize seeds were surface sterilized with 1% sodium

hypochloride solution for 10 min and rinsed five times with sterile distilled water. Seeds were

then placed in 30 ml of sterile distilled water and shaken for 48 hours before planting. From 5

days after emergence (DAE), plants were watered every other day with 200mL of ⅓ Hoagland

solution, modified to a final P concentration of 25 µM (Hoagland and Broyer, 1936). Plants were

harvested at 56 DAE.

Evaluation of plant growth

Leaf length and width were measured manually, and leaf area estimated as 0.75 x length x width

(Mollier and Pellerin, 1999). Shoot and root fresh weights were measured after cutting the aerial

part and washing the root. Dry weight was determined by oven drying tissue at 70º for 48 hrs.

Root volume was measured by submergence in a test tube and recording of the volume of water

displaced.

Estimation of fungal colonization in seedling roots

Lateral root segments were collected at random from the upper 15 cm of the root system, placed

in 10% KOH solution, autoclaved for 10 min, and stained with 0.05% trypan blue in

acetoglycerol. The percentage of root length colonized was quantified in 15 root pieces per plant

using a modified grid-line intersect method (McGonigle et al., 1990).

Characterization of root system architecture by image analysis

The cleaned root system was placed in a water-filled tub and photographed using a digital Nikon

camera D3500. Raw images were individually processed using Adobe® Photoshop® CC (Version

14.0) to remove background and increase contrast. Processed images were scaled and analyzed

using GiA Roots software (Galkovskyi et al. 2012). After scale calibration, the images were

segmented using the adaptive thresholding method and basic parameter settings were manually

optimized. Finally, all binary images were analyzed to quantify the root system traits. Stacked

(overlay) root system images were generated using the stacks tool in ImageJ/Fiji (version 2.0.0).

Root traits are described in Supplementary Table S1.

Analysis of the leaf ionome

The third youngest leaf (variously, leaf 5 or 6 for different individuals) was collected for ionomic

analysis using inductively coupled plasma mass spectrometry (ICP-MS) as described previously

(Ziegler et al. 2013, Ramírez-Flores et al., 2017). Briefly, weighed tissue samples were digested

in 2.5 ml of concentrated nitric acid (AR Select Grade, VWR) with an added internal standard

(20 p.p.b. In, BDH Aristar Plus). The concentration of the elements B, Na, Mg, Al, P, S, K, Ca,

Mn, Fe, Co, Ni, Cu, Zn, As, Se, Rb, Sr, Mo and Cd was measured using an Elan 6000 DRC-e

mass spectrometer (Perkin-Elmer SCIEX) connected to a PFA microflow nebulizer (Elemental

Scientific) and Apex HF desolvator (Elemental Scientific). A control solution was run every 10th

sample to correct for machine drift both during a single run and between runs. Given that

samples were digested in concentrated nitric acid prior to analysis and run in a 70% N

atmosphere, it was not possible to accurately estimate N concentration.

Statistical analyses and data visualization

All statistical analyses were performed in R (R Core Team, 2019). From the initial planting, a

small number of plants did not germinate. In addition, we removed clear outliers (assessed by

visual inspection of plant growth and health) and any plants assigned to the non-inoculated group

that showed evidence of root-internal hyphal structures. The final dataset consisted of 53

individuals (Table S2). Leaf surface area estimates were square root transformed for analysis. A

linear fixed-effect model was applied on a trait-by-trait basis to control for differences between

the four planting dates. Principal component (PC) analysis was performed on Gia Roots traits

along with root fresh (RFW) and dry weight (RDW), using R/ade4::dudi.pca (Dray and Dufour,

2007) with centered and scaled values. Linear Discriminant (LD) analysis on PC scores was

performed with R/MASS::lda (Venables and Ripley, 2002). Ionomic analysis generated element

concentration data; in addition, total element content was estimated as the product of

concentration and shoot dry weight (SDW).

Data were analyzed using R/stats::lm to fit inoculation status and genotype in a complete

model. After adjustment for multiple testing (R/stats::p.adjust; method = "BH"), no significant (p

< 0.05) interaction between inoculation x genotype was detected for any trait, and we present

here the additive model for simplicity. We also performed a one-way ANOVA with a single

four-level treatment factor (B73.NC, B73.M, W22.NC and W22.M) that was used to assign

means groups with R/agricolae::HSD.test (de Mendiburu, 2019) for presentation in Fig. S1. Box

plots were generated with base R using default settings. Root images were pre-processed using

imagemagick (www.imagemagick.org), converting them to png format, setting the background

to transparent and reducing the alpha level. Images were imported into R using R/png and plotted

onto the PCA/LD space using the rasterImage function. Pearson's correlation coefficients were

calculated using R/Hmisc::rcorr (Harrell, 2019) and visualized using R/gplots::heatmap.2

(Warnes et al., 2019). The impact of root system size and inoculation on element content, was

assessed using the model *Content = NSA + Inoculation*, evaluating the *Inoculation* term using

sequential (Type I) sum-of-squares (implemented by default in R). For comparison, partial (Type

III) sum-of-squares were calculated using R/car::Anova (Fox & Weisberg, 2011), setting the

"contrasts" option to c("contr.sum", "contr.poly").

RESULTS

Growth of maize seedlings increased when inoculated with *Rhizophagus irregularis*

To evaluate the relationship between root system architecture (RSA) and nutrient acquisition in

mycorrhiza plants, we grew the two maize inbred lines B73 and W22 under low phosphorus (P)

conditions, with (M) or without (NC) inoculation with the AM fungus *Rhizophagus irregularis*

(Fig. 1A). These two widely used inbred lines were selected to provide a reference for future

work, and to allow greater generalization than would be possible based on a single genotype.

B73 is the reference for the most complete maize genome assembly. W22 is the background for

the primary publicly available maize reverse genetics collections. Inoculation with AM fungus

significantly enhanced plant growth in both genotypes (Fig. 1B, S1, S2. Table 1). Fungal

inoculation and plant genotype were both significant ($p < 0.001$) predictors of plant biomass but

there was no evidence of an interaction between the two (Fig. S2, Table 1; S2). By harvest (55

DAE), the marginal effect of inoculation on shoot dry weight (SDW) was an increase of 142%

(Table 1). In both M and NC treatments, B73 tended to be smaller than W22, the range of SDW

overlapping between B73 M and W22 NC treatments (Fig. 1, S1, S2). The inclusion of plants

similar in size, yet differing in inoculation status, was important for our subsequent interpretation

of element content.

To quantify mycorrhizal colonization, the percentage of root length containing different

fungal structures was estimated by microscopic inspection. NC plants were confirmed to be free

from colonization (fungal structures were observed in three plants in the NC group; these

individuals were not included in the analysis), while inoculated plants showed fungal structures

typical of the symbiosis (hyphae, arbuscules and vesicles). M plants were well colonized (Fig.

1C-E), and there was no difference in colonization rate between the two plant genotypes (Fig S1.

Table S2).

The root system was modified by inoculation with *Rhizophagus irregularis*

Root fresh and dry weight (RFW and RDW) and root volume (RV) increased significantly in

inoculated plants (Fig. 2A. Table 1). To characterize RSA, we photographed the plants and

analyzed the images with GiA Roots (General Image Analysis of Roots; Galkovskyi et al.,

2012). Ten GiA Roots traits showed a significant response to inoculation (Table 1. Fig. S1). In a

principal component (PC) analysis using all 19 GiA Roots traits, the first four PCs explained

90% of the total variance (Fig. 2B; 2C). PC1 (explaining 59% of the total variance) was

dominated by variables associated with overall root system and plant size (Fig. 2; 4A). PC2

(explaining 12% of the total variance) was associated with overall root system shape and aspect

ratio (Fig. 2; 4A). PC3 (7%) and PC4 (7%) captured aspects of root branching and root system

solidity (Fig. 3; 4A). NC and M plants were well differentiated by PC1 and, to a lesser extent, by

PC3, indicating a shift towards larger, more branched/solid root systems in M plants. GiA Roots

evaluated solidity as the total network area divided by the network convex area. Although,

greater solidity was clear in M W22, the root system in the smallest B73 NC plants was also

considered relatively solid because the convex area itself was so low. We performed a linear

discriminant (LD) analysis using the root PC values, reinforcing the separation of M and NC

plants by size and solidity of the root system and better distinguishing the W22 M and B73 NC

treatments (Fig. 3).

Nutrient content was correlated with root system size and mycorrhizal colonization

AM fungi can increase plant nutrient uptake by delivery of nutrients via the hyphal network and

as a secondary consequence of promoting root growth. To better understand the relative

importance of these two factors, we considered the relationship between root system size and

mineral nutrient content. We determined the concentration of 20 different elements, including P,

in leaf tissue using inductively coupled plasma mass spectrometry (ICP-MS). Although leaf

element concentration provides an endpoint readout of various aspects of plant nutrient relations

(Salt et al., 2008), it was used here as a proxy for nutrient uptake. We estimated nutrient content

as the product of leaf concentration and total shoot dry weight and investigated the relationship

with the GiaRoots trait NSA (Network Surface Area), used as our measure of root system size.

As expected, an increase in NSA (associated with larger plants) was correlated with greater

nutrient content (Fig. 4B). Of greater interest, however, was the degree to which this relationship

was modified by AM colonization; *i.e.* for any given nutrient, was NSA sufficient to explain

element content in M plants? We ran a linear model using NSA and inoculation status as

predictors, assessing inoculation based on sequential (Type I) sum-of-squares, evaluating any

fungal effect beyond that explained by NSA alone (Fig. 5). A significant positive effect of

inoculation was associated with boron (B), Ca, Mg, P, S and strontium (Sr), indicating a unit of

root surface area to be associated with a greater leaf content of these elements in inoculated

plants.

DISCUSSION

Inoculation with *Rhizophagus irregularis* resulted in increased growth of maize grown under low

P availability. In comparison with a previous small pot evaluation (Sawers et al., 2017), plants

were larger, and the proportional increase in M plants was greater, presumably reflecting less

growth inhibition in the larger tubes that we used here. A difference in leaf surface area was

observed by day 40 - 45 after emergence, broadly consistent with the timing reported in rice for

the first observation of arbuscules and the accumulation of transcripts encoding mycorrhiza-

associated P transporters (Gutjahr et al., 2008). AM colonization was correlated not only with an

increase in root system size, but also in the degree of branching and solidity of the root system

(as captured by the GiA Roots traits MaxNR, MNR, NS). Inoculation with AM fungi has

previously been shown to promote root growth and branching in diverse plant hosts (Berta et al.,

1990; Paszkowski and Boller, 2002; Oláh et al., 2005; Gutjahr et al., 2009).

Increased growth was accompanied by greater concentration of Ca, Mn, P and Sr in the

leaves of M plants. The leaf concentration of K and Se was reduced in M plants, but for all

elements, taking biomass into account, total content increased. The content of B, Ca, Mg, P, S

and Sr following inoculation exceeded expectation based on root surface area alone. We interpret

this to reflect the impact of hyphal foraging, although we do not discount additional

contributions of enhanced root function (*e.g.* greater density or length of root hairs; stimulation

of the plant uptake pathways, production of exudates by plant or fungus) or changes in nutrient

partitioning. For other elements, the content in M plants could be explained by the larger root

system alone, although there is the possibility of a "hidden" mycorrhizal contribution (*i.e.* fungal

nutrient delivery balanced by an equivalent reduction in direct root uptake. Smith et al., 2003).

The growth increase in inoculated plants was likely driven by increased P uptake, given

that the experiment was conducted under low P availability. The route of P from soil to fungus,

and subsequent delivery to the plant host is well characterized (Chiu and Paszkowski, 2019). We

saw clear evidence that a unit of mycorrhizal root translates to a greater quantity of P obtained

than an equivalent unit of non-colonized root, as would be anticipated as a consequence of

hyphal foraging (Fig. 5). In comparison to P, the impact of AM colonization on the uptake of

Ca, Mg and S (the other macronutrients for which our observations were consistent with hyphal

foraging) is less well characterized (Gerlach et al., 2015; Ramírez-Flores et al., 2017). In the case

of S, it has been shown that AM fungi have the capacity to transfer S from soil to their plant

hosts (Gray and Gerdemann, 1973; Allen and Shachar-Hill, 2009), and that accumulation of

plant sulphate transporter transcripts increases in colonized roots (Casieri et al., 2012;

Giovannetti et al., 2014). Furthermore, the promoter of the *Lotus japonicus* sulphate transporter

gene *LjSULTR1;2* is active in arbusculated cells (Giovannetti et al., 2014), suggesting a function

in uptake from the peri-arbuscular space analogous to that of the PT4 high-affinity P transporter.

With regard to Mg, transcripts encoding Mg transporters are known to accumulate to higher

levels in wheat following mycorrhizal inoculation (Li et al., 2018). In common with P, the

elements S, Ca and Mg are often poorly available due to low-mobility and formation of

conjugates with other soil compounds (Kelly and Barber, 1991; Scherer, 2001; Lynch, 2019). As

such, hyphal foraging and delivery of these nutrients by mycorrhizal fungi would be predicted to

be of potential benefit to plants under field conditions. In the future, it will be informative to

combine evaluation of RSA in the field with assessment of AM fungal communities and

quantification of elements in reproductive stage plants and grain.

LITERATURE CITED

Allen JW, Shachar-Hill Y (2009) Sulfur transfer through an arbuscular mycorrhiza. Plant
Physiol **149**: 549–560

Aloni R, Aloni E, Langhans M, Ullrich CI (2006) Role of cytokinin and auxin in shaping root
architecture: regulating vascular differentiation, lateral root initiation, root apical dominance
and root gravitropism. Ann Bot **97**: 883–893

Bago B, Pfeffer PE, Abubaker J, Jun J, Allen JW, Brouillette J, Douds DD, Lammers PJ,
Shachar-Hill Y (2003) Carbon export from arbuscular mycorrhizal roots involves the

translocation of carbohydrate as well as lipid. Plant Physiol **131**: 1496–1507

Berta G, Fusconi A, Trotta A, Scannerini S (1990) Morphogenetic modifications induced by
the mycorrhizal fungus *Glomus* strain E3 in the root system of *Allium porrum* L. New
Phytologist **114**: 207–215

Berta G, Trotta A, Fusconi A, Hooker JE, Munro M, Atkinson D, Giovannetti M, Morini S,
Fortuna P, Tisserant B, Gianinazzi-Pearson V, Gianinazzi S (1995) Arbuscular
mycorrhizal induced changes to plant growth and root system morphology in *Prunus*
cerasifera. Tree Physiol **15**: 281–293

Bouranis DL, Buchner P, Chorianopoulou SN, Hopkins L, Protonotarios VE, Siyiannis VF,
Hawkesford MJ (2008) Responses to sulfur limitation in Maize. *In* NA Khan, S Singh, S
Umar, eds, Sulfur assimilation and abiotic stress in plants. Springer Berlin Heidelberg,
Berlin, Heidelberg, pp 1–19

Campbell BD, Grime JP, Mackey JML (1991) A trade-off between scale and precision in
resource foraging. Oecologia **87**: 532–538

Casieri L, Gallardo K, Wipf D (2012) Transcriptional response of *Medicago truncatula*
sulphate transporters to arbuscular mycorrhizal symbiosis with and without sulphur stress.
Planta **235**: 1431–1447

Chiu CH, Paszkowski U (2019) Mechanisms and impact of symbiotic phosphate acquisition.
Cold Spring Harb Perspect Biol. doi: 10.1101/cshperspect.a034603

Farley RA, Fitter AH (1999) The responses of seven co-occurring woodland herbaceous
perennials to localized nutrient-rich patches. J Ecol **87**: 849–859

Fox J & Weisberg S (2011). An {R} companion to applied regression Second Edition.
Thousand Oaks CA

Fukaki H, Tasaka M (2009) Hormone interactions during lateral root formation. Plant Mol Biol
69: 437–449

Galkovskyi T, Mileyko Y, Bucksch A, Moore B, Symonova O, Price CA, Topp CN, Iyer-
Pascuzzi AS, Zurek PR, Fang, Harer J, Benfey PN, Weitz JS (2012) GiA Roots:
software for the high throughput analysis of plant root system architecture. BMC Plant Biol
12: 116

Gamuyao R, Chin JH, Pariasca-Tanaka J, Pesaresi P, Catausan S, Dalid C, QLAmet-
Loedin I, Tecson-Mendoza EM, Wissuwa M, Heuer S (2012) The protein kinase Pstol1
from traditional rice confers tolerance of phosphorus deficiency. Nature **488**: 535–539

Garay-Arroyo A, De La Paz Sánchez M, García-Ponce B, Azpeitia E, Alvarez-Buylla ER
(2012) Hormone symphony during root growth and development. Dev Dyn **241**: 1867–1885

Gerlach N, Schmitz J, Polatajko A, Schlüter U, Fahnenstich H, Witt S, Fernie AR, Uroic K,

Scholz U, Sonnewald U, Bucher M (2015) An integrated functional approach to dissect systemic responses in maize to arbuscular mycorrhizal symbiosis. Plant, Cell & Environment **38**: 1591–1612

Giovannetti M, Tolosano M, Volpe V, Kopriva S, Bonfante P (2014) Identification and functional characterization of a sulfate transporter induced by both sulfur starvation and mycorrhiza formation in *Lotus japonicus*. New Phytol **204**: 609–619

González-Guerrero M, Azcón-Aguilar C, Mooney M, Valderas A, MacDiarmid CW, Eide DJ, Ferrol N (2005) Characterization of a *Glomus intraradices* gene encoding a putative Zn transporter of the cation diffusion facilitator family. Fungal Genet Biol **42**: 130–140

Govindarajulu M, Pfeffer PE, Jin H, Abubaker J, Douds DD, Allen JW, Bücking H, Lammers PJ, Shachar-Hill Y (2005) Nitrogen transfer in the arbuscular mycorrhizal symbiosis. Nature **435**: 819–823

Gray LE, Gerdemann JW (1973) Uptake of sulphur-35 by vesicular-arbuscular mycorrhizae. Plant and Soil **39**: 687–689

Grossman JD, Rice KJ (2012) Evolution of root plasticity responses to variation in soil nutrient distribution and concentration. Evol Appl **5**: 850–857

Gruber BD, Giehl RFH, Friedel S, von Wirén N (2013) Plasticity of the *Arabidopsis* root system under nutrient deficiencies. Plant Physiol **163**: 161–179

Guan JC, Koch KE, Suzuki M, Wu S, Latshaw S, Petruff T, Goulet C, Klee HJ, McCarty DR (2012) Diverse roles of strigolactone signaling in maize architecture and the uncoupling of a branching-specific subnetwork. Plant Physiol **160**: 1303–1317

Gutjahr C, Banba M, Croset V, An K, Miyao A, An G, Hirochika H, Imaizumi-Anraku H, Paszkowski U (2008) Arbuscular mycorrhiza-specific signaling in rice transcends the common symbiosis signaling pathway. Plant Cell **20**: 2989–3005

Gutjahr C, Casieri L, Paszkowski U (2009) *Glomus intraradices* induces changes in root system architecture of rice independently of common symbiosis signaling. New Phytol **182**: 829–837

Harrell Jr F with contributions from Charles Dupont and many others (2019). Hmisc: Harrell Miscellaneous. R package version 4.2-0.

Hoagland DR, Broyer TC (1936) General nature of the process of salt accumulation by roots with description of experimental methods. Plant Physiol **11**: 471–507

Hochholdinger F, Feix G (1998) Early post-embryonic root formation is specifically affected in the maize mutant *lrt1*. Plant J **16**: 247–255

Hochholdinger F, Tuberosa R (2009) Genetic and genomic dissection of maize root development and architecture. Curr Opin Plant Biol **12**: 172–177

Ho MD, McCannon BC, Lynch JP (2004) Optimization modeling of plant root architecture for
water and phosphorus acquisition. J Theor Biol **226**: 331–340

Jung JKH, McCouch S (2013) Getting to the roots of it: Genetic and hormonal control of root
architecture. Front Plant Sci **4**: 186

Kelly JM, Barber SA (1991) Magnesium uptake kinetics in loblolly pine seedlings. Plant and
Soil **134**: 227–232

Koltai H (2011) Strigolactones are regulators of root development. New Phytol **190**: 545–549

Kutz A, Muller A, Hennig P, Kaiser WM, Piotrowski M, Weiler EW (2002) A role for
nitrilase 3 in the regulation of root morphology in sulphur-starving *Arabidopsis thaliana*.
Plant J **30**: 95–106

Li M, Wang R, Tian H, Gao Y (2018) Transcriptome responses in wheat roots to colonization
by the arbuscular mycorrhizal fungus *Rhizophagus irregularis*. Mycorrhiza **28**: 747–759

Liu A, Hamel C, Hamilton RI, Ma BL, Smith DL (2000) Acquisition of Cu, Zn, Mn and Fe by
mycorrhizal maize (*Zea mays* L.) grown in soil at different P and micronutrient levels.
Mycorrhiza **9**: 331–336

López-Bucio J, Cruz-Ramírez A, Herrera-Estrella L (2003) The role of nutrient availability
in regulating root architecture. Current Opinion in Plant Biology **6**: 280–287

López-Bucio J, Hernández-Abreu E, Sánchez-Calderón L, Nieto-Jacobo MF, Simpson J,
Herrera-Estrella L (2002) Phosphate availability alters architecture and causes changes in
hormone sensitivity in the Arabidopsis root system. Plant Physiol **129**: 244–256

López-Pedrosa A, González-Guerrero M (2006) *GintAMT1* encodes a functional high-affinity
ammonium transporter that is expressed in the extraradical mycelium of *Glomus*
intraradices. Fungal Genet. Biol.

Lynch JP (1995) Root architecture and plant productivity. Plant Physiol **109**: 7–13

Lynch JP (2019) Root phenotypes for improved nutrient capture: an underexploited opportunity
for global agriculture. New Phytol. doi: 10.1111/nph.15738

Lynch JP, Brown KM (2008) Root strategies for phosphorus acquisition. Plant Ecophysiology
83–116

Malamy JE, Ryan KS (2001) Environmental regulation of lateral root initiation in *Arabidopsis*.
Plant Physiol **127**: 899–909

de Mendiburu F (2019). agricolae: Statistical procedures for agricultural research. R package
version 1.3-0.

McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA (1990) A new method which

gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal
fungi. New Phytol **115**: 495–501

Mollier A, Pellerin S (1999) Maize root system growth and development as influenced by
phosphorus deficiency. J Exp Bot **50**: 487–497

Mora-Macías J, Ojeda-Rivera JO, Gutiérrez-Alanís D, Yong-Villalobos L, Oropeza-Aburto
A, Raya-González J, Jiménez-Domínguez G, Chávez-Calvillo G, Rellán-Álvarez R,
Herrera-Estrella L (2017) Malate-dependent Fe accumulation is a critical checkpoint in
the root developmental response to low phosphate. Proc Natl Acad Sci U S A **114**: E3563–
E3572

Nouri E, Breuillin-Sessoms F, Feller U, Reinhardt D (2014) Phosphorus and nitrogen regulate
arbuscular mycorrhizal symbiosis in *Petunia hybrida*. PLoS One **9**: e90841

Oláh B, Brière C, Bécard G, Dénarié J, Gough C (2005) Nod factors and a diffusible factor
from arbuscular mycorrhizal fungi stimulate lateral root formation in *Medicago truncatula*
via the DMI1/DMI2 signalling pathway. Plant J **44**: 195–207

Osmont KS, Sibout R, Hardtke CS (2007) Hidden branches: developments in root system
architecture. Annu Rev Plant Biol **58**: 93–113

Parniske M (2008) Arbuscular mycorrhiza: the mother of plant root endosymbioses. Nat Rev
Microbiol **6**: 763–775

Paszkowski U, Boller T (2002) The growth defect of *lrt1*, a maize mutant lacking lateral roots,
can be complemented by symbiotic fungi or high phosphate nutrition. Planta **214**: 584–590

Pearson JN, Jakobsen I (1993) Symbiotic exchange of carbon and phosphorus between
cucumber and three arbuscular mycorrhizal fungi. New Phytologist **124**: 481–488

Postma JA, Dathe A, Lynch JP (2014) The optimal lateral root branching density for maize
depends on nitrogen and phosphorus availability. Plant Physiol **166**: 590–602

Ramírez-Flores MR, Rellán-Álvarez R, Wozniak B, Gebreselassie M-N, Jakobsen I,
Olalde-Portugal V, Baxter I, Paszkowski U, Sawers RJH (2017) Co-ordinated changes
in the accumulation of metal ions in maize (*Zea mays* ssp. *mays* L.) in response to
inoculation with the arbuscular mycorrhizal fungus *Funneliformis mosseae*. Plant Cell
Physiol **58**: 1689–1699

Rangarajan H, Postma JA, Lynch JP (2018) Co-optimization of axial root phenotypes for
nitrogen and phosphorus acquisition in common bean. Ann Bot **122**: 485–499

Rubio G, Liao H, Yan X, Lynch JP (2003) Topsoil foraging and its role in plant
competitiveness for phosphorus in common bean. Crop Science **43**: 598

Salt DE, Baxter I, Lahner B (2008) Ionomics and the study of the plant ionome. Annu Rev
Plant Biol **59**: 709–733

Sawers RJH, Svane SF, Quan C, Grønlund M, Wozniak B, Gebreselassie M-N, González-Muñoz E, Chávez Montes RA, Baxter I, Goudet J, Jakobsen I, Paszkowski U (2017) Phosphorus acquisition efficiency in arbuscular mycorrhizal maize is correlated with the abundance of root-external hyphae and the accumulation of transcripts encoding PHT1 phosphate transporters. New Phytol **214**: 632–643

Scherer HW (2001) Sulphur in crop production. European Journal of Agronomy **14**: 81–111

Schlicht M, Ludwig-Müller J, Burbach C, Volkmann D, Baluska F (2013) Indole-3-butyric acid induces lateral root formation via peroxisome-derived indole-3-acetic acid and nitric oxide. New Phytol **200**: 473–482

Schüßler A, Schwarzott D, Walker C (2001) A new fungal phylum, the *Glomeromycota*: phylogeny and evolution. Mycol Res **105**: 1413–1421

Shahzad Z, Amtmann A (2017) Food for thought: how nutrients regulate root system architecture. Curr Opin Plant Biol **39**: 80–87

Smith SE, Read DJ (2010) Mycorrhizal Symbiosis. Academic Press

Smith SE, Smith FA, Jakobsen I (2003) Mycorrhizal fungi can dominate phosphate supply to plants irrespective of growth responses. Plant Physiol **133**: 16–20

Tisserant B, Gianinazzi S, Gianinazzi-Pearson V (1996) Relationships between lateral root order, arbuscular mycorrhiza development, and the physiological state of the symbiotic fungus in *Platanus acerifolia*. Canadian Journal of Botany **74**: 1947–1955

Venables, WN & Ripley, BD (2002) Modern Applied Statistics with S. Fourth Edition. Springer, New York

Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, Lumley T, Maechler M, Magnusson A, Moeller S, Schwartz M, Venables W (2019). gplots: Various R programming tools for plotting data. R package version 3.0.1.1.

Williamson LC, Ribrioux SP, Fitter AH, Leyser HM (2001) Phosphate availability regulates root system architecture in *Arabidopsis*. Plant Physiol **126**: 875–882

Wissuwa M, Gatdula K, Ismail A (2005) Candidate gene characterization at the *Pup1* locus: a major QTL increasing tolerance of phosphorus deficiency.

Zhang H, Jennings A, Barlow PW, Forde BG (1999) Dual pathways for regulation of root branching by nitrate. Proc Natl Acad Sci U S A **96**: 6529–6534

Zhang Y, von Behrens I, Zimmermann R, Ludwig Y, Hey S, Hochholdinger F (2015) *LATERAL ROOT PRIMORDIA 1* of maize acts as a transcriptional activator in auxin signalling downstream of the *Aux/IAA* gene *rootless* with undetectable meristem 1. J Exp Bot **66**: 3855–3863

513   **Ziegler G, Terauchi A, Becker A, Armstrong P, Hudson K, Baxter I** (2013) Ionomic
514   screening of field-grown soybean identifies mutants with altered seed elemental
515   composition. *The Plant Genome* 6: 0

516
517
518
519

520 **Table 1. Traits responsive to inoculation with *Rhizophagus irregularis***

Trait[1]	Description[2]	NC		M		MR[3] (%)
		mean	SE	mean	SE	
QLA45	Sqrt leaf surface area, day 45 (cm)	11.6	0.449	14.8	0.497	28
QLA50	Sqrt leaf surface area, day 50 (cm)	12.2	0.426	17.4	0.433	43
QLA55	Sqrt leaf surface area, day 55 (cm)	13.6	0.471	19.6	0.676	44
SFW	Shoot fresh weight (g)	14	1.3	35.8	2.16	156
RFW	Root fresh weight (g)	17.6	1.64	44.5	2.84	153
SDW	Shoot dry weight (g)	1.78	0.184	4.29	0.284	142
RDW	Root dry weight (g)	1.39	0.122	3.28	0.185	136
RV	Root volume (ml)	18.6	1.75	48.4	3.17	160
MaxNR	Maximum number of roots	36.3	1.78	53.9	2.03	49
MNR	Median number of roots	23.4	1.35	39.8	1.57	70
NB	Network bushiness	1.59	0.054	1.37	0.028	-14
NetA	Network area (cm^2)	95.1	6.89	181	9.72	90
NL	Network length (cm)	1870	150	3610	206	93
NLD	Network Length Distribution	0.937	0.058	0.661	0.033	-29
NP	Network perimeter (cm)	3640	281	6870	390	89

NS	Network solidity	0.129	0.004	0.185	0.006	43
NSA	Network surface area (cm^2)	361	27.1	698	38.4	93
NV	Network volume (cm^3)	6.84	0.536	13.3	0.808	95
Ca	Leaf Ca concentration (ppm)	9470	435	12900	589	36
K	Leaf K concentration (ppm)	24000	612	18300	1130	-24
Mn	Leaf Mn concentration (ppm)	115	5.25	149	6.26	29
P	Leaf P concentration (ppm)	612	19.3	1160	28.9	89
Se	Leaf Se concentration (ppm)	0.663	0.017	0.498	0.017	-25
Sr	Leaf Sr concentration (ppm)	39	1.86	51.3	2.85	32

522   [1] Traits listed showed a significant (adjP < 0.05) main effect of fungal inoculation.

523   [2] GiA Roots traits as described in full by Galkovskyi et al. 2012

524   [3] Mycorrhiza response calculated as M-NC/NC x 100

 FIGURE LEGENDS

Figure 1. Experimental set-up and plant growth response. The maize inbred lines B73 and
W22 were grown with (M) or without (NC) inoculation with the AM fungus *Rhizophagus*
irregularis. A) Overall view of the growth system, the block in the foreground are inoculated
plants. B) Square root of leaf surface area (QLA) for M and NC individuals, quantified every 5
days from 5 days after emergence until harvest at day 55. Boxes show 1st quartile, median and
3rd quartile. Whiskers extend to the most extreme points within 1.5x box length; outlying values
beyond this range are shown as filled circles. Days at which M and NC groups were significantly
different for a given inbred (Tukey HSD; $p < 0.05$) are indicated with an asterisk. C) Root crown
on an M plant, illustrating the profusion of lateral roots and characteristic yellow pigmentation.
D) Trypan-blue stained root section of an M plant, showing root-internal hyphae (hyp), vesicles
(ves), and arbuscules (arb). Scale bar, 15 μm. E) Colonization (% root length colonized) for B73
and W22, determined with respect to fungal hyphae (H), arbuscules (A) and vesicles (V). Points
indicate individual plants. Boxes and whiskers as in B).

Figure 2. Principal component analysis describes variation in root system architecture. A)
Stacked (overlay) images of the root systems of all plants in the NC/M B73/W22 treatment
groups. Scale bar, 30 cm. B) The contribution of root system traits to principal components (PCs)
1 to 4. The variance explained by each PC is given in parentheses. See Table S1 for trait
abbreviations. C) Root system images arranged by loading on each of PCs 1 to 4. Coloured
points at the base of the image indicate genotype and fungal treatment as described in the key.
Traits with a major positive or negative contribution to each PC are listed on the right or left of
the plot, respectively. A single word description of the extremes of each PC is given at the side
of each plot.

Figure 3. Inoculation with AM fungi is correlated with an increase in size and branching of
the root system. Root system images projected by Linear Discriminants (LDs) 1 and 2. Colored
points indicate genotype and fungal treatment as described in the key.

Figure 4. Root system architecture is correlated with total element content in the leaf. A)
Heatmap representation of pairwise correlations of root system principal components (PC) with
their contributing GiARoot traits (Abbreviated as described in Materials and Methods. SFW,
shoot fresh weight and SDW, shoot dry weight also shown). B) Heatmap representation of
pairwise correlations of root system principal components (PC) and colonization measures (%
root length of hyphae (Hyp), vesicles (Ves) and arbuscules (Arb)) with total element content. In
both A) and B) significant correlations are indicated by asterisks (*, $p < 0.05$; **, $p < 0.01$; ***,
$p < 0.001$).

Figure 5. AM colonization modifies the relationship between root system size and element
content. Scatter plot representation of the relationship between element content (shown in mg or

μg) and root system size (estimated as the GiaRoots trait NSA) for twenty named elements.
Colored points indicate genotype and fungal treatment of individual plants as described in the
key. Black and red lines indicate the fit for the additive model *Total ~ NSA + Fungus* for levels
NC and M of *Fungus*, respectively. The p-value for an effect of *Fungus* (calculated using Type I
SS; adjusted for multiple tests) is given in parentheses following the element name in the plot
title.
572

Figure S1. Impact of AM colonization of growth, root architecture and nutrient levels. The
maize inbred lines B73 and W22 were grown with (M) or without (NC) inoculation with the AM
fungus *Rhizophagus irregularis*. Traits are described in main text and Table S1, S2, S3. Boxes
show 1st quartile, median and 3rd quartile. Whiskers extend to the most extreme points within
1.5x box length; outlying values beyond this range are not shown. Line segments show the
reaction norm for each genotype, connecting median values for NC and M.
579

ACKNOWLEDGEMENTS

We thank Greg Ziegler and Ivan Baxter (Danforth Center, MO) for ionomic analysis and

Benjamin Barrales Gámez for assistance in the greenhouse experiment. We thank the reviewers

for constructive feedback on the manuscript. M. Rosario Ramírez-Flores was supported by the

Mexican National Council of Science and Technology (CONACYT) through its PhD scholarship

program. This work was supported by the Mexican National Commission for the Study and Use

of Biodiversity (CONABIO) project *Impact of native arbuscular mycorrhizal fungi on maize*

performance.

AUTHOR CONTRIBUTIONS

RRF, RJHS and VOP designed the study. MRRF conducted the experiments. MRRF, EBB and

RRA performed image analysis. All authors contributed to data analysis and writing of the

manuscript.

M. Rosario Ramírez-Flores[1], Elohim Bello-Bello[2], Rubén Rellán-Álvarez[2, 3], Ruairidh J. H.

Sawers[2,4*] and Víctor Olalde-Portugal[1*]

PREPRINT

This article is available as a preprint at https://www.biorxiv.org/content/10.1101/695411v1

CONFLICT OF INTEREST STATEMENT

Rubén Rellán-Álvarez is a review editor for Plant Direct.

LD1 (85%)

large ← → small

sparse

LD2 (11%)

branched

B73 NC ■   B73 M ■   W22 NC ■   W22 M ■