ORP5 REGULATES TRANSPORT OF LIPIDS AND CALCIUM TO

MITOCHONDRIA AT ENDOPLASMIC RETICULUM-MITOCHONDRIA MEMBRANE

CONTACT SITES

Leila Rochin[1], Cécile Sauvanet[1]*, Eeva Jääskeläinen[2]*, Audrey Houcine[3]*,

Annukka Kivelä[2], Xingjie MA[5], Eyra Marien[4], Jonas Dehairs[4], Julie Neveu[1],

Romain Le Bars[1], Johannes Swinnen[4], David Bernard[5], David Tareste[6], Vesa M.

Olkkonen[2,7], and Francesca Giordano[1#]

[1] Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud,

Université Paris-Saclay, Gif-sur-Yvette cedex 91198, France.

[2] Minerva Foundation Institute for Medical Research, Biomedicum 2U, FI-00290

Helsinki, Finland

[3] Institut Jacques Monod, CNRS, UMR7592, Université Paris Diderot, Sorbonne

Paris Cité, F-75013 Paris, France

[4] Laboratory of lipid metabolism and cancer, Department of Oncology, KU Leuven,

Leuven B-3000, Belgium.

[5] Centre de Recherche en Cancérologie de Lyon, UMR INSERM U1052/CNRS 5286,

Centre Léon Bérard, 69373 LYON Cedex 08, France

[6] Institute of Psychiatry and Neuroscience of Paris (IPNP), UMR-S1266 INSERM-

Paris Descartes University, Paris, France

[7] Department of Anatomy, Faculty of Medicine, FI-00014 University of Helsinki,

Finland

* These authors contributed equally to this work

[#] Correspondence to: francesca.giordano@i2bc.paris-saclay.fr

ABSTRACT

Mitochondria are dynamic organelles essential for cell survival whose structural and

functional integrity rely on selective and regulated transport of lipids from/to the

endoplasmic reticulum (ER) and across the two mitochondrial membranes. As they are

not connected by vesicle transport, the exchange of lipids between ER and mitochondria

occurs at sites of close organelle apposition called membrane contact sites. However,

the mechanisms and proteins involved in these processes are only beginning to emerge.

Here, we show that ORP5/8 mediate non-vesicular transport of Phosphatidylserine (PS)

from the ER to mitochondria in mammalian cells. We also show that ER-mitochondria

contacts where ORP5/8 reside are physically and functionally linked to the MIB/MICOS

complexes that bridge the mitochondria membranes, cooperating with them to facilitate

PS transfer from the ER to the mitochondria. Finally, we show that ORP5 but not ORP8,

additionally regulates import of calcium to mitochondria and consequently cell

senescence.

KEYWORDS

Membrane contact sites, mitochondria, ORP, lipid transfer, calcium

ABBREVIATIONS

APOO: Apolipoprotein O

APOOL: Apolipoprotein O-like

CHCHD3/6: Coiled-Coil-Helix-Coiled-Coil-Helix Domain Containing 3/6

Calcium: Ca^{2+}

CCCP: Carbonyl cyanide 3-chlorophenylhydrazone

CJ: cristae junctions

CV-ATP5A: complex V alpha subunit

EM: electron microscopy

ER: endoplasmic reticulum

FFAT: two phenylalanines (FF) in an Acidic Tract

GRP75: glucose-regulated protein 75

HRP: horseradish peroxidase

IF: immunofluorescence

IMM: inner mitochondrial membrane

IMS: intermembrane space

IP3R: Inositol trisphosphate receptor

LAR-GECO: low-affinity red fluorescent genetically encoded Ca2+ indicators for

optical imaging

LTPs: lipid transfer proteins

MAM: mitochondria-associated ER membrane

MCU: mitochondrial calcium uniporter

MIB: Mitochondrial intermembrane space bridging complex

MICOS: Mitochondrial Contact Sites and Cristae Junction Organizing System

MINOS1: mitochondrial inner membrane organizing system 1

MS: mass spectrometry

mt-Pericam: mitochondrial permuted YFP and a CaM (calmodulin)

OMM: outer mitochondrial membrane

ORD: OSBP-related lipid-binding/transfer domain

ORP: Oxysterol binding protein (OSBP)-related proteins

Osh: oxysterol-binding homology

PC: Phosphatidylcholine

PDI: Protein disulfide isomerase

PE: phosphatidylethanolamine

PH: pleckstrin homology

PI4P: Phosphatidylinositol-4-phosphate

PIP2: Phosphatidylinositol 4,5-bisphosphate

POPC: Palmitoyloleoylphosphatidylcholine

PS: phosphatidylserine

PTPIP51: Protein tyrosine phosphatase interacting protein 51

QIL1: MICOS complex subunit MIC13

R-GECO: Red fluorescent, genetically-encoded Ca2+ indicator for optical imaging

SA β-gal: Senescence-associated β –galactosidase

SAM: Sorting and Assembly Machinery

SAM50: Sorting Assembly Machinery 50

TLC: thin layer chromatography

TMRE: Tetramethylrhodamine ethyl ester

VAPB: Vesicle-associated membrane protein-associated protein B

VDAC: Voltage Dependent Anion Channel

WB: western blotting

INTRODUCTION

Vesicular trafficking is the major pathway for transport of proteins and metabolites,

such as lipids, between membranes. However, an alternative route, which is vesicle-

independent, occurs at regions of close inter-organelle membrane proximity (within less

than 30 nm) also called membrane contact sites (Scorrano, De Matteis et al., 2019).

This route is particularly important to preserve membrane composition, integrity and

identity of the intracellular organelles such as the mitochondria that are largely excluded

from the classical vesicle-mediated trafficking pathways. Like other organelles,

mitochondria can be closely associated with the endoplasmic reticulum (ER), the major

site of lipid synthesis and the major intracellular calcium (Ca^{2+}) store. ER membrane

subdomains closely apposed to the mitochondria are called mitochondria-associated ER

membranes (MAMs) and they facilitate the exchange of Ca^{2+} and lipids between the two

organelles (Herrera-Cruz & Simmen, 2017, Tatsuta, Scharwey et al., 2014, Vance,

2014)

Mitochondria are involved in a plethora of cellular processes including energy

production, lipid metabolism, Ca^{2+} homeostasis and apoptosis. To fulfill their numerous

functions, mitochondria need to maintain a defined membrane composition by receiving

essential lipids and lipid precursors from the ER through membrane contact sites

(Giordano, 2018, Vance & Tasseva, 2013).

Increasing lines of evidence suggest that lipid transfer proteins (LTPs) play a major

role in regulating the lipid composition of membranous organelles by facilitating non-

vesicular lipid transport at membrane contact sites. In recent years, several tethering

complexes with lipid transfer activity have been identified at membrane contact sites

between the ER and other intracellular organelles including the plasma membrane (PM)

in yeast as well as in mammalian cells. However, the LTPs of ER-mitochondria

membrane contact sites remain still mysterious. The best-studied lipid transfer/tethering

complex at ER-mitochondria contact sites is the yeast ER-mitochondria encounter

structure (ERMES) (Kornmann et al 2009, Lang et al 2015) that bridges the ER and

the mitochondrial membranes and also facilitates the exchange of phospholipids (in

particular phosphatidylcoline, PC) between them. Despite its important functions in

yeast, the ERMES complex is not conserved in metazoans. In metazoans, very little

is known on how lipids are exchanged at ER-mitochondria membrane contact sites

and about the proteins involved in this process. Some tethers at mammalian ER-

mitochondria contact sites have started to emerge in the past years (Gatta and

Levine 2017), including the Mitochondrial Ca^{2+} Uniporter (MCU)–Voltage Dependent

Anion Channel (VDAC)–Glucose Regulated Protein 75 (GRP75)–Inositol

Triphosphate Receptor (IP3R) machinery involved in Ca^{2+} import from the ER to

mitochondria (Rizzuto et al. 2009) and the recently identified Protein tyrosine

phosphatase interacting protein 51 (PTPIP51)–Vesicle-associated membrane

protein-associated protein B (VAPB) complex (Stoica et al 2014) alteration of which

induces defects in ER-mitochondria tethering and Ca^{2+} exchanges. However, none of

these proteins has been directly involved in non-vesicular lipid transport between ER

and mitochondria membranes.

The Oxysterol binding protein (OSBP)-related proteins constitute a large family of

LTPs conserved from yeast (Osh) to humans (ORP) and localized to different

subcellular sites, shown in several cases to be membrane contact sites. A common

feature of all ORPs is the presence of an OSBP-related lipid-binding/transfer (ORD)

domain. Most ORP proteins contain an FFAT domain that binds ER-localized VAP

proteins and a pleckstrin homology (PH) domain that interacts with lipids or proteins

in distinct non-ER organelle membranes. Two members of this family, ORP5 and

ORP8, do not contain a FFAT domain but are directly anchored to the ER through

their C-terminal domain. We have recently shown that ORP5 and ORP8 localize to

ER-mitochondria contact sites where they play a key role in maintaining

mitochondrial morphology and respiratory function (Galmes et al. 2016). ORP5 and

ORP8 have been previously shown to transfer phosphatidylserine (PS) from the

cortical ER to the PM, in counter-exchange with the phosphoinositides

Phosphatidylinositol-4-phosphate (PI4P) and Phosphatidylinositol 4,5-bisphosphate

(PIP_2) (Chung et al. 2015; Ghai et al, 2017). Interestingly, transport of PS occurs also

at ER-mitochondria contact sites. Newly synthesized PS, by the ER-localized PS-

Synthase 1 (PSS1), is shuttled from the ER to the outer mitochondrial membrane

(OMM) and from OMM to inner mitochondrial membrane (IMM) where it is rapidly

converted to phosphatidylethanolamine (PE) (Vance, 1990, Vance & Tasseva, 2013). At

the IMM, PE plays crucial roles in maintaining mitochondrial tubular morphology and

therefore in mitochondrial respiratory functions (Joshi, Thompson et al., 2012,

Steenbergen, Nanowski et al., 2005). Regardless of extensive studies on PS transport

between ER and mitochondria since its first discovery more than 20 years ago (Vance,

1990), the mechanisms underlying this transport and the proteins involved are still

elusive.

Here we show that ORP5 and ORP8 are the LTPs that transfer PS from the ER

to the mitochondria at ER-mitochondria contact sites in mammalian cells. We also

show that they are physically and functionally connected with Sorting Assembly

Machinery 50 (SAM50) and Mitofilin, two components of the Mitochondrial

Intermembrane space Bridging (MIB) and Mitochondrial Contact sites and Cristae

junction Organizing System (MICOS) complexes, that anchors the IMM to the OMM.

The MICOS complex (composed of seven proteins: Mitofilin, CHCHD3, CHCHD6,

APOO, QIL1, APOOL and MINOS1) is preferentially located at Cristae Junctions (CJ),

tubular structures that connect the IMM to the cristae, and it is necessary for CJ

formation, cristae morphology and mitochondria function (Harner, Korner et al., 2011,

Huynen, Muhlmeister et al., 2016, Ott, Dorsch et al., 2015, Wollweber, von der

Malsburg et al., 2017). The integral IMM protein Mitofilin is the central component of

the MICOS complex and carries a large domain exposed to the mitochondria

intermembrane space (IMS) that interacts with the OMM Sorting and Assembly

Machinery (SAM) to form the MIB complex (Friedman, Mourier et al., 2015, Guarani,

McNeill et al., 2015). The SAM complex is constituted of SAM50 (a pore-forming β-

barrel protein), Metaxin1 and 2, and is involved in the membrane insertion and

assembly of mitochondrial β-barrel proteins (Hohr, Lindau et al., 2018, Kozjak,

Wiedemann et al., 2003, Kozjak-Pavlovic, Ross et al., 2007) (see cartoon in Figure

8).

Here we show that ORP5/8 interact and cooperate with SAM50 and Mitofilin to

facilitate the movement of PS from the ER to the mitochondria across ER-

mitochondria contact sites and the IMS and consequently the synthesis of

mitochondrial PE. Finally, we uncover an additional role for ORP5, but not ORP8, in

regulating Ca^{2+} import to mitochondria, through regulation of SAM50-VDAC protein

levels. Our findings reveal a novel link between lipid and Ca^{2+} transport at ER-

mitochondria contact sites and provide a mechanistic insight of the specific role of

ORP5 in modulating cellular senescence.

RESULTS

ORP5 and ORP8 transport PS from the ER to the mitochondria at ER-mitochondria contact sites

We have recently reported that ORP5 and ORP8, shown to transfer lipids at ER-PM contacts, localize also to ER-mitochondria contact sites (Galmes, Houcine et al., 2016). However, their role in lipid transport at ER-mitochondria contacts still remains to be established. Taking into account their role in counter-exchanging PS with the PM phosphoinositides PI4P and PIP_2 at ER-PM contacts (Chung et al. 2015; Ghai et al, 2017), we hypothesized that ORP5 and ORP8 could also mediate PS transport at ER-mitochondria contact sites.

The ORD domain of ORP8 has been shown to transfer PS (in counter-transport with PI4P or PIP_2) between liposome membranes *in vitro* (Chung et al. 2015; Ghai et al, 2017). Since PI4P and PIP_2 are not present on mitochondrial membranes it is possible that at ER-mitochondria contact sites ORP5/8 might transport back to the ER other phospholipids, such as PE. However, no study so far has ever addressed the ability of ORP5 and ORP8 ORD domains (ORD5, ORD8) to transfer other phospholipids between liposomes *in vitro.* First, to compare and study the role of ORD5 and ORD8 in the transfer of phospholipids *in vitro*, we purified the recombinant ORD5 (aa 265-703) and ORD8 (aa 328-767) in bacteria (Escherichia coli) (Fig. S1a) and analyzed their lipid transfer ability by measuring the transport of fluorescent phospholipids (TopFluor-PS, -PC or -PE) from donor to acceptor liposomes *in vitro*. Donor liposomes containing fluorescent phospholipids and biotinylated lipids (liposomes composed of 1 mol% TopFluor-PS, -PC or -PE and 2 mol% of biotinylated-PE, and 97 mol% POPC) were first immobilized on streptavidin beads and then mixed with acceptor liposomes (composed of 100 mol % POPC) in

presence or absence of ORP5/8 ORD domains (Fig. 1a). After 1 hour at 37°C,

acceptor liposomes were recovered from the supernatant and their fluorescence was

measured (Fig. 1a). Our results show that both ORD5 and ORD8 transfer PS, but not

PC and PE, from donor to acceptor liposomes (Fig. 1b). They also reveal that ORD5

and ORD8 share equivalent abilities to transfer PS *in vitro*.

Next, we addressed whether ORP5 and ORP8 could mediate non-vesicular

transfer of PS at ER-mitochondria contact sites *in situ*. As the ER-derived PS is the

major precursor for mitochondrial PE, if ORP5 and ORP8 mediate non-vesicular

transport of PS from the ER to the mitochondria, then their absence should lead to a

reduction of mitochondrial PE. To test this, we used Percoll gradient-based

subcellular fractionation (Galmes et al. 2016) to isolate pure mitochondria from HeLa

cells where ORP5 or ORP8 were transiently silenced by RNAi. We chose to use a

transient knockdown as it overcomes the limits and/or compensatory effects on lipid

transport/biosynthetic pathways that other stable approaches could induce. The

purity of mitochondria and of the other subcellular fractions was verified in control,

ORP5 and ORP8 knockdown conditions by western blotting (WB) (Fig. S1b). As

controls for the purity of subcellular fractions, the samples were probed for

cytochrome c as mitochondrial marker and IP3R-3 as a MAM-enriched marker. All

markers were highly enriched in their respective fractions and were absent in the

others. In accord with our previous study (Galmes et al., 2016), ORP5 and ORP8

were enriched in the MAM fraction and absent in the mitochondria fraction of control

cells. On the contrary, they were strongly suppressed in ORP5 and ORP8

knockdown cell lysates and in the respective MAM fractions (Fig. S1b). The PE

content of purified mitochondria from ORP5 and ORP8 knockdown was then

analyzed by mass spectrometry (MS)-based lipidomics, revealing a reduction of PE

levels in mitochondria isolated from ORP5 and ORP8 knockdown cells of 34% and

20%, respectively, as compared to control cells (Fig. 1c). Interestingly, PE levels of

the total cells were unchanged, indicating a specific effect of ORP5 or ORP8

depletion on mitochondrial PE.

In subsequent experiments, we measured the levels of mitochondrial PE newly

synthesized from the ER-derived PS by using a radiometric PS-to-PE conversion

assay *in situ* (Shiao, Lupo et al., 1995) in silenced or control HeLa cells (Fig. S1c-1d).

This assay allows the monitoring of PS transfer from the ER to mitochondria by

measuring the levels of radioactive PS and PS-derived PE by thin layer

chromatography (TLC) after 18h of incorporation of radioactive L-[^{3}H(G)]-serine into

the cells. A significant decrease in the levels of newly synthetized PE was found in

ORP5 knockdown and in ORP5+ORP8 double-knockdown cells (Fig. 1d). The

decrease was stronger in ORP5+ORP8 double-knockdown cells, indicating a

cooperative effect of ORP5 and ORP8 in this process. A slight decrease was also

found in ORP8 knockdown cells, although not statistically significant, suggesting a

major role of ORP5 compared to ORP8 in PS transfer at ER-mitochondria contact

sites *in situ*.

The decrease in mitochondria PE is in accord with its essential role in

maintaining the morphology of mitochondrial cristae and with our previous electron

microscopy (EM) observations that knockdown of ORP5 or ORP8 alters the

architecture of the cristae, while ER-mitochondria contact sites with normal

morphology were still visible (Galmes et al., 2016). However, our previous study did

not rule out that these contact sites could be affected in their number/size. To test

whether the effects on PS transport at ER-mitochondria contacts were specific of

ORP5 and ORP8 loss of function or simply due to a decrease of ER-mitochondria

contacts induced by their knockdown, we quantified the abundance of ER-mitochondria contact sites by EM in control and ORP5, ORP8, ORP5+ORP8 knockdown cells. To facilitate the visualization of the ER we transfected the cells with a HRP-KDEL construct (carrying a horseradish peroxidase (HRP) tagged with an ER retention motif) that stains the ER with a dark signal. Our quantifications revealed that ORP5, ORP8 or ORP5+ORP8 knockdowns did not affect the extent of ER-mitochondria contact sites (Fig. 1e-1f). These results indicate that ORP5 and ORP8 silencing affect non-vesicular PS transport at ER-mitochondria contacts but not their extent and integrity, confirming that at these sites ORP5 and ORP8 act exclusively as LTPs and not as tether (Fig. 1h). Additionally, 52% of mitochondria in ORP5+ORP8 double-knockdown cells display aberrant cristae morphology versus 9% in control cells (Fig. 1e-1g). These defects in cristae morphology were stronger than what we had previously shown in the case of ORP5 and ORP8 individual knockdowns (Galmes et al., 2016), and most likely reflect the stronger effect of ORP5+ORP8 double-knockdown on PS transport at ER-mitochondria contact sites.

ER-mitochondria contact sites marked by ORP5/8, are physically and functionally connected with the mitochondrial intermembrane space bridging (MIB) complex

To investigate whether ORP5/8 localize to specific ER-mitochondria contact site subdomains we performed a morphological analysis of ORP5 localization by immuno-EM (IEM) on ultrathin cryosections from HeLa cells transfected with HA-ORP5. The advantage of analyzing ORP5 localization is its preferential localization to contact sites as compared to ORP8 (Galmes et al. 2016), which remains also largely present within the reticular ER. Interestingly, the majority of ORP5 gold particles was

found to localize to ER elements in a very close proximity (86% within 0-100 nm

distances, 50% of which within 50 nm) to the CJ (arrow, Fig. 2a-b), tubular structures

that connect the IMM to the cristae. In yeast, CJ were shown to be closely associated

to OMM-IMM contact sites tethered by the MICOS complex (Harner et al., 2011). IEM

analysis using Mitofilin-EGFP, an EGFP-tagged construct of the human orthologue of

the central component of the MICOS complex, confirmed that human Mitofilin,

similarly to its yeast orthologue, preferentially localizes to the IMM in close proximity

of CJ and in the cristae that arise from them (arrow, Fig. 2a).

To identify specific binding partners of ORP5 at ER-mitochondria contact sites

we carried out a MS-analysis on GFP-pull downs from cells expressing EGFP-ORP5,

EGFP-ORP5ΔPH (an ORP5 mutant lacking the PM-targeting PH domain that is

localized at ER-mitochondria but not at ER-PM contact sites), or EGFP alone as a

control (Fig. S2a). As expected, the highest hit detected in both EGFP-ORP5 and

EGFP-ORP5ΔPH pull-downs was its binding partner ORP8, with which ORP5 forms

a complex at contact sites (Chung, Torta et al., 2015, Galmes et al., 2016). Other

major hits are listed in Fig. S2a and comprise several outer mitochondrial membrane

proteins, including the mitochondrial outer membrane protein PTPIP51, a previously

reported interacting partner of ORP5 (Galmes et al., 2016).

Interestingly, among these proteins the Sorting Assembly Machinery Subunit 50

(SAM50), a central component of the SAM protein complex involved in the import

and assembly of mitochondria β-barrel proteins in the OMM (Hohr et al., 2018), had

one of the highest scores. SAM50 is part of the Mitochondrial intermembrane space

bridging (MIB) complex, composed of metaxin-1, metaxin-2 and the MICOS complex

that anchors the OMM to the IMM at CJ (Huynen et al., 2016, Ott et al., 2015).

SAM50 is also known to directly bind the central component of the MICOS complex

Mitofilin (Ott et al., 2015). Consistently, Mitofilin was also detected in the MS of

EGFP-ORP5 and EGFP-ORP5ΔPH pull-downs (Fig. S2a). Interestingly, SAM50 and

Mitofilin showed a higher interaction score in EGFP-ORP5ΔPH immunoprecipitates,

as compared to EGFP-ORP5. Of note, metaxin-2, another component of the MIB

complex, localized to the outer mitochondrial membrane, was also detected in the

MS of immunoprecipitated EGFP-ORP5ΔPH, whereas its score in the EGFP-ORP5

was lower than the assigned threshold (50) (Fig. S2a). To verify that the

overexpression of ORP5 or ORP5ΔPH did not affect the protein levels of SAM50 and

Mitofilin, WB analysis using anti-actin as loading control was carried out on cells

expressing EGFP-ORP5, EGFP-ORP5ΔPH or EGFP alone. Our results show that

neither EGFP-ORP5 nor EGFP-ORP5ΔPH overexpression alters the amount of

SAM50 and Mitofilin proteins as compared to the overexpression of EGFP alone (Fig.

S2b).

To confirm ORP5/8 interaction with SAM50 and Mitofilin, GFP-pull down

experiments from HeLa cells expressing EGFP-ORP5, EGFP-ORP8 or EGFP alone

were carried out (Fig. 2d). Consistent with the MS data, endogenous SAM50 and

Mitofilin were recovered with both EGFP-ORP5 and EGFP-ORP8 but not with EGFP

alone, confirming specific interaction of ORP5 and ORP8 with SAM50 and Mitofilin.

Of interest, the interaction of SAM50 and Mitofilin with EGFP-ORP5 was stronger

than with EGFP-ORP8.

Parallel GFP-pull down experiments were carried out from cells expressing

EGFP-tagged ORP5 (EGFP-ORP5ΔPH) or ORP8 (EGFP-ORP8ΔPH) PH domain

deleted constructs, and compared to the full-length proteins (EGFP-ORP5 and

EGFP-ORP8) or to the EGFP alone. In accord with the MS data, the deletion of the

PH domain increased ORP5 interaction with SAM50, as compared to the full-length

protein (Fig. 2e). In addition, EGFP-ORP8ΔPH interaction with SAM50 was higher

than of EGFP-ORP8, although still lower than the interaction of EGFP-ORP5ΔPH or

even EGFP-ORP5 with SAM50.

As the PH domain is not required for the interaction with SAM50 we further

investigated the role of the other domains of ORP5 in such interaction by

immunoprecipitating ORP5 deletion mutants for the ORD and the transmembrane

(TM) domains (EGFP-ORP5ΔORD, EGFP-ORP5ΔTM). While the deletion of the

ORD domain did not affect the interaction between ORP5 and SAM50, the deletion of

the TM domain decreased the amount of SAM50 co-immunoprecipitated with ORP5,

indicating that ORP5 should be properly anchored to the ER to localize at ER-

mitochondria contact sites and to interact with SAM50 (Fig. 2d).

To observe ORP5 and SAM50 localization we performed confocal analysis on

cells expressing EGFP-ORP5 or EGFP-ORP5ΔPH and stained with an anti-SAM50

antibody to look at the endogenous protein (Fig. 2e). As expected, endogenous

SAM50 localizes on mitochondria in cells stained with Mitotracker and, as previously

shown (Galmes et al., 2016), EGFP-ORP5 localizes to both ER-PM and ER-

mitochondria contact sites while the EGFP-ORP5ΔPH loses its localization to ER-PM

contacts. Both full-length and ΔPH constructs co-localized with endogenous SAM50

at ER elements in contact with mitochondria (Fig. 2e). However, this was more

evident in the case of EGFP-ORP5ΔPH (bottom image, Fig. 2e).

To test whether ORP5 and the MIB complex functionally cooperate in non-

vesicular transfer of PS we depleted ORP5, SAM50 or Mitofilin alone or in

combination by RNAi and analyzed the content in mitochondrial PE newly

synthesized from PS using the same radiometric PS-to-PE conversion assay *in*

situ described above. Robust knockdown of ORP5, SAM50 or Mitofilin was

confirmed by western blotting after 48 hours (Fig. 3a). Analysis of PS-derived

newly synthetized PE revealed a significant decrease in PE in ORP5 (Fig. 1d and

Fig. 3b) and Mitofilin knockdown cells (Fig. 3b). Moreover, the double-knockdown

of ORP5 and Mitofilin had an additive effect, supporting a cooperative function of

these two proteins in the same process (PS transport/PE synthesis). However, the

levels of PE were not changed in SAM50 knockdown cells as compared to control.

This can be explained by the fact that other subunits of the MIB complex (i.e.

Mitofilin) might compensate for its depletion. Indeed, levels of Mitofilin are

increased in SAM50 knockdown cells (Fig. 3a). Accordingly, the double-silencing

of SAM50 and either ORP5 or Mitofilin had a significant impact on PE synthesis

(Fig. 3b). Moreover, the reduction in PE was even stronger as compared to the

individual knockdowns indicating that disruption of two components of PS

transport pathways (*via* direct transport at ER-mitochondria contact sites and/or

via OMM-IMM tethering) has a significant impact on PE synthesis.

To verify the possibility of an indirect effect of Mitofilin or SAM50 silencing on the

morphology and abundance of ER-mitochondria contacts we carried out a

morphological analysis by conventional EM in all these knockdown conditions and

quantified ER-mitochondria contacts by HRP-KDEL EM (to stain the ER) in Mitofilin

or SAM50 silenced cells. Morphological analysis by conventional EM showed that

transient (48 hours) knockdowns of SAM50 and Mitofilin, as well as double-

knockdowns of ORP5+Mitofilin, ORP5+SAM50 and SAM50+Mitofilin, induce

formation of multilamellar cristae, almost devoid of CJ (Fig. S3a-b), complementing

previous observations obtained by other groups through stable disruption of the

MICOS/MIB functions (Ding, Wu et al., 2015, Ott et al., 2015). However, in all

knockdown conditions analyzed ER-mitochondria contact sites were still present and

their morphology not altered. Quantitative morphological analysis by HRP-KDEL EM

in control and Mitofilin or SAM50 silenced cells confirmed that the abundance of ER-

mitochondria contact sites was not altered by Mitofilin or SAM50 knockdown (Fig. 3c-

d) indicating that the effects on PS-derived PE synthesis were specifically due to

Mitofilin or SAM50 loss of function effects on intra-mitochondrial membrane bridges.

Overall our results indicate that ER-mitochondria contacts where ORP5/8 localize

are physically and functionally linked to intra-mitochondria contacts where the

components of the MIB/MICOS complex reside, and this association facilitates the

transfer of PS from the OMM to the IMM for synthesis of PE.

ORP5 additionally regulates VDAC-mediated Ca^{2+} import to mitochondria

through its interaction with SAM50

To get further insights into the functional link between ORP5/8 and the

MIB/MICOS complex we explored whether ORP5 and ORP8 knockdowns might

affect the expression and/or the localization of SAM50 and Mitofilin. Interestingly

when we analyzed cell lysates by WB, we found that SAM50 and Mitofilin protein

expression was significantly and specifically increased in ORP5- but not in ORP8

knockdown cells, as compared to control cells (Fig. 4a-b). The effects of ORP5

silencing were specific for SAM50 and Mitofilin, as other mitochondrial proteins

(Cytochrome c, CV-ATP5A) or proteins of the general ER (PDI) and of the MAMs

(VAPB) were not affected (Fig. S2c). This could also be observed by IF analysis

where the increase in SAM50 staining is specific to ORP5 knockdown (Fig. 4c).

Intriguingly, such increase in SAM50 was accompanied by a dramatic

accumulation in the voltage-dependent anion channel VDAC, a 19-stranded

transmembrane β-barrel protein (Colombini, 2012) that requires SAM50 for its

biogenesis and insertion in the OMM (Hohr et al., 2018). These results suggest that the

accumulation of VDAC protein in ORP5 knockdown cells is due to the parallel

increase of SAM50. The simultaneous knockdowns of ORP5 and SAM50 rescued

the increase in VDAC, confirming that its accumulation in ORP5 knockdown cells

was due to the accumulation of SAM50. Interestingly, ORP5 and SAM50 double-

knockdown did not rescue the increased protein levels of Mitofilin (Fig. 4a-b), in

accord with the fact that SAM50 knockdown alone also increased the protein levels

of Mitofilin. To verify that the increase in their protein amounts did not depend on the

increase in their transcripts we quantified the transcript levels of ORP5, SAM50 and

the three VDAC isoforms (VDAC1, 2 and 3) by real-time PCR upon ORP5 or SAM50

knockdown (Fig. 4d). The quantity of SAM50 and Mitofilin transcripts was unaffected

by ORP5 silencing, and VDAC1 transcript levels were even slightly reduced, possibly

reflecting a compensatory mechanism induced by the increase in VDAC protein.

Similarly, ORP5, VDAC and Mitofilin transcript levels were not affected in the

absence of SAM50. Overall, these data indicate that ORP5 knockdown induces an

accumulation of SAM50 and VDAC (SAM50-dependent) proteins in the absence of

major effects on the corresponding mRNAs.

As VDAC is one of the major mitochondrial players in mediating Ca^{2+} transport

and signaling between ER and mitochondria (Rapizzi, Pinton et al., 2002), we next

investigated if ORP5 knockdown could regulate Ca^{2+} influx from the ER to

mitochondria in living cells. We first used a ratiometric Ca^{2+} indicator selectively

targeted to mitochondria (mt-Pericam) which is based on a circularly permuted yellow

fluorescent protein fused to calmodulin (Nagai, Sawano et al., 2001). Upon binding to

Ca^{2+}, the excitation peak of mt-Pericam shifts from 415 nm to 494 nm while the

emission spectrum remains unchanged at 515 nm, allowing us to quantify rapid

changes in mitochondrial Ca^{2+}. We thus transfected ORP5, ORP8 or Ctrl siRNA-

treated cells with mt-Pericam and then we measured mitochondrial Ca^{2+} fluxes, in

response to histamine, by live-cell spinning disk microscopy (Fig. 5a-b). Soon after

the addition of histamine mt-Pericam fluorescence ratio (488/405) rapidly increased

in all conditions, reflecting the increase in mitochondrial Ca^{2+} influx triggered by the

histamine treatment. However, interestingly, ORP5 knockdown cells showed a

significantly higher increase as compared to ORP8 knockdown and to control cells.

Quantification of the maximum amplitude of the response after histamine treatment

(expressed as R_{max}/R_0, where R is the 488/405 ratio, R_{max} is the maximum

fluorescence ratio after histamine addition and R_0 the basal ratio before histamine

addition) confirmed that ORP5 but not ORP8 silencing induced a statistically relevant

increase of mitochondrial Ca^{2+} influx, triggered by the addition of histamine, as

compared to control cells (Fig. 5b).

Passive entry of Ca^{2+} into mitochondria can be driven by large negative

membrane potential of mitochondria and occurs through the mitochondrial Ca^{2+}

uniporter MCU spanning the IMM (Patron, Raffaello et al., 2013). To be sure that the

increase of mitochondrial Ca^{2+} upon ORP5 knockdown was not indirectly due to a

change of mitochondrial membrane potential, we performed flow cytometric analysis

in ORP5 or ORP8 knockdown cells. Knockdown cells were loaded with the

mitochondrial membrane potential sensitive dye TMRE (Tetramethylrhodamine ethyl

ester), in the presence (or absence) of the mitochondrial oxidative phosphorylation

uncoupler CCCP (Carbonyl cyanide 3-chlorophenylhydrazone) to induce membrane

depolarization (Fig. S4a). Neither ORP5 nor ORP8 knockdown altered mitochondrial

membrane potential as compared to control cells, even in presence of CCCP that

strongly decreases TMRE fluorescence intensity (Fig. S4a), showing that the effect of

ORP5 knockdown on Ca^{2+} is not due to a change of mitochondrial membrane

potential.

To further study the role of ORP5 in the regulation of Ca^{2+} homeostasis a parallel

set of experiments was performed using another type of Ca^{2+}-probe, the red

fluorescent genetically encoded Ca^{2+} indicator (R-GECO) and its two variants

selectively-targeted to mitochondria (mito-LAR-GECO) or ER (ER-LAR-GECO) to

respectively measure cytosolic, mitochondrial and ER Ca^{2+} in living cells (Fig. 5c-e).

These experiments confirmed that ORP5 silencing increases mitochondrial Ca^{2+}

uptake (Fig. 5c). The increase in mitochondrial Ca^{2+} uptake triggered by histamine

treatment was accompanied by a decrease in ER Ca^{2+} and an increase in cytosolic

Ca^{2+} in all conditions (Fig. 5d-e). Interestingly, no statistically relevant differences of

the decrease ER Ca^{2+} were measured between ORP5 silenced and control cells.

This indicates that the effects on mitochondrial Ca^{2+} fluxes observed in ORP5 do not

depend on the modification of ER Ca^{2+} storage or release. An increase in cytosolic Ca^{2+}

was measured in ORP5 and ORP8 silenced cells as compared to control. As this

increase was not specific of ORP5 silencing, it likely reflects a common role of

ORP5/8 in Ca^{2+} homeostasis at ER-PM contact sites, as recently proposed (Pulli,

Lassila et al., 2018).

So far, our data show that absence of ORP5 increases mitochondrial Ca^{2+}

possibly acting on the mitochondrial proteins SAM50 and VDAC and this is supported

by their specific accumulation in the absence of ORP5. To confirm that the increase

of mitochondrial Ca^{2+} in ORP5-silenced cells was indeed dependent on VDAC and

SAM50, we measured mitochondrial Ca^{2+} using mt-Pericam in VDAC- (all three

isoforms), ORP5+VDAC- and ORP5+SAM50- doubly-silenced cells. A significant

decrease in VDAC and SAM50 as well as ORP5 levels was detected by WB in the

corresponding silenced cells (Fig. 6a and Fig. 4a). Even though the increased level of

VDAC protein induced by the absence of ORP5 was not completely rescued in

ORP5+VDAC or ORP5+SAM50 knockdowns, the increase in mitochondrial Ca^{2+}

phenotype was significantly rescued to control levels and partially rescued in

ORP5+VDAC and ORP5+SAM50-silenced cells respectively (Fig. 6b and c). Overall,

these results demonstrated that ORP5 plays an additional role in regulating import of

mitochondrial Ca^{2+} through regulation of SAM50 and VDAC proteins.

ORP5 knockdown induces cell senescence in a SAM50-dependent manner

Accumulation of mitochondrial Ca^{2+} has been shown to be involved in both

apoptotic cell death and senescence (permanent cell proliferation arrest) (Wiel,

Lallet-Daher et al., 2014). Interestingly, it also has been shown that ORP5 expression

promotes cell proliferation (Du, Zadoorian et al., 2018). As our data show a new role

of ORP5 in regulating Ca^{2+} import to mitochondria at ER-mitochondria contact sites

(Fig. 6), we were wondering if ORP5 could mediate cellular processes such as

apoptosis and senescence. To address this question, we decided to study both

apoptosis and senescence in ORP5 depleted cells.

For apoptosis, we performed immunofluorescence and stained with anti-cleaved

caspase-3 (to detect apoptotic cells) HeLa cells transfected with control siRNA or

siRNA against ORP5 or ORP8. We also treated the knockdown cells with

staurosporine to induce apoptosis or with DMSO as control (Fig. S4b). Quantification

of the percentage of cleaved caspase-3 positive cells (% of total cells) in all

conditions showed no significant difference between ORPs knockdown and control

cells (Fig. S4b), even in presence of staurosporine, showing that in these conditions

(transient silencing) ORP5/8 are not mediating apoptosis.

To study cellular senescence we performed knockdown of ORP5 in MRC5

normal human fibroblasts that are a good model for senescence studies (Ma,

Warnier et al., 2018). We first looked at cell proliferation using crystal violet staining

in both control and ORP5 knockdown cells (Fig. 7a). A significant decrease of the cell

number was detected in ORP5 depleted cells compared to control (Fig. 7a). This

decrease of the cell number in ORP5 knockdown cells could be correlated with an

increase of the Senescence-associated β –galactosidase (SA β-gal) staining in this

condition as compared to control (Fig. 7a). Accordingly, these results were

corroborated by qRT-PCR analysis showing a significant increase of senescence

markers p16 and p21 and a significant decrease of the proliferation marker Ki67 in

ORP5 knockdown cells compared to control (Fig. 7b). Because we have shown that

ORP5 knockdown induces an elevation of mitochondrial Ca^{2+} via a SAM50-

dependent accumulation of VDAC, we hypothesized that the induction of senescence

in MRC5 cells was also due to the effect of ORP5 on SAM50. Thus, we decided to

perform a double-knockdown of ORP5 and SAM50 in MCR5 cells. In accord with our

hypothesis, double-silencing of ORP5 and SAM50 partially rescued the senescence

phenotype found in ORP5 knockdown cells. ORP5+SAM50 double-knockdown cells

displayed increased proliferation and underwent less senescence as compared to

ORP5 single knockdown cells (Fig. 7c). To verify that this effect on cellular

senescence was specific to ORP5, given its role on mitochondria Ca^{2+}, we performed

in parallel a knockdown of ORP8 in MRC5 cells. Differently to ORP5, ORP8

knockdown did not induce cell senescence (Fig. 7c). Taken together, these data

reveal a novel specific role of ORP5 and involving SAM50 in regulating cellular

senescence in MRC5 cells.

DISCUSSION

In this study, by using a combination of *in vitro* and *in situ* biochemical and imaging approaches, we describe a new function of ORP5/8 in the maintenance of mitochondrial levels of PE, an essential phospholipid of mitochondria, in mammalian cells. Importantly, we provide the first evidence of LTPs (ORP5 and ORP8) directly mediating non-vesicular transfer of PS at ER-mitochondria contact sites, from the ER to the mitochondria where it is converted into PE.

In yeast, large complexes such as the ERMES and the ER-Membrane protein Complex (EMC) have been previously involved in phospholipid trafficking (PC and PS, respectively) between the ER and the mitochondrial membranes (Tamura & Endo, 2017) (Lang, John Peter et al., 2015). For instance, three subunits of the ERMES complex contain a Synaptotagmin-like Mitochondria lipid binding Protein (SMP) domain that forms a hydrophobic lipid transport cavity shared by other proteins of the tubular lipid-binding protein TULIP superfamily (Kopec, Alva et al., 2010, Reinisch & De Camilli, 2016). However, a direct role (independent of its tethering function) of ERMES in lipid transport at ER-mitochondria contact sites is still questioned. In addition, the ERMES complex is not conserved in metazoans. The EMC complex, which has been involved in PS shuttling at ER-mitochondria contact sites in yeast, is instead highly conserved in metazoans (Wideman, 2015). However, no lipid-binding/transfer domain has been found in the EMC proteins, indicating that their implication in lipid transfer at ER-mitochondria contacts is linked to their tethering function rather than to a direct lipid transfer activity. Recently, the mammalian LTP VPS13A has been shown to localize to contact sites including ER-mitochondria contacts. VPS13A contains a lipid-binding domain (VPS13α) that has the ability to harbor multiple phospholipids at once and transfer them between

liposomes *in vitro*. However its role in lipid transfer at membrane contact sites *in situ* has not been established yet. Differently from the SMP and VPS13α that can simultaneously host multiple phospholipids, the ORD domain of Osh/ORPs forms a cavity that can host only one lipid at a time (Maeda, Anand et al., 2013, Wang, Ma et al., 2019).

ORP5 and ORP8 have recently been shown to counter-exchange PS with PM phosphoinositides PI4P and PIP_2 at ER-PM contact sites in HeLa cells (Chung et al., 2015, Ghai, Du et al., 2017). However PI4P and PIP_2 are not present on the mitochondrial membranes, while PE is highly abundant on these membranes, in addition to being an essential lipid of all biological membranes. Our *in vitro* data show that the ORD domains of ORP5 and ORP8 transport PS, but not other phospholipids such as PE and PC, indicating a specific role of ORP5/8 in PS transport and excluding the possibility that ORP5/8 might also participate in the transport of a fraction of PE back to the ER. It is possible that ORP5/8 cooperate with other LTPs, such as VPS13A, for the exchange of other lipids (including PE) at ER-mitochondria contact sites. Importantly, we have confirmed the role of ORP5/8 in PS transfer by measuring a decrease of PS-derived mitochondrial PE in ORP5 (and even more in ORP5+8) depleted HeLa cells *in situ*. Accordingly, the knockdown of ORP5/8 affects cristae morphology (Fig.1e) and the respiratory function of mitochondria (Galmes et al., 2016), all phenotypes that are expected in the case of decrease in mitochondrial PE (Joshi et al., 2012, Steenbergen et al., 2005). Our data suggest that the gradient of PS at ER-mitochondria contacts is sufficient to trigger the ORP5/8-mediated unidirectional transport of PS from the MAMs, where it is highly enriched, to the mitochondria membranes, where it is rapidly converted into PE and is therefore present at a very low amount. Our findings have important implications in the general

field of LTPs, as they suggest that the same LTP can use different means to transfer

lipids depending on the local gradients present at the specific membrane contact

sites where it is localized. They also suggest a possible new function in PS transport at

ER-mitochondria contact sites for the yeast orthologues of ORP5/8, Osh6/7p, whose

localization/function at ER-mitochondria contact sites still remain to be addressed

(Moser von Filseck, Copic et al., 2015).

Another important feature of our work is the observation that ORP5 and ORP8

knockdowns do not affect the extent of ER-mitochondria contact sites. These data,

consistent with our previous observation that ORP5/8 overexpression does not alter

the length of ER-mitochondria contacts (Galmes et al., 2016), confirm that the main

function of these proteins at MAMs is lipid transfer and not membrane tethering. This

is a unique feature among the LTPs that have been identified so far at MAMs. For

instance, modulation of the protein levels of other LTPs such as ERMES in yeast, or

VPS13A in mammals, results in an alteration of the extent of ER-mitochondria

contact sites (Kornmann, Currie et al., 2009, Kumar, Leonzino et al., 2018). Thus,

ORP5/8 represent so far a unique tool to study lipid transport at ER-mitochondria

contact sites, independently of membrane tethering.

Importantly, our study also reveals that ORP5/8 physically and functionally

interact with SAM50 and Mitofilin, two key components of the MIB/MICOS complex

that anchor the IMM to the OMM at the level of CJ (Huynen et al., 2016, Ott et al., 2015,

Wollweber et al., 2017). The biochemical interaction between ORP5 and

SAM50/Mitofilin suggests the existence of a physical link between ER-mitochondria

contact sites involved in lipid transport and intra-mitochondria contacts. ORP5

localization by IEM at ER-mitochondria contact sites near the CJ, where Mitofilin and

MICOS complex also reside, confirms the existence of such tripartite membrane

structure. Importantly, we also show that ORP5 cooperates with SAM50 and Mitofilin to facilitate the movement of PS from the ER to the IMM across ER-mitochondria contact sites and the IMS. Taken together these findings provide the first evidence of a physical and functional link between ER-OMM and OMM-IMM contact sites to facilitate transport of PS from the ER to the mitochondria.

Finally, we uncover a specific role for ORP5, but not ORP8, in regulating Ca^{2+} import to mitochondria, via a SAM50-VDAC axis. SAM50, in addition to be part of the MIB complex, is also the major subunit of the Sorting and Assembly Machinery (SAM) that mediates membrane insertion and assembly of mitochondrial β-barrel proteins (Hohr et al., 2018, Kozjak et al., 2003). One of the most abundant mitochondria β-barrel proteins is VDAC, which forms a pore that allows the passage of small metabolites and ions such as ATP and Ca^{2+} across the OMM (Colombini, 2012). VDAC exists as three isoforms (VDAC1, 2 and 3) that share common channeling properties. VDAC interacts with the IP3Rs located in the ER membrane via the molecular chaperone GRP75 that links the two proteins together. The interaction between IP3Rs and VDAC acts as a bridge to allow the transfer of Ca^{2+} from the ER to the OMM, and then from the OMM to the mitochondrial matrix through the mitochondrial Ca^{2+} uniporter MCU. Ca^{2+} release from the ER to mitochondria is triggered by IP_3 binding to IP3Rs upon histamine treatment (Rizzuto, Marchi et al., 2009). Here we show that knockdown of ORP5, but not ORP8, increases the protein levels of SAM50 and VDAC as well as the levels of histamine-induced Ca^{2+} import to mitochondria. Importantly, double-knockdown of ORP5 and VDAC rescues the mitochondrial Ca^{2+} increase induced by ORP5 silencing, confirming that this increase in Ca^{2+} is indeed VDAC-dependent. Mitochondrial membrane potential has been proposed to influence VDAC open-closed state and consequently import of Ca^{2+} to mitochondria (Tan & Colombini, 2007). However, we did

not detect any alteration in mitochondria membrane potential in ORP5 knockdown cells,

supporting our idea that the increase in mitochondrial Ca^{2+} observed in ORP5-silenced

cells is related to the increase in VDAC protein levels. This is also in accord with

previous studies showing that the overexpression of VDAC in HeLa cells enhances

mitochondrial Ca^{2+} uptake upon histamine treatment (Rapizzi et al., 2002). Our data

also indicate that the effect of VDAC accumulation on mitochondrial Ca^{2+} fluxes does not

depend on the modification of ER Ca^{2+} storage or release. Indeed, we observe a similar

decrease of ER Ca^{2+} in ORP5 depleted and control cells, upon histamine treatment. Our

results correlate well with those of others (Rapizzi et al., 2002) who also observed similar

decrease of ER Ca^{2+} in VDAC-overexpressing and control cells.

Accumulation of Ca^{2+} in mitochondria, induced by release of Ca^{2+} through IP3R

channel, has been recently shown to be involved in cellular senescence in normal

cells (Martin & Bernard, 2018, Wiel et al., 2014). Interestingly, we found that

knockdown of ORP5, but not ORP8, induces cellular senescence in MCR5 cells.

These findings are highly relevant because ORP5 overexpression has been

associated to several tumors (Du, Turner et al., 2017). We also show that double-

knockdown of ORP5 and SAM50 restores the levels of VDAC protein, as well as of

mitochondrial Ca^{2+}, and enables cells to escape from the senescence fate induced by

ORP5 depletion. These data confirm that all these phenotypes are dependent on

SAM50, and possibly on its known role in the biogenesis of VDAC (Hohr et al., 2018).

Future studies are required to further dissect the mechanisms underlying the role of

ORP5 in modulating SAM50 protein levels in mitochondria.

In conclusion, our data reveal that: 1) ORP5/8 constitute the molecular machinery

mediating PS transfer at ER-mitochondria contact sites; 2) ER-mitochondria contacts

where ORP5/8 localize are physically and functionally associated with intra-

mitochondrial contacts to facilitate the transport of PS from the ER to mitochondria

membranes; 3) ORP5 plays a novel specific role in regulating cellular senescence in

MCR5 cells by modulating mitochondrial Ca^{2+} fluxes at ER-mitochondria contact sites

via SAM50/VDAC (Fig. 8). Our study uncovers a novel link between lipid and Ca^{2+}

transport at ER-mitochondria contact sites and provides a first molecular clue on how

lipids are transported at ER-mitochondria contact sites. Furthermore, we reveal a

specific role of ORP5 in regulating mitochondrial Ca^{2+} homeostasis and cellular

senescence in mammalian cells.

METHODS

Cell culture, siRNA and transfection.

HeLa cells were cultured in DMEM (Life Technologies) containing GlutaMax (Life

Technologies) and supplemented with 10% FBS (Life Technologies) and 1%

penicillin/streptomycin (Life Technologies) at 37°C and 5% CO_2. Transfection of

plasmids and siRNA oligos (Dharmacon, GE Healthcare) was carried out with

lipofectamine 2000 and oligofectamine (Life Technologies) according to

manufacturer's instructions.

For senescence studies, MRC5 normal human fibroblasts (ATCC) were cultured in

Dulbecco's modified Eagle's medium (Life Technologies) containing GlutaMax (Life

Technologies) and supplemented with 10% FBS (Sigma-Aldrich) and 1%

penicillin/streptomycin (Life Technologies). MRC5 were transfected with siRNAs

using Dharmafect 1 transfection reagent according to manufacturer's instructions

(Dharmacon, GE Healthcare).

siRNAs oligonucleotides

For knockdowns, HeLa cells were transfected with siRNA oligos by using

oligofectamine (Life Technologies) and cultured for 48 hours prior to analysis.

For senescence studies, MRC5 cells were transfected with siRNAs using Dharmafect

1 transfection reagent according to manufacturer's instructions (Dharmacon, GE

Healthcare).

Double-stranded siRNAs were derived from the following references:

siRNA	Company, Reference
OSBPL8	Dharmacon, J-009508-06 (Galmes et al., 2016)
OSBPL8	Dharmacon, J-009508-05 (Galmes et al., 2016)

OSBPL5	Dharmacon, J-009274-10 (Galmes et al., 2016)
OSBPL5	Dharmacon, J-009274-11 (Galmes et al., 2016)
SAMM50	Dharmacon, J-017871-18
SAMM50	Dharmacon, J-017871-19
Non-targeting	Dharmacon, D-001810-10
hVDAC1	5'-AAGCGGGAGCACATTAACCTG-3' (De Stefani, Bononi et al., 2012)
hVDAC2	5'-AAGGATGATCTCAACAAGAGC-3' (De Stefani et al., 2012)
VDAC3	5'-AAGGGTGGCTTGCTGGCTATC-3' (De Stefani et al., 2012)
siMitofilin	5' -AAUUGCUGGAGCUGGCCUUTT-3' (John, Shang et al., 2005)

Plasmids and cDNA clones

EGFP-ORP5, EGFP-ORP8, EGFP-ORP5ΔPH, EGFP-ORP8ΔPH, EGFP-

ORP5ΔORD and EGFP-ORP5ΔTM were described in (Galmes et al., 2016). The

following reagents were kind gifts: 2mt RP ratiometric Pericam from Paolo Pinton

(University of Ferrara) (Filippin, Abad et al., 2005); ssHRP-KDEL from T. Schikorski

(Schikorski, Young et al., 2007); GST-ORD8 (ORD ORP8, corresponding to aa 328-

767) from P. De Camilli (Chung et al., 2015). GECO constructs were from Robert

Campbell (Addgene: CMV-R-GECO1 # 32444, CMV-mito-R-GECO1 # 46021, CMV-

ER-LAR-GECO1 # 61244, CMV-mito-LAR-GECO1.2 # 61245).

Cloning of HA-ORP5, Mitofilin-GFP and GST- ORD5 (ORD ORP5)

cDNAs of ORP5 (full-length), Mitoflin (full-length from FLAG-mitofilin (Ott et al.,

2015)) and GST-ORD5 (corresponding to aa 265-703), were amplified by PCR. In all

PCR reactions, Herculase II fusion DNA polymerase (Agilent) was used. Primers

used were (coding sequence shown in lowercase):

5' AgeI-HA-ORP5_Fw GGCGGC *ACCGGT* cgccacc ATGTACCCATACGATGTTCCA

GATTACGCT atgaaggaggaggccttcctc

3' XhoI-STOP-ORP5_Rv GGC *CTCGAG* ctatttgaggatgtggttaatg

5' KpnI-Mitofilin_Fw AGACCCAAGCTT *GGTACC* atg

3' BamHI-GC-Mitofilin_Rv GTAATC *GGATTC* GC ctctggct

5' SalI-TC-ORD5_Fw GCACAG GTCGAC TC gagacccctggggccccggt

3' NotI-STOP-ORD5_Rv GCACA *GCGGCCGC* ctactgtggccggagggctggtcg

For the HA-ORP5 cloning the PCR product (carrying the HA tag at the N-terminus of

ORP5) was ligated between AgeI and XhoI in the pEGFP-C1 vector (Clontech) and

replacing the GFP- with the HA-tag. For the other clonings the PCR products were

ligated between KpnI and BamHI for Mitofilin, and between SalI and NotI for ORD5,

in the pEGFP-N1 vector (Clontech) to generate Mitofilin-EGFP or in the pGEX-6P-1

to generate GST-ORD5.

mRNA analyses by quantitative by reverse transcriptase PCR (qPCR)

Total RNA was isolated from HeLa cells transfected with siRNAs for 48 hours as

described above, by using a Purelink™ kit (Ambion/Thermo Scientific, Foster City,

CA). The RNA (0.5 μg per specimen) was reverse transcribed with a SuperScript

VILO™ cDNA synthesis kit (Invitrogen/Thermo Scientific, Carlsbad, CA) according to

the manufacturer's protocol. Quantification of the mRNAs of interest was carried out

on a Roche Lightcycler™ 480 II instrument by using SYBR-Green I Master mix

(Roche, Basel, Switzerland) and primers specified in Table SX. Succinate

dehydrogenase complex. subunit A, the mRNA of which remained markedly stable

under the present conditions, was employed as a reference housekeeping mRNA.

Relative mRNA levels were calculated by using the $-\Delta\Delta$Ct method.

Sequences of the primers used for qPCR:

mRNA	Sequence 5'-3'
SDHA* (housekeeping)	Fw: CATGCTGCCGTGTTCCGTGTGGG
	Rv: GGACAGGGTGTGCTTCCTCCAGTGCTCC
ORP5	Fw: GTGCCGCTGGAGGAGCAGAC
	Rv: AGGGGCTGTGGTCCTCGTATC
SAMM50	Fw: CAAGTGGACCTGATTTTGGAGG
	Rv: AGACGGAGCAATTTTTCACGG
Mitofilin	Fw: GTTGTATCTCAGTATCATGAGCTGG
	Rv: GTTCAGCTGATCAATACGACGA
VDAC1	Fw: ACGTATGCCGATCTTGGCAAA
	Rv: TCAGGCCGTACTCAGTCCATC
VDAC2	Fw: GCTACAGGACTGGGGACTTC
	Rv: AATGCCAAAACGAGTGCAGTT
VDAC3	Fw: GTAAATAATGCCAGCCTGATTG
	Rv: CTTCCAGGGACAAATCTGATG

*succinate dehydrogenase complex, subunit A

For senescence studies, MCR5 cells RNAs were extracted using RNA Extracol lysis

buffer (Dutscher). cDNAs were synthesized from total RNA using the Maxima first

strand cDNA synthesis kit (Life Technologies). Quantitative PCR mixture contained

cDNA, TaqMan mix (Roche), 100 µM of a Universal Probe Library probe (Roche) and

200 nM of primers (Sigma-Aldrich) according to manufacturer's instructions. qPCRs

were carried out on a FX96 Thermocycler (Bio-Rad). qPCR reactions were as

follows: 95°C 10 min, followed with 40 cycles of 95°C 10s, 59°C 30s. The reactions

were performed at least in duplicate. The relative amount of mRNA (ORP5, p16, p21,

Ki67, PGK1 and HPRT1) was calculated using the comparative Ct ($\Delta\Delta$Ct) method

and data were normalized using two housekeeping genes (PGK1 and HPRT1).

Sequences of the primers and UPL probes used for qPCR:

mRNA/UPL probe	Sequence 5'-3'
CDKN2A (p16) / UPL 34	Fw: GTGGACCTGGCTGAGGAG
	Rv: CTTTCAATCGGGGATGTCTG
p21(CDKN1A) / UPL 32	Fw: TCACTGTCTTGTACCCTTGTGC
	Rv: GGCGTTTGGAGTGGTAGAAAT
Ki67 / UPL 32	Fw: TCAAGGAACTGATTCAGGAGAAG
	Rv: GTGCACTGAAGAACACATTTCC
ORP5 / UPL 17	Fw: GTTGTATCTCAGTATCATGAGCTGG
	Rv: GTTCAGCTGATCAATACGACGA
PGK1 / UPL probe 67	Fw: CAGCTGCTGGGTCTGTCAT
	Rv: GCTGGCTCGGCTTTAACC
HPRT1 / UPL probe 73	Fw: TGACCTTGATTTATTTTGCATACC
	Rv: CGAGCAAGACGTTCAGTCCT

781   **Antibodies, dyes and reagents list**

782   Primary antibodies used in this study were:

	Antibody	Company & Reference
WB	IP3R-3	BD Transduction Laboratories, 610312
WB	ORP5	SIGMA, HPA038712
WB	ORP5	SIGMA, HPA058727
WB	beta-Actin	Abcam, ab8226
WB	ORP8	GeneTex, GTX121273
IF, WB	SAMM50	SIGMA, HPA034537
WB	Mitofilin	Proteintech, 10179-1-AP
WB	VDAC1	Abcam, ab15895
WB	Cytochrome C	BD Pharmingen, 556433
WB	Total OXPHOS CV-ATP5A	Abcam, ab110413

WB	PDI	GeneTex, GTX30716
WB	VAPB	SIGMA, HPA013144
IF, WB	GFP	Roche, 11814460001
IEM	GFP	Life technologies, A11122
IF	HA	SIGMA, H3663
IF	Cleaved caspase-3	Cell Signaling, 9661
IF	FLAG M2	SIGMA, F1804
IF	MitoTracker® Red CMXRos	M7512, Invitrogen
IF	MitoTracker® Deep Red FM	M22426, Invitrogen

Biochemical analyses

Western blot

For immunoblotting, cells were resuspended in lysis buffer [50 mM Tris, 150 mM NaCl, 1% Triton X-100, 10 mM EDTA, pH 7.2, and protease inhibitor cocktail (Roche)]. Cell lysates were then centrifuged at 21 000 g for 20 min at 4°C. The supernatants were boiled in reducing SDS sample buffer and proteins were separated using 10% SDS-PAGE and immunoblot was carried using standard methods.

For Western blot quantification, bands of protein of interest were detected using ChemiDoc™ Imaging Systems (Life Science Research, Bio-Rad) and analyzed using Image Lab™ Software. All data are presented as mean ±SEM of three experimental replicates.

Immunoprecipitation of ORPs

HeLa cells transfected with EGFP-tagged ORPs were washed in cold PBS and lysed on ice in lysis buffer [50 mM Tris, 120 mM NaCl, 40 mM Hepes, 0,5% digitonin, 0,5%

CHAPS, pH 7.36, and protease inhibitor cocktail (Roche). Cell lysates were then

centrifuged at 21 000 g for 20 min at 4°C. Supernatants were then incubated with

Chromotek GFP-trap agarose beads (Allele Biotech) for 1 hour at 4°C under rotation.

Subsequently beads were washed in 0.1 M phosphate buffer. After extensive washes

in cold lysis buffer, immunoprecipitated proteins bound to the beads were processed

for Mass Spectrometry analysis (see below) or incubated in sample buffer (containing

2% SDS) and then boiled for 1 min. In the latter case immunoprecipitates were

loaded and separated in 10% SDS–PAGE gel and immunoblotting was carried out.

Cell fractionation

HeLa cells (100×10^6 cells) were harvested 48 hours after transfection with siRNA

oligos and washed with PBS by centrifugation at 600 g for 5 min. The cell pellet was

resuspended in starting buffer (225 mM mannitol, 75 mM sucrose and 30 mM Tris-

HCl pH 7.4) and homogenized using a Tissue Grinder Dura-Grind®, Stainless Steel,

Dounce (Wheaton). The homogenate was centrifuged three times at 600 g for 5 min

to remove nuclei and unbroken cells. The crude mitochondria was pelleted by

centrifugation at 10 000 g for 10 min. To separate MAM and pure mitochondria

fractions, the pellet was resuspended in MRB buffer (250 mM mannitol, 5 mM

HEPES and 0.5 mM EGTA, pH 7.4) and layered on top of different concentrations of

Percoll gradient (225 mM mannitol, 25 mM HEPES, 1 mM EGTA pH 7.4 and 30% or

15% Percoll). After centrifugation at 95 000 g for 30 min, two dense bands containing

either the pure mitochondria or MAM fraction were recovered and washed twice with

MRB buffer by centrifugation at 6300 g for 10 min to remove residual Percoll and

residual contamination. MAM was pelleted by centrifugation at 100 000 g for 1 hour.

MAMs and pure mitochondria pellets were resuspended in Lysis Buffer (50 mM Tris,

150 mM NaCl, 1% Triton X-100, 10 mM EDTA, pH 7.2, and protease inhibitor

cocktail) and protein concentrations were determined by Bradford assay. Equal

amount of proteins were loaded on 4-20% gradient SDS-PAGE gels (Biorad) and

immunoblotting was carried out. Pure mitochondria were processed for MS-lipidomic

analysis (see below).

Mass spectrometry analysis

Mass Spectrometry (MS) analysis was carried out by the proteomics/mass

spectrometry platform in IJM (http://www.ijm.fr/plateformes/spectrometrie-de-masse).

Briefly, after washes with binding buffer, immunoprecipitations beads were rinsed

with 100 µl of NH4HCO3 25 mmol/l. Proteins on beads were digested overnight at

37°C by sequencing grade trypsin (12,5 µg/ml; Promega Madison, Wi, USA) in 20 µl

of NH4HCO3 25 mmol/l. Digests were analysed by an Orbitrap Fusion (Thermo

Fisher Scientific, San Jose, CA) equipped with a Thermo Scientific EASY-Spray

nanoelectrospray ion source and coupled to an Easy nano-LC Proxeon 1000 system

(Thermo Fisher Scientific, San Jose, CA). MS/MS data were processed with

Proteome Discoverer 1.4 software (Thermo Fisher scientific, San Jose, CA) coupled

to an in house Mascot search server (Matrix Science, Boston, MA ; version 2.4.2).

MS/MS datas were searched against SwissProt databases with Homo sapiens

taxonomy.

Mass Spectrometry-lipidomic analysis

700 µl of homogenized cells were mixed with 800 µl 1 N HCl:CH_3OH 1:8 (v/v), 900 µl

$CHCl_3$ and 200 µg/ml of the antioxidant 2,6-di-tert-butyl-4-methylphenol (BHT; Sigma

Aldrich). The organic fraction was evaporated using a Savant Speedvac spd111v

(Thermo Fisher Scientific). Lipid pellets were reconstituted in running solution

(CH_3OH:$CHCl_3$:NH_4OH; 90:10:1.25; v/v/v). Phospholipid species were analyzed by

electrospray ionization tandem mass spectrometry (ESI-MS/MS) on a hybrid triple

quadrupole/linear ion trap mass spectrometer (4000 QTRAP system; Applied

Biosystems SCIEX) equipped with a TriVersa NanoMate (Advion Biosciences)

robotic nanosource. Phospholipid profiling was executed by (positive or negative)

precursor ion or neutral loss scanning at a collision energy of 35 eV for neutral loss

141 (phosphatidylethanolamine (PE)). Phospholipid quantification was performed by

multiple reaction monitoring (MRM), the transitions being based on the neutral losses

or the typical product ions as described above. The MRM dwell time was set to 100

ms and typically the signal was averaged over 20 cycles. Lipid standards used were

PE25:0 and PE43:6 (Avanti Polar Lipids). The data were corrected for isotope effects

as described by (Liebisch, Lieser et al., 2004).

Lipid Transfer assay

ORP5 and ORP8 ORD domain purification

Escherichia coli BL21DE3 RILP (Invitrogen) cells were transformed with plasmids

encoding for GST tagged ORP5 or ORP8 ORD domains following the manufacturer's

instruction. Bacteria were then grown overnight at 37°C and used to inoculate a

large-scale volume (1L). When the OD_{600} reached 0.4, cultures were cool down and

incubate at 18°C until they reached O_{D600} = 0.65. Cells were induced by addition

of isopropyl β-D-1-thiogalactopyranoside to a final concentration of 0.1 mM and were

incubated overnight at 18°C before harvesting. Cells were resuspended in 35 ml

binding buffer (1X PBS, 1 mM EDTA, 1 mM DTT, Protease inhibitor) then 250 units

of benzonase nuclease (Sigma) were added to the resuspension. Cells were lysed by

sonication and the supernatant was recover after 20 min centrifugation at 184 000g

and 4°C. Supernatant containing GST tagged proteins was incubated with 2 ml of

Glutathione Sepharose 4 fast flow for 1 hour at 4°C under nutation. Beads were

washed using a serie of wash buffer: 1st (1X PBS, 1 mM EDTA, 1 mM DTT), 2nd HSP-

removal buffer (50 mM Tris pH7.5, 50 mM KCl, 20 mM MgCl2, 5 mM ATP) then

cleavage buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM DTT).

Cleavage of the GST tag was realized overnight at 4°C using Prescission protease.

Untagged proteins were eluted with cleavage buffer, flash frozen and stored at -80°C

until lipid transfer assay was performed.

Lipids

1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1,2-dioleoyl-sn-glycero-3-

phosphoethanolamine-N-(cap   biotinyl)   (Biotinyl   Cap   PE),   1-palmitoyl-2-

(dipyrrometheneboron   difluoride)undecanoyl-sn-glycero-3-phosphoethanolamine

(TopFluor-PE),   1-palmitoyl-2-(dipyrrometheneboron   difluoride)undecanoyl-sn-

glycero-3-phospho-L-serine   (TopFluor-PS),   1-palmitoyl-2-(dipyrrometheneboron

difluoride)undecanoyl-sn-glycero-3-phosphocholine (TopFluor-PC) were purchased

from Avanti Polar Lipids as chloroform solutions.

Liposome preparation

1 µmol of the appropriate lipid mixtures in chloroform solution was dried in glass

tubes for 10 min under a gentle stream of argon, and then for 1 hour under vacuum.

The dried lipid films were resuspended in 1 ml of buffer H (25 mM HEPES/KOH, pH

7.7; 150 mM KCL; 10% Glycerol) by vigorously vortexing for 30 min at room

temperature. Unilamellar liposomes were produced by seven freeze-thaw cycles (30

sec in liquid nitrogen followed by 5 min in a 37°C water bath) and extrusion (at least

21 times) through a polycarbonate filter with 100 nm pore size (polycarbonate

membranes from Avanti Polar Lipids). The liposomes were then stored on ice.

 Lipid Transfer assay in vitro

The lipid transfer assays were realized with liposomes prepared as described above.

The donor liposomes contained 1% mol TopFluor lipids (-PS, -PC or -PE) and 2%

mol of Biotinyl Cap PE. The acceptor liposomes contained only POPC. For each

reaction, 25 µl of streptavidin-coated magnetic beads (DynabeadsMyOne

Streptavidin T1, Invitrogen) were washed in buffer H and mixed with 25 µl of 1 mM

donor liposomes. The mixture was incubated for 1 hour at 25°C with intermittent

gentle mixing. Bead-bound donor liposomes were then washed, resuspended in 25µl

and mixed with 25µl of 1 mM acceptor liposomes and 50µl of buffer H or protein (0.3

µM protein and 2.5 µM TopFluor lipids in the reaction solution). The mixture was

incubated at 37°C for 1 hour with intermittent gentle mixing. Supernatant containing

acceptor liposomes was recovered after binding of bead-bound donor liposomes to a

magnetic rack. TopFluor fluorescence of acceptor and donor liposomes was

measured (after solubilization with 0.4% (w/v) n-dodecyl-β-D-maltoside) in a

SpectraMax M5 plate reader (Molecular Device) equilibrated to 30°C (excitation: 450

nm; emission: 510 nm; cutoff: 475 nm; low gain). The percentage of lipids transferred

from donor to acceptor liposomes was determined using the following formula:

$100*F_{acceptor}/(F_{acceptor}+F_{donor})$.

Radiometric assay for the conversion of PS to PE

Hela cells were seeded on 6-well plates and transfected for 48 hours with the non-

targeting, ORP5 or ORP8-specific siRNAs specified above by using Oligofectamine

(Thermo Fisher Scientific). The cells were then washed and shifted into Hanks

balanced salt solution (Gibco) supplemented with a serine-free MEM amino acid

mixture and MEM vitamins (Gibco), followed by 4 or 18 hours labeling with 2 µl/well

L-[^{3}H(G)]serine (30.9 Ci/mmol, NET24800, Perkin-Elmer). After the labeling, the cells

were scraped into 0.9 ml 2% NaCl per well, a 0.1 ml aliquot was withdrawn for

protein analysis with the BCA assay (Thermo Fisher Scientific), and, after adding 50

nmol of unlabeled PS as carrier, the remaining 0.8 ml was subjected to lipid

extraction by an acid modification of the Folch method (Kim, Song et al., 2017). After

drying, the lipids were resolved in 50 μl CHCl$_3$ and applied on Merck TLC Silica gel

60^{TM} plates, followed by separation by using CHCl$_3$-methanol-acetic acid-H2O

(50:30:8:3.5) as solvent. The PS and PE spots identified from the mobility of

standards run on the same plates were scraped into scintillation vials for analysis of

[^{3}H] radioactivity. The DPM values were corrected for total cell protein, and the ratio

of [^{3}H] in PE vs. PS calculated.

Fluorescence Microscopy

Immunofluorescence and Confocal Microscopy

HeLa cells were seeded on 13 mm glass bottom coverslips (Agar Scientific).

Immunofluorescence was carried out one day after transfection. Transfected cells

were fixed with 4% PFA/PBS for 15 min at room temperature, washed in PBS and

incubated with 50 mM NH4Cl/PBS for 15 min at room temperature. After washing

with PBS and blocking buffer (1% BSA/ 0,1% Saponin in PBS), cells were incubated

with primary antibodies diluted in blocking buffer for 1 hour at room temperature and

then with fluorescently-labeled secondary antibodies. After washing with blocking

buffer and then PBS, coverslips were mounted on microscopy slides and images

were acquired on Confocal inverted microscope SP8-X (DMI 6000 Leica). Optical

sections were acquired with a 63x/1.4 Oil immersion objective using the LAS-X

software and fluorescent pictures were collected with a PMTs GaAsP hybride camera

(Hamamatsu). Images from a mid-focal plane are shown. Images were processed

using Image J.

Calcium (Ca^{2+}) indicators

Ca^{2+} intake assay using Geco constructs

HeLa cells were seeded on glass bottom ibidi chambers (μ-slide 2 wells) 3 days

before imaging. Knockdown of the proteins of interest was realized 2 days before

imaging, as described above. The day after knockdown, 0.8 μg of GECO plasmid

were transfected with 6 μl of lipofectamine 2000/well (Invitrogen), according to the

manufacturer's instructions. The day of imaging, cells were washed 4-5 hours before

the imaging with fresh DMEM. Cell imaging was performed on an inverted Nikon Ti

Eclipse E microscope coupled with a Spinning Disk (Yokogawa CSU-X1-A1) and

cage incubator to control both temperature and CO_2 (37 °C, 5% CO2). After

excitation with a 561 nm laser (Coherent, 100 mW), fluorescence from GECO

constructs was detected with a 40X oil immersion objective (PLAN FLUOR; NA: 1.30;

Nikon) and a Prime 95B sCMOS camera (Photometrics). Images were captured

every 5 sec during 10 min. Approximatively 1 min after the start of captures,

histamine was added to a final concentration of 100 μM to trigger Ca^{2+} release from

intracellular stores. Image analysis was performed using Image J by measuring the

fluorescence intensity over time (time-lapse) of cells expressing the GECO

constructs. Histograms represent the response of the cell to the addition of histamine

shown as the maximum fluorescence (Fmax) after treatment / mean fluorescence

before histamine F0.

Ratiometric Ca^{2+} imaging using a mt-Pericam construct

HeLa cells were seeded on glass bottom ibidi chambers (µ-slide 2 wells) 3 days

before imaging. Knockdown of the proteins of interest was realized 2 days before

imaging, as described above. The day after knockdown, 2.4 µg of mt-Pericam

plasmid were transfected with 6 µl of lipofectamine 2000/well (Invitrogen), according

to the manufacturer's instructions. The day of imaging, cells were washed 4-5 hours

before the imaging with fresh DMEM.

Imaging of mt-pericam was performed on an inverted Nikon Ti Eclipse Eclipse-E

microscope coupled with a Spinning Disk (Yokogawa, CSU-X1-A1), a 40x CFI Plan

Fluor objective (Nikon, NA 1.30, oil immersion) and a sCMOS camera (Photometrics,

Prime 95B) with a binning of 2. Knock-down cells expressing pericam constructs

were imaged in a controlled atmosphere (37°C, 5% CO_2). For each field of view, a

time-lapse was recorded with an image every 8 sec during 10 min. Approximatively 3

min after the start of captures, histamine was added to a final concentration of 100

µM. For each time point mt-pericam fluorescence was collected using a 525/45

bandpass filter (Semrock) after alternative excitation at 405 nm (Vortran, 100mW

laser) and 488 nm (Vortran, 150mW laser). A perfect focus system (Nikon) was used

to keep the focus constant along the timelapse, the whole system was driven by

Metamorph (version 7.7, Molecular Devices).

All the image processing and quantification was performed using a custom script in

Image J. First, images were corrected for flat-field and dark noise using reference

images respectively acquired on a fluorescent plastic slide (Chroma) and with no light

reaching the detector. Regions of interest were also drawn on an empty area of every

stack to perform a background subtraction on every image. Then hyperstacks

containing two channels were registered to correct any lateral drift of the sample

during the acquisition using the Image J plugin "HyperStackReg (v05)" with the

"translation" algorithm. Once the stacks were processed, regions of interest were

drawn around individual cells, a threshold applied (Mean) and the mean intensity of

each cell recorded on each image pair along the time-lapse. For every cell the ratio

488/405 was plotted along the time and only the cells with a stable ratio during the

first 3 min time points and with a sharp increase observed after addition of histamine

were kept for the final analysis.

1003

Flow cytometry for TMRE analysis

Membrane potential studies using TMRE- Mitochondrial Membrane Potential

Assay Kit

HeLa cells were seeded on 6-well plates 3 days before TMRE detection and the day

after they were transfected for 48 hours with the non-targeting, ORP5 or ORP8-

specific siRNAs specified above. The day before analysis, the medium was replaced

with DMEM 10% FBS with or without 20 µM CCCP (Carbonyl cyanide m-

chlorophenyl hydrazine) at 37°C and 5% CO_2 to induce mitochondrial depolarization.

After 24 hours, 200 nM of TMRE in DMEM 10% FBS (Mitochondrial Membrane

Potential Assay Kit, abcam ab113852) was added to cells and replaced after 30 min.

Cells were kept at 37°C and 5% CO_2 for additional 15 min and then trypsinized and

resuspended in DMEM (1-2x105 cells in 500 µl). TMRE was then analysed by flow

cytometry.

Flow cytometry

Multicolor flow cytometry was conducted on Beckman Coulter's CytoFLEX S

cytometer. Analyses were performed using Beckman Coulter's Kaluza and CytExpert

software. TMRE (Ex/Em = 549/575) was excited by a 561 nm laser and detected

through a 585/542 nm filter channel. All acquisitions were made on 10 000 cells. All

data are presented as mean ±SEM of three independent experimental replicates.

1023

Apoptosis assay using Cleaved-caspase-3 expression

HeLa cells were seeded on glass bottom 13 mm coverslips in a 24 well plates. The

next day, 48 hours knockdown of the proteins of interest were realized as described

above. 2 days after the knockdown, cells were treated with 1 μM staurosporine or

with DMSO to induce apoptosis. After 4 hours apoptotic cells were analyzed by

confocal immunofluorescence, using cleaved Caspase-3 antibody and Mitotracker

red to label mitochondria. Cleaved Caspase-3 positive cells were quantified using the

Cell Counter plugin from Image J Software. All data are presented as mean ±SEM of

three experimental replicates.

1033

Senescence studies using SA-β-Galactosidase assay and crystal violet

staining

For SA-β-galactosidase assay, cells were washed twice with PBS and fixed 5 min

with 2% formaldehyde / 0.2% glutaraldehyde. Cells were then rinsed twice in PBS

and incubated at 37°C overnight in SA-β-Gal staining solution as previously

described (Itahana, Campisi et al., 2007).

For crystal violet staining, cells were washed with PBS, fixed for 15 min in 3.7%

formaldehyde and then stained with 0.5% crystal violet solution. After drying, plates

were scanned.

Relative cell count was determined by dissolving crystal violet in acetic acid solution

and measuring optical density at 595 nM.

1045

Electron Microscopy Analysis

Conventional EM

For conventional EM, cells grown on 13 mm glass bottom coverslips (Agar Scientific) were fixed with 2.5% glutaraldehyde and 2% PFA in 0.1 M cacodylate, 0.05% $CaCl_2$ buffer for 24 hours. After several washes with 0.1 M cacodylate buffer, the cells were postfixed with 1% OsO_4, 1.5% potassium ferricyanide in 0.1M Cacodylate for 1 hour. After several washes with 0.1 M cacodylate buffer and H_2O, the cells were stained with 0.5% uranyl acetate for 24 hours. After several washes with H_2O, the cells were dehydrated in ethanol and embedded in Epon while on the coverslips. Ultrathin sections were prepared, counterstained with uranyl acetate and observed under a MET JEOL 1400 equipped with a Orius High speed (Gatan) camera.

HRP Detection

HeLa cells expressing HRP-KDEL were fixed on coverslips with 1.3% glutaraldehyde in 0.1 M cacodylate buffer, washed in 0.1 M ammonium phosphate [pH 7.4] buffer for 1 hour and HRP was visualized with 0.5 mg/ml DAB and 0.005% H_2O_2 in 0.1 M Ammonium Phosphate [pH 7.4] buffer. Development of HRP (DAB dark reaction product) took between 5 min to 20 min and was stopped by extensive washes with cold water. Cells were postfixed in 2% OsO_4+1% $K_3Fe(CN)_6$ in 0.1 M cacodylate buffer at 4°C for 1 hour, washed in cold water and then contrasted in 0.5% uranyl acetate for 2 hours at 4°C, dehydrated in an ethanol series and embedded in epon as for conventional EM. Ultrathin sections were counterstained with 2% uranyl acetate and observed under a FEI Tecnai 12 microscope equipped with a CCD (SiS 1kx1k keenView) camera.

Immunogold labelling

HeLa cells were fixed with a mixture of 2%PFA and 0.125% glutaraldehyde in 0.1 M

phosphate buffer [pH 7.4] for 2 hours, and processed for ultracryomicrotomy as

described previously (Slot & Geuze, 2007). Ultrathin cryosections were single- or

double-immunogold-labeled with antibodies and protein A coupled to 10 or 15 nm

gold (CMC, UMC Utrecht, The Netherlands), as indicated in the legends to the

figures. Immunogold-labeled cryosections were observed under a FEI Tecnai 12

microscope equipped with a CCD (SiS 1kx1k keenView) camera.

Quantifications

For the quantification of the number of cristae junction in Epon sections, about 200

mitochondria were analyzed in randomly selected cell profiles and cristae junctions

were counted in each of the mitochondria profile and reported as number of

cristae/mitochondria profile. All data are presented as mean ±SEM of three

experimental replicates.

For the quantification of ER-mitochondria contact sites in HRP-stained Epon

sections, the total circumference of each mitochondria and the length of the multiple

HRP-positive ER segments closely associated (<30 nm) with them were measured

by iTEM software (Olympus) on acquired micrographs of HeLa cells for each of 20-

30 cell profiles, as indicated in the figure legends. Cells were randomly selected for

analysis without prior knowledge of transfected plasmid or siRNA. All data are

presented as mean (%) ±SEM of three experimental replicates.

For the quantification of immunogold labeling on ultrathin cryosections, 150 gold

particles were counted on acquired micrographs of randomly selected cell profiles at

specific ranges of distance from CJ (0-50, 50-100, 100-150, 150-200 nm) in each of

three experiments. All data are presented as mean (%) ±SEM of three experimental

replicates.

ACKNOWLEDGEMENTS

We thank Dr V. Kozjak-Pavlovic and P. Pinton for kindly sharing reagents and advices with us. We also thank Dr. R. Legouis for discussion; Claire Boulogne, Mickael Bourge, and Remi Le Borgne for help and consultation with microscopy experiments; Blandine Bourigault, Riikka Kosonen and Liisa Arala for technical assistance. The present work has benefited from Imagerie-Gif core facility supported by l' Agence Nationale de la Recherche (ANR-11-EQPX-0029/Morphoscope, ANR-10-INBS-04/FranceBioImaging; ANR-11-IDEX-0003-02/ Saclay Plant Sciences). For the Immuno-EM and HRP-KDEL EM analysis/quantifications we acknowledge the ImagoSeine core facility of the Institut Jacques Monod, member of IBiSA and the France-BioImaging (ANR-10-INBS-04) infrastructure. We also thank Jean-Michel Camadro and Thibault Leger (IJM proteomics platform) for proteomics analysis. This work was supported by ANR Jeune Chercheur (ANR0015TD and ATIP-Avenir Program to F.G.; the Academy of Finland (grants 285223 and 322647), the Sigrid Juselius Foundation, the Finnish Foundation for Cardiovascular Research and the Magnus Ehrnrooth Foundation to V.M.O.; the Fondation ARC pour la recherche sur le cancer and INCA (N°2018-144) to D.B.; the "Association Française contre les Myopathies" (AFM Research grant 20123) and the ITMO-Inserm Plan Cancer 2014-2019 to D.T.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Chung J, Torta F, Masai K, Lucast L, Czapla H, Tanner LB, Narayanaswamy P, Wenk MR, Nakatsu F, De Camilli P (2015) INTRACELLULAR TRANSPORT. PI4P/phosphatidylserine countertransport at ORP5- and ORP8-mediated ER-plasma membrane contacts. Science 349: 428-32

Colombini M (2012) VDAC structure, selectivity, and dynamics. Biochimica et biophysica acta 1818: 1457-65

De Stefani D, Bononi A, Romagnoli A, Messina A, De Pinto V, Pinton P, Rizzuto R (2012) VDAC1 selectively transfers apoptotic Ca2+ signals to mitochondria. Cell Death Differ 19: 267-73

Ding C, Wu Z, Huang L, Wang Y, Xue J, Chen S, Deng Z, Wang L, Song Z, Chen S (2015) Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae structure. Sci Rep 5: 16064

Du X, Turner N, Yang H (2017) The role of oxysterol-binding protein and its related proteins in cancer. Semin Cell Dev Biol

Du X, Zadoorian A, Lukmantara IE, Qi Y, Brown AJ, Yang H (2018) Oxysterol-binding protein-related protein 5 (ORP5) promotes cell proliferation by activation of mTORC1 signaling. The Journal of biological chemistry 293: 3806-3818

Filippin L, Abad MC, Gastaldello S, Magalhaes PJ, Sandona D, Pozzan T (2005) Improved strategies for the delivery of GFP-based Ca2+ sensors into the mitochondrial matrix. Cell calcium 37: 129-36

Friedman JR, Mourier A, Yamada J, McCaffery JM, Nunnari J (2015) MICOS coordinates with respiratory complexes and lipids to establish mitochondrial inner membrane architecture. eLife 4

Galmes R, Houcine A, van Vliet AR, Agostinis P, Jackson CL, Giordano F (2016) ORP5/ORP8 localize to endoplasmic reticulum-mitochondria contacts and are involved in mitochondrial function. EMBO reports 17: 800-10

Ghai R, Du X, Wang H, Dong J, Ferguson C, Brown AJ, Parton RG, Wu JW, Yang H (2017) ORP5 and ORP8 bind phosphatidylinositol-4, 5-biphosphate (PtdIns(4,5)P 2) and regulate its level at the plasma membrane. Nature communications 8: 757

Giordano F (2018) Non-vesicular lipid trafficking at the endoplasmic reticulum-mitochondria interface. Biochemical Society transactions 46: 437-452

Guarani V, McNeill EM, Paulo JA, Huttlin EL, Frohlich F, Gygi SP, Van Vactor D, Harper JW (2015) QIL1 is a novel mitochondrial protein required for MICOS complex stability and cristae morphology. eLife 4

Harner M, Korner C, Walther D, Mokranjac D, Kaesmacher J, Welsch U, Griffith J, Mann M, Reggiori F, Neupert W (2011) The mitochondrial contact site complex, a determinant of mitochondrial architecture. The EMBO journal 30: 4356-70

Herrera-Cruz MS, Simmen T (2017) Of yeast, mice and men: MAMs come in two flavors. Biol Direct 12: 3

Hohr AIC, Lindau C, Wirth C, Qiu J, Stroud DA, Kutik S, Guiard B, Hunte C, Becker T, Pfanner N, Wiedemann N (2018) Membrane protein insertion through a mitochondrial beta-barrel gate. Science 359

Huynen MA, Muhlmeister M, Gotthardt K, Guerrero-Castillo S, Brandt U (2016) Evolution and structural organization of the mitochondrial contact site (MICOS) complex and the mitochondrial intermembrane space bridging (MIB) complex. Biochimica et biophysica acta 1863: 91-101

Itahana K, Campisi J, Dimri GP (2007) Methods to detect biomarkers of cellular senescence: the senescence-associated beta-galactosidase assay. Methods Mol Biol 371: 21-31

John GB, Shang Y, Li L, Renken C, Mannella CA, Selker JM, Rangell L, Bennett MJ, Zha J (2005) The mitochondrial inner membrane protein mitofilin controls cristae morphology. Molecular biology of the cell 16: 1543-54

Joshi AS, Thompson MN, Fei N, Huttemann M, Greenberg ML (2012) Cardiolipin and mitochondrial phosphatidylethanolamine have overlapping functions in mitochondrial fusion in Saccharomyces cerevisiae. The Journal of biological chemistry 287: 17589-97

Kim SH, Song HE, Kim SJ, Woo DC, Chang S, Choi WG, Kim MJ, Back SH, Yoo HJ (2017) Quantitative structural characterization of phosphatidylinositol phosphates from biological samples. J Lipid Res 58: 469-478

Kopec KO, Alva V, Lupas AN (2010) Homology of SMP domains to the TULIP superfamily of lipid-binding proteins provides a structural basis for lipid exchange between ER and mitochondria. Bioinformatics 26: 1927-31

Kornmann B, Currie E, Collins SR, Schuldiner M, Nunnari J, Weissman JS, Walter P (2009) An ER-mitochondria tethering complex revealed by a synthetic biology screen. Science 325: 477-81

Kozjak V, Wiedemann N, Milenkovic D, Lohaus C, Meyer HE, Guiard B, Meisinger C, Pfanner N (2003) An essential role of Sam50 in the protein sorting and assembly machinery of the mitochondrial outer membrane. The Journal of biological chemistry 278: 48520-3

Kozjak-Pavlovic V, Ross K, Benlasfer N, Kimmig S, Karlas A, Rudel T (2007) Conserved roles of Sam50 and metaxins in VDAC biogenesis. EMBO reports 8: 576-82

Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, Wheeler H, Reinisch KM, De Camilli P (2018) VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact sites. The Journal of cell biology 217: 3625-3639

Lang A, John Peter AT, Kornmann B (2015) ER-mitochondria contact sites in yeast: beyond the myths of ERMES. Current opinion in cell biology 35: 7-12

Liebisch G, Lieser B, Rathenberg J, Drobnik W, Schmitz G (2004) High-throughput quantification of phosphatidylcholine and sphingomyelin by electrospray ionization tandem mass spectrometry coupled with isotope correction algorithm. Biochimica et biophysica acta 1686: 108-17

Ma X, Warnier M, Raynard C, Ferrand M, Kirsh O, Defossez PA, Martin N, Bernard D (2018) The nuclear receptor RXRA controls cellular senescence by regulating calcium signaling. Aging Cell 17: e12831

Maeda K, Anand K, Chiapparino A, Kumar A, Poletto M, Kaksonen M, Gavin AC (2013) Interactome map uncovers phosphatidylserine transport by oxysterol-binding proteins. Nature 501: 257-61

Martin N, Bernard D (2018) Calcium signaling and cellular senescence. Cell calcium 70: 16-23

Moser von Filseck J, Copic A, Delfosse V, Vanni S, Jackson CL, Bourguet W, Drin G (2015) INTRACELLULAR TRANSPORT. Phosphatidylserine transport by ORP/Osh proteins is driven by phosphatidylinositol 4-phosphate. Science 349: 432-6

Nagai T, Sawano A, Park ES, Miyawaki A (2001) Circularly permuted green fluorescent proteins engineered to sense Ca2+. Proceedings of the National Academy of Sciences of the United States of America 98: 3197-202

Ott C, Dorsch E, Fraunholz M, Straub S, Kozjak-Pavlovic V (2015) Detailed analysis of the
human mitochondrial contact site complex indicate a hierarchy of subunits. PloS one 10:
e0120213
Patron M, Raffaello A, Granatiero V, Tosatto A, Merli G, De Stefani D, Wright L,
Pallafacchina G, Terrin A, Mammucari C, Rizzuto R (2013) The mitochondrial calcium
uniporter (MCU): molecular identity and physiological roles. The Journal of biological
chemistry 288: 10750-8
Pulli I, Lassila T, Pan G, Yan D, Olkkonen VM, Tornquist K (2018) Oxysterol-binding
protein related-proteins (ORPs) 5 and 8 regulate calcium signaling at specific cell
compartments. Cell calcium 72: 62-69
Rapizzi E, Pinton P, Szabadkai G, Wieckowski MR, Vandecasteele G, Baird G, Tuft RA,
Fogarty KE, Rizzuto R (2002) Recombinant expression of the voltage-dependent anion
channel enhances the transfer of Ca2+ microdomains to mitochondria. The Journal of
cell biology 159: 613-24
Reinisch KM, De Camilli P (2016) SMP-domain proteins at membrane contact sites:
Structure and function. Biochimica et biophysica acta 1861: 924-927
Rizzuto R, Marchi S, Bonora M, Aguiari P, Bononi A, De Stefani D, Giorgi C, Leo S, Rimessi
A, Siviero R, Zecchini E, Pinton P (2009) Ca(2+) transfer from the ER to mitochondria:
when, how and why. Biochimica et biophysica acta 1787: 1342-51
Schikorski T, Young SM, Jr., Hu Y (2007) Horseradish peroxidase cDNA as a marker for
electron microscopy in neurons. Journal of neuroscience methods 165: 210-5
Scorrano L, De Matteis MA, Emr S, Giordano F, Hajnoczky G, Kornmann B, Lackner LL,
Levine TP, Pellegrini L, Reinisch K, Rizzuto R, Simmen T, Stenmark H, Ungermann C,
Schuldiner M (2019) Coming together to define membrane contact sites. Nature
communications 10: 1287
Shiao YJ, Lupo G, Vance JE (1995) Evidence that phosphatidylserine is imported into
mitochondria via a mitochondria-associated membrane and that the majority of
mitochondrial phosphatidylethanolamine is derived from decarboxylation of
phosphatidylserine. The Journal of biological chemistry 270: 11190-8
Slot JW, Geuze HJ (2007) Cryosectioning and immunolabeling. Nature protocols 2: 2480-
91
Steenbergen R, Nanowski TS, Beigneux A, Kulinski A, Young SG, Vance JE (2005)
Disruption of the phosphatidylserine decarboxylase gene in mice causes embryonic
lethality and mitochondrial defects. The Journal of biological chemistry 280: 40032-40
Tamura Y, Endo T (2017) Role of Intra- and Inter-mitochondrial Membrane Contact
Sites in Yeast Phospholipid Biogenesis. Advances in experimental medicine and biology
997: 121-133
Tan W, Colombini M (2007) VDAC closure increases calcium ion flux. Biochimica et
biophysica acta 1768: 2510-5
Tatsuta T, Scharwey M, Langer T (2014) Mitochondrial lipid trafficking. Trends in cell
biology 24: 44-52
Vance JE (1990) Phospholipid synthesis in a membrane fraction associated with
mitochondria. The Journal of biological chemistry 265: 7248-56
Vance JE (2014) MAM (mitochondria-associated membranes) in mammalian cells: lipids
and beyond. Biochimica et biophysica acta 1841: 595-609
Vance JE, Tasseva G (2013) Formation and function of phosphatidylserine and
phosphatidylethanolamine in mammalian cells. Biochimica et biophysica acta 1831:
543-54

Wang H, Ma Q, Qi Y, Dong J, Du X, Rae J, Wang J, Wu WF, Brown AJ, Parton RG, Wu JW,
Yang H (2019) ORP2 Delivers Cholesterol to the Plasma Membrane in Exchange for
Phosphatidylinositol 4, 5-Bisphosphate (PI(4,5)P2). Mol Cell 73: 458-473 e7
Wideman JG (2015) The ubiquitous and ancient ER membrane protein complex (EMC):
tether or not? F1000Res 4: 624
Wiel C, Lallet-Daher H, Gitenay D, Gras B, Le Calve B, Augert A, Ferrand M, Prevarskaya
N, Simonnet H, Vindrieux D, Bernard D (2014) Endoplasmic reticulum calcium release
through ITPR2 channels leads to mitochondrial calcium accumulation and senescence.
Nature communications 5: 3792
Wollweber F, von der Malsburg K, van der Laan M (2017) Mitochondrial contact site and
cristae organizing system: A central player in membrane shaping and crosstalk.
Biochimica et biophysica acta 1864: 1481-1489
1276

a

Streptavidin-coated bead

Donor liposomes
Biotinylated,
TopFluor-labeled

Acceptor
liposomes

Fluorescent
lipid transfert
+/- ORP5/8 ORD

b

c

Lipidomic analysis of PE content:

d

Radiometric measurement of
newly synthetised PE:

e

HRP-KDEL

siCtrl

siORP5 + siORP8

f

ER-mitochondria contacts:

g

■ siCtrl
■ siORP5 + siORP8

normal   altered

h

mito

PS

PE   PS

ORP5/8   ER

Figure 1

Figure 1. ORP5 and ORP8 mediate PS transfer *in vitro* and at ER-mitochondria contact sites

(a) Schematic cartoon showing the *in vitro* assay used to study ORP5/8 ORD-mediated lipid transport between liposomes. (b) Donor liposomes containing fluorescent lipids (97 mol% POPC, 1 mol% TopFluor-PS, -PC or -PE and 2 mol% of biotinylated-PE) and pre-bound to streptavidin beads were mixed at a 1:1 molar ratio with acceptor liposomes (100 mol% POPC) in the presence or absence of ORP5 or ORP8 ORD domains (250 µM of donor and acceptor liposomes and 0,3 µM of proteins in the reaction solution). The increase in fluorescence in the acceptor liposomes, which remain unbound in the solution, was measured after 1 hour of incubation at 37° C. Data are presented as % of transferred lipid $\pm$SEM and are the mean of six independent replicates. **P<0,001 (c) Mass spectrometry (MS)-based quantification of the PE content (nmol/mg protein) of mitochondria isolated from Ctrl, ORP5 or ORP8 knockdown cells and of Ctrl, ORP5 or ORP8 knockdown intact cells. Data are shown as mean of three independent replicates $\pm$SEM. *P<0,05. (d) Ctrl, ORP5, ORP8 or ORP5+ORP8 double-knockdown cells were incubated with L-[^{3}H(G)]serine (30.9 Ci/mmol) for 18 hour. After extraction and separation of lipids by TLC, PS and PE spots were scraped and analysed for [^{3}H] radioactivity, as described under "Methods". Each condition was analyzed in triplicate in each of three independent biological replicates. Data are presented as mean of PE:PS ratio $\pm$SEM. *P<0,05, **P<0,01 compared to Ctrl. (e) Electron micrographs of HRP-KDEL-expressing HeLa cells treated with Ctrl siRNAs or siRNAs against ORP5 and ORP8 (siORP5+siORP8). Red arrows indicate ER-mitochondria contact sites. Scale bar, 500 nm. (f) Quantifications of the extent of ER-mitochondria contact sites in Ctrl, ORP5, ORP8 and ORP5+ORP8 knockdown cells expressing HRP-KDEL. Data are shown as % of the ER in contact with mitochondria (mitochondria occupancy) $\pm$SEM, n = 20 cell profiles and $\pm$900 mitochondria; n.s; not significant. (g) Quantifications of the number of mitochondria with aberrant cristae morphology in the indicated siRNA conditions. Data are shown as % of mitochondria $\pm$SEM, n = 20 cell profiles and $\pm$700 mitochondria. **P<0,01 compared to siCtrl. (h) Schematic representation of non-vesicular PS transfer mediated by ORP5/8 at ER-mitochondria contact sites. PS is transported to mitochondrial membranes where it is rapidly converted into mitochondrial PE.

Figure 2

Figure 2. ORP5 and ORP8 interact with the MIB/MICOS complex at ER-mitochondria contacts near CJ

(a) Electron micrograph of ultrathin cryosection of HeLa cells transfected with HA-ORP5 or Mitofilin-EGFP and immunogold stained with anti-HA or anti-GFP (10 or 15 nm gold) showing HA-ORP5 localization at ER-mitochondria contacts in close proximity to CJ (arrow) and the localization of the MICOS complex (Mitofilin) at CJ (arrows). Scale bar 150 nm. (b) Quantification of the proximity of HA-ORP5 gold particles to the CJ. Results are presented as the percentage of ORP5 gold particles (n = 23 cell profiles) at specific ranges of distance from CJ (0-50, 50-100, 100-150, 150-200 nm). (c) EGFP-ORP5, EGFP-ORP8 or EGFP alone were transfected in HeLa cells then immuno-precipitated from lysates and analyzed by western blot using antibodies against GFP (for ORP5 or ORP8), Mitofilin or SAM50. (d) EGFP-ORP5, EGFP-ORP5ΔPH, EGFP-ORP5ΔORD, EGFP-ORP5ΔTM, EGFP-ORP8, EGFP-ORP8ΔPH or EGFP alone were transfected in HeLa cells then immuno-precipitated from lysates and analyzed by western blot using antibodies against GFP (for ORP5 or ORP8) or against SAM50. (e) Fluorescent images of HeLa cells transfected with either EGFP-ORP5 or EGFP-ORP5ΔPH and stained with MitoTracker and anti-SAM50 antibody. Insets show magnifications of the boxed regions. Scale bar, 10 μm.

a

siCtrl
siMitofilin
siMitofilin + siSAM50
siSAM50
siORP5
siMitofilin + siORP5

ORP5

SAM50

Mitofilin

Actin

b

Radiometric measurement of newly synthetised PE

PE:PS ratio

1,2
1,1
1
0,9
0,8

siCtrl
siORP5 *
siMitofilin *
siSAM50
siORP5+siMitofilin **
siMitofilin+siSAM50 **
siORP5+siSAM50 **

c

siCtrl

siMitofilin

siSAM50

d

ER-mitochondria contacts

Mitochondria occupancy (%)

n.s.

siCtrl   siMitofilin   siSAM50

Figure 3

Figure 3. ORP5 and the MIB/MICOS complex function together to mediate non-vesicular transport of PS from the ER to mitochondrial membranes

(a) Western analysis showing ORP5, SAM50, Mitofilin and Actin levels in protein lysates from HeLa cells treated with siRNA against Ctrl, ORP5, Mitofilin or SAM50. Arrow indicates the specific band for Mitofilin. (b) Radiometric measurement of PS-to-PE conversion in the indicated siRNAs. Data are presented as mean of PE:PS ratio $\pm$SEM. Each condition was performed in triplicate in each of the independent biological replicates (n = 5 for siCtrl and siORP5; n = 4 for siSAM50; n = 3 for the other siRNAs conditions). *P<0,05, **P<0,01 compared to Ctrl. (c) Representative electron micrographs of HeLa cells treated with Ctrl siRNAs or siRNAs against Mitofilin or SAM50 and transfected with HRP-KDEL. Red arrows indicate ER-mitochondria contact sites. Scale bar, 500 nm. (d) Quantifications of the extent of ER-mitochondria contact sites in Ctrl, Mitofilin and SAM50 knockdown cells expressing HRP-KDEL. Data are shown as % of the ER in contact with mitochondria (mitochondria occupancy) $\pm$SEM, n = 30 for siCtrl, n = 20 cell profiles for siMitofilin and siSAM50 and 1 000 mitochondria; n.s; not significant.

Figure 4

Figure 4. Silencing of ORP5 but not ORP8 specifically increases VDAC protein level in a SAM50-dependent manner

(a) Western analysis showing ORP5, ORP8, SAM50, Mitofilin, VDAC and Actin levels in protein lysates from HeLa cells treated with either Ctrl siRNAs, or with siRNAs against ORP5, ORP8, or SAM50. (b) Quantification of ORP5, SAM50, VDAC and Mitofilin protein expression in the indicated knockdown conditions. Normalized data are shown as mean $\pm$ SEM (n = 3). (c) Confocal images of Ctrl, ORP5 and ORP8 knockdown HeLa cells immunostained using anti-SAM50 antibody. Scale bar, 10 μm. (d) Quantitative RT-PCR analysis of ORP5, SAM50, Mitofilin, VDAC1, VDAC2 and VDAC3 in ORP5 (left graph) or in SAM50 (right graph) knockdown cells versus control HeLa cells. y axis: $2^{(-\Delta\Delta Ct)}$ value represents differences between the mean Ct (Cycle threshold) values of tested genes and those of reference gene (SDHA).

Figure 5

Figure 5. Silencing of ORP5 but not ORP8 increases mitochondrial calcium

(a) Pseudocolor 488/405 ratio time-lapse images of a representative Ctrl, ORP5 or ORP8 knockdown HeLa cell expressing mt-Pericam before and after histamine treatment. Scale bar 10 µm. (b) Representative curves of Ctrl, ORP5 and ORP8 knockdown HeLa cells expressing mt-Pericam showing the ratio of Pericam fluorescence before and after histamine treatment (+His). The change in the ratio of Pericam fluorescence after histamine treatment under each condition was compared (Rmax/R0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, $p< 0.001$; n.s; not significant. (c) Representative curves of Ctrl and ORP5 knockdown HeLa cells expressing the GECO construct Mito-LAR showing the fluorescence before and after histamine treatment. The change in fluorescence after histamine treatment under each condition was compared (Fmax/F0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, $p< 0.001$; n.s; not significant. (d) Representative curves of Ctrl and ORP5 knockdown HeLa cells expressing the GECO construct ER-LAR showing the fluorescence before and after histamine treatment. The change in fluorescence after histamine treatment under each condition was compared (Fmax/F0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, n.s; not significant. (e) Representative curves of Ctrl, ORP5 and ORP8 knockdown HeLa cells expressing the GECO construct R-GECO showing the fluorescence before and after histamine treatment. The change in fluorescence after histamine treatment under each condition was compared (Fmax/F0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, $p< 0.001$ and ****, $p< 0.0001$; n.s; not significant.

Figure 6

Figure 6. Calcium increase induced by ORP5 knockdown is dependent on VDAC

(a) Western analysis showing ORP5 and VDAC protein levels in Ctrl, ORP5 and VDAC siRNA-treated HeLa cells. (b). Representative curves of Ctrl, ORP5, VDAC and ORP5+VDAC knockdown HeLa cells expressing mt-Pericam showing the ratio of Pericam fluorescence before and after histamine treatment (+His). The change in the ratio of Pericam fluorescence after histamine treatment under each condition was compared (R_{max}/R_0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, p< 0.001; n.s; not significant. (c). Representative curves of Ctrl, ORP5 and ORP5+SAM50 knockdown HeLa cells expressing mt-Pericam showing the ratio of Pericam fluorescence before and after histamine treatment (+His). The change in the ratio of Pericam fluorescence after histamine treatment under each condition was compared (R_{max}/R_0). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, p< 0.001; n.s; not significant.

Figure 7

Figure 7. ORP5 knockdown promotes cell senescence

(a) MCR5 cells transfected with control siRNA or with siRNA against ORP5 were fixed and then stained with crystal violet solution (CA) or with SA-β-galactosidase (SA-β-Gal) staining solution. The graph on the top right shows the relative cell number determined by crystal violet staining. The graph on the bottom right shows the relative percentage of β-gal positive cells determined by SA-β-galactosidase assay. Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, **, p< 0.01; *, p< 0.05. (b) Quantitative RT-PCR analysis of ORP5, p16, p21 and Ki67 in ORP5 knockdown cells and control HeLa cells. y axis:$2^{(-\Delta\Delta Ct)}$ value represents differences between the mean Ct (Cycle threshold) values of tested genes and those of reference genes (PGK1 and HPRT1). Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, **, p< 0.01; *, p< 0.05. (c) MCR5 cells transfected with control siRNA or with siRNA against ORP5, SAM50 or ORP8 were fixed and then stained with crystal violet solution or with SA-β-Gal staining solution. The graph on the left shows the relative cell number determined by crystal violet staining. The graph on the right shows the relative percentage of β-gal positive cells determined by SA-β-galactosidase assay. Data are presented as normalized mean $\pm$ SEM, n = 3 per condition, ***, p< 0.001; *, p< 0.05; n.s; not significant.

a

Model:

Figure 8. Lipid and calcium transport occur at the same ER-mitochondria contact site microdomains and are regulated by a ORP5-SAM50-VDAC pathway

ORP5/8 are directly involved in the transfer of PS from ER to mitochondria at ER-mitochondria membrane contact sites. This transfer occurs at the level of cristae junctions where ORP5/8 localize and interacts with the proteins of the MIB complex SAM50 and Mitofilin. This interaction facilitates the transfer of PS from ER to IMM at the level of the CJ where the PS is then converted into PE that play a critical role in cristae organization and mitochondrial function. ORP5 is also involved in calcium homeostasis by regulating the level of VDAC *via* SAM50 interaction and regulation.

Figure 8

Figure S1. (a) Coomassie stained SDS-PAGE of the recombinant ORD domain of ORP5/8 proteins purified in BL21DE3 RILP cells. (b) Crude mitochondria, mitochondria, and MAM fractions were purified from Ctrl, ORP5 and ORP8 siRNA-treated HeLa cells. Equal amounts of protein from each fraction were loaded on a 4–20% gradient SDS–PAGE gel and immunoblotted using anti-ORP5, anti-ORP8, anti-IP3R-3 (MAM protein), and anti-cytochrome c (mitochondrial protein). Mito, mitochondria; MAM, mitochondria-associated ER membrane. (c) WB analysis showing ORP5, ORP8 and Actin levels in protein lysates from HeLa cells treated with either Ctrl siRNAs or with siRNAs against ORP5 or/and ORP8.

Figure S1

a

		IP EGFP	IP EGFP-ORP5	IP EGFP-ORP5ΔPH
Protein	**Localization**		**Mascot Score**	
ORP5	MAMs	0	9689	12192
ORP8	MAMs	31	930	690
Mitofilin	IMM	43	579	1150
SAM50	OMM	0	237	516
PSS1	MAMs	47	175	167
TOM22	OMM	0	77	95
PTPIP51	OMM	0	76	73
Metaxin2	OMM	0	32	175

b

c

Figure S2. (a) Identification of proteins that interact with EGFP-tagged ORP5 constructs by mass spectrometry. Proteins associated to MAMs and mitochondrial outer and inner membranes (OMM, IMM) are interacting partners of EGFP-ORP5 constructs. Note the presence of some proteins of the MIB complex: Mitofilin, SAM50 and Metaxin2. The interaction of Mitofilin, SAM50 and Metaxin2 with the EGFP-ORP5ΔPH construct is stronger than with EGFP-ORP5. (b) WB analysis showing GFP (EGFP-tagged constructs), Mitofilin, VDAC and SAM50 and Actin levels in protein lysates from HeLa cells transfected with either EGF construct (Control) or with EGFP-ORP5 or EGFP-ORP5ΔPH constructs. (c) WB analysis showing PDI (ER), VAPB (ER), CV-ATP5A (IMM), Cytochrome C (Cyt C, IMM) and Actin levels in protein lysates from HeLa cells treated with either Ctrl siRNAs or with siRNAs against ORP5, ORP8, or SAM50.

Figure S2

Figure S3. (a) Representative EM micrographs showing the morphology of mitochondria in HeLa cells treated with siRNA against ORP5, ORP8, Mitofilin and SAM50. Scale bar, 200 nm. Red arrows indicate ER elements in contact with mitochondria; arrowheads indicate CJ; m, mitochondria. (b) Quantifications of the number of CJ per mitochondria profile in the indicated siRNA conditions. % of CJ ±SEM, n = 170-260 mitochondria, ***, p< 0.001; **, p< 0.01.

Figure S3

Figure S4

Figure S4. (a) Representative graphs of flow cytometry analysis of HeLa cells transfected with siRNA against ORP5 or ORP8 (n = 10 000 cells) and loaded with the mitochondrial membrane potential sensitive dye TMRE, in presence (right graph) or absence (left graph) of CCCP. Flow cytometry histograms are showing the relative TMRE fluorescence intensity; the average of three independent replicates is shown. Data are presented as normalized mean $\pm$ SEM. (b) Fluorescent images of HeLa cells transfected with siRNA against ORP5, ORP8 or control, treated with DMSO or staurosporine and stained with anti-cleaved Caspase-3 (apoptotic cells) and Mitotracker. Scale bar, 20 μm. The graphs represent the means $\pm$ SEM of the percentage of cleaved Caspase-3 positive cells in the different conditions; the average of three independent replicates is shown, n.s; not significant.