Unlocking the potential of historical abundance datasets to study

biomass change in flying insects

Rebecca S. Kinsella[1], Chris D. Thomas[1,2], Terry J. Crawford[1], Jane K. Hill[1,2], Peter J.

Mayhew[1] and Callum J. Macgregor[1,2,3]*

[1]: Department of Biology, University of York, Wentworth Way, York, YO10 5DD

[2]: Leverhulme Centre for Anthropocene Biodiversity, University of York, Wentworth Way,

York, YO10 5DD

[3]: Energy and Environment Institute, University of Hull, Cottingham Road, Hull, HU6 7RX

Running head: Predicting moth biomass from abundance

*Corresponding author:

Callum J. Macgregor; tel: (+44) 07886 149722; email: callumjmacgregor@gmail.com.

Abstract

1. Trends in insect abundance are well-established in some datasets, but far less is

known about how abundance measures translate into biomass trends. Moths

(Lepidoptera) provide particularly good opportunities to study trends and drivers of

biomass change at large spatial and temporal scales, given the existence of long-

term abundance datasets. However, data on the body masses of moths are required

for these analyses, but such data do not currently exist.

2. To address this data gap, we collected empirical data in 2018 on the forewing length

and dry mass of field-sampled moths, and used these to train and test a statistical

model that predicts the body mass of moth species from their forewing lengths (with

refined parameters for Crambidae, Erebidae, Geometridae and Noctuidae).

3. Modelled biomass was positively correlated, with high explanatory power, with

measured biomass of moth species (R^2 = 0.886 ± 0.0006, across 10,000

bootstrapped replicates) and of mixed-species samples of moths (R^2 = 0.873 ±

0.0003), showing that it is possible to predict biomass to an informative level of

accuracy, and prediction error was smaller with larger sample sizes.

4. Our model allows biomass to be estimated for historical moth abundance datasets,

and so our approach will create opportunities to investigate trends and drivers of

insect biomass change over long timescales and broad geographic regions.

Keywords

Biodiversity decline, body mass, forewing length, Lepidoptera, moths, predictive model

Background

Multiple recent studies have reported that insect biomass, abundance and diversity are in decline over recent decades (Conrad *et al.*, 2006; Hallmann *et al.*, 2017, 2019; Lister & Garcia, 2018; Harris *et al.*, 2019; Seibold *et al.*, 2019; van Strien *et al.*, 2019; Wepprich *et al.*, 2019; Bell *et al.*, 2020; Roth *et al.*, 2020; Salcido *et al.*, 2020), but with substantial spatial, temporal and taxonomic variation in the existence and strength of such declines (Shortall *et al.*, 2009; Macgregor *et al.*, 2019b; Outhwaite *et al.*, 2020). The reasons for such variation are not yet known, and it is therefore possible that declines in insect biomass are not always symptomatic of equivalent declines in abundance, or *vice versa*. Biomass could remain stable even in the face of declining abundance if communities became increasingly comprised of larger-bodied species (Pöyry *et al.*, 2017). Likewise, changes in community-level biomass could be attributable to changes in community composition, even in the absence of an overall abundance change. This might occur if insect communities became more biased towards larger- or smaller-bodied species, e.g. through size-bias in strength of selection for or against particular traits (Coulthard *et al.*, 2019), such as faster or slower life-histories, degree of habitat specialization (Mattila *et al.*, 2011; Davis *et al.*, 2013), or strength of attraction to artificial light at night (van Langevelde *et al.*, 2011). However, the dynamics of biomass and of biomass change, as well as the relationships between biomass, abundance, species richness, and community composition, remain poorly understood at large spatial and temporal scales because of a lack of suitable data on insect biomass (Macgregor *et al.*, 2019b).

Opportunities to investigate changes over time and space in insect communities are provided by several large-scale, long-term abundance datasets for moths (Lepidoptera) in the UK, including the Rothamsted Insect Survey (RIS; Storkey *et al.*, 2016), the National Moth Recording Scheme (NMRS; Fox *et al.*, 2011), and the Garden Moth Scheme (GMS; Bates *et al.*, 2014; Wilson *et al.*, 2018), and elsewhere (e.g. Groenendijk & Ellis, 2011;

Valtonen *et al.*, 2014, 2017). Two-thirds of British species of macro-moths show negative

abundance trends in the long-term (Conrad *et al.*, 2006), with similar patterns observed

elsewhere in Europe (e.g. Groenendijk & Ellis, 2011; Valtonen *et al.*, 2017). The potential

drivers of these declines are diverse (Fox, 2013), and likely to include habitat loss and

fragmentation, agricultural intensification and associated agrochemical use, increased

prevalence of artificial light at night and other factors associated with urbanisation, and

climate change (Wickramasinghe *et al.*, 2004; Morecroft *et al.*, 2009; Fox, 2013; Fox *et al.*,

2014; Gilburn *et al.*, 2015; van Langevelde *et al.*, 2018). Moths contribute to important

ecosystem functions, including nocturnal pollination (Macgregor *et al.*, 2015, 2019a) and

energy transfer from producers to higher-level consumers (e.g. Franklin *et al.*, 2003; Hooks

et al., 2003; Singer *et al.*, 2012). Thus, moths can be important to the conservation of their

predators, such as bats (Vaughan, 1997; Threlfall *et al.*, 2012) and some birds (Sierro *et al.*,

2001; Denerley *et al.*, 2018). In transferring energy, the quantity of vegetation consumed by

caterpillars and the biomass of insects available to predators may be functionally important

determinants of ecosystem processes (Brose *et al.*, 2005). Similarly, the body size of

individual species can play a substantial role in structuring networks of interspecific

interactions (Woodward *et al.*, 2005). All of these factors make moths a valuable taxon in

which to study long-term biomass change at the community level, but biomass data are

currently lacking for these analyses.

Existing long-term moth population and distribution datasets are potentially a very valuable

resource for understanding biomass changes, but these datasets record abundance, not

measurements of body mass or size, and in most cases do not retain specimens (preventing

biomass information from being obtained retrospectively). To address questions of biomass

change using these abundance datasets requires reliable body mass data for all species, but

such empirical data are currently available for only a limited set of species (García-Barros,

2015). An alternative approach is to use empirical data from a subset of species to model

the expected body mass of all species, using some other, more readily-available, trait. Such

models have previously been formulated to predict the body mass of moths and other

invertebrates from their body length (Sage, 1982; Sample *et al.*, 1993; Sabo *et al.*, 2002;

Höfer & Ott, 2009) and variants thereof (García-Barros, 2015), chosen because it is easily

measurable from museum specimens (García-Barros, 2015). However, for moths, body

length data are not widely available and in any case may be influenced to a greater degree

by contraction in dried specimens than other traits (García-Barros, 2015). The only

morphological trait for which existing data on many species are readily available is forewing

length: for example, an expected range of forewing lengths is included for all British species

of macro-moths, and most British species of micro-moths, in standard field guides (Sterling &

Parsons, 2012; Waring & Townsend, 2017), and it may therefore be possible to predict body

mass based on forewing length (Miller, 1997). The existence of substantial interfamilial

variation in body plan (e.g. between Saturniidae and Sphingidae; Janzen, 1984) may provide

opportunities to use taxonomy to fine-tune models, but no previous model has included any

refinement based on taxonomic relationships between moths.

In this study, we develop a statistical model to estimate the body mass of individual moths

from their forewing length, and hence quantify the biomass of samples of moths for which

species abundances only have been recorded. We have four aims: (i) collection of empirical

data (during 2018 on the University of York campus, UK) to establish the relationship

between forewing length and body mass in moths; (ii) construction of a predictive model for

estimating body mass from species identity and associated forewing length, (iii) testing the

accuracy of this model's predictions and how accuracy changes with increasing moth

abundance, and (iv) using existing data on forewing lengths to predict the body masses of all

British macro-moths, thus providing a resource to users of moth population data and to

comparative biologists.

Materials and methods

Field sampling, identification and measurement of moths

We sampled moths at three sites (Fig. S1) on the University of York campus, UK (53°56'41"

N 1°2'2" W; vice-county 61, South-east Yorkshire) between 11th June and 20th July 2018

(Appendix S1.1). Moths were collected using Heath-style moth traps (Heath, 1965), each

operating a 15 W actinic fluorescent tube and powered by a 12 V battery (Anglian

Lepidopterist Supplies, Hindolveston, UK). Moths were euthanized and returned to the

laboratory for identification and measurement. Moths were identified to species-level where

possible using standard field guides (Sterling & Parsons, 2012; Waring & Townsend, 2017).

Where species-level identification would have required dissection of the genitalia,

identification was made to aggregate level (e.g. Common Rustic agg. *Mesapamea*

secalis/didyma). After identification, moths were allowed to air-dry at room temperature for a

minimum of one week, which was sufficient for the dry body mass of even the largest

individuals to stabilize (Appendix S1.2, Fig. S2). After drying, we measured the forewing

length and dry mass of each moth. Forewing length was measured from wing base to wing-

tip, using calipers and a ruler, to the nearest 1 mm. Dry mass was measured using an A&D

HR-202 balance (A&D Instruments Ltd., Abingdon, UK), to the nearest 0.01 mg.

Measurements were precise to within ± 6 % of the true value (Appendix S1.2).

Modelling forewing length – body mass relationship from empirical data

To investigate the relationship between forewing length (mm) and body mass (mg) in moths,

we constructed generalized linear mixed-effects models (GLMMs) using our 2018 field data,

with species as a random effect, and body mass explained by the interaction between

forewing length and taxonomic family. We selected between three candidate model

structures by comparison of Bayesian Information Criterion (BIC) scores: (i) linear predictor

(i.e. ln(body mass) ~ wing length × family); (ii) non-linear predictor (i.e. ln(body mass) ~

ln(wing length) × family); and (iii) segmented predictor (as for model i, but permitting the

slope of the model to change once as forewing length increases). Finally, we tested the

significance of independent variables, including the interaction between wing length and

family, using Likelihood Ratio Tests.

To reduce the risk of our predictive model overfitting for families represented by only a few

species in our dataset (and therefore to allow accurate predictions of body mass to be

made), we refitted this model with a simplified family variable, in which seven families

represented by fewer than five species in our dataset were grouped together as 'other'

(effectively reducing the family variable from 11 categories to 5). The four retained families

(each with ≥ 5 species sampled) were Crambidae, Erebidae, Geometridae and Noctuidae,

allowing the predictive model's parameters to be refined for these families, whilst also

making overall predictions for all other families. We fitted a GLMM to the dataset as above,

using this reduced version of the family variable, and extracted all fitted parameters from the

GLMM to form the predictive model. We did not include information on whether individuals

were male or female, even though male and female moths can differ substantially in size in

some species, because this information is not recorded in historical abundance datasets.

Our model therefore used overall slope and intercept to predict body mass from forewing

length for all moths, with a refined prediction for moths from the most speciose (and

therefore data-rich) four families in our dataset.

Testing model accuracy

To test the accuracy of this general predictive modelling approach when making predictions

based on forewing length data from field guides, we estimated the body mass of each of the

94 moth species in our dataset from its expected forewing length (obtained by taking the

midpoint between minimum and maximum forewing lengths given by field guides for micro-

moths (Sterling & Parsons, 2012) and macro-moths (Waring & Townsend, 2017); Table S1),

and used these estimates to calculate the estimated biomass of each mixed-species sample

of moths (where one sample = all moths that were captured at the same site on the same

day, across multiple traps; n = 44 samples). We compared between these estimates of

biomass, and the empirically-measured biomass of the moths in question. We conducted

this testing at both species- and sample-levels, because rare species from rare families are

likely to have the least accurate predictions from our model, but may also have the least

impact on the accuracy of sample-level predictions.

We first compared between measured and estimated biomass for the full set of 600 moths.

At both species- and sample-level, we tested the relationship between measured and

predicted biomass, using model II regressions with a Major Axis approach because neither

biomass variable was dependent upon the other (Legendre & Legendre, 2012). Significance

of relationships from random was tested using one-tailed permutation tests (with 100

permutations), and relationships were also compared to the desired y = x (i.e. estimated =

measured) relationship by calculation of 95% confidence intervals around the estimated

slope. The strength of the relationships between measured and estimated biomass at

species- and sample-level was determined by model R^2 values.

However, because in this case comparisons were not independent of the predictive model

(i.e. model accuracy was tested with the same data that had been used to fit the model), we

also used a resampling approach to further test the accuracy of our general predictive

modelling approach. We split our full dataset 10,000 times into training and testing subsets.

In each replicate, we randomly selected 480 individual moths (80 % of the 600 total

individuals) without replacement to form a training subset, with the remaining 120 individuals

forming an independent testing subset. We trained a model with the same structure as the

full predictive model (above) on the training dataset, and from its parameters, extracted

estimates of species- and sample-level biomass as above for the 120 moths included in the

testing dataset. We tested the relationship between measured and predicted biomass for

each replicate as above. Across the results of all 10,000 replicates (and at both species- and

sample-levels), we then calculated the proportion of replicates for which measured and

estimated biomass were significantly correlated, the mean and standard error of model R^2

values, and the proportion of replicates for which the modelled relationship was significantly

different from y = x.

Finally, we used a resampling approach to assess the influence of moth abundance (i.e.

sample size) on prediction error. We randomly sampled sets of individuals (with

replacement, from the full set of 600 measured individuals) at sample sizes between 10 and

1000 in steps of 10, taking 1000 replicates at each sample size for a total of 100,000

replicates. For each replicate sample, we calculated the measured biomass and the

estimated biomass (based on the parameters of the final predictive model). We then

calculated the prediction error for each sample as a proportion of the true biomass,

normalizing by subtracting the known prediction error of 3.40 % in the full dataset (i.e. the

total predicted biomass of all 600 moths was 3.40 % lower than their total measured

biomass), such that:

$$prediction\ error\ = \left(100\ \times \frac{predicted\ biomass\ -\ measured\ biomass}{measured\ biomass}\right) - 3.4$$

Grouping sample sizes into windows of 100, we calculated the mean, standard error and

range of prediction errors observed across all replicates in each window.

All statistical analyses were conducted in R version 3.6.1 (R Core Team, 2019) using the

following packages: *lme4* to fit and assess linear mixed-effects models (Bates *et al.*, 2015);

lmodel2 to conduct model II regressions (Legendre, 2018); and *ggplot2* to plot figures

(Wickham, 2016). All R scripts and data used in the analysis are archived online at Zenodo

(doi: 10.5281/zenodo.3734274).

Results

Field sampling, identification and measurement of moths

We sampled 614 individual moths, of which 13 could not be confidently identified beyond

family level. One micro-moth (*Narycia duplicella* [Goeze, 1783], Psychidae) could not be

detected by our balance (and therefore weighed less than 0.005 mg). These 14 individual

moths were excluded from further analyses. The remaining dataset contained exactly 600

individual moths, representing 94 species from 11 families (6.6 % of all species, or 13.7 % of

macro-moth species, ever recorded in vice-county 61; Table S2). Among these moths,

forewing lengths ranged from 7 mm (individuals of *Eudonia pallida* (Crambidae) and *Agapeta*

hamana (Tortricidae)) to 40 mm (an individual of *Laothoe populi* (Sphingidae)) and dry body

masses ranged from 1.1 mg (an individual of *Eupithecia tenuiata* (Geometridae)) to 753.2

mg (an individual of *Smerinthus ocellata* (Sphingidae)).

Modelling forewing length – body mass relationship from empirical data

From the three candidate model structures described above, we selected the non-linear

predictor (model ii) as the best-fitting model (BIC: 360.7, compared to 431.1 and 494.3 for

models i and iii respectively). The natural logarithms of body mass and forewing length were

significantly related to each other at both species and individual levels (Fig. 1), with variation

among the 11 families in the slope and intercept of this relationship (individual level: χ^2 =

35.9, d.f. = 10, $P < 0.001$; marginal R^2 = 0.819) revealing that interfamilial variation in body

plan significantly influences the scaling of forewing length to body mass.

The significance of the model (and almost all of its explained variance) was retained when

fitting the simplified model (in which seven families represented by < 5 species were

grouped as 'other'; χ^2 = 30.7, d.f. = 4, $P < 0.001$; marginal R^2 = 0.812), resulting in a set of

parameters from which body mass could be predicted based on forewing length (Table 1).

All four families retained as independent levels (Crambidae, Erebidae, Geometridae and

Noctuidae) had larger intercepts and shallower slopes than the overall prediction across the

other families (Table 1). Thus we conclude that the non-linear model with simplified family

variable has the greatest potential for estimating body mass.

Testing model accuracy

We then used our best-fitting model to estimate body masses for all 94 species as described

above, and compared between measured and estimated biomass for the full sample of 600

individual moths. We found that our estimates of biomass significantly correlated with

measured biomass at both species- and sample-levels (Fig. 2), even though body mass

varied widely both within and between species (within-species s.d. of body mass = 34.6 mg,

between-species s.d. of body mass = 74.7 mg). At sample-level, the relationship between

estimated and measured biomass was not significantly different from a 1:1 relationship

(Table 2), with 91.5 % of variation explained. At species-level, estimated biomass explained

91.1 % of variation. The relationship was less steep than the expected 1:1 relationship

(Table 2) with all moths included; however, the 1:1 relationship was recovered when we

excluded the 34 smallest species from models (i.e. only included species weighing > 15 mg,

n = 60 species). These results indicate that our predictive model may slightly overestimate

the body mass of very small species of moths, but that this does not substantially bias

estimates of sample-level biomass.

To test whether this general predictive modelling approach can accurately estimate biomass

beyond the sampled individuals and species, we split our data 10,000 times into random

training (480 individuals in each case) and testing (120 individuals) subsets. We refitted our

final model to the training subset in each case and predicted the body masses of individuals

in each testing subset. We found again that our estimates of biomass significantly correlated

with the measured biomass at both species- and sample-levels in 100% of replicates (Table

3). At sample-level, estimated biomass explained on average 88.4 % (± s.e. 0.07) of

variation in measured biomass, and was not significantly different from a 1:1 relationship in

75.6 % of cases (Table 3), despite predictive models being built on a substantially reduced

dataset compared to our final model. At species-level, estimated biomass explained 87.4 %

(± s.e. 0.03) of variation in measured biomass, but the relationship was not significantly

different from the expected 1:1 relationship in 19.5 % of cases (being significantly less steep

than the expected relationship in the remaining 80.5 %: Table 3). As above, when we

excluded the species weighing < 15 mg from testing subsets, the relationship was not

significantly different from 1:1 in 81.3 % of cases (Table 3). These results indicate that

predictions made using this approach are likely to remain accurate even when predicting

beyond the training dataset.

Testing the influence of sample size on prediction error, we found that prediction error

decreased initially as sample size increased, but remained relatively stable for samples

larger than approximately 250 moths (Fig. S4). For samples of 10-100 moths, the standard

error of prediction error was 0.13 (range -61.79 to 83.42 %), whereas for samples of 910-

1000 moths, the prediction error was much less variable (s.e. 0.03, range -9.20 to 10.23 %).

This indicates that sample-level estimates of biomass are especially accurate for samples

containing > 250 moths.

Discussion

Findings from our analyses show a strong relationship between forewing length and body

mass in moths, which enables prediction (to an informative level of accuracy) of the biomass

of samples of moths when such data are not available (e.g. because historical specimens

have not been kept). Generating biomass data using this approach will provide an additional

tool to ongoing investigation of the nature and consequences of changes in insect

populations (Hallmann *et al.*, 2017; Macgregor *et al.*, 2019b; Didham *et al.*, 2020b) using

long-term recording datasets. It may also permit the inclusion of estimates of moth body

mass in comparative studies and trait-based analyses, despite the general lack of empirical

data of this nature (García-Barros, 2015). In particular, these data will facilitate studies of the

relationships between biomass, abundance and community composition (Appendix S2),

including important ecological questions such as: do biomass declines indicate a general

decline in the abundance of the majority of species, a severe decline in the biomass of a few

key species (e.g. Shortall *et al.*, 2009), a shift in community composition towards smaller-

bodied species, all of the above, or something else entirely?

Evaluation of the predictive model's current and future utility

Overall, the estimates of body mass calculated using the predictive model's parameters

performed relatively well during testing, with ~ 90 % of variation in measured biomass

explained by predicted biomass at both species- and sample-levels, and prediction error

decreasing as sample size increased (Fig. S4). Therefore, using estimated body masses

from the model (Table S1) to calculate the combined biomass of large samples of moths

should yield accurate results. Our sampled dataset was only sufficiently data-rich to allow

refined parameter estimates for four families (Crambidae, Erebidae, Geometridae and

Noctuidae). Further improvement of the model's accuracy may be possible by means of

family-specific parameter estimates for other families, at sub-family level (to better account

for within-family variation in body plan), or potentially even through a phylogenetic imputation

approach (Penone *et al.*, 2014). However, all of these approaches require the collection of

additional data on body masses from a greater number of species from families which are

less abundant in moth-trap samples than the dominant family, Noctuidae, and so collection

of sufficient data using field sampling may be challenging. An alternative may be to use

museum collections to measure individuals of a much wider range of species and families,

where methods exist to account for the mass of entomological pins when taking such

measurements (Gilbert, 2011). Taking such an approach might allow for more data to be

collected even from rarely-trapped families (e.g. Sphingidae), or those which are speciose

globally but have few (e.g. Saturniidae) or no (e.g. Hedylidae) species extant in Britain,

thereby potentially allowing refined predictions to be made for many more families,

potentially even at a global scale. Nevertheless, estimates of British moth biomass made

using our approach (Macgregor *et al.*, 2019b) revealed that 93.3% of total biomass is

comprised of the three macro-moth families for which we made refined predictions

(Erebidae, Geometridae and Noctuidae), so improving prediction accuracy for other families

may have limited impact on the sample-level accuracy of the overall model.

One source of potential error when using published forewing lengths to estimate biomass is

that 19% of individuals in our 2018 dataset had a measured forewing length which was

outside the expected range given by field guides. Nevertheless, there was an overall

correlation (R^2 = 0.942) between the mean forewing length at species-level derived from our

2018 empirical measurements and the midpoint of the range of forewing lengths for each

species, taken from the published field guides (Fig. S5). This suggests sufficient accuracy in

our approach, particularly considering that our largest measured species had a forewing

length 571 % larger than that of our smallest species. Similarly, the approaches we took to

measuring forewing lengths (i.e. with analogue callipers and a ruler, to the nearest 1mm)

and dry body masses (i.e. after one week of air-drying) mean that our dataset may not be

fully comparable to datasets collected under other conditions or using other approaches (e.g.

using digital callipers with higher resolution to measure forewing length, or measuring dry

body mass after oven-drying). However, since all air-drying took place, and all

measurements were taken, by the same person in the same laboratory over the same 6-

week period (and air-drying for one week was shown to be sufficient for the mass of even

the largest moths to stabilise: Fig. S2), these measurements are adequate to accurately

establish the relative relationships between species for both forewing length and dry body

mass. Therefore, our models can also be safely used to estimate relative change in moth

biomass over time, or in space, assuming only that the average body mass of each

individual species does not substantially change over the same scales.

An additional source of possible error in our models is sexual dimorphism in moths. Some

moth species, including some sampled in our study (e.g. Drinker *Euthrix potatoria*;

Lasiocampidae), exhibit substantial sexual dimorphism in wing length (Waring & Townsend,

2017) and in body mass (Allen *et al.*, 2011). However, we did not quantify or adjust for

sexual dimorphism in this study because long-term recording schemes rarely include

information on sex of individual moths, even for dimorphic species, although the majority of

such records are likely to be males (Altermatt *et al.*, 2009). Therefore, for estimation of

sample-level biomass, it will be of most use to provide a single average estimate of body

mass per species, regardless of size dimorphism.

Future research using our predictive model to study biomass change

Questions remain regarding temporal, spatial, and taxonomic variation in observed biomass

declines (Shortall *et al.*, 2009; Macgregor *et al.*, 2019b; Didham *et al.*, 2020b), the potential

drivers of these declines (Grubisic *et al.*, 2018; Komonen *et al.*, 2019; Didham *et al.*, 2020a),

and the challenges of extrapolating across data types, geographic locations, and temporal

and spatial scales (Thomas *et al.*, 2019; Wagner, 2019; Didham *et al.*, 2020b). Our study

illustrates the power of predictive models of body mass to tackle these challenges. Applying

these estimates in the same way to RIS datasets across the UK over longer time-scales, or

to other long-term moth abundance datasets, such as the National Moth Recording Scheme

or the Garden Moth Scheme (Fox *et al.*, 2011; Bates *et al.*, 2014), will facilitate investigation

of declines over longer time-periods and broader geographical scales than has previously

been feasible. Moreover, the same general approach could be used to estimate body mass

of moths in other databases, including macro-moth recording schemes from other regions

(e.g. the Noctua database; Groenendijk & Ellis, 2011) and micro-moths, which were

incorporated into the NMRS in 2016. Combining with similar existing models for other insect

families and invertebrate taxa (Sage, 1982; Sample *et al.*, 1993; Sabo *et al.*, 2002; Höfer &

Ott, 2009) could facilitate comparison of biomass losses across multiple datasets and taxa at

a global scale. However, a possibility that cannot be accounted for using this approach is

change over time in the mean body mass of individual moths (Wu *et al.*, 2019), e.g. in

response to climate warming (Gardner *et al.*, 2011).

Our approach may also be of use for conducting trait-based analyses of moths (e.g. van

Langevelde *et al.*, 2018), where it is important that trait data have high precision (Middleton-

Welling *et al.*, 2018). Our predictive model offers a means to estimate body mass

reproducibly, potentially across multiple data sources, using a trait (forewing length) that is

straightforward to measure using basic equipment, and therefore can be robustly applied to

other datasets. Previous trait-based analyses have used forewing length as a proxy for body

size, but we have shown that there is interfamilial variation in this relationship (Fig. 1), which

can be incorporated by using our approach. However, an appropriate level of caution is

advised before applying our specific estimates of body mass to systems where the moth

fauna is markedly different from that of the UK (e.g. tropical forests). Under such

circumstances, using a parallel approach to fit a new, regionally-specific predictive model

may be the most robust approach.

Conclusions

We have developed a predictive model to estimate the dry body mass of moths based on

their forewing length, using it to generate body masses for all British species of macro-moth.

The predictions of sample biomass made by our model correlated strongly with measured

biomass of the same samples ($R^2 = 0.915$), indicating that this approach provides a robust

way to estimate the biomass of samples of moths identified to species level. Our approach

unlocks new opportunities to study trends in moth biomass over time and over large

geographic regions.

Acknowledgements

This study was funded by the Department of Biology, University of York, through an internal

grant awarded to C.J.M. and R.S.K. We are grateful to collaborators and staff who have

contributed data from the Rothamsted Insect Survey, a BBSRC-supported National

Capability. C.D.T., J.K.H. and C.J.M. were additionally supported by the Natural

Environment Research Council (grant no. NE/N015797/1).

Author contributions

The study was conceived by C.J.M. and C.D.T., and designed by those authors in

discussion with J.K.H. Field and laboratory work was conducted by R.S.K., who also carried

out the statistical analysis with C.J.M. The first draft of the manuscript was written by R.S.K.

and C.J.M., and all authors contributed to subsequent revisions.

Data Accessibility

All R scripts and data used in the analysis are archived online at Zenodo (doi:

10.5281/zenodo.3734274).

Competing Interests

The authors declare no competing interests.

References

Allen, C.E., Zwaan, B.J. & Brakefield, P.M. (2011) Evolution of sexual dimorphism in the

Lepidoptera. *Annual Review of Entomology*, **56**, 445–464.

Altermatt, F., Baumeyer, A. & Ebert, D. (2009) Experimental evidence for male biased flight-

to-light behavior in two moth species. *Entomologia Experimentalis et Applicata*, **130**, 259–

265.

Bates, A.J., Sadler, J.P., Grundy, D., Lowe, N., Davis, G., Baker, D., *et al.* (2014) Garden

and landscape-scale correlates of moths of differing conservation status: significant effects

of urbanization and habitat diversity. *PLoS One*, **9**, e86925.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, **67**, 1–48.

Bell, J.R., Blumgart, D. & Shortall, C.R. (2020) Are insects declining and at what rate? An

analysis of standardised, systematic catches of aphid and moth abundances across Great

Britain. *Insect Conservation and Diversity*, **13**, 115–126.

Brose, U., Berlow, E.L. & Martinez, N.D. (2005) Scaling up keystone effects from simple to

complex ecological networks. *Ecology Letters*, **8**, 1317–1325.

Conrad, K.F., Warren, M.S., Fox, R., Parsons, M.S. & Woiwod, I.P. (2006) Rapid declines of

common, widespread British moths provide evidence of an insect biodiversity crisis.

Biological Conservation, **132**, 279–291.

Coulthard, E., Norrey, J., Shortall, C. & Harris, W.E. (2019) Ecological traits predict

population changes in moths. *Biological Conservation*, **233**, 213–219.

Davis, R.B., Õunap, E., Javoiš, J., Gerhold, P. & Tammaru, T. (2013) Degree of

specialization is related to body size in herbivorous insects: a phylogenetic confirmation.

Evolution, **67**, 583–589.

Denerley, C., Redpath, S.M., Wal, R. van der, Newson, S.E., Chapman, J.W. & Wilson, J.D.

(2018) Breeding ground correlates of the distribution and decline of the Common Cuckoo

Cuculus canorus at two spatial scales. *The Ibis*, **354**, 547.

Didham, R.K., Barbero, F., Collins, C.M., Forister, M.L., Hassall, C., Leather, S.R., *et al.*

(2020a) Spotlight on insects: trends, threats and conservation challenges. *Insect*

Conservation and Diversity, **13**, 99–102.

Didham, R.K., Basset, Y., Collins, C.M., Leather, S.R., Littlewood, N.A., Menz, M.H.M., *et al.*

(2020b) Interpreting insect declines: seven challenges and a way forward. *Insect*

Conservation and Diversity, Nature and Environment Series, **13**, 103–114.

Fox, R. (2013) The decline of moths in Great Britain: a review of possible causes: The

decline of moths in Great Britain. *Insect Conservation and Diversity*, **6**, 5–19.

Fox, R., Oliver, T.H., Harrower, C., Parsons, M.S., Thomas, C.D. & Roy, D.B. (2014) Long-

term changes to the frequency of occurrence of British moths are consistent with opposing

and synergistic effects of climate and land-use changes. *Journal of Applied Ecology*, **51**,

949–957.

Fox, R., Randle, Z., Hill, L., Anders, S., Wiffen, L. & Parsons, M.S. (2011) Moths count:

recording moths for conservation in the UK. *Journal of Insect Conservation*, **15**, 55–68.

Franklin, A.J., Liebhold, A.M., Murray, K. & Donahue, C. (2003) Canopy herbivore

community structure: large-scale geographical variation and relation to forest composition.

Ecological Entomology, **28**, 278–290.

García-Barros, E. (2015) Multivariate indices as estimates of dry body weight for

comparative study of body size in Lepidoptera. *Nota Lepidopterologica*, **38**, 59–74.

Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L. & Heinsohn, R. (2011) Declining body

size: a third universal response to warming? *Trends in Ecology & Evolution*, **26**, 285–291.

Gilbert, J.D.J. (2011) Insect dry weight: Shortcut to a difficult quantity using museum

specimens. *The Florida Entomologist*, **94**, 964–970.

Gilburn, A.S., Bunnefeld, N., Wilson, J.M., Botham, M.S., Brereton, T.M., Fox, R., *et al.*

(2015) Are neonicotinoid insecticides driving declines of widespread butterflies? *PeerJ*, **3**,

e1402.

Groenendijk, D. & Ellis, W.N. (2011) The state of the Dutch larger moth fauna. *Journal of*

Insect Conservation, **15**, 95–101.

Grubisic, M., Grunsven, R.H.A. van, Kyba, C.C.M., Manfrin, A. & Hölker, F. (2018) Insect

declines and agroecosystems: does light pollution matter?: Insect declines and

agroecosystems. *Annals of Applied Biology*, **8**, e67798.

Hallmann, C.A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., *et al.* (2017)

More than 75 percent decline over 27 years in total flying insect biomass in protected areas.

PLoS One, **12**, e0185809.

Hallmann, C.A., Zeegers, T., Klink, R., Vermeulen, R., Wielink, P., Spijkers, H., *et al.* (2019)

Declining abundance of beetles, moths and caddisflies in the Netherlands. *Insect*

Conservation and Diversity, **6**, 24451.

Harris, J.E., Rodenhouse, N.L. & Holmes, R.T. (2019) Decline in beetle abundance and

diversity in an intact temperate forest linked to climate warming. *Biological Conservation*,

240, 108219.

Heath, J. (1965) A genuinely portable MV light trap. *Entomologist's Record and Journal of*

Variation, **77**, 236–238.

Höfer, H. & Ott, R. (2009) Estimating biomass of Neotropical spiders and other arachnids

(Araneae, Opiliones, Pseudoscorpiones, Ricinulei) by mass-length regressions. *Journal of*

Arachnology, 160–169.

Hooks, C.R.R., Pandey, R.R. & Johnson, M.W. (2003) Impact of avian and arthropod

predation on lepidopteran caterpillar densities and plant productivity in an ephemeral

agroecosystem. *Ecological Entomology*, **28**, 522–532.

Janzen, D.H. (1984) Two ways to be a tropical big moth: Santa Rosa saturniids and

sphingids. *Oxford Surveys in Evolutionary Biology*, **1**, 140.

Komonen, A., Halme, P. & Kotiaho, J.S. (2019) Alarmist by bad design: Strongly popularized

unsubstantiated claims undermine credibility of conservation science. *Rethinking Ecology*, **4**,

17–19.

Langevelde, F. van, Braamburg-Annegarn, M., Huigens, M.E., Groendijk, R., Poitevin, O.,

Deijk, J.R. van, *et al.* (2018) Declines in moth populations stress the need for conserving

dark nights. *Global Change Biology*, **24**, 925–932.

Langevelde, F. van, Ettema, J.A., Donners, M., WallisDeVries, M.F. & Groenendijk, D.

(2011) Effect of spectral composition of artificial light on the attraction of moths. *Biological*

Conservation, **144**, 2274–2281.

Legendre, P. (2018) *lmodel2: Model II Regression*. R package version 1.7-3. https://cran.R-

project.org/package=lmodel2.

Legendre, P. & Legendre, L.F.J. (2012) *Numerical Ecology*. Elsevier.

Lister, B.C. & Garcia, A. (2018) Climate-driven declines in arthropod abundance restructure

a rainforest food web. *Proceedings of the National Academy of Sciences of the United*

States of America, **115**, E10397–E10406.

Macgregor, C.J., Kitson, J.J.N., Fox, R., Hahn, C., Lunt, D.H., Pocock, M.J.O., *et al.* (2019a)

Construction, validation, and application of nocturnal pollen transport networks in an agro-

ecosystem: a comparison using light microscopy and DNA metabarcoding. *Ecological*

Entomology, **44**, 17–29.

Macgregor, C.J., Pocock, M.J.O., Fox, R. & Evans, D.M. (2015) Pollination by nocturnal

Lepidoptera, and the effects of light pollution: a review. *Ecological Entomology*, **40**, 187–198.

Macgregor, C.J., Williams, J.H., Bell, J.R. & Thomas, C.D. (2019b) Moth biomass increases

and decreases over 50 years in Britain. *Nature Ecology & Evolution*, **3**, 1645–1649.

Mattila, N., Kaitala, V., Komonen, A., Päivinen, J. & Kotiaho, J.S. (2011) Ecological

correlates of distribution change and range shift in butterflies. *Insect Conservation and*

Diversity, **4**, 239–246.

Middleton-Welling, J., Wade, R.A., Dennis, R.L.H., Dapporto, L. & Shreeve, T.G. (2018)

Optimising trait and source selection for explaining occurrence and abundance changes: A

case study using British butterflies. *Functional Ecology*, **111**, 543.

Miller, W.E. (1997) Body weight as related to wing measure in hawkmoths (Sphingidae).

Journal of the Lepidopterists' Society, **51**, 91–92.

Morecroft, M.D., Bealey, C.E., Beaumont, D.A., Benham, S., Brooks, D.R., Burt, T.P., *et al.*

(2009) The UK Environmental Change Network: Emerging trends in the composition of plant

and animal communities and the physical environment. *Biological Conservation*, **142**, 2814–

2832.

Outhwaite, C.L., Gregory, R.D., Chandler, R.E., Collen, B. & Isaac, N.J.B. (2020) Complex

long-term biodiversity change among invertebrates, bryophytes and lichens. *Nature Ecology*

& Evolution, 1–9.

Penone, C., Davidson, A.D., Shoemaker, K.T., Di Marco, M., Rondinini, C., Brooks, T.M., *et*

al. (2014) Imputation of missing data in life-history trait datasets: which approach performs

the best? *Methods in Ecology and Evolution*, **5**, 961–970.

Pöyry, J., Carvalheiro, L.G., Heikkinen, R.K., Kühn, I., Kuussaari, M., Schweiger, O., *et al.*

(2017) The effects of soil eutrophication propagate to higher trophic levels: Effects of soil

eutrophication on herbivores. *Global Ecology and Biogeography*, **26**, 18–30.

R Core Team. (2019) *R: A language and environment for statistical computing.* R

Foundation for Statistical Computing, Vienna, Austria.

Roth, N., Zoder, S., Zaman, A.A., Thorn, S. & Schmidl, J. (2020) Long-term monitoring

reveals decreasing water beetle diversity, loss of specialists and community shifts over the

past 28 years. *Insect Conservation and Diversity*, **13**, 140–150.

Sabo, J.L., Bastow, J.L. & Power, M.E. (2002) Length–mass relationships for adult aquatic

and terrestrial invertebrates in a California watershed. *Journal of the North American*

Benthological Society, **21**, 336–343.

Sage, R.D. (1982) Wet and dry-weight estimates of insects and spiders based on length.

The American Midland Naturalist, **108**, 407–411.

Salcido, D.M., Forister, M.L., Garcia Lopez, H. & Dyer, L.A. (2020) Loss of dominant

caterpillar genera in a protected tropical forest. *Scientific Reports*, **10**, 422.

Sample, B.E., Cooper, R.J., Greer, R.D. & Whitmore, R.C. (1993) Estimation of insect

biomass by length and width. *The American Midland Naturalist*, 234–240.

Seibold, S., Gossner, M.M., Simons, N.K., Blüthgen, N., Müller, J., Ambarlı, D., *et al.* (2019)

Arthropod decline in grasslands and forests is associated with landscape-level drivers.

Nature, **574**, 671–674.

Shortall, C.R., Moore, A., Smith, E., Hall, M.J., Woiwod, I.P. & Harrington, R. (2009) Long-

term changes in the abundance of flying insects. *Insect Conservation and Diversity*, **2**, 251–

260.

Sierro, A., Arlettaz, R., Naef-Daenzer, B., Strebel, S. & Zbinden, N. (2001) Habitat use and

foraging ecology of the nightjar (*Caprimulgus europaeus*) in the Swiss Alps: towards a

conservation scheme. *Biological Conservation*, **98**, 325–331.

Singer, M.S., Farkas, T.E., Skorik, C.M. & Mooney, K.A. (2012) Tritrophic interactions at a

community level: effects of host plant species quality on bird predation of caterpillars. *The*

American Naturalist, **179**, 363–374.

Sterling, P. & Parsons, M. (2012) *Field Guide to the Micro-Moths of Great Britain and*

Ireland. First Edition. British Wildlife Publishing Ltd, Gillingham.

Storkey, J., Macdonald, A.J., Bell, J.R., Clark, I.M., Gregory, A.S., Hawkins, N.J., *et al.*

(2016) Chapter One - The Unique Contribution of Rothamsted to Ecological Research at

Large Temporal Scales. In *Advances in Ecological Research* (ed. by Dumbrell, A.J., Kordas,

R.L. & Woodward, G.). Academic Press, pp. 3–42.

Strien, A.J. van, Swaay, C.A.M. van, Strien-van Liempt, W.T.F.H. van, Poot, M.J.M. &

WallisDeVries, M.F. (2019) Over a century of data reveal more than 80% decline in

butterflies in the Netherlands. *Biological Conservation*, **234**, 116–122.

Thomas, C.D., Jones, T.H. & Hartley, S.E. (2019) "Insectageddon": A call for more robust

data and rigorous analyses. *Global Change Biology*, **25**, 1891–1892.

Threlfall, C.G., Law, B. & Banks, P.B. (2012) Sensitivity of insectivorous bats to urbanization:

Implications for suburban conservation planning. *Biological Conservation*, **146**, 41–52.

Valtonen, A., Hirka, A., Szőcs, L., Ayres, M.P., Roininen, H. & Csóka, G. (2017) Long-term

species loss and homogenization of moth communities in Central Europe. *Journal of Animal*

Ecology, **86**, 730–738.

Valtonen, A., Leinonen, R., Pöyry, J., Roininen, H., Tuomela, J. & Ayres, M.P. (2014) Is

climate warming more consequential towards poles? The phenology of Lepidoptera in

Finland. *Global Change Biology*, **20**, 16–27.

Vaughan, N. (1997) The diets of British bats (Chiroptera). *Mammal Review*, **27**, 77–94.

Wagner, D.L. (2019) Global insect decline: Comments on Sánchez-Bayo and Wyckhuys

(2019). *Biological Conservation*, **233**, 332–333.

Waring, P. & Townsend, M. (2017) *Field Guide to the Moths of Great Britain and Ireland*

(Field Guides). 3rd Revised Edition. Bloomsbury Natural History.

Wepprich, T., Adrion, J.R., Ries, L., Wiedmann, J. & Haddad, N.M. (2019) Butterfly

abundance declines over 20 years of systematic monitoring in Ohio, USA. *PLoS One*, **14**,

e0216270.

Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New

York.

Wickramasinghe, L.P., Harris, S., Jones, G. & Jennings, N.V. (2004) Abundance and

species richness of nocturnal insects on organic and conventional farms: Effects of

agricultural intensification on bat foraging. *Conservation Biology*, **18**, 1283–1292.

Wilson, J.F., Baker, D., Cheney, J., Cook, M., Ellis, M., Freestone, R., *et al.* (2018) A role for

artificial night-time lighting in long-term changes in populations of 100 widespread macro-

moths in UK and Ireland: a citizen-science study. *Journal of Insect Conservation*, **22**, 189–

196.

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J.M., Olesen, J.M., Valido, A., *et al.*

(2005) Body size in ecological networks. *Trends in Ecology & Evolution*, **20**, 402–409.

Wu, C.-H., Holloway, J.D., Hill, J.K., Thomas, C.D., Chen, I.-C. & Ho, C.-K. (2019) Reduced

body sizes in climate-impacted Borneo moth assemblages are primarily explained by range

shifts. *Nature Communications*, **10**, 4612.

Figures and tables

Figure 1: Relationship between forewing length (mm) and dry mass (mg). In the top panel,

the mean forewing length and dry mass of each species sampled in the study is shown on

logarithmic axes, with error bars showing standard errors and family indicated by the

combination of point colour and shape. In panel (b), the forewing length and dry mass of

every individual moth sampled in the study is shown on logarithmic axes, with the four most

speciose families in our sample (Crambidae, Erebidae, Geometridae and Noctuidae)

indicated as above by point colour and shape.

Figure 2: Accuracy of predicted biomass of moth species and samples of moths compared

to the true, measured biomass. (a) Predicted dry mass of species (mg) is plotted against

mean measured dry mass (mg); the 1:1 relationship is plotted as a blue line, and points are

coloured by the number of individual moths from which the measured mean was calculated.

(b) The absolute difference between mean measured dry mass and predicted dry mass of

each moth species is plotted against the number of individuals from which the measured

mean was calculated; a horizontal line is plotted at y = 0. (c) Predicted dry mass of samples

(mg) is plotted against measured dry mass (mg); the 1:1 relationship is plotted as a blue line,

and points are coloured by the number of individual moths contained in the sample. (d) The

absolute difference between measured and predicted dry mass of each sample of moths is

plotted against measured dry mass (mg); a horizontal line is plotted at y = 0.

Table 1: Parameters of the predictive model, extracted by fitting a GLMM with the fixed-

effects structure: ln(body mass) ~ ln(forewing length) × family, to data from 600 individual

moths. The number of measured individuals and species on which each parameter estimate

was based is given. Overall model parameters are given, including the χ^2 and *P*-values of a

Likelihood Ratio Test of the model's overall significance. Family-specific slope and intercept

values are refinements to be added to the parameters for 'other families' (rather than taken

in isolation). To predict body mass of a moth from its forewing length, these parameters

should be applied to the following formula: ln(body mass) = (ln(forewing length) × ('other

families' slope + family slope adjustment)) + ('other families' intercept + family intercept

adjustment).

Family adjustment	*n* species (*n* individuals)	χ^2, d.f. (*P*)	Slope estimate (s.e.)	Intercept estimate (s.e.)
Overall model	94 (600); 11 families	30.7, 4 (<0.001)	-	-
'Other families' (no adjustment)	15 (67); 7 families	-	3.056 (0.180)	-5.016 (0.540)
Crambidae	11 (38)	-	-0.904 (0.311)	1.361 (0.813)
Erebidae	10 (79)	-	-0.601 (0.360)	1.294 (1.029)
Geometridae	22 (52)	-	-0.492 (0.322)	0.344 (0.891)
Noctuidae	36 (364)	-	-1.297 (0.239)	3.788 (0.694)

689 **Table 2**: Details of statistical models testing the relationships between measured biomass

690 and estimated biomass at species- and sample-level for the final model. Relationships were

691 tested using a model II regression, and significance was determined by a one-tailed

692 permutation test with 100 permutations. The R^2 of each model is also given, alongside the

693 estimated intercept and slope of each model, with associated 95% confidence intervals.

Level	Data subset	n	Model R^2	Model intercept (95% CI)	Model slope (95% CI)	P
Sample	Full dataset	44	0.915	0.275 (-0.310 - 0.810)	0.952 (0.865 - 1.047)	0.010
Species	Full dataset	94	0.911	0.557 (0.382 - 0.723)	0.874 (0.819 - 0.932)	0.010
	Species weighing > 15 mg only	60	0.823	0.168 (-0.311 - 0.595)	0.964 (0.853 - 1.090)	0.010

Table 3: Details of bootstrap testing (over 10,000 replicates) of statistical models testing the

relationships between measured biomass and estimated biomass at species- and sample-

level. Each model was fitted to a training dataset consisting of 480 randomly-selected

individuals, and tested on the remaining 120 individuals. Relationships were tested using a

model II regression, and significance was determined by a one-tailed permutation test with

100 permutations. The R^2 of each model was also taken, alongside the 95% confidence

intervals for the estimated slope. Here, the number of replicates (/10,000) for which

measured and estimated biomass were significantly related is given, as well as the number

of replicates for which the 95% confidence intervals for the estimated slope contained 1 (i.e.

y = x). The mean model R^2 (and standard error) across all 10,000 replicates is also given.

For tests of the slope's relationship to 1, all models were re-tested with the species weighing

> 15 mg excluded from the testing dataset.

Level	Data subset	Bootstrap replicates	% models significant	% models slope not equal to 1	Mean model R^2 (s.e.)
Sample	Full dataset	10,000	100	24.4	0.8843 (0.0007)
Species	Full dataset	10,000	100	80.5	0.8736 (0.0003)
	Species weighing > 15 mg only	10,000	-	19.5	-

712