Finding missing links in interaction networks

J. Christopher D. Terry (1) james.terry@zoo.ox.ac.uk Corresponding author

Owen T. Lewis (1) owen.lewis@zoo.ox.ac.uk

1: Department of Zoology, University of Oxford, Oxford, OX1 3PS, UK.

Running title: Finding missing links in interaction networks

Keywords: ecological network, bipartite network, missing links, sampling completeness, network

metrics

Data accessibility statement

R functions to run the predictive models are publicly available in the R package *cassandRa* on CRAN.

All code and necessary data to generate simulated datasets and carry out analyses are available on

GitHub (github.com/jcdterry/PredictingMissingLinks). Should the manuscript be accepted, this will

also be archived in a public repository and the data doi included.

Statement of Authorship

JCDT initiated the research, in discussion with OTL. JCDT conducted all analyses and wrote the first

draft of the manuscript. Both authors contributed to revisions.

Abstract

Documenting which species interact within ecological communities is challenging and labour-

intensive. As a result, many interactions remain unrecorded, potentially distorting our understanding

of network structure and dynamics. We test the utility of four structural models and a new coverage-

deficit model for predicting missing links in both simulated and empirical bipartite networks. We find

they can perform well, but that the predictive power of structural models varies with the underlying

network structure. Predictions can be improved by ensembling multiple models. Sample-coverage

estimators of the number of missed interactions are highly correlated with the number of missed

interactions, but strongly biased towards underestimating the true number of missing links.

Augmenting observed networks with most-likely missing links improves estimates of qualitative

network metrics. Simple to implement tools to identify likely missing links can enable prioritisation

of research effort and more robust assessment of network properties.

Introduction

Networks documenting trophic, competitive or mutualistic interactions among species are an

essential tool for understanding the structure and dynamics of ecological communities (Memmott

2009; Poisot *et al.* 2016b; Dormann *et al.* 2017; Delmas *et al.* 2019). Interaction networks can be a

better indicator of the resilience of communities than species counts (Valiente-Banuet *et al.* 2015)

and are increasingly applied in conservation (Tylianakis *et al.* 2010) and biomonitoring (Gray *et al.*

2014).

Documenting the species present in a system can be difficult enough (Gotelli & Colwell 2010);

accurately enumerating their interactions is even more challenging. As a result, ecological

interaction networks are routinely under-sampled, with consequences for our understanding of

communities (Goldwasser & Roughgarden 1997; Blüthgen 2010; Chacoff *et al.* 2012; Jordano 2016).

Ecological networks are sparse and hence any unobserved interactions could be either 'missing links'

(interactions that would be observed with complete sampling), or true negatives. Differences or

deficiencies in sampling can lead to divergent conclusions about underlying ecological mechanisms

(Lee & Guénard 2019). Furthermore, under-sampling will skew the apparent prevalence of

interactions towards frequent interactions and may miss infrequent interactions entirely (Dormann

et al. 2017).

Common solutions to this problem have included rarefying networks to a consistent level of

sampling (Morris *et al.* 2014), sampling until metric values stabilise (Rivera-Hutinel *et al.* 2012; Costa

et al. 2016), explicit observation models (Weinstein & Graham 2017), and the use of network metrics

that are comparatively robust to under sampling (Vizentin-Bugoni *et al.* 2016). There has been an

extensive literature testing these different approaches (Nielsen & Bascompte 2007; Blüthgen *et al.*

2008; Poisot *et al.* 2012b; Fründ *et al.* 2016; de Aguiar 2018; Henriksen *et al.* 2019), often arguing for

an emphasis on quantitative rather than qualitative metrics, which reduce the influence of missed

infrequent interactions.

Nonetheless, infrequent interactions may play an important role in ecological communities. Since

networks are not static, there is value in distinguishing infrequent interactions from those that are

genuinely absent. Rarely-observed links have the potential to grow in importance with changes in

biotic (Terry *et al.* 2017) or abiotic conditions (Staniczenko *et al.* 2017). At equilibrium, interactions

may occur at low frequency, yet be dynamically important. For instance, while interactions involving

a species suppressed to a low level by a consumer may be observed infrequently, removal of the

consumer can allow prey abundance to increase rapidly (Paine 1980). At a network level, the

prevalence and distribution of apparently weak or infrequent interactions can have marked

consequences for network stability and dynamics (McCann *et al.* 1998; Berlow 1999; Gellner &

McCann 2016; Jacquet *et al.* 2016; Kadoya *et al.* 2018).

Ecological networks are manifestly non-random (Dunne & Pascual 2006; Montoya *et al.* 2006;

Vázquez *et al.* 2009; Bascompte & Jordano 2013). In principle, it should be possible to use this

underlying structure to help infer where links are most likely to have been missed (Bartomeus 2013;

Morales-Castilla *et al.* 2015; Valdovinos 2019). To this end, a number of missing link inference

approaches have been tested on ecological networks. These have included hierarchical structuring

models (Clauset *et al.* 2008), linear filtering (Stock *et al.* 2017), matching-centrality models (Rohr *et*

al. 2016), stochastic block models (Guimera & Sales-Pardo 2009) and *k*-nearest neighbour

recommenders (Desjardins-Proulx *et al.* 2017). These models have shown that missing links can be

inferred from partially-observed structured networks. However, although ecological networks are

used as part of benchmarking tests, often the inference scenarios are not orientated towards

ecological needs and to date there has been limited uptake of these methods from the ecological

community.

Furthermore, it is common in empirical ecological networks for species to have been sampled to

markedly different extents (although see Novotny *et al.* 2010). This has the potential to give further

information about the potential location of missing links, since missing links are more likely for

poorly-sampled species (Blüthgen *et al.* 2008).

Here, we explore the potential to identify missing links in a wide range of ecological bipartite

networks with a range of models, including a new species-level coverage-deficit approach which

incorporates information on the completeness of sampling for individual species. We show that

inferential models can perform well, especially when used in ensemble, and demonstrate how they

could inform empirical work to document ecological networks and help evaluate hypotheses.

Methods

Predictive Models

We seek to infer the true interaction frequency matrix composed of elements a_{ij} starting from an

observed bipartite adjacency matrix **O**. This has n rows representing each focal layer species and m

columns detailing each potential interaction partner. Statistical models can be used to assign to each

unobserved interaction in a network a relative probability $P(a_{ij} > 0)$ of being a 'missing link' (false

negative), rather than a true negative.

Here, we introduce and test a new coverage deficit model and four structural models described in

the literature, singularly and in combination. All models were fit using the *cassandRa* R package

(Terry 2019). Initially we focus on identifying the qualitative network (i.e. where $a_{ij} \neq 0$), but we

later test the potential to augment networks with likely missed links and the quantitative

relationship between $P(a_{ij} > 0)$ and a_{ij}. We assume throughout that the species observed (the

dimensions of **O**) are a complete list – we do not attempt here to predict interactions with

unobserved species.

Figure 1. Demonstration of the fits of predictive models to a subsampled simulated bipartite network. Black dots (●)

indicate observed interactions, red targets (⊙) indicate unobserved missing links that we wish to identify. Brighter colours

indicate a higher probability of that interaction being present according to that model. Species are sorted to highlight the

features of each model. In the coverage deficit model, observed interactions (grey) are not assigned a likelihood and the

size of the dot is proportional to the number of observations. AUC values for this example were as follows: Degree: 0.80,

Latent-Trait: 0.76, Matching-Centrality: 0.81, SBM: 0.88, Coverage-Deficit: 0.7, Averaging the SBM and the Coverage-Deficit

model 0.92.

Coverage-Deficit Model

When ecological bipartite networks are constructed, it is common that one level (the focal level) of a

network is directly sampled and the interaction partners in the other level counted. Examples

include gathering insect hosts and rearing parasitoids from them, or observing pollinators that visit a

set of plant species. This sampling may be opportunistic, matched to the local abundance, or

standardised in some other manner. The result is a set of observed interaction counts with each

focal layer species. Although these interaction counts may be later rescaled to better represent local

abundances, here we assume the raw observation data is available. The method relies on

discretisable relationships where individual interaction events can be observed. This includes many

ecological interactions such as parasitism, pollination visits and feeding observations, but excludes

diffuse facilitatory or competitive interactions.

An extensive literature has developed to assess the sample completeness of species inventories

(Chao & Colwell 2017), based on the observation that the majority of information about the number

of missing observations is contained in the most infrequent observations. Due to its deep roots in

information theory these approaches can be applied to any case of estimating sample coverage,

including interaction sample coverage (Chacoff *et al.* 2012; Jordano 2016). The core of our coverage-

deficit model estimates the interaction partner sample completeness for each focal layer species,

based on the observed occurrence frequencies and the 'Chao1' estimator (Chao 1984). This estimate

of *coverage deficit* ($\hat{C}_{def}$), is a lower bound estimator of the proportion of interactions that occur

between as-yet-unobserved partners, or alternatively the probability that the next interaction

partner observed would not yet have been observed with that species. It is defined as:

$$\hat{C}_{def} = \frac{f_1}{n} \frac{f_1\,(n-1)}{f_1(n-1)+2f_2}$$

where f_1 is the number of interactions observed only once, f_2 is the number of interactions

observed exactly twice, and n is the total observed interactions involving that focal species.

Unfortunately, ecological interaction networks typically contain a number of focal layer species that

are extremely poorly observed, resulting in situations where the Chao1 estimator is undefined or

inappropriate. For very poorly sampled species, the distribution of interactions cannot itself provide

much information (Colwell & Coddington 1994). Where observations are exclusively singletons the

index will estimate zero coverage. At the other end of the scale, if a focal species is observed with no

singleton interactions the index will estimate that there is zero sample coverage deficit.

In these cases, we use a simple binomial model to estimate $\hat{C}_{def}$, the probability that the next

interactor drawn will not yet have been observed interacting with the focal species. The likelihood of

not having yet observed any of the unobserved missing links is: $L = \left(1 - \hat{C}_{def}\right)^n$. While the

maximum likelihood estimate of $\hat{C}_{def}$ will be 0, with a Bayesian approach and assuming a flat prior

the posterior mean can be found directly as: $\mathbb{E}(\hat{C}_{def}|n) = 1 - \frac{1}{n+2}$. We use this alternative estimate

when $n \leq 5$, all the observations are singletons ($f_1 = n$), or if there are no singletons or doubletons

($f_1 + f_2 = 0$). We use the $\hat{C}_{def_i}$ estimate for each focal layer species i to assign a relative probability

to each potential missing interaction as:

143
$$P\left(a_{ij} > 0\right) \sim \frac{\hat{C}_{def_i}}{\sum_j (O_{ij} = 0)}.$$

Since coverage deficit is strongly correlated with sample size, we also tested a simpler model that

assigns the relative probability of missed interactions based solely on the number of observations of

each focal layer-species: $P\left(a_{ij} > 0\right) \sim \frac{1}{\sum_j (O_{ij})}$.

Latent-Trait Model

Species do not interact at random with each other – similar species tend to interact with a

similar sets of interaction partners. It should be possible to use this information to guide our

expectations of missing links through a trait-based framework (Morales-Castilla *et al.* 2015;

Bartomeus *et al.* 2016; Laigle *et al.* 2018). These similarities can in directly measured traits, such as

corolla and tongue length (Vizentin-Bugoni *et al.* 2014; Klumpers *et al.* 2019) or through inferred

'latent' traits that are not directly measurable. Latent traits can be identified whether the

community is structured by phylogeny, microhabitat or physiological differences. A survey of

published networks found that the number of latent traits required may be quite low (Eklöf *et al.*

2013).

Intraspecific trait variation caused by spatial, temporal or ontogenetic differences can pose

considerable challenges (González-Varo & Traveset 2016). Nonetheless, such approaches have had

successes and a family of approaches have been developed including matching models (Rohr *et al.*

2016), Dirichlet-multinomial regression (Crea *et al.* 2016), matrix factorisation (Seo & Hutchinson

2018) and fourth-corner analyses (Spitz *et al.* 2014). While it is possible to fit any number of latent

traits, here we examine using a simple single latent trait model (detailed in SI 3) to reduce the

potential of overfitting (Rohr *et al.* 2016).

Degree Model

It is readily observable that some species have many interaction partners while other species

interact with very few. This manifests itself in skewed degree distributions within ecological

networks (Jordano *et al.* 2003). All else being equal, it would be less surprising to discover that a

missed interaction involving a generalist species, than one with few observed interactions. Many

bipartite interaction networks, especially mutualistic networks, are observed to be 'nested' - the

range of specialists is a subset of the generalists (Bascompte *et al.* 2003; Ulrich *et al.* 2009). Simple

degree-models can fit binary nestedness and preferential patterns well. Although nestedness can be

generated by a variety of processes (Song *et al.* 2017) and patterns observable in binary networks

may not be reflected in the quantitative networks (Staniczenko *et al.* 2013), our principle objective

here is to identify missing links. For our degree model (detailed in SI 3), we assign each species a

connectance term. When both species have high observed degrees, the probability of those species

interacting will be high.

Matching – Centrality Model

The matching – centrality approach (Rohr *et al.* 2016, 2018) combines a latent-trait and a degree

model and is fit in a similar manner (SI 3). This dual approach has been shown to have a high

capacity to tightly to diverse network structures (Rohr *et al.* 2016).

Stochastic Block Model

Ecological networks, especially antagonistic networks, frequently show comparatively discrete

compartments or modularisation, where subsets of species interact more strongly within their group

than with outsiders (Olesen *et al.* 2007, 2008; Allesina & Pascual 2009; Schleuning *et al.* 2014). This

may reflect fundamental incompatibilities in physiological traits, temporal mismatch (such as

nocturnal/diurnal partitioning) or spatial segregation.

This grouping can be represented by stochastic block models (SBMs). These have been shown to

perform well on ecological data in comparison to other clustering algorithms (Leger *et al.* 2015) and

are seeing increased use in the ecological literature (Sander *et al.* 2015; Kéfi *et al.* 2016). In an SBM,

each species is assigned to a group in a defined set. The probability of interaction between two

species is defined based on their group membership. We find optimal group assignations and fit the

model using a degree-corrected bipartite-SBM specific algorithm (Larremore *et al.* 2014, SI 3).

Combinations of Models

The aforementioned models can be combined into ensembles to harness multiple sources of

information and structure. For example, if a generalist species was also comparatively poorly

sampled, this could indicate even more strongly the location of likely missing links. Probabilities can

be combined by either averaging or multiplying with distinct consequences. Multiplying emphasises

the extreme probabilities each model finds, while averaging highlights interactions that are

consistently identified by multiple models. Here we test ensembling the Matching-Centrality model

with an SBM, and each of the structural models with the coverage-deficit model, either averaged or

multiplied. Before combining models we standardise each set of probabilities assigned to

unobserved interactions to sum to 1.

Testing Predictions

Datasets

Testing the efficacy of extrapolations requires knowledge of the 'true' network. Since the efficacy of

the method will depend on the network structure we take two complimentary approaches to

defining our 'true' networks. First, we generate a set of simulated networks that vary over a wide

range of network properties. Second, we use a large and diverse set of empirical networks from

literature sources. In both cases 'observed' networks were generated by taking draws from a

multinomial distribution parameterised by the 'true' interaction frequency. This replicates ecological

sampling processes where the most frequent interactions are highly likely to have been observed

and the rarest missed.

We initially generated 2000 simulated networks from a probabilistic two-trait niche model (modified

from Fründ _et al._ 2016) described in full in SI 1. Key network properties were calculated using the

bipartite R package (Dormann _et al._ 2008): connectance, weighted nestedness (Galeano _et al._ 2009),

cluster coefficient (Watts & Strogatz 1998), distribution of interaction frequency (Shannon diversity,

Bersier _et al._ 2002) and network level specificity H2' (Blüthgen _et al._ 2006). Distributions of these

metrics and the (low) correlations between them are depicted in SI 1.

Our objective was to generate interaction matrices that represent a wide range of ecologically

plausible bipartite networks. However, models used to generate plausible interaction networks have

many similarities to the predictive models described above. This is for a good reason –models are

chosen because they are thought to parsimoniously represent ecological networks. To reduce

circularity, our generative model is considerably more complex than our predictive models and we

examine how predictive model performance changes in response to network properties.

Using the 'true' networks we took 300-2000 samples per network to generate our 'observed'

networks. We discarded occasional cases where all true interactions were observed. This resulted in

a low mean true sample coverage deficit of 2.4%, corresponding to an average network

completeness of 64% due to skewed interaction strength distributions. Our systems therefore

represent reasonably 'well-sampled' systems that are nonetheless missing a large proportion of the

interactions present.

We collated a diverse set of 113 empirical networks representing antagonistic, mutualistic and

commensalistic interaction types (SI 2). We collated quantitative single-class bipartite networks from

the *Web of Life* (www.web-of-life.es) repository with 30-200 species and 40-431 observed

interactions. This included 48 plant-pollinator (19 sources), 23 mammal-flea (one source, but across

a wide geographic range), 2 plant-ant mutualism (2 sources) and 10 seed disperser (from 7 sources)

networks. We supplemented this with 25 mammal-dung beetle interaction networks (Frank *et al.*

2018, selecting those with ≥30 total species, ≥40 observed interactions, and ≥5 mammal species) and

5 host-parasitoid networks (Tylianakis *et al.* 2007).

These networks are not exhaustively sampled (SI 2). Nonetheless, we took the empirical interaction

frequency to be the 'true' network to define a multinomial distribution. We subsampled each

empirical network 20 times, each time taking a different fraction between 0.2 and 1 of the original

sample size, taken to be the sum of the observations. We excluded cases where the observed web

included more than 95% of the 'true' interactions.

Testing model capacity

Using each model or combination of models we assign each unobserved interaction a relative

probability of being a missing link. We assume that all observations are true positives (i.e. there have

been no misidentifications) although in principle similar methods could be used to identify

candidates for re-evaluation. Where a species in the 'true' network was not included in the

'observed' network, we did not attempt to predict its interactions and reduced the size of the 'true'

network correspondingly for all purposes.

We assess performance at identifying missing links with the area under receiver operating

characteristic curve (AUC) metric to assess the information content of a signal. AUC can be

considered the chance that a 'false negative' (missing link) was assigned a higher relative probability

than a 'true negative'. A value of 0.5 indicates no useful signal while a value of 1 indicates perfect

discrimination. To determine if $P(a_{ij} > 0)$ is reflective of true relative interaction frequency, we

calculated the mean Spearman's rank-correlation with the 'true' relative interaction strengths of the

unobserved interactions for each model.

We used the *EstimateR* function in the *vegan* R package (Oksanen *et al.* 2018) to estimate the

number of missing links based on the total distribution of observed interaction counts using the bias-

corrected Chao1 index. We then augmented the observed interaction networks by adding the

estimated number of the most probable missing links to the network based on the average of the

SBM, matching-centrality and coverage-deficit models. Each selected missing link was taken to be

'observed' once and network-level metrics calculated for the observed, augmented and true

networks. We tested three qualitative network metrics that had been identified (Fründ *et al.* 2016)

as being susceptible to misestimation in under-sampled networks (connectance, nestedness and

niche overlap), alongside two metrics considered to be comparatively resistant to under-sampling

(H2' and weighted NODF nestedness).

<u>Results</u>

Locating missing links in simulated networks

In general, the network models showed impressive performance at identifying likely missing links.

The overall best performing model was the stochastic block model, closely followed by the other

three network models (Figure 2a). The coverage deficit model lagged behind structural models, but

was more informative than simply ranking focal species by sample size. Combinations of models

tended to perform better than individual models on the simulated data (Table 1), with the overall

best performing model averaging the stochastic block model, the matching-centrality model and the

coverage deficit model. However, the marginal improvements from combining models were often

small.

Figure 2. a) Model performance across 1985 simulated networks. Mean differences between the statistical models were

small, although given the large sample size all pairwise differences were statistically significant (Holm-corrected t-tests,

paired by network ID, p<0.0001). **b)** Model performance on the empirical data sets, subdivided by network class. Each

empirical network was subsampled 20 times to different fractions of the original sample size. Models perform differently

on different classes of data (Model comparison, accounting for dataset ID as a random effect, shows model:class

interaction term is significant: p< 0.0001, n_{obs} = 12672). In all cases boxes show first and third quartiles.

Model	Simulated Dataset			Empirical Dataset		
		$+C_{def}$ (Multiplied)	$+C_{def}$ (Averaged)		$+C_{def}$ (Multiplied)	$+C_{def}$ (Averaged)
Focal Layer Sampling						
Coverage-Deficit	0.600			0.589[abc]		
Observation Count	0.580			0.603[de]		
Network Models						
Latent-Trait	0.660 [a]	0.649	0.654 [a]	0.631[f]	0.588[abc]	0.590[abc]
Degree	0.676	0.708[efg]	0.683 [b]	0.721 [i]	0.651[h]	0.663
Matching-Centrality	0.671	0.702[de]	0.684 [b]	0.718 [i]	0.642[g]	0.658
SBM	0.691 [bcd]	0.689 [bc]	0.695 [cd]	0.630[f]	0.584[a]	0.590[c]
Combined Models						
SBM and Matching-Centrality (Multiplied)	0.702 [de]	0.705[ef]	0.711[f]	0.647 [gh]	0.594[c]	0.604[de]
SBM and Matching-Centrality (Averaged)	0.705[ef]	0.709[f]	0.713 [g]	0.648 [gh]	0.600[d]	0.603[d]

|

Table 1 Mean AUC values using different predictive models or combinations of models across the whole

simulated (n= 1985) and empirical (n=113, 2112 subsamples) network sets. Incorporating coverage deficit

tends to lead to modest improvements in the simulated networks but not in the empirical. Sets of results not

found to be significantly different (Holm-correct t-tests, paired by network ID) are indicated with grouping

letters, separate for simulated and empirical results.

Model performance and relative hierarchy was affected by the underlying network structure (Figure

3). On average, missing links were identified best in nested, unclustered networks where

interactions were evenly distributed. An SBM was generally either as-good-as, or better than, the

other models across the networks, but particularly at large matrix size, low connectance, low (or

negative) nestedness. The coverage deficit and focal layer sample size models were the least

responsive to network structure, since our simulated network generator did not result in a

pronounced abundance-generality correlation. Average model performance did not appreciably

change with increased sample size, overall interaction coverage deficit or samples per host (Figure

S4). However, with an increased fraction of observed interactions, the structural models increased

their performance, while the coverage deficit model performance remained constant (Figure S4).

Figure 3. Response of model performance to 'true' network structure. Smoothed line shows a GAM fit for each

predictive model. Lower black line shows the local density distribution of that property across the simulated

test dataset.

Locating missing links in empirical networks

The performance of the different models varied considerably between the different classes of

networks (Figure 2b). In line with our expectations, the degree model performed best in the

mutualistic networks, while the latent-trait model showed its best performance on antagonistic

networks. In all six network classes the matching-centrality model performed the best on average.

The 'true network' properties (Fig S3) or the degree of sampling rarefaction (Fig S4) had relatively

little effect on the predictive performance. An exception was networks that show 'anti-nestedness' -

in these cases the predictive capacity of the centrality model was low, although the matching-

centrality model maintained its advantage.

The coverage-deficit model was very poor at predicting missing links in the empirical data and

combining it with the network-structure models lead to a *reduction* in overall predictive power

(Table 1). Ranking by the focal species observation count performed better overall.

Identifying frequency and number of missing interactions

Correlations between probabilities assigned to missing links ($P(a_{ij} > 0)$) and their 'true' interaction

frequency were stronger in the simulated dataset than the empirical (Table S2). Of the structural

models, the degree-model was on average the most strongly positively correlated with interaction

strength (ρ_{Sim} = 0.51, ρ_{Emp} = 0.15), while the SBM was the least (ρ_{Sim} = 0.28, ρ_{Emp} = 0.06). The

coverage deficit model was negatively correlated with true frequency (ρ_{Sim} = -0.35, ρ_{Emp} = -0.05)

i.e. missing links successfully identified due to poor sampling tended to be infrequent.

The Chao1 estimate of the number of missing interactions appeared to perform well on the

empirical networks but consistently underestimated the true interaction count and coverage deficit

in the simulated data (Figure 4).

Figure 4. Relationships between **a)** estimated sample coverage and 'true' sample coverage and **b)** estimated

number of missing interactions and the 'true' number of missed interactions. Results are shown using the bias-

corrected Chao1 estimator, the ACE estimator had very similar results. Line shows a 1:1 relationship that an

unbiased metric would follow.

Improving estimates of network metrics

The 'augmented' networks showed increased correlation, reduced bias and reduced error in

qualitative metrics, compared to the original observed networks (Figure 5a). The quantitative

network metrics (H2'and weighted NODF, Figure 5b) were already well estimated by the under-

sampled network, and the inclusion of additional interactions made little difference.

Figure 5. Relationship between network metrics calculated on the observed (black) and 'augmented' (red) interaction

343 networks to the true network value in the simulated networks. Line shows 1:1 relationship, Pearson's correlation

344 coefficient (r) shown for each metric. Qualitative metrics (a) show notable improvements in correlation with the true

345 values while quantitative (b) networks are largely unchanged.

Discussion

We have shown that is possible to infer missing links in range of ecological networks with a

reasonable degree of accuracy and this approach can improve estimates of network metrics. The

challenge now is to put these tools into practice to leverage additional ecological insight.

Choosing between predictive models

Ideally, a model used to infer missed interactions would capture the true ecological drivers, but this

is not essential for all purposes. Given the diversity of ecological networks, there will never be a

single 'best' model and victors in comparisons will be dependent on the data set. In our simulated

data it would appear that a stochastic block model had the best individual performance. In our

empirical datasets, the matching-centrality model performed consistently well (as found by Rohr *et*

al. 2016). However, we have reason to be distrustful of the empirical datasets (discussed below). The

pronounced, likely artefactual, abundance-generality relationship will favour degree-models.

Nonetheless, we would argue that splitting hairs over the best performing model is not necessarily

the most productive route. Networks will be structured by multiple processes and we note that in

our simulated data sets that the very best performance comes from the combination of different

models. These identify different missing interactions – structure-based models pick out the more

frequent interactions while coverage deficit models highlight comparatively infrequent interactions

that would be the hardest to determine through further undirected sampling. Future progress will

come from operationalising estimated missing links, rather than from further incremental model

refinements.

Distinctions between empirical and simulated networks

The empirical and simulated datasets substantially overlapped in key network metrics. The main

relevant difference between our datasets is increased correlation in empirical networks between

species marginal totals and generality (mean Spearman's ρ_{Sim} = 0.43, ρ_{Emp} = 0.76). This can account

for the poor performance of the coverage-deficit model in these cases. Disentangling the extent to

which apparent specialism of rarely observed species is a sampling artefact is challenging (Blüthgen

et al. 2006), since there are ecological explanations for a relationship (Fort *et al.* 2016; Dormann *et*

al. 2017; Simmons *et al.* 2019). For our purposes here, this likely bias in the structure of empirical

ecological networks has the consequence that predictive models may identify both missing links

introduced by our subsampling procedure and missing links due to gaps in the original dataset. We

would therefore place more weight on the results on the simulated data, while noting that, despite

the obstacles, the structural models are still able to perform reasonably well on the sparse empirical

data.

Estimating the quality of sampling and the number of missing interactions

Sample coverage estimators offer promise for estimating the number of missed interactions (Chacoff

et al. 2012; Traveset *et al.* 2015; Jordano 2016; MacGregor *et al.* 2017). The distinction between

sample-coverage and network completeness is worth emphasising. Apparently high values of sample

coverage alone should not be used as a simple proxy for 'network completeness', although they are

related to the reliability of certain quantitative network metrics (Henriksen *et al.* 2019). We found

the whole-network-level Chao1 estimate of missed interactions to be strongly correlated with the

'true' number, but notably underestimated the true number of missing links in our simulated

networks, a pattern also observed by Fründ *et al.* (2016). Although there appeared to be no

systematic bias in estimated network completeness in the empirical datasets, these 'true' networks

are themselves estimated to be significantly incomplete, suggesting an overall underestimation here

also. Our focus here was on the location, not the number of missing links. Future work should

investigate how long tails of infrequent interactions relate to this underestimation and consider

whether species rank abundance distribution estimators (Chao *et al.* 2014) can be usefully

generalised to assigning a frequency distribution to estimated missing links.

The way forward

Most of the models used here, and indeed our fitting approaches, likely have further room for

refinement, and there are many further classes of predictive models that we did not test, such as

nearest neighbour algorithms (Desjardins-Proulx *et al.* 2017). Furthermore, we based our models

entirely on the information contained in the network itself. In most cases, ecologists constructing a

network have access to (sometimes considerable) additional information and expertise on the study

system. Introducing pre-existing knowledge about likely or forbidden interactions, or pooled

knowledge from multiple sample sites offer considerable opportunities to maximise the information

in a network (Gray *et al.* 2015; Pomeranz *et al.* 2018). External information would be particularly

useful in separating 'forbidden links' (Olesen *et al.* 2011) that never occur due to fundamental

incompatibilities from those that occur elsewhere but would not within those spatio-temporal

boundaries. For instance, species may not be observed to interact because of a fundamental

mismatch in body-sizes (a forbidden link) or because they interact at night but data was only

gathered in the day.

Because the naïve models used here do not use external information, or the judgment of the

ecologists constructing the network, the quality of the estimation very much represents a lower

bound of what is possible with the approach in practice. Due to the underlying idiosyncrasies of

ecological systems, often euphemistically labelled 'species identity effects', there will always be an

upper limit to the predictive capacity of any statistical model. We suggest, rather than developing

new models to gain additional marginal improvements in predictive capacity, focus instead needs to

switch to developing frameworks to make use of this information to test the robustness of

conclusions derived from ecological networks.

Complete network inventories are not a realistic or necessarily useful goal (Jordano 2016). Enormous

sample sizes and strictly deliminated boundaries are required to exhaustively sample all interactions

(Martinez *et al.* 1999). There is undoubtedly a long tail of extremely infrequent interactions that will

never be observed in a realistic sampling regime. Furthermore, there is rarely a clear line defining

network boundaries (Poisot *et al.* 2015). Ecological networks are a constantly moving target

(Rasmussen *et al.* 2013; CaraDonna *et al.* 2017) and there may well be little value in including certain

rare, occasional visitors and unusual interactions. It can be challenging to determine if infrequent

interactions are simply a curiosity (Mercer 1966) or may have significant consequences (Dudley *et al.*

2016). The utility of increasing the accuracy with which one type of interaction is understood will

plateau as the impact of other interaction types and ecological guilds predominate (Lafferty *et al.*

2008). Despite the challenges, we see two principle uses for inferred missing links:

Firstly, directing further sampling where the goal is a descriptive network. In many cases the

topography of the network is of principle interest, given the potential for interaction strengths to

vary through time and in response to perturbation. For example, when tracing the relationships of

species that may be the target for eradication measures, such as disease-vector mosquitoes (Collins

et al. 2019), a critical first step is an inventory of interactions in the community. Since in most cases

investment in sampling is a finite resource, directing sampling towards high-likelihood interactions

could more efficiently increase the overall network completeness. However, we caution that

directed sampling will introduce biases in the generated network (Gibson *et al.* 2011; Falcão *et al.*

2016) and may hinder cross-community comparability (Poisot *et al.* 2012a). In certain cases

independently sampled networks, even if less complete, may be more useful than comparisons

between 'augmented' networks. The use of information from other networks may lead them to

appear more homogenous than they really are and lead to underestimations of interaction β-

diversity through time and space (Poisot *et al.* 2015; Graham & Weinstein 2018).

Secondly, the identification of probable missing links can allow better testing of the reliability of

conclusions drawn from networks, potentially through explicitly probabilistic networks (Poisot et al.

2016a). Statistical resampling and rarefaction procedures that can only work backwards or adding

random interactions may both introduce biases. Whilst it is established that certain quantitative

metrics are relatively robust to under-sampling (Blüthgen *et al.* 2008; Fründ *et al.* 2016; Henriksen *et*

al. 2019), these metrics may not align with the goals of the study. If the conclusions are robust to the

inclusion of probable missing links then confidence in the results would be increased. Conversely, if

the inclusion of most-likely missing links erased the differences this would signal that the conclusions

are less robust than hoped. In these cases, care should be taken when selecting a predictive model

to avoid a clash with the properties under investigation.

Conclusion

The identification of likely missing links is a powerful tool to include in the arsenal of network

ecologists. We have shown that they can perform well across a wide range of network types.

Importantly, inferential models need not be technically challenging to implement – for example the

cassandRa package allows all the models discussed here to be fit with a single function call. We hope

that such approaches become more widely used in guiding ecological sampling and testing the

robustness of conclusions drawn from bipartite networks.

Acknowledgments

Thanks to the CERO group for comments on this project. Funding was provided by NERC grant

NE/N010221/1.

References

1.de Aguiar, M.A.M. (2018). Revealing biases in the sampling of ecological interaction networks. *bioRxiv*

2.Allesina, S. & Pascual, M. (2009). Food web models: a plea for groups. *Ecol. Lett.*, 12, 652–662

3.Bartomeus, I. (2013). Understanding Linkage Rules in Plant-Pollinator Networks by Using Hierarchical Models

That Incorporate Pollinator Detectability and Plant Traits. *PLoS One*, 8, 1–8

4.Bartomeus, I., Gravel, D., Tylianakis, J.M., Aizen, M.A., Dickie, I.A. & Bernard-Verdier, M. (2016). A common

framework for identifying linkage rules across different types of interactions. *Funct. Ecol.*, 30, 1894–1903

5.Bascompte, J. & Jordano, P. (2013). *Mutualistic Networks*. Princeton University Press, Princeton

6.Bascompte, J., Jordano, P., Melian, C.J. & Olesen, J.M. (2003). The nested assembly of plant-animal

mutualistic networks. *Proc. Natl. Acad. Sci.*, 100, 9383–9387

7.Berlow, E.L. (1999). Strong effects of weak interactions in ecological communities. *Nature*, 398, 330–334

8.Bersier, L.-F., Banašek-Richter, C. & Cattin, M.-F. (2002). Quantitative descriptors of food web matrices.

Ecology, 83, 2917–2926

9.Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: A critique and an

ecologist's guide. *Basic Appl. Ecol.*, 11, 185–195

10.Blüthgen, N., Fründ, J., Vázquez, D.P. & Menzel, F. (2008). What Do Interaction Network Metrics Tell Us

about Specialization and Biological Traits. *Ecology*, 89, 3387–3399

11.Blüthgen, N., Menzel, F. & Blüthgen, N. (2006). Measuring specialization in species interaction networks.

BMC Ecol., 6, 9

12.CaraDonna, P.J., Petry, W.K., Brennan, R.M., Cunningham, J.L., Bronstein, J.L., Waser, N.M., *et al.* (2017).

Interaction rewiring and the rapid turnover of plant–pollinator networks. *Ecol. Lett.*, 20, 385–394

13.Chacoff, N.P., Vázquez, D.P., Lomáscolo, S.B., Stevani, E.L., Dorado, J. & Padrón, B. (2012). Evaluating

sampling completeness in a desert plant-pollinator network. *J. Anim. Ecol.*, 81, 190–200

14.Chao, A. (1984). Nonparametric estimation of the number of classes in a population. *Scand. J. Stat.*, 11,

265–270

15.Chao, A. & Colwell, R.K. (2017). Thirty years of progeny from Chao's inequality: Estimating and comparing

richness with incidence data and incomplete sampling. *Stat. Oper. Res. Trans.*, 41, 3–54

16.Chao, A., Hsieh, T.C., Chazdon, R.L. & Colwell, R.K. (2014). Unveiling the Species-Rank Abundance

Distribution by Generalizing Good-Turing Sample Coverage Theory. *Ecology*, 96, 1189–1201

17.Clauset, A., Moore, C. & Newman, M.E.J. (2008). Hierarchical structure and the prediction of missing links in

networks. *Nature*, 453, 98–101

18.Collins, C.M., Bonds, J.A.S., Quinlan, M.M. & Mumford, J.D. (2019). Effects of the removal or reduction in

density of the malaria mosquito, Anopheles gambiae s.l ., on interacting predators and competitors in local

ecosystems. *Med. Vet. Entomol.*, 33, 1–15

19.Colwell, R.K. & Coddington, J.A. (1994). Estimating terrestrial biodiversity through extrapolation. *Philos.*

Trans. R. Soc. B-Biological Sci., 345, 101–118

20.Costa, J.M., da Silva, L.P., Ramos, J.A. & Heleno, R.H. (2016). Sampling completeness in seed dispersal

networks: When enough is enough. *Basic Appl. Ecol.*, 17, 155–164

21.Crea, C., Ali, R.A. & Rader, R. (2016). A new model for ecological networks using species-level traits.

Methods Ecol. Evol., 7, 232–241

22.Delmas, E., Besson, M., Brice, M.-H., Burkle, L.A., Dalla Riva, G. V., Fortin, M.-J., *et al.* (2019). Analysing

ecological networks of species interactions. *Biol. Rev.*, 94, 16–36

23.Desjardins-Proulx, P., Laigle, I., Poisot, T. & Gravel, D. (2017). Ecological interactions and the Netflix

problem. *PeerJ*, 5, e3644

24.Dormann, C.F., Frund, J. & Martin Schaefer, H. (2017). Identifying Causes of Patterns in Ecological Networks:

Opportunities and Limitations. *Annu. Rev. Ecol. Evol. Syst*, 48, 559–84

25.Dormann, C.F., Gruber, B. & Frund, J. (2008). Introducing the bipartite Package: Analysing Ecological

Networks. *R News*, 8, 8–11

26.Dudley, J.P., Hang'Ombe, B.M., Leendertz, F.H., Dorward, L.J., de Castro, J., Subalusky, A.L., *et al.* (2016).

Carnivory in the common hippopotamus Hippopotamus amphibius: implications for the ecology and

epidemiology of anthrax in African landscapes. *Mamm. Rev.*, 46, 191–203

27.Dunne, J.A. & Pascual, M. (2006). *Ecological Networks: Linking Structure to Dynamics in Food Webs*. Oxford

University Press, Oxford

28.Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N.P., *et al.* (2013). The dimensionality of

ecological networks. *Ecol. Lett.*, 16, 577–583

29.Falcão, J.C.F., Dáttilo, W. & Rico-Gray, V. (2016). Sampling effort differences can lead to biased conclusions

on the architecture of ant-plant interaction networks. *Ecol. Complex.*, 25, 44–52

30.Fort, H., Vázquez, D.P. & Lan, B.L. (2016). Abundance and generalisation in mutualistic networks: Solving

the chicken-and-egg dilemma. *Ecol. Lett.*, 19, 4–11

31.Frank, K., Krell, F.-T., Slade, E.M., Raine, E.H., Chiew, L.Y., Schmitt, T., *et al.* (2018). Global dung webs: high

trophic generalism of dung beetles along the latitudinal diversity gradient. *Ecol. Lett.*, 21, 1229–1236

32.Fründ, J., Mccann, K.S. & Williams, N.M. (2016). Sampling bias is a challenge for quantifying specialization

and network structure: Lessons from a quantitative niche model. *Oikos*, 125, 502–513

33.Galeano, J., Pastor, J.M. & Iriondo, J.M. (2009). Weighted-Interaction Nestedness Estimator (WINE): A new

estimator to calculate over frequency matrices. *Environ. Model. Softw.*, 24, 1342–1346

34.Gellner, G. & McCann, K.S. (2016). Consistent role of weak and strong interactions in high- and low-diversity

trophic food webs. *Nat. Commun.*, 7, 11180

35.Gibson, R.H., Knott, B., Eberlein, T. & Memmott, J. (2011). Sampling method influences the structure of

plant-pollinator networks. *Oikos*, 120, 822–831

36.Goldwasser, L. & Roughgarden, J. (1997). Sampling Effects and the Estimation of Food-Web Properties.

Ecology, 78, 41–54

37.González-Varo, J.P. & Traveset, A. (2016). The Labile Limits of Forbidden Interactions. *Trends Ecol. Evol.*, 31,

700–710

38.Gotelli, N.J. & Colwell, R.K. (2010). Estimating species richness. In: *Biological Diversity: Frontiers in*

Measurement and Assessment (eds. Magurran, A. & McGill, B.J.). Open University Press, pp. 416–422

39.Graham, C.H. & Weinstein, B.G. (2018). Towards a predictive model of species interaction beta diversity.

Ecol. Lett., 21, 1299–1310

40.Gray, C., Baird, D.J., Baumgartner, S., Jacob, U., Jenkins, G.B., O'Gorman, E.J., *et al.* (2014). Ecological

networks: the missing links in biomonitoring science. *J. Appl. Ecol.,* 51, 1444–1449

41.Gray, C., Figueroa, D.H., Hudson, L.N., Ma, A., Perkins, D. & Woodward, G. (2015). Joining the dots: An

automated method for constructing food webs from compendia of published interactions. *Food Webs,* 5, 11–

20

42.Guimera, R. & Sales-Pardo, M. (2009). Missing and spurious interactions and the reconstruction of complex

networks. *Proc. Natl. Acad. Sci.,* 106, 22073–22078

43.Henriksen, M. V., Chapple, D.G., Chown, S.L. & McGeoch, M.A. (2019). The effect of network size and

sampling completeness in depauperate networks. *J. Anim. Ecol.,* 88, 211–222

44.Jacquet, C., Moritz, C., Morissette, L., Legagneux, P., Massol, F., Archambault, P., *et al.* (2016). No

complexity–stability relationship in empirical ecosystems. *Nat. Commun.,* 7, 12573

45.Jordano, P. (2016). Sampling networks of ecological interactions. *Funct. Ecol.,* 30, 1883–1893

46.Jordano, P., Bascompte, J. & Olesen, J.M. (2003). Invariant properties in coevolutionary networks of plant-

animal interactions. *Ecol. Lett.,* 6, 69–81

47.Kadoya, T., Gellner, G. & McCann, K.S. (2018). Potential oscillators and keystone modules in food webs.

Ecol. Lett., 21, 1330–1340

48.Kéfi, S., Miele, V., Wieters, E.A., Navarrete, S.A. & Berlow, E.L. (2016). How Structured Is the Entangled

Bank? The Surprisingly Simple Organization of Multiplex Ecological Networks Leads to Increased Persistence

and Resilience. *PLOS Biol.*, 14, e1002527

49.Klumpers, S.G.T., Stang, M. & Klinkhamer, P.G.L. (2019). Foraging efficiency and size matching in a plant-

pollinator community: the importance of sugar content and tongue length. *Ecol. Lett.*, 469–479

50.Lafferty, K.D., Allesina, S., Arim, M., Briggs, C.J., De Leo, G., Dobson, A.P., *et al.* (2008). Parasites in food

webs: the ultimate missing links. *Ecol. Lett.*, 11, 533–546

51.Laigle, I., Aubin, I., Digel, C., Brose, U., Boulangeat, I. & Gravel, D. (2018). Species traits as drivers of food

web structure. *Oikos*, 127, 316–326

52.Larremore, D.B., Clauset, A. & Jacobs, A.Z. (2014). Efficiently inferring community structure in bipartite

networks. *Phys. Rev. E*, 90, 012805

53.Lee, R.H. & Guénard, B. (2019). Choices of sampling method bias functional components estimation and

ability to discriminate assembly mechanisms. *Methods Ecol. Evol.*, 2019, 1–12

54.Leger, J.-B., Daudin, J.-J. & Vacher, C. (2015). Clustering methods differ in their ability to detect patterns in

ecological networks. *Methods Ecol. Evol.*, 6, 474–481

55.MacGregor, C.J., Evans, D.M. & Pocock, M.J.O. (2017). Estimating sampling completeness of interactions in

quantitative bipartite ecological networks: incorporating variation in species specialisation. *bioRxiv*, 195917

56.Martinez, N.D., Hawkins, B.A., Dawah, H.A. & Feifarek, B.P. (1999). Effects of Sampling Effort on

Characterization of Food-Web Structure. *Ecology*, 80, 1044–1055

57.McCann, K., Hastings, A. & Huxel, G.R. (1998). Weak trophic interactions and the balance of nature. *Nature,*

395, 794–798

58.Memmott, J. (2009). Food webs: a ladder for picking strawberries or a practical tool for practical problems?

Philos. Trans. R. Soc. B Biol. Sci., 364, 1693–1699

59.Mercer, A.J. (1966). Turnstones feeding on human corpse. *Br. Birds*, 59, 307

60.Montoya, J.M., Pimm, S.L. & Solé, R. V. (2006). Ecological networks and their fragility. *Nature*, 442, 259–64

61.Morales-Castilla, I., Matias, M.G., Gravel, D. & Araújo, M.B. (2015). Inferring biotic interactions from

proxies. *Trends Ecol. Evol.*, 30, 347–356

62.Morris, R.J., Gripenberg, S., Lewis, O.T. & Roslin, T. (2014). Antagonistic interaction networks are structured

independently of latitude and host guild. *Ecol. Lett.*, 17, 340–349

63.Nielsen, A. & Bascompte, J. (2007). Ecological networks, nestedness and sampling effort. *J. Ecol.*, 95, 1134–

1141

64.Novotny, V., Miller, S.E., Baje, L., Balagawi, S., Basset, Y., Cizek, L., *et al.* (2010). Guild-specific patterns of

species richness and host specialization in plant-herbivore food webs from a tropical forest. *J. Anim. Ecol.*, 79,

1193–203

65.Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., *et al.* (2018). vegan:

Community Ecology Package

66.Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H., Rasmussen, C. & Jordano, P. (2011). Missing and

forbidden links in mutualistic networks. *Proc. R. Soc. B Biol. Sci.*, 278, 725–732

67. Olesen, J.M., Bascompte, J., Dupont, Y.L. & Jordano, P. (2007). The modularity of pollination networks. *Proc.*

Natl. Acad. Sci., 104, 19891–19896

68. Olesen, J.M., Bascompte, J., Elberling, H. & Jordano, P. (2008). Temporal dynamics in a pollination network.

Ecology, 89, 1573–1582

69. Paine, R.T. (1980). Food Webs: Linkage, Interaction Strength and Comunity Infrastructure. *J. Anim. Ecol.*, 49,

666–685

70. Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012a). The dissimilarity of species interaction

networks. *Ecol. Lett.*, 15, 1353–1361

71. Poisot, T., Canard, E., Mouquet, N. & Hochberg, M.E. (2012b). A comparative study of ecological

specialization estimators. *Methods Ecol. Evol.*, 3, 537–544

72. Poisot, T., Cirtwill, A.R., Cazelles, K., Gravel, D., Fortin, M.-J. & Stouffer, D.B. (2016a). The structure of

probabilistic networks. *Methods Ecol. Evol.*, 7, 303–312

73. Poisot, T., Stouffer, D.B. & Gravel, D. (2015). Beyond species: why ecological interaction networks vary

through space and time. *Oikos*, 124, 243–251

74. Poisot, T., Stouffer, D.B. & Kéfi, S. (2016b). Describe, understand and predict: why do we need networks in

ecology? *Funct. Ecol.*, 30, 1878–1882

75. Pomeranz, J.P.F., Thompson, R.M., Poisot, T. & Harding, J.S. (2018). Inferring predator–prey interactions in

food webs. *Methods Ecol. Evol.*, 2018, 1–12

76. Rasmussen, C., Dupont, Y.L., Mosbacher, J.B., Trøjelsgaard, K. & Olesen, J.M. (2013). Strong Impact of

Temporal Resolution on the Structure of an Ecological Network. *PLoS One*, 8, e81694

77.Rivera-Hutinel, A., Bustamante, R.O., Marín, V.H. & Medel, R. (2012). Effects of sampling completeness on

the structure of plant-pollinator networks. *Ecology*, 93, 1593–1603

78.Rohr, R.P., Naisbit, R.E., Mazza, C. & Bersier, L.-F. (2016). Matching–centrality decomposition and the

forecasting of new links in networks. *Proc. R. Soc. B Biol. Sci.*, 283, 20152702

79.Rohr, R.P., Naisbit, R.E., Mazza, C. & Bersier, L.-F. (2018). Statistical Approaches for Inferring and Predicting

Food-Web Architecture. In: *Adaptive Food Webs, Stability and Transitions of Real and Model Ecosystems* (eds.

Moore, J.C., de Ruiter, P.C., McCann, K.S. & Wolters, V.). Cambridge University Press, pp. 178–192

80.Sander, E.L., Wootton, J.T. & Allesina, S. (2015). What Can Interaction Webs Tell Us About Species Roles?

PLoS Comput. Biol., 11, e1004330

81.Schleuning, M., Ingmann, L., Strauß, R., Fritz, S.A., Dalsgaard, B., Matthias Dehling, D., *et al.* (2014).

Ecological, historical and evolutionary determinants of modularity in weighted seed-dispersal networks. *Ecol.*

Lett., 17, 454–463

82.Seo, E. & Hutchinson, R.A. (2018). Predicting Links in Plant-Pollinator Interaction Networks Using Latent

Factor Models With Implicit Feedback. *AAAI*

83.Simmons, B.I., Vizentin-Bugoni, J., Maruyama, P.K., Cotton, P.A., Marín-Gómez, O.H., Lara, C., *et al.* (2019).

Abundance drives broad patterns of generalisation in plant–hummingbird pollination networks. *Oikos*,

oik.06104

84.Song, C., Rohr, R.P. & Saavedra, S. (2017). Why are some plant–pollinator networks more nested than

others? *J. Anim. Ecol.*, 86, 1417–1424

85.Spitz, J., Ridoux, V. & Brind'Amour, A. (2014). Let's go beyond taxonomy in diet description: Testing a trait-

based approach to prey-predator relationships. *J. Anim. Ecol.*, 83, 1137–1148

86.Staniczenko, P.P.A., Kopp, J.C. & Allesina, S. (2013). The ghost of nestedness in ecological networks. *Nat.*

Commun., 4, 1391

87.Staniczenko, P.P.A., Lewis, O.T., Tylianakis, J.M., Albrecht, M., Coudrain, V., Klein, A.-M., *et al.* (2017).

Predicting the effect of habitat modification on networks of interacting species. *Nat. Commun.*, 8, 792

88.Stock, M., Poisot, T., Waegeman, W. & De Baets, B. (2017). Linear filtering reveals false negatives in species

interaction data. *Sci. Rep.*, 7, 45908

89.Terry, J.C.D. (2019). cassandRa v.0.1.0, *https://cran.r-project.org/package=cassandRa*

90.Terry, J.C.D., Morris, R.J. & Bonsall, M.B. (2017). Trophic interaction modifications: an empirical and

theoretical framework. *Ecol. Lett.*, 20, 1219–1230

91.Traveset, A., Olesen, J.M., Nogales, M., Vargas, P., Jaramillo, P., Antolín, E., *et al.* (2015). Bird–flower

visitation networks in the Galápagos unveil a widespread interaction release. *Nat. Commun.*, 6, 6376

92.Tylianakis, J.M., Laliberté, E., Nielsen, A. & Bascompte, J. (2010). Conservation of species interaction

networks. *Biol. Conserv.*, 143, 2270–2279

93.Tylianakis, J.M., Tscharntke, T. & Lewis, O.T. (2007). Habitat modification alters the structure of tropical

host-parasitoid food webs. *Nature*, 445, 202–5

94.Ulrich, W., Almeida-Neto, M. & Gotelli, N.J. (2009). A consumer's guide to nestedness analysis. *Oikos*, 118,

3–17

95.Valdovinos, F.S. (2019). Mutualistic networks: moving closer to a predictive theory. *Ecol. Lett.*

96.Valiente-Banuet, A., Aizen, M.A., Alcántara, J.M., Arroyo, J., Cocucci, A., Galetti, M., *et al.* (2015). Beyond

species loss: The extinction of ecological interactions in a changing world. *Funct. Ecol.*, 29, 299–307

97.Vázquez, D.P., Bluthgen, N., Cagnolo, L. & Chacoff, N.P. (2009). Uniting pattern and process in plant-animal

mutualistic networks: A review. *Ann. Bot.*, 103, 1445–1457

98.Vizentin-Bugoni, J., Maruyama, P.K., Debastiani, V.J., Duarte, L. da S., Dalsgaard, B. & Sazima, M. (2016).

Influences of sampling effort on detected patterns and structuring processes of a Neotropical plant-

hummingbird network. *J. Anim. Ecol.*, 85, 262–272

99.Vizentin-Bugoni, J., Maruyama, P.K. & Sazima, M. (2014). Processes entangling interactions in communities:

forbidden links

are more important than abundance in a hummingbird-plant network. *Proc. R. Soc. B Biol. Sci.*, 281, 20132397

100.Watts, D.J. & Strogatz, S.H. (1998). Collective dynamics of "small-world" networks. *Nature*, 393, 440–442

101.Weinstein, B.G. & Graham, C.H. (2017). On comparing traits and abundance for predicting species

interactions with imperfect detection. *Food Webs*, 11, 17–25