Finding missing links in interaction networks

J. Christopher D. Terry (1) james.terry@zoo.ox.ac.uk Corresponding author

Owen T. Lewis (1) owen.lewis@zoo.ox.ac.uk

1: Department of Zoology, University of Oxford, Oxford, OX1 3PS, UK.

Running title: Finding missing links in interaction networks

Abstract

Documenting which species interact within ecological communities is challenging and labour-

intensive. As a result, many interactions remain unrecorded, potentially distorting our

understanding of network structure and dynamics. We test the utility of four structural models

and a new coverage-deficit model for predicting missing links in both simulated and empirical

bipartite networks. We find they can perform well, but that the predictive power of structural

models varies with the underlying network structure. Predictions can be improved by ensembling

multiple models. Sample-coverage estimators of the number of missed interactions are highly

correlated with the number of missed interactions, but strongly biased towards underestimating

the true number of missing links. Augmenting observed networks with most-likely missing links

improves estimates of qualitative network metrics. Tools to identify likely missing links can be

simple to implement, allowing the prioritisation of research effort and more robust assessment of

network properties.

Keywords: ecological network, bipartite network, missing links, sampling completeness,

network metrics

Introduction

Networks documenting trophic, competitive or mutualistic interactions among species are an essential tool for understanding the structure and dynamics of ecological communities (Memmott 2009, Poisot et al. 2016b, Dormann et al. 2017, Delmas et al. 2019). Interaction networks can be a better indicator of the resilience of communities than species counts (Valiente-Banuet et al. 2015) and are increasingly applied in conservation (Tylianakis et al. 2010) and biomonitoring (Gray et al. 2014).

Documenting the species present in a system can be difficult enough (Gotelli and Colwell 2010); accurately enumerating their interactions is even more challenging. As a result, ecological interaction networks are routinely under-sampled, with consequences for our understanding of communities (Goldwasser and Roughgarden 1997, Blüthgen 2010, Chacoff et al. 2012, Jordano 2016). Ecological networks are sparse and hence any unobserved interactions could be either 'missing links' (interactions that would be observed with complete sampling), or true negatives. Differences or deficiencies in sampling can lead to divergent conclusions about underlying ecological mechanisms (Lee and Guénard 2019). Furthermore, under-sampling will skew the apparent prevalence of interactions towards frequent interactions and may miss infrequent interactions entirely (Dormann et al. 2017).

Common solutions to this problem have included rarefying networks to a consistent level of sampling (Morris et al. 2014), sampling until metric values stabilise (Rivera-Hutinel et al. 2012, Costa et al. 2016), explicit observation models (Weinstein and Graham 2017), and the use of network metrics that are comparatively robust to under sampling (Vizentin-Bugoni et al. 2016). There has been an extensive literature testing these different approaches (Nielsen and Bascompte 2007, Blüthgen et al. 2008, Poisot et al. 2012b, Fründ et al. 2016, Henriksen et al. 2019, de

Aguiar et al. 2019), often arguing for an emphasis on quantitative rather than qualitative metrics,

which reduce the influence of missed infrequent interactions.

Nonetheless, infrequent interactions may play an important role in ecological communities.

Since networks are not static, there is value in distinguishing infrequent interactions from those

that are genuinely absent. Rarely-observed links have the potential to grow in importance with

changes in biotic (Terry et al. 2017) or abiotic conditions (Staniczenko et al. 2017). At

equilibrium, interactions may occur at low frequency, yet be dynamically important. For

instance, while interactions involving a species suppressed to a low level by a consumer may be

observed infrequently, removal of the consumer can allow prey abundance to increase rapidly

(Paine 1980). At a network level, the prevalence and distribution of apparently weak or

infrequent interactions can have marked consequences for network stability and dynamics

(McCann et al. 1998, Berlow 1999, Gellner and McCann 2016, Jacquet et al. 2016, Kadoya et al.

2018).

Ecological networks are manifestly non-random (Dunne and Pascual 2006, Montoya et al. 2006,

Vázquez et al. 2009, Bascompte and Jordano 2013). In principle, it should be possible to use this

underlying structure to help infer where links are most likely to have been missed (Bartomeus

2013, Morales-Castilla et al. 2015, Valdovinos 2019). To this end, a number of missing link

inference approaches have been tested on ecological networks. These have included hierarchical

structuring models (Clauset et al. 2008), linear filtering (Stock et al. 2017), matching-centrality

models (Rohr et al. 2016), stochastic block models (Guimera and Sales-Pardo 2009) and *k*-

nearest neighbour recommenders (Desjardins-Proulx et al. 2017). These models have shown that

missing links can be inferred from partially-observed structured networks. However, although

ecological networks are used as part of benchmarking tests, often the inference scenarios are not

orientated towards ecological needs and to date there has been limited uptake of these methods

from the ecological community.

Furthermore, it is common in empirical ecological networks for species to have been sampled to

markedly different extents (although see Novotny *et al.* 2010). This has the potential to give

further information about the potential location of missing links, since missing links are more

likely for poorly-sampled species (Blüthgen et al. 2008).

Here, we explore the potential to identify missing links in a wide range of ecological bipartite

networks with a range of models, including a new species-level coverage-deficit approach which

incorporates information on the completeness of sampling for individual species. We show that

inferential models can perform well, especially when used in ensemble, and demonstrate how

they could inform empirical work to document ecological networks and help evaluate

hypotheses.

Methods

First we present a set of predictive models to detect missing links and introduce a suite of

simulated and empirical networks. These are used to explore each model's capacity to identify

missing links and improve the estimation of network-level metrics.

Predictive Models

Our predictive models seek to identify the true missing links within an observed bipartite

adjacency matrix $\mathbf{O}$ composed of observation counts o_{ij}. This has rows representing each focal

layer species i and columns detailing each potential interaction partner j. We assume that this set

of observations is derived from a set of true interaction frequencies, where the relative rate of

interaction between two species is given by a_{ij}. We use our statistical models to assign to each

unobserved interaction in a network a relative probability of being a 'missing link' (false

negative), rather than a true negative: $P(a_{ij} > 0 \mid \mathbf{O})$ for the cases where $\mathbf{O}_{ij} = 0$. We assume

throughout that the species observed (the dimensions of $\mathbf{O}$) are a complete list – we do not

attempt to predict the identity of interactions with unobserved species. Here we introduce and

detail a new 'coverage deficit' model and four structural models described in the literature. All

models were fit using the *cassandRa* R package (Terry 2019). Example predictions on a

simulated network of each of the core models are shown in Figure 1.

97   *Sample size and Coverage-Deficit Models*

Interactions are more likely to have been missed between species that are relatively poorly

sampled. When ecological bipartite networks are constructed, it is common that one level (the

focal level) of a network is sampled directly and the interaction partners in the other level

counted. Examples include gathering insect hosts and rearing parasitoids from them, or

observing pollinators that visit a set of plant species. This sampling may be opportunistic,

matched to the local abundance, or standardised in some other manner. The result is a set of

observed interaction counts with each focal layer species. Although interaction counts may be

later rescaled to better represent local abundances, here we assume the raw observation data are

available of discretisable relationships where individual interaction events can be observed. This

includes many ecological interactions such as parasitism, pollination visits and feeding

observations, but excludes diffuse facilitatory or competitive interactions. A simple approach

assigns a relative probability of missed interactions based solely on the number of observations

of each focal layer-species:

111
$$P(a_{ij} > 0 \mid \mathbf{O}) \sim \frac{1}{\sum_j \mathbf{O}_{ij}}.$$

We refer to this as our 'sample-size' model.

This approach can be refined using techniques from the extensive literature developed to assess

the sample completeness of species inventories using information about the most infrequent

observations to estimate the number of missing observations (Chao and Jost 2012). Rooted in

information theory, these approaches can be applied to any case of estimating sample coverage,

including interaction sample coverage (Chacoff et al. 2012, Jordano 2016). The core of our

coverage-deficit model estimates the interaction partner sample completeness for each focal

layer species, based on the observed occurrence frequencies and the 'Chao1' estimator (Chao

1984, Chao & Jost, 2012). Where data is sufficient, this coverage deficit estimator ($\hat{C}_{def}$) is an

estimate of the proportion of interactions that occur between as-yet-unobserved partners, or

alternatively the probability that the next interaction partner observed would not yet have been

observed interacting with that species. It is defined as:

$$\hat{C}_{def} = \frac{f_1}{n} \frac{f_1\,(n-1)}{f_1\,(n-1) + 2f_2}$$

124   where f_1 is the number of interactions observed only once, f_2 is the number of interactions

125   observed exactly twice, and n is the total observed interactions involving that focal species. For

126   each focal layer species i, we use a $\hat{C}_{def_i}$ estimate to assign a relative probability to each

127   potential missing interaction as:

128
$$P\big(a_{ij} > 0 \,|\, \mathbf{O}\big) \sim \frac{\hat{C}_{def_i}}{\sum_j (\mathbf{O}_{ij}=0)} .$$

Unfortunately, ecological interaction networks typically contain a number of focal layer species

that are extremely poorly observed, resulting in situations where the Chao1 estimator for

coverage deficit is undefined or inappropriate. For very poorly sampled species, the distribution

of interactions cannot itself provide much information (Colwell and Coddington 1994). Where

observations are exclusively singletons ($f_1 = n$) the index will estimate zero coverage. At the

other end of the scale, if a focal species is observed with no singleton interactions the index will

estimate that there is zero sample coverage deficit.

In these cases, we use a simple binomial model to estimate $\hat{C}_{def}$, the probability that the next

interactor drawn will not yet have been observed interacting with the focal species. The

likelihood of not having yet observed any of the unobserved missing links is: $L = \left(1 - \hat{C}_{def}\right)^n$.

While the maximum likelihood estimate of $\hat{C}_{def}$ will be 0, with a Bayesian approach and

assuming a flat prior the posterior mean can be found directly as: $\mathbb{E}(\hat{C}_{def}|n) = 1 - \frac{1}{n+2}$. We use

this alternative estimate for $\hat{C}_{def}$ when $n \leq 5$ (cut-off selected via accuracy tests on simulated

samples), all the observations are singletons ($f_1 = n$), or if there are no singletons or doubletons

($f_1 + f_2 = 0$).

Latent-Trait Model

Species do not interact at random with each other – similar species tend to interact with a

similar set of interaction partners. It should be possible to use this information to guide our

expectations of missing links through a trait-based framework (Morales-Castilla et al. 2015,

Bartomeus et al. 2016, Laigle et al. 2018). These similarities can be in directly-measured traits,

such as corolla and tongue length (Vizentin-Bugoni et al. 2014, Klumpers et al. 2019) or through

inferred 'latent' traits that are not directly measurable. Latent traits can be identified whether the

community is structured by phylogeny, microhabitat or physiological differences. A survey of

published networks found that the number of latent traits required may be quite low (Eklöf et al.

2013).

Intraspecific trait variation caused by spatial, temporal or ontogenetic differences can pose

considerable challenges (González-Varo and Traveset 2016). Nonetheless, such approaches have

had successes and a suite of approaches has been developed including matching models (Rohr et

al. 2016), Dirichlet-multinomial regression (Crea et al. 2016), matrix factorisation (Seo and

Hutchinson 2018) and fourth-corner analyses (Spitz et al. 2014). While it is possible to fit any

number of latent traits, here we explore use of a simple single latent trait model to reduce the

potential of overfitting (Rohr et al. 2016).

In this model, each focal layer species i is assigned a trait value m_i and each partner-layer

species j a value m_j. The probability of the interaction existing between each i and j is

determined from the difference in trait values through a logistic model:

$$\text{logit}\left(P\left(a_{ij} > 0 \mid \mathbf{O}\right)\right) = -\lambda \left(m_i - m_j\right)^2 + k_M .$$

where λ, k_M and the vector of trait parameters $\mathbf{m}$ are found through maximum likelihood

optimisation (details given in SI 3). We constrain λ to be positive, such that greater trait

differences between potential interactors reduce the probability of an interaction. k_M is an

intercept parameter to capture the average probability of an interaction between two species with

perfectly matching traits. We penalise strongly divergent traits by introducing a Cauchy-

distributed prior with mean 0 on $\mathbf{m}$.

Multiple trait distributions may have very similar likelihoods, especially with several specialised

interactions or where species bridge network components. To account for this, we optimise 10

differently initialised models and average the predictions of the five models with the greatest

likelihood. This model averaging approach is conceptually similar to, but computationally far

cheaper than, sampling from posterior distributions of trait vectors.

Degree Model

It is readily observable that some species have many interaction partners while other species

interact with very few. This manifests itself in skewed degree distributions within ecological

networks (Jordano et al. 2003). All else being equal, it would be less surprising to discover a

missed interaction involving a known generalist species than one with few observed interactions.

Many bipartite interaction networks, especially mutualistic networks, are observed to be 'nested'

- the range of specialists is a subset of the generalists (Bascompte et al. 2003, Ulrich et al. 2009).

Simple degree-models can fit binary nestedness and preferential patterns well. Although

nestedness can be generated by a variety of processes (Song et al. 2017) and patterns observable

in binary networks may not be reflected in the quantitative networks (Staniczenko et al. 2013),

our principle objective here is to identify missing links.

To fit a degree model, we assign each species a connectance term, c, and determine the

probability of an interaction between focal species i and interactor j as:

$$\text{logit}\left(P\left(a_{ij} > 0 \,\middle|\, \mathbf{O}\right)\right) = c_i + c_j + k_c$$

where k_c is a constant intercept term and all parameters are found by maximum likelihood.

When both species have high degrees, their connectance parameters will be high, and the

probability of those species interacting will be high.

Matching – Centrality Model

The matching – centrality approach (Rohr et al. 2016, 2018) combines a latent-trait and a degree

model and is fit in a similar manner. This dual approach has been shown to have a high capacity

to fit tightly to diverse network structures (Rohr et al. 2016). The probability of each interaction

existing is found by:

$$\text{logit}\left(P\left(a_{ij} > 0 \mid \mathbf{O}\right)\right) = c_i + c_j - \lambda\left(m_i - m_j\right)^2 + k_B.$$

As in the latent-trait model, λ is defined to be positive, k_B is a constant intercept term and all

parameters are found by maximum likelihood. We again apply a weak Cauchy prior centred at 0

onto the latent trait terms **m** and fit 10 models, averaging predictions of the best five.

Block Model

Ecological networks, especially antagonistic networks, frequently show comparatively discrete

compartments or modularisation, where subsets of species interact more strongly within their

group than with outsiders (Olesen et al. 2007, 2008, Allesina and Pascual 2009, Schleuning et al.

2014). This may reflect fundamental incompatibilities in physiological traits, temporal mismatch

(such as nocturnal/diurnal partitioning) or spatial segregation.

This grouping can be represented by stochastic block models (SBMs), which have been shown to

perform well on ecological data in comparison to other clustering algorithms (Leger et al. 2015)

and are used increasingly in ecology (Sander et al. 2015, Kéfi et al. 2016). In SBMs, each species

is assigned to a group, G_x, in a defined set: $x \in \{1, \dots, g\}$. The probability of interaction

between two species is determined based on their group membership:

$$P(a_{ij} > 0 \mid \mathbf{O}) = \omega_{G_i G_j}.$$

The elements of ω are the between-group interaction probabilities and are directly specified as

the fraction of observed interactions between each of the groups. We find optimal group

assignations and fit the model using a degree-corrected bipartite-SBM specific algorithm

(Larremore *et al.* 2014, detailed in SI 3). To capture inherent indeterminacy in underlying group

structure, we fit 10 differently-initialised models, and use the average prediction of the five

models with the highest likelihood.

Combinations of Models

Individually the above models each capture discrete pieces of information about the identity of

missing links. However, the structure of ecological networks is the product of many separate

drivers. To capture this diversity, we combine the predictions of multiple models into ensembles.

We test combining the matching-centrality model with the block model, and each of the

'structural' models with the coverage-deficit model.

We test two ensembling approaches, multiplication and averaging. Multiplying the relative

probabilities assigned to each putative missing link, $P_{ens}(a_{ij} > 0) \sim P_1(a_{ij} > 0) \times P_2(a_{ij} > 0)$,

emphasises the extreme probabilities of the constituent models. Averaging the relative

probabilities, $P_{ens}(a_{ij} > 0) \sim P_1(a_{ij} > 0) + P_2(a_{ij} > 0)$, highlights possible interactions that

are consistently identified by multiple models. Before combining models we standardise each set

of probabilities assigned to unobserved interactions to sum to 1.

<u>*Datasets*</u>

Testing the efficacy of extrapolations requires knowledge of the 'true' network. We take two

complimentary approaches to defining our 'true' networks. First, we generate a set of simulated

networks that vary over a wide range of network properties. Second, we use a large and diverse

set of empirical networks from literature sources. In both cases 'observed' networks were

generated by taking draws from a multinomial distribution parameterised by the 'true' interaction

frequency. This replicates ecological sampling processes where the most frequent interactions

are highly likely to have been observed and the rarest missed.

We initially generated 2000 simulated networks from a probabilistic two-trait niche model

(modified from Fründ *et al.* 2016) described in full in SI 1. Key network properties were

calculated using the *bipartite* R package (Dormann et al. 2008): connectance, weighted

nestedness (Galeano et al. 2009), cluster coefficient (Watts and Strogatz 1998), distribution of

interaction frequency (Shannon diversity, Bersier *et al.* 2002) and network level specificity H2'

(Blüthgen et al. 2006). Distributions of these metrics and the (low) correlations between them are

depicted in SI 1.

Our objective was to generate interaction matrices that represent a wide range of ecologically

plausible bipartite networks. However, models used to generate plausible interaction networks

have many similarities to the predictive models described above. This is for a good reason –

models are chosen because they are thought to represent ecological networks parsimoniously. To

reduce circularity, our generative model is considerably more complex than our predictive

models and we examine how predictive model performance changes in response to network

properties.

Using the 'true' networks we took 300-2000 samples per network to generate our 'observed'

networks. We discarded occasional cases where all true interactions were observed. This resulted

in a low mean true sample coverage deficit of 2.4%, corresponding to an average network

completeness of 64% due to skewed interaction strength distributions. Our 'observed' networks

therefore represent reasonably 'well-sampled' systems that are nonetheless missing a large

proportion of the interactions present.

We collated a diverse set of 113 empirical networks representing antagonistic, mutualistic and

commensalistic interaction types (SI 2). We collated quantitative single-class bipartite networks

from the *Web of Life* (www.web-of-life.es) repository with 30-200 species and 40-431 observed

interactions. This included 48 plant-pollinator (19 sources), 23 mammal-flea (one source, but

across a wide geographic range), 2 plant-ant mutualism (2 sources) and 10 seed disperser (from 7

sources) networks. We supplemented this with 25 mammal-dung beetle interaction networks

(Frank *et al.* 2018, selecting those with ≥30 total species, ≥40 observed interactions, and ≥5

mammal species) and 5 host-parasitoid networks (Tylianakis et al. 2007).

These empirical networks are not exhaustively sampled - the Chao1 estimator indicates that the

observed interactions represent on average only 76% of the underlying network (SI 2).

Nonetheless, we took the empirical interaction frequency to be the 'true' network to define a

multinomial distribution. We subsampled each empirical network 20 times, each time taking a

different fraction between 0.2 and 1 of the original sample size, taken to be the sum of the

observations. We excluded cases where the observed web included more than 95% of the 'true'

interactions.

Model Testing

Using each model or combination of models we assign each unobserved interaction a relative

probability of being a missing link. We assume that all observations are true positives (i.e. there

have been no misidentifications) although in principle similar methods could be used to identify

candidates for re-evaluation. Where a species in the 'true' network was not included in the

'observed' network, we did not attempt to predict its interactions and reduced the size of the

'true' network correspondingly for all purposes.

Identifying missing links

We assess performance at identifying missing links with the area under receiver operating

characteristic curve (AUC) metric to assess the information content of a signal, using the pROC

R package (Robin et al. 2011). AUC can be considered the chance that a 'false negative'

(missing link) was assigned a higher relative probability than a 'true negative'. A value of 0.5

indicates no useful signal while a value of 1 indicates perfect discrimination. To determine if

$P(a_{ij} > 0)$ is reflective of true relative interaction frequency, we calculated the mean

Spearman's rank-correlation with the 'true' relative interaction strengths of the unobserved

interactions for each model.

Improving estimates of network metrics

We used the *estimateR* function in the *vegan* R package (Oksanen et al. 2018) to estimate the

number of missing links based on the total distribution of observed interaction counts using the

bias-corrected Chao1 index. We then augmented the observed interaction networks by adding the

estimated number of the most probable missing links to the network based on the average of the

block, matching-centrality and coverage-deficit models. Each selected missing link was taken to

be 'observed' once, and network-level metrics calculated for the observed, augmented and true

networks. We tested three qualitative network metrics (connectance, nestedness and niche

overlap) identified as being susceptible to misestimation in under-sampled networks (Fründ et al.

2016), alongside two metrics considered to be comparatively resistant to under-sampling (H2'

and weighted NODF nestedness).

<u>Results</u>

Identifying missing links - Simulated networks

In general, the network models showed impressive performance at identifying likely missing

links. The best-performing model overall was the block model, closely followed by the other

three network models (Fig. 2a). The coverage deficit model lagged behind the structural models,

but was more informative than simply ranking focal species by sample size. Combining the

predictions of multiple models tended to improve performance relative to individual models

(Table 1). The overall best-performing model averaged the predictions of the block model, the

matching-centrality model and the coverage-deficit model. However, the marginal improvements

from combining models were often small.

Model performance and the relative hierarchy of models was affected by the underlying true

network structure (Fig. 3). On average, missing links were identified best in nested, unclustered

networks where interaction frequencies were more evenly distributed. The coverage-deficit and

sample size models were the least responsive to network structure, since our simulated network

generator did not result in a pronounced abundance-generality correlation. Average model

performance did not change appreciably with increased sample size, overall interaction coverage

deficit or samples per host (Fig. S4). However, when a greater fraction of interactions had been

observed the structural models increased their performance, while performance of the coverage

deficit model remained unchanged (Fig. S4).

Identifying missing links - Empirical networks

The performance of the different predictive models varied considerably between the different

categories of empirical network (Fig. 2b). As expected, the degree model performed best for

mutualistic networks, and the latent-trait model for antagonistic networks. In all six network

categories the matching-centrality model had the best overall performance. The 'true network'

properties (Fig. S3) or the degree of sampling rarefaction (Fig. S4) had relatively little effect on

predictive performance, except for networks that show 'anti-nestedness'. In these cases, the

predictive capacity of the degree model was low, although the matching-centrality model

maintained its advantage. The coverage-deficit model was very poor at predicting missing links

in the empirical datasets and combining it with the network-structure models led to a *reduction*

in overall predictive power (Table 1).

Identifying interaction frequency of missing links

Correlations between probabilities assigned to missing links $P\left(a_{ij} > 0 \middle| \mathbf{O}\right)$ were related to their

'true' interaction frequency. The relationship was stronger in the simulated dataset than the

empirical dataset (Table S2). Of the structural models, the degree model showed the strongest

positive correlation ($\rho_{Sim} = 0.51$, $\rho_{Emp} = 0.15$), while the block model showed the weakest (ρ_{Sim}

$= 0.28$, $\rho_{Emp} = 0.06$). The coverage-deficit model was negatively correlated with true frequency

(ρ_{Sim} = -0.35, ρ_{Emp} = -0.05) i.e. missing links successfully identified via this model tended to be

infrequent.

Improving estimates of network metrics

Integrating inferred interactions into observed networks could improve estimated of key metrics.

The 'augmented' networks showed increased correlation, reduced bias and reduced error in

qualitative metrics, compared to the original observed networks (Fig. 4a). The quantitative

network metrics (H2'and weighted NODF, Fig. 4b) were already well-estimated by the under-

sampled network, and the inclusion of additional interactions made little difference. The Chao1

estimator for the number of missing interactions appeared to perform well for the empirical

networks but consistently underestimated the true interaction count and coverage deficit in the

simulated data (Fig. S5).

Discussion

We have shown that is possible to infer missing links in range of ecological networks with a

reasonable degree of accuracy and this approach can improve estimates of network metrics. The

challenge now is to put these tools into practice to leverage additional ecological insight.

Choosing between predictive models

Ideally, a model used to infer missed interactions would capture the true ecological drivers, but

this is not essential for all purposes. Given the diversity of ecological networks, there will never

be a single 'best' model and victors in comparisons will depend on the data set. In our simulated

data our block model appears to perform best. In our empirical datasets, the matching-centrality

model performed consistently well (as found by Rohr *et al.* 2016). However, we have reason to

distrust the empirical datasets (discussed below). The pronounced, likely artefactual, abundance-

generality relationship will favour degree-models.

Nonetheless, splitting hairs over the best-performing model is not necessarily a productive route.

Networks are structured by multiple processes and in our simulated data sets the very best

performance comes from combining different models, which identify different missing

interactions. Structure-based models pick out the more frequent interactions while coverage

deficit models highlight comparatively infrequent interactions that would be the hardest to

determine through further undirected sampling. Future progress will come from operationalising

estimated missing links, rather than from further incremental model refinements.

Distinctions between empirical and simulated networks

The empirical and simulated datasets overlapped substantially in key network metrics. The main relevant difference between these datasets is the stronger correlation in empirical networks between species' marginal totals and generality (mean Spearman's $\rho_{Sim} = 0.43$, $\rho_{Emp} = 0.76$). This can account for the poor performance of the coverage-deficit model in these cases. Disentangling the extent to which apparent specialism of rarely observed species is a sampling artefact is challenging (Blüthgen et al. 2006), since there are ecological explanations for such a relationship (Fort et al. 2016, Dormann et al. 2017, Simmons et al. 2019). For our purposes, this likely bias in the structure of empirical ecological networks has the consequence that predictive models may identify missing links introduced both by our subsampling procedure, and due to gaps in the original dataset. We therefore place more weight on the results from the simulated data, while noting that, despite the obstacles, the structural models are still able to perform reasonably well on the sparse empirical data.

Estimating the quality of sampling and the number of missing interactions

Sample coverage estimators offer promise for estimating the number of missed interactions (Chacoff et al. 2012, Traveset et al. 2015, Jordano 2016, MacGregor et al. 2017). The distinction between sample-coverage and network completeness is worth emphasising. Apparently high values of sample coverage alone should not be used as a simple proxy for 'network completeness', although they are related to the reliability of certain quantitative network metrics (Henriksen et al. 2019). We found that the whole-network-level Chao1 estimate of missed interactions was strongly correlated with the 'true' number of missing links in our simulated networks, but notably underestimated this number, a pattern also observed by Fründ *et al.* (2016). Although there appeared to be no systematic bias in estimated network completeness in

the empirical datasets, these 'true' networks are themselves highly incomplete, suggesting an

overall underestimation also in these cases. Our focus in this paper is on the location rather than

the number of missing links. Future work should investigate how long tails of infrequent

interactions relate to this underestimation, and consider whether species rank abundance

distribution estimators (Chao et al. 2014) can be usefully generalised to assign a frequency

distribution to estimated missing links.

The way forward

Most of the models used here, and indeed our fitting approaches, likely have further room for

refinement, and there are many further classes of predictive models that we did not test, such as

nearest neighbour algorithms (Desjardins-Proulx et al. 2017). Furthermore, we based our models

entirely on the information contained in the network itself. In most cases, ecologists constructing

a network have access to (sometimes considerable) additional information and expertise on the

study system. Introducing pre-existing knowledge about likely or forbidden interactions, or

pooled knowledge from multiple sample sites offers considerable opportunities to maximise the

information in a network (Gray et al. 2015, Pomeranz et al. 2018). External information would

be particularly useful in separating 'forbidden links' (Olesen et al. 2011) that never occur due to

fundamental incompatibilities from those that occur elsewhere but not within those spatio-

temporal boundaries. For instance, species may not be observed to interact because of a

fundamental mismatch in body-sizes (a forbidden link) or because they interact at night but data

were gathered only during the day.

Because the naïve models used here do not use external information, or the judgment of the

ecologists constructing the network, the quality of the estimation very much represents a lower

bound of what is possible with the approach in practice. As a result of the underlying

idiosyncrasies of ecological systems, often euphemistically labelled 'species identity effects',

there will always be an upper limit to the predictive capacity of any statistical model. We suggest

that, rather than developing new models to gain marginal improvements in predictive capacity, a

more productive focus would be to develop frameworks exploiting this information to test the

robustness of conclusions derived from ecological networks.

Complete network inventories are not a realistic or necessarily useful goal (Jordano 2016).

Enormous sample sizes and strictly deliminated boundaries are required to exhaustively sample

all interactions (Martinez et al. 1999). There is undoubtedly a long tail of extremely infrequent

interactions that will never be observed in a realistic sampling regime. Furthermore, there is

rarely a clear line defining network boundaries (Poisot et al. 2015). Ecological networks are a

constantly moving target (Rasmussen et al. 2013, CaraDonna et al. 2017) and there may be little

value in including certain rare, occasional visitors and unusual interactions. It can be challenging

to determine if infrequent interactions are simply a curiosity (e.g. Mercer 1966) or have

significant consequences (e.g. Dudley et al. 2016). The utility of increasing the accuracy with

which one type of interaction is understood will plateau as the impact of other interaction types

and ecological guilds predominate (Lafferty et al. 2008). Despite the challenges, we see two

principle uses for inferred missing links.

First, inferring missing links will direct further sampling where the goal is a descriptive network.

In many cases the topography of the network is of principle interest, given the potential for

interaction strengths to vary through time and in response to perturbation. For example, when

tracing the relationships of species that may be the target for eradication measures, such as

disease-vector mosquitoes (Collins et al. 2019), a critical first step is an inventory of interactions

in the community. Since in most cases investment in sampling is a finite resource, directing

sampling towards high-likelihood interactions could more efficiently increase the overall

network completeness. However, we caution that directed sampling will introduce biases in the

generated network (Gibson et al. 2011, Falcão et al. 2016) and may hinder cross-community

comparability (Poisot et al. 2012a). In certain cases, independently-sampled networks (even if

less complete) may be more useful than comparisons between 'augmented' networks. The use of

information from other networks may lead them to appear more homogenous than they really are

and lead to underestimates of interaction β-diversity through time and space (Poisot et al. 2015,

Graham and Weinstein 2018).

Second, the identification of probable missing links can improve tests of the reliability of

conclusions drawn from networks, potentially through explicitly probabilistic networks (Poisot et

al. 2016a). Statistical resampling and rarefaction procedures that can only work backwards or by

adding random interactions may both introduce biases. While particular quantitative metrics are

relatively robust to under-sampling (Blüthgen et al. 2008, Fründ et al. 2016, Henriksen et al.

2019), these metrics may not align with the goals of the study. If the conclusions are robust to the

inclusion of probable missing links, then confidence in the results would be increased.

Conversely, if including most-likely missing links alters conclusions, this would signal that these

conclusions are unlikely to be robust. In these cases, care should be taken when selecting a

predictive model to avoid a clash with the properties under investigation.

Conclusion

The identification of likely missing links is a powerful tool to include in the arsenal of network

ecologists. We have shown that models to identify missing links can perform well across a wide

range of network types. Importantly, inferential models need not be technically challenging to

implement – for example the *cassandRa* package allows all the models discussed here to be fit

with a single function call. We hope that such approaches become more widely used in guiding

ecological sampling and testing the robustness of conclusions drawn from bipartite networks.

Acknowledgments

Thanks to the CERO group for comments on this project, and two anonymous reviewers for their

constructive comments. Funding was provided by NERC Standard Grant NE/N010221/1.

Data accessibility

R functions to run the predictive models are publicly available in the R package *cassandRa* on

CRAN. All code and necessary data to generate simulated datasets and carry out analyses are

available on GitHub (github.com/jcdterry/PredictingMissingLinks). Should the manuscript be

accepted, this will also be archived in a public repository and the data doi included.

Statement of Authorship

JCDT initiated the research, in discussion with OTL. JCDT conducted all analyses and wrote the

first draft of the manuscript. Both authors contributed to revisions.

References

de Aguiar, M. A. M., E. A. Newman, M. M. Pires, J. D. Yeakel, C. Boettiger, L. A. Burkle, D.

Gravel, P. R. Guimarães, J. L. O'Donnell, T. Poisot, M.-J. Fortin, and D. H. Hembry. 2019.

Revealing biases in the sampling of ecological interaction networks. PeerJ 7:e7566.

Allesina, S., and M. Pascual. 2009. Food web models: a plea for groups. Ecology Letters

12:652–662.

Bartomeus, I. 2013. Understanding Linkage Rules in Plant-Pollinator Networks by Using

Hierarchical Models That Incorporate Pollinator Detectability and Plant Traits. PLoS ONE

8:1–8.

Bartomeus, I., D. Gravel, J. M. Tylianakis, M. A. Aizen, I. A. Dickie, and M. Bernard-Verdier.

2016. A common framework for identifying linkage rules across different types of

interactions. Functional Ecology 30:1894–1903.

Bascompte, J., and P. Jordano. 2013. Mutualistic Networks. Princeton University Press,

Princeton.

Bascompte, J., P. Jordano, C. J. Melian, and J. M. Olesen. 2003. The nested assembly of plant-

animal mutualistic networks. Proceedings of the National Academy of Sciences 100:9383–

9387.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. Nature

398:330–334.

Bersier, L.-F., C. Banašek-Richter, and M.-F. Cattin. 2002. Quantitative descriptors of food web

matrices. Ecology 83:2917–2926.

Blüthgen, N. 2010. Why network analysis is often disconnected from community ecology: A

critique and an ecologist's guide. Basic and Applied Ecology 11:185–195.

Blüthgen, N., J. Fründ, D. P. Vázquez, and F. Menzel. 2008. What Do Interaction Network

Metrics Tell Us about Specialization and Biological Traits. Ecology 89:3387–3399.

Blüthgen, N., F. Menzel, and N. Blüthgen. 2006. Measuring specialization in species interaction

networks. BMC Ecology 6:9.

CaraDonna, P. J., W. K. Petry, R. M. Brennan, J. L. Cunningham, J. L. Bronstein, N. M. Waser,

and N. J. Sanders. 2017. Interaction rewiring and the rapid turnover of plant–pollinator

networks. Ecology Letters 20:385–394.

Chacoff, N. P., D. P. Vázquez, S. B. Lomáscolo, E. L. Stevani, J. Dorado, and B. Padrón. 2012.

Evaluating sampling completeness in a desert plant-pollinator network. Journal of Animal

Ecology 81:190–200.

Chao, A. 1984. Nonparametric estimation of the number of classes in a population. Scandinavian

Journal of Statistics 11:265–270.

Chao, A., T. C. Hsieh, R. L. Chazdon, and R. K. Colwell. 2014. Unveiling the Species-Rank

Abundance Distribution by Generalizing Good-Turing Sample Coverage Theory. Ecology

96:1189–1201.

Chao, A., and L. Jost. 2012. Coverage-based rarefaction and extrapolation: standardizing

samples by completeness rather than size. Ecology 93:2533–2547.

Clauset, A., C. Moore, and M. E. J. Newman. 2008. Hierarchical structure and the prediction of

missing links in networks. Nature 453:98–101.

Collins, C. M., J. A. S. Bonds, M. M. Quinlan, and J. D. Mumford. 2019. Effects of the removal

or reduction in density of the malaria mosquito, Anopheles gambiae s.l ., on interacting

predators and competitors in local ecosystems. Medical and Veterinary Entomology 33:1–

15.

Colwell, R. K., and J. A. Coddington. 1994. Estimating terrestrial biodiversity through

extrapolation. Philosophical Transactions of the Royal Society B-Biological Sciences

345:101–118.

Costa, J. M., L. P. da Silva, J. A. Ramos, and R. H. Heleno. 2016. Sampling completeness in

seed dispersal networks: When enough is enough. Basic and Applied Ecology 17:155–164.

Crea, C., R. A. Ali, and R. Rader. 2016. A new model for ecological networks using species-

level traits. Methods in Ecology and Evolution 7:232–241.

Delmas, E., M. Besson, M.-H. Brice, L. A. Burkle, G. V. Dalla Riva, M.-J. Fortin, D. Gravel, P.

R. Guimarães, D. H. Hembry, E. A. Newman, J. M. Olesen, M. M. Pires, J. D. Yeakel, and

T. Poisot. 2019. Analysing ecological networks of species interactions. Biological Reviews

94:16–36.

Desjardins-Proulx, P., I. Laigle, T. Poisot, and D. Gravel. 2017. Ecological interactions and the

Netflix problem. PeerJ 5:e3644.

Dormann, C. F., J. Fründ, and H. M. Schaefer. 2017. Identifying Causes of Patterns in Ecological

Networks: Opportunities and Limitations. Annual Review of Ecology, Evolution, and

Systematics 48:559–584.

Dormann, C. F., B. Gruber, and J. Frund. 2008. Introducing the bipartite Package: Analysing

Ecological Networks. R News 8:8–11.

Dudley, J. P., B. M. Hang'Ombe, F. H. Leendertz, L. J. Dorward, J. de Castro, A. L. Subalusky,

and M. Clauss. 2016. Carnivory in the common hippopotamus Hippopotamus amphibius□:

implications for the ecology and epidemiology of anthrax in African landscapes. Mammal

Review 46:191–203.

Dunne, J. A., and M. Pascual. 2006. Ecological Networks: Linking Structure to Dynamics in

Food Webs. Oxford University Press, Oxford.

Eklöf, A., U. Jacob, J. Kopp, J. Bosch, R. Castro-Urgal, N. P. Chacoff, B. Dalsgaard, C. de Sassi,

M. Galetti, P. R. Guimarães, S. B. Lomáscolo, A. M. Martín González, M. A. Pizo, R.

Rader, A. Rodrigo, J. M. Tylianakis, D. P. Vázquez, and S. Allesina. 2013. The

dimensionality of ecological networks. Ecology Letters 16:577–583.

Falcão, J. C. F., W. Dáttilo, and V. Rico-Gray. 2016. Sampling effort differences can lead to

biased conclusions on the architecture of ant-plant interaction networks. Ecological

Complexity 25:44–52.

Fort, H., D. P. Vázquez, and B. L. Lan. 2016. Abundance and generalisation in mutualistic

networks: Solving the chicken-and-egg dilemma. Ecology Letters 19:4–11.

Frank, K., F.-T. Krell, E. M. Slade, E. H. Raine, L. Y. Chiew, T. Schmitt, C. S. Vairappan, P.

Walter, and N. Blüthgen. 2018. Global dung webs: high trophic generalism of dung beetles

along the latitudinal diversity gradient. Ecology Letters 21:1229–1236.

Fründ, J., K. S. Mccann, and N. M. Williams. 2016. Sampling bias is a challenge for quantifying

specialization and network structure: Lessons from a quantitative niche model. Oikos

125:502–513.

Galeano, J., J. M. Pastor, and J. M. Iriondo. 2009. Weighted-Interaction Nestedness Estimator

(WINE): A new estimator to calculate over frequency matrices. Environmental Modelling

& Software 24:1342–1346.

Gellner, G., and K. S. McCann. 2016. Consistent role of weak and strong interactions in high-

and low-diversity trophic food webs. Nature Communications 7:11180.

Gibson, R. H., B. Knott, T. Eberlein, and J. Memmott. 2011. Sampling method influences the

structure of plant-pollinator networks. Oikos 120:822–831.

Goldwasser, L., and J. Roughgarden. 1997. Sampling Effects and the Estimation of Food-Web

Properties. Ecology 78:41–54.

González-Varo, J. P., and A. Traveset. 2016. The Labile Limits of Forbidden Interactions.

Trends in Ecology and Evolution 31:700–710.

Gotelli, N. J., and R. K. Colwell. 2010. Estimating species richness. Pages 416–422 *in* A.

Magurran and B. J. McGill, editors. Biological Diversity: Frontiers in Measurement and

Assessment. Open University Press.

Graham, C. H., and B. G. Weinstein. 2018. Towards a predictive model of species interaction

beta diversity. Ecology Letters 21:1299–1310.

Gray, C., D. J. Baird, S. Baumgartner, U. Jacob, G. B. Jenkins, E. J. O'Gorman, X. Lu, A. Ma,

M. J. O. Pocock, N. Schuwirth, M. Thompson, and G. Woodward. 2014. Ecological

networks: the missing links in biomonitoring science. Journal of Applied Ecology 51:1444–

1449.

Gray, C., D. H. Figueroa, L. N. Hudson, A. Ma, D. Perkins, and G. Woodward. 2015. Joining the

dots: An automated method for constructing food webs from compendia of published

interactions. Food Webs 5:11–20.

Guimera, R., and M. Sales-Pardo. 2009. Missing and spurious interactions and the reconstruction

of complex networks. Proceedings of the National Academy of Sciences 106:22073–22078.

Henriksen, M. V., D. G. Chapple, S. L. Chown, and M. A. McGeoch. 2019. The effect of

network size and sampling completeness in depauperate networks. Journal of Animal

Ecology 88:211–222.

Jacquet, C., C. Moritz, L. Morissette, P. Legagneux, F. Massol, P. Archambault, and D. Gravel.

2016. No complexity–stability relationship in empirical ecosystems. Nature

Communications 7:12573.

Jordano, P. 2016. Sampling networks of ecological interactions. Functional Ecology 30:1883–

1893.

Jordano, P., J. Bascompte, and J. M. Olesen. 2003. Invariant properties in coevolutionary

networks of plant-animal interactions. Ecology Letters 6:69–81.

Kadoya, T., G. Gellner, and K. S. McCann. 2018. Potential oscillators and keystone modules in

food webs. Ecology Letters 21:1330–1340.

Kéfi, S., V. Miele, E. A. Wieters, S. A. Navarrete, and E. L. Berlow. 2016. How Structured Is the

Entangled Bank? The Surprisingly Simple Organization of Multiplex Ecological Networks

Leads to Increased Persistence and Resilience. PLOS Biology 14:e1002527.

Klumpers, S. G. T., M. Stang, and P. G. L. Klinkhamer. 2019. Foraging efficiency and size

matching in a plant-pollinator community: the importance of sugar content and tongue

length. Ecology Letters 22:469–479.

Lafferty, K. D., S. Allesina, M. Arim, C. J. Briggs, G. De Leo, A. P. Dobson, J. A. Dunne, P. T.

J. Johnson, A. M. Kuris, D. J. Marcogliese, N. D. Martinez, J. Memmott, P. A. Marquet, J.

P. McLaughlin, E. A. Mordecai, M. Pascual, R. Poulin, and D. W. Thieltges. 2008.

Parasites in food webs: the ultimate missing links. Ecology Letters 11:533–546.

Laigle, I., I. Aubin, C. Digel, U. Brose, I. Boulangeat, and D. Gravel. 2018. Species traits as

drivers of food web structure. Oikos 127:316–326.

Larremore, D. B., A. Clauset, and A. Z. Jacobs. 2014. Efficiently inferring community structure

in bipartite networks. Physical Review E 90:012805.

Lee, R. H., and B. Guénard. 2019. Choices of sampling method bias functional components

estimation and ability to discriminate assembly mechanisms. Methods in Ecology and

Evolution 10:867–878.

Leger, J.-B., J.-J. Daudin, and C. Vacher. 2015. Clustering methods differ in their ability to

detect patterns in ecological networks. Methods in Ecology and Evolution 6:474–481.

MacGregor, C. J., D. M. Evans, and M. J. O. Pocock. 2017. Estimating sampling completeness

of interactions in quantitative bipartite ecological networks: incorporating variation in

species specialisation. bioRxiv:195917.

Martinez, N. D., B. A. Hawkins, H. A. Dawah, and B. P. Feifarek. 1999. Effects of Sampling

Effort on Characterization of Food-Web Structure. Ecology 80:1044–1055.

McCann, K., A. Hastings, and G. R. Huxel. 1998. Weak trophic interactions and the balance of

nature. Nature 395:794–798.

Memmott, J. 2009. Food webs: a ladder for picking strawberries or a practical tool for practical

problems? Philosophical Transactions of the Royal Society B: Biological Sciences

364:1693–1699.

Mercer, A. J. 1966. Turnstones feeding on human corpse. British Birds 59:307.

Montoya, J. M., S. L. Pimm, and R. V Solé. 2006. Ecological networks and their fragility. Nature

442:259–64.

Morales-Castilla, I., M. G. Matias, D. Gravel, and M. B. Araújo. 2015. Inferring biotic

interactions from proxies. Trends in Ecology & Evolution 30:347–356.

Morris, R. J., S. Gripenberg, O. T. Lewis, and T. Roslin. 2014. Antagonistic interaction networks

are structured independently of latitude and host guild. Ecology Letters 17:340–349.

Nielsen, A., and J. Bascompte. 2007. Ecological networks, nestedness and sampling effort.

Journal of Ecology 95:1134–1141.

Novotny, V., S. E. Miller, L. Baje, S. Balagawi, Y. Basset, L. Cizek, K. J. Craft, F. Dem, R. a I.

Drew, J. Hulcr, J. Leps, O. T. Lewis, R. Pokon, A. J. a Stewart, G. A. Samuelson, and G. D.

Weiblen. 2010. Guild-specific patterns of species richness and host specialization in plant-

herbivore food webs from a tropical forest. The Journal of animal ecology 79:1193–203.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.

B. O'Hara, G. L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs, and H. Wagner.

2018. vegan: Community Ecology Package.

Olesen, J. M., J. Bascompte, Y. L. Dupont, H. Elberling, C. Rasmussen, and P. Jordano. 2011.

Missing and forbidden links in mutualistic networks. Proceedings of the Royal Society B:

Biological Sciences 278:725–732.

Olesen, J. M., J. Bascompte, Y. L. Dupont, and P. Jordano. 2007. The modularity of pollination

networks. Proceedings of the National Academy of Sciences 104:19891–19896.

Olesen, J. M., J. Bascompte, H. Elberling, and P. Jordano. 2008. Temporal dynamics in a

pollination network. Ecology 89:1573–1582.

Paine, R. T. 1980. Food Webs: Linkage, Interaction Strength and Comunity Infrastructure. The

Journal of Animal Ecology 49:666–685.

Poisot, T., E. Canard, D. Mouillot, N. Mouquet, and D. Gravel. 2012a. The dissimilarity of

species interaction networks. Ecology Letters 15:1353–1361.

Poisot, T., E. Canard, N. Mouquet, and M. E. Hochberg. 2012b. A comparative study of

ecological specialization estimators. Methods in Ecology and Evolution 3:537–544.

Poisot, T., A. R. Cirtwill, K. Cazelles, D. Gravel, M.-J. Fortin, and D. B. Stouffer. 2016a. The

structure of probabilistic networks. Methods in Ecology and Evolution 7:303–312.

Poisot, T., D. B. Stouffer, and D. Gravel. 2015. Beyond species: why ecological interaction

networks vary through space and time. Oikos 124:243–251.

Poisot, T., D. B. Stouffer, and S. Kéfi. 2016b. Describe, understand and predict: why do we need

networks in ecology? Functional Ecology 30:1878–1882.

Pomeranz, J. P. F., R. M. Thompson, T. Poisot, and J. S. Harding. 2018. Inferring predator–prey

interactions in food webs. Methods in Ecology and Evolution 2018:1–12.

Rasmussen, C., Y. L. Dupont, J. B. Mosbacher, K. Trøjelsgaard, and J. M. Olesen. 2013. Strong

Impact of Temporal Resolution on the Structure of an Ecological Network. PLoS ONE

8:e81694.

Rivera-Hutinel, A., R. O. Bustamante, V. H. Marín, and R. Medel. 2012. Effects of sampling

completeness on the structure of plant-pollinator networks. Ecology 93:1593–1603.

Robin, X., N. Turck, A. Hainard, N. Tiberti, F. Lisacek, J.-C. Sanchez, and M. Müller. 2011.

pROC: an open-source package for R and S+ to analyze and compare ROC curves. BMC

Bioinformatics 12:77.

Rohr, R. P., R. E. Naisbit, C. Mazza, and L.-F. Bersier. 2016. Matching–centrality

decomposition and the forecasting of new links in networks. Proceedings of the Royal

Society B: Biological Sciences 283:20152702.

Rohr, R. P., R. E. Naisbit, C. Mazza, and L.-F. Bersier. 2018. Statistical Approaches for Inferring

and Predicting Food-Web Architecture. Pages 178–192 *in* J. C. Moore, P. C. de Ruiter, K.

S. McCann, and V. Wolters, editors. Adaptive Food Webs, Stability and Transistions of

Real and Model Ecosystems. Cambridge University Press.

Sander, E. L., J. T. Wootton, and S. Allesina. 2015. What Can Interaction Webs Tell Us About

Species Roles? PLoS Computational Biology 11:e1004330.

Schleuning, M., L. Ingmann, R. Strauß, S. A. Fritz, B. Dalsgaard, D. Matthias Dehling, M. Plein,

F. Saavedra, B. Sandel, J.-C. Svenning, K. Böhning-Gaese, and C. F. Dormann. 2014.

Ecological, historical and evolutionary determinants of modularity in weighted seed-

dispersal networks. Ecology Letters 17:454–463.

Seo, E., and R. A. Hutchinson. 2018. Predicting Links in Plant-Pollinator Interaction Networks

Using Latent Factor Models With Implicit Feedback. Pages 808–815 Proceedings of the

Thirty-Second AAAI Conference on Artificial Intelligence (AAAI-18).

Simmons, B. I., J. Vizentin-Bugoni, P. K. Maruyama, P. A. Cotton, O. H. Marín-Gómez, C.

Lara, L. Rosero-Lasprilla, M. A. Maglianesi, R. Ortiz-Pulido, M. A. Rocca, L. C.

Rodrigues, B. Tinoco, M. F. Vasconcelos, M. Sazima, A. M. M. González, J. Sonne, C.

Rahbek, L. V. Dicks, B. Dalsgaard, and W. J. Sutherland. 2019. Abundance drives broad

patterns of generalisation in plant–hummingbird pollination networks. Oikos:oik.06104.

Song, C., R. P. Rohr, and S. Saavedra. 2017. Why are some plant–pollinator networks more

nested than others? Journal of Animal Ecology 86:1417–1424.

Spitz, J., V. Ridoux, and A. Brind'Amour. 2014. Let's go beyond taxonomy in diet description:

Testing a trait-based approach to prey-predator relationships. Journal of Animal Ecology

83:1137–1148.

Staniczenko, P. P. A., J. C. Kopp, and S. Allesina. 2013. The ghost of nestedness in ecological

networks. Nature Communications 4:1391.

Staniczenko, P. P. A., O. T. Lewis, J. M. Tylianakis, M. Albrecht, V. Coudrain, A.-M. Klein, and

F. Reed-Tsochas. 2017. Predicting the effect of habitat modification on networks of

interacting species. Nature Communications 8:792.

Stock, M., T. Poisot, W. Waegeman, and B. De Baets. 2017. Linear filtering reveals false

negatives in species interaction data. Scientific Reports 7:45908.

Terry, J. C. D. 2019. cassandRa R package.

Terry, J. C. D., R. J. Morris, and M. B. Bonsall. 2017. Trophic interaction modifications: an

empirical and theoretical framework. Ecology Letters 20:1219–1230.

Traveset, A., J. M. Olesen, M. Nogales, P. Vargas, P. Jaramillo, E. Antolín, M. M. Trigo, and R.

Heleno. 2015. Bird–flower visitation networks in the Galápagos unveil a widespread

interaction release. Nature Communications 6:6376.

Tylianakis, J. M., E. Laliberté, A. Nielsen, and J. Bascompte. 2010. Conservation of species

interaction networks. Biological Conservation 143:2270–2279.

Tylianakis, J. M., T. Tscharntke, and O. T. Lewis. 2007. Habitat modification alters the structure

of tropical host-parasitoid food webs. Nature 445:202–5.

Ulrich, W., M. Almeida-Neto, and N. J. Gotelli. 2009. A consumer's guide to nestedness

analysis. Oikos 118:3–17.

Valdovinos, F. S. 2019. Mutualistic networks: moving closer to a predictive theory. Ecology

Letters 22:1517–1534.

Valiente-Banuet, A., M. A. Aizen, J. M. Alcántara, J. Arroyo, A. Cocucci, M. Galetti, M. B.

García, D. García, J. M. Gómez, P. Jordano, R. Medel, L. Navarro, J. R. Obeso, R. Oviedo,

N. Ramírez, P. J. Rey, A. Traveset, M. Verdú, and R. Zamora. 2015. Beyond species loss:

The extinction of ecological interactions in a changing world. Functional Ecology 29:299–

307.

Vázquez, D. P., N. Bluthgen, L. Cagnolo, and N. P. Chacoff. 2009. Uniting pattern and process

in plant-animal mutualistic networks: A review. Annals of Botany 103:1445–1457.

Vizentin-Bugoni, J., P. K. Maruyama, V. J. Debastiani, L. da S. Duarte, B. Dalsgaard, and M.

Sazima. 2016. Influences of sampling effort on detected patterns and structuring processes

of a Neotropical plant-hummingbird network. Journal of Animal Ecology 85:262–272.

Vizentin-Bugoni, J., P. K. Maruyama, and M. Sazima. 2014. Processes entangling interactions in

communities: forbidden links are more important than abundance in a hummingbird-plant

network. Proceedings of the Royal Society B: Biological Sciences 281:20132397.

Watts, D. J., and S. H. Strogatz. 1998. Collective dynamics of "small-world" networks. Nature

393:440–442.

Weinstein, B. G., and C. H. Graham. 2017. On comparing traits and abundance for predicting

species interactions with imperfect detection. Food Webs 11:17–25.

Model	Simulated Dataset Accuracy Score (AUC)			Empirical Dataset Accuracy Score (AUC)		
	Structural Model-Only	Ensembled with Coverage-Deficit model		Structural Model-Only	Ensembled with Coverage-Deficit model	
		($\times$)	(+)		($\times$)	(+)
Focal Layer Sampling						
Coverage-Deficit	0.600			0.589[abc]		
Observation Count	0.580			0.603[de]		
Network Models						
Latent-Trait	0.660 [a]	0.649	0.654 [a]	0.631[f]	0.588[abc]	0.590[abc]
Degree	0.676	0.708[efg]	0.683 [b]	0.721 [i]	0.651[h]	0.663
Matching-Centrality	0.671	0.702[de]	0.684 [b]	0.718 [i]	0.642[g]	0.658
Block	0.691 [bcd]	0.689 [bc]	0.695 [cd]	0.630[f]	0.584[a]	0.590[c]
Combined Models						
Block and Matching-Centrality (Multiplied)	0.702 [de]	0.705[ef]	0.711[f]	0.647 [gh]	0.594[c]	0.604[de]
Block and Matching-Centrality (Averaged)	0.705[ef]	0.709[f]	0.713 [g]	0.648 [gh]	0.600[d]	0.603[d]

Table 1 Mean AUC values using different predictive models or combinations of models across the whole

simulated (n=1985) and empirical (n=113, 2112 subsamples) network sets. Within each dataset, results

are shown with just the network-structure based model (left column), ensembled with the coverage-deficit

model, by multiplying the probabilities (centre column) or by averaging the assigned probabilities (right

column). Incorporating coverage deficit tends to lead to modest improvements in the simulated networks

but not in the empirical networks. Sets of results found not to differ significantly (Holm-correct t-tests,

paired by network ID) are indicated with grouping letters, applied separately for simulated and empirical

results.

Figure 1. Demonstration of the fits of predictive models to a subsampled simulated bipartite

network. Brighter colours indicate a higher probability of an unobserved interaction being

present according to that model. Black dots (●) indicate observed interactions and red targets (⌖)

indicate unobserved missing links that we wish to identify. AUC values for this example were as

follows: Degree: 0.80, Latent-Trait: 0.76, Matching-Centrality: 0.81, Block: 0.88, Coverage-

Deficit: 0.7, Averaging the Block and the Coverage-Deficit model 0.92.

Figure 2. a) Model performance across 1985 simulated networks. Mean differences between the

statistical models were small, although given the large sample size all pairwise differences were

statistically significant (Holm-corrected t-tests, paired by network ID, p<0.0001). **b)** Model

performance on the empirical data sets, subdivided by network class. Each empirical network

was subsampled 20 times to different fractions of the original sample size. Models perform

differently on different classes of data (Model comparison, accounting for dataset ID as a

random effect, shows model:class interaction term is significant: $p < 0.0001$, $n_{obs} = 12672$). In all

cases boxes show first and third quartiles.

Figure 3. Response of model performance to 'true' network structure. Smoothed line shows a

GAM fit for each predictive model. Lower black line shows the local density distribution of that

property across the simulated test dataset.

Figure 4. Relationship between network metrics calculated on the observed (black) and

'augmented' (red) interaction networks to the true network value in the simulated networks. The

line shows the 1:1 relationship. Pearson's correlation coefficient (r) is shown for each metric.

Qualitative metrics (a) show notable improvements in correlation with the true values when

augmented, while quantitative metrics (b) are largely unchanged.

a) Artificial Dataset

b) Empirical Dataset

Predictive Model: ⊟ Coverage Deficit ⊟ Sample Size ⊟ Block ⊟ Degree ⊟ Latent-Trait ⊟ Matching-Centrality

a)

Connectance

Augmented r: 0.635
Observed r: 0.474

Nestedness

Augmented r: 0.794
Observed r: 0.734

Upper Level Niche Overlap

Augmented r: 0.948
Observed r: 0.938

Lower Level Niche Overlap

Augmented r: 0.941
Observed r: 0.937

b)

Specialisation, H2'

Augmented r: 0.99
Observed r: 0.993

Weighted Nestedness, NODF

Augmented r: 0.811
Observed r: 0.838

True Value Of Full Network

Estimated Metric Value