Title: kin recognition and genetic variation for competitive ability in the annual legume

Medicago minima

Authors: Sara Tomiolo[1]*, Claire Thomas[2], Michael K. Jespersen[1], Christian F. Damgaard[1], Bodil

K. Ehlers[1]

Affiliation:

Department of Bioscience, Aarhus University. Vejlsovej 25, 8600 Silkeborg, Denmark

INRA - Montpellier SupAgro. 2 Place Pierre Viala, 34060 Montpellier, France

*Corresponding author: sara.tomiolo@gmail.com

Keywords:

Intraspecific interactions, kin-competition, mini-communities, competitive ability, root behaviour

Abstract

Knowing which mechanisms drive the outcome of intraspecific interactions is highly relevant for

understanding diversity maintenance. Many plant species exhibit strong population genetic

substructure over small spatial scales due to more shared genes between conspecific close

neighbours. This results in frequent interactions among genetically related plants. Predictions of

how genetic similarity may drive the outcome of intraspecific interactions are based on two

contrasting theories: the resource partitioning hypothesis and kin selection theory. The first predicts

that competition will be stronger among closely related conspecific (i.e. kin) because similar

genotypes have similar resource requirements. The second predicts instead that competition will be

reduced among kin, in order to maximize the inclusive fitness.

Although efforts have been made to reconcile these two theories as non-mutually exclusive,

outcome of intraspecific interaction studies are frequently interpreted as the results of either one or

the other.

We used an annual legume, *Medicago minima*, to conduct two greenhouse experiments testing

changes in root behaviour, above-ground growth and biomass in response to neighbour identity. We

found evidence of both genetic variation in competitive ability, and reduced competition among kin

in some genotypes. Reduced competitive growth towards kin was found in the most competitive

genotypes, suggesting that kin avoidance and competitive ability were not independent. With

presence of both kin competition avoidance and variation for competitive ability, the outcome of

intraspecific interactions will strongly depend on the local spatial genetic substructure. This is

highly relevant to predict how intraspecific competition affect diversity maintenance.

Introduction

The last decade has seen an increasing interest in understanding the mechanisms driving the outcome of intraspecific interactions and whether new insights can be integrated into species coexistence theory (e.g. Bolnick et al., 2011; Hart, Schreiber, & Levine, 2016). In plants, genetic variation for competitive ability for traits like growth rate, resource acquisition and allocation can create variation in competitive ability within a species. This often results in an asymmetric competition where competitively superior genotypes outperform their neighbours when competing for the same limiting resource (Weiner & Thomas, 1986; Weiner, Campbell, Pino, & Echarte, 2009).

Studies on intraspecific interactions generally test either of two theoretical predictions: the resource partitioning hypothesis (Young, 1981) and kin selection theory (Hamilton, 1964). The resource partitioning hypothesis, predicts that closely related species have similar resource requirements and therefore compete more harshly (MacArthur & Levins, 1967; Burns & Strauss, 2012). At the intraspecific level, where genetically similar individuals are expected to have similar resource requirements, the resource-partitioning hypothesis predicts that competition between close relatives should be stronger than between distantly related conspecifics (Young, 1981; Kelly, 1996). Conversely, kin selection theory predicts that kin should avoid strong competition with relatives, as it reduces the inclusive fitness (Hamilton, 1964) and should either reduce competitive interference with kin (West, Pen, & Griffin, 2002) or disperse to avoid competition (Ronce, 2007).

In plant ecology, resource partitioning has long been the traditional view on plant competition. More recently, kin selection theory received support from several studies, which found plants that exhibit a plastic growth response with reduced competitive growth toward kin neighbours compared

to unrelated conspecifics (Dudley & File, 2007; Bhatt, Khandelwal, & Dudley, 2010; Crepy & Casal, 2014).

However, there is currently no consensus, as empirical studies have produced contradicting results, finding evidence for both increased and reduced competition in plants growing with kin compared to non-kin (Donohue, 2003; Cheplick & Kane, 2004; Milla, Forero, Escudero, & Iriondo, 2009; Masclaux et al., 2010; Simonsen, Chow, & Stinchcombe, 2014). This is consistent with the view that only under a specific set of conditions would the evolution of reduced competition among kin be advantageous (Crepy & Casal, 2014; Ehlers & Bilde, 2019). For example, life-history traits such as barochory and synaptospermy are associated with a clustered distribution of kin or siblings (Lepik, Abakumova, Zobel, & Semchenko, 2012), and in this case, it would be favourable to discriminate between siblings and strangers and modulate the intensity of competition accordingly.

One issue is that it can be difficult to determine if different outcomes of intraspecific interactions between kin and non-kin are caused by "actual" kin responses or by genetic similarity. With genetic variation for competitive ability, symmetric competition among kin may be the result of similar competitive ability among genetically identical individuals, but may also result from reduced competition toward plants recognized as kin. These two explanations can lead to similar outcomes in terms of mean fitness differences between kin and non-kin interactions, making it difficult to distinguish between the explanatory mechanisms (e.g. Milla et al., 2009; Simonsen et al., 2014).

Recent conceptual papers on this topic propose that , rather than interpreting intraspecific interactions as the results of either genetic variation in competitive abilities or avoidance of kin competition, both mechanisms may act simultaneously within a species and possibly even within a population (File, Murphy, & Dudley, 2011; Ehlers & Bilde, 2019). However, this interpretation of intraspecific interactions has so far received little empirical attention (but see Biernaskie, 2010).

If both mechanisms co-occur, the outcome of intraspecific interactions in natural populations may

depend greatly on the small-scale population genetic structure. For example, the same genotype

could experience reduced intraspecific competition when growing close to kin, and varying

intensity of intraspecific competition when growing with non-kin, which depend on differences in

competitive ability and resource requirements among genotypes.

Understanding what drives the outcome of genotype-dependent interactions among conspecifics

may help explain the maintenance of intraspecific genetic variation, and determine the relative

strength of within versus between species competition that governs species coexistence (Chesson,

2000; Vellend & Geber, 2005; Laird & Schamp, 2006; Ehlers et al., 2016).

In this study, we used the annual legume *Medicago minima* to investigate the outcome of

intraspecific interactions, by growing different genotypes with either kin or non-kin neighbours. We

report the results of two experiments in which we tested for a) different responses in root growth

toward kin and non-kin neighbours, and b) different responses in above-ground growth and biomass

of plants grown in mini-communities with neighbours of varying relatedness.

We used these two experiments to test the following hypotheses: 1) *M. minima* genotypes reduce

competition among kin by decreasing root growth towards kin vs. non-kin neighbours, and 2) the

outcome of competitive interactions among genotypes are affected by both genotype-specific

variation in competitive ability among non-kin and reduced competition with kin.

Material and methods

Study species:

Medicago minima L. is an annual legume widespread in the Mediterranean basin.

This species is mostly autogamous and produces spiny dehiscent fruits typically containing 5-8

seeds each. Fruits mostly disperse close to the maternal individual, as they remain on the withered

mother plant until the end of the life cycle, but they can also be passively dispersed by grazers when

attaching to their fur.

Previous studies on a closely related annual species, *Medicago truncatula*, which has a similar

reproductive and dispersal strategy as *M. minima* and shares its same habitat, found that *M.*

truncatula is characterized by a well-defined population structure in which densely populated kin

patches alternate with more heterogeneous matrix of different genotypes (Bonnin, Ronfort,

Wozniak, & Olivieri, 2001; Siol, Bonnin, Olivieri, Prosperi, & Ronfort, 2007). Due to the strong

similarity in life-history traits between the two species, we expect that they also exhibit a similar

population structure.

For the current experiment, four genotypes of *M. minima* were sampled at four sites in the garrigue

vegetation North of Montpellier, southern France. Distance between sampling sites ranged between

500 m and 2 km (Ehlers et al., 2016). The same genotypes were previously found to have higher

survival in kin groups than non-kin groups, indicative of reduced kin competition (Ehlers et al.,

2016). Seeds from fruits sampled in the field were grown for one generation in a greenhouse to

remove maternal effects. Because *M. minima* is selfing, seedlings emerging from the same seed

pod, which we refer to as kin, are full sibs.

Experimental set up:

We conducted two greenhouse experiments to test the effects of neighbour plant relatedness (i.e. kin

or non-kin) on below- and aboveground growth of *Medicago minima* plants. In both experiments,

plants were grown in a mix of sand, greenhouse soil and vermiculite in proportions 1:1:1. To ensure

germination, seeds were scarified with sandpaper and placed on wet filter paper for 48 hours before

sowing on to seed trays placed in a greenhouse. Greenhouse temperature was set to a minimum

temperature of 15 C degrees. After germination, seedlings were transplanted into pots, and care was

taken to match similar sized seedlings for the same pot.

Experiment 1: We set up a pairwise comparison experiment in which each of the four target

genotypes was grown with either a kin or a non-kin neighbour. To account for the effect that genetic

variation in competitive ability may have on the outcome of genotype-by-genotype interactions, we

combined each genotype with two different non-kin genotypes, thus resulting in one kin treatment

and two non-kin treatments for each focal genotype (Table 1). Each of these genotype combinations

was replicated three times.

Table 1: combinations of kin (K) and non-kin (NK) treatments used for experiment 1 and experiment 2.

Additional kin and two non-kin treatments where activated carbon was added are marked with a "c". . The top

row indicates the focal genotypes (in bold) while the column indicates the identity of the neighbour or

surrounding genotypes

	Focal	102	114	121	130
	102	K + c	NK + c		NK
Neighbour	**114**	NK	K	NK	
	121	NK		K	NK
	130		NK	NK + c	K +c

Plants' pairs were sowed at a distance of 2.5 cm in transparent plexiglass containers (30 cm x 25 cm

x 3 cm) that allowed to observe and measure the direction of root growth. We kept the plexiglass

containers tilted at a 50°angle, thus forcing roots to grow against the downward-facing surface. To

avoid exposure of roots to light, we kept the downward-facing side of each container covered with a

black plastic slate that could be removed when recording root growth. Growth and direction of roots

were monitored twice a week. A transparent millimetre plastic foil glued over the container's

surface allowed for accurately measuring root growth.

After 6 weeks, we harvested above- and belowground biomass of each plant. Roots were carefully

separated from soil, washed, oven-dried at 70 C degrees for 72 hours. Above- and belowground

biomass was weighed separately on a high precision scale.

Experiment 2: In this experiment, we sowed four individuals of *M. minima* in 20 cm diameter pots.

One focal genotype was placed at the centre of the pot and was surrounded by three equidistant

individuals. Focal genotypes were exposed to three different treatments based on surrounding

plants' identity: 1) kin, where the surrounding individuals belonged to the same genotype as the

focal; 2) mix, where the surrounding individuals belonged to three different genotypes (i.e.

surrounding genotypes were different from each other and the focal); 3) non-kin, where the three

surrounding individuals belonged to the same genotype, but were different from the focal.

Each of the four genotypes was exposed to kin, mix and two different combinations of non-kin

treatments. The genotype combinations used for non-kin treatments were the same used for

experiment 1 (Table 1). Treatments were replicated four times for each genotype (N pots = 64).

Additionally, we exposed two selected kin and two non-kin treatments (three replicates for each) to

the application of activated carbon, by adding 20 ml activated carbon/l of soil (N pots = 12).

Activated carbon is known to dampen the effects of root exudates (Callaway & Aschehoug, 2000;

Callaway, Ridenour, Laboski, Weir, & Vivanco, 2005), which mediate neighbour identity

recognition (Semchenko et al., 2014). Thus, by comparing pots treated with activated carbon and

pots that were untreated, we aimed at testing whether plants modify their growth when unable to

recognize the identity of their neighbours.

We recorded diameter and number of leaves of focal plants weekly to estimate early growth rate.

For the first three weeks, we measured diameter for both focal and surrounding plants, and

afterwards, we restricted measures only to focal plants. Due to technical difficulties, we were able

to count the number of leaves on focal and surrounding plants only for the first three weeks. After

16 weeks, aboveground biomass of all plants (central and surrounding) was harvested individually,

dried at 70 C degrees for 72 hours and weighed using a high-precision scale (Mettled Toledo

AX504, d = 0.1mg)

Statistical analyses

Experiment 1: For each focal plant, we estimated root behaviour of focal plants by calculating the

difference between the number of lateral roots growing away from and growing towards the

neighbour, standardized by the plant's total number of roots. The same calculation was applied to

root length, which was measured in cm. We applied linear models to test the effect of focal

genotype, treatment (kin vs. non-kin) and their interaction on root behaviour (creating separate

models for the number of roots and root length).

Subsequently, we tested the effects of focal genotype identity and treatment (kin vs. non-kin) on

above- and below-ground biomass (using generalized linear models with gamma distribution and

identity link function) and on the roots-to-total biomass ratio (using linear models).

Experiment 2: We calculated early radial growth and early leaves growth as the difference in

respectively maximum diameter and number of leaves between week 2 and week 1. We applied

linear regression models to test how early leaves growth of focal and surrounding individuals varied

across treatments. We also applied linear models to test for the effects of focal genotype, treatment

and their interaction on early radial growth and final biomass of focal plants.

To test how total pot biomass and within-pot variance in biomass among plants changed across

treatments, we used linear models and applied log-transformation to both total biomass and biomass

variance. We also used linear models to test how total pot biomass and within-pot variance in

biomass changed across treatments (kin vs. non-kin) in response to activated carbon. For this

analysis, total biomass was square-root transformed and within-pot variance was log-transformed.

Effects of focal genotype, treatment and surrounding genotype identity on focal plants biomass

were tested using linear models.

All statistical analyses were conducted using the statistical software R, version 3.5.3 (R Core Team,

2019). Post-hoc tests were carried out using the package "emmeans" (Lenth, 2019).

Results

Experiment 1: We found that root growth (both number and length of roots) away vs. toward a

neighbour varied across focal genotypes and treatments (Table 2, Figure 1A). Two genotypes (114

and 121) grew more roots away from their neighbour when this was kin compared to non-kin, a

response consistent with avoiding root competition with kin. In contrast, genotype 102 grew more

roots toward kin neighbours compared to non-kin, whereas no sign of a differential root growth in

response to neighbour identity was observed for the fourth genotype (130). These results were

consistent for the number of roots (Table 2, Figure 1A) and the length of roots (Table 2, Figure 1B).

Figure 1: Boxplots of A) Relative number of roots growing away *vs*. toward the neighbour, B) Relative length of roots growing away *vs*. toward the neighbour (experiment 1). Each dot represents one individual replicate. Positive values indicate that higher number of roots and higher root growth (length measured in cm) were recorded away *vs*. toward the neighbour. Panels represent focal genotypes (102, 114, 121, 130) separately, and on each panel we report the results of post-hoc tests for the effect of treatment (kin *vs*. non-kin).

212 **Table 2: Results of linear models conducted on the number and the total length of roots growing away *vs.* toward**

213 **a neighbour, in response to identity of focal genotype, treatment (kin or non-kin) and their interaction**

214 **(Experiment 1)**

Number of roots			
term	**df**	**F test**	**p.value**
focal	3	4.42	0.012
treatment	1	13.25	0.001
focal: treatment	3	7.68	0.001
Length of roots			
term	**df**	**F test**	**p.value**
focal	3	3.4	0.031
treatment	1	14.6	0.001
focal: treatment	3	8.37	<0.001

216 We did not find any effect of treatment (kin vs. non-kin) on the production of aboveground biomass

217 (Wald-Chi-Square = 1.39, p = 0.23), belowground biomass (Wald-Chi-Square = 0.35, p = 0.55) or

218 on root-to-total biomass ratio (F1 = 0.14, p = 0.70). Identity of focal genotype had an effect only on

219 belowground biomass (Wald-Chi-Square = 14.40, p = 0.002) where focal genotypes 102 and 130

220 produced the lowest, and 114 and 121 the largest amount of root biomass (Figure 2).

Figure 2: Boxplots of A) aboveground biomass, B) belowground biomass, C) root-to-total biomass ratio for the

focal genotypes (Experiment 1). Dots represent single focal plants.

Experiment 2: Linear regression for growth of focal and surrounding leaves (Figure 3A-C) revealed

different slopes across treatments. We found a significant positive relationship between leaves

growth of focal and surrounding plants in kin treatments (F_1=8.94, p = 0.009), whereas we did not

find any significant relationship in mix (F_1= 0.19, p = 0.64) and non-kin (F_1= 0.15, p = 0.69)

treatments. Early radial growth was different across genotypes (Table 3, Figure 3C). Genotype 102

attained the lowest growth rate and genotype 121 the fastest growth rate, but no significant effect of

treatment was found (Table 3).

232   **Table 3: results of linear models for early radial growth of focal individuals in response to focal genotype and**

233   **treatment (Experiment 2)**

term	df	F test	p.value
focal	3	3.03	0.037
treatment	2	2.32	0.108
focal: treatment	6	0.95	0.469

Figure 3: A-C) Regression for number of leaves grown during the first two weeks (experiment 2) on focal and

surrounding genotypes across treatments (kin, non-kin and mix); D) early radial growth of focal individuals

across focal genotypes.

Analysing pot-level aboveground biomass, we found that total pot biomass (Figure 4A) was very

similar across treatments ($F_2 = 0.26$, $p = 0.7$), in contrast the within-pot variance in biomass (Figure

4B) varied substantially across treatments ($F_2 = 3.59$, $p = 0.03$). The variance in biomass among

plants within pots in the mix treatments was three times higher than in kin treatments.

Table 4: results of linear models on within pot biomass variance and total pot biomass in response to treatment

(kin *vs.* non-kin), activated carbon application, and their interaction (Experiment 2)

Within-pot variance in biomass

term	df	F test	p.value
treatment	1	0.4	0.539
carbon	1	4.7	0.048
treatment: carbon	1	3.37	0.088

Total pot biomass

term	df	F test	p.value
treatment	1	0.5	0.492
carbon	1	2.64	0.126
treatment: carbon	1	5.97	0.028

Figure 4: A, B) Plots representing mean total biomass (±1SE) and mean within-pot variance in biomass (±1SE) for untreated soil; C, D) Boxplots comparing total pot biomass and within-pot variance in biomass for kin and non-kin treatments in untreated soil vs. soil with activated carbon. Dots indicate single pots (experiment 2).

The differences among treatments in within-pot variance changed remarkably in pots treated with activated carbon. Total biomass in pots treated with activated carbon was higher compared to untreated pots (Table 4, Figure 4C). However, this effect was observed only in kin treatments as indicated by the significant treatment x carbon interaction. Similarly, activated carbon was associated with a large shift in within-pot variance in biomass for kin treatments (Table 4, Figure 4D). In kin treatments, within-pot variance in biomass for pots treated with activated carbon was on

average 2.7 times larger than in untreated kin pots, and comparable to within-pot variance in

biomass found in mix treatments in untreated soil (Figure 4B, D).

We found no effect of focal genotype identity ($F_3 = 2.31$, $p = 0.08$) or treatment ($F_2 = 0.17$, $p = 0.83$)

on aboveground biomass of focal plants. However, to understand if biomass of focal plants varied

with genotype identity of surrounding plants (rather than treatment), we analysed the effect of

surrounding and focal genotype identity on the biomass of focal plants, in kin and non-kin

treatments. This showed a significant effect of identity of surrounding genotype ($F_3 = 2.93$, $p =$

0.04), but not of focal genotype ($F_3 = 0.59$, $p = 0.62$). Focal plants surrounded by genotype 102

attained the highest biomass and focal plant surrounded by genotype 121 the lowest (Figure 5).

Figure 5: Boxplot representing biomass of focal plants in response to the identity of surrounding genotypes in

kin and non-kin treatments. Dots represent single focal plants (experiment 2).

Discussion

The results of our experiments provide support for both reduced competition among kin and genetic

variation in competitive ability. Importantly, had we not specifically observed root behaviour in

experiment 1, and added activated carbon to pots in the mini-communities of experiment 2, we

would not have been able to conclude that kin recognition alters competitive growth.

Here we discuss separately the results pointing toward either mechanism (i.e. reduced competition

among kin and genetic variation for competitive ability), and how these results may help better

understand the different outcomes of intraspecific interactions. Our plants did not produce flowers

before the end of either experiment. Hence, we were not able to test for fitness differences in seed

set among genotypes, and we based our interpretations on the observed variations in root growth

behaviour, biomass and growth among interacting plants.

Reduced competition among kin

Our first experiment tested whether plants of *Medicago minima* altered their root growth toward

their neighbour depending on whether the neighbour was kin or non-kin. We found that the

response to neighbour relatedness was genotype-specific (Figure 1). In general, all genotypes grew

more roots away from than toward their neighbours, which can be attributed to our experimental

design, where two plants were grown close to each other leaving most space for root growth in the

direction away from the neighbour.

Two of the focal genotypes showed a response consistent with avoiding kin root competition.

Genotype 121 and genotype 114 showed differential growth response to kin vs. non-kin and grew

proportionally fewer roots toward their kin neighbours. In contrast, genotype 102 grew more roots

toward its kin neighbour relative to non-kin. This genotype was also characterised by the slowest

growth rate (Figure 2), and it is possible that for this genotype, growing with kin of similar slow

growth rate, simply resulted in more space to proliferate roots in direction of kin relative to fast-

growing non-kin neighbours. Our findings of reduced root growth toward kin neighbours are

consistent with previous studies also finding reduced root competition among kin (Dudley & File,

2007; Bhatt et al., 2010).

In our second experiment, we exposed focal genotypes located at the centre of a pot, to three

neighbours that were either kin, non-kin or belonged to different genotypes (mix). We found a

positive relationship in early leaves growth of focal and surrounding plants in kin treatments, but

not in non-kin and mix treatments (Figure 3 A-C). This positive relationship, found only in kin

treatments, could be the result of reduced competition among kin. However, such response could

also be due to low variation in competitive ability among identical genotypes, leading to similar

growth rates of focal and surrounding plants in kin treatments, and in a weak or even negative

relationship in growth rates in non-kin and mixed treatments, where fast-growing and potentially

more competitive genotypes could impede the growth of slow-growing ones.

Further support for reduced competition among kin came from the results of total pot biomass and

within-pot variance in biomass. We found no differences in total pot biomass among treatments

(Figure 4A), indicating that regardless of treatment, plants grew to fill all the space and resource

available within pots. However, we found large differences in how biomass was distributed among

individual plants within the same pot (i.e. within-pot variance in biomass) depending on treatment

(Figure 4B). This within-pot variance was the smallest in kin treatments and the largest in mix

treatments, again suggesting reduced or very similar competition among plants in kin pots.

Importantly, the application of activated carbon revealed a very different response in both total

biomass and particularly in within-pot variance in biomass. Application of activated carbon is

known to inhibit the action of root exudates (Callaway & Aschehoug, 2000; Callaway et al., 2005),

which have been demonstrated to mediate neighbour identity recognition in plants (Yang, Li, Xu, &

Kong, 2018). In kin treatments, the application of activated carbon led to 2.7 fold increase of

within-pot biomass variance as opposed to a 1.7 fold decrease in biomass variance for non-kin

treatments (Figure 4D). Similarly, the total pot biomass in kin treatments increased by 19% pots

treated with active carbon compared to untreated soil, whereas no change was observed for non-kin

treatments (Figure 4C).

The within-pot variance in biomass observed for kin treatments in activated carbon was comparable

to the within-pot variance of mix treatments in untreated soil. This strongly suggests that when

plants cannot detect the identity of their kin neighbours, they become more competitive. Therefore,

reduced variance in biomass among plants in untreated kin pots is most likely caused by kin

recognition and reduced competition among kin. If this was a mere consequence of similarity in

competitive ability among identical genotypes, we would not except such a strong shift in the

variance of biomass among plants in kin pots treated with activated carbon.

Although the reliability of activated carbon has been criticized as it may stimulate plants growth

(Lau et al., 2008), our results are in line with previous studies (e.g. Biedrzycki, Jilany, Dudley, &

Bais, 2010; Semchenko, Saar, & Lepik, 2014), which show that suppression of communication via

root-exudates affects plant competition and alters the response to neighbour identity.

Genetic variation in competitive ability

Our results also show the presence of genotypic variation in competitive ability. For early radial

growth of plants (Figure 3D), genotype 121 showed the fastest and genotype 102 the slowest

growth rates. Consistent with this, we found that biomass of focal plants varied significantly

depending on genotype identity of the surrounding plants (Figure 5). Focal plants attained the

highest biomass when surrounded by genotype 102 (the genotype with the slowest growth), and the

lowest biomass when surrounded by genotype 121 (the fastest-growing genotype). This is

consistent with other studies reporting intraspecific variation for competitive ability via variation in

growth rate and ability to suppress the growth of neighbours (e.g. Masclaux et al., 2010; Cahill,

Kembel, & Gustafson, 2005).

Additional support for genotypic variation for competitive ability came from the significantly

higher variance in biomass among plants within pots in mix treatments. This suggests a high

asymmetric competition among different genotypes, consistent with genetic variation in competitive

ability (Donohue, 2003; Simonsen et al., 2014).

Intraspecific interactions as a result of both reduced kin competition and variation in competitive

ability

Our results suggest that reduced competition among kin may not be independent of competitive

ability. Across both experiments, genotype 102 showed the weakest response in terms of reduced

competition towards kin, while attaining the lowest values of belowground biomass (experiment 1)

and the smallest competitive effect on non-kin focal plants (experiment 2). Conversely, genotype

121, which showed the strong signals of reduced competition among kin, showed the highest

growth rate, obtained the larger biomass (experiment 1), and exerted the strongest competitive

effect on focal plants (experiment 2). Similarly, genotype 114 also showed signals for reduced

competition among kin and was among the fastest-growing and more competitive genotypes

(experiment 1). This suggests that reduced competition among kin could be more pronounced in

strongly competitive genotypes that are also more likely to interfere with their kin neighbours. This

would be consistent with previous studies showing that kin recognition is a density-dependent

response (Lepik et al., 2012).

Our study is one of few (see also Biernaskie, 2010) showing experimentally that reduced kin

competition and genetic variation in competitive ability act simultaneously to shape the outcome of

intraspecific interactions.

In plant populations characterized by a strong spatial genetic structure, kin recognition and reduced

kin competition may favour the growth of kin genotypes, creating positive density-dependent kin

interactions. This can result in higher resistance to the invasion of more distantly related

conspecifics and even of other species (Molofsky & Bever, 2002). Indeed, it was previously found

that *M. minima* had higher survival in kin interactions compared to both non-kin and heterospecific

interactions (Ehlers et al., 2016). Positive density-dependent interactions in plants can influence

both the distribution of intraspecific genetic variation and local species co-existence (Molofsky &

Bever, 2002; Laird & Schamp, 2006) and can be important for invasion success and range

expansion (Crandall & Knight, 2015).

Intraspecific interactions with more distantly related conspecifics should, on the other hand, be

governed by variation in competitive ability among interacting genotypes (Fridley, Grime, &

Bilton, 2007; Fridley & Grime, 2010). If highly competitive genotypes do not reduce competition

with kin, they may ultimately be subject to negative density-dependence, when increasing success

results in more encounters with highly competitive kin genotypes.

Our finding of genotypes that are both highly competitive against non-kin, and reduce their

competition towards kin suggests a potentially highly successful strategy unless curbed by trade-

offs and costs of plastic growth response to neighbour's identity (Crepy & Casal, 2014).

In conclusion, we find that within a species, intraspecific interactions are shaped by both reduced

competition with kin and by variation in growth rate in response to non-kin neighbours. Had we not

observed root behaviour, and suppressed neighbour recognition by adding activated carbon, the

specific kin response might have gone unnoticed as the symmetric growth in kin pots could also be

explained by genetic variation in competitive ability. When predicting the outcome of intraspecific

interactions, these findings emphasize the importance of knowing the degree of genetic substructure

in populations as local positive density-dependent interactions may be expected in clusters of kin,

but not of non-kin.

Acknowledgements: This work was funded by a grant from the Danish Council of Independent

Research (grant number 6108-00200B) to BKE. We are grateful to L. Lauridsen and J. Rytter for

their help in the greenhouse.

Authors contribution: BKE, MJ and CT planned the experiments, MJ and CT carried out the

experiments, ST carried out the statistical analyses and wrote the first draft of the manuscript, all

authors contributed to reviewing following versions.

References:

Bhatt, M. V., Khandelwal, A., & Dudley, S. A. (2010). Kin recognition, not competitive

interactions, predicts root allocation in young *Cakile edentula* seedling pairs. *New Phytologist*,

189(4), 1135–1142. doi:10.1111/j.1469-8137.2010.03548.x

Biedrzycki, M. L., Jilany, T. A., Dudley, S. A., & Bais, H. P. (2010). Root exudates mediate kin

recognition in plants. *Communicative & Integrative Biology*, *3*(1), 28–35.

Biernaskie, J. M. (2010). Evidence for competition and cooperation among climbing plants.

Proceedings of the Royal Society B: Biological Sciences, *278*(1714), 1989–1996.

doi:10.1098/rspb.2010.1771

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., Vasseur, D.

A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology &*

Evolution, *26*(4), 183–192. doi:10.1016/j.tree.2011.01.009

Bonnin, I., Ronfort, J., Wozniak, F., & Olivieri, I. (2001). Spatial effects and rare outcrossing

events in *Medicago truncatula* (Fabaceae). *Molecular Ecology*, *10*(6), 1371–1383.

Burns, J. H., & Strauss, S. Y. (2012). Effects of competition on phylogenetic signal and phenotypic

plasticity in plant functional traits. *Ecology*, *93*(sp8), S126–S137.

Cahill, F. J. J., Kembel, S. W., & Gustafson, D. J. (2005). Differential genetic influences on

competitive effect and response in *Arabidopsis thaliana*. *Journal of Ecology*, *93*(5), 958–967.

Callaway, R. M., & Aschehoug, E. T. (2000). Invasive plants versus their new and old neighbors: a

mechanism for exotic invasion. *Science*, *290*(5491), 521–523.

Callaway, R. M., Ridenour, W. M., Laboski, T., Weir, T., & Vivanco, J. M. (2005). Natural

selection for resistance to the allelopathic effects of invasive plants. *Journal of Ecology*, *93*(3), 576–

583.

Cheplick, G. P., & Kane, K. H. (2004). Genetic relatedness and competition in *Triplasis purpurea*

(Poaceae): resource partitioning or kin selection? *International Journal of Plant Sciences*, *165*(4),

623–630.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual Review of Ecology*

and Systematics, *31*(1), 343–366. doi:10.1146/annurev.ecolsys.31.1.343

Crandall, R., & Knight, T. M. (2015). Positive frequency dependence undermines the success of

restoration using historical disturbance regimes. *Ecology Letters*, *18*(9), 883–891.

Crepy, M. A., & Casal, J. J. (2014). Photoreceptor-mediated kin recognition in plants. *New*

Phytologist, *205*(1), 329–338. doi:10.1111/nph.13040

Donohue, K. (2003). The influence of neighbor relatedness on multilevel selection in the Great

Lakes sea rocket. *The American Naturalist*, *162*(1), 77–92. doi:10.1086/375299

Dudley, S. A., & File, A. L. (2007). Kin recognition in an annual plant. *Biology Letters*, *3*(4), 435–

438.

Ehlers, B. K., & Bilde, T. (2019). Inclusive fitness, asymmetric competition and kin selection in

plants. *Oikos*, *128*(6), 765–774. doi:10.1111/oik.06390

Ehlers, B. K., Damgaard, C. F., & Laroche, F. (2016). Intraspecific genetic variation and species

coexistence in plant communities. *Biology Letters*, *12*(1), 20150853.

Ehlers, B. K., David, P., Damgaard, C. F., & Lenormand, T. (2016). Competitor relatedness,

indirect soil effects and plant coexistence. *Journal of Ecology*, *104*(4), 1126–1135.

doi:10.1111/1365-2745.12568

File, A. L., Murphy, G. P., & Dudley, S. A. (2011). Fitness consequences of plants growing with

siblings: reconciling kin selection, niche partitioning and competitive ability. *Proceedings of the*

Royal Society B: Biological Sciences, *279*(1727), 209–218.

Fridley, J. D., & Grime, J. P. (2010). Community and ecosystem effects of intraspecific genetic

diversity in grassland microcosms of varying species diversity. *Ecology*, *91*(8), 2272–2283.

Fridley, J. D., Grime, J. P., & Bilton, M. (2007). Genetic identity of interspecific neighbours

mediates plant responses to competition and environmental variation in a species-rich grassland.

Journal of Ecology, *95*(5), 908–915. doi:10.1111/j.1365-2745.2007.01256.x

Hamilton, W. D. (1964). The genetical evolution of social behavior, I and II. J. *Theor Biol*, *7*, 1–52.

Hart, S. P., Schreiber, S. J., & Levine, J. M. (2016). How variation between individuals affects

species coexistence. *Ecology Letters*, *19*(8), 825–838. doi:10.1111/ele.12618

Kelly, J. K. (1996). Kin selection in the annual plant *Impatiens capensis*. *The American Naturalist*,

147(6), 899–918.

Laird, R. A., & Schamp, B. S. (2006). Competitive intransitivity promotes species coexistence. *The*

American Naturalist, *168*(2), 182–193. doi:10.1086/506259

Lau, J. A., Puliafico, K. P., Kopshever, J. A., Steltzer, H., Jarvis, E. P., Schwarzländer, M.,

Hufbauer, R. A. (2008). Inference of allelopathy is complicated by effects of activated carbon on

plant growth. *New Phytologist*, *178*(2), 412–423. doi:10.1111/j.1469-8137.2007.02360.x

Lenth, R. (2019). *Emmeans: Estimated marginal means, aka least-squares means*. Retrieved from

https://CRAN.R-project.org/package=emmeans

Lepik, A., Abakumova, M., Zobel, K., & Semchenko, M. (2012). Kin recognition is density-

dependent and uncommon among temperate grassland plants. *Functional Ecology*, *26*(5), 1214–

1220. doi:10.1111/j.1365-2435.2012.02037.x

MacArthur, R., & Levins, R. (1967). The limiting similarity, convergence, and divergence of

coexisting species. *The American Naturalist*, *101*(921), 377–385.

Masclaux, F., Hammond, R. L., Meunier, J., Gouhier-Darimont, C., Keller, L., & Reymond, P.

(2010). Competitive ability not kinship affects growth of *Arabidopsis thaliana* accessions. *New*

Phytologist, *185*(1), 322–331. doi:10.1111/j.1469-8137.2009.03057.x

Milla, R., Forero, D. M., Escudero, A., & Iriondo, J. M. (2009). Growing with siblings: a common

ground for cooperation or for fiercer competition among plants? *Proceedings of the Royal Society*

B: Biological Sciences, *276*(1667), 2531–2540. doi:10.1098/rspb.2009.0369

Molofsky, J., & Bever, J. D. (2002). A novel theory to explain species diversity in landscapes:

Positive frequency dependence and habitat suitability. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *269*(1508), 2389–2393.

R Core Team. (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R

Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Ronce, O. (2007). How does it feel to be like a rolling stone? Ten questions about dispersal

evolution. *Annu. Rev. Ecol. Evol. Syst.*, *38*, 231–253.

Semchenko, M., Saar, S., & Lepik, A. (2014). Plant root exudates mediate neighbour recognition

and trigger complex behavioural changes. *New Phytologist*, *204*(3), 631–637.

doi:10.1111/nph.12930

Simonsen, A. K., Chow, T., & Stinchcombe, J. R. (2014). Reduced plant competition among kin

can be explained by Jensen's inequality. *Ecology and Evolution*, *4*(23), 4454–4466.

doi:10.1002/ece3.1312

Siol, M., Bonnin, I., Olivieri, I., Prosperi, J. M., & Ronfort, J. (2007). Effective population size

associated with self-fertilization: lessons from temporal changes in allele frequencies in the selfing

annual *Medicago truncatula*. *Journal of Evolutionary Biology*, *20*(6), 2349–2360.

doi:10.1111/j.1420-9101.2007.01409.x

Vellend, M., & Geber, M. A. (2005). Connections between species diversity and genetic diversity.

Ecology Letters, *8*(7), 767–781.

Weiner, J., Campbell, L. G., Pino, J., & Echarte, L. (2009). The allometry of reproduction within

plant populations. *Journal of Ecology*, *97*(6), 1220–1233.

Weiner, J., & Thomas, S. C. (1986). Size variability and competition in plant monocultures. *Oikos*,

211–222.

West, S. A., Pen, I., & Griffin, A. S. (2002). Cooperation and competition between relatives.

Science, *296*(5565), 72–75. doi:10.1126/science.1065507

Yang, X.-F., Li, L.-L., Xu, Y., & Kong, C.-H. (2018). Kin recognition in rice (*Oryza sativa*) lines.

New Phytologist, *220*(2), 567–578. doi:10.1111/nph.15296

Young, J. (1981). Sib competition can favour sex in two ways. *Journal of Theoretical Biology*,

88(4), 755–756.