Assessing auditory processing endophenotypes associated with Schizophrenia in individuals with 22q11.2 Deletion Syndrome

Ana A. **Francisco** [1,2], John J. **Foxe** [1,2,3], Douwe J. **Horsthuis** [1], Danielle **DeMaio** [1], Sophie **Molholm** [1,2,3]

[1] *The Cognitive Neurophysiology Laboratory, Department of Pediatrics, Albert Einstein College of Medicine, Bronx, New York, USA.*

[2] *Department of Neuroscience, Rose F. Kennedy Center, Albert Einstein College of Medicine, Bronx, New York, USA.*

[3] *The Cognitive Neurophysiology Laboratory, Department of Neuroscience, The Ernest J. Del Monde Institute for Neuroscience, School of Medicine and Dentistry, University of Rochester, Rochester, New York, USA.*

ABSTRACT

22q11.2 Deletion Syndrome (22q11.2DS) is the strongest known molecular risk factor for schizophrenia. Scalp recorded electrical brain responses to auditory stimuli have been studied extensively in schizophrenia and described as potential biomarkers of vulnerability to psychosis. Here we sought to understand whether these brain responses, along with cognitive factors associated with schizophrenia, might aid in differentiating individuals with 22q11.2DS as a function of psychotic symptoms, and ultimately serve as early signals of higher risk for schizophrenia in this population. Using a passive duration oddball paradigm and high-density electrophysiology, we tested basic auditory processing in a group of 21 adolescents and adults with 22q11.2DS with varying degrees of psychotic symptomatology and neurotypical age-matched controls. We examined whether the N1 and mismatch negativity (MMN), typically reduced in idiopathic schizophrenia, significantly correlated with psychotic symptomatology, IQ, and working memory. At the group level of analysis, individuals diagnosed with 22q11.2DS presented *increased* N1 responses and *similar* MMN responses to their neurotypical peers. Correlation analyses revealed that poorer working memory was associated with decreased MMN amplitude, and that this was largely driven by the 22q11.2DS group. We suggest that enhanced auditory sensory processing is general to 22q11.2DS, whilst poorer working memory is related to vulnerability for psychosis and smaller MMNs in 22q11.2DS.

Keywords: EEG, Auditory Evoked potential, Copy Number Variation, Event-related potential, Mismatch Negativity, Psychosis.

INTRODUCTION

22q11.2 Deletion Syndrome (22q11.2DS henceforth; also named DiGeorge syndrome or velo-cardio-facial syndrome) results from a hemizygous microdeletion of approximately 1.5 to 3 megabases on the long arm of chromosome 22. The deleted region contains about 60 known genes, some of which are highly expressed in the brain and known to affect early neuronal migration and cortical development[1, 2]. 22q11.2DS is the most common chromosomal microdeletion disorder, with prevalence estimates ranging from 1 per 1000 to 1 per 4000 live births[3, 4].

The phenotypic expression of 22q11.2DS is highly variable and ranges from severely life-threatening to less severe conditions[5]. Its clinical presentation includes variable developmental delays, cognitive deficits and neuropsychiatric conditions, and multi-organ dysfunction such as cardiac and palatal abnormalities[6, 7]. Cognitively, 22q11.2DS is characterized by deficits in executive function[8-11], nonverbal memory[12, 13], visuospatial[14-16] and visual-motor[17] processing, and working memory[18]. Approximately 60% of individuals diagnosed with 22q11.2DS meet criteria for at least one psychiatric diagnosis[19-21]—and the development of psychosis is one of the most significant concerns for parents of children with 22q11.2DS: About 30% of individuals identified with the deletion go on to develop schizophrenia[22-24]–an approximately 10-fold increase in risk compared to individuals with other developmental disabilities[25]—and more than half of adolescents with the syndrome experience schizotypical traits and transient psychotic experiences[26].

Idiopathic and 22q11.2DS-associated schizophrenia present a similar clinical path leading to psychosis[27], with subtle social deficits and poor neurocognitive performance preceding its onset both in the general population[28, 29] and in 22q11.2DS[30]. The clinical presentation of schizophrenia does not appear to differ either[31], with deficits in cognition and perception, and positive (hallucinations, delusions, thought disorder) and negative (flat affect, alogia, avolition and anhedonia) symptoms[32]. However, neurocognitive features are more severe in 22q11.2DS-associated schizophrenia, likely reflecting poorer baseline cognitive function[33]. These clinical/cognitive similarities are accompanied by high concordance of neuroanatomic correlates[34, 35 36-39], suggesting that common cerebral alterations may underlie psychotic symptomatology in both populations.

Reducing the duration of untreated psychosis, before irreversible pathological brain changes take place[40], is known to lead to better functional outcomes in patients with psychotic disorders such as schizophrenia[41, 42], strongly motivating the goal of early identification prior to frank disease onset. Since clearly observable behavioral changes are often the product of processes that have started occurring in the brain significantly earlier, the direct and reliable measurement of functional brain activity and the consequent potential to detect neural vulnerability prior to behaviorally observable symptoms is key to enabling interventions focused on prevention rather than on treatment. Sensory and cognitive auditory evoked potentials (AEP) such as the N1 and the mismatch negativity (MMN) have been identified as

potential biomarkers for vulnerability, conversion, and progression of psychosis in schizophrenia. The auditory N1 is the first prominent negative AEP[43], and reflects neural activity generated in and around primary auditory cortex[44]. Several studies demonstrate a reduction of the auditory N1 in schizophrenia, not only in chronic patients, but also in individuals at-risk for psychosis and in first-episode patients[45-49]; but see[50] for a review on contradictory evidence). The MMN occurs when a repeating stimulus (the standard) in an auditory stream is replaced by a deviant stimulus. Occurring typically 100 to 200 ms following the deviant event, the MMN is thought to reflect largely preattentive neural processes underlying detection of a pattern violation and updating of a representation of a regularity in the auditory environment[51-53]. MMN amplitude has been shown to be reduced in schizophrenia (for reviews, see[54, 55]) in at-risk individuals[56-60], recent onset[57, 58, 61-63] and chronic patients[57, 61, 64-71].

While previous studies have measured the N1 and MMN in 22q11.2DS, the limited results available have yielded inconsistent findings[72, 73]. Such inconsistencies might reflect the relatively high degree of phenotypic variability that is inherent to this population. Here, we sought to measure the relationship between auditory brain function and cognitive function, focusing on cognitive abilities known to be vulnerable in this population. We also set out to test for potential associations between MMN and N1 amplitudes and the number of psychotic symptoms expressed in the 22q11.2DS cohort. Given the working memory difficulties previously described in 22q11.2DS, stimulus onset asynchronies (SOAs) were parametrically varied to impose different levels of demand on the sensory memory system[74, 75].

METHODS

Participants

Twenty-one participants diagnosed with 22q11.2DS (age range: 14-35 years old; 12 with at least one psychotic symptom) and 21 neurotypical age-matched controls (NT) (age range: 14-38 years old) were recruited. Exclusionary criteria for the neurotypical group included hearing impairment, developmental and/or educational difficulties, neurological problems, and the presence of psychotic symptomatology. Exclusionary criteria for the 22q11.2DS group included hearing impairment and current neurological problems. Participants passed a hearing test (below 25dB HL for 500, 100, 2000, 4000Hz) performed on both ears using a *Beltone Audiometer* (Model 112). All participants signed an informed consent approved by the Institutional Review Board of the Albert Einstein College of Medicine and were monetarily compensated for their time. All aspects of the research conformed to the tenets of the Declaration of Helsinki.

Experimental Procedure and Stimuli

Testing was carried out over a 2-day period and included cognitive testing and EEG recording. IQ and working memory measures were computed from the Wechsler Adult Intelligence Scale, WAIS-IV[76] or the Wechsler Intelligence Scale for Children, WISC-V[77]. To quantify the number of psychotic symptoms, either the Structured Clinical Interview for DSM-5, SCID-V[78] or the Structured Clinical Interview for DSM-IV Childhood Diagnoses, Kid-SCID[79] was performed.

The EEG protocol focused on auditory processing, utilizing a duration-MMN oddball paradigm. Participants sat in a sound- and electrically-shielded booth (Industrial Acoustics Company Inc, Bronx, NY) and watched a muted movie of their choice on a laptop (Dell Latitude E6430 ATG or E5420M) while passively listening to regularly (85%) occurring standard tones interspersed with deviant tones (15%). These tones had a frequency of 1000 Hz with a rise and fall time of 10 ms, and were presented at an intensity of 75dB SPL using a pair of Etymotic insert earphones (Etymotic Research, Inc., Elk Grove Village, IL, USA). Standard tones had a duration of 100 ms while deviant tones were 180 ms in duration. These tones were presented in a random oddball configuration (excepting that at least two standards preceded a deviant) to yield an MMN. In three blocked conditions, the stimulus onset asynchrony (SOA) was either 450 ms, 900 ms or 1800 ms. Each SOA was presented in blocks of 4 minutes long, composed of 500, 250 or 125 trials respectively. Participants were presented with 14 blocks (2*450 ms, 4*900 ms and 8*1800 ms), resulting in a possible 1000 trials (and 150 deviants) per SOA.

Data acquisition and analysis

EEG data were acquired continuously at a sampling rate of 512 Hz from 64 locations using scalp electrodes mounted on an elastic cap and seven external channels (mastoids, temples, and nasion) (Active 2 system; Biosemi[tm], The Netherlands; 10-20 montage). Preprocessing was done using EEGLAB (version 14.1.1)[80] toolbox for MATLAB (version 2017a; MathWorks, Natick, MA). Data were downsampled to 256 Hz, re-referenced to TP8 and filtered using a 1 Hz high pass filter (0.5 Hz transition bandwidth, filter order 1690) and a 45 Hz low pass filter (5 Hz transition bandwidth, filter order 152). Both were zero-phase Hamming windowed sinc FIR filters. Bad channels were automatically detected based on kurtosis measures and rejected after visual confirmation. Artifacts were removed by running an Independent Component Analysis (ICA) to exclude components accounting for motor artifacts, eye blinks and saccades and bad channels. After ICA, the excluded channels were interpolated, using the spherical spline method. Data were segmented into epochs of -100ms to 400ms using a baseline of -100ms to 0ms. These epochs went through an artifact detection algorithm (moving window peak-to-peak threshold at 120 µV). No significant differences were found between number of trials included in the analyses per group (22q11.2DS group: average trial rejection = 5.36%; NT group: average trial rejection = 3.30%; $t = -1.08$, $df = 35.12$, $p = .29$).

The definition of the N1 and the MMN windows was based on the typical time of occurrence of the N1 and duration-MMN components, and on visual confirmation that amplitudes were maximal in these intervals. N1 amplitude was measured between 80 and 120 ms. The MMN is the difference between deviants and standards and was here measured between 200 and 240 ms (100 to 140 ms post deviance onset). Amplitude measures were taken at FCz, where signal was maximal for both groups. For the correlation analyses, a composite measure of the response was used to reduce the number of tests. This was computed by taking the average of the three ISI conditions for each individual. All *p-values* were submitted to Holm-Bonferroni corrections for multiple comparisons[81], using the *p.adjust* of the *stats* package in R[82].

RESULTS

Table 1 shows a summary of the included participants' IQ, performance on the working memory task, and number of psychotic symptoms. Working memory scores were only available for a subset of the individuals in the neurotypical group. Two-sample independent-means *t* tests were used to test for group differences. When the assumption of the homogeneity of variances was violated, *Welch* corrections were applied to adjust degrees of freedom. Statistical analyses confirmed that the groups differed significantly in IQ, working memory, and number of psychotic symptoms presented. There were significantly more females than males in each group, but no differences in sex between groups. Age did not differ between groups.

Table 1. Characterization of neurotypical and 22q11.2DS individuals included in the analyses: age, sex, IQ, working memory, and number of psychotic symptoms presented.

	Neurotypical controls	22q11.2DS	t-test	Cohen's d
Age	M=22.43; SD=7.27	M=22.67; SD=7.20	t=-0.11, df=39.40, p=.92	d=-0.03
Sex	8 males, 13 females	6 males, 15 females	-	-
IQ	M=113.75; SD=15.35	M=72.95; SD=13.19	t=9.02, df=37.16, p<.01	d=2.85
Working memory	M=103.46; SD=13.81	M=78.50; SD=15.77	t=4.94, df=25.80, p<.01	d=1.68
N symptoms	0	M=0.95; SD=1.12	t=-3.91, df=20.00, p<.01	d=-1.21

Figure 1 shows correlations between number of psychotic symptoms presented, IQ and working memory. Lower IQ (r = -.56, p < .01) and poorer working memory (r = -.59, p < .01) scores were observed in those individuals presenting more psychotic symptoms.

Figure 1. Pearson correlations between IQ scores and number of psychotic symptoms, and working memory performance and number of psychotic symptoms.

Figure 2 shows the averaged ERPs and topographies for the time windows of interest (N1 and MMN), per SOA and by group. Overall, 22q11.2DS data were characterized by increased amplitude auditory evoked responses (Figure 2B) and greater variability (Figure 2C).

Figure 2. Panel A: Upper left, averaged Standard and Deviant ERPs and Standard topographies per SOA for the NT group at FCz; Lower right, smaller plot, shows MMN, i.e., differences between standards and deviants; topographies are organized from the shorter (450 ms) to the longer (1800 ms) SOA. Panel B: Averaged ERPs and topographies per SOA for the 22q11.2DS group at FCz (smaller plot shows MMN, i.e., differences between standards and deviants); topographies are organized from the shorter (450 ms) to the longer (1800 ms) SOA. Panel C: Violin plots showing distribution of amplitudes for N1 and MMN per group and SOA at FCz. Black square indicates median.

Separate mixed-effects models were implemented to analyze the N1 and MMN data, using the *lmer* function in the *lme4* package[83] in R[82]. Mean amplitude at FCz was the numeric dependent variable. For the N1, only standards were considered. For the MMN, mean amplitude represents the difference between standards and deviants. Group (NT = −0.5, 22q11.2DS = 0.5) was a contrast-coded fixed factor, and SOA a numeric fixed factor. Subjects and SOA were added as random factors. Models were fit using the maximum likelihood criterion. *P* values were estimated using *Satterthwaite* approximations.

For the N1, no significant main effect of group was found. There was an effect of SOA: Both 900 ms ($\beta = -1.41$, SE = 0.07, $p < .001$) and 1800 ms ($\beta = -2.97$, SE =, 0.07 $p < .001$) SOAs elicited larger

responses than the 450 ms SOA. There was also an interaction between group and SOA, with the difference between the 450 ms and the 1800 ms SOA being larger in 22q11.2DS than in the neurotypical control group (β = -0.41, SE = 0.14, $p < .01$). *Post-hoc* analyses conducted to clarify the basis for this interaction revealed differences between the groups for all SOAs: 450 ms SOA (t = 2.86, df = 407.89, $p <$ .01, d = 0.29), 900 ms SOA (t = 2.68, df = 362.91, $p < .01$, d = 0.26) and 1800 ms SOA (t = 3.73, df = 414.12, $p < .01$, d = 0.36), with individuals diagnosed with 22q11.2DS presenting with increased amplitudes compared to their neurotypical peers. N1 amplitudes did not correlate significantly with IQ, working memory, or number of psychotic symptoms.

As observed in the N1 time window, there was no significant effect of group for the MMN. Once again, there was an effect of SOA, though in the opposite direction of what was expected: Both the 900 ms SOA (β = -0.46, SE = 0.08, $p < .001$) and the 1800 ms SOA (β = -0.68, SE = 0.08, $p < .001$) conditions elicited larger MMN responses than the 450 ms SOA. The effect of the interaction between group and SOA was also significant here: The difference between the 450 ms and the 1800 ms SOA was larger in the 22q11.2DS group than in the neurotypical control group (β = -0.55, SE = 0.15, $p < .001$). *Post-hoc* analyses conducted to clarify this interaction revealed differences between the groups in the 1800 ms SOA, with individuals diagnosed with 22q11.2DS appearing to present increased amplitudes when compared to their neurotypical peers. However, this result did not hold after correcting for multiple comparisons (t = 2.13, df = 333.90, p = .102). In contrast to what was observed for the N1 component, MMN amplitudes correlated significantly with working memory performance (r = -.42, p = .05), with individuals who scored lower on the working memory tasks presenting decreased MMN amplitudes (Figure 3). *Post-hoc* correlation tests examining the two groups separately were performed to clarify this relationship. This correlation appears to have been mainly driven by the 22q11.2DS group, as can be appreciated in Figure 3 B and C: While the association between working memory and MMN amplitudes was significant for the 22q11.2DS (r = -.48, p < .05), that was not the case for the neurotypical controls (r = -.56, p > .05).

Figure 3. Pearson correlation between MMN amplitude and working memory performance across groups (A) and separately for neurotypical controls (B) and individuals diagnosed with 22q11.2DS (C).

DISCUSSION

Here, we characterized early auditory sensory processing in a group of adolescents and adults with 22q11.2DS, with and without psychotic symptomatology, and related those measures to cognitive and clinical characteristics in order to assess their potential informativeness with regard to vulnerability, conversion, and progression of psychosis.

First, when compared to their neurotypical peers, individuals diagnosed with 22q11.2DS presented increased N1 amplitudes. Larger auditory evoked potentials have also been described in a

22q11.2DS mouse model: Loudness dependent P1-N1-P2 amplitudes were increased in mice with the deletion[84]. Likewise, two human studies looking at auditory processing in 22q11.2DS showed enhanced early sensory responses in 22q11.2DS as reflected in an increased N1[73, 85]. This increase has been associated to elevated activity in the anterior cingulate and dorsomedial frontal cortex[73]. It could additionally be explained by alterations in the cortical glutamate N-methyl-D-aspartate (NMDA) receptors[85]: Increased N1 responses have been reported after administration of ketamine, an NMDA antagonist[86, 87]. This enhanced N1 in 22q11.2DS sits in stark contrast with findings in schizophrenia, where N1 is instead typically reduced compared to controls[45-48, 88, 89]. These findings suggest that enlarged N1 amplitude is a marker of 22q11.2DS, rather than vulnerability for psychosis. Moreover, an enhanced N1 could reflect a lack of adaptation in 22q11.2DS. Typically, repeated presentation of an auditory stimulus leads to reductions in amplitude of the response. Inhibitory processes could thus be impaired in 22q11.2DS, leading to an overall amplified sensory response.

Second, the groups did not differ in MMN amplitude, with the exception of a larger difference between the shortest and the longest SOAs in 22q11.2DS, when compared to the control group. Previous studies on MMN in 22q11.2DS have yielded somewhat inconsistent evidence (for a review, see[90]): Whereas some studies have found reduced pitch, duration and frequency MMNs[85, 91], others failed to show differences between individuals with 22q11.2DS and their neurotypical peers[72, 92]. Such inconsistencies in MMN differences may reflect the phenotypic heterogeneity that is characteristic of 22q11.2DS, as well as, possibly, methodological and stimulus related differences. Adding to the range of findings, here we observed that whereas at characteristic stimulation rates for MMN studies the response appeared typical, at the longest SOA it increased in amplitude relative to the shortest SOA. This contrasts with an expected attenuation of MMN amplitude with increasing SOAs in this group, due to weaker memory traces (see[75]). Previous work from our lab using the same paradigm showed that increasing SOA effectively indexed weakness in maintenance of the memory trace in Rett Syndrome[74]. An increase in MMN amplitude for the 22q11.2DS group at the longest SOA is thus hard to reconcile with the extant literature and, to our knowledge, has not been previously reported.

That individuals with 22q11.2DS did not show reduced MMN responses is also at odds with evidence showing decreased amplitudes in those at-risk of developing schizophrenia[45-48, 56-60, 88]. A plausible hypothesis is that reduced auditory sensory amplitudes are only characteristic of those 22q11.2DS individuals with greatest risk for psychosis and therefore not clearly observable here, given the mixed nature of this 22q11.2DS sample (individuals with and without psychotic symptoms). And, indeed, analyses taking into consideration individual differences revealed an association between MMN amplitudes and working memory: Worse performance on working memory measures was associated with reduced MMNs. Importantly, those individuals with more psychotic symptoms presented poorer working memory and lower IQ, which may suggest an association between worse cognitive and psychiatric

profiles and MMN reductions. This would accord with evidence in 22q11.2DS studies showing associations between MMN amplitudes and working memory[91] and between working memory and psychotic symptomatology severity[93]. Moreover, it would be consistent with the conceptualization of cognitive functioning as a vulnerability marker for psychosis[94]. In particular, working memory deficits are cardinal in schizophrenia[95, 96] and have been argued as a risk factor for psychosis[97, 98].

MMN reductions, observed here for those individuals with 22q11.2DS with lower working memory scores, could be caused by deficient auditory sensory memory-trace formation[99], which has been hypothesized to reflect disturbances of glutamatergic NMDA-receptor-mediated neurotransmission[100]. One of the genes responsible for glutamatergic transmission in the brain is the proline dehydrogenase (PRODH) gene, located on chromosome 22q11. Disruptions in NMDA-mediated glutamatergic transmission have been demonstrated in rats with elevated concentrations of L-proline[101], as well as in Prodh-deficient mice[102]. A mouse model of 22q11.2DS found those mice to display increased locomotor activity in response to NMDA-receptor antagonists, but only after puberty, suggesting the potential of this specific mouse to model the impaired NMDA neurotransmission in idiopathic schizophrenia and in those individuals with 22q11.2DS who go on to develop a psychotic disorder[84]. Second, the reduced MMN may be suggestive of a dampened frontal attention-orienting function[103]. Attentional orienting has been argued to be impaired in 22q11.2DS and to be one of the mechanisms mediating the cognitive decline during the so-called prodromal stage in those individuals with the deletion presenting psychotic symptomatology[92]. Third, loss of grey-matter volume in the left-hemisphere Heschl's gyrus could also be reflected in an MMN reduction and explain speech perception deficits and speech sound-related hallucinations[104]. Reductions in temporal grey matter have also been reported in 22q11.2DS[38].

Lastly, individuals diagnosed with 22q11.2DS had, as expected, lower IQ scores and impaired working memory [18, 105-107]. Though intellectual disability is quite prevalent in 22q11.2DS, its causes are not well understood. Several genes have been identified as potential candidates: For instance, *Tbx1* has been implicated in brain development and may thus have a role in cognitive deficits[108]; PRODH mutations are associated with intellectual disability, and are found in about one-third of the individuals diagnosed with 22q11.2DS[109]. Regarding working memory, the copy number elevations of catechol-O-methyl-transferase (COMT) have been associated with such impairments, not only in 22q11.2DS[110], but also in schizophrenia[111] (but see[112] for different findings). COMT mutations have been argued to result in increased dopamine degradation in the frontal lobes, which could potentially provide a molecular basis for some of the symptomatology associated with both schizophrenia and 22q11.2DS[113, 114]. Additionally, increased variability was observed in the 22q11.2DS group, in line with the remarkable variability in expression described in the syndrome[6,115, 116]. Though here we focus on the distinction between individuals with and without psychotic symptoms, other subgroups with different cognitive and neural

profiles may co-exist within 22q11.2DS. The definition of such subgroups would not only add to the understanding of the phenotype, but also, potentially, impact intervention.

In summary, using a passive oddball paradigm, we show an enhanced N1 in 22q11.2DS. Interestingly, MMN amplitudes correlated with working memory scores: Those who showed relatively decreased amplitudes, presented poorer working memory performance. The simultaneous presence of an increased N1 and a decreased MMN (for a subset of the 22q11.2DS group), may suggest that two different mechanisms are at play, impacting the neural response patterns reported here: One intrinsic to 22q11.2DS which results in increased sensory responses and another associated with psychosis and leading to decreased MMN responses. That reduced MMNs were observed only for those individuals with poorer working memory performance (and, importantly, psychotic symptomatology) and not for the 22q11.2DS group as whole, is in accordance with previous evidence of unimpaired MMN in first-degree relatives of individuals diagnosed with schizophrenia and the consequent hypothesis that the MMN deficit observed in schizophrenia might be a consequence of the disorder and not endophenotypic[69].

Funding

This work was supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), under award number U54 HD090260.

Acknowledgements

We wish to thank Drs. Juliana Bates, Katherine Behar, and Pamela Counts who performed the clinical assessments, and Elise Taverna and Danielle Newbury for their help with data collection. We would also like to acknowledge the role of the Montefiore-Einstein Regional Center for 22q11.2 Deletion Syndrome in recruitment, and the Rose F. Kennedy Intellectual and Developmental Disability Research Center for all of its support. We extend our most sincere gratitude to the participants and their families for their interest, their involvement, and their time.

REFERENCES

1. Karayiorgou M, Simon TJ, Gogos JA. 22q11.2 microdeletions: Linking DNA structural variation to brain dysfunction and schizophrenia. *Nature Reviews Neuroscience* Jun 2010;11(6):402-416.
2. Meechan DW, Tucker ES, Maynard TM, LaMantia A. Diminished dosage of 22q11 genes disrupts neurogenesis and cortical development in a mouse model of 22q11 deletion/DiGeorge syndrome. *Proceedings of the National Academy of Sciences* 2009;106(38):16434-16445.
3. Botto LD, May K, Fernhoff PM, et al. A population-based study of the 22q11.2 deletion: Phenotype, incidence, and contribution to major birth defects in the population. *Pediatrics* 2003;112(1):101-107.
4. Grati FR, Molina Gomes D, Ferreira JC, et al. Prevalence of recurrent pathogenic microdeletions and microduplications in over 9500 pregnancies. *Prenatal Diagnosis* Aug 2015;35(8):801-809.
5. Philip N, Bassett AS. Cognitive, behavioural and psychiatric phenotype in 22q11.2 deletion syndrome. *Behavior Genetics* 2011;41(3):403-412.
6. McDonald-McGinn DM, Sullivan KE, Marino B, et al. 22q11.2 deletion syndrome. *Nature Reviews Disease Primers* Nov 19 2015;1:15071.
7. Shprintzen RJ, Goldberg R, Golding-Kushner KJ, Marion RW. Late-onset psychosis in the velo-cardio-facial syndrome. *American Journal of Medical Genetics* 1992;42:141-142.
8. Bish JP, Ferrante SM, McDonald-McGinn D, Zackai E, Simon TJ. Maladaptive conflict monitoring as evidence for executive dysfunction in children with chromosome 22q11.2 deletion syndrome. *Developmental Science* 2005;8(1):36-43.
9. Rockers K, Ousley O, Sutton T, Schoenberg E, Coleman K, Walker E, Cubells JF. Performance on the modified card sorting test and its relation to psychopathology in adolescents and young adults with 22q11.2 deletion syndrome. *Journal of Intellectual Disabilities Research* Jul 2009;53(7):665-676.
10. Sobin C, Kiley-Brabeck K, Daniels S, Blundell M, Anyane-Yeboa K, Karayiorgou M. Networks of attention in children with the 22q11 deletion syndrome. *Developmental Neuropsychology* 2004;26(2):611-626.
11. Van Aken K, Swillen A, Beirinckx M, Janssens L, Caeyenberghs K, Smits-Engelsman B. Prospective control abilities during visuo-manual tracking in children with 22q11.2 Deletion syndrome compared to age- and IQ-matched controls. *Research in Developmental DIsabilities* May-Jun 2010;31(3):634-641.
12. Bearden CE, Woodin MF, Wang PP, Moss E, McDonald-McGinn D, Zackai E, Emannuel B, Cannon TD. The neurocognitive phenotype of the 22q11.2 deletion syndrome: selective deficit in visual-spatial memory. *Journal of Clinical and Experimental Neuropsychology* Aug 2001;23(4):447-464.
13. Swillen A, Devriendt K, Legius E, Prinzie P, Vogels A, Ghesquière P, Fryns JP. The behavioural phenotyoe in velo-cardio-facial syndrome (VCFS): From infancy to adolescence. *Genetic Counceling* 1999;10(1):79-88.
14. Antshel KM, Fremont W, Kates WR. The neurocognitive phenotype in velo-cardio-facial syndrome: a developmental perspective. *Developmental Disabilities Research Reviews* 2008;14(1):43-51.
15. Cabaral M, Beaton EA, Stoddard J, Simon TJ. Impaired multiple object tracking in children with chromosome 22q11.2 deletion syndrome. *Journal of Neurodevelopmental Disorders* 2012;4(6):1-8.
16. Simon TJ, Bish JP, Bearden CE, et al. A multilevel analysis of cognitive dysfunction and psychopathology associated with chromossome 22q11.2 deletion syndrome in children. *Developmental Psychopathology* 2005;17(3):753-784.
17. Howley SA, Prasad SE, Pender NP, Murphy KC. Relationship between reaction time, fine motor control, and visual-spatial perception on vigilance and visual-motor tasks in 22q11.2 Deletion Syndrome. *Research in Developmental DIsabilities* Sep-Oct 2012;33(5):1495-1502.
18. Lajiness-O'Neill RR, Beaulieu I, Titus JB, Asamoah A, Bigler ED, Bawle EV, Pollack R. Memory and learning in children with 22q11.2 deletion syndrome: evidence for ventral and dorsal stream disruption? *Child Neuropsychology* Feb 2005;11(1):55-71.
19. Murphy KC. Schizophrenia and velo-cardio-facial syndrome. *The Lancet* 2002;359(9304):426-430.
20. Bassett AS, Chow EWC, Weksberg R. Chromosomal abnormalities and schizophrenia. *American Journal of Medical Genetics* 2000;97:45-51.
21. Baker K, Vorstman JA. Is there a core neuropsychiatric phenotype in 22q11.2 deletion syndrome? *Current Opinion in Neurology* Apr 2012;25(2):131-137.
22. Bassett AS, Chow EWC. 22q11 deletion syndrome: A genetic subtype of schizophrenia. *Biological Psychiatry* 1999;10:882-891.

23. Monks S, Niarchou M, Davies AR, Walters JT, Williams N, Owen MJ, van den Bree MB, Murphy KC. Further evidence for high rates of schizophrenia in 22q11.2 deletion syndrome. *Schizophrenia Research* Mar 2014;153(1-3):231-236.

24. Murphy KC, Jones LA, Owen MJ. High rates of schizophrenia in adults with velo-cardio-facial syndrome. *Archives of General Psychiatry* 1999;56:940-945.

25. Hemmings CP. Schizophrenia spectrum disorders in people with intellectual disabilities. *Current Opinion in Psychiatry* 2006;19:470-474.

26. Baker K, Skuse DH. Adolescents and young adults with 22q11 deletion syndrome: Psychopathology in an at-risk group. *The British Journal of Psychiatry* 2005;186:115-120.

27. Schneider M, Armando M, Pontillo M, Vicari S, Debbane M, Schultze-Lutter F, Eliez S. Ultra high risk status and transition to psychosis in 22q11.2 deletion syndrome. *World Psychiatry* 2016;15(3):259-265.

28. Hans SL, Auerbach JG, Asarnow JR, Styr B, Marcus J. Social adjustment of adolescents at risk for schizophrenia: the Jerusalem Infant Development Study. *Journal of the American Academy of Child and Adolescent Psychiatry* 2000;39:1406-1414.

29. Welham J, Isohanni M, Jones P, McGrath J. The antecedents of schizophrenia: A review of birth cohort studies. *Schizophrenia Bulletin* May 2009;35(3):603-623.

30. Yuen T, Chow EW, Silversides CK, Bassett AS. Premorbid adjustment and schizophrenia in individuals with 22q11.2 deletion syndrome. *Schizophrenia Research* Dec 2013;151(1-3):221-225.

31. Tang SX, M. MT, Calkins ME, et al. The Psychosis Spectrum in 22q11.2 Deletion Syndrome Is Comparable to That of Nondeleted Youths. *Biological Psychiatry* 2017;82:17-25.

32. Bassett AS, Chow E. Schizophrenia and 22q11.2 Deletion Syndrome. *Current Psychiatry Reports* 2008;10(2):148-157.

33. Chow EW, Watson M, Young DA, Bassett AS. Neurocognitive profile in 22q11 deletion syndrome and schizophrenia. *Schizophrenia Research* Oct 2006;87(1-3):270-278.

34. Kwon JS, Shenton ME, Hirayasu Y, et al. MRI study of cavum septi pellucidi in schizophrenia, affective disorder, and schizotypal personality disorder. *American Journal of Psychiatry* 1998;155(4):509-515.

35. Nopoulos P, Swayze V, Flaum M, Ehrhardt JC, Yuh WTC, Andreasen NC. Cavum septi pellucidi in normals and patients with schizophrenia as detected by magnetic resonance imaging. *Biological Psychiatry* 1997;41:1102-1108.

36. Bish JP, Pendyal A, Ding L, Ferrante H, Nguyen V, McDonald-McGinn D, Zackai E, Simon TJ. Specific cerebellar reductions in children with chromosome 22q11.2 deletion syndrome. *Neuroscience Letters* May 22 2006;399(3):245-248.

37. Chow EWC, Mikulis DJ, Zipursky RB, Scutt LE, Weksberg R, Bassett AS. Qualitative MRI findings in adults with 22q11 deletion syndrome and schizophrenia. *Biological Psychiatry* 1999;46:1436-1442.

38. Eliez S, Schmitt JE, White CD, Wellis VG, Riss AL. A Quantitative MRI study of posterior fossa development in velocardiofacial syndrome. *Biological Psychiatry* 2001;49:540-546.

39. Van Amelsvoort T, Daly E, Henry J, Robertson D, Ng V, Owen M, Murphy KC, Murphy DGM. Brain anatomy in adults with velocardiofacial syndrome with and without schizophrenia. *Archives of General Psychiatry* 2004;61:1085-1096.

40. Takahashi T, Suzuki M, Tanino R, et al. Volume reduction of the left planum temporale gray matter associated with long duration of untreated psychosis in schizophrenia: a preliminary report. *Psychiatry Research* Apr 15 2007;154(3):209-219.

41. Chang WC, Hui CLM, Tang JYM, Wong GHY, Lam MML, Chan SKW, Chen EYH. Persistent negative symptoms in first-episode schizophrenia: A prospective three-year follow-up study. *Schizophrenia Research* 2011;133(1-3):22-28.

42. Kato T, Higuchi Y, Sumiyoshi T, Seo T, Miyanishi T, Kawasaki Y, Suzuki M. Mismatch negativity and cognitive performance for the prediction of psychosis in subjects with at-risk mental state. *PLoS ONE* 2013;8(1):e54080.

43. Näätänen R, Picton T. The N1 wave of the human electric and magnetic response to sound: A review and an analysis of the component structure. *Psychophysiology* 1987;24(4):375-425.

44. Giard MH, Perrin F, Echallier JF, Thevenet M, Froment JC, Pernier J. Dissociation of temporal and frontal components in the human auditory N1 wave: A scalp current density and dipole model analysis. *Electroencephalography and Clinical Neurophysiology* 1994;92:238-253.

45. Strik WK, Dierks T, Boning J, Osterheider M, Caspari A, Korber J. Disorders of smooth pursuit eye movement and auditory N100 in schizphrenic patients. *Psychiatry Research* 1992;41:227-235.

46. Brockhaus-Dumke A, Schultze-Lutter F, Mueller R, Tendolkar I, Bechdolf A, Pukrop R, Klosterkoetter J, Ruhrmann S. Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. *Biological Psychiatry* Sep 1 2008;64(5):376-384.

47. Salisbury DF, Collins KC, McCarley RW. Reductions in the N1 and P2 auditory event-related potentials in first-hospitalized and chronic schizophrenia. *Schizophrenia Bulletin* Sep 2010;36(5):991-1000.
48. Foxe JJ, Yeap S, Snyder AC, Kelly SP, Thakore JH, Molholm S. The N1 auditory evoked potential component as an endophenotype for schizophrenia: High-density electrical mapping in clinically unaffected first-degree relatives, first-episode, and chronic schizophrenia patients. *European Archives of Psychiatry and Clinical Neuroscience* Aug 2011;261(5):331-339.
49. Leavitt VM, Molholm S, Ritter W, Shpaner M, Foxe JJ. Auditory processing in schizophrenia during the middle latency period (10–50 ms): high-density electrical mapping and source analysis reveal subcortical antecedents to early cortical deficits. *Journal of Psychiatry & Neuroscience* 2007;32(5):339-353.
50. Rosburg T, Boutros NN, Ford JM. Reduced auditory evoked potential component N100 in schizophrenia: A critical review. *Psychiatry Research* Dec 15 2008;161(3):259-274.
51. Näätänen R, Alho K. Mismatch negativity-A unique measure of sensory processing in audition. *International Journal of Neuroscience* 1995;80(1-4):317-337.
52. Ritter W, Deacon D, Gomes H, Javitt DC, Vaughan JH. The mismatch negativity of event-related potentials as a probe of transient auditory memory: A review. *Ear and Hearing* 1995;16(1):52-67.
53. Ritter W, Sussman E, Molholm S, Foxe JJ. Memory reactivation or reinstatement and the mismatch negativity. *Psychophysiology* 2002;39:158-165.
54. Erickson MA, Ruffle A, Gold JM. A meta-analysis of mismatch negativity in schizophrenia: From clinical risk to disease specificity and progression. *Biological Psychiatry* Jun 15 2016;79(12):980-987.
55. Damaso KA, Michie PT, Todd J. Paying attention to MMN in schizophrenia. *Brain Research* Nov 11 2015;1626:267-279.
56. Bodatsch M, Ruhrmann S, Wagner M, et al. Prediction of psychosis by mismatch negativity. *Biological Psychiatry* May 15 2011;69(10):959-966.
57. Jahshan C, Cadenhead KS, Rissling AJ, Kirihara K, Braff DL, Light GA. Automatic sensory information processing abnormalities across the illness course of schizophrenia. *Psychological Medicine* 2011;42(1):85-97.
58. Atkinson RJ, Michie PT, Schall U. Duration mismatch negativity and P3a in first-episode psychosis and individuals at ultra-high risk of psychosis. *Biological Psychiatry* Jan 15 2012;71(2):98-104.
59. Shaikh M, Valmaggia L, Broome MR, et al. Reduced mismatch negativity predates the onset of psychosis. *Schizophrenia Research* 2012;134(1):42-48.
60. Shin KS, Kim JS, Kang DH, Koh Y, Choi JS, O'Donnell BF, Chung CK, Kwon JS. Pre-attentive auditory processing in ultra-high-risk for schizophrenia with magnetoencephalography. *Biological Psychiatry* Jun 15 2009;65(12):1071-1078.
61. Umbricht DS, Bates JA, Lieberman JA, Kane JM, Javitt DC. Electrophysiological indices of automatic and controlled auditory information processing in first-episode, recent-onset and chronic schizophrenia. *Biological Psychiatry* Apr 15 2006;59(8):762-772.
62. Javitt DC, Shelley AM, Silipo G, Lieberman JA. Schizophrenia spectrum disorders in people with intellectual disabilities. *Archives of General Psychiatry* 2000;57:1131-1137.
63. Todd J, Michie PT, Schall U, Karayanidis F, Yabe H, Naatanen R. Deviant matters: Duration, frequency, and intensity deviants reveal different patterns of mismatch negativity reduction in early and late schizophrenia. *Biological Psychiatry* Jan 1 2008;63(1):58-64.
64. Michie PT, Budd TW, Todd J, Rock D, Wichmann H, Box J, Jablensky AV. Duration and frequency mismatch negativity in schizophrenia. *Clinical Neurophysiology* 2000;111:1054-1065.
65. Oades RD, Wild-Wall N, Juran SA, Sachsse J, Oknina LB, Ropcke B. Auditory change detection in schizophrenia: Sources of activity, related neuropsychological function and symptoms in patients with a first episode in adolescence, and patients 14 years after an adolescent illness-onset. *BMC Psychiatry* Feb 8 2006;6(7):1-14.
66. Brockhaus-Dumke A, Tendolkar I, Pukrop R, Schultze-Lutter F, Klosterkotter J, Ruhrmann S. Impaired mismatch negativity generation in prodromal subjects and patients with schizophrenia. *Schizophrenia Research* Mar 1 2005;73(2-3):297-310.
67. Oknina LB, Wild-Wall N, Oades RD, et al. Frontal and temporal sources of mismatch negativity in healthy controls, patients at onset of schizophrenia in adolescence and others at 15 years after onset. *Schizophrenia Research* Jul 1 2005;76(1):25-41.
68. Light GA, Braff DL. Mismatch negativity deficits are associated with poor functioning in schizophrenia patients. *Archives of General Psychiatry* 2005;62:127-136.
69. Magno E, Yeap S, Thakore JH, Garavan H, De Sanctis P, Foxe JJ. Are auditory-evoked frequency and duration mismatch negativity deficits endophenotypic for schizophrenia? High-density electrical mapping

in clinically unaffected first-degree relatives and first-episode and chronic schizophrenia. *Biological Psychiatry* Sep 1 2008;64(5):385-391.

70. Kiang M, Braff DL, Sprock J, Light GA. The relationship between preattentive sensory processing deficits and age in schizophrenia patients. *Clinical Neurophysiology* Nov 2009;120(11):1949-1957.

71. Leitman DI, Sehatpour P, Higgins BA, Foxe JJ, Silipo G, Javitt DC. Sensory deficits and distributed hierarchical dysfunction in schizophrenia. *Am J Psychiatry* Jul 2010;167(7):818-827.

72. Larsen KM, Morup M, Birknow MR, et al. Altered auditory processing and effective connectivity in 22q11.2 deletion syndrome. *Schizophrenia Research* Jan 30 2018;197:328-336.

73. Rihs TA, Tomescu MI, Britz J, et al. Altered auditory processing in frontal and left temporal cortex in 22q11.2 deletion syndrome: A group at high genetic risk for schizophrenia. *Psychiatry Research* May 30 2013;212(2):141-149.

74. Brima T, Molholm S, Molloy CJ, Sysoeva OV, Nicholas E, Djukic A, Freedman EG, Foxe JJ. Auditory sensory memory span for duration is severely curtailed in females with Rett syndrome. *Transl Psychiatry* Apr 9 2019;9(1):130.

75. Bartha-Doering L, Deuster D, Giordano V, am Zehnhoff-Dinnesen A, Dobel C. A systematic review of the mismatch negativity as an index for auditory sensory memory: From basic research to clinical and developmental perspectives. *Psychophysiology* 2015;52:1115-1130.

76. Wechsler D. *WAIS-IV Administration and Scoring Manual*. San Antonio, TX: Psychological Corporation; 2008.

77. Wechsler D. *WISC-V: Technical and Interpretive Manual*. Bloomington, MN: Pearson; 2014.

78. First MB, Williams JBW, Karg RS, Spitzer RL. *Structured Clinical Interview for DSM-5—Research Version (SCID-5 for DSM-5, research version; SCID-5-RV)*. Arlington, VA: American Psychiatric Association; 2015.

79. Hien D, Matzner FJ, First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV-Child Edition (Version 1.0)*. New York: Columbia University; 1994.

80. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods* Mar 15 2004;134(1):9-21.

81. Holm S. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 1979;6(2):65-70.

82. *R: A language and environment for statistical computing* [computer program]. Version 3.3.2. Vienna, Austria: R Foundation for Statistical Computing; 2014.

83. Bates D, Mächler M, Bolker BM, Walker SC. Ime4: Linear mixed-effects models using Eigen and S4 (R package Version 1.1-7) [Computer software]. *Retrieved from http://CRANR-projectorg/package=lme4* 2014.

84. Didriksen M, Fejgin K, Nilsson SRO, et al. Persistent gating deficit and increased sensitivity to NMDA receptor antagonism after puberty in a new mouse model of the human 22q11.2 microdeletion syndrome: a study in male mice. *Journal of Psychiatry & Neuroscience* 2017;42(1):48-58.

85. Cantonas LM, Tomescu MI, Biria M, Jan RK, Schneider M, Eliez S, Rihs TA, Michel CM. Abnormal development of early auditory processing in 22q11.2 Deletion Syndrome. *Translational Psychiatry* 2019;9(138):1-12.

86. Oranje B, van Berckel BNM, Kemner C, van Ree JM, Kahn RS, Verbaten MN. The effects of a sub-anaesthetic dose of ketamine on human selective attention. *Neuropsychopharmacology* 2000;22(3):293-302.

87. Umbricht D, Schmid L, Koller R, Vollenweider FX, Hell D, Javitt DC. Ketamine-induced deficits in auditory and visual context-dependent processing in healthy volunteers. *Archives of General Psychiatry* 2000;57:1139-1147.

88. Turetsky BI, Calkins ME, Light GA, Olincy A, Radant AD, Swerdlow NR. Neurophysiological endophenotypes of schizophrenia: The viability of selected candidate measures. *Schizophrenia Bulletin* Jan 2007;33(1):69-94.

89. Leavitt VM, Molholm S, Ritter W, Shpaner M, Foxe JJ. Auditory processing in schizophrenia during the middle latency period (10-50 ms): high-density electrical mapping and source analysis reveal subcortical antecedents to early cortical deficits. *J Psychiatry Neurosci* Sep 2007;32(5):339-353.

90. Larsen KM, Dzafic I, Siebner HR, Garrido MI. Alteration of functional brain architecture in 22q11.2 deletion syndrome-Insights into susceptibility for psychosis. *NeuroImage* Sep 5 2018:1-18.

91. Baker K, Baldeweg T, Sivagnanasundaram S, Scambler P, Skuse D. COMT Val108/158 Met modifies mismatch negativity and cognitive function in 22q11 deletion syndrome. *Biological Psychiatry* Jul 1 2005;58(1):23-31.

92. Zarchi O, Carmel M, Avni C, et al. Schizophrenia-like neurophysiological abnormalities in 22q11.2 deletion syndrome and their association to COMT and PRODH genotypes. *Journal of Psychiatric Research* Nov 2013;47(11):1623-1629.

93. Montojo CA, Ibrahim A, Karlsgodt KH, Chow C, Hilton AE, Jonas RK, Vesagas TK, Bearden CE. Disrupted working memory circuitry and psychotic symptoms in 22q11.2 deletion syndrome. *NeuroImage: Clinical* 2014;4:392-402.

94. Bowie CR, Harvey PD. Cognition in schizophrenia: Impairments, determinants, and functional Importance. *Psychiatric Clinics of North America* 2005;28:613-633.

95. Forbes NF, Carrick LA, McIntosh AM, Lawrie SM. Working memory in schizophrenia: A meta-analysis. *Psychological Medicine* 2009;39:889-905.

96. Lee J, Park S. Working memory impairments in schizophrenia: A meta-analysis. *Journal of Abnormal Psychology* 2005;114(4):599-611.

97. Lencz T, Smith CW, McLaughlin D, Auther A, Nakayama E, Hovey L, Cornblatt BA. Generalized and specific neurocognitive deficits in prodromal schizophrenia. *Biological Psychiatry* 2006;59:863-871.

98. Smith CW, Park S, Cornblatt B. Spatial working memory deficits in adolescents at clinical high risk for schizophrenia. *Schizophrenia Research* 2006;81:211-215.

99. Javitt DC, Grochowski S, Shelley A-M, Ritter W. Impaired mismatch negativity (MMN) generation in schizophrenia as a function of stimulus deviance, probability, and interstimulus/interdeviant interval. *Electroencephalography and Clinical Neurophysiology* 1998;108:143-153.

100. Kantrowitz J, Javitt DC. Glutamatergic transmission in schizophrenia: from basic research to clinical practice. *Current Opinion in Psychiatry* Mar 2012;25(2):96-102.

101. Henzi V, Reichling DB, Helm SW, MacDermott AB. L-proline activates glutamate and glycine receptors in cultured rat dorsal horn neurons. *Molecular Pharmacology* 1992;41:793-801.

102. Paterlini M, Zakharenko SS, Lai W, et al. Transcriptional and behavior interaction between 22q1.2 orthologs modulates schizophrenia-related phenotypes in mice. *Nature Neuroscience* 2005;8(11):1586-1594.

103. Baldeweg T, Klugman A, Gruzelier JH, Hirsch SR. Impairment in frontal but not temporal components of mismatch negativity in schizophrenia. *International Journal of Psychophysiology* 2002;43:111-122.

104. Näätänen R, Kahkonen S. Central auditory dysfunction in schizophrenia as revealed by the mismatch negativity (MMN) and its magnetic equivalent MMNm: A review. *International Journal of Neuropsychopharmacology* Feb 2009;12(1):125-135.

105. Swillen A, Devriendt K, Legius E, Eyskens B, Dumoulin M, Gewillig M, Fryns JP. Intelligence and psychosocial adjustment in velocardiofacial syndrome: A study of 37 children and adolescents with VCFS. *Journal of Medical Genetics* 1997;34:453-458.

106. De Smedt B, Devriendt K, Fryns JP, Vogels A, Gewillig M, Swillen A. Intellectual abilities in a large sample of children with Velo-Cardio-Facial Syndrome: An update. *J Intellect Disabil Res* Sep 2007;51(9):666-670.

107. Maeder J, Schneider M, Bostelmann M, Debbane M, Glaser B, Menghetti S, Schaer M, Eliez S. Developmental trajectories of executive functions in 22q11.2 deletion syndrome. *Journal of Neurodevelopmental Disorders* 2016;8(10):1-12.

108. Paylor R, Glaser B, Mupo A, et al. Tbx1 haploinsufficiency is linked to behavioral disorders in mice and humans: Implications for 22q11 deletion syndrome. *Proceedings of the National Academy of Sciences* 2006;103(20):7729-7734.

109. Goodman BK, Rutberg J, Lin WW, Pulver AE, Thomas GH, Geraghty MT. Hyperprolinaemia in patients with deletion (22)(q11.2) syndrome. *Journal of Inherited Metabolic Disease* 2000;23:847-848.

110. Boku S, Izumi T, Abe S, et al. Copy number elevation of 22q11.2 genes arrests the developmental maturation of working memory capacity and adult hippocampal neurogenesis. *Molecular Psychiatry* Apr 2018;23(4):985-992.

111. Diaz-Asper CM, Goldberg TE, Kolachana BS, Straub RE, Egan MF, Weinberger DR. Genetic variation in catechol-O-methyltransferase: effects on working memory in schizophrenic patients, their siblings, and healthy controls. *Biological Psychiatry* Jan 1 2008;63(1):72-79.

112. Ho BC, Wassink TH, O'Leary DS, Sheffield VC, Andreasen NC. Catechol-O-methyl transferase Val158Met gene polymorphism in schizophrenia: Working memory, frontal lobe MRI morphology and frontal cerebral blood flow. *Molecular Psychiatry* Mar 2005;10(3):287-298.

113. Bearden CE, Jawad AF, Lynch DR, et al. Effects of a functional COMT polymorphism on prefrontal cognitive function in patients with 22q11.2 deletion syndrome. *American Journal of Psychiatry* 2004;161:1700-1702.

114. Gothelf D, Eliez S, Thompson T, et al. COMT genotype predicts longitudinal cognitive decline and psychosis in 22q11.2 deletion syndrome. *Nature Neuroscience* Nov 2005;8(11):1500-1502.
115. McDonald-McGinn DM, Sullivan KE. Chromosome 22q11.2 deletion syndrome (DiGeorge syndrome/velocardiofacial syndrome). *Medicine* Jan 2011;90(1):1-18.
116. Yamagishi H, Ishii C, Maeda J, et al. Phenotypic discordance in monozygotic twins with 22q11.2 Deletion. *American Journal of Medical Genetics* 1998;78:319-321.