Accelerated viral dynamics in bat cell lines, with implications for zoonotic emergence

Cara E. Brook[1,2†], Mike Boots[1], Kartik Chandran[3], Andrew P. Dobson[2], Christian Drosten[4],

Andrea L. Graham[2], Bryan T. Grenfell[2,5], Marcel A. Müller[4,6], Melinda Ng[3], Lin-Fa Wang[7],

Anieke van Leeuwen[8,2]

[1] Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, USA.

[2] Department of Ecology & Evolutionary Biology, Princeton University, Princeton, NJ, USA.

[3] Department of Microbiology and Immunology, Albert Einstein College of Medicine, Bronx,

NY, USA.

[4] Institute of Virology, Charité-Universitätsmedizin Berlin, corporate member of Freie

Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin,

Germany.

[5] Fogarty International Center, National Institutes of Health, Bethesda, Maryland, USA.

[6] Martsinovsky Institute of Medical Parasitology, Tropical and Vector Borne Diseases, Sechenov

University, Moscow, Russia.

[7] Emerging Infectious Diseases Program, Duke-National University of Singapore Medical

School, Singapore.

[8] Royal Netherlands Institute for Sea Research, Department of Coastal Systems, and Utrecht

University, the Netherlands

Abstract

Bats host virulent zoonotic viruses without experiencing disease. A mechanistic understanding of the impact of bats' virus hosting capacities, including uniquely constitutive immune pathways, on cellular-scale viral dynamics is needed to elucidate zoonotic emergence. We carried out virus infectivity assays on bat cell lines expressing induced and constitutive immune phenotypes, then developed a theoretical model of our *in vitro* system, which we fit to empirical data. Best fit models recapitulated expected immune phenotypes for representative cell lines, supporting robust antiviral defenses in bat cells that correlated with higher estimates for within-host viral propagation rates. In general, heightened immune responses limit pathogen-induced cellular morbidity, which can facilitate the establishment of rapidly-propagating persistent infections within-host. Rapidly-replicating viruses that have evolved with bat immune systems will likely cause enhanced virulence following emergence into secondary hosts with immune systems that diverge from those unique to bats.

Introduction

Bats have received much attention in recent years for their role as reservoir hosts for emerging viral zoonoses, including rabies and related lyssaviruses, Hendra and Nipah henipaviruses, Ebola and Marburg filoviruses, and SARS coronavirus (Calisher et al. 2006; Wang and Anderson 2019). In most non-Chiropteran mammals, henipaviruses, filoviruses, and coronaviruses induce substantial morbidity and mortality, display short durations of infection, and elicit robust, long-term immunity in hosts surviving infection (Nicholls et al. 2003; Hooper et al. 2001; Mahanty and Bray 2004). Bats, by contrast, demonstrate no obvious disease symptoms upon infection with pathogens that are highly virulent in non-volant mammals (Schountz et al. 2017) but may, instead, support viruses as long-term persistent infections, rather than transient, immunizing pathologies (Plowright et al. 2016).

Recent research advances are beginning to shed light on the molecular mechanisms by which bats avoid pathology from these otherwise virulent pathogens (Brook and Dobson 2015). Bats leverage a suite of species-specific mechanisms to limit viral load, which include host receptor sequence incompatibilities for some bat-virus combinations (Ng et al. 2015; Igarashi et al. 2020) and constitutive expression of the antiviral cytokine, IFN-α, for others (Zhou et al. 2016). Typically, the presence of viral RNA or DNA in the cytoplasm of mammalian cells will induce secretion of type I interferon proteins (IFN-α and IFN-β), which promote expression and translation of interferon-stimulated genes (ISGs) in neighboring cells and render them effectively antiviral (Stetson and Medzhitov 2006). In some bat cells, the transcriptomic blueprints for this IFN response are expressed constitutively, even in the absence of stimulation by viral RNA or DNA (Zhou et al. 2016). In non-flying mammals, constitutive IFN expression would likely elicit widespread inflammation and concomitant immunopathology upon viral infection, but bats

support unique adaptations to combat inflammation (Zhang et al. 2013; Ahn et al. 2019; Xie et

al. 2018; Pavlovich et al. 2018) that may have evolved to mitigate metabolic damage induced

during flight (Kacprzyk et al. 2017). The extent to which constitutive IFN-α expression signifies

constitutive antiviral defense in the form of functional IFN-α protein remains unresolved. In bat

cells constitutively expressing IFN-α, some protein-stimulated, downstream ISGs appear to be

also constitutively expressed, but additional ISG induction is nonetheless possible following viral

challenge and stimulation of IFN-β (Zhou et al. 2016; Xie et al. 2018). Despite recent advances

in molecular understanding of bat viral tolerance, the consequences of this unique bat immunity

on within-host virus dynamics—and its implications for understanding zoonotic emergence—

have yet to be elucidated.

The field of 'virus dynamics' was first developed to describe the mechanistic

underpinnings of long-term patterns of steady-state viral load exhibited by patients in chronic

phase infections with HIV, who appeared to produce and clear virus at equivalent rates (Nowak

and May 2000; Ho et al. 1995). Models of simple target cell depletion, in which viral load is

dictated by a bottom-up resource supply of infection-susceptible host cells, were first developed

for HIV (Perelson 2002) but have since been applied to other chronic infections, including

hepatitis-C virus (Neumann et al. 1998), hepatitis-B virus (Nowak et al. 1996) and

cytomegalovirus (Emery et al. 1999). Recent work has adopted similar techniques to model the

within-host dynamics of acute infections, such as influenza A and measles, inspiring debate over

the extent to which explicit modeling of top-down immune control can improve inference

beyond the basic resource limitation assumptions of the target cell model (Baccam et al. 2006;

Pawelek et al. 2012; Saenz et al. 2010; Morris et al. 2018).

To investigate the impact of unique bat immune processes on *in vitro* viral kinetics, we

first undertook a series of virus infection experiments on bat cell lines expressing divergent

interferon phenotypes, then developed a theoretical model elucidating the dynamics of within-

host viral spread. We evaluated our theoretical model analytically independent of the data, then

fit the model to data recovered from *in vitro* experimental trials in order to estimate rates of

within-host virus transmission and cellular progression to antiviral status under diverse

assumptions of absent, induced, and constitutive immunity. Finally, we confirmed our findings in

spatially-explicit stochastic simulations of fitted time series from our mean field model. We

hypothesized that top-down immune processes would overrule classical resource-limitation in

bat cell lines described as constitutively antiviral in the literature, offering a testable prediction

for models fit to empirical data. We further predicted that the most robust antiviral responses

would be associated with the most rapid within-host virus propagation rates but also protect cells

against virus-induced mortality to support the longest enduring infections in tissue culture.

Results

Virus infection experiments in antiviral bat cell cultures yield reduced cell mortality and

elongated epidemics.

We first explored the influence of innate immune phenotype on within-host viral

propagation in a series of infection experiments in cell culture. We conducted plaque assays on

six-well plate monolayers of three immortalized mammalian kidney cell lines: [1] Vero (African

green monkey) cells, which are IFN-defective and thus limited in antiviral capacity (Desmyter,

Melnick, and Rawls 1968); [2] RoNi/7.1 (*Rousettus aegyptiacus*) cells which demonstrate

idiosyncratic induced interferon responses upon viral challenge (Kuzmin et al. 2017; Arnold et

al. 2018; Biesold et al. 2011; Pavlovich et al. 2018); and [3] PaKiT01 (*Pteropus alecto*) cells

which constitutively express IFN-α (Zhou et al. 2016; Crameri et al. 2009). To intensify cell

line-specific differences in constitutive immunity, we carried out infectivity assays with GFP-

tagged, replication-competent vesicular stomatitis Indiana viruses: rVSV-G, rVSV-EBOV, and

rVSV-MARV, which have been previously described (Miller et al. 2012; Wong et al. 2010).

Two of these viruses, rVSV-EBOV and rVSV-MARV, are recombinants for which cell entry is

mediated by the glycoprotein of the bat-evolved filoviruses, Ebola (EBOV) and Marburg

(MARV), thus allowing us to modulate the extent of structural, as well as immunological,

antiviral defense at play in each infection. Previous work in this lab has demonstrated

incompatibilities in the NPC1 filovirus receptor which render PaKiT01 cells refractory to

infection with rVSV-MARV (Ng and Chandran 2018), making them structurally antiviral, over

and above their constitutive expression of IFN-α. All three cell lines were challenged with all

three viruses at two multiplicities of infection (MOI): 0.001 and 0.0001. Between 18-39 trials

were run at each cell-virus-MOI combination, excepting rVSV-MARV infections on PaKiT01

cells at MOI=0.001, for which only 8 trials were run (see Materials and Methods; Figure 1-figure

supplement 1-3, Supplementary File 1).

Because plaque assays restrict viral transmission neighbor-to-neighbor in two-

dimensional cellular space (Howat et al. 2006), we were able to track the spread of GFP-

expressing virus-infected cells across tissue monolayers via inverted fluorescence microscopy.

For each infection trial, we monitored and re-imaged plates for up to 200 hours of observations

or until total monolayer destruction, processed resulting images, and generated a time series of

the proportion of infectious-cell occupied plate space across the duration of each trial (see

Materials and Methods). We used generalized additive models to infer the time course of all cell

culture replicates and construct the multi-trial dataset to which we eventually fit our mechanistic

transmission model for each cell line-virus-specific combination (Figure 1; Figure 1-figure

supplement 1-5).

All three recombinant vesicular stomatitis viruses (rVSV-G, rVSV-EBOV, and rVSV-

MARV) infected Vero, RoNi/7.1, and PaKiT01 tissue cultures at both focal MOIs. Post-

invasion, virus spread rapidly across most cell monolayers, resulting in virus-induced epidemic

extinction. Epidemics were less severe in bat cell cultures, especially when infected with the

recombinant filoviruses, rVSV-EBOV and rVSV-MARV. Monolayer destruction was avoided in

the case of rVSV-EBOV and rVSV-MARV infections on PaKiT01 cells: in the former, persistent

viral infection was maintained throughout the 200-hour duration of each experiment, while, in

the latter, infection was eliminated early in the time series, preserving a large proportion of live,

uninfectious cells across the duration of the experiment. We assumed this pattern to be the result

of immune-mediated epidemic extinction (Figure 1). Patterns from MOI=0.001 were largely

recapitulated at MOI = 0.0001, though at somewhat reduced total proportions (Figure 1-figure

supplement 5).

A theoretical model fit to *in vitro* data recapitulates expected immune phenotypes for bat

cells.

We next developed a within-host model to fit to these data in order to elucidate the

effects of induced and constitutive immunity on the dynamics of viral spread in host tissue

(Figure 1). The compartmental within-host system mimicked our two-dimensional cell culture

monolayer, with cells occupying five distinct infection states: susceptible (S), antiviral (A),

exposed (E), infectious (I), and dead (D). We modeled exposed cells as infected but not yet

infectious, capturing the 'eclipse phase' of viral integration into a host cell which precedes viral

replication. Antiviral cells were immune to viral infection, in accordance with the "antiviral

state" induced from interferon stimulation of ISGs in tissues adjacent to infection (Stetson and

Medzhitov 2006). Because we aimed to translate available data into modeled processes, we did

not explicitly model interferon dynamics but instead scaled the rate of cell progression from

susceptible to antiviral (ρ) by the proportion of exposed cells (globally) in the system. In systems

permitting constitutive immunity, a second rate of cellular acquisition of antiviral status (ε)

additionally scaled with the global proportion of susceptible cells in the model. Compared with

virus, IFN particles are small and highly diffusive, justifying this global signaling assumption at

the limited spatial extent of a six well plate and maintaining consistency with previous modeling

approximations of IFN signaling in plaque assay (Howat et al. 2006).

To best represent our empirical monolayer system, we expressed our state variables as

proportions (P_S, P_A, P_E, P_I, and P_D), under assumptions of frequency-dependent transmission in a

well-mixed population (Keeling and Rohani 2008), though note that the inclusion of P_D

(representing the proportion of dead space in the modeled tissue) had the functional effect of

varying transmission with infectious cell density. This resulted in the following system of

ordinary differential equations:

$\frac{dP_S}{dt} = bP_D(P_S + P_A) - \beta P_S P_I - \mu P_S - \rho P_E P_S - \varepsilon P_S + cP_A$ (1)

$\frac{dP_A}{dt} = \rho P_E P_S + \varepsilon P_S - cP_A - \mu P_A$ (2)

$\frac{dP_E}{dt} = \beta P_S P_I - \sigma P_E - \mu P_E$ (3)

$\frac{dP_I}{dt} = \sigma P_E - \alpha P_I - \mu P_I$ (4)

$\quad \frac{dP_D}{dt} = \mu(P_S + P_E + P_I + P_A) + \alpha P_I - bP_D(P_S + P_A)$ (5)

We defined "induced immunity" as complete, modeling all cells as susceptible to viral

invasion at disease free equilibrium, with defenses induced subsequent to viral exposure through

the term ρ. By contrast, we allowed the extent of constitutive immunity to vary across the

parameter range of $\varepsilon > 0$, defining a "constitutive" system as one containing *any* antiviral cells at

disease free equilibrium. In fitting this model to tissue culture data, we independently estimated

both ρ and ε, as well as the cell-to-cell transmission rate, β, for each cell-virus combination.

Since the extent to which constitutively-expressed IFN-α is constitutively translated into

functional protein is not yet known for bat hosts (Zhou et al. 2016), this approach permitted our

tissue culture data to drive modeling inference: even in PaKiT01 cell lines known to

constitutively express IFN-α, the true constitutive extent of the system (i.e. the quantity of

antiviral cells present at disease free equilibrium) was allowed to vary through estimation of ε.

For the purposes of model-fitting, we fixed the value of c, the return rate of antiviral cells to

susceptible status, at 0. The small spatial scale and short time course (max 200 hours) of our

experiments likely prohibited any return of antiviral cells to susceptible status in our empirical

system; nonetheless, we retained the term c in analytical evaluations of our model because

regression from antiviral to susceptible status is possible over long time periods *in vitro* and at

the scale of a complete organism (Samuel and Knutson 1982; Rasmussen and Farley 1975;

Radke et al. 1974).

Before fitting to empirical time series, we undertook bifurcation analysis of our

theoretical model and generated testable hypotheses on the basis of model outcomes. From our

within-host model system (equations 1-5), we derived the following expression for R_0, the

pathogen basic reproduction number (Supplementary File 2):

$R_0 = \frac{\beta\sigma(b-\mu)(c+\mu)}{b(\sigma+\mu)(\alpha+\mu)(c+\mu+\varepsilon)}$ (6)

Pathogens can invade a host tissue culture when $R_0 > 1$. Rapid rates of constitutive antiviral

acquisition (ε) will drive $R_0 < 1$: tissue cultures with highly constitutive antiviral immunity will

be therefore resistant to virus invasion from the outset. Since, by definition, induced immunity is

stimulated following initial virus invasion, the rate of induced antiviral acquisition (ρ) is not

incorporated into the equation for R_0; while induced immune processes can control virus after

initial invasion, they cannot prevent it from occurring to begin with. In cases of fully induced or

absent immunity ($\varepsilon = 0$), the R_0 equation thus reduces to a form typical of the classic SEIR

model:

$R_0 = \frac{\beta\sigma(b-\mu)}{b(\alpha+\mu)(\sigma+\mu)}$ (7)

At equilibrium, the theoretical, mean field model demonstrates one of three infection

states: endemic equilibrium, stable limit cycles, or no infection (Figure 2). Respectively, these

states approximate the persistent infection, virus-induced epidemic extinction, and immune-

mediated epidemic extinction phenotypes previously witnessed in tissue culture experiments

(Figure 1). Theoretically, endemic equilibrium is maintained when new infections are generated

at the same rate at which infections are lost, while limit cycles represent parameter space under

which infectious and susceptible populations are locked in predictable oscillations. Endemic

equilibria resulting from cellular regeneration (i.e. births) have been described *in vivo* for HIV

(Coffin 1995) and *in vitro* for herpesvirus plaque assays (Howat et al. 2006), but, because they so

closely approach zero, true limit cycles likely only occur theoretically, instead yielding stochastic

extinctions in empirical time series.

Bifurcation analysis of our mean field model revealed that regions of no infection

(pathogen extinction) were bounded at lower threshold (Branch point) values for β, below which

the pathogen was unable to invade. We found no upper threshold to invasion for β under any

circumstances (i.e. β high enough to drive pathogen-induced extinction), but high β values

resulted in Hopf bifurcations, which delineate regions of parameter space characterized by limit

cycles. Since limit cycles so closely approach zero, high βs recovered in this range would likely

produce virus-induced epidemic extinctions under experimental conditions. Under more robust

representations of immunity, with higher values for either or both induced (ρ) and constitutive

(ε) rates of antiviral acquisition, Hopf bifurcations occurred at increasingly higher values for β,

meaning that persistent infections could establish at higher viral replication rates (Figure 2).

Consistent with our derivation for R_0, we found that the Branch point threshold for viral invasion

was independent of changes to the induced immune parameter (ρ) but saturated at high values of

ε that characterize highly constitutive immunity (Figure 3).

We next fit our theoretical model by least squares to each cell line-virus combination,

under absent, induced, and constitutive assumptions of immunity. In general, best fit models

recapitulated expected outcomes based on the immune phenotype of the cell line in question, as

described in the general literature (Table 1; Supplementary File 4). The absent immune model

offered the most accurate approximation of IFN-deficient Vero cell time series, the induced

immune model best recovered the RoNi/7.1 cell trials, and, in most cases, the constitutive

immune model most closely recaptured infection dynamics across constitutively IFN-α-

expressing PaKiT01 cell lines (Figure 1; Figure 1-figure supplement 4-5, Supplementary File 4).

Ironically, the induced immune model offered a slightly better fit than the constitutive to rVSV-

MARV infections on the PaKiT01 cell line (the one cell line-virus combination for which we

know a constitutively antiviral cell-receptor incompatibility to be at play). Because constitutive

immune assumptions can prohibit pathogen invasion ($R_0 < 1$), model fits to this time series

under constitutive assumptions were handicapped by overestimations of ε, which prohibited

pathogen invasion. Only by incorporating an exceedingly rapid rate of induced antiviral

acquisition could the model guarantee that initial infection would be permitted and then rapidly

controlled.

Robust immunity is linked to rapid within-host virus transmission rates in fitted models.

In fitting our theoretical model to *in vitro* data, we estimated the within-host virus

transmission rate (β) and the rate(s) of cellular acquisition to antiviral status (ρ or $\rho + \varepsilon$) (Table

1; Supplementary File 4). Under absent immune assumptions, ρ and ε were fixed at 0 while β

was estimated; under induced immune assumptions, ε was fixed at 0 while ρ and β were

estimated; and under constitutive immune assumptions, all three parameters (ρ, ε, and β) were

simultaneously estimated for each cell-virus combination. Best fit parameter estimates for

MOI=0.001 data are visualized in conjunction with $\beta - \rho$ and $\beta - \varepsilon$ bifurcations in Figure 4; all

general patterns were recapitulated at lower values for β on MOI=0.0001 trials (Figure 4-figure

supplement 1).

As anticipated, the immune absent model (a simple target cell model) offered the best fit

to IFN-deficient Vero cell infections (Figure 4; Table 1; Supplementary File 4). Among Vero

cell trials, infections with rVSV-G produced the highest β estimates, followed by infections with

rVSV-EBOV and rVSV-MARV. Best fit parameter estimates on Vero cell lines localized in the

region of parameter space corresponding to theoretical limit cycles, consistent with observed

virus-induced epidemic extinctions in stochastic tissue cultures.

In contrast to Vero cells, the induced immunity model offered the best fit to all RoNi/7.1

data, consistent with reported patterns in the literature and our own validation by qPCR (Table 1;

Figure 1-figure supplement 6; Biesold et al. 2011; Kuzmin et al. 2017; Arnold et al. 2018;

Pavlovich et al. 2018). As in Vero cell trials, we estimated highest β values for rVSV-G

infections on RoNi/7.1 cell lines but here recovered higher β estimates for rVSV-MARV than for

rVSV-EBOV. This reversal was balanced by a higher estimated rate of acquisition to antiviral

status (ρ) for rVSV-EBOV versus rVSV-MARV. In general, we observed that more rapid rates

of antiviral acquisition (either induced, ρ, constitutive, ε, or both) correlated with higher

transmission rates (β). When offset by ρ, β values estimated for RoNi/7.1 infections maintained

the same amplitude as those estimated for immune-absent Vero cell lines but caused gentler

epidemics and reduced cellular mortality (Figure 1). RoNi/7.1 parameter estimates localized in

the region corresponding to endemic equilibrium for the deterministic, theoretical model (Figure

4), yielding less acute epidemics which nonetheless went extinct in stochastic experiments.

Finally, rVSV-G and rVSV-EBOV trials on PaKiT01 cells were best fit by models

assuming constitutive immunity, while rVSV-MARV infections on PaKiT01 were matched

equivalently by models assuming either induced or constitutive immunity—with induced models

favored over constitutive in AIC comparisons because one fewer parameter was estimated

(Figure 1-figure supplement 4-5; Supplementary File 4). For all virus infections, PaKiT01 cell

lines yielded β estimates a full order of magnitude higher than Vero or RoNi/7.1 cells, with each

β balanced by an immune response (either ρ, or ρ combined with ε) also an order of magnitude

higher than that recovered for the other cell lines (Figure 4; Table 1). As in RoNi/7.1 cells,

PaKiT01 parameter fits localized in the region corresponding to endemic equilibrium for the

deterministic theoretical model. Because constitutive immune processes can actually prohibit

initial pathogen invasion, constitutive immune fits to rVSV-MARV infections on PaKiT01 cell

lines consistently localized at or below the Branch point threshold for virus invasion ($R_0 = 1$).

During model fitting for optimization of ε, any parameter tests of ε values producing $R_0 < 1$

resulted in no infection and, consequently, produced an exceedingly poor fit to infectious time

series data. In all model fits assuming constitutive immunity, across all cell lines, parameter

estimates for ρ and ε traded off, with one parameter optimized at values approximating zero,

such that the immune response was modeled as almost entirely induced or entirely constitutive

(Table 1; Supplementary File 4). For RoNi/7.1 cells, even when constitutive immunity was

allowed, the immune response was estimated as almost entirely induced, while for rVSV-G and

rVSV-EBOV fits on PaKiT01 cells, the immune response optimized as almost entirely

constitutive. For rVSV-MARV on PaKiT01 cells, however, estimation of ρ was high under all

assumptions, such that any additional antiviral contributions from ε prohibited virus from

invading at all. The induced immune model thus produced a more parsimonious recapitulation of

these data because virus invasion was always permitted, then rapidly controlled.

Antiviral cells safeguard live cells against rapid cell mortality to elongate epidemic

duration *in vitro.*

In order to compare the relative contributions of each cell line's disparate immune

processes to epidemic dynamics, we next used our mean field parameter estimates to calculate

the initial 'antiviral rate'—the initial accumulation rate of antiviral cells upon virus invasion for

each cell-virus-MOI combination—based on the following equation:

$Antiviral\ Rate = \rho P_E P_S - \varepsilon P_S$ (8)

where P_E was calculated from the initial infectious dose (MOI) of each infection experiment and

P_S was estimated at disease free equilibrium:

$$P_E = 1 - e^{-MOI} \tag{9}$$

$$P_S = \frac{(b-\mu)(c+\mu)}{b(c+\mu+\varepsilon)} \tag{10}$$

Because ρ and ε both contribute to this initial antiviral rate, induced and constitutive immune

assumptions are capable of yielding equally rapid rates, depending on parameter fits. Indeed,

under fully induced immune assumptions, the induced antiviral acquisition rate (ρ) estimated for

rVSV-MARV infection on PaKiT01 cells was so high that the initial antiviral rate exceeded even

that estimated under constitutive assumptions for this cell-virus combination (Supplementary

File 4). In reality, we know that NPC1 receptor incompatibilities make PaKiT01 cell lines

constitutively refractory to rVSV-MARV infection (Ng and Chandran 2018) and that PaKiT01

cells also constitutively express the antiviral cytokine, IFN-α. Model fitting results suggest that

this constitutive expression of IFN-α may act more as a rapidly inducible immune response

following virus invasion than as a constitutive secretion of functional IFN-α protein.

Nonetheless, as hypothesized, PaKiT01 cell lines were by far the most antiviral of any in our

study—with initial antiviral rates estimated several orders of magnitude higher than any others in

our study, under either induced or constitutive assumptions (Table 1; Supplementary File 4).

RoNi/7.1 cells displayed the second-most-pronounced signature of immunity, followed by Vero

cells, for which the initial antiviral rate was essentially zero even under forced assumptions of

induced or constitutive immunity (Table 1; Supplementary File 4).

Using fitted parameters for β and ε, we additionally calculated R_0, the basic reproduction

number for the virus, for each cell line-virus-MOI combination (Table 1; Supplementary File 4).

We found that R_0 was essentially unchanged across differing immune assumptions for RoNi/7.1

and Vero cells, for which the initial antiviral rate was low. In the case of PaKiT01 cells, a high

initial antiviral rate under either induced or constitutive immunity resulted in a correspondingly

high estimation of β (and, consequently, R_0) which still produced the same epidemic curve that

resulted from the much lower estimates for β and R_0 paired with absent immunity. These

findings suggest that antiviral immune responses protect host tissues against virus-induced cell

mortality and may facilitate the establishment of more rapid within-host transmission rates.

Total monolayer destruction occurred in all cell-virus combinations excepting rVSV-

EBOV infections on RoNi/7.1 cells and rVSV-EBOV and rVSV-MARV infections on PaKiT01

cells. Monolayer destruction corresponded to susceptible cell depletion and epidemic turnover

where R-effective (the product of R_0 and the proportion susceptible) was reduced below one

(Figure 5). For rVSV-EBOV infections on RoNi/7.1, induced antiviral cells safeguarded remnant

live cells, which birthed new susceptible cells late in the time series. In rVSV-EBOV and rVSV-

MARV infections on PaKiT01 cells, this antiviral protection halted the epidemic (Figure 5; R-

effective <1) before susceptibles fully declined. In the case of rVSV-EBOV on PaKiT01, the

birth of new susceptibles from remnant live cells protected by antiviral status maintained late-

stage transmission to facilitate long-term epidemic persistence. Importantly, under fixed

parameter values for the infection incubation rate (σ) and infection-induced mortality rate (α),

models were unable to reproduce the longer-term infectious time series captured in data from

rVSV-EBOV infections on PaKiT01 cell lines without incorporation of cell births, an

assumption adopted in previous modeling representations of IFN-mediated viral dynamics in

tissue culture (Howat et al. 2006). In our experiments, we observed that cellular reproduction

took place as plaque assays achieved confluency.

Finally, because the protective effect of antiviral cells is more clearly observable

spatially, we confirmed our results by simulating fitted time series in a spatially-explicit,

stochastic reconstruction of our mean field model. In spatial simulations, rates of antiviral

acquisition were fixed at fitted values for ρ and ε derived from mean field estimates, while

transmission rates (β) were fixed at values ten times greater than those estimated under mean

field conditions, accounting for the intensification of parameter thresholds permitting pathogen

invasion in local spatial interactions (see Materials and Methods; Videos 1-3; Figure 5-figure

supplement 3; Supplementary File 5; Webb, Keeling, and Boots 2007). In immune capable time

series, spatial antiviral cells acted as 'refugia' which protected live cells from infection as each

initial epidemic wave 'washed' across a cell monolayer. Eventual birth of new susceptibles from

these living refugia allowed for sustained epidemic transmission in cases where some infectious

cells persisted at later timepoints in simulation (Videos 1-3; Figure 5- figure supplement 3).

<u>Discussion</u>

Bats are reservoirs for several important emerging zoonoses but appear not to experience

disease from otherwise virulent viral pathogens. Though the molecular biological literature has

made great progress in elucidating the mechanisms by which bats tolerate viral infections (Zhou

et al. 2016; Ahn et al. 2019; Xie et al. 2018; Pavlovich et al. 2018; Zhang et al. 2013), the impact

of unique bat immunity on virus dynamics within-host has not been well-elucidated. We used an

innovative combination of *in vitro* experimentation and within-host modeling to explore the

impact of unique bat immunity on virus dynamics. Critically, we found that bat cell lines

demonstrated a signature of enhanced interferon-mediated immune response, of either

constitutive or induced form, which allowed for establishment of rapid within-host, cell-to-cell

virus transmission rates (β). These results were supported by both data-independent bifurcation

analysis of our mean field theoretical model, as well as fitting of this model to viral infection

time series established in bat cell culture. Additionally, we demonstrated that the antiviral state

induced by the interferon pathway protects live cells from mortality in tissue culture, resulting in

in vitro epidemics of extended duration that enhance that probability of establishing a long-term

persistent infection. Our findings suggest that viruses evolved in bat reservoirs possessing

enhanced IFN capabilities could achieve more rapid within-host transmission rates without

causing pathology to their hosts. Such rapidly-reproducing viruses would likely generate extreme

virulence upon spillover to hosts lacking similar immune capacities to bats.

To achieve these results, we first developed a novel, within-host, theoretical model

elucidating the effects of unique bat immunity, then undertook bifurcation analysis of the

model's equilibrium properties under immune absent, induced, and constitutive assumptions. We

considered a cell line to be constitutively immune if possessing any number of antiviral cells at

disease free equilibrium but allowed the extent of constitutive immunity to vary across the

parameter range for ε, the constitutive rate of antiviral acquisition. In deriving the equation for

R_0, the basic reproduction number, which defines threshold conditions for virus invasion of a

tissue ($R_0 > 1$), we demonstrated how the invasion threshold is elevated at high values of

constitutive antiviral acquisition, ε. Constitutive immune processes can thus prohibit pathogen

invasion, while induced responses, by definition, can only control infections *post-hoc*. Once

thresholds for pathogen invasion have been met, assumptions of constitutive immunity will limit

the cellular mortality (virulence) incurred at high transmission rates. Regardless of mechanism

(induced or constitutive), interferon-stimulated antiviral cells appear to play a key role in

maintaining longer term or persistent infections by safeguarding susceptible cells from rapid

infection and concomitant cell death.

Fitting of our model to *in vitro* data supported expected immune phenotypes for different

bat cell lines as described in the literature. Simple target cell models that ignore the effects of

immunity best recapitulated infectious time series derived from IFN-deficient Vero cells, while

models assuming induced immune processes most accurately reproduced trials derived from

RoNi/7.1 (*Rousettus aegyptiacus*) cells, which possesses a standard virus-induced IFN-response.

In most cases, models assuming constitutive immune processes best recreated virus epidemics

produced on PaKiT01 (*Pteropus alecto*) cells, which are known to constitutively express the

antiviral cytokine, IFN-α (Zhou et al. 2016). Model support for induced immune assumptions in

fits to rVSV-MARV infections on PaKiT01cells suggests that the constitutive IFN-α expression

characteristic of *P. alecto* cells may represent more of a constitutive immune priming process

than a perpetual, functional, antiviral defense. Results from mean field model fitting were

additionally confirmed in spatially explicit stochastic simulations of each time series.

As previously demonstrated in within-host models for HIV (Coffin 1995; Perelson et al.

1996; Nowak et al. 1995; Bonhoeffer et al. 1997; Ho et al. 1995), assumptions of simple target-

cell depletion can often provide satisfactory approximations of viral dynamics, especially those

reproduced in simple *in vitro* systems. Critically, our model fitting emphasizes the need for

incorporation of top-down effects of immune control in order to accurately reproduce infectious

time series derived from bat cell tissue cultures, especially those resulting from the robustly

antiviral PaKiT01 *P. alecto* cell line. These findings indicate that enhanced IFN-mediated

immune pathways in bat reservoirs may promote elevated within-host virus replication rates

prior to cross-species emergence. We nonetheless acknowledge the limitations imposed by *in*

vitro experiments in tissue culture, especially involving recombinant viruses and immortalized

cell lines. Future work should extend these cell culture studies to include measurements of

multiple state variables (i.e. antiviral cells) to enhance epidemiological inference.

The continued recurrence of Ebola epidemics across central Africa highlights the

importance of understanding bats' roles as reservoirs for virulent zoonotic disease. The past

decade has born witness to emerging consensus regarding the unique pathways by which bats

resist and tolerate highly virulent infections (Brook and Dobson 2015; Xie et al. 2018; Zhang et

al. 2013; Ahn et al. 2019; Zhou et al. 2016; Ng et al. 2015; Pavlovich et al. 2018). Nonetheless,

an understanding of the mechanisms by which bats support endemic pathogens at the population

level, or promote the evolution of virulent pathogens at the individual level, remains elusive.

Endemic maintenance of infection is a defining characteristic of a pathogen reservoir (Haydon et

al. 2002), and bats appear to merit such a title, supporting long-term persistence of highly

transmissible viral infections in isolated island populations well below expected critical

community sizes (Peel et al. 2012). Researchers debate the relative influence of population-level

and within-host mechanisms which might explain these trends (Plowright et al. 2016), but

increasingly, field data are difficult to reconcile without acknowledgement of a role for persistent

infections (Peel et al. 2018; Brook et al. 2019). We present general methods to study cross-scale

viral dynamics, which suggest that within-host persistence is supported by robust antiviral

responses characteristic of bat immune processes. Viruses which evolve rapid replication rates

under these robust antiviral defenses may pose the greatest hazard for cross-species pathogen

emergence into spillover hosts with immune systems that differ from those unique to bats.

Materials and Methods

Key Resources Table[*]				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
cell line (Vero)	Kidney (normal, epithelial, adult)	ATCC	CCL-81	
cell line (*Rousettus aegyptiacus)*	Kidney (normal, epithelial, adult)	(Biesold et al. 2011; Kühl et al. 2011)	RoNi/7.1	
cell line (*Pteropus alecto*)	Kidney (normal, epithelial, adult)	(Crameri et al. 2009)	PaKiT01	
virus strain	Replication competent, recombinant vesicular stomatitis Indiana virus expressing eGFP	(Miller et al. 2012; Wong et al. 2010)	rVSV-G	
virus strain	Replication competent, recombinant vesicular stomatitis Indiana virus expressing eGFP & EBOV GP in place of VSV G	(Miller et al. 2012; Wong et al. 2010)	rVSV-EBOV	
virus strain	Replication competent, recombinant vesicular stomatitis Indiana virus expressing eGFP & MARV GP in place of VSV G	(Miller et al. 2012; Wong et al. 2010)	rVSV-MARV	
reagent	Hoechst 33342 Fluorescent Stain	ThermoFisher	cat #: 62249	
reagent	L-Glutamine Solution	ThermoFisher	cat #: 25030081	
reagent	Gibco HEPES	ThermoFisher	cat #: 15630080	

reagent	iTaq Universal SYBR Green Supermix	BioRad	cat #: 1725120	
commercial assay or kit	Quick RNA Mini Prep Kit	Zymo	cat #: R1054	
commercial assay or kit	Invitrogen Superscript III cDNA Synthesis Kit	ThermoFisher	cat #: 18080051	
software	MatCont (version 2.2)	(Dhooge et al. 2008)	MatCont	
R	R version 3.6.0	(R Core Team 2019)	R	

[*]Note that primers for *R. aegyptiacus* and *P. alecto* β-Actin, IFN-α, and IFN-β genes are listed in the Supplementary File 6.

Cell Culture Experiments.

Cells.

All experiments were carried out on three immortalized mammalian kidney cell lines:

Vero (African green monkey), RoNi/7.1 (*Rousettus aegyptiacus*) (Kühl et al. 2011; Biesold et al.

2011) and PaKiT01 (*Pteropus alecto*) (Crameri et al. 2009). The species identification of all bat

cell lines were confirmed morphologically and genetically in the publications in which they were

originally described (Kühl et al. 2011; Biesold et al. 2011; Crameri et al. 2009). Vero cells were

obtained from ATCC.

Monolayers of each cell line were grown to 90% confluency (~9 x 10^5 cells) in 6-well

plates. Cells were maintained in a humidified 37˚C, 5% CO_2 incubator and cultured in

Dulbecco's modified Eagle medium (DMEM) (Life Technologies, Grand Island, NY),

supplemented with 2% fetal bovine serum (FBS) (Gemini Bio Products, West Sacramento, CA),

and 1% penicillin-streptomycin (Life Technologies). Cells were tested monthly for mycoplasma

contamination while experiments were taking place; all cells assayed negative for contamination

at every testing.

Previous work has demonstrated that all cell lines used are capable of mounting a type I

IFN response upon viral challenge, with the exception of Vero cells, which possess an IFN-β

deficiency (Desmyter, Melnick, and Rawls 1968; Rhim et al. 1969; Emeny and Morgan 1979).

RoNi/7.1 cells have been shown to mount idiosyncratic induced IFN defenses upon viral

infection (Pavlovich et al. 2018; Kuzmin et al. 2017; Arnold et al. 2018; Kühl et al. 2011;

Biesold et al. 2011), while PaKiT01 cells are known to constitutively express the antiviral

cytokine, IFN-α (Zhou et al. 2016). This work is the first documentation of IFN signaling

induced upon challenge with the particular recombinant VSVs outlined below. We verified

known antiviral immune phenotypes via qPCR. Results were consistent with the literature,

indicating a less pronounced role for interferon defense against viral infection in RoNi/7.1 versus

PaKiT01 cells.

Viruses.

Replication-capable recombinant vesicular stomatitis Indiana viruses, expressing filovirus

glycoproteins in place of wild type G (rVSV-G, rVSV-EBOV, and rVSV-MARV) have been

previously described (Wong et al. 2010; Miller et al. 2012). Viruses were selected to represent a

broad range of anticipated antiviral responses from host cells, based on a range of past

evolutionary histories between the virus glycoprotein mediating cell entry and the host cell's

entry receptor. These interactions ranged from the total absence of evolutionary history in the

case of rVSV-G infections on all cell lines to a known receptor-level cell entry incompatibility in

the case of rVSV-MARV infections on PaKiT01 cell lines.

To measure infectivities of rVSVs on each of the cell lines outlined above, so as to

calculate the correct viral dose for each MOI, NH_4Cl (20 mM) was added to infected cell

cultures at 1–2 hours post-infection to block viral spread, and individual eGFP-positive cells

were manually counted at 12–14 hours post-infection.

Innate Immune Phenotypes via qPCR of IFN Genes.

Previously published work indicates that immortalized kidney cell lines of *Rousettus*

aegyptiacus (RoNi/7.1) and *Pteropus alecto* (PaKiT01) exhibit different innate antiviral immune

phenotypes through, respectively, induced (Biesold et al. 2011; Pavlovich et al. 2018; Kühl et al.

2011; Arnold et al. 2018) and constitutive (Zhou et al. 2016) expression of type I interferon

genes. We verified these published phenotypes on our own cell lines infected with rVSV-G,

rVSV-EBOV, and rVSV-MARV via qPCR of IFN-α and IFN-β genes across a longitudinal time

series of infection.

Specifically, we carried out multiple time series of infection of each cell line with each of

the viruses described above, under mock infection conditions and at MOIs of 0.0001 and

0.001—with the exception of rVSV-MARV on PaKiT01 cell lines, for which infection was only

performed at MOI=0.0001 due to limited viral stocks and the extremely low infectivity of this

virus on this cell line (thus requiring high viral loads for initial infection). All experiments were

run in duplicate on 6-well plates, such that a typical plate for any of the three viruses had two

control (mock) wells, two MOI=0.0001 wells and two MOI=0.001 wells, excepting PaKiT01

plates, which had two control and four MOI=0.0001 wells at a given time. We justify this

PaKiT01 exemption through the expectation that IFN-α expression is constitutive for these cells,

and by the assumption that any expression exhibited at the lower MOI should also be present at

the higher MOI.

For these gene expression time series, four 6-well plates for each cell line–virus

combination were incubated with virus for one hour at 37˚C. Following incubation, virus was

aspirated off, and cell monolayers were washed in PBS, then covered with an agar plaque assay

overlay to mimic conditions under which infection trials were run. Plates were then harvested

sequentially at timepoints of roughly 5, 10, 15, and 20 hours post-infection (exact timing varied

as multiple trials were running simultaneously). Upon harvest of each plate, agar overlay was

removed, and virus was lysed and RNA extracted from cells using the Zymo Quick RNA Mini

Prep kit, according to the manufacturer's instructions and including the step for cellular DNA

digestion. Post-extraction, RNA quality was verified via nanodrop, and RNA was converted to

cDNA using the Invitrogen Superscript III cDNA synthesis kit, according to the manufacturer's

instructions. cDNA was then stored at 4˚C and as a frozen stock at -20˚C to await qPCR.

We undertook qPCR of cDNA to assess expression of the type I interferon genes, IFN-α

and IFN-β, and the housekeeping gene, β-Actin, using primers previously reported in the

literature (Supplementary File 6). For qPCR, 2ul of each cDNA sample was incubated with 7ul

of deionized water, 1ul of 5UM forward/reverse primer mix and 10ul of iTaq Universal SYBR

Green, then cycled on a QuantStudio3 Real-Time PCR machine under the following conditions:

initial denaturation at 94˚C for 2 min followed by 40 cycles of: denaturation at 95˚C (5 sec),

annealing at 58˚C (15 sec), and extension at 72˚C (10 sec).

We report simple δ-Ct values for each run, with raw Ct of the target gene of interest

(IFN-α or IFN-β) subtracted from raw Ct of the β-Actin housekeeping gene in Figure 1-figure

supplement 6. Calculation of fold change upon viral infection in comparison to mock using the

δ-δ-Ct method (Livak and Schmittgen 2001) was inappropriate in this case, as we wished to

demonstrate constitutive expression of IFN-α in PaKiT01 cells, whereby data from mock cells

was identical to that produced from infected cells.

Plaque Assays and Time Series Imaging.

After being grown to ~90% confluency, cells were incubated with pelleted rVSVs

expressing eGFP (rVSV-G, rVSV-EBOV, rVSV-MARV). Cell lines were challenged with both a

low (0.0001) and high (0.001) multiplicity of infection (MOI) for each virus. In a cell monolayer

infected at a given MOI (m), the proportion of cells (P), infected by k viral particles can be

described by the Poisson distribution: $P(k) = \frac{e^{-m}m^k}{k!}$, such that the number of initially infected

cells in an experiment equals: $1 - e^{-m}$. We assumed that a ~90% confluent culture at each

trial's origin was comprised of ~9×10^5 cells and conducted all experiments at MOIs of 0.0001

and 0.001, meaning that we began each trial by introducing virus to, respectively, ~81 or 810

cells, representing the state variable 'E' in our theoretical model. Low MOIs were selected to

best approximate the dynamics of mean field infection and limit artifacts of spatial structuring,

such as premature epidemic extinction when growing plaques collide with plate walls in cell

culture.

Six well plates were prepared with each infection in duplicate or triplicate, such that a

control well (no virus) and 2-3 wells each at MOI 0.001 and 0.0001 were incubated

simultaneously on the same plate. In total, we ran between 18-39 trials at each cell-virus-MOI

combination, excepting r-VSV-MARV infections on PaKiT01 cells at MOI=0.001, for which we

ran only 8 trials due to the low infectivity of this virus on this cell line, which required high viral

loads for initial infection. Cells were incubated with virus for one hour at 37°C. Following

incubation, virus was aspirated off, and cell monolayers were washed in PBS, then covered with

a molten viscous overlay (50% 2X MEM/L-glutamine; 5% FBS; 3% HEPES; 42% agarose),

cooled for 20 minutes, and re-incubated in their original humidified 37˚C, 5% CO_2 environment.

After application of the overlay, plates were monitored periodically using an inverted

fluorescence microscope until the first signs of GFP expression were witnessed (~6-9.5 hours

post-infection, depending on the cell line and virus under investigation). From that time forward,

a square subset of the center of each well (comprised of either 64- or 36-subframes and

corresponding to roughly 60 and 40% of the entire well space) was imaged periodically, using a

CellInsight CX5 High Content Screening (HCS) Platform with a 4X air objective

(ThermoFisher, Inc., Waltham, MA). Microscope settings were held standard across all trials,

with exposure time fixed at 0.0006 sec for each image. One color channel was imaged, such that

images produced show GFP-expressing cells in white and non-GFP-expressing cells in black

(Figure 1-figure supplement 1).

Wells were photographed in rotation, as frequently as possible, from the onset of GFP

expression until the time that the majority of cells in the well were surmised to be dead, GFP

expression could no longer be detected, or early termination was desired to permit Hoechst

staining.

In the case of PaKiT01 cells infected with rVSV-EBOV, where an apparently persistent

infection established, the assay was terminated after 200+ hours (8+ days) of continuous

observation. Upon termination of all trials, cells were fixed in formaldehyde (4% for 15 min),

incubated with Hoechst stain (0.0005% for 15 min) (ThermoFisher, Inc., Waltham, MA), then

imaged at 4X on the CellInsight CX5 High Content Screening (HCS) Platform. The machine was

allowed to find optimal focus for each Hoechst stain image. One color channel was permitted

such that images produced showed live nuclei in white and dead cells in black.

Hoechst Staining.

Hoechst stain colors cellular DNA, and viral infection is thought to interfere with the

clarity of the stain (Dembowski and DeLuca, 2015). As such, infection termination, cell fixation,

and Hoechst staining enables generation of a rough time series of uninfectious live cells (i.e.

susceptible + antiviral cells) to complement the images which produced time series of

proportions infectious. Due to uncertainty over the exact epidemic state of Hoechst-stained cells

(*i.e.* exposed but not yet infectious cells may still stain), we elected to fit our models only to the

infectious time series derived from GFP-expressing images and used Hoechst stain images as a

post hoc visual check on our fit only (Figure 5; Figure 5-figure supplement 1-2).

Image Processing.

Images recovered from the time series above were processed into binary ('infectious' vs.

'non-infectious' or, for Hoechst-stained images, 'live' vs. 'dead') form using the EBImage

package (Pau et al. 2010) in R version 3.6 for MacIntosh, after methods further detailed in

Supplementary File 7. Binary images were then further processed into time series of infectious

or, for Hoechst-stained images, live cells using a series of cell counting scripts. Because of

logistical constraints (i.e. many plates of simultaneously running infection trials and only one

available imaging microscope), the time course of imaging across the duration of each trial was

quite variable. As such, we fitted a series of statistical models to our processed image data to

reconstruct reliable values of the infectious proportion of each well per hour for each distinct

trial in all cell line–virus-MOI combinations (Figure 1-figure supplement 2-3), as well as for

declining live cell counts from control well data derived from the Hoestch time series

(Supplementary File 1; Supplementary File 7; Figure 1-figure supplement 7). All original and

processed images, image processing and counting code, and resulting time series data are freely

available for download at the following FigShare repository: DOI:

10.6084/m9.figshare.8312807.

Mean Field Model.

Theoretical Model Details.

To derive the expression for R_0, the basic pathogen reproductive number *in vitro*, we

used Next Generation Matrix (NGM) techniques (Diekmann, Heesterbeek, and Metz 1990;

Heffernan, Smith, and Wahl 2005), employing Wolfram Mathematica (version 11.2) as an

analytical tool. R_0 describes the number of new infections generated by an existing infection in a

completely susceptible host population; a pathogen will invade a population when $R_0 >$

1 (Supplementary File 2). We then analyzed stability properties of the system, exploring

dynamics across a range of parameter spaces, using MatCont (version 2.2) (Dhooge et al. 2008)

for Matlab (version R2018a) (Supplementary File 3).

Theoretical Model Fitting.

The birth rate, b, and natural mortality rate, μ, balance to yield a population-level growth

rate, such that it is impossible to estimate both b and μ simultaneously from total population size

data alone. As such, we fixed b at .025 and estimated μ by fitting an infection-absent version of

our mean field model to the susceptible time series derived via Hoechst staining of control wells

for each of the three cell lines (Figure 1- figure supplement 7). This yielded a natural mortality

rate, μ, corresponding to a lifespan of approximately 121, 191, and 84 hours, respectively, for

Vero, RoNi/7.1, and PaKiT01 cell lines (Figure 1-figure supplement 7). We then fixed the virus

incubation rate, σ, as the inverse of the shortest observed duration of time from initial infection

to the observation of the first infectious cells via fluorescent microscope for all nine cell line –

virus combinations (ranging 6 to 9.5 hours). We fixed α, the infection-induced mortality rate, at

$\frac{1}{6}$, an accepted standard for general viral kinetics (Howat et al. 2006), and held c, the rate of

antiviral cell regression to susceptible status, at 0 for the timespan (<200 hours) of the

experimental cell line infection trials.

We estimated cell line–virus-MOI-specific values for β, ρ, and ε by fitting the

deterministic output of infectious proportions in our mean field model to the full suite of

statistical outputs of all trials for each infected cell culture time series (Figure 1-figure

supplement 2-3). Fitting was performed by minimizing the sum of squared differences between

the deterministic model output and cell line–virus-MOI-specific infectious proportion of the data

at each timestep. We optimized parameters for MOI = 0.001 and 0.0001 simultaneously to

leverage statistical power across the two datasets, estimating a different transmission rate, β, for

trials run at each infectious dose but, where applicable, estimating the same rates of ρ and ε

across the two time series. We used the differential equation solver lsoda() in the R package

deSolve (Soetaert, Petzoldt, and Setzer 2010) to obtain numerical solutions for the mean field

model and carried out minimization using the 'Nelder-Mead' algorithm of the optim() function

in base R. All model fits were conducted using consistent starting guesses for the parameters, β

(β= 3), and where applicable, ρ (ρ = 0.001) and ε (ε = 0.001). In the case of failed fits or

indefinite hessians, we generated a series of random guesses around the starting conditions and

continued estimation until successful fits were achieved.

All eighteen cell line–virus-MOI combinations of data were fit by an immune absent ($\varepsilon =$

$\rho = 0$) version of the theoretical model and, subsequently, an induced immunity ($\varepsilon = 0$; $\rho > 0$)

and constitutive immunity ($\varepsilon > 0$; $\rho > 0$) version of the model. Finally, we compared fits across

each cell line–virus-MOI combination via AIC. In calculating AIC, the number of fitted

parameters in each model (k) varied across the immune phenotypes, with one parameter (β)

estimated for absent immune assumptions, two (β and ρ) for induced immune assumptions, and

three (β, ρ, and ε) for constitutive immune assumptions. The sample size (n) corresponded to the

number of discrete time steps across all empirical infectious trials to which the model was fitted

for each cell-line virus combination. All fitting and model comparison script is freely available

for download at the following FigShare repository: DOI: 685 10.6084/m9.figshare.8312807.

Spatial Model Simulations.

Finally, we verified all mean field fits in a spatial context, in order to more thoroughly

elucidate the role of antiviral cells in each time series. We constructed our spatial model in C++

implemented in R using the packages Rcpp and RcppArmadillo (Eddelbuettel and Francois

2011; Eddelbuettel and Sanderson 2017). Following Nagai and Honda (2001) and Howat et al.

(2006), we modeled this system on a two-dimensional hexagonal lattice, using a ten-minute

epidemic timestep for cell state transitions. At the initialization of each simulation, we randomly

assigned a duration of natural lifespan, incubation period, infectivity period, and time from

antiviral to susceptible status to all cells in a theoretical monolayer. Parameter durations were

drawn from a normal distribution centered at the inverse of the respective fixed rates of μ, σ, α,

and *c*, as reported with our mean field model. Transitions involving the induced (ρ) and

constitutive (ε) rates of antiviral acquisition were governed probabilistically and adjusted

dynamically at each timestep based on the global environment. As such, we fixed these

parameters at the same values estimated in the mean field model, and multiplied both ρ and ε by

the global proportion of, respectively, exposed and susceptible cells at a given timestep.

In contrast to antiviral acquisition rates, transitions involving the birth rate (*b*) and the

transmission rate (β) occurred probabilistically based on each cell's local environment. The birth

rate, *b*, was multiplied by the proportion of susceptible cells within a six-neighbor circumference

of a focal dead cell, while β was multiplied by the proportion of infectious cells within a thirty-

six neighbor vicinity of a focal susceptible cell, thus allowing viral transmission to extend

beyond the immediate nearest-neighbor boundaries of an infectious cell. To compensate for

higher thresholds to cellular persistence and virus invasion which occur under local spatial

conditions (Webb, Keeling, and Boots 2007), we increased the birth rate, *b*, and the cell-to-cell

transmission rate, β, respectively, to six and ten times the values used in the mean field model

(Supplementary File 4). We derived these increases based on the assumption that births took

place exclusively based on pairwise nearest-neighbor interactions (the six immediately adjacent

cells to a focal dead cell), while viral transmission was locally concentrated but included a small

(7.5%) global contribution, representing the thirty-six cell surrounding vicinity of a focal

susceptible. We justify these increases and derive their origins further in Supplementary File 5.

We simulated ten stochastic spatial time series for all cell-virus combinations under all

three immune assumptions at a population size of 10,000 cells and compared model output with

data in Figure 5-figure supplement 3. Spatial model code is available for public access at the

following FigShare repository: DOI: 10.6084/m9.figshare.8312807.

Acknowledgments

CEB was supported by a National Science Foundation Graduate Research Fellowship at

Princeton University and a Miller Institute for Basic Research Fellowship at UC Berkeley.

Tissue culture experiments were funded by an NIH grant R01 AI134824 to KC. Work in LFW's

lab was funded by the Singapore National Research Foundation grants (NRF2012NRF-CRP001-

056 and NRF2016NRF-NSFC002-013). CD was supported by the German Research

Council (DFG) grant DFG SPP1596 (DR 772/10–2), the Federal Ministry of Education and

Research (BMBF) grant RAPID (#01KI1723A) and the EU Horizon 2020 grant EVAg

(#653316). The authors thank the Chandran lab at Albert Einstein College of Medicine – in

particular, Cecilia Harold, Megan Slough, Rohit Jangra, and Tanwee Alkutkar – for technical

support during tissue culture experiments. The authors further thank Jessica Metcalf and the

Graham lab at Princeton for conceptual guidance throughout the project's development.

<u>References</u>

Ahn, Matae, Danielle E. Anderson, Qian Zhang, Chee Wah Tan, Beng Lee Lim, Katarina Luko,

Ming Wen, et al. 2019. "Dampened NLRP3-Mediated Inflammation in Bats and

Implications for a Special Viral Reservoir Host." *Nature Microbiology* 4: 789–99.

https://doi.org/10.1038/s41564-019-0371-3.

Arnold, Catherine E, Jonathan C Guito, Louis A Altamura, Sean P Lovett, Elyse R Nagle,

Gustavo F Palacios, and Mariano Sanchez-lockhart. 2018. "Transcriptomics Reveal

Antiviral Gene Induction in the Egyptian Rousette Bat Is Antagonized in Vitro by Marburg

Virus Infection." *Viruses* 10 (607). https://doi.org/10.3390/v10110607.

Baccam, Prasith, Catherine Beauchemin, Catherine A Macken, Frederick G Hayden, and Alan S

Perelson. 2006. "Kinetics of Influenza A Virus Infection in Humans." *Journal of Virology*

80 (15): 7590–99. https://doi.org/10.1128/JVI.01623-05.

Biesold, Susanne E., Daniel Ritz, Florian Gloza-Rausch, Robert Wollny, Jan Felix Drexler,

Victor M. Corman, Elisabeth K V Kalko, Samuel Oppong, Christian Drosten, and Marcel

A. Müller. 2011. "Type I Interferon Reaction to Viral Infection in Interferon-Competent,

Immortalized Cell Lines from the African Fruit Bat Eidolon Helvum." *PLoS ONE* 6 (11).

https://doi.org/10.1371/journal.pone.0028131.

Bonhoeffer, S., R. M. May, G. M. Shaw, and M. A. Nowak. 1997. "Virus Dynamics and Drug

Therapy." *Proceedings of the National Academy of Sciences* 94 (13): 6971–76.

https://doi.org/10.1073/pnas.94.13.6971.

Brook, Cara E., and Andrew P. Dobson. 2015. "Bats as 'special' Reservoirs for Emerging

Zoonotic Pathogens." *Trends in Microbiology* 23 (3): 172–80.

https://doi.org/10.1016/j.tim.2014.12.004.

Brook, Cara E., Hafaliana Christian Ranaivoson, Christopher C. Broder, Andrew A.

Cunningham, Jean-Michel Héraud, Alison J. Peel, Louise Gibson, James L.N. Wood, C.

Jessica Metcalf, and Andrew P. Dobson. 2019. "Disentangling Serology to Elucidate

Henipa- and Filovirus Transmission in Madagascar Fruit Bats." *Journal of Animal Ecology*

00: 1– 16. https://doi.org/10.1111/1365-2656.12985.

Calisher, Charles H, James E Childs, Hume E Field, Kathryn V Holmes, and Tony Schountz.

2006. "Bats: Important Reservoir Hosts of Emerging Viruses." *Clinical Microbiology*

Reviews 19 (3): 531–45. https://doi.org/10.1128/CMR.00017-06.

Coffin, John M. 1995. "HIV Population Dynamics in Vivo: Implications for Genetic Variation,

Pathogenesis, and Therapy." *Science (New York, N.Y.)* 267 (5197): 483–89.

Crameri, Gary, Shawn Todd, Samantha Grimley, Jennifer A. McEachern, Glenn A. Marsh, Craig

Smith, Mary Tachedjian, et al. 2009. "Establishment, Immortalisation and Characterisation

of Pteropid Bat Cell Lines." *PLoS ONE* 4 (12): e8266.

https://doi.org/10.1371/journal.pone.0008266.

Desmyter, J, J L Melnick, and W E Rawls. 1968. "Defectiveness of Interferon Production and of

Rubella Virus Interference in a Line of African Green Monkey Kidney Cells (Vero)."

Journal of Virology 2 (10): 955–61. https://doi.org/J. Virol. 1968 2:1955.

Dhooge, A., W. Govaerts, Yu A. Kuznetsov, H. G E Meijer, and B. Sautois. 2008. "New

Features of the Software MatCont for Bifurcation Analysis of Dynamical Systems."

Mathematical and Computer Modelling of Dynamical Systems 14 (2): 147–75.

https://doi.org/10.1080/13873950701742754.

Diekmann, O., J.A.P. Heesterbeek, and J.A.J Metz. 1990. "On the Definition and Computation of

the Basic Reproduction Ratio R0 in Models for Infectious Diseases in Heterogenous

Populations." *Journal of Mathematical Biology* 28: 365–82.

Eddelbuettel, Dirk, and Romain Francois. 2011. "Rcpp: Seamless R and C++ Integration."

Journal of Statistical Software 40: 1–18. https://doi.org/10.1007/978-1-4614-6868-4.

Eddelbuettel, Dirk, and Conrad Sanderson. 2017. "RcppArmadillo: Accelerating R with High-

Performance C++ Linear Algebra." *Computational Statistics and Data Analysis* 71 (2014):

1–16.

Emeny, J. M., and M. J. Morgan. 1979. "Regulation of the Interferon System: Evidence That

Vero Cells Have a Genetic Defect in Interferon Production." *Journal of General Virology*

43 (1): 247–52. https://doi.org/10.1099/0022-1317-43-1-247.

Emery, V.C., A.V. Cope, E.F. Bowen, D. Gor, and P.D. Griffiths. 1999. "The Dynamics of

Human Cytomegalovirus Replication in Vivo." *The Journal of Experimental Medicine* 190

(2): 177–82. https://doi.org/10.1084/jem.190.2.177.

Haydon, Daniel T, Sarah Cleaveland, Louise H Taylor, and M Karen Laurenson. 2002.

"Identifying Reservoirs of Infection: A Conceptual and Practical Challenge." *Emerging*

Infectious Diseases 8 (12): 1468–73. https://doi.org/10.3201/eid0812.010317.

Heffernan, J.M., R.J. Smith, and L.M. Wahl. 2005. "Perspectives on the Basic Reproductive

Ratio." *Journal of The Royal Society Interface* 2 (4): 281–93.

https://doi.org/10.1098/rsif.2005.0042.

Ho, David D., Avidan U. Neumann, Alan S. Perelson, Wen Chen, John M. Leonard, and Martin

Markowitz. 1995. "Rapid Turnover of Plasma Virions and CD4 Lymphocytes in HIV-1

Infection." *Nature* 373: 123–26. https://doi.org/10.1038/373123a0.

Hooper, Peter, Sherif Zaki, Peter Daniels, and Deborah Middleton. 2001. "Comparative

Pathology of the Diseases Caused by Hendra and Nipah Viruses." *Microbes and Infection* 3

(4): 315–22. https://doi.org/10.1016/S1286-4579(01)01385-5.

Howat, Tom J, Cristina Barreca, Peter O'Hare, Julia R Gog, and Bryan T Grenfell. 2006.

"Modelling Dynamics of the Type I Interferon Response to in Vitro Viral Infection."

Journal of the Royal Society, Interface 3 (10): 699–709.

https://doi.org/10.1098/rsif.2006.0136.

Igarashi, Manabu, Heinz Feldmann, Yoshihiro Takadate, Tatsunari Kondoh, Manabu Igarashi,

Junki Maruyama, Rashid Manzoor, and Hirohito Ogawa. 2020. "Niemann-Pick C1

Heterogeneity of Bat Cells Controls Filovirus Tropism." *Cell Reports* 30 (2): 308-319.e5.

https://doi.org/10.1016/j.celrep.2019.12.042.

Kacprzyk, Joanna, Graham M. Hughes, Eva M. Palsson-McDermott, Susan R. Quinn, Sébastien

J. Puechmaille, Luke A. J. O'Neill, and Emma C. Teeling. 2017. "A Potent Anti-

Inflammatory Response in Bat Macrophages May Be Linked to Extended Longevity and

Viral Tolerance." *Acta Chiropterologica* 19 (2): 219–28.

https://doi.org/10.3161/15081109ACC2017.19.2.001.

Keeling, Matt J., and Pejman Rohani. 2008. *Modeling Infectious Diseases in Humans and*

Animals. Princeton, NJ: Princeton University Press.

Kühl, Annika, Markus Hoffmann, Marcel A. Müller, Vincent J. Munster, Kerstin Gnirß, Miriam

Kiene, Theodros Solomon Tsegaye, et al. 2011. "Comparative Analysis of Ebola Virus

Glycoprotein Interactions with Human and Bat Cells." *Journal of Infectious Diseases* 204:

S840–S849. https://doi.org/10.1093/infdis/jir306.

Kuzmin, I.V., Toni M. Schwarz, Philipp A. Ilinykh, Ingo Jordan, Thamas G. Ksiazek, Ravi

Sachidanandam, Christopher F. Basler, and Alexander Bukreyev. 2017. "Innate Immune

Response of Bat and Human Cells to Filoviruses: Commonalities and Distinctions." *Journal*

of Virology 91 (8): e02471-16. https://doi.org/10.1128/JVI.02471-16.

Livak, K J, and T D Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using Real-

Time Quantitative PCR and the $2^{\wedge}(-\Delta\Delta CT)$ Method." *Methods* 25 (4): 402–8.

https://doi.org/10.1006/meth.2001.1262.

Mahanty, Siddhartha, and Mike Bray. 2004. "Pathogenesis of Filoviral Haemorrhagic Fevers."

Lancet Infectious Diseases 4 (8): 487–98. https://doi.org/10.1016/S1473-3099(04)01103-X.

Miller, Emily Happy, Gregor Obernosterer, Matthijs Raaben, Andrew S Herbert, Maika S

Deffieu, Anuja Krishnan, Esther Ndungo, et al. 2012. "Ebola Virus Entry Requires the

Host-Programmed Recognition of an Intracellular Receptor." *The EMBO Journal* 31 (8):

1947–60. https://doi.org/10.1038/emboj.2012.53.

Morris, Sinead E., Andrew J. Yates, Rik L. de Stwart, Rory D. de Vries, Michael J. Mina, Ashley

N. Nelson, Wen-Hsuan W. Lin, Roger D. Kouyos, Diane E. Griffin, and Bryan T. Grenfell.

2018. "Modeling the Measles Paradox Reveals the Importance of Cellular Immunity in

Regulating Viral Clearance." *PLoS Pathogens* 14 (12): e1007493.

Neumann, AU, NP Lam, H Dahari, DR Gretch, TE Wiley, TJ Layden, and AS Perelson. 1998.

"Hepatitis C Viral Dynamics in Vivo and the Antiviral Efficacy of Interferon-Alpha

Therapy." *Science* 282: 103–7.

Ng, Melinda, and Kartik Chandran. 2018. "Unpublished Results."

Ng, Melinda, Esther Ndungo, Maryska Kaczmarek, Andrew S. Herbert, Tabea Binger, Rebekah

James, Rohit K. Jangra, et al. 2015. "NPC1 Contributes to Species-Specific Patterns of

Ebola Virus Infection in Bats." *ELife* 4: e11785. https://doi.org/10.7554/eLife.11785.

Nicholls, J M, L L Poon, K C Lee, W F Ng, S T Lai, C Y Leung, C M Chu, et al. 2003. "Lung

Pathogy of Fatal Severe Acute Respiratory Syndrom." *Lancet* 361: 1773–78.

837   https://doi.org/10.1016/S0140-6736(03)13413-7.

838 Nowak, M. A., and R. M. May. 2000. *Virus Dynamics: Mathematical Principles of Immunology*

839   *and Virology*. Oxford, UK: Oxford University Press.

840 Nowak, Martin A., Robert M. May, Rodney E. Phillips, Sarah Rowland-Jones, David G. Lalloo,

841   Steven McAdam, Paul Klenerman, et al. 1995. "Antigenic Oscillations and Shifting

842   Immunodominance in HIV-1 Infections." *Nature* 375: 606–11.

843 Nowak, Martin A, Sebastian Bonhoeffer, Andrew M Hill, Richard Boehme, Howard C Thomas,

844   and Hugh McDade. 1996. "Viral Dynamics in Hepatitis B Virus Infection." *Proceedings of*

845   *the National Academy of Sciences* 93: 4398–4402.

846   http://www.pnas.org/content/pnas/93/9/4398.full.pdf.

847 Pau, Gregoire, Florian Fuchs, Oleg Sklyar, Michael Boutros, and Wolfgang Huber. 2010.

848   "EBImage-an R Package for Image Processing with Applications to Cellular Phenotypes."

849   *Bioinformatics* 26 (7): 979–81. https://doi.org/10.1093/bioinformatics/btq046.

850 Pavlovich, Stephanie S., Sean P. Lovett, Galina Koroleva, Jonathan C. Guito, Catherine E.

851   Arnold, Elyse R. Nagle, Kirsten Kulcsar, et al. 2018. "The Egyptian Rousette Genome

852   Reveals Unexpected Features of Bat Antiviral Immunity." *Cell* 173 (5): 1098–1110.

853   https://doi.org/10.1016/j.cell.2018.03.070.

854 Pawelek, Kasia A., Giao T. Huynh, Michelle Quinlivan, Ann Cullinane, Libin Rong, and Alan S.

855   Perelson. 2012. "Modeling Within-Host Dynamics of Influenza Virus Infection Including

856   Immune Responses." *PLoS Computational Biology* 8 (6): e1002588.

857   https://doi.org/10.1371/journal.pcbi.1002588.

858 Peel, Alison J., Kate S. Baker, David T. S. Hayman, Christopher C. Broder, Andrew A.

859   Cunningham, Anthony R. Fooks, Romain Garnier, James L. N. Wood, and Olivier Restif.

2018. "Support for Viral Persistence in Bats from Age-Specific Serology and Models of

Maternal Immunity." *Scientific Reports* 8 (1): 3859. https://doi.org/10.1038/s41598-018-

22236-6.

Peel, Alison J, Kate S Baker, Gary Crameri, Jennifer A Barr, David TS Hayman, Edward

Wright, Christopher C Broder, et al. 2012. "Henipavirus Neutralising Antibodies in an

Isolated Island Population of African Fruit Bats." *PloS One* 7 (1): e30346.

https://doi.org/10.1371/journal.pone.0030346.

Perelson, A S, A U Neumann, M Markowitz, J M Leonard, and D D Ho. 1996. "HIV-1

Dynamics in Vivo: Virion Clearance Rate, Infected Cell Life-Span, and Viral Generation

Time." *Science (New York, N.Y.)* 271 (5255): 1582–86.

https://doi.org/10.1126/science.271.5255.1582.

Perelson, Alan S. 2002. "Modelling Viral and Immune System Dynamics." *Nature Reviews

Immunology* 2 (1): 28–36. https://doi.org/10.1038/nri700.

Plowright, Raina K., Alison J. Peel, Daniel G. Streicker, Amy Gilbert, Hamish McCallum, James

Wood, Michelle L. Baker, and Olivier Restif. 2016. "Transmission or Within-Host

Dynamics Driving Pulses of Zoonotic Viruses in Reservoir-Host Populations." *PLoS

Neglected Tropical Diseases* 10 (8): e0004796.

https://doi.org/10.1371/journal.pntd.0004796.

Radke, K L, C Colby, J R Kates, H M Krider, and D M Prescott. 1974. "Establishment and

Maintenance of the Interferon-Induced Antiviral State: Studies in Enucleated Cells."

Journal of Virology 13 (3): 623–30.

http://www.ncbi.nlm.nih.gov/pubmed/4362865%0Ahttp://www.pubmedcentral.nih.gov/arti

clerender.fcgi?artid=PMC355347.

Rasmussen, L., and L. B. Farley. 1975. "Inhibition of *Herpesvirus Hominis* Replication by

Human Interferon." *Infection and Immunity* 12 (1): 104–8.

Rhim, J. S., K. Schell, B. Creasy, and W. Case. 1969. "Biological Characteristics and Viral

Susceptibility of an African Green Monkey Kidney Cell Line (Vero)." *Proceedings of the*

Society for Experimental Biology and Medicine 132 (2): 670-678.

Saenz, Roberto A, Michelle Quinlivan, Debra Elton, Shona MacRae, Anthony S Blunden,

Jennifer A Mumford, Janet M Daly, et al. 2010. "Dynamics of Influenza Virus Infection and

Pathology." *Journal of Virology* 84 (8): 3974–83. https://doi.org/10.1128/JVI.02078-09.

Samuel, C. E., and G. S. Knutson. 1982. "Mechanism of Interferon Action." *Journal of*

Biological Chemistry 257 (19): 11791–95.

Schountz, Tony, Michelle L Baker, John Butler, and Vincent Munster. 2017. "Immunological

Control of Viral Infections in Bats and the Emergence of Viruses Highly Pathogenic to

Humans." *Frontiers in Immunology* 8 (September): Article 1098.

https://doi.org/10.3389/fimmu.2017.01098.

Soetaert, Karline, Thomas Petzoldt, and R Woodrow Setzer. 2010. "Package DeSolve: Solving

Initial Value Differential Equations in R." *Journal of Statistical Software* 33 (9): 1–25.

https://doi.org/10.18637/jss.v033.i09.

Stetson, Daniel B., and Ruslan Medzhitov. 2006. "Type I Interferons in Host Defense." *Immunity*

25 (3): 373–81. https://doi.org/10.1016/j.immuni.2006.08.007.

Team, R Core. 2019. "R: A Language and Environment for Statistical Computing." *R*

Foundation for Statistical Computing. Vienna, Austria.

Wang, Lin-Fa, and Danielle E Anderson. 2019. "Viruses in Bats and Potential Spillover to

Animals and Humans." *Current Opinion in Virology* 34: 79–89.

https://doi.org/10.1016/j.coviro.2018.12.007.

Webb, Steven D., Matt J. Keeling, and Mike Boots. 2007. "Host-Parasite Interactions between

the Local and the Mean-Field: How and When Does Spatial Population Structure Matter?"

Journal of Theoretical Biology 249 (1): 140–52. https://doi.org/10.1016/j.jtbi.2007.06.013.

Wong, Anthony C, Rohini G Sandesara, Nirupama Mulherkar, Sean P Whelan, and Kartik

Chandran. 2010. "A Forward Genetic Strategy Reveals Destabilizing Mutations in the

Ebolavirus Glycoprotein That Alter Its Protease Dependence during Cell Entry." *Journal of*

Virology 84 (1): 163–75. https://doi.org/10.1128/JVI.01832-09.

Xie, Jiazheng, Yang Li, Xurui Shen, Jiazheng Xie, Yang Li, Xurui Shen, Geraldine Goh, et al.

2018. "Dampened STING-Dependent Interferon Activation in Bats." *Cell Host and*

Microbe 23: 297–301.

Zhang, Guojie, Christopher Cowled, Zhengli Shi, Zhiyong Huang, Kimberly a Bishop-Lilly,

Xiaodong Fang, James W Wynne, et al. 2013. "Comparative Analysis of Bat Genomes

Provides Insight into the Evolution of Flight and Immunity." *Science* 339 (6118): 456–60.

https://doi.org/10.1126/science.1230835.

Zhou, Peng, Mary Tachedjian, James W Wynne, Victoria Boyd, Jie Cui, Ina Smith, Christopher

Cowled, et al. 2016. "Contraction of the Type I IFN Locus and Unusual Constitutive

Expression of IFN-α in Bats." *Proceedings of the National Academy of Sciences* 113 (10):

2696–2701. https://doi.org/10.1073/pnas.1518240113.

Tables

Table 1. Optimized parameters from best fit deterministic model and spatial approximation at MOI=.001

Cell Line	Virus	Immune Assumption	δ AIC Reduction†	Antiviral Rate	ε [lci – uci] *	ρ [lci – uci] *	β [lci – uci] *	mean field R_0	spatial β
Vero	rVSV-G	Absent	2	0	0 [0-0]	0 [0-0]	2.44 [1.52-3.36]	8.73	24.418
	rVSV-EBOV	Absent	2	0	0 [0-0]	0 [0-0]	1.5 [1.06-1.94]	5.42	14.996
	rVSV-MARV	Absent	2	0	0 [0-0]	0 [0-0]	0.975 [0.558-1.39]	3.45	9.752
RoNi/7.1	rVSV-G	Induced	2	7.03×10^{-5}	0 [0-0]	0.089 [0-0.432]	2.47 [1.49-3.45]	10.91	24.705
	rVSV-EBOV	Induced	2.01	2.87×10^{-5}	0 [0-0]	0.0363 [0-0.343]	0.685 [0.451-0.919]	3.04	6.849
	rVSV-MARV	Induced	2	1.40×10^{-5}	0 [0-0]	0.0177 [0-0.257]	1.23 [0.917-1.55]	5.48	12.324
PaKiT01	rVSV-G	Constitutive	29.9	.00209	0.00602 [0-0.019]	8.26×10^{-8} [0-4.75×10^{-7}]	3.45 [1.07-5.84]	6.20	34.516
	rVSV-EBOV	Constitutive	27.9	.00499	0.0478 [0-0.0958]	4.46×10^{-8} [0-4.37×10^{-7}]	34.5 [28.7-40.2]	18.82	344.821
	rVSV-MARV	Induced	2	.00687	0 [0-0]	13.1 [0-37.9]	3.25 [0-41.3]	8.83	32.452

†Improvement in AIC from next best model for same cell line-virus-MOI combination. All δ-AIC are reported in Supplementary File 4.
* lci = lower and uci = upper 95% confidence interval. No confidence interval is shown for spatial β which was fixed at 10 times the estimated mean for the mean field model fits when paired with equivalent values of ε and ρ.
All other parameters were fixed at: b=.025 (mean field), 0.15 (spatial); α = 1/6; c=0; µ= 1/121 (Vero), 1/191 (RoNi/7.1), and 1/84 (PaKiT01)

Main Figure Captions

Figure 1. Fitted time series of infectious cell proportions from mean field model for rVSV-G,

rVSV-EBOV, and rVSV-MARV infections (columns) on Vero, RoNi/7.1, and PaKiT01 cell

lines (rows) at MOI=0.001. Results are shown for the best fit immune absent model on Vero

cells, induced immunity model on RoNi/7.1 cells, and constitutive (for rVSV-VSVG and rVSV-

EBOV) and induced (for rVSV-MARV) immunity models on PaKiT01 cells. Raw data across all

trials are shown as open circles (statistical smoothers from each trial used for fitting are available

in Figure 1-figure supplement 2-3). Model output is shown as a solid crimson line (95%

confidence intervals by standard error = red shading). Panel background corresponds to

empirical outcome of the average stochastic cell culture trial (persistent infection = white; virus-

induced epidemic extinction = gray; immune-mediated epidemic extinction = black). Parameter

values are listed in Table 1 and Supplementary File 4. Results for absent/induced/constitutive

fitted models across all cell lines are shown in Figure 1-figure supplement 4 (MOI=0.001) and

Figure 1-figure supplement 5 (MOI=0.0001).

Figure 2. Two parameter bifurcations of the mean field model, showing variation in the

transmission rate, β, against variation in the pathogen-induced mortality rate, α, under diverse

immune assumptions. Panel (A) depicts dynamics under variably constitutive immunity, ranging

from absent (left: $\varepsilon = 0$) to high (right: $\varepsilon = .0025$). In all panel (A) plots, the rate of induced

immune antiviral acquisition (ρ) was fixed at 0.01. Panel (B) depicts dynamics under variably

induced immunity, ranging from absent (left: ρ=0) to high (right: ρ=1). In all panel (B) plots, the

rate of constitutive antiviral acquisition (ε) was fixed at 0.0001. Branch point curves are

represented as solid lines and Hopf curves as dashed lines. White space indicates endemic

equilibrium (persistence), gray space indicates limit cycles, and black space indicates no

infection (extinction). Other parameter values for equilibrium analysis were fixed at: b = .025, μ

= .001, σ = 1/6, c = 0. Special points from bifurcations analyses are listed in Supplementary File

3.

Figure 3. Two parameter bifurcations of the mean field model, showing variation in the

transmission rate, β, against variation in: (A) the induced immunity rate of antiviral acquisition

(ρ) and (B) the constitutive immunity rate of antiviral acquisition (ε). Panels show variation in

the extent of immunity, from absent (left) to high (right). Branch point curves are represented as

solid lines and Hopf curves as dashed lines. White space indicates endemic equilibrium

(persistence), gray space indicates limit cycling, and black space indicates no infection

(extinction). Other parameter values for equilibrium analysis were fixed at: b = .025, μ = .001, σ

= 1/6, α = 1/6, c = 0. Special points from bifurcations analyses are listed in Supplementary File

3.

Figure 4. Best fit parameter estimates for β and ρ or ε from mean-field model fits to MOI=0.001

time series data, atop (A,B) β – ρ and (C) β – ε bifurcation. Fits and bifurcations are grouped by

immune phenotype: (A) absent; (B) induced; (C) constitutive immunity, with cell lines differentiated

by shape (Vero=circles; RoNi/7.1 = triangles; PaKiT01=squares) and viral infections by color

(rVSV-G = green, rVSV-EBOV = magenta, rVSV-MARV = blue). Note that y-axis values are ten-

fold higher in panel (C). Branch point curves (solid lines) and Hopf curves (dashed lines) are

reproduced from Figure 3. White space indicates endemic equilibrium (pathogen persistence), gray

space indicates limit cycling (virus-induced epidemic extinction), and black space indicates no

infection (immune-mediated pathogen extinction). In panel (A) and (B), ε is fixed at 0; in panel (C),

ρ is fixed at 5×10^{-8} for bifurcation curves and estimated at 4×10^{-8} and 8×10^{-8} for rVSV-EBOV and

rVSV-G parameter points, respectively. Other parameter values were fixed at: $b = .025$, $\mu = 0.001$, σ

$= 1/6$, $\alpha = 1/6$, and $c = 0$ across all panels. Raw fitted values and corresponding 95% confidence

intervals for β, ρ, and ε, background parameter values, and AIC recovered from model fit, are

reported in Supplementary File 4. Parameter fits at MOI=0.0001 are visualized in Figure 4-figure

supplement 1.

Figure 5. Fitted time series of susceptible (green shading) and antiviral (blue shading) cell

proportions from the mean field model for rVSV-G, rVSV-EBOV, and rVSV-MARV infections

(columns) on Vero, RoNi/7.1, and PaKiT01 cell lines (rows) at MOI=0.001. Results are shown

for the best fit immune absent model on Vero cells, induced immunity model on RoNi/7.1 cells

and constitutive (rVSV-G and rVSV-EBOV) and induced (rVSV-MARV) immune models on

PaKiT01 cells. Combined live, uninfectious cell populations (S + A + E) are shown in tan

shading, with raw live, uninfectious cell data from Hoechst stains visualized as open circles. The

right-hand y-axis corresponds to R-effective (pink solid line) across each time series; R-effective

=1 is a pink dashed, horizontal line. Panel background corresponds to empirical outcome of the

average stochastic cell culture trial (persistent infection = white; virus-induced epidemic

extinction = gray; immune-mediated epidemic extinction = black). Parameter values are listed in

Supplementary File 4 and results for absent/induced/constitutive fitted models across all cell

lines in Figure 5-figure supplement 1 (MOI=0.001) and Figure 5-figure supplement 2

(MOI=0.0001).

1013

   Figure Supplement Captions

1015   **Figure 1 – figure supplement 1.** Cell culture models of viral propagation. (A), (B), and (C)

1016   show raw, original images of rVSV-EBOV propagation across Vero cell lines at, respectively,

1017   17, 21, and 28 hours post-infection (timesteps 2, 3, and 5 from trial Ver6_B1). (D), (E), and (F)

1018   show corresponding, binary images processed in the R package, EBImage. Cells expressing viral

1019   eGFP are depicted in white and uninfected/dead cells in black.

1020

Figure 1 – figure supplement 2. Time series data to which mean field mechanistic models were

fit, across rVSV-G (left), rVSV-EBOV (middle), and rVSV-MARV (right) infections on Vero,

RoNi/7.1, and PaKiT01 cell lines, at MOI=0.001. Open circles show raw data across all trials,

while red, dashed line gives the statistical mean of each trials, established from GAM model

incorporating random effects per trial. Results for MOI=0.0001 are shown in Figure 1 – figure

supplement 3.

1027

Figure 1 – figure supplement 3. Time series data to which mean field mechanistic models were

fit, across rVSV-G (left), rVSV-EBOV (middle), and rVSV-MARV (right) infections on Vero,

RoNi/7.1, and PaKiT01 cell lines, at MOI=0.0001. Open circles show raw data across all trials,

while red, dashed line gives the statistical mean of each trials, established from GAM model

incorporating random effects per trial. Results for MOI=0.001 are shown in Figure 1 – figure

supplement 2.

1034

Figure 1 – figure supplement 4. Figure replicates Figure 1 (main text) but includes all output

across mean field model fits assuming (A) absent immunity, (B) induced immunity, and (C)

constitutive immunity. Figure shows fitted time series of infectious cell proportions for rVSV-G,

rVSV-EBOV, and rVSV-MARV infections (columns) on Vero, RoNi/7.1, and PaKiT01 cell

lines (rows) at MOI=0.001. Raw data across all trials are shown as open circles and model output

as the solid crimson line (95% confidence intervals by standard error = red shading). Panel

background corresponds to empirical outcome of the average stochastic cell culture trial

(persistent infection = white; virus-induced epidemic extinction = gray; immune-mediated

epidemic extinction = black).

1044

1045   **Figure 1 – figure supplement 5.** Figure replicates Figure 1 – figure supplement 4 exactly but

1046   shows model fits and data for all cell-virus combinations at MOI=0.0001.

1047

Figure 1 – figure supplement 6. (A) IFN-α and (B) IFN-β gene expression profiles from qPCR

for rVSV infections on RoNi/7.1 and PaKiT01 cell lines. Panels show δ-Ct (raw Ct of IFN gene

assay subtracted from raw Ct of β-Actin housekeeping gene assay) across a time series for mock

(left), MOI=0.0001 (middle) and MOI=0.001 (right) infections across a time series. Viruses are

represented by color (rVSV-G=green, rVSV-EBOV=magenta, rVSV-MARV=blue). The red

dashed line at δ-Ct=37 corresponds to no expression; higher expression is indicated at lower

values for δ-Ct. qPCR was carried out using primers summarized in Supplementary File 6.

1055

1056   **Figure 1 – figure supplement 7.** Curve fits to control data for standard birth ($b = .025$) and

1057   natural mortality ($\mu = \frac{1}{121}, \frac{1}{191}, \frac{1}{84}$ hours for, respectively, Vero, RoNi/7.1, and PaKiT01 cell

1058   lines) rates across all three cell lines. Raw data from multiple trials are shown as open circles,

1059   statistical means as dashed black lines, with the output from the mean field model, using the

1060   fixed birth rate and estimated mortality rate, in solid green.

1061

Figure 4 – figure supplement 1. Best fit parameter estimates for β and ρ or ε from mean-field

model fits to MOI=0.0001 time series data, atop (A,B) β – ρ and (C) β – ε bifurcation. Fits and

bifurcations are grouped by immune phenotype: (A) absent; (B) induced; (C) constitutive

immunity, with cell lines differentiated by shape (Vero=circles; RoNi/7.1 = triangles;

PaKiT01=squares) and viral infections by color (rVSV-G = green, rVSV-EBOV = magenta,

rVSV-MARV = blue). Note that y-axis values are ten-fold higher in panel (C). Branch point

curves (solid lines) and Hopf curves (dashed lines) are reproduced from Figure 3 (main text).

White space indicates endemic equilibrium (pathogen persistence), gray space indicates limit

cycling (virus-induced epidemic extinction), and black space indicates no infection (immune-

mediated pathogen extinction). In panel (A) and (B), ε is fixed at 0; in panel (C), ε is fixed at

5×10^{-8} for bifurcation curves and estimated at 4×10^{-8} and 8×10^{-8} for rVSV-EBOV and rVSV-G

parameter points, respectively. To construct bifurcation curves, other parameter values were

fixed at: $b = 0.025$, $\mu = 0.001$, $\alpha = \frac{1}{6}$, and $c = 0$ across all panels. Raw fitted values and

corresponding 95% confidence intervals for β, ρ, and ε, background parameter values, and AIC

recovered from model fit, are reported in Supplementary File 4. Parameter fits at MOI=0.0001

are visualized in Figure 4 of the main text.

1078

Figure 5 – figure supplement 1. Figure replicates Figure 5 (main text) but includes all output

across mean field model fits assuming (A) absent immunity, (B) induced immunity, and (C)

constitutive immunity. Figure shows fitted time series of susceptible (green shading) and

antiviral (blue shading) cell proportions from the mean field model for rVSV-G, rVSV-EBOV,

and rVSV-MARV infections (columns) on Vero, RoNi/7.1, and PaKiT01 cell lines (rows) at

MOI=0.001. Combined live, uninfectious cell populations (S + A + E, summed across the time

series) is shown in tan shading, with raw live, uninfectious cell data from Hoechst stains of

terminal time series visualized as open circles. The right-hand y-axis corresponds to R-effective

(pink solid line) across each time series; R-effective = 1 is given as a pink dashed, horizontal

line. Panel background corresponds to empirical outcome of the average stochastic cell culture

trial (persistent infection = white; virus-induced epidemic extinction = gray; immune-mediated

epidemic extinction = black).

1091

1092   **Figure 5 – figure supplement 2.** Figure replicates Figure 5 – figure supplement 1 exactly but

1093   shows model fits and data for all cell-virus combinations at MOI=0.0001.

1094

Figure 5 – figure supplement 3. Spatial model state variable outputs, fit to MOI=0.001 data

only, for all 27 unique cell line - virus - immune assumption combinations: (A) absent immunity,

(B) induced immunity, and (C) constitutive immunity. Values for ρ and ε were fixed at

equivalent values to those optimized in mean field trials and β fixed at ten times the value

estimated under mean field conditions. Figure shows mean output from 10 runs of the spatial

stochastic model, on a 10,000 cell lattice for MOI=0.001 infections of rVSV-G, rVSV-EBOV,

and rVSV-MARV (columns) on Vero, RoNi/7.1, and PaKiT01 (rows) cell lines. Mean state

variable outputs are plotted as colored lines with 95% confidence intervals by standard error

shown in corresponding shading (infectious = red; susceptible = green; antiviral = blue). Raw

infectious cell data across all time trials are plotted as open red circles, with the Hoechst-stained

live cell population as open black circles. Modeled live, uninfectious cell populations (S+A+E)

are shown in tan shading in the background. Panel background shading corresponds to the mean

spatial model outcome for each cell line – virus combination (persistent infection = white; virus-

induced epidemic extinction = gray; immune-mediated epidemic extinction = black). All

parameter values are reported in Supplementary File 4.

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

 Video Captions

1129 **Video 1.** Two hundred hour time series of spatial stochastic model for rVSV-EBOV infection on

1130 10,000 cell grid for PaKiT01, assuming conditions of *absent immunity*: (A) spatial spread of

1131 infection, (B) time series of state variables. Parameter values are listed in Supplementary File 4.

1132

1133 Video 2. Two hundred hour time series of spatial stochastic model for rVSV-EBOV infection on

1134 10,000 cell grid for PaKiT01, assuming conditions of *induced immunity*: (A) spatial spread of

1135 infection, (B) time series of state variables. Parameter values are listed in Supplementary File 4.

1136

1137 Video 3. Two hundred hour time series of spatial stochastic model for rVSV-EBOV infection on

1138 10,000 cell grid for PaKiT01, assuming conditions of *constitutive immunity*: (A) spatial spread of

1139 infection, (B) time series of state variables. Parameter values are listed in Supplementary File 4.

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

Supplementary File Captions

Supplementary File 1. (A) Raw proportion infectious from cell culture images. Dataset gives raw proportion of infectious cells and time elapsed for all trials of all cell line/virus/MOI combinations, derived from image processing of binary images. (B) Raw proportion uninfectious from cell culture images. Dataset gives raw proportion of uninfectious cells and time elapsed for all trials of all cell line/virus/MOI combinations, derived from image processing of binary Hoechst-stained images. (C) Statistical mean of infectious time series for all trials of each cell line/virus/MOI experiment, from GAM fitted model incorporating random effects by trial. Data were smoothed to yield the proportion infectious per hourly timestep for each trial, and mean field mechanistic models were fit to the smoothed mean of all compiled trials for each cell line/virus/MOI combination. (D) Statistical mean of uninfectious time series for all eighteen cell line/virus/MOI experiments, from generalized linear model fit to Hoechst stain data reported on tab B. Note that these means were not used in epidemic model fitting but natural mortality rates for each cell line were derived from fitting an infection-absent model to the trajectory of susceptible decline for control trials for each cell line, as shown in Figure 1 – figure supplement 7. All original raw image files, processed binary images, and image processing code are available freely for download at the following FigShare repository: DOI: 10.6084/m9.figshare.8312807.

Supplementary Files 2-7 are all supplementary tables or text which do not require captions.

Legend: persistent infection · virus–induced epidemic extinction · immune–mediated epidemic extinction · modeled infectious time series

Columns: rVSV–G, rVSV–EBOV, rVSV–MARV

Rows: Vero, RoNi/7.1, PaKiT01

y-axis: proportion

x-axis: hrs post–infection

Legend: persistent infection | virus–induced epidemic extinction | immune–mediated epidemic extinction | modeled infectious time series

A. Absent Immunity

Columns: rVSV–G | rVSV–EBOV | rVSV–MARV
Rows: Vero | RoNi/7.1 | PaKiT01

B. Induced Immunity

Columns: rVSV–G | rVSV–EBOV | rVSV–MARV
Rows: Vero | RoNi/7.1 | PaKiT01

C. Constitutive Immunity

Columns: rVSV–G | rVSV–EBOV | rVSV–MARV
Rows: Vero | RoNi/7.1 | PaKiT01

y-axis: proportion
x-axis: hrs post-infection

A. Absent Immunity

rVSV−G   rVSV−EBOV   rVSV−MARV

B. Induced Immunity

C. Constitutive Immunity

persistent infection   virus−induced epidemic extinction   immune−mediated epidemic extinction   modeled infectious time series

Vero   RoNi/7.1   PaKiT01

proportion

hrs post−infection

A.

transmission, β

ρ

absent
immunity

B.

induced immunity

transmission, β

induced antiviral acquisition, ρ

.00001   .001   .1   1   10

Cell

● Vero

▲ RoNi/7.1

■ PaKiT01

Curve

— Branch

- - Hopf

Virus

● rVSV−G

● rVSV−EBOV

● rVSV−MARV

C.

constitutive immunity

transmission, β

constitutive antiviral acquisition, ϵ

.00001   .001   .1   1   10

Legend (top):
persistent infection   virus–induced epidemic extinction   immune–mediated epidemic extinction
proportion live cell   proportion susceptible   proportion antiviral

A. Absent Immunity

rVSV–G   rVSV–EBOV   rVSV–MARV

(panels labeled at right: Vero, RoNi/7.1, PaKiT01)

R-eff: ––

B. Induced Immunity

(panels labeled at right: Vero, RoNi/7.1, PaKiT01)

C. Constitutive Immunity

(panels labeled at right: Vero, RoNi/7.1, PaKiT01)

proportion

hrs post–infection

Legend (top of figure):
- persistent infection
- virus-induced epidemic extinction
- immune-mediated epidemic extinction
- proportion live cell
- proportion susceptible
- proportion antiviral

A. Absent Immunity

rVSV-G | rVSV-EBOV | rVSV-MARV

Vero | RoNi/7.1 | PaKiT01

R-eff: --

B. Induced Immunity

rVSV-G | rVSV-EBOV | rVSV-MARV

Vero | RoNi/7.1 | PaKiT01

C. Constitutive Immunity

rVSV-G | rVSV-EBOV | rVSV-MARV

Vero | RoNi/7.1 | PaKiT01

proportion

hrs post-infection

Supplementary File 2. Derivation of R_0

From the main text, we recall that our mean field model takes on the following form:

$$\frac{dP_S}{dt} = bP_D(P_S + P_A) - \beta P_S P_I - \mu P_S - \rho P_E P_S - \varepsilon P_S + cP_A \tag{1}$$

$$\frac{dP_A}{dt} = \rho P_E P_S + \varepsilon P_S - cP_A - \mu P_A \tag{2}$$

$$\frac{dP_E}{dt} = \beta P_S P_I - \sigma P_E - \mu P_E \tag{3}$$

$$\frac{dP_I}{dt} = \sigma P_E - \alpha P_I - \mu P_I \tag{4}$$

$$\frac{dP_D}{dt} = \mu(P_S + P_E + P_I + P_A) + \alpha P_I - bP_D(P_S + P_A) \tag{5}$$

Parameters are defined according to:

- b = natural birth rate; fixed at .025 to yield stable population sizes in *in vitro* models (Figure 1-figure supplement 5)
- μ = natural mortality rate; estimated for each cell line by fitting a simple birth-death model to susceptible declines from control trials; values at $\frac{1}{121}, \frac{1}{191}, \frac{1}{84}$ hours for, respectively, Vero, RoNi/7.1, and PaKiT01 cell lines
- β = transmission coefficient; estimated for each cell line/virus/MOI combination by fitting model to data
- α = infection-induced mortality rate (virulence); fixed at $\frac{1}{6}$ hours
- σ = incubation rate of cell transitioning from an exposed (E) to an infectious (I) state; measured for each cell line-virus combination as the inverse of the time from initial infection to first observation of infectious virus by GFP under the fluorescent micropscope. Respectively, fixed at the following rates: $\frac{1}{8.38}, \frac{1}{7.07}, \frac{1}{9.58}$ hours for rVSV-G, rVSV-EBOV, and rVSV-MARV infections on Vero cells; $\frac{1}{8}, \frac{1}{6.72}, \frac{1}{6.78}$ hours for rVSV-G, rVSV-EBOV, and rVSV-MARV infections on RoNi/7.1 cells; and $\frac{1}{7.17}, \frac{1}{6.03}, \frac{1}{6.42}$ hours for rVSV-G, rVSV-EBOV, and rVSV-MARV infections on PaKiT01 cells
- ρ = rate of infection-induced acquisition to antiviral state, based on the global proportion of exposed cells in the tissue culture; fixed at 0 under conditions of absent immunity and estimated under conditions of induced and constitutive immunity for all cell line/virus combinations by fitting model to joint data over both MOI-0.001 and 0.0001 data.
- ε = rate of constitutive acquisition to antiviral state, based on the global proportion of susceptible cells in the tissue culture; fixed at 0 under assumptions of absent and induced immunity; estimated under conditions of constitutive immunity for all cell line/virus combinations by fitting model to joint data over both MOI=0.001 and 0.0001 data.
- c = rate of return to susceptibility from antiviral status; fixed at 0 in all model fits under all assumptions of immunity. Regression from antiviral status to susceptibility occurs on longer time scales than the 200 hour duration of our cell culture trials
- We assume that $P_S + P_A + P_E + P_I + P_D = 1$

We rewrote the equations to model explicit cell densities within a defined well volume. Note that these two systems of equations are equivalent and that parameter values can be used interchangeably across both forms. In the following system, N indicates the total cell capacity of the area modeled:

$$\frac{dS}{dt} = bD\frac{(S+A)}{N} - \frac{\beta SI}{N} - \mu S - \frac{\rho ES}{N} - \varepsilon S + cA \tag{6}$$

$$\frac{dA}{dt} = \frac{\rho ES}{N} + \varepsilon S - cA - \mu A \tag{7}$$

$$\frac{dE}{dt} = \frac{\beta SI}{N} - \sigma E - \mu E \tag{8}$$

$$\frac{dI}{dt} = \sigma E - \alpha I - \mu I \tag{9}$$

$$\frac{dD}{dt} = \mu(S + E + I + A) + \alpha I - bD\frac{(S+A)}{N} \tag{10}$$

To solve for R_0, we adopted a next generation matrix approach, after Heffernan, Smith, and Wahl 2005, which we applied to the proportional system of equations (1-5, above). Step one of this process necessitated constructing two transition matrices: the F matrix, which describes how new infections are created and the V matrix which represents host transitions between exposed and infected states.

To build these F and V matrices, we first calculated the value of P_S at the disease free equilibrium (DFE, whereby P_I and P_E are equal to zero. We allow for the presence of antiviral cells (P_A) at disease free equilibrium, though their equilibrium quantity (and, by consequence, the extent of constitutive immunity at play in the system) depends on certain parameter values. When $\varepsilon = 0$, P_A will always be 0 at disease free equilibrium. By allowing the extent of constitutive immunity to vary with parameter values, we maintain consistency with the published literature, which suggests that, in constitutively antiviral bat cell lines (i.e. PaKiT01), some interferon-stimulated genes (ISGs) are perpetually expressed, like their IFN-α precursor, while others still require induction upon infection. At DFE, our antiviral population takes on the following form:

$$P_A^* = \frac{\varepsilon P_S}{(c+\mu)} \tag{11}$$

And, by extension, our susceptible population can be represented as:

$$P_S^* = \frac{(b-\mu)(c+\mu)}{b(c+\mu+\varepsilon)} \tag{12}$$

For cells with either absent or fully induced immunity, for which $\varepsilon = 0$, this contracts to:

$$P_S^* = \frac{(b-\mu)}{b} \tag{13}$$

Because we cannot have negative cell proportions at equilibrium, we impose the following constraint on the above parameters:

$$(b - \mu)(c + \mu) > 0 \tag{14}$$

When we substitute the above value for P_S* into the infectious equations, we find:

$$\frac{dP_E}{dt} = 0 = \frac{\beta(b-\mu)(c+\mu)}{b(c+\mu+\varepsilon)} P_I - \sigma P_E - \mu P_E \tag{15}$$

$$\frac{dP_I}{dt} = 0 = \sigma P_E - \alpha P_I - \mu P_I \tag{16}$$

Our F matrix then takes on the following form:

$$F = \begin{bmatrix} 0 & \frac{\beta(b-\mu)(c+\mu)}{b(c+\mu+\varepsilon)} \\ 0 & 0 \end{bmatrix} \tag{17}$$

And the V matrix takes on the following form:

$$V = \begin{bmatrix} -(\sigma + \mu) & 0 \\ \sigma & -(\alpha + \mu) \end{bmatrix} \tag{18}$$

After Heffernan, Smith, and Wahl 2005, it follows that $G = -FV^{-1}$. Thus, it follows that:

$$V^{-1} = \begin{bmatrix} \frac{1}{(\sigma+\mu)} & 0 \\ \frac{\sigma}{(\sigma+\mu)(\alpha+\mu)} & \frac{1}{(\alpha+\mu)} \end{bmatrix} \tag{19}$$

When this F matrix is multiplied by matrix $-V^{-1}$, we find:

$$G = \begin{bmatrix} \frac{\beta\sigma(b-\mu)(c+\mu)}{b(\sigma+\mu)(\alpha+\mu)(c+\mu+\varepsilon)} & \frac{\beta(b-\mu)(c+\mu)}{b(\alpha+\mu)(c+\mu+\varepsilon)} \\ 0 & 0 \end{bmatrix} \tag{20}$$

which, after matrix algebra, yields the following equation for R_0:

$$R_0 = \frac{\beta\sigma(b-\mu)(c+\mu)}{b(\sigma+\mu)(\alpha+\mu)(c+\mu+\varepsilon)} \tag{21}$$

Pathogens can invade a host tissue under conditions of $R_0 > 1$, or when the system satisfies the following inequality:

$$\beta\sigma(b - \mu)(c + \mu) > b(\sigma + \mu)(\alpha + \mu)(c + \mu + \varepsilon) \tag{22}$$

Finally, we note that for all cells lacking in constitutive antiviral properties ($\varepsilon = 0$), R_0 reduces to:

$$R_0 = \frac{\beta\sigma(b-\mu)}{b(\alpha+\mu)(\sigma+\mu)} \tag{23}$$

Supplementary File 3. Special Points from Bifurcation Analysis

Special Point	ε	ρ	β	P_I
Branch Point	0	.01	0.176	0
Hopf	0	.01	1.76	0.0133
Branch Point	0.0001	.01	0.193	0
Hopf	0.0001	.01	2.747	0.00944
Branch Point[†]	0.0025	.01	0.615	0
Branch Point	0.0001	0	0.193	0
Hopf	0.0001	0	1.407	0.0156
Branch Point	0.0001	.1	0.193	0
Hopf	0.0001	.1	16.918	0.00178
Branch Point[†]	0.0001	1	0.193	0

Note: All other parameters in this bifurcation analysis were fixed at the following
values: $b = .025$; $\mu = 0.001$; $\sigma = \frac{1}{6}$; $\alpha = \frac{1}{6}$; $c = 0$

[†]For these scenarios with high antiviral rates (either induced, ρ, or constitutive, ε), no
Hopf bifurcation was observed.

Supplementary File 4. Optimized parameters from all deterministic model outputs and spatial approximations

Assumption	Cell Line	Virus	δ AIC+	ε [lci – uci] *	ρ [lci – uci] *	MOI	Antiviral Rate	β [lci – uci] *	Mean Field R_0	Spatial β
Absent	Vero	rVSV-G	0	0	0 [0-0]	0.0001	0	3.14 [1.86-4.41]	11.211	--
						0.001	0	2.44 [1.52-3.36]	8.729	24.418
		rVSV-EBOV	0	0	0 [0-0]	0.0001	0	1.89 [1.42-2.36]	6.823	--
						0.001	0	1.5 [1.06-1.94]	5.416	14.996
		rVSV-MARV	0	0	0 [0-0]	0.0001	0	1.19 [0.624-1.76]	4.223	--
						0.001	0	0.975 [0.558-1.39]	3.454	9.752
	RoNi/7.1	rVSV-G	32.7	0	0 [0-0]	0.0001	0	2.33 [1.39-3.27]	10.278	--
						0.001	0	2.42 [1.54-3.3]	10.686	24.205
		rVSV-EBOV	222	0	0 [0-0]	0.0001	0	0.609 [.367-0.851]	2.707	--
						0.001	0	0.675 [.466-0.885]	3.001	6.753
		rVSV-MARV	3.36	0	0 [0-0]	0.0001	0	1.22 [0.891-1.54]	5.405	--
						0.001	0	1.23 [0.924-1.53]	5.457	12.284
	PaKiT01	rVSV-G	175	0	0 [0-0]	0.0001	0	2.48 [1.38-3.58]	6.689	--
						0.001	0	2.56 [1.58-3.55]	6.917	25.639
		rVSV-EBOV	27.9	0	0 [0-0]	0.0001	0	0.663 [.399-0.927]	1.811	--
						0.001	0	0.837 [0.215-1.46]	2.287	8.37
		rVSV-MARV	665	0	0 [0-0]	0.0001	0	0.393 [0-2.2]	1.068	--
						0.001	0	0.379 [0-1.66]	1.03	3.785
Induced	Vero	rVSV-G	2	0 [0-0]	4.55×10^{-10} [0-3.92×10^{-7}]	0.0001	3.04×10^{-14}	3.14 [1.86-4.41]	11.211	--
						0.001	3.04×10^{-13}	2.44 [1.52-3.36]	8.727	24.413
		rVSV-EBOV	2	0 [0-0]	4.64×10^{-8} [0-4.38×10^{-7}]	0.0001	3.11×10^{-12}	1.89 [1.42-2.36]	6.824	--
						0.001	3.10×10^{-11}	1.5 [1.06-1.94]	5.416	14.996
		rVSV-MARV	2	0 [0-0]	2.05×10^{-7} [0-5.97×10^{-7}]	0.0001	1.37×10^{-11}	1.19 [0.624-1.76]	4.223	--
						0.001	1.37×10^{-10}	0.975 [0.558-1.39]	3.454	9.752
	RoNi/7.1	rVSV-G	0	0 [0-0]	0.089 [0-0.432]	0.0001	7.03×10^{-6}	2.38 [1.37-3.39]	10.504	--
						0.001	7.03×10^{-5}	2.47 [1.49-3.45]	10.907	24.705
		rVSV-EBOV	0	0 [0-0]	0.0363 [0-0.343]	0.0001	2.87×10^{-6}	.622 [.336-.907]	2.763	--
						0.001	2.87×10^{-5}	.685 [.451-.919]	3.043	6.849
		rVSV-MARV	0	0 [0-0]	0.0177 [0-0.257]	0.0001	1.40×10^{-6}	1.22 [.882-1.56]	5.424	--
						0.001	1.40×10^{-5}	1.23 [.917-1.55]	5.475	12.324
	PaKiT01	rVSV-G	29.9	0 [0-0]	0.311 [0-1.02]	0.0001	1.63×10^{-5}	2.8 [1.43-4.17]	7.562	--
						0.001	0.000162	2.72 [1.4-4.03]	7.334	27.183
		rVSV-EBOV	1020	0 [0-0]	0.881 [0-3.32]	0.0001	0.0000461	1.08 [0-2.65]	2.947	--
						0.001	0.000461	1.55 [0-4.08]	4.233	15.495
		rVSV-MARV	0	0 [0-0]	13.1 [0-37.9]	0.0001	0.000687	0.698 [0-10.1]	1.9	--
						0.001	0.00687	3.25 [0-41.3]	8.828	32.452

Constitutive	Vero	rVSV-G	4	2.86x10^{-24} [0-0.00237]	3.05x10^{-8} [0-0.00237]	0.0001	2.04x10^{-12}	3.14 [1.81-4.47]	11.211	--
						0.001	2.04x10^{-11}	2.44 [1.49-3.4]	8.73	24.42
		rVSV-EBOV	4	5.42x10^{-20} [0-0.00202]	2.39x10^{-8} [0-0.00202]	0.0001	1.60x10^{-12}	1.89 [1.39-2.39]	6.824	--
						0.001	1.60x10^{-11}	1.5 [1.03-1.97]	5.416	14.996
		rVSV-MARV	247	0.00725 [0.00325-0.0113]	4.84x10^{-10} [0-3.92x10^{-7}]	0.0001	0.00259	2.42 [0.672-4.16]	4.565	--
						0.001	0.00259	1.83 [0.626-3.04]	3.464	18.347
	RoNi/7.1	rVSV-G	2	6.14x10^{-10} [0-3.93x10^{-7}]	0.089 [0-0.432]	0.0001	7.03x10^{-6}	2.38 [1.37-3.39]	10.504	--
						0.001	7.03x10^{-5}	2.47 [1.49-3.45]	10.906	24.703
		rVSV-EBOV	2.01	1.08x10^{-9} [0-3.93x10^{-7}]	0.0366 [0-0.346]	0.0001	2.90x10^{-6}	0.622 [0.336-0.908]	2.763	--
						0.001	2.89x10^{-5}	0.685 [0.45-0.919]	3.042	6.846
		rVSV-MARV	2	5.54x10^{-10} [0-3.93x10^{-7}]	0.0176 [0-0.257]	0.0001	1.40x10^{-6}	1.22 [0.882-1.56]	5.424	--
						0.001	1.39x10^{-5}	1.23 [0.917-1.55]	5.475	12.324
	PaKiT01	rVSV-G	0	0.00602 [0-0.019]	8.26x10^{-8} [0-4.75x10^{-7}]	0.0001	0.00209	3.68 [0.919-6.44]	6.593	--
						0.001	0.00209	3.45 [1.07-5.84]	6.189	34.516
		rVSV-EBOV	0	0.0478 [0-0.0958]	4.46x10^{-8} [0-4.37x10^{-7}]	0.0001	0.00499	15.6 [12.7-18.5]	8.518	--
						0.001	0.00499	34.5 [28.7-40.2]	18.823	344.821
		rVSV-MARV	2	3.99x10^{-7} [6.86x10^{-9}-7.91x10^{-7}]	13.1 [1.48-24.8]	0.0001	0.000687	0.699 [0-10.1]	1.902	--
						0.001	0.00687	3.24 [0-14.5]	8.815	32.406

[†]Best fit models indicated at δ-AIC = 0 are presented in Table 1 and Figure 1 and 3 (main text)

* lci = lower and uci = upper 95% confidence interval. No confidence interval is shown for spatial β which was fixed at 10 times the estimated mean for the mean field model fits when paired with equivalent values of ε and ρ.

All other parameters were fixed at the following values: b=.025 (mean field), .15 (spatial); α = 1/6; c=0; μ= 1/121 (Vero), 1/191(RoNi/7.1, and 1/84 (PaKiT01)

Supplementary File 5. Justification for parameter increase from mean field to spatial model.

Because spatial configurations elevate thresholds for pathogen invasion, as well for host cell persistence (Webb, Keeling, and Boots 2007), we were forced to elevate both birth rates (b) and transmission rates (β) above those used in the mean field model to apply to the spatial context. We here justify our chosen values for parameter increase:

In Webb, Keeling, and Boots 2007, the authors approximate spatial dynamics using a series of differential equations tracking the pairwise neighbor-neighbor interactions of a regular square lattice with a Von Neumann neighborhood. Webb *et al.* represent local reproduction as $b(1 - L_b)/z$ and local transmission as $\beta(1 - L_I)/z$ where z is the number of near-neighbor pairs, L_b is the proportion of global reproduction (births), and L_I is the proportion of global transmission in the system. Note that we altered the reproduction parameter r to b to reflect our own nomenclature.

Cumulatively, we can represent total reproduction (b) and total transmission (β_{tot}) in a system with both local and global effects as:

$$b_{tot} = \frac{b(1-L_b)}{z} + bL_b \tag{1}$$

$$\beta_{tot} = \frac{\beta(1-L_I)}{z} + \beta L_I \tag{2}$$

Our goal is to find a scalar (q) by which to multiply the mean field birth (q_b) and transmission (q_β) rates to values appropriate for the spatial context, such that:

$$b_{spatial} = q_b b_{mean} \tag{3}$$

$$\beta_{spatial} = q_\beta \beta_{mean} \tag{4}$$

Since we know that L_b and L_I both equal 1 in the mean field model, we can represent these interactions as:

$$b_{mean} = \frac{q_b b_{mean}(1-L_b)}{z} + q_b b_{mean} L_b \tag{5}$$

$$\beta_{mean} = \frac{q_\beta \beta_{mean}(1-L_b)}{z} + q_\beta \beta_{mean} L_b \tag{6}$$

And then solve for (q_b) and (q_β):

$$q_b = \frac{z}{zL_b+1-L_b} \tag{7}$$

$$q_\beta = \frac{z}{zL_I+1-L_I} \tag{8}$$

It is easy to see how, in the perfect pair approximation, when L_b or $L_I = 0$, $q = z$ meaning that the mean field transmission rate is simply multiplied by the number of nearest neighbors.

Equally, in the mean field context, when L_b or $L_I = 1$, $q = 1$.

From equation (7), it is straightforward to calculate (q_b) for our spatial model, since births are permitted only at the nearest-neighbor interface, and each cell has six adjacent neighbors. As such:

$$q_b = \frac{6}{6*0+1-0} = 6 \tag{9}$$

To this end, we multiplied all mean field birth rates (b=0.025) by 6 to equal .15 in the spatial model.

The dynamics are slightly more complicated in the case of the transmission rate modifier (q_β) because our spatial model allows cells to influence infection up to three 'rings' out from the nearest neighbor, such that each cell is affected by the proportion infectious in a 36-cell vicinity. These dynamics accurately reflect viral transmission, which, even in a plaque assay, can diffuse beyond the immediate neighbor-neighbor boundary, especially in a ten-minute timestep.

As such, we chose to represent transmission processes in the spatial model under assumptions of $L_I = .075$, allowing ~7.5% of transmission to be modeled globally and leading to a less extreme multiplication of the mean field transmission rate than assumed under the perfect pair approximation ($q = z$) illustrated in equation (9) above. Under these new assumptions, we modify the mean field transmission rate for the spatial context as follows:

$$q_\beta = \frac{36}{36*.075+1-.075} = 9.93 \tag{10}$$

Since this is an approximation and all transitions occur stochastically, we round q_β up to multiply all mean field transmission rates by ten in the spatial context.

Supplementary File 6. Primers for qPCR

Species	Cell Line	Gene	Primer	Sequence	Original Publication
Rousettus aegyptiacus	RoNi/7.1	β-Actin	Fwd	GGCTCCCAGCACAATGAAGA	Biesold *et al.*, 2011
			Rev	GGAGCCGCCGATCCA	Biesold *et al.*, 2011
		IFN-α	Fwd	GAGACTCCCCTGCTGGATGA	Cowled *et al.*, 2011
			Rev	ATAGAGGGTGATTCTCTGGAAGTATTTC	Cowled *et al.*, 2011
		IFN-β	Fwd	CAGCTATTTCCATGAGCTACAACTTG	Biesold *et al.*, 2011
			Rev	TTAACTGCCACAGGAGCTTCAG	Biesold *et al.*, 2011
Pteropus alecto	PaKiT01	β-Actin	Fwd	GGCTCCCAGCACAATGAAGA	Biesold *et al.*, 2011
			Rev	GGAGCCGCCGATCCA	Biesold *et al.*, 2011
		IFN-α	Fwd	GAGACTCCCCTGCTGGATGA	Cowled *et al.*, 2011
			Rev	ATAGAGGGTGATTCTCTGGAAGTATTTC	Cowled *et al.*, 2011
		IFN-β	Fwd	CTCTAGCACTGGCTGGAATGAA	Cowled *et al.*, 2011
			Rev	TGCCCACCGAGTGTCTCA	Cowled *et al.*, 2011

Supplementary File 7. Detailed methods for image and image data processing.

Image Processing.

       All image processing and data analysis was carried out in R version 3.6 for MacIntosh (R Core Team 2019). Original images were imported into R and processed via the package EBImage (Pau et al. 2010). Composite images of each well were first split into the 36 or 64-subframes from which they were composed (each subframe represents the visual region of focus for the microscope at the time of imaging). Each subframe was trimmed (to remove border effects), processed individually, and recompiled post-processing into binary form, such that light-colored regions of the original image were assigned a value of 1 (white), and dark regions were assigned a value of 0 (black). In the case of images of GFP-expressing cells, these white regions corresponded to "infectious" cells, while in images of Hoechst-stained nuclei, they indicated live, "uninfected" cells.

       Microscope focus was poor for control wells and for subframes early in the time series of each trial before GFP expression became apparent, and the original versions of any such subframes were light gray and grainy. Our image processing code identified these subframes as any which possessed a mean pixel value greater than .25 (a value substantially higher than any subframes in which true GFP-expressing or Hoechst-stained cells were visible) and subsequently converted the entire frame to 0 (black).

       All other subframes were processed following thresholding methods that have been previously described by the authors of EBImage (Pau et al. 2010). As a first pass, all pixels excepting the top 25% lightest pixels tallied were converted to 0 (black). Next, each image frame was walked through a series of secondary thresholding steps using if-else statements in R, such that the lightness threshold for "infectious" status was elevated in frames which were lighter overall due to manual variation in imaging and focusing. Processed subframes were then reconstructed into composite binary images, which were manually checked against original images to ensure consistent and reliable results.

       Post-processing into binary form, the number of discrete shapes with value of 1 were tabulated within each image, using the max(bwlabel(X)) function in EBimage, to determine a cell count per image, again corresponding to a count of infectious cells for GFP-expressing images and to a count of uninfected cells for Hoechst stain images. All image processing and counting scripts, in addition to the resulting data, are freely available for download at the following FigShare repository: DOI: 10.6084/m9.figshare.8312807.

Image Data processing.

       GFP-expressing images were processed and cells counted across the duration of each infection trial, thus generating a time series of infectious cells. For a subset of plates, infection was terminated, and cells were fixed, Hoechst stained, and imaged at periodic intervals across the duration of the time series. Processing of these images thus allowed for construction of a corresponding time series of live, uninfected cells. Because of logistical constraints (i.e. many plates of simultaneously running infection trials and only one available imaging microscope), the time course of imaging across the duration of each trial was quite variable. As such, we fitted a series of statistical models to our raw image data to reconstruct reliable values of the infectious proportion of each well per hour for each distinct trial in all cell line–virus-MOI combinations (Figure 1-figure supplement 2-3).

There was considerable natural variation in initial cell counts across each trial, resulting from subtle differences in the seeding density and growth duration of time until the trial was initiated (when wells were subjectively deemed to have reached "90% confluency"). Baseline cell counts were also different across our three cell lines, which varied substantially in natural size. To correct for this variation, we opted to model the proportion of infectious cell spaces per hour for each well, rather than rely on the raw count data. To this end, we collected the maximum number of live cells counted in susceptible control wells at timepoint 0 and set this count to a rough measure of 100% well occupancy for the cell line in question. Using this method, maximum cell counts were, respectively, 103712, 82308, and 92233 for Vero, RoNi/7.1, and PaKiT01 cell lines, reflecting innate variation in cell sizes. We then converted all cell counts tabulated via our image processing code across the infectious time trials into proportions by dividing the counts by the total number of possible cell spaces for the cell line in question. Though clearly subject to some error, these methods nonetheless maintained internal consistency in our counting methods and generated reasonable time series. We originally experimented with directly tabulating the proportion of infected versus uninfected space in our binary images; however, this approach impaired our ability to generalize across more or less densely seeded trials, as well as trials on cell lines of disparate sizes. As such, we adopted the count-to-proportion methods described here.

To generate an infectious time series of evenly distributed time steps against which to fit our mean field mechanistic model, we next fit a series of statistical models to the proportional data produced from the image processing methods described above. For the GFP-expressing data, we used the mgcv package in R (Wood 2001) to fit generalized additive models (GAMs) in the Gaussian family, with time elapsed (in hours) post infection as a predictor variable for proportion of infectious cells (the response variable). We fit a separate GAM model to each unique cell – virus – MOI combination, incorporating a random effect of well ID (such that each trial was modeled individually), and we fixed the smoothing parameter at k=7 for all trials, as recommended by the package author (Wood 2001). The gam.predict() function was used to return an estimate of infectious proportions per hour across the duration of each time series for each cell-virus-MOI combination.

The uninfected counts from the Hoechst stain data were much less numerous since each count required termination of the infection trial and fixation of cells; by definition, only one data point could be produced per trial. Due to this relative scarcity, we opted to fit a more standard linear regression model, again in the Gaussian family, to these data, rather than using the data-hungry GAM methods employed above. As before, we set time elapsed post infection as the predictor for the Hoechst stain data and produced a unique estimate of the proportion of uninfected cells per hour across the duration of the longest-observed trial. No random effects were included in this model, and the resulting time series were used to estimate natural mortality rates for each cell line, when fit to control well data depicting natural susceptible decline (Figure 1-figure supplement 7).