An optimized competitive-aging method reveals gene-drug interactions underlying the chronological lifespan of *Saccharomyces cerevisiae*

J. Abraham Avelar-Rivas[1], Michelle Munguía-Figueroa[1], Alejandro Juárez-Reyes[1], Erika Garay[1], Sergio E. Campos[1], Noam Shoresh[2] & Alexander DeLuna[1,*]

[1] *Centro de Investigación y de Estudios Avanzados del IPN, Unidad de Genómica Avanzada (Langebio), 36824 Irapuato, Guanajuato, Mexico.*
[2] *Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA.*

*Corresponding author: alexander.deluna@cinvestav.mx

Author's emails:
J. Abraham Avelar-Rivas (jesus.avelar@cinvestav.mx)
Michelle Munguía-Figueroa (michelle.munguia@cinvestav.mx)
Alejandro Juárez-Reyes (alejandro.juarez@cinvestav.mx)
Erika Garay (egaray.gar@gmail.com)
Sergio E. Campos (sergio.e.campos@cinvestav.mx)
Noam Shoresh (noamshoresh@gmail.com)
Alexander DeLuna (alexander.deluna@cinvestav.mx)

Keywords: Post-mitotic aging; Genetic analysis; High-throughput screening; Gene-drug interactions; Saccharomyces cerevisiae

ABSTRACT

The chronological lifespan of budding yeast is a model of aging and age-related diseases. This paradigm has recently allowed genome-wide screening of genetic factors underlying post-mitotic viability in a simple unicellular system, which underscores its potential to provide a comprehensive view of the aging process. However, results from different large-scale studies show little overlap and typically lack quantitative resolution to derive interactions among different aging factors. We previously introduced a sensitive, parallelizable approach to measure the chronological-lifespan effects of gene deletions based on the competitive aging of fluorescence-labeled strains. Here, we present a thorough description of the method, including an improved multiple-regression model to estimate the association between death rates and fluorescent signals, which accounts for possible differences in growth rate and experimental batch effects. We illustrate the experimental procedure—from data acquisition to calculation of relative survivorship—for ten deletion strains with known lifespan phenotypes, which is achieved with high technical replicability. We apply our method to screen for gene-drug interactions in an array of yeast deletion strains, which reveals a functional link between protein glycosylation and lifespan extension by metformin. Competitive-aging screening coupled to multiple-regression modeling provides a powerful, straight-forward way to identify aging factors in yeast and their interactions with pharmacological interventions.

INTRODUCTION

A major challenge in aging research is to describe the way in which different genetic pathways and biochemical processes mediating aging are interconnected to one another (Kirkwood, 2008; López-Otín et al., 2013). Simple cellular models provide a starting point to grant a systems-level understanding of aging, in which the lifespan phenotype is addressed as a complex trait resulting from the action of multiple genes, cellular processes, environmental factors, and their interactions.

The chronological lifespan (CLS) of *Saccharomyces cerevisiae* is used to describe genetic, nutrimental, and pharmacological factors underlying survivorship of post-mitotic, non-dividing cells (Longo and Fabrizio, 2012). The budding yeast's replicative-lifespan and CLS are simple experimental models that have been used to reveal the conserved lifespan-extending effects of reduced TOR and RAS/PKA signaling, as well as the anti-aging effect of rapamycin, spermidine, and caloric restriction (Wei et al., 2008; Eisenberg et al., 2009; Longo and Fabrizio, 2012; Gems and Partridge, 2013). Traditionally, the CLS of a yeast-cell population is measured by counting colony-forming units from samples of a long-term stationary-phase culture (Longo et al., 2012; Hu et al., 2013). More recently, large-scale screening approaches have been implemented to screen for genetic aging factors in yeast. These studies provide unbiased catalogues of CLS mutant phenotypes (Powers et al., 2006; Fabrizio et al., 2010; Matecic et al., 2010; Burtner et al., 2011; Garay et al., 2014), mutants with diminished or enhanced response to dietary restriction or nutrient limitation (Gresham et al., 2011; Campos et al., 2018), and CLS phenotypes of collections of wild isolates and lines derived from biparental crosses (Jung et al., 2018; Barré et al., 2019).

A current limitation in the field is that large-scale CLS-phenotyping screens have

resulted in a large number of false positive hits when further confirmed by smaller-scale

approaches, ranging from 50% to 94% (Powers et al., 2006; Fabrizio et al., 2010;

Matecic et al., 2010; Burtner et al., 2011; Garay et al., 2014). In addition, comparisons

of different large-scale studies show that there is little overlap among the identified

genetic factors, which could be explained in part by differences in genotypic

background, media composition, and subtle environmental variations (Smith et al.,

2016). In addition, changes in controlled or uncontrolled environmental conditions are

known to be important modifiers of CLS phenotypes and confounding causes of aging

(Burtner et al., 2009, 2011; Santos et al., 2015; Smith et al., 2016; Campos and

DeLuna, 2019). In this context, a combination of high throughput and resolution is much

needed to correctly determine not only genetic aging factors, but also to quantitatively

derive their interactions with nutrimental, chemical, or pharmacological environments.

In an effort to improve the throughput of CLS screening without sacrificing phenotyping

sensitivity, we previously introduced a competition-based method for quantitative large-

scale genetic analysis (Garay et al., 2014). In brief, each RFP-labeled single-deletion

strain is mixed with a CFP wild-type reference and grown to saturation; fluorescence

signal in outgrowth cultures is used to estimate the relative number of viable cells in the

non-dividing culture at different time points in stationary phase. This approach

recapitulates known CLS factors and suggests new lifespan phenotypes in yeast (Garay

et al., 2014). More recently, we have used this approach to screen for dietary restriction

factors, namely CLS gene-deletion phenotypes that are aggravated or alleviated when

yeast populations are aged under a poor nitrogen source (Campos et al., 2018).

In this study, we describe an optimized multiple regression modeling strategy to analyze

measurements from our competition-based approach for CLS genetic analysis in yeast,

by accounting for possible differences in growth rate and experimental batch effects. In

addition, we provide a systematic analysis of the method's replicability and data-

analysis scripts. For ten knockout strains, we compare the replicability of our results

with those obtained with a useful parallelizable approach based on outgrowth kinetics

(Murakami et al., 2008; Jung et al., 2015). Importantly, we take advantage of our

improved data-analysis method to derive gene-drug interactions by measuring the

relative effects on survival of metformin in 76 deletion strains of widely conserved

genes. We discuss the potential of competitive-aging screening to describe large

numbers of genetic and environmental interactions underlying aging and longevity in

aging cells.

MATERIALS AND METHODS

Strains and media. Ten single-gene deletion strains targeting *ATG1*, *HAP3*, *MSN2*,

MSN4, *RAS2*, *RIM15*, *RPS16A*, *STE12*, *GLN3*, and *SWR1* were generated *de novo* by

PCR-based gene replacement in the YEG01-RFP background(Garay et al., 2014) using

the *natMX4* module from pAG25 (Euroscarf). In addition, two isogenic reference strains

were generated over the YEG01-RFP and YEG01-CFP backgrounds by deleting the

neutral *HO* locus. Lithium-acetate transformation and 100 µg/ml clonNAT (Werner

BioAgents) were used. The resulting strains were *MAT-α xΔ::natMX4 PDC1-*

RFP.CaURA3MX4 can1Δ::STE2pr-SpHIS5 lyp1Δ ura3Δ0 his3Δ1 LEU2 MET15; strong

constitutive expression of fluorescent proteins during exponential growth is achieved by

carboxyl-terminal fusion to the Pdc1 protein.

For gene-drug interactions, a collection of RFP-tagged gene-deletion strains was

generated by mating an array of 85 strains from the yeast deletion collection to the *hoΔ*

YEG01-RFP SGA-starter strain, as previously described(Garay et al., 2014). Resulting

prototrophic strains were *MAT*-a *xΔ::kanMX4 PDC1-RFP.CaURA3MX4 can1Δ::STE2pr-*

SpHIS5 lyp1Δ ura3Δ0 his3Δ1 LEU2 MET15. The CFP reference strain was the neutral

marker reference *his3Δ::kanMX4*. Double-marker strains and CLS data were

successfully obtained for 76 deletion strains (Supplementary Data S1).

Aging medium was synthetic complete (SC) with ammonium, 2% glucose, and 0.2%

amino acid supplement mix (CSHL Manual, 2005), without buffering. 40mM metformin

(Sigma D150959) was used where indicated. Single-culture and competitive outgrowth

kinetics were done in low fluorescence YNB medium (YNB-lf) with 2% glucose and

complete amino-acids supplement mix (Sigma Y1501 completed with Sigma

U0750)(Sheff and Thorn, 2004). All cultures were incubated in a growth chamber at 30

ºC and 80%-90% relative humidity without shaking; aging co-cultures were vigorously

shaken at every sampling point. For media recipes, see Supplementary Table S1.

Competitive-aging culture setup and outgrowth measurements. To obtain relative

CLS measurements, mutant ($x\Delta_{RFP}$) and reference strains (WT_{RFP} and WT_{CFP}) were pre-

cultured separately until saturation. Saturated cultures were mixed in a 2:1

mutant:reference ratio and ~1.5 µL aliquots were transferred with a 96 solid-pin

replicator (V&P Scientific VP 407) onto 96 semi deepwell plates (Nunc 260251) with 750

µL of fresh aging medium and disposable plastic covers.

Outgrowth sampling of the competitive-aging culture began 3 days after initial

inoculation (time zero) and was repeated initially every 24 h and 48-72 h afterwards for

up to 16 days after time zero. At each sampling day, T_i, 12 µL of shaken aging cultures

were inoculated onto 140µL fresh YNB-lf medium in clear polystyrene 96-well plates

(Corning 3585). Outgrowth was monitored every 1-3 h until cultures reached stationary

phase, by measuring raw fluoresence of mCherry (*RFP*, Ex 587/5 nm and Em 610/5

nm), Cerulean (*CFP*, Ex 433/5 nm and Em 475/5 nm), and absorbance at 600 nm

(*OD_{600}*) using a Tecan Infinite M1000 reader integrated to a robotic platform (Tecan

Freedom EVO200). Outgrowth cultures were resuspended by vigorous shaking right

before every measurement. To increase fluorescence-signal dynamic range, an

outgrowth culture with the samples was measured one day before the actual experiment

to calculate the optimal-gain at late exponential-growth, when fluorescence signal is at

its maximum level. Optimal gain values were fixed for all measurements in the

experiment; values used were 140-167 and 131-137 for *RFP* and *CFP*, respectively.

Death-rate calculation from competitive-aging outgrowth data. The $\frac{RFP}{CFP}$ signal ratio

was used to estimate the number of cells expressing each fluorescent signal. Auto

fluorescence background was defined as the *RFP* and *CFP* signal of WT$_{CFP}$ and WT$_{RFP}$

monocultures, respectively, and was subtracted from all competitive aging cultures.

Using data from different measurements in the outgrowth cultures (t_j, in hours) of each

stationary-phase sampling point (T_i, in days), the signal ratio $ln\left(\frac{RFP}{CFP}\right)_{T_i,t_j}$ of each sample

(w) was fitted to the linear model $A_w + S_w \cdot T_i + G_w \cdot t_j + C_{T_i, t_j}$. The ratio of viable cells at

the beginning of the experiment is modeled in A, while S (relative survivorship) is the

death-rate difference of the mutant and wild-type reference, and G (relative growth) is

their growth-rate difference in the outgrowth co-culture. In addition, the term C_{T_i, t_j} was

introduced to consider the systematic variation of each plate at each stationary-phase

sampling point $T_i t_j$. A complete description of the model and its implementation is

provided in Supplementary Note S1.

Deriving gene-drug interactions from CLS phenotypes. To identify gene-drug

interactions, namely cases in which the CLS phenotype of a gene-deletion strain is

significantly aggravated or alleviated by treatment with a drug, the relative survivorship

of a set of 76 deletion strains was measured with and without 40 mM metformin.

Deletion strains were randomly selected from the yeast deletion collection, considering

only genes with a mammalian ortholog (Ensembl). Relative survivorship values were

rescaled to a dimensionless parameter, given that metformin increases yeast CLS and

that the relative survivorship S expressed in days is constrained by the death rate of the

wild-type reference, r_{wt}, which is different in each condition. For each mutant under a

given condition, rescaled survivorship (rS) was defined as $rS = -\ln\left(1 - \frac{S}{r_{wt}}\right)$, where r_{wt}

is the average death rate of all control wild-type competitions in each plate. Gene-drug

interactions were defined as cases in which a mutant's rS in SC was significantly

different to the rS in SC + metformin ($p<0.05$; t-test).

Measuring death rate in monoculture from outgrowth kinetics. CLS estimates

based on outgrowth kinetics of single-population aging cultures were adapted from a

well established high-throughput method(Murakami et al., 2008; Jung et al., 2015).

Culture density was monitored by measuring absorbance at OD_{600nm}; background

signal—OD_{600} at outgrowth inoculation—was subtracted to each data point. At each

aging sampling point T_i (days), the time-shift in hours, t, to reach a fixed cell density of

OD_{600}=0.35 reports for the remaining fraction of viable cells in the population, as

previously reported(Murakami et al., 2008). Specifically, for each successive age time

point, the percent of viability (V_T) was calculated using the equation $V_T = \frac{1}{2(\Delta t_n/\delta)}$, where

Δt_n is the time shift and δ is the doubling time. Death rates were the exponential decay

rates calculated by fitting all V_T data points as a function of time in stationary phase to a

first-order exponential model (Matlab, *fit*).

Live/dead cell viability assays. Stationary-phase cultures (5 μL) were transferred to

50 μL of BD FACSFlow (BD #342003) with 0.1% propidium iodide (PI) and 0.1%

SYTO9 green (ThermoFisher L34952) in 96-well plates (Corning 3585), shaken for 1

min at 1,000 rpm in a plate shaker (Heidolph Titramax 1000), and incubated at room

temperature in the dark for 20 minutes. An analytical flow cytometer with a high-

throughput sampler (LSRFortessa-HTS, Becton Dickinson) was used to capture 10,000

events. SYTO9 fluorescence was excited with a 488nm blue laser and collected through

a 525/50 nm band-pass filter and a 505LP emission filter, while PI signal was excited

with a 488nm blue laser and collected through a 712/21 band-pass filter. Events with a

increased PI/SYTO9 signal were counted as alive.

RESULTS

Relative survivorship in stationary phase can be estimated from bulk fluorescence signal of two populations in co-culture

We sought to optimize data analysis and systematically test a competition-based method aimed at describing CLS phenotypes of viable gene-deletion strains in budding yeast. To directly measure the lifespan effects, we tracked changes in the relative abundances of RFP- deletion ($x\Delta_{RFP}$) and CFP-tagged wild-type (WT_{CFP}) strains in co-culture, as a function of time in stationary phase (T, days) (**Figure 1A**). To this end, we inoculated stationary-phase cells at different time points into fresh medium and monitored the outgrowth at multiple times (t, hours) by measuring absorbance at 600nm (OD_{600}), bulk RFP signal (RFP), and bulk CFP signal (CFP), until the outgrowth co-cultures reached saturation.

We characterized the CLS of ten deletion strains that are known to show increased or decreased lifespan (Fabrizio and Longo, 2003; Wei et al., 2008; Alvers et al., 2009; Laschober et al., 2010; Burtner et al., 2011; Garay et al., 2014; Campos et al., 2018). Each $x\Delta_{RFP}$ was co-cultured until saturation with the WT_{CFP} reference in up to seven replicates in a single deep-well plate (see Materials and Methods). Competitive-aging cultures were monitored for ~15 days in stationary phase. As expected, outgrowth kinetics measured by OD_{600} showed a clear shift with time (days) in stationary phase; aging co-cultures gradually took a longer time in outgrowth (hours) to reach a given cell density (**Figure 1B**). This prevalent shift in growth kinetics reflects the loss of viability with culture age, as previously described (Murakami et al., 2008).

In terms of fluorescence-signal kinetics, we observed that the WT_{RFP} or WT_{CFP}

monocultures mostly recapitulated OD_{600} kinetics, suggesting that loss of viability can

also be measured by the shift of the fluorescence signal over the days (**Figure 1B**, red

and blue lines). WT_{RFP}+WT_{CFP} populations competed in co-culture showed similar shifts

in fluorescence signals, suggesting that loss of viability occurred at similar rates in both

wild-type populations, as expected (**Figure 1B**, gray lines). In contrast, some

$x\Delta_{RFP}$+WT_{CFP} co-cultures showed delayed or exacerbated shifts in fluorescence

kinetics. For instance, the $hap3\Delta_{RFP}$+WT_{CFP} co-culture showed a steep decrease in bulk

RFP signal as a function of days in stationary phase, along with a slight increase in CFP

signal (**Figure 1B**, orange lines), suggesting that loss of viability occurred at a rate

faster than the WT_{CFP} reference (short-lifespan phenotype). In contrast, the

$swr1\Delta_{RFP}$+WT_{CFP} co-culture showed a small increase in RFP signal along with steady

CFP signal (**Figure 1B,** green lines), suggesting slower loss of viability of $swr1\Delta$

compared to the WT (long-lifespan phenotype). These examples illustrate, from raw

data, that changes in bulk fluorescence in co-culture can be used to estimate loss of cell

population viability as a function of time in stationary phase.

A model to adjust relative survivorship in co-culture from changes in relative
fluorescence signal

To obtain a quantitative phenotypic value from our experimental measurements, we

developed a model that provides a relative survivorship parameter (S). Specifically, we

established a multiple linear regression analysis where each experimental

measurement in the outgrowth culture is modeled as:

$$ln\left(\frac{RFP}{CFP}\right)_{T_i,\,t_j} = A_w + S_w \cdot T_i + G_w \cdot t_j + C_{T_i,\,t_j}$$

Using a system of linear equations, we obtained the regression coefficients using all

outgrowth measurements in a 96-well plate. The contribution of relative survivorship (S)

along with that of the other three parameters for competing populations are illustrated in

Figure 2. The expected value, $ln\left(\frac{RFP}{CFP}\right)_{T_i,\,t_j}$, is the logarithmic quotient of the sizes of the

populations of $x\Delta_{RFP}$ and WT$_{CFP}$ from an outgrowth inoculated at day T_i in stationary

phase and measured after t_j hours in the outgrowth culture (Supplementary Note S1).

The parameter S is the difference in death rates (**Figure 2A**, linear regression, open

circles), while G is the difference in growth rates (linear regression, crosses in **Fig2A,B**).

The parameter A_w is the logarithmic proportion of the sizes of both populations at the

beginning of the experiment (**Figure 2B** , solid vertical lines at $T_1 = 0$). The term $C_{T_i,\,t_j}$

is the error in each measurement, which is mostly determined by the deviation from

zero change in WT$_{RFP}$+WT$_{CFP}$ reference competitions, and is similar to the deviation of

measurements in $x\Delta_{RFP}$+WT$_{CFP}$ competitions (**Figure 2B**). The rationale here is that

relative survivorship (S) and relative growth rate (S) are by definition equal to zero in

reference competitions (WT$_{RFP}$+WT$_{CFP}$), and therefore consistent changes in $ln\left(\frac{RFP_{ref}}{CFP_{ref}}\right)$

are likely due to systematic errors in the measurements. It must be noted that there are

usually survivorship and growth-rate differences between the two wild-type strains

(WT$_{RFP}$ and WT$_{CFP}$), which are taken into account while fitting the data. For instance, in

this experiment the WT$_{RFP}$ (and isogenic deletion strains) die faster than the WT$_{CFP}$

reference used, which is evident from the negative slope of data points in **Fig 2A**; this

lifespan difference is corrected by fitting data to the model in which the WT has S=0.

In the following sections, we show that competitive-aging experiments with multiple-

regression modeling provide reliable and replicable quantifications of relative

survivorship. We also show that this procedure is useful to identify CLS phenotypes and

to score their interactions with pharmacological factors.

Competitive-aging experiments provide replicable estimates of survivorship

To systematically assess the technical replicability of the competitive-aging method, we

measured the CLS of ten mutants and reference strains in 96-well plates with multiple

independent replicate wells. Specifically, we measured relative survivorship of up to

seven replicate samples of each one of the ten deletion mutants together with 31 wild-

type reference competitions in a 96-well plate (**Figure 3A**); the entire experiment was

carried two times: one experiment with three replicate plates and the second one with

two replicate plates. We validated our results with another large-scale method that

provides precise estimates of survivorship (**Figure 3B**). In particular, we measured CLS

of the same array of mutants and wild-type strains using an established high-throughput

method that is based in the changes of outgrowth kinetics of aging monocultures

(Murakami et al., 2008; Jung et al., 2015). Qualitative inspection showed that both

methods correctly scored those mutants with known CLS effect (Fabrizio et al., 2001;

Fabrizio and Longo, 2003). Specifically *rim15Δ* and *hap3Δ* had reduced survivorship,

while *ras2Δ* and *gln3Δ* showed increased lifespan compared to wild type.

To quantitatively contrast the replicability of both experimental approaches, we fitted

decay curves from the outgrowth kinetics experiments to an exponential model. We

then compared the difference of adjusted exponential death rates of wild-type and

mutant strains from monoculture aging to the *S* parameter obtained from competitive-

aging. We observed that both methods performed similarly when comparing the

technical variation within each of the plate replicates; there was no significant difference

in the typical standard error of the mean in outgrowth kinetics and competitive aging

(Supplementary Figure S1). Likewise, when looking at the correlation of quantitative

data resulting from independent replicates, we found equivalent correlation coefficients

between replicates of each of the two approaches, showing that competitive-aging is as

replicable as the outgrowth-kinetics method (Supplementary Figure S1). We note that

gln3Δ was not included in one of the experiments, and that the *ras2Δ* was atypically

noisy in one of the outgrowth-kinetics replicates (not shown); hence these strains were

conservatively excluded from this analysis to prevent overestimation of the intra- and

inter-batch variability of the outgrowth-kinetics approach. Together, these results

indicate that the competition-based method is highly and at least as replicable as the

established method, despite inherent variation of different experimental batches.

We looked closer into the distribution of effects of all deletion mutants in five replicates,

as determined by the two experimental approaches (**Figure 2C**). Competitive-aging

screening showed higher dynamic range but also higher variability. Deletion of *rim15Δ*

consistently showed strong short-lived phenotypes with both approaches, while milder

short-lived phenotypes of *hap3Δ*, *ste12Δ*, *msn4Δ*, and *msn2Δ* were also scored in most

of the plate replicates. Virtually all long-lived phenotypes were correctly scored by both

approaches, except for one replicate of *swr1Δ* by the OD-kinetics method. The short-

and long-lived phenotypes of *atg1Δ* and *rps16aΔ*, respectively, were not recapitulated

by any of the two methods, most likely due to differences in our experimental conditions

or strain background.

Having multiple reference samples (WT$_{RFP}$+WT$_{CFP}$) in each 96-well plate improves the

fit of the model, but necessarily reduces the number of samples in large-scale analyses,

which usually require running many batches in parallel. To maximize experimental

throughput without losing quantitative resolution, we used this experiment to estimate

the optimal number of reference samples per plate to estimate of S. To do so, we

quantified how the number of WT-reference samples (from 1 to 31 in this experiment)

used to fit S=0 affected the variation of mutant's S. We observed that both the average

standard deviation of S and the confidence intervals of the fit of S showed a steep

decline from one to five reference samples, after which variation kept decreasing with

diminishing returns (Supplementary Figure S2). Thus, we conclude that including 6 to

10 reference samples is enough to provide a robust description of relative-CLS

phenotypes in large-scale genetic analyses of mutants, environments, and their

interactions.

Finally, we used this data set to evaluate the influence of the growth rate of the mutants

in the context of our competitive-aging method and multiple-regression model, which

includes parameter G. Importantly, we observed that the model correctly identified

mutants with growth defects, as measured by independent culture outgrowths

(Supplementary Figure S3); hence, this parameter may enhance a more accurate

quantification of relative survivorship of deletion strains. To compare the estimation of

the effects on survivorship upon introduction of the G parameter, the data was fitted

assuming G=0 in all mutant-wild-type competitions; we observed that the difference in

the estimation of S when G was included was modest, but mostly explainable by G. As

expected, the relative survivorship of slow-growth mutants (*rim15Δ*, *gln3Δ*, and *ras2Δ*)

was underestimated when differences in growth rate were not taken into account

(Supplementary Figure S3). These results indicate that multiple-regression model

optimizes data analysis, especially in screens including mutants with impaired

exponential growth.

Competitive-quantification of CLS under different conditions successfully

describes gene-drug interactions

Lifespan is a complex trait determined by different cellular pathways, hundreds of

genes, and environmental variables[2,31]. A current challenge in the field is to understand

how different factors are integrated with one another to control cell survivorship. Our

competitive-aging method provides high-resolution and replicable data, which enables

an accurate quantitative description of CLS-phenotype interactions. As a proof of

principle experiment using our optimized experimental design and data analysis, we

screened for gene-environment interactions in an array of knockout mutants aged with

and without the lifespan-extending drug metformin.

We confirmed that the CLS of WT reference samples of yeast increased significantly

from a half life of 10.7 to 15.5 days when treated with 40mM metformin (**Figure 4A**;

$p<10^{-9}$, Wilcoxon rank sum test). Next, we used our competitive-aging approach to

measure the CLS of an array of 76 knockout strains aged with or without metformin

(random selection of strains from the yeast deletion collection). In both conditions, we

observed high quantitative correlation of the phenotypes between replicate plates in the

same experiment and between two independent experiments (Supplementary Figure

S4). We also confirmed that parameter G predicted actual growth rates and resulted in a

modest correction of S, as expected (Supplementary Figure S5).

A direct quantitative comparison of CLS phenotypes under both conditions is shown in

Figure 4B (see Supplementary Figure S6 for data rescaling). Most samples were found

to fall close to the diagonal; namely the phenotypic effect of the knockout relative to WT

was similar under both conditions. The phenotypes of 21 of the 76 knockouts (27%)

were significantly different when treated with metformin (Supplementary Figure S7; *t*-

test, *p*<0.05). Mutants with significantly-different phenotypes are potential gene-drug

interactions, especially those instances with large deviations from the expectation. In

many cases, we observed that metformin alleviated or even reverted the short-lifespan

effects of the gene deletion, with *pep1Δ*, *die2Δ*, and *alg3Δ* being the most extreme

instances. In only one mutant, *lsc2Δ*, the drug significantly aggravated the relative short-

lived gene-deletion phenotype. On the opposite scenario, the nominal long-lived

phenotype of certain mutants was rendered neutral or closer to neutral with the

metformin treatment (eg. *hsv2Δ*, *ubp13Δ*, and *ubp5Δ*).

We aimed to validate some of the potential gene-drug interactions identified in our

screen, focusing on eight mutants with large deviation from the expectation. First, we

measured the CLS of wild-type and gene-deletion strains with or without metformin,

using the outgrowth OD-kinetics approach (Supplementary Figure S7). A clear effect of

metformin treatment on the CLS phenotype was evident for *pep1Δ*, *die2Δ*, and *alg3Δ*; in

these cases the short-lifespan phenotype of the mutant was suppressed or alleviated by

metformin. We carried out live/dead staining of stationary-phase cells on these three

strains, which confirmed gene-drug interactions (**Figure 4C**). The identified interactions,

with specific quantitative information on the magnitude and sign of the effects, indicate a

functional association between protein glycosylation and metformin treatment, providing

novel information to understand the mechanisms of longevity by metformin in yeast.

Together, results in this section show that competitive-aging yields accurate and

replicable large-scale CLS data in a straight-forward manner to shed light on the

mechanisms of pharmacological interventions that extend lifespan.

DISCUSSION

Genetic analysis of the CLS of budding yeast has led to the genome-wide identification

of genes involved in aging; recent efforts have sought to describe interactions between

genetic and environmental modulators of the phenotype (Garay et al., 2014; Smith et

al., 2016; Campos et al., 2018; Jung et al., 2018). A previous report from our group

showed advantages of using a competitive-aging approach, in which fluorescently-

labeled strains in co-culture provide high resolution in parallel setups (Garay et al.,

2014). Competitive-aging has also allowed scoring gene-environment interactions at the

genomewide level, specifically interactions with dietary restriction (Campos et al., 2018).

Here, we have presented an optimized model to calculate the relative survivorship of

deletion strains, taking into account the possible confounding effects coming from

growth-rate differences and systematic batch effects. It is certain that major findings in

our previous reports would hold given the relatively mild contribution of the *G* and *C*

parameters in most samples (the experimental design of such studies precludes fitting data to the model herein presented). Nonetheless, the optimized procedure herein presented is a more exhaustive description of the actual competitive-aging setup, which could improve the scoring CLS phenotypes in specific cases, particularly strains with strong growth defects or experiments with strong batch effects. In addition, we directly compared the performance of a competitive-aging setup to an established approach (Murakami et al., 2008; Jung et al., 2015). Finally, using this enhanced method and data analysis in a proof-of-principle experiment, we were able to unravel significant gene-drug interactions in an array of gene-deletions strains subjected to the lifespan-extending drug metformin.

Early CLS genome-wide screens were based on large pools of gene deletions followed by molecular-barcode hybridization or sequencing (Fabrizio et al., 2010; Matecic et al., 2010; Gresham et al., 2011). These studies provided important insight into which genetic factors mediate stationary phase survival, such as autophagy, vacuolar protein sorting, and regulation of translation. However, the high rates of false positives—specially in the cases of long-lived phenotypes—and low overlap among the sets of genes from different studies (Smith et al., 2016) suggest that systematic errors in barcode detection or major experimental batch effects result in poor experimental replicability. On the other side of the spectrum, an ingenious outgrowth-kinetics approach of yeast monocultures increases the feasibility of percent-survivorship estimates, compared to the conventional colony-forming units method (Murakami et al., 2008); but throughput is still limited with this approach. To overcome this limitation, Jung and co-workers scaled-up this strategy using monocultures in multi-well plates, whereby

more strains can be tested in parallel (Jung et al., 2015, 2018). There is still the issue

that mild environmental variation can affect separate cultures differently, that could lead

to low reproducibility (Burtner et al., 2011); for instance, strains may reach stationary

phase at different times after inoculation. In this regards, competitive-aging provides a

direct phenotypic comparison and, arguably, more consistent results, given that the

mutant population of interest is aged with an internal reference strain under the exact

same microenvironment. Importantly, competitive-aging can also be carried out in multi-

well plates, enabling high-throughput experimental setups.

One of the inherent drawbacks of the competitive-aging method is that no absolute

death-rate information is provided, which can be easily solved thorough characterization

of the wild-type strain in monoculture, as we have shown. Likewise, the dynamic range

of this method depends on the actual death rate of the reference strain used. For

instance, quantitative phenotypic descriptions are limited if $S >> 0$; in other words only

semi-quantitative estimates are possible for extremely long-lived strains ($S \approx r_{wt}$). The

use of different reference strains, eg. specific gene deletions with known long-lifespan

phenotypes, may help to overcome this limitation. In addition, just as most population-

based CLS methods, competitive aging depends on cells being able to re-enter the cell

cycle, which can only be distinguished from actual death using outgrowth-independent

methods, such as live/dead staining.

We have illustrated the potential of competitive-aging screening by characterizing an

array of yeast deletion strains exposed to metformin. A number of lifespan-extending

pharmacological interventions are already being tested for age-related diseases in

humans, even when the mechanisms underlying their beneficial effects are frequently

not fully understood (Mullard, 2018; Zimmermann et al., 2018). Given that lifespan is a complex phenotype, identifying conserved gene-drug interactions could shed light on the modes of action and to pinpoint genetic modifiers of the drug's effects (Zimmermann et al., 2018). By screening an array of 76 gene deletions aged with metformin, we found a number of cases in which metformin buffers both short- or long-lifespan mutant phenotypes. The interacting genes suggest a role of protein glycosylation and protein homeostasis, which is in line with previous evidence showing that metformin alters glycation, protein transport, and protein degradation in yeast (Borklu-Yucel et al., 2015; Kazi et al., 2017). Our results also uncovered interactions between metformin and genes involved in mitochondrial function (*erp6Δ*, *lsc2Δ, ema35Δ*, and *ylh47Δ*), which is a known player in the cellular response to metformin (Borklu-Yucel et al., 2015; Stynen et al., 2018). It remains to be addressed how these proteins are specifically related to the known response involving the mitochondrial electron transport chain and homeostasis of copper and iron.

Competitive-aging can readily be adapted to screen double mutants at large scale and to score genetic interactions underlying CLS phenotypes. Genetic interactions (epistasis) are a powerful tool to describe the architecture of phenotypes and the functional relationships of different genetic pathways (Segrè et al., 2005; Collins et al., 2007; Onge et al., 2007; Phillips, 2008; Kuzmin et al., 2018). While epistasis-network analyses in yeast have granted deep knowledge of the genetic landscape of mitotically-active, proliferating cells, less is known about how genetic interactions shape the genetic architecture of post-mitotic survivorship. Large-scale genetic analysis of double-mutants aged in competition with their single-mutant references is an attractive

439 experimental setup to identify interactions among different genes and pathways

440 underlying CLS in yeast. Our competitive-aging screening method and quantitative

441 analysis provide a powerful systematic tool to shed light on the complex genetic,

442 environmental, and pharmacological wiring of aging cells.

Acknowledgements

We are grateful to Erika Cruz-Bonilla, Nelly Selem, and Judith Ulloa for critical reading

of the manuscript. This work was funded by the Consejo Nacional de Ciencia y

Tecnología de México (CONACYT grants CB-2015/164889 and PN-2016/2370). J.A.A.-

R. received a doctoral fellowship from CONACYT (#264529). The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript. This manuscript has been released as a pre-print at BioRxiv (Avelar-Rivas

et al., 2019).

Author Contributions

452 Conceived and designed the study: JAA-R, AD. Performed the experiments: JAA-R,

453 MM-F, AJ-R, SEC. Developed and implemented the model: JAA-R, EG, NS. Analyzed

454 the data: JAA-R, SEC, AD. Wrote the paper: JAA-R, AD. Acquired funding: AD. All

455 authors read and approved the final version of the submitted manuscript.

Conflict of interest

457 The authors declare that they have no competing interests.

FIGURE LEGENDS

Figure 1. Competitive-aging of fluorescently labeled budding yeast strains. *A*, Schematic of the experimental setup. A co-culture of fluorescence-tagged gene deletion ($x\Delta_{RFP}$) and wild-type reference (WT_{CFP}) strains in stationary phase is sampled regularly at days T. Outgrowth cultures in fresh medium are monitored at t hours with simultaneous measurement of absorbance at 600nm (OD_{600}), and raw *RFP* and *CFP* signals. Change in fluorescent-signal ratio in the outgrowth cultures is used to estimate the relative number of viable cells after T days in stationary phase. Possible differences in growth rate (G) are taken into account (see multiple linear modeling). ***B*,** Raw data of OD_{600}, *RFPraw*, and *CFPraw* signal obtained from outgrowth-culture kinetics started after T days in stationary phase. Selected examples are shown in five columns, from left to right: Only WT_{RFP} (red lines), only WT_{CFP} (blue lines), both WT populations in co-culture (WT_{RFP} +WT_{CFP}, gray lines), a short-lived deletion strain in co-culture with the WT (*hap3*Δ_{RFP} +WT_{CFP}, orange lines), and a long-lived strain with the WT (*swr1*Δ_{RFP} +WT_{CFP}, green lines). Samples are measured throughout the outgrowth culture after inoculation, but analysis only considers data points at exponential growth (circles in grayscale).

Figure 2. Multiple linear modeling fits relative survivorship and relative growth rate from competitive-aging cultures. Examples of a reference competition (gray) and two deletion strains in co-culture with the WT_{CFP} (*hap3*Δ_{RFP}, orange; *swr1*Δ_{RFP}, green). In both panels, dots indicate the experimental *ln*(*RFP/CFP*) measurements. ***A*,** Fitted data is shown in the T timescale (days). Open circles are the data fitted at T_i and error bars at those time points are the 95% CI of the fit; their linear regression has slope S (the relative survivorship, our parameter of interest). For each well, parameter A indicates the deviation from ln(*RFP/CFP*)=0 at T=0, as indicated by the vertical line at T=0. ***B*,** Fitted data is shown in the t timescale (hours); nine panels correspond to outgrowths at different age times, *T*. Crosses shown directly below or above experimental data are the fitted data $ln\left(\frac{RFP}{CFP}\right)_{T_i, t_j}$ at the time of each measurement; their linear regression has slope G (the relative growth rate). All samples (wells) in the same

96-well plate measurement have the same $C_{T,t}$ (systematic batch error, gray vertical lines).

Figure 3. The competitive-aging method is accurate and replicable. *A*, Relative survivorship estimated by competitive aging. The modeled *ln*(*RFP/CFP*) is shown for the WT and ten mutant strains at each T_i point, averaged for at least six technical replicates along with the CI 95% (shaded area). ***B*,** Percent of surviving cells in monoculture over time measured by the shift in outgrowth kinetics; the mean and CI 95% are shown. ***C*,** Average relative survivorship of three to five experimental replicates is shown, expressed either as the difference of individual death rates for monoculture outgrowth kinetics ($r_{wt} - r_x$, black) or as survivorship coefficients for competitive aging (S, red). Each data point is the average of at least six technical replicates of monocultures or competitions in an independent experimental replicate. Horizontal lines are the mean of each data series. Stacked bars below the plot indicate the number of replicate plates per sample; solid fraction indicates the number of significant samples compared with the WT distribution (*p*<0.01, *t*-test; each experimental batch has 31 reference and at least six deletion samples).

Figure 4. Identification of gene-drug interactions by competitive-aging screening. *A*, Half life of WT-reference samples with (*n*=39) or without metformin (*n*=40). ***B*,** Scatter plot comparing the CLS phenotypes of 76 gene-deletion strains with or without metformin; the average rescaled relative survivorship rS shown of four replicates is shown. Color scale indicates the *p*-value of paired t-tests between the rS in SC and SC with metformin. ***C*,** Gene-drug interactions confirmed by live/dead staining. Survival curves of wild-type (light colors) and gene-deletion (dark colors) strains in nominal SC medium (green) or SC with metformin (orange); 95% CI were calculated from at least three replicates (shaded area). The interaction between metformin and all three gene-deletions was scored significant by two-way ANOVA tests of the death rates (******p< 0.01).

SUPPLEMENTARY ITEMS (ZIP)

Note S1. Development and implementation of a linear model to calculate relative survival of mutant strains in competitive-aging cultures (PDF)

Table S1. Recipes for media used in this study (PDF)

Figure S1. Comparing replicability of CLS measurements from two methods (PDF)

Figure S2. Variation in S decreases with the number of reference samples used (PDF)

Figure S3. The parameter G corrects effects of differential growth rates that otherwise affect relative survivorship, S (PDF)

Figure S4. Relative survivorship (S) is well replicated between plates within the same experimental batch, as well as between different batches (PDF)

Figure S5. The parameter G corrects effects of differential growth rates that otherwise affect relative survivorship, S (PDF)

Figure S6. Relative survivorship (S) data rescaling (PDF)

Figure S7. Candidate gene-drug interactions and validation (PDF)

Data S1. Array of mutants used for gene-drug interactions, with their relative survivorship, rescaled survivorship, and relative growth rates (XLSX)

REFERENCES

Alvers, A. L., Fishwick, L. K., Wood, M. S., Hu, D., Chung, H. S., Dunn Jr, W. A., et al. (2009). Autophagy and amino acid homeostasis are required for chronological longevity in *Saccharomyces cerevisiae*. *Aging Cell* 8, 353–369. doi:10.1111/j.1474-9726.2009.00469.x.

Avelar-Rivas, J. A., Munguía-Figueroa, M., Juárez-Reyes, A., Garay, E., Campos, S. E., Shoresh, N., and DeLuna, A. (2019). A competitive-aging method for quantitative genetic analysis of the chronological lifespan of Saccharomyces cerevisiae. *bioRxiv*, 696682. doi: 10.1101/696682.

Barré, B. P., Hallin, J., Yue, J.-X., Persson, K., Mikhalev, E., Irizar, A., et al. (2019). Intragenic repeat expansions control yeast chronological aging. *bioRxiv*, 653006. doi:10.1101/653006.

Borklu-Yucel, E., Eraslan, S., and Ulgen, K. O. (2015). Transcriptional remodeling in response to transfer upon carbon-limited or metformin-supplemented media in S. cerevisiae and its effect on chronological life span. *Appl. Microbiol. Biotechnol.* 99, 6775–6789. doi:10.1007/s00253-015-6728-5.

Burtner, C. R., Murakami, C. J., Kennedy, B. K., and Kaeberlein, M. (2009). A molecular mechanism of chronological aging in yeast. *Cell Cycle* 8, 1256–1270. doi:10.4161/cc.8.8.8287.

Burtner, C. R., Murakami, C. J., Olsen, B., Kennedy, B. K., and Kaeberlein, M. (2011). A genomic analysis of chronological longevity factors in budding yeast. *Cell Cycle* 10, 1385–1396. doi:10.4161/cc.10.9.15464.

Campos, S. E., Avelar-Rivas, J. A., Garay, E., Juárez-Reyes, A., and DeLuna, A. (2018). Genomewide mechanisms of chronological longevity by dietary restriction in budding yeast. *Aging Cell* 17, e12749. doi:10.1111/acel.12749.

Campos, S. E., and DeLuna, A. (2019). Functional genomics of dietary restriction and longevity in yeast. *Mech. Ageing Dev.* 179, 36–43. doi:10.1016/J.MAD.2019.02.003.

Collins, S. R., Miller, K. M., Maas, N. L., Roguev, A., Fillingham, J., Chu, C. S., et al. (2007). Functional dissection of protein complexes involved in yeast chromosome biology using a genetic interaction map. *Nature* 446, 806–810. doi:10.1038/nature05649.

Eisenberg, T., Knauer, H., Schauer, A., Büttner, S., Ruckenstuhl, C., Carmona-Gutierrez, D., et al. (2009). Induction of autophagy by spermidine promotes longevity. *Nat. Cell Biol.* 11, 1305–1314. doi:10.1038/ncb1975.

Fabrizio, P., Hoon, S., Shamalnasab, M., Galbani, A., Wei, M., Giaever, G., et al. (2010). Genome-Wide Screen in Saccharomyces cerevisiae Identifies Vacuolar Protein Sorting, Autophagy, Biosynthetic, and tRNA Methylation Genes Involved in Life Span Regulation. *PLoS Genet.* 6, e1001024. doi:10.1371/journal.pgen.1001024.

Fabrizio, P., and Longo, V. D. (2003). The chronological life span of Saccharomyces cerevisiae. *Aging Cell* 2, 73–81. doi:10.1046/j.1474-9728.2003.00033.x.

Fabrizio, P., Pozza, F., Pletcher, S. D., Gendron, C. M., and Longo, V. D. (2001). Regulation of longevity and stress resistance by Sch9 in yeast. *Science (80-.).* 292, 288–290. doi:10.1126/science.1059497.

Garay, E., Campos, S. E., González de la Cruz, J., Gaspar, A. P., Jinich, A., and DeLuna, A. (2014). High-Resolution Profiling of Stationary-Phase Survival Reveals Yeast Longevity Factors and Their Genetic Interactions. *PLoS Genet.* 10, e1004168. doi:10.1371/journal.pgen.1004168.

Gems, D., and Partridge, L. (2013). Genetics of Longevity in Model Organisms: Debates and Paradigm Shifts. *Annu. Rev. Physiol.* 75, 621–644. doi:10.1146/annurev-physiol-030212-183712.

Gresham, D., Boer, V. M., Caudy, A., Ziv, N., Brandt, N. J., Storey, J. D., et al. (2011). System-level analysis of genes and functions affecting survival during nutrient starvation in Saccharomyces cerevisiae. *Genetics* 187, 299–317. doi:10.1534/genetics.110.120766.

Hu, J., Wei, M., Mirisola, M. G., and Longo, V. D. (2013). "Assessing Chronological Aging in Saccharomyces cerevisiae," in (Humana Press, Totowa, NJ), 463–472. doi:10.1007/978-1-62703-239-1_30.

Jung, P. P., Christian, N., Kay, D. P., Skupin, A., and Linster, C. L. (2015). Protocols and Programs for High-Throughput Growth and Aging Phenotyping in Yeast. *PLoS One* 10, e0119807. doi:10.1371/journal.pone.0119807.

Jung, P. P., Zhang, Z., Paczia, N., Jaeger, C., Ignac, T., May, P., et al. (2018). Natural variation of chronological aging in the Saccharomyces cerevisiae species reveals diet-dependent mechanisms of life span control. *npj Aging Mech. Dis.* 4, 3. doi:10.1038/s41514-018-0022-6.

Kazi, R. S., Banarjee, R. M., Deshmukh, A. B., Patil, G. V., Jagadeeshaprasad, M. G., and Kulkarni, M. J. (2017). Glycation inhibitors extend yeast chronological lifespan by reducing advanced glycation end products and by back regulation of proteins involved in mitochondrial respiration. *J. Proteomics* 156, 104–112. doi:10.1016/J.JPROT.2017.01.015.

Kirkwood, T. B. L. (2008). A systematic look at an old problem. *Nature* 451, 644–647. doi:10.1038/451644a.

Kuzmin, E., VanderSluis, B., Wang, W., Tan, G., Deshpande, R., Chen, Y., et al. (2018). Systematic analysis of complex genetic interactions. *Science* 360, eaao1729. doi:10.1126/science.aao1729.

Laschober, G. T., Ruli, D., Hofer, E., Muck, C., Carmona-Gutierrez, D., Ring, J., et al. (2010). Identification of evolutionarily conserved genetic regulators of cellular aging. *Aging Cell* 9, 1084–1097. doi:10.1111/j.1474-9726.2010.00637.x.

Longo, V. D., and Fabrizio, P. (2012). Chronological aging in Saccharomyces cerevisiae. *Subcell. Biochem.* 57, 101–21. doi:10.1007/978-94-007-2561-4_5.

Longo, V. D., Shadel, G. S., Kaeberlein, M., and Kennedy, B. (2012). Replicative and Chronological Aging in Saccharomyces cerevisiae. *Cell Metab.* 16, 18–31. doi:10.1016/J.CMET.2012.06.002.

López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The

Hallmarks of Aging. *Cell* 153, 1194–1217. doi:10.1016/J.CELL.2013.05.039.

Matecic, M., Smith, D. L., Pan, X., Maqani, N., Bekiranov, S., Boeke, J. D., et al. (2010). A Microarray-Based Genetic Screen for Yeast Chronological Aging Factors. *PLoS Genet.* 6, e1000921. doi:10.1371/journal.pgen.1000921.

Mullard, A. (2018). Anti-ageing pipeline starts to mature. *Nat. Rev. Drug Discov.* 17, 609–612. doi:10.1038/nrd.2018.134.

Murakami, C. J., Burtner, C. R., Kennedy, B. K., and Kaeberlein, M. (2008). A Method for High-Throughput Quantitative Analysis of Yeast Chronological Life Span. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* 63, 113–121. doi:10.1093/gerona/63.2.113.

Onge, R. P. S., Mani, R., Oh, J., Proctor, M., Fung, E., Davis, R. W., et al. (2007). Systematic pathway analysis using high-resolution fitness profiling of combinatorial gene deletions. *Nat. Genet.* 39, 199–206. doi:10.1038/ng1948.

Phillips, P. C. (2008). Epistasis — the essential role of gene interactions in the structure and evolution of genetic systems. *Nat. Rev. Genet.* 9, 855–867. doi:10.1038/nrg2452.

Powers, R. W., Kaeberlein, M., Caldwell, S. D., Kennedy, B. K., Fields, S., and Fields, S. (2006). Extension of chronological life span in yeast by decreased TOR pathway signaling. *Genes Dev.* 20, 174–84. doi:10.1101/gad.1381406.

Santos, J., Leitão-Correia, F., Sousa, M. J., and Leão, C. (2015). Ammonium is a key determinant on the dietary restriction of yeast chronological aging in culture medium. *Oncotarget* 6, 6511–23. doi:10.18632/oncotarget.2989.

Segrè, D., DeLuna, A., Church, G. M., and Kishony, R. (2005). Modular epistasis in yeast metabolism. *Nat. Genet.* 37, 77–83. doi:10.1038/ng1489.

Sheff, M. A., and Thorn, K. S. (2004). Optimized cassettes for fluorescent protein tagging inSaccharomyces cerevisiae. *Yeast* 21, 661–670. doi:10.1002/yea.1130.

Smith, D. L., Maharrey, C. H., Carey, C. R., White, R. A., Hartman, J. L., Hartman, J. L., et al. (2016). Gene-nutrient interaction markedly influences yeast chronological lifespan. *Exp. Gerontol.* 86, 113–123. doi:10.1016/j.exger.2016.04.012.

Stynen, B., Abd-Rabbo, D., Kowarzyk, J., Miller-Fleming, L., Aulakh, S. K., Garneau, P., et al. (2018). Changes of Cell Biochemical States Are Revealed in Protein Homomeric Complex Dynamics. *Cell* 175, 1418-1429.e9. doi:10.1016/J.CELL.2018.09.050.

Wei, M., Fabrizio, P., Hu, J., Ge, H., Cheng, C., Li, L., et al. (2008). Life span extension by calorie restriction depends on Rim15 and transcription factors downstream of Ras/PKA, Tor, and Sch9. *PLoS Genet.* 4, 0139–0149. doi:10.1371/journal.pgen.0040013.

Zimmermann, A., Hofer, S., Pendl, T., Kainz, K., Madeo, F., and Carmona-Gutierrez, D. (2018). Yeast as a tool to identify anti-aging compounds. *FEMS Yeast Res.* 18. doi:10.1093/femsyr/foy020.

A

$x\Delta_{RFP}$ + WT$_{CFP}$

Co-culture in stationary phase

T, days

T_1 T_2 $T_{...}$ T_n

t_m OD_{600}
$t_{...}$ RFP_{raw}
t_2 G
t_1 CFP_{raw}

t, hours

B

T, days
0   14

— WT$_{RFP}$ — WT$_{CFP}$ — WT$_{RFP}$+WT$_{CFP}$ — $hap3\Delta_{RFP}$+WT$_{CFP}$ — $swr1\Delta_{RFP}$+WT$_{CFP}$

OD_{600}

CFP_{raw}

RFP_{raw}

t, hours

Supplementary Note 1. Development and implementation of a linear model to calculate relative survival of mutant strains in competitive-aging cultures

To obtain relative survivorship of competitive-aging cultures we modeled WT and mutant populations with initial sizes $N_{wt\,T0}$ and $N_{x\,T0}$ and declining trough time in stationary phase at constant exponential rates r_{wt} and r_x. After T_i days, the sizes of the remaining viable populations are $N_{wt\,T_i} = N_{wtT_0} \cdot e^{-r_{wt}\cdot T_i}$ and $N_{x\,T_i} = N_{x\,T_0} \cdot e^{-r_x T_i}$. In every outgrowth sampling, we inoculated small volumes of this aged co-culture onto fresh medium, in which both populations resume cell division and grow exponentially at rates g_{wt} and g_x. At any outgrowth, the population sizes after t_j hours are $N_{wt\,Tt_j} = N_{wt\,T} \cdot e^{g_{wt}\cdot t_j}$ and $N_{x\,Tt_j} = N_{x\,T} \cdot e^{g_x\cdot t_j}$. Given that both strains are aged and grown in co-culture, the relationship between the sizes of their populations at any time, t_j, during the outgrowth T_i is:

$$ln\,(\frac{N_x}{N_{wt}})\;_{T_i,\,t_j} = ln\,(\frac{N_x}{N_{wt}})\;_{T_0} + (r_{wt} - r_x)T_i + (g_x - g_{wt})(t_j)$$

As shown in Figure S1, the measurements of *CFP* and *RFP* signals are proportional to $N_{wt\,Tt}$ and $N_{x\,Tt}$, therefore, the experimental measurements of each well, *w,* are equal to the linear equation:

$$ln\,(\frac{RFP}{CFP})\;_{T_i,\,t_j} = A_w + S_w \cdot T_i + G_w \cdot t_j + C_{T_i,\,t_j}$$

We run a multiple linear regression where we fit the data obtained from measuring bulk RFP signal (*RFP*), and bulk CFP signal (*CFP*) in the outgrowth cultures in fresh medium sampled by inoculating from aging stationary phase co-cultures. The first step is to subtract auto-fluorescence RFP_{raw} from WT$_{CFP}$ as a function of the OD_{600} from all RFP_{raw} measurements; the same is done for CFP_{raw} from WT$_{RFP}$. We keep outgrowth measurements for days in which we observe an increase in both RFP$_{raw}$ and CFP_{raw}. This leaves us with the *RFP* and *CFP* quantities that should be a function of the death and growth rates (r and g). For each measured well, *w*, we

expressed such relationship using the model $ln\left(\frac{RFP}{CFP}\right)_{T_i,t_j} = A_w + S_w \cdot T_i + G_w \cdot t_j + C_{T_i,t_j}$ in which A_w reflects the proportion $ln\left(\frac{RFP}{CFP}\right)_{T_0,t_0}$ at the beginning of the experiment, this number will depend on the amount of Δx_{RFP} and WT_{CFP} strains that are mixed together when the aging co-cultures are set up. The relative survivorship is modeled in $S_w = (r_{wt} - r_x)$, which is expressed in terms of the number of the time T (in days) that the co-culture has been in stationary phase. Meanwhile the difference in growth rates of both strains is modeled in $G_w = (g_x - g_{wt})$, which is expressed in terms of the time t (in hours) within every outgrowth. Finally, the model takes into account the batch error of each plate read, as C_{T_i,t_j}.

To fit the data, we generate a system of linear equations. For well w_1, there are as many equations as the total number of measurements of that well across the entire experiment. For instance, day number one, T_1, has the following equations:

Eq1.1.1 $\quad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1,t_1} = A_{w_1} + S_{w_1} \cdot T_1 + G_{w_1} \cdot t_{1.1} + C_{T_1,t_1}$

Eq1.2.1 $\quad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1,t_2} = A_{w_1} + S_{w_1} \cdot T_1 + G_{w_1} \cdot t_{1.2} + C_{T_1,t_2}$

Eq1.3.1 $\quad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1,t_3} = A_{w_1} + S_{w_1} \cdot T_1 + G_{w_1} \cdot t_{1.3} + C_{T_1,t_3}$

Eq1.m.1 $\quad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1,t_m} = A_{w_1} + S_{w_1} \cdot T_1 + G_{w_1} \cdot t_{1.m} + C_{T_1,t_m}$

In these equations, the term $ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1,t_m}$ is the empirical determination of the logarithmic quotient of both fluorescent signals at each measurement of the outgrowth in exponential phase. A_{w_1} is the same variable for all equations of w_1, likewise for $S_{w_1} \cdot T_1$ because the relative death rate of the well (S_{w_1}) is the same and is multiplied by the same number of days of the experiment, T_1. By definition, the first day of measurement is considered time $T_1 = 0$ for all plates and wells. One of the two terms that change between these equations is, $G_{w_1} \cdot t_{1.m}$. Although the

relative growth rate is the same for each well throughout the entire experiment, the time at which the measurement is done changes and that is reflected in the different coefficients that these equations will get for those terms. For example, if measurements are done every 1.5 hours and the exponential phase starts at the third measurement, coefficients would be $t_{1.1} = 3$; $t_{1.2} = 4.5$; $t_{1.3} = 6$. Finally, the other term that changes in these equations is C_{T_1, t_m} which reflects that there are batch effects influencing measurements; such effects appear any time a plate is read, and so it has similar effects in all of the wells within the same plate. For that reason C_{T_1, t_m} is different at each measurement of a well, but is shared in all wells from the same plate read. Therefore, the measurements of different wells in one plate with k wells lead to equations:

$$Eq1.1.1 \qquad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_1, t_1} = A_{w_1} + S_{w_1} \cdot T_1 + G_{w_1} \cdot t_{1.1} + C_{T_1, t_1}$$

$$Eq1.1.2 \qquad ln\left(\frac{RFP_{w_2}}{CFP_{w_2}}\right)_{T_1, t_1} = A_{w_2} + S_{w_2} \cdot T_1 + G_{w_2} \cdot t_{1.1} + C_{T_1, t_1}$$

$$Eq1.1.3 \qquad ln\left(\frac{RFP_{w_3}}{CFP_{w_3}}\right)_{T_1, t_1} = A_{w_3} + S_{w_3} \cdot T_1 + G_{w_3} \cdot t_{1.1} + C_{T_1, t_1}$$

$$Eq1.1.k \qquad ln\left(\frac{RFP_{w_k}}{CFP_{w_k}}\right)_{T_1, t_1} = A_{w_k} + S_{w_k} \cdot T_1 + G_{w_k} \cdot t_{1.1} + C_{T_1, t_1}$$

The same wells measured the second day will have the equations:

$$Eq2.1.1 \qquad ln\left(\frac{RFP_{w_1}}{CFP_{w_1}}\right)_{T_2, t_1} = A_{w_1} + S_{w_1} \cdot T_2 + G_{w_1} \cdot t_{2.1} + C_{T_2, t_1}$$

$$Eq2.1.2 \qquad ln\left(\frac{RFP_{w_2}}{CFP_{w_2}}\right)_{T_2, t_1} = A_{w_2} + S_{w_2} \cdot T_2 + G_{w_2} \cdot t_{2.1} + C_{T_2, t_1}$$

$$Eq2.1.1 \qquad ln\left(\frac{RFP_{w_3}}{CFP_{w_3}}\right)_{T_2, t_1} = A_{w_3} + S_{w_3} \cdot T_2 + G_{w_3} \cdot t_{2.1} + C_{T_2, t_1}$$

$$Eq2.1.k \qquad ln\left(\frac{RFP_{w_k}}{CFP_{w_k}}\right)_{T_2, t_1} = A_{w_k} + S_{w_k} \cdot T_2 + G_{w_k} \cdot t_{2.1} + C_{T_2, t_1}$$

If the second outgrowth occurs after one day then $T_2 = 1$. It is worth pointing out that this $t_{2.1}$ is different from $t_{1.1}$ because the outgrowths of each day are not measured at identical points in time. For example, it could be that $t_{2.1} = 3.6$ and $t_{2.2} = 4.1$.

We verified the quality of the data and set some restrictions. For instance, in every valid measurement both fluorescent signals must be above the maximum auto-fluorescence detection, as indicated by wells with monocultures of WT_{CFP} or WT_{RFP} strain with auto-fluorescence of the opposite signal. This step removes all monoculture wells from the system of equations.

Our model fitting has the assumption that $WT_{CFP}+WT_{RFP}$ competitions have $S_x = 0$ and $G_x = 0$, so deviations from such values in wells ref, containing such mixture of strains depend on the well specific variable $A_{w_{ref.i}}$ or to the plate variables C_{T_n,t_m}. To accomplish that, we eliminate S and G from the reference-well equations, so every measurement from those wells will feed the system of equations as follows:

$$Eq1.r.1 \qquad ln\left(\frac{RFP_{w_{ref.i}}}{CFP_{ref.i}}\right)_{T_1,t_1} = A_{w_{ref.i}} + C_{T_1,t_1}$$

$$Eq1.r.2 \qquad ln\left(\frac{RFP_{ref.i}}{CFP_{w_{ref.i}}}\right)_{T_1,t_2} = A_{w_{ref.i}} + C_{T_1,t_2}$$

$$Eq2.r.1 \qquad ln\left(\frac{RFP_{w_{ref.i}}}{CFP_{w_{ref.i}}}\right)_{T_2,t_1} = A_{w_{ref.i}} + C_{T_2,t_1}$$

To find an approximation to the solution of the entire set of equations, we built a predictor matrix that contains all the valid measurements of the plate and the logarithmic ratio of the actual experimental measurements as the responses vector. In this example, we show the matrix for wells w_1,w_2, w_{ref} and w_k outgrown at days T_1, T_2 and T_n with 3 measurements per outgrowth $t_{n.1}$, $t_{n.2}$ and $t_{n.m}$:

Aw_1	Sw_1	Gw_1	Aw_2	Sw_2	Gw_2	Aw_{ref}	Aw_k	Sw_k	Gw_k	CT_1t_1	CT_1t_2	CT_1t_m	CT_2t_1	CT_2t_2	CT_2t_m	CT_nt_1	CT_nt_2	CT_nt_m
1	T_1	$t_{1.1}$	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
1	T_1	$t_{1.2}$	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0
1	T_1	$t_{1.m}$	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0
1	T_2	$t_{2.1}$	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
1	T_2	$t_{2.2}$	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0
1	T_2	$t_{2.m}$	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0
1	T_n	$t_{n.1}$	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
1	T_n	$t_{n.2}$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
1	T_n	$t_{n.m}$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	1	T_1	$t_{1.1}$	0	0	0	0	1	0	0	0	0	0	0	0	0
0	0	0	1	T_1	$t_{1.2}$	0	0	0	0	0	1	0	0	0	0	0	0	0
0	0	0	1	T_1	$t_{1.m}$	0	0	0	0	0	0	1	0	0	0	0	0	0
0	0	0	1	T_2	$t_{2.1}$	0	0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	1	T_2	$t_{2.2}$	0	0	0	0	0	0	0	0	1	0	0	0	0
0	0	0	1	T_2	$t_{2.m}$	0	0	0	0	0	0	0	0	0	1	0	0	0
0	0	0	1	T_n	$t_{n.1}$	0	0	0	0	0	0	0	0	0	0	1	0	0
0	0	0	1	T_n	$t_{n.2}$	0	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	1	T_n	$t_{n.m}$	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	0	0	0	1	T_1	$t_{1.1}$	1	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	T_1	$t_{1.2}$	0	1	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	T_1	$t_{1.m}$	0	0	1	0	0	0	0	0	0
0	0	0	0	0	0	0	1	T_2	$t_{2.1}$	0	0	0	1	0	0	0	0	0
0	0	0	0	0	0	0	1	T_2	$t_{2.2}$	0	0	0	0	1	0	0	0	0
0	0	0	0	0	0	0	1	T_2	$t_{2.m}$	0	0	0	0	0	1	0	0	0
0	0	0	0	0	0	0	1	T_n	$t_{n.1}$	0	0	0	0	0	0	1	0	0
0	0	0	0	0	0	0	1	T_n	$t_{n.2}$	0	0	0	0	0	0	0	1	0
0	0	0	0	0	0	0	1	T_n	$t_{n.m}$	0	0	0	0	0	0	0	0	1

If there are k wells in a plate, the plate was measured $m * n$ times then there are $k * m * n$ rows in the matrix and there are $3k + m * n - 2Nw_{ref}$ variables (or columns)

to fit (Nw_{ref} is the number of wells that contain reference co-cultures); hence, for the previous matrix there are 36 rows and 19 predictor variables.

We applied a multiple linear regression to approximate the solution. The Matlab function *regress* was used, as it provides useful statistics about the fit such as the 95% confidence interval for the coefficient estimates.

Scripts and documentation to run this model are available in GitHub: https://abrahamavelar.github.io/LinearModelCLS/

Table S1. Media recipes

Aging medium (SC)

Reagents
Difco™ Yeast Nitrogen Base w/o Amino Acids (ref 291940)
D-(+)- Glucose (Sigma G2870)
Yeast Synthetic Drop-out Medium Supplements **without uracil** (Sigma Y1501)
Uracil (Sigma, U0750)

Recipe
1. Before mixing any of the components 1.79 g of Uracil were added to the commercial Synthetic Drop-out mix (20 g) to obtain a complete amino acid mix
2. For 1 L of SC medium mix together:
3. Add the components to 500 mL of deionized water and dissolve
4. Add deionized water to make 1 L of medium.
5. Filter sterilize
Note: do not buffer pH

YNB low fluorescence medium (YNB-lf)

Recipe
1. For 1 L of YNB-lf medium, mix together:

-5 g	$(NH_4)_2SO_4$
-1 g	KH_2PO_4
- 0.5 g	$MgSO_4$
- 0.1 g	NaCl
- 0.1 g	$CaCl_2$
- 20 g	D-(+)- Glucose (Sigma G2870)
- 0.79 g	CSM (complete supplement mixture, MP)

2. Add the components to 500 mL of deionized water and dissolve thoroughly
3. Add 1 mL of trace element stock*
4. Add 1 mL of vitamin stock**
5. Add deionized water to make 1 L of medium
6. Filter sterilize

*Trace-element stock
- Mix the indicated amount of components together:

50mg	H_3BO_4
4mg	$CuSO_4$
10mg	KI
20mg	$FeCl_3$
40mg	$MnSO_4$
20mg	Na_2MoO_4
40mg	$ZnSO_4$

- Bring to 100 mL with deionized water

(...)

** Vitamin stock
- Mix the indicated amount of components together:

0.2mg	Biotin
40mg	Calcium pantothenate
200mg	Inositol
40mg	Niacin
20mg	Para-amino benzoic acid
40mg	Pyroxidine HCl
40mg	Thiamine HCl

- Bring to 100 mL with deionized water

Supplementary Figure S1. Comparing replicability of CLS measurements from two methods. **A,** Paired Pearson's *r* correlation coefficient among the relative survivorship, *S*, from three (experiment #1, plates A, B, and C) or two replicates (experiment #2, plates C and D). Competitive-aging correlations are shown above, while OD-kinetics in monoculture are shown below the diagonal. **B,** Cumulative S.E.M. (intra-experiment) of all mutant´s *S*. Data of the five experimental batches are shown in each series (*n*=47 and *n*=48 for monoculture outgrowth kinetics and competitive aging, respectively). The median S.E.M. of *S* from both methods are statistically indistinguishable (*p*=0.154, Wilcoxon rank sum test).

Supplementary Figure S2. Variation in S decreases with the number of reference samples used. Thirty-one $WT_{RFP}+WT_{CFP}$ reference competitions were used in each plate. To estimate the effects of increasing number of reference samples, the model was fed with different numbers of reference wells and the reminder control samples were treated as additional mutant samples. For each given number of reference samples, 100 permutations were randomly selected and the selection coefficient (S) of all mutant strains was calculated accordingly. **A**, The variation using different permutations was quantified as the standard deviation of each mutant sample; the average standard deviation of S of all mutant samples is shown as a function of the number of reference samples in the model. **B,** The fit to the model was calculated along with the 95% CIs; the difference between the upper and lower CI bounds is plotted against the number of reference samples added to the model.

Supplementary Figure S3. The parameter G corrects effects of differential growth rates that otherwise affect relative survivorship, S. **A**, The actual growth rate at exponential phase of gene-deletion strains measured as average increase in OD_{600} with time (n=6-7), compared to the modeled parameter G. **B**, Scatter plot showing the average S value from 3-5 replicates with parameter G is included in the multiple linear regression (horizontal axis) and when the regression is run without G (S_{woG}, vertical axis). **C,** The mild difference between calculations of S showed in panel B is mostly explained by changes in the parameter G.

Supplementary Figure S4. The relative survivorship, S, is well replicated within the same experiment (replicates 'A' and 'B', or 'C' and 'D') and between different experiments (#1 and #2), both without (green) and with metformin (orange). Left and central panels show single data points comparing two replicate plates, while the right panel shows the comparison of the averaged data of two replicates within each independent experiment. The Pearson correlation coefficients are shown in each panel.

Supplementary Figure S5. The parameter G corrects effects of differential growth rates that otherwise affect relative survivorship, S. **A**, Box plots of the G calculated for mutants with reduced fitness or without effects as previously reported (Costanzo et al. 2010). Deleterious knockouts, (fitness $f<0.95$, $n=10$) usually have a more negative G value than neutral knockouts without known growth defects ($f\geq0.95$, $n=67$) ($p=0.0014$, Wilcoxon rank sum test). **B**, Scatter plot showing the obtained S value with the parameter G is included in the multiple linear regression (horizontal axis) and when the regression runs without G (S_{woG}, vertical axis). **C,** The difference between calculations of S showed in panel A is mostly explained by average changes in the parameter G.

Supplementary Figure S6. Relative survivorship (rS) data rescaling. To quantitatively compare the relative phenotypes between two conditions of different WT phenotype, data were rescaled to a dimensionless relative survival parameter, rS. **A**, The death rates of the WT without (green) and with metformin (orange) were calculated by fitting to an exponential function the percent survival at different days of aging. Data points were from ten replicate samples in four replicate experiments (n=39 with metformin and n=40 for SC); lines are the average fits. **B**, The relative survival S shows a bias between the two conditions (left); the rescaled survivorship parameter rS provides a better comparison under two conditions in which the WT dies at different rates (right). Specifically, relative survivorship was rescaled with the formula $rS = -\log\left(1 - \frac{S}{r_{wt}}\right)$ using the average wild-type death rate obtained in each plate.

Supplementary Figure S7. Candidate gene-drug interactions and validation. ***A,*** Plot shows individual and average measurements of rS of strains with differential phenotypic effects ($p<0.05$, *t*-test). Gray diagonal lines are used to visualize the sign of the gene-drug interaction. ***B,*** Validation of gene-drug interactions using the OD kinetics method. Survival curves were determined through aging monocultures of eight gene-deletion strains with some of the most extreme short- or long-lived phenotypes in SC, including candidate gene-drug interactions. Average percentage of survivorship is shown for the wild-type (solid lines) and gene-deletion strains (dashed lines) in both conditions, SC (green) and SC+Metformin (orange); error bars indicate the S.E.M (*n*=4).