Title: Gamma oscillations weaken with age in healthy elderly in human EEG

Authors: Dinavahi V. P. S. Murty[1], Keerthana Manikandan[1], Wupadrasta Santosh Kumar[1], Ranjini Garani Ramesh[1], Simran Purokayastha[1], Mahendra Javali[2], Naren Prahalada Rao[3] and Supratim Ray[1,a].

Affiliations:

[1]Centre for Neuroscience, Indian Institute of Science, Bangalore, India, 560012,

Telephone +91 80 2293 3437, Facsimile +91 80 2360 3323

[2]MS Ramaiah Medical College & Memorial Hospital, Bengaluru, India

[3]National Institute of Mental Health and Neurosciences, Bengaluru, India

[a]**Corresponding Author:** sray@iisc.ac.in

Keywords: Gamma, EEG, Alzheimer's

Conflict of interest: The authors declare no competing financial interests.

Funding Disclosure: This work was supported by Wellcome Trust/DBT India Alliance (500145/Z/09/Z; Intermediate fellowship to SR), Tata Trusts Grant and DBT-IISc Partnership Programme.

Abstract

Gamma rhythms (~20-70 Hz) have been reported to be abnormal in mental disorders such as autism and schizophrenia in humans, and Alzheimer's disease (AD) models in rodents. However, the effect of normal aging on these oscillations is not known, especially for elderly subjects (>49 years) for which AD is most prevalent. In a first large-scale (236 subjects; 104 females) electroencephalogram (EEG) study on gamma oscillations on elderly subjects (aged 50-88 years), we presented full-screen Cartesian gratings that induced two distinct gamma oscillations (slow: 20-34 Hz and fast: 36-66 Hz). Power and centre frequency significantly decreased with age for both slow and fast gamma, but not alpha (8-12 Hz). Reduction was more salient for fast gamma than slow. These results were independent of microsaccades and pupillary reactivity to stimulus, as well as variations in power spectral density with age. Steady-state visual evoked potentials (SSVEPs) also reduced with age. These results are crucial first steps towards using gamma/SSVEPs as biomarkers of cognitive decline in elderly.

Significance statement

No study in humans has examined visual narrow-band gamma oscillations with healthy aging in elderly subjects. In a first large-scale (236 subjects) EEG study on stimulus-induced narrow band gamma in cognitively normal elderly (>49 years) humans, we show that both power and centre frequency of slow and fast gamma (20-34 Hz and 36-66 Hz respectively) decrease with age, but alpha power does not. Steady-state visual evoked potential (SSVEPs) in gamma range also decrease with age. Any EEG-based biomarker is accessible and affordable to patients of a wide socio-economic spectrum. Our results are important steps for developing such screening/diagnostic tests for aging-related diseases, like Alzheimer's disease.

Introduction

Gamma rhythms are narrow-band oscillations often observed in the electrical activity of the brain, with centre frequency occupying ~20-70 Hz frequency range. Previous studies have proposed involvement of these rhythms in certain higher cognitive functions like feature binding (Gray et al., 1989), attention (Gregoriou et al., 2009; Chalk et al., 2010) and working memory (Pesaran et al., 2002). Further, some studies have shown that these rhythms may be abnormal in neuropsychiatric disorders such as schizophrenia (Tada et al., 2014; Hirano et al., 2015), autism (Uhlhaas and Singer, 2007; Wilson et al., 2007; An et al., 2018) and Alzheimer's disease (AD; Palop and Mucke, 2016; Mably and Colgin, 2018).

Gamma rhythms can be induced in the occipital areas by presenting appropriate visual stimuli such as bars and gratings, and their magnitude and centre frequency critically depend on the properties of the stimulus such as contrast, size, orientation, spatial frequency and drift rate (Jia et al., 2013; Ray and Maunsell, 2015; Murty et al., 2018). Recently, we showed that large (full-screen) gratings induce two distinct narrow-band gamma oscillations in local field potentials (LFP) in macaque V1 and posterior electrodes in human EEG, which we termed slow (~20-40 Hz) and fast (~40-70 Hz) gamma (Murty et al., 2018). Fast gamma was not a harmonic of slow, but instead these rhythms were differently tuned to stimulus properties. Importantly, slow gamma was observed only when the grating size was sufficiently large (diameter >8º of visual angle for humans). Two distinct gamma rhythms have also been recently reported in human MEG (Pantazis et al., 2018) and in visual cortex (Veit et al., 2017) and hippocampus (Colgin et al., 2009) in rodents. These rhythms have been suggested to be generated from excitatory-inhibitory interactions of pyramidal cell and interneuron networks (Buzsáki and Wang, 2012), specifically involving parvalbumin (PV) and somatostatin (SOM) interneurons (Cardin et al., 2009; Sohal et al., 2009; Veit et al., 2017).

A recent study has reported PV interneuron dysfunction in parietal cortex of AD patients

and transgenic models of mice (Verret et al., 2012); and aberrant gamma activity in parietal

cortex in such mice. However, our knowledge about these rhythms in healthy aging in humans

is limited. Studies in human MEG have observed that the centre frequency of gamma

oscillations is negatively correlated with age of healthy subjects in the range of 8-45 years

(Muthukumaraswamy et al., 2010; Gaetz et al., 2012), but our understanding of these rhythms

in elderly humans (>49 years), which is clinically more relevant for studying diseases of

abnormal aging like AD, is lacking.

Further, visual stimulation of wild type and AD models of mice using light flickering at

40 Hz rescued AD pathology in visual cortex (Iaccarino et al., 2016). Such stimulation is

known to entrain brain rhythms and generate steady-state visual evoked potentials (SSVEPs)

at 40 Hz. However, to our knowledge, no previous study has examined SSVEPs in gamma

band in healthy elderly. Furthermore, a recent study has shown flattening of power spectral

density (PSD) in 2-24 Hz range in elderly subjects compared to younger subjects (Voytek et.

al., 2015). However, how this flattening affects gamma rhythms in elderly has not been

examined.

In this study, we described the variation of the two gamma rhythms in healthy elderly

subjects. We first used a battery of cognitive tests to identify a large cohort (236 subjects; 104

females) of cognitively healthy elderly subjects aged between 50-88 years. For comparison, we

also included 47 younger subjects (aged 20-48 years, 16 females). We induced gamma

oscillations using full-screen static Cartesian gratings while monitoring their eye movements

and recording EEG, and studied how slow and fast gamma and alpha oscillations, as well as

slope of the PSD, varied with age in elderly subjects. In a subset of subjects, we examined

SSVEPs in gamma frequency range (32 Hz) as well.

Materials and Methods

Human subjects

We recruited 236 elderly subjects (104 females) aged 50-88 years from the Tata Longitudinal Study of Aging cohort from urban communities in Bangalore through awareness talks on healthy aging and dementia. Recruitment was done by trained psychologists, who also collected their demographic details. Psychiatrists and neurologists at National Institute of Mental Health and Neurosciences (NIMHANS), Bangalore and M S Ramaiah Hospital, Bangalore assessed the cognitive function of these subjects using a combination of Clinical Dementia Rating scale (CDR), Addenbrook's Cognitive Examination-III (ACE-III), Hindi Mental State Examination (HMSE), and other structured and semi-structured interviews. We considered only those subjects who were labelled as cognitively healthy for this study. Out of 236 cognitively healthy subjects thus recruited, we discarded data of 9 subjects (3 females) due to noise (see Artifact Rejection section below for details). We were thus left with 227 subjects (101 females) aged 50-88 years (mean±SD: 66.8±8.2 years) for analysis.

Further, we also recruited 47 younger subjects (16 females) aged 20-48 years (mean±SD: 30.4±7.1 years) from the student and staff community of Indian Institute of Science. We screened them orally for any past history of neurological/psychiatric illness. We had presented data from 10 of these younger subjects in an earlier study (Murty et al., 2018).

All subjects participated in the study voluntarily and against monetary compensation. We obtained informed consent from all subjects for performing the experiment. The Institute Human Ethics Committees of Indian Institute of Science, NIMHANS, Bangalore and M S Ramaiah Hospital, Bangalore approved all procedures.

 EEG recordings

Experimental setup, EEG recordings and analysis were similar to what we had described

in our previous study (Murty et al., 2018). We recorded raw EEG signals from 64 active

electrodes (actiCAP) using BrainAmp DC EEG acquisition system (Brain Products GmbH).

We placed the electrodes according to the international 10-10 system. We filtered raw signals

online between 0.016 Hz (first-order filter) and 1000 Hz (fifth-order Butterworth filter),

sampled at 2500 Hz and digitized at 16-bit resolution (0.1 µV/bit). We rejected electrodes

whose impedance was more than 25 KΩ. This led to a rejection of 3.9% of electrodes in elderly

age-group (1.1% in younger subjects). However, most of these electrodes were frontal/central,

and specifically, none were the ten parieto-occipital/occipital electrodes used for analyses (see

Data Analysis subsection). Impedance of the final set of electrodes was 5.5±4.2 KΩ (mean±SD)

for elderly subjects and 3.7±3.4 KΩ for younger subjects. We referenced EEG signals to FCz

during acquisition (unipolar reference scheme).

Experimental setting and behavioral task

All subjects sat in a dark room in front of an LCD screen with their head supported by a

chin rest. The screen (BenQ XL2411) had a resolution of 1280 x 720 pixels and a refresh rate

of 100 Hz. It was gamma-corrected and was placed at a mean±SD distance of 58.1±0.9 cm

from the subjects (53.9-63.0 cm for all 274 subjects, 54.9-61.0 cm for the 227 elderly subjects)

according to their convenience (thus subtending a width of at least 52º and height of at least

30º of visual field for full-screen gratings). We calibrated the stimuli to the viewing distance in

all cases.

Subjects performed a visual fixation task. Stimulus presentation was done by a custom

software running on MAC OS that also controlled the task flow. Every trial started with the

onset of a fixation spot (0.1°) shown at the center of the screen, on which the subjects were

instructed to hold and maintain fixation. After an initial blank period of 1000 ms, a series of

stimuli (2 to 3) were randomly shown for 800 ms each with an inter-stimulus interval of 700

ms. Stimuli were sinusoidal luminance gratings presented full screen at full contrast. For the

main "Gamma" experiment, these were presented at one of three spatial frequencies (SFs): 1,

2, and 4 cycles per degree (cpd) and one of four orientations: 0°, 45°, 90° and 135°. Subjects

performed this task in 2-3 blocks (total 597 blocks across 283 subjects) during a single session

according to their comfort. For an initial subset of subjects, stimuli with SF of 0.5 and/or 8

were also presented, but we discarded these SFs from further analysis to maintain uniformity.

We also tested 32-Hz SSVEPs on a subset of the subjects who had analyzable data for the

gamma experiment (221/227 elderly and 46/47 younger subjects) according to their willingness

for the SSVEP experiment. Gratings (with SF and orientation that showed high change in slow

and fast gamma power for each subject) counter-phased at 16 cycles per second (cps) in a

similar stimulus presentation paradigm as described above, randomly interleaved with static

gratings of the same SF and orientation.

Eye position analysis

We recorded eye signals (pupil position and diameter data) using EyeLink 1000 (SR

Research Ltd., sampled at 500 Hz) during the entire trial for all but one subject. We calibrated

the eye-tracker for pupil position and monitor distance for each subject before the start of the

session. All the subjects were able to maintain fixation with a standard deviation of less than

0.6° (elderly, eye-data for Gamma experiment shown in Figure 7a) and 0.2° (young, data not

shown). We defined fixation breaks as eye-blinks or shifts in eye-position outside a square

window of width 5º centered on the fixation spot. We rejected stimulus repeats with fixation

breaks during -0.5s to 0.75s of stimulus onset, either online (and repeated the stimulus thus

discarded), or offline (we took a few additional trials to compensate for possible offline

rejection), according to the subjects' comfort. This led to rejection of 16.7±14.2% (mean±SD)

and 16.7±15.1% stimulus repeats for elderly subjects (for Gamma and SSVEP experiments

respectively), most of who preferred offline rejection. For younger subjects, for many of whom

we used online eye-monitoring, the rate of rejection due to fixation breaks was low (4.9±5.7%

and 4.2±7.0%).

<u>Artifact rejection</u>

We first estimated bad stimulus repeats for each unipolar electrode separately. We applied

a trial-wise thresholding process on both raw waveforms (high-pass filtered at 1.6 Hz to

eliminate slow trends if any) and multi-tapered PSD (computed between -500 ms to 750 ms of

stimulus onset using the Chronux toolbox (Mitra and Bokil, 2008, http://chronux.org/,

RRID:SCR_005547)). Any stimulus repeat for which either the waveform or the PSD deviated

by 6 times the standard deviation from the mean at any time bin (between -500 ms to 750 ms)

or frequency point (between 0-200 Hz) was considered a bad repeat for that electrode. We then

created a common set of bad repeats across all 64 unipolar electrodes by first discarding those

electrodes that had more than 30% of all repeats marked as bad, and subsequently assigning

any repeat as bad if it occurred in more than 10% of total number of remaining electrodes.

Finally, any repeat that was marked bad in any of the ten unipolar electrodes used for analysis

(P3, P1, P2, P4, PO3, POz, PO4, O1, Oz, and O2; see Data Analysis subsection) was

unconditionally included in the common bad repeats list, providing a final list of common bad

repeats for each block for each subject. In spite of these stringent conditions, these led to a

rejection of less than 20% of data (18.4±6.4% and 17.0±5.1% for elderly and younger subjects).

In addition, we calculated slopes (see Data Analysis subsection) of PSD (calculated with

1 taper and averaged across repeats, after removal of bad repeats) for each block in 56 Hz to

84 Hz range (to include the fast gamma range) for each unipolar electrode. Previous studies

have shown that in clean electrophysiological data, PSD slopes are typically between 0.5 to 4.5 (Podvalny et al., 2015; Shirhatti et al., 2016; Muthukumaraswamy and Liley, 2018; Sheehan et al., 2018). We therefore discarded those electrodes (5.0±5.9% for elderly and 5.2±7.7% for younger subjects) that had PSD slopes less than 0. We further discarded any block (53/497 and 5/100 for elderly and younger subjects) that did not have at least a single clean bipolar electrode pair in any of the three groups of bipolar electrodes used for analysis (depicted in Figure 3d, see Data Analysis subsection for details): PO3-P1, PO3-P3, POz-PO3 (left anterolateral group); PO4-P2, PO4-P4, POz-PO4 (right anterolateral group) and Oz-POz, Oz-O1, Oz-O2 (posteromedial group). We then pooled data across all good blocks for every subject separately for final analysis. Those subjects who did not have any analyzable blocks (9/236 and 0/47 for elderly and younger subjects respectively) were discarded from further analysis, leaving 227 elderly (aged 50-88 years, mean±SD: 66.8±8.2 years, females: 101) and 47 young subjects (aged 20-48 years, mean±SD: 30.4±7.1 years, females: 16) for analysis. The total number of repeats per electrode that were finally analyzed were 276.2±87.2 for elderly subjects and 270.4±67.4 for younger subjects.

We applied a similar artifact rejection procedure for SSVEP experiment. Out of subjects with analyzable blocks for the Gamma experiment, 197 elderly (mean±SD: 66.8±7.8 years, females: 93) and 43 young (mean±SD: 30.4±7.3 years, females: 15) subjects had analyzable blocks (242/270) for SSVEP experiment. Using similar selection criteria as before, we rejected 7.7±5.2% of repeats for elderly subjects and 6.6±4.1% for younger subjects. The total number of analyzed repeats per electrode for counter-phasing condition were 30.2±6.9 and 29.7±6.6 for elderly and younger subjects respectively.

    EEG data analysis

For all analyses (unless otherwise mentioned), we used bipolar reference scheme. We re-

referenced data at each electrode offline to its neighboring electrodes. We thus obtained 112

bipolar pairs out of 64 unipolar electrodes (Murty et al., 2018, depicted in Figure 3e). We

considered the following bipolar combinations for analysis, except for scalp maps: PO3-P1,

PO3-P3, POz-PO3 (left anterolateral group); PO4-P2, PO4-P4, POz-PO4 (right anterolateral

group) and Oz-POz, Oz-O1, Oz-O2 (posteromedial group), depicted in Figure 3d. We

discarded a bipolar electrode if either of its constituting unipolar electrodes was marked bad as

described in the previous section. Data was pooled for the rest of the bipolar combinations in

each of the electrode groups for further analysis.

We analyzed all data using custom codes written in MATLAB (The MathWorks, Inc,

RRID:SCR_001622). We computed PSD and the time-frequency power spectrograms using

multi-taper method with a single taper using Chronux toolbox. We chose baseline period

between -500 ms to 0 ms of stimulus onset, while stimulus period between 250 ms to 750 ms

to avoid stimulus-onset related transients, yielding a frequency resolution of 2 Hz for the PSDs.

We calculated time frequency power spectra using a moving window of size 250 ms and step

size of 25 ms, giving a frequency resolution of 4 Hz.

We calculated change in power in alpha rhythm and the two gamma rhythms as follows:

$$\Delta Power = 10(log_{10}\frac{\sum_f ST(f)}{\sum_f BL(f)})$$

Where ST and BL are stimulus and baseline power spectra (across frequency f) averaged

across repeats for all stimulus conditions and analyzable bipolar electrodes. For slow gamma,

$f \in [20\ 34]$ Hz and for fast gamma, $f \in [36\ 66]$ Hz. Note that due to this averaging, on

average more than 250 stimulus repeats were available for analysis, yielding very reliable

estimates. We defined the center frequency for a gamma rhythm as the frequency at which the

change in power (in these averaged PSDs) was maximum within that gamma range. For SSVEP

experiment, we analyzed only the counter-phasing condition. Hence, we took the power at 32

Hz (twice the counter-phasing frequency, i.e. 16 cps) for analysis.

We generated scalp maps using the topoplot.m function of EEGLAB toolbox (Delorme

and Makeig, 2004, RRID:SCR_007292), modified to show each electrode as a colored disc,

with color representing the change in power of slow gamma/fast gamma/SSVEP from baseline

in decibels (dB).

We calculated slopes for rejecting noisy electrodes (as described in Artifact Rejection

subsection) by fitting PSD across all analyzable repeats for each individual unipolar electrode

with a power-law function as follows:

$$P(f) = A. f^{-\beta}$$

where P is the PSD across frequencies $f \in [56\ 84]$ Hz. A (scaling factor) and β (slope) are

free parameters obtained using least square minimization using the program *fminsearch* in

MATLAB. We similarly estimated slopes for PSDs averaged across analyzable unipolar or

bipolar electrodes during baseline period (-0.5 to 0 ms) for Supplementary Figure 1.

<u>Microsaccades and pupil data analysis</u>

We detected microsaccades using a threshold-based method described earlier (Murty et

al., 2018), initially proposed by (Engbert, 2006). In brief, we categorized eye movements with

velocities that crossed a specified threshold for at least a specified duration of time as

microsaccades. We set the velocity threshold between 3-6 times the standard deviation of eye-

velocities and minimum microsaccade duration between 10-15 ms for every subject so as to

maximize the correlation between peak velocity and amplitude of all microsaccades for that

subject (also called a "main sequence", see Engbert, 2006 for details), while maintaining the

minimum microsaccade velocity at 10º/s and the microsaccade rate between 0.5/s and 3.0/s.

The above algorithm was applied for the analysis period of -0.5 s to 0.75 s of stimulus

onset. After removing the microsaccade-containing repeats, there were 128.1±71.1 (mean±SD,

minimum 5) repeats for elderly subjects (n=226, excluding 1 subject for whom eye-data could

not be collected) for anterolateral electrodes reported in Figure 7c. Results did not change when

we discarded 13 elderly subjects with less than 30 repeats without microsaccades from analysis

(data not shown).

EyeLink 1000 system recorded pupil data in arbitrary units for every subject since pupil

data cannot be calibrated for this tracker. Hence, instead of directly comparing time-series of

pupil data, we used coefficient of variation (CV, ratio of standard deviation to mean) for every

repeat as a measure of pupillary reactivity to stimulus of that repeat. This simple measure scales

standard deviation of a distribution with respect to its mean. This allows comparison of

variation in different distributions without getting affected by the mean of the distributions. We

calculated CV for each analyzable trial separately and calculated mean CV across trials for

every subject for comparison.

<u>Statistical analysis</u>

Our findings were based mainly on PSD plots and we used appropriate statistical methods

(Pearson correlation, linear regression and ANOVA) to confirm our interpretations. We used

one-way (or two-way, as necessary) ANOVA to compare means of bar plots in Figures 4c, 4d,

6a and 8c, although non-parametric tests on medians instead of means using Kruskal-Wallis

test (not reported) yielded qualitatively similar results. For two-way ANOVA, we considered

age-group and gender as independent factors although including their interaction effect in the

model yielded qualitatively similar results (not reported). We used Bonferroni correction for

multiple tests/comparisons wherever necessary.

Results

We recorded EEG from 236 elderly subjects aged 50-88 years and 47 subjects aged 20-48 years while presenting full-screen sinusoidal grating stimuli on a computer monitor. The stimuli differed in spatial frequency and orientation. Each stimulus was presented for 800 ms following a baseline period of 700 ms. The subjects were required to maintain fixation throughout the trial (containing 2-3 stimuli), while their eye-data was recorded for offline analysis. For analysis, we pooled the data across the four orientations (0º, 45º, 90º, 135º) for all subjects, as orientation selectivity was reported to be very low for humans for both slow and fast gamma bands (Murty et al., 2018). Similarly, we pooled across three spatial frequencies (1-4 cpd) which were reported to elicit salient gamma oscillations in humans (Murty et al., 2018).

Figure 1 shows the results of an example subject, a 53 years old female. Trial-averaged evoked potentials were plotted (Figure 1a) for electrodes P3, P1, P2, P4, PO3, POz, PO4, O1, Oz, O2 for unipolar reference (Figure 1a, left column) and PO3-P1, PO3-P3, POz-PO3, PO4-P2, PO4-P4, POz-PO4, Oz-POz, Oz-O1 and Oz-O2 (shown as dots in scalp maps in Figure 1c) for bipolar reference (Figure 1a, right column). These traces revealed a transient in the first 250 ms of stimulus onset and after the stimulus offset (i.e. after 800 ms). For the same set of electrodes, trial-wise power spectrograms were averaged to generate raw spectrogram and change in power spectrogram (w.r.t. a baseline period of -500 ms to 0 ms of stimulus onset), plotted in Figure 1a. Although not noticeable in the evoked potential traces and raw spectrograms, these stimuli elicited prominent gamma band responses as seen in the change in power spectrograms. These responses were in slow gamma (~20-34 Hz) and fast gamma (~36-66 Hz) range. Consistent with previous results (Murty et al., 2018), these responses were seen during the stimulus period (after the onset-transient) and were best noticed for bipolar reference (compared to unipolar reference). Also, slow gamma power showed a gradual build-up whereas

fast gamma power showed a decreasing trend with stimulus duration (Figure 1a, bottom row).

Alpha (8-12 Hz) power suppression was very weak in this subject. We also plotted power

spectral densities (PSD) in the baseline period (dotted black trace in Figure 1b) and stimulus

period (250 ms to 750 ms; solid black trace in Figure 1b) and change in power spectrum (blue

trace in Figure 1b). Prominent 'bumps' in the slow and fast gamma range were noticeable in

PSD in the stimulus period as well as change in spectrum. Also, no 'bump' was noticeable in

the baseline PSD in the alpha range for this subject. These changes were most prominent in the

parieto-occipital and occipital electrodes, as seen in the scalp maps for the bipolar reference

case in Figure 1c.

Our primary emphasis was to characterize gamma and other spectral signatures as a

function of age within the elderly population (>49 years), for which we divided these subjects

into two groups: 50-64 years (95 subjects; 51 female) and >64 years (141 subjects, 53 female).

For completeness, we also show results from a cohort of younger subjects aged between 20-49

years (47 subjects; 16 female).

Baseline power of slow and fast gamma did not vary with age

A recent study (Voytek et al., 2015) has suggested rotation of PSDs with age such that

power at frequencies above ~15 Hz is higher in elderly subjects compared to younger subjects,

leading to flatter PSDs in elderly subjects. This could potentially bias the estimation of change

in power in slow and fast gamma range in subjects of different age groups as higher baseline

power in these rhythms in older subjects may lead to lower estimates of change in power.

Hence, we first checked whether there was any difference in baseline PSDs across age. We

calculated mean baseline PSDs of 10 unipolar electrodes and 9 bipolar electrodes separately,

as mentioned above. We compared PSDs between 2-200 Hz in the two elderly groups as well

as the younger group (Figures 2a and 2b for males and females), and males versus females

(averaged across all ages; Figure 2c).

Because our primary emphasis was comparison within the elderly group, we first

compared the PSDs between the two elderly subgroups (dark and light gray traces in Figures

2a and 2b). The PSDs indeed appeared to become flatter with age (light gray trace was above

the dark gray trace), but this effect was prominent only at frequencies above ~50 Hz. In the

slow and fast gamma ranges (indicated by coloured bars on the abscissa of plots in Figure 2),

the two gray traces were largely overlapping. To quantify this, we performed a two-way

ANOVA on baseline powers of alpha, slow gamma and fast gamma with age-group (50-64 or

>64 years) and gender as factors and found that effect of age group was not significant for

power in any band (p>0.05 in all cases except for fast gamma in bipolar case where p=0.03,

which was not significant at Bonferroni corrected significance level of 0.05/3 = 0.016). Results

were not qualitatively different when we performed one-way ANOVA for baseline power of

alpha/slow/fast gamma across age-groups for males and females separately (p>0.05 for all

cases except for fast gamma in females for bipolar case where p=0.03).

We obtained similar trends for comparisons (one-way ANOVA separately for males

and females) between younger (<50 years) and elderly subjects (50 years and above). Baseline

powers in alpha/slow/fast gamma ranges were not significantly different for younger and

elderly male subjects in either reference schemes (Figure 2a, p>0.05 for all cases). However,

elderly females had more baseline fast gamma power compared to younger females

($F(1,115)$=7.9, p=0.006) in bipolar case and lesser alpha power in both unipolar

($F(1,115)$=17.6, p=5.5*10^{-5}) and bipolar ($F(1,115)$=6.5, p=0.012) cases (Figure 2b). These

differences could be due to a small sample size of females in the younger age-group (n=16).

Across genders, females had significantly higher baseline slow gamma power than

males (Figure 2c, data pooled across all 274 subjects; one-way ANOVA across gender:

$F(1,272)=24.5/27.9$, p=$1.3*10^{-6}/2.6*10^{-7}$ for unipolar/bipolar reference schemes) and higher

alpha power ($F(1,272)=4.6/8.4$, p=0.03/0.004 for unipolar/bipolar conditions). However,

baseline fast gamma power was not significantly different (p>0.05 for both reference schemes).

Amongst the elderly subjects (n=227, data not shown), females had only higher slow gamma

compared to males ($F(1,225)=16.4/21.3$, p=$7.1*10^{-5}/6.4*10^{-6}$ for unipolar/bipolar conditions

for slow gamma, $F(1,225)=5.3$, p=0.022 for alpha in bipolar case and p>0.05 for all other

cases).

To test for the rotation of PSDs with age as suggested by Voytek *et al.* (2015), we

computed the slopes between 16-44 Hz (Supplementary Figure 1; see Methods for details; this

range was chosen to avoid the bump in the alpha band at the lower end and the 50 Hz noise at

the higher end). Two-way ANOVA with age (young and elderly) and gender (male and female)

as factors showed no significant difference in the slopes between young and elderly subjects

for either unipolar or bipolar reference scheme case (p>0.05). However, females had steeper

slopes compared to males ($F(1,271)=7.9$, p=0.005 and $F(1,271)=31.4$, p=$5.1*10^{-8}$ for unipolar

and bipolar cases respectively, Supplementary Figure 1a). Since females had higher baseline

alpha power compared to males (Figures 2c), we tested whether any differences in baseline

PSD slopes could be because of differences in baseline alpha power. We divided baseline PSDs

of all subjects (young and elderly pooled together) into terciles based on alpha power (Figure

2d). Subjects who had higher baseline alpha power also had steeper PSD slopes. Regression of

PSD slopes in 16-44 Hz frequency range with baseline alpha power was significant for both

reference schemes (Supplementary Figure 1b). Further, when we performed partial correlation

of slopes with age and baseline alpha power, slopes were significantly correlated with alpha

power (rho=0.57, p=$5.4*10^{-25}$ and rho=0.58, p=$3.1*10^{-26}$ for unipolar and bipolar cases

respectively) but not with age (rho=0.07 and -0.12 for unipolar and bipolar, p>0.05 for both).

Thus, PSD slope was not influenced by age, but by baseline alpha power. We discuss these

results in the context of the findings of Voytek and colleagues in the Discussion.

<u>Gamma was observed in more than 80% of subjects</u>

As reported in our earlier study (Murty et al., 2018) and as in Figure 1, gamma was best

observed, as a response to full-screen 100% contrast Cartesian visual gratings, in bipolar

referencing scheme compared to unipolar. Hence, we limited further analysis to bipolar

referencing. We divided the 9 bipolar electrodes mentioned above into 3 groups (Figure 3d):

PO3-P1, PO3-P3, POz-PO3 (left anterolateral group); PO4-P2, PO4-P4, POz-PO4 (right

anterolateral group) and Oz-POz, Oz-O1, Oz-O2 (posteromedial group). For each subject, we

chose the electrode group that had maximum change in power in slow and fast gamma ranges

added together. We labelled a subject as having either of the gamma rhythms if the change in

power in these rhythms during stimulus period (calculated from data pooled across electrodes

chosen for the subject) exceeded an arbitrarily chosen threshold of 0.5 dB from baseline. Figure

3a shows scatter plot of slow versus fast gamma change in power for all subjects. Based on our

threshold, ~84% of subjects had at least one gamma (slow: ~77% and fast: ~64%), while ~57%

of subjects had both the gammas, which could be observed as distinct "bumps" in the change

in PSD from baseline (Figure 3b). Figure 3c shows the percentage of subjects in each age-

group who had no/slow/fast/both gammas based on our threshold. The percentage of subjects

who had only fast gamma or both gammas was highest in 20-49 years age-group and lowest in

>64 years age-group.

Figure 3e shows change in power in slow (top row) and fast (bottom row) gamma

rhythms across all electrodes (plotted as disks) for the young (left column) and the two elderly

age-groups (middle and right columns). Both gamma rhythms were best observed in the same

9 bipolar electrodes mentioned above and depicted in Figures 3d and 3e. Further, power in both

gamma bands appeared to decrease with age across the two elderly age-groups, although the

results were more prominent for fast gamma.

<u>Change in gamma power was negatively correlated with age</u>

To quantify this difference, we tested how gamma oscillations correlated with age in

these electrode groups. We tested for anterolateral and posteromedial groups separately

(Figures 4 and Supplementary Figure 2 respectively). For Figure 4, out of the left and right

anterolateral groups, we chose that group which had maximum slow and fast gamma power

change summed together. Figures 4a and 4b show mean change in spectrograms and PSDs

respectively for the three age-groups separately for males and females. These plots highlight

all the major results discussed later. First, both slow and fast gamma power reduced with age.

This was observed between young and elderly groups (black versus the other two traces), and

also within the two elderly sub-groups (dark and light gray traces). Second, peak frequencies

of both slow and fast gamma reduced with age. Third, alpha suppression (change in 8-12 Hz

power from baseline) in the stimulus period was more pronounced in young versus elderly, but

there was no difference between the two elderly sub-groups.

The first observation was also reflected in the gamma power computed within the pre-

specified ranges (as shown in the bar plots shown in Figure 4c and 4d), but there were some

caveats. We computed the total power within a pre-specified band, which typically has larger

contribution from lower frequencies because the absolute power is larger compared to higher

frequencies within the band (which is not reflected in Figure 4b because it only shows the

change in power). Consequently, if the traces are overlapping at lower frequencies within the

band and diverge at higher frequencies, which was the case in the slow gamma range for both

males and females (Figure 4b), the total power in the band may not be significantly different.

In particular, for young females, the power at the start of the slow gamma band (20-26 Hz) was

slightly lower than the elderly subgroups (Figure 4b, bottom plot, black versus gray traces), but

became higher at higher frequencies within the slow gamma band (28-34 Hz). However,

because the absolute power is higher between 20-26 Hz than 28-34 Hz, the total slow gamma

power was actually lower for young females compared to elderly (Figure 4c, black versus gray

bars). These issues can be partially addressed by changing the frequency range over gamma is

computed (dependent on age and potentially even across subjects), but then the results are

dependent on the level of customization of ranges, which we wanted to minimize. We observed

that younger subjects had significantly more fast but not slow gamma than elderly subjects

(two-way ANOVA with age-group (20-49 and >49 years) and gender as factors,

$F(1,271)=1.3/35.6$, p=$0.2/7.6*10^{-9}$ for slow/fast gamma across age-groups). Also, females had

more slow and fast gamma than males (same two-way ANOVA, $F(1,271)=4.7/37.9$,

p=$0.03/2.5*10^{-9}$ for slow/fast gamma).

Among the elderly subjects, visual inspection of change in spectrograms and spectra

revealed that both slow and fast gamma power was less in subjects of >64 years age-group

compared to 50-64 years age-group. This trend was also noticeable in the bar plots in Figures

4c and 4d for both genders. As before, it was significant only for fast gamma (two-way

ANOVA with age-group (50-64 and >64 years) and gender as factors; $F(1,224)=2.4/11.4$,

p=$0.12/8.4*10^{-4}$ for slow/fast gamma across age-group). Females had higher slow and fast

gamma compared to males (same two-way ANOVA, $F(1,224)=7.4/21.7$, p=$0.007/5.4*10^{-6}$ for

slow/fast gamma across gender). We further quantified this observation by regressing change

in slow and fast gamma power across age (scatter plots in Figures 4c and 4d). When the

regression was done separately for males and females, the slopes were always negative (males:

β=-0.008/-0.018 and females: β=-0.018/-0.016 for slow/fast gamma) but did not reach

significance except for fast gamma in elderly males (p=$2.4*10^{-4}$). When we pooled data across

both genders, the results were significant (linear regression, β=-0.02, R^2=0.02, p=0.022 and β=-

0.02, R^2=0.08, p=$1.4*10^{-5}$ for slow and fast gamma respectively). Similar, albeit weaker results

were observed in the posteromedial group of electrodes for slow gamma (Supplementary

Figure 2c; linear regression, β=-0.016, R^2=0.02, p=0.04) as well as fast gamma (Supplementary

Figure 2d; β=-0.02, R^2=0.04, p=0.001).

<u>Centre frequency of slow and fast gamma was negatively correlated with age</u>

Gamma peak centre frequency was shown to decrease with age in an age group between

8-45 years (Muthukumaraswamy et al., 2010; Gaetz et al., 2012). This was observed in our

data as well as noted above. To examine the change in centre frequency of slow and fast gamma

rhythms in elderly in more detail, we plotted the change in power spectra (frequencies

mentioned on abscissa) vs age (on ordinate, arranged in increasing order from top to bottom)

of all 227 elderly subjects, separately for anterolateral (left column) and posteromedial (right

column) group of electrodes. We defined centre frequency for each gamma as the frequency

that had maximum change in power in the frequency range of that gamma, provided the total

change in power in that gamma band was greater than our threshold of 0.5 dB (represented by

circles and triangles for slow and fast gamma in Figure 5a; number of subjects having slow and

fast gamma power change above this threshold is mentioned in Figure 5b). Figure 5b shows

the same result as scatter plots of centre frequencies of slow (top row) and fast gamma (bottom

row) plotted against the age of the subjects. Solid lines in Figures 5a and 5b indicate regression

fits of centre frequencies against age (showing decreasing trend). This trend was significant for

both slow and fast gamma in the anterolateral group (linear regression for centre frequency vs

age: β=-0.08, R^2=0.04, p=0.008 for slow gamma and β=-0.16, R^2=0.06, p=0.008 for fast

gamma). Similar, albeit weaker results were observed for the posteromedial group (fast

gamma: β=-0.17, R^2=0.06, p=0.008; slow gamma: β=-0.06, R^2=0.02, p=0.052). Note that

because our analysis was done over 500 ms of data, the frequency resolution was 2 Hz, which

limited our ability to observe small shifts in the peak frequency.

<u>Alpha suppression did not reduce with age in elderly subjects</u>

We noticed prominent alpha suppression for younger as well as elderly subjects, as

noted above (and in Supplementary Figures 2a and 2b). Alpha suppression was stronger in

younger subjects compared to elderly subjects (data for anterolateral group is shown in Figure

6a; two-way ANOVA with age-groups (20-49 and >49 years) and gender as factors;

$F(1,271)$=33.2, p=$2.2*10^{-8}$ across age-groups), but did not differ significantly between genders

($F(1,271)$=0.5, p=0.49 across gender). Amongst elderly subjects however, alpha suppression

did not differ across age-groups (50-64 and >64 years) and gender (two-way ANOVA, p>0.05

for both age-group and gender). We further confirmed this observation by regressing alpha

suppression across age for all the elderly subjects (scatter plot in Figure 6a). Alpha suppression

was not significantly correlated with age for either gender or for data pooled across genders

(p>0.05 for all cases). The trends were not qualitatively different when we repeated the analysis

for the posteromedial group of electrodes. This is also observed in the scalp maps shown in

Figure 6b. We did not test for alpha frequency as 2 Hz frequency resolution of our PSDs left

us with only 3 frequency points in the alpha range (8, 10 and 12 Hz).

 Microsaccades and pupillary reactivity did not contribute to negative correlation between

 change in gamma power and age

Next, we studied the potential contribution of eye-movement (including microsaccades)

and pupillary diameter on our results. Figure 7a shows mean eye-position for each of the elderly

age-groups in horizontal (top row) and vertical (middle row) directions (n=226, eye data was

unavailable in one subject; thickness represent SEM). Eye-position did not vary in the two age-

groups in either direction. Further, we extracted microsaccades for every analysed trial for

every subject in the two age-groups (see Materials and Methods), yielding a total of 27350 and

38373 microsaccades for the two age-groups. Figure 7b shows a scatter plot of peak velocity

versus maximum displacement for each microsaccade (a plot called "main sequence", see

Engbert, 2006). These microsaccade clouds were highly overlapping for these two groups and

had similar microsaccade rates (0.80±0.05/s and 0.88±0.05/s). Histograms of microsaccade rate

during -0.5 − 0.75 s of stimulus onset for both the elderly age-groups were also highly

overlapping (Figure 7a, bottom row), although we see a trend of slightly higher microsaccade

rate for subjects aged >64 years compared to 50-64 years age-group. We then computed power

after removing trials containing microsaccades (see Materials and Methods for details), and

could replicate the results in Figure 4: change in both slow and fast gamma power decreased

with age significantly (β=-0.02, R^2=0.03, p=0.015 for slow gamma and β=-0.02, R^2=0.08,

p=2.7*10^{-5} for fast gamma, Figure 7c).

We next tested if pupillary reactivity to stimulus presentation affected change in gamma

power with age. We calculated mean coefficient of variation (CV) of pupil diameter across

every analysable trial for all 226 subjects (eye data was unavailable in one subject). We

observed that mean CV decreased significantly with age in the elderly subjects, possibly

because of senile miosis (Pearson correlation, r=-0.24, p=3.5*10^{-4}, Figure 7d top row).

However, neither slow nor fast gamma power varied with mean CV of pupil diameter

(slow/fast: r=0.07/0.1, p=0.31/0.14 and r=0.09/0.12, p=0.19/0.06 for anterolateral (Figure 7d

middle and bottom rows) and posteromedial electrodes respectively (data not shown)).

<u>32 Hz SSVEP power was negatively correlated with age</u>

Finally, we checked whether SSVEPs in the gamma range were affected by healthy

aging. Specifically, we tested 32-Hz SSVEPs elicited by gratings counter-phasing at 16 cps.

Figure 8a and 8b show change in power spectrograms and spectra respectively for males and

females separately for the two elderly and the younger age-groups for the anterolateral group

of electrodes, with same conventions as in Figures 4. We saw clear peaks at 32 Hz in both

change in power spectrograms and PSDs. Insets in Figure 8b show a zoomed-in image of the

respective change in PSDs to show the difference in these peaks for the three age-groups.

Amongst the elderly age-groups, the mean SSVEP change in power was less in the >64 years

age-group compared to 50-64 years age-group in both males and females. We regressed the

SSVEP power change with age (scatter plot in Figure 8c bottom row, shown separately for

males and females). Change in SSVEP power at 32 Hz decreased significantly with age for

both males and females separately (males: β=-0.17, R^2=0.09, p=0.002 and females: β=-0.18,

R^2=0.08, p=0.007) as well as when the data were pooled across genders (β=-0.19, R^2=0.11,

p=1.4 X 10^{-6}, regression fit indicated by black line in bottom row of Figure 8c).

We repeated this analysis for posteromedial group of electrodes and noticed similar

results (regression of change in 32 Hz SSVEP power versus age for males: β=-0.16, R^2=0.11,

p=0.0008, females: β=-0.14, R^2=0.06, p=0.02 and for data pooled across gender: β=-0.17,

R^2=0.11, p=1.5 X 10^{-6}). We noticed this decrease of 32 Hz SSVEP power with age also in the

mean scalp maps for all analysable electrodes across all subjects in the three age-groups, as

depicted in Figure 8d.

Discussion

We tested for age-dependent variation of stimulus-induced change in power and centre frequency of narrow-band gamma oscillations in both slow and fast gamma frequency ranges in healthy elderly subjects aged 50-88 years. We observed a decrease in power of both slow and fast gamma oscillations with age, although the decrease in fast gamma was more salient than slow gamma. On the other hand, level of alpha suppression did not change with age in elderly subjects. Finally, centre frequency of both gamma rhythms decreased with age in these subjects. As there was no significant change in baseline slow/fast gamma power, eye-position and microsaccade rate across age, we ruled out the possibility that the age-related variations in gamma could be because of such factors. Further, we also studied variation of 32 Hz SSVEP power with age and observed a negative correlation. All these results were also analysed for a cohort of younger subjects (aged 20-48), and the results were consistent with previous reports on gamma oscillations and some of the trends observed in the elderly population.

Baseline alpha power and alpha suppression

Previous studies have suggested reduction in baseline alpha power in elderly subjects compared to younger subjects (Babiloni et al., 2006). Also, task-related modulation of alpha power was seen to be reduced in older adults compared to younger subjects (Vaden et al., 2012). Our results were similar to these previous reports: baseline alpha power was significantly higher in younger females versus elderly (Figure 2b), and showed a decreasing trend with age in males (although not significant, Figure 2a). Similarly, stimulus-induced alpha suppression was stronger for younger subjects compared to elderly subjects (Figure 6a). This is notwithstanding the different recording paradigms from previous studies: in our study, baseline alpha was recorded during eyes-open state (as opposed to resting, eyes-closed state in

Babiloni and colleagues, (2006)) and alpha suppression was measured during passive fixation

(as opposed to an active memory task in Vaden and colleagues (2012)). Among the elderly

subjects, however, neither baseline alpha power (Figures 2a and 2b) nor alpha suppression

(Figure 6a) varied with age. Different results for alpha versus gamma power (which decreased

with age) in elderly subjects suggest different biophysical mechanisms of these oscillations.

<u>Baseline PSD slopes</u>

Some authors have suggested that power-law distribution ($1/f^{\beta}$, where β is the PSD

slope) of brain electrical activity represents broadband scale-free activity of brain that is

dependent on behavioral states (He et al., 2010; He, 2014; Podvalny et al., 2015) and cognitive

abilities (Voytek et al., 2015; Sheehan et al., 2018). Specifically, Voytek and colleagues had

suggested that flattening of PSD slopes might be a hallmark of senile physiological cognitive

decline. In our study however, we did not notice any significant correlation between baseline

PSD slopes and age in the unipolar reference scheme (as used by Voytek and colleagues),

especially for elderly subjects. There are several reasons that could have led to this discrepancy.

First, we estimated broadband slopes in the range of 16-44 Hz as opposed to 2-24 Hz (as in

Voytek et al.). This is to avoid the contribution of baseline alpha power (8-12 Hz), against

which we were testing for slopes (Figure 2d and Supplementary Figure 1b). Second, the sample

size of Voytek and colleagues was small (11 young and 13 elderly) with a larger proportion of

females in the younger group (male:female = 4:7 and 8:5 in young and elderly groups). Because

females had steeper slopes than males (Figure 2c), underrepresentation of females in the elderly

group could have led to flatter PSDs in their data. Finally, we found that PSD slopes were

correlated with baseline alpha power (which was higher in younger versus elderly), but there

was no dependence of slope on age when controlled for baseline alpha power (using partial

correlation). Note that a similar correlation of slopes with alpha power in human MEG and

EEG as well as monkey ECoG has also been reported by Muthukumaraswamy and Liley

(2018).

We note, however, that the PSDs did tend to become flatter with age, albeit at a higher

frequency range (>50 Hz; Figure 2a and 2b), consistent with the ECoG results of Voytek and

colleagues and consistent with the neural noise hypothesis proposed by them. Further, our

"spontaneous activity" used for PSD computation was during the fixation task itself, and PSDs

were computed using segments of 500 ms, much less than the 2 second segments used by

Voytek and colleagues. Consequently, the frequency resolution was 4 times higher in the study

of Voytek and colleagues, which could have led to the identification of small changes in slopes

better than ours. Longer stimulus-free epochs (at least 2 seconds or more), preferentially in

both eyes closed and eyes open conditions are required to test whether the flattening of PSD

slope occurs at lower frequencies as well.

<u>Possible confounds from ocular factors</u>

Broadband induced gamma responses have been proposed to be correlated with

occurrence of microsaccades (Yuval-Greenberg et al., 2008). However, in our previous study,

we did not note any effect of microsaccades on orientation tuning of narrow-band slow and fast

gamma oscillations in macaques (Murty et al., 2018). Consistently, we did not find any effect

of microsaccades on age-dependent decrease of slow and fast gamma power in this study.

It is possible that retinal illuminance is reduced due to senile pupillary miosis, which is

indirectly reflected in the reduced pupillary reactivity to stimulus presentation across age

(Figure 7c). Other abnormalities of peripheral visual system like age-related increase in density

of crystalline lens, age-related macular degeneration, etc. could have had affected our results

(Owsley, 2011). The subjects did not undergo a thorough ophthalmic examination due to time

limitations. However, we argue that the results presented here are likely due to

neurophysiological effects of aging on two grounds. First, in addition to a reduction in gamma

power, there is a reduction in gamma centre frequency with age, which is harder to explain

based on the abnormalities listed above. Second, slow/fast gamma power was not dependent

on pupillary reactivity to stimulus (Figure 7d). Nonetheless, we observed that the percentage

of variance in the gamma power/frequency or SSVEP power explained by age is very less

(maximum R^2 among all cases was 0.11 for decrease in SSVEP power across age in

posteromedial electrodes). Hence, we recognise that age is one of the many possible factors

that influence gamma power/frequency and do not completely rule out the possibility that any

hidden physiological variables could have had contributed to this variance.

<u>Possible mechanisms of age-related reductions in gamma frequency and change in power</u>

Previous studies in MEG had reported significant positive correlations between (fast)

gamma frequency and cortical thickness as well as volume of cuneus (Gaetz et al., 2012) and

thickness of pericalcarine area (Muthukumaraswamy et al., 2010), measured through structural

MRI. Further, age-related decreases in cortical volume, thickness and/or surface area were

observed in various regions of the brain like precuneus, cuneus, lingual, pericalcarine and

lateral occipital areas of the occipital cortex (Salat et al., 2004; Lemaitre et al., 2012). In line

with these observations, these MEG studies have also noted decrease of centre frequency of

gamma with age, esp. in young and middle-aged adults. Similarly, (fast) gamma peak

frequency has been positively correlated with brain GABA levels (Edden et al., 2009;

Muthukumaraswamy et al., 2009). Although we have not tested directly for these variables in

our study, we speculate that these morphological and neurochemical factors are the reasons for

our observations regarding fast gamma frequency. However, the factors behind reduction of

slow/fast gamma power and slow gamma frequency with age remain open questions.

To conclude, our study throws light on various features of baseline spectra (like baseline

alpha power and its relation to PSD slopes) and spectral responses to Cartesian gratings (alpha

suppression, slow and fast gamma) in a large cohort of healthy elderly. Our study could thus

act as normative for future gamma and SSVEP studies in the elderly age-group. Further, based

on observations in previous rodent studies (for example, Verret et al., 2012; Iaccarino et al.,

2016) as described before, some authors have suggested a causative role of (fast) gamma

disruption in neurodegenerative disorders of aging such as AD (Palop and Mucke, 2016).

Alternatively, our results suggest that gamma and SSVEPs suffer reduction in power with age

even in the absence of cognitive decline. These studies taken together, decrease in

gamma/SSVEP power may represent a continuum of healthy aging – preclinical cognitive

decline – dementia spectrum and may act as a harbinger to senile or pathological cognitive

decline, a hypothesis that needs to be tested in future studies.

References

An K, Ikeda T, Yoshimura Y, Hasegawa C, Saito DN, Kumazaki H, Hirosawa T, Minabe Y, Kikuchi M
(2018) Altered Gamma Oscillations during Motor Control in Children with Autism Spectrum
Disorder. J Neurosci 38:7878–7886.

Babiloni C, Binetti G, Cassarino A, Forno GD, Percio CD, Ferreri F, Ferri R, Frisoni G, Galderisi S, Hirata
K, Lanuzza B, Miniussi C, Mucci A, Nobili F, Rodriguez G, Romani GL, Rossini PM (2006)
Sources of cortical rhythms in adults during physiological aging: A multicentric EEG study.
Hum Brain Mapp 27:162–172.

Buzsáki G, Wang X-J (2012) Mechanisms of Gamma Oscillations. Annu Rev Neurosci 35:203–225.

Cardin JA, Carlén M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai L-H, Moore CI (2009) Driving
fast-spiking cells induces gamma rhythm and controls sensory responses. Nature 459:663–
667.

Chalk M, Herrero JL, Gieselmann MA, Delicato LS, Gotthardt S, Thiele A (2010) Attention Reduces
Stimulus-Driven Gamma Frequency Oscillations and Spike Field Coherence in V1. Neuron
66:114–125.

Colgin LL, Denninger T, Fyhn M, Hafting T, Bonnevie T, Jensen O, Moser M-B, Moser EI (2009)
Frequency of gamma oscillations routes flow of information in the hippocampus. Nature
462:353–357.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics
including independent component analysis. J Neurosci Methods 134:9–21.

Edden RAE, Muthukumaraswamy SD, Freeman TCA, Singh KD (2009) Orientation Discrimination
Performance Is Predicted by GABA Concentration and Gamma Oscillation Frequency in
Human Primary Visual Cortex. J Neurosci 29:15721–15726.

Engbert R (2006) Microsaccades: a microcosm for research on oculomotor control, attention, and
visual perception. In: Progress in Brain Research (Martinez-Conde S, Macknik SL, Martinez
LM, Alonso J-M, Tse PU, eds), pp 177–192 Visual Perception. Elsevier. Available at:
http://www.sciencedirect.com/science/article/pii/S0079612306540099 [Accessed
November 12, 2017].

Gaetz W, Roberts TPL, Singh KD, Muthukumaraswamy SD (2012) Functional and structural correlates
of the aging brain: Relating visual cortex (V1) gamma band responses to age-related
structural change. Hum Brain Mapp 33:2035–2046.

Gray CM, König P, Engel AK, Singer W (1989) Oscillatory responses in cat visual cortex exhibit inter-
columnar synchronization which reflects global stimulus properties. Nature 338:334–337.

Gregoriou GG, Gotts SJ, Zhou H, Desimone R (2009) High-Frequency, Long-Range Coupling Between
Prefrontal and Visual Cortex During Attention. Science 324:1207–1210.

He BJ (2014) Scale-free brain activity: past, present, and future. Trends Cogn Sci 18:480–487.

He BJ, Zempel JM, Snyder AZ, Raichle ME (2010) The Temporal Structures and Functional Significance
of Scale-free Brain Activity. Neuron 66:353–369.

Hirano Y, Oribe N, Kanba S, Onitsuka T, Nestor PG, Spencer KM (2015) Spontaneous Gamma Activity
in Schizophrenia. JAMA Psychiatry 72:813–821.

Iaccarino HF, Singer AC, Martorell AJ, Rudenko A, Gao F, Gillingham TZ, Mathys H, Seo J, Kritskiy O,
Abdurrob F, Adaikkan C, Canter RG, Rueda R, Brown EN, Boyden ES, Tsai L-H (2016) Gamma
frequency entrainment attenuates amyloid load and modifies microglia. Nature 540:230.

Jia X, Tanabe S, Kohn A (2013) Gamma and the Coordination of Spiking Activity in Early Visual Cortex.
Neuron 77:762–774.

Lemaitre H, Goldman AL, Sambataro F, Verchinski BA, Meyer-Lindenberg A, Weinberger DR, Mattay
VS (2012) Normal age-related brain morphometric changes: nonuniformity across cortical
thickness, surface area and gray matter volume? Neurobiol Aging 33:617.e1-617.e9.

Mably AJ, Colgin LL (2018) Gamma oscillations in cognitive disorders. Curr Opin Neurobiol 52:182–
187.

Mitra P, Bokil H (2008) Observed brain dynamics. Oxford ; New York: Oxford University Press.

Murty DVPS, Shirhatti V, Ravishankar P, Ray S (2018) Large Visual Stimuli Induce Two Distinct
Gamma Oscillations in Primate Visual Cortex. J Neurosci 38:2730–2744.

Muthukumaraswamy SD, Edden RAE, Jones DK, Swettenham JB, Singh KD (2009) Resting GABA
concentration predicts peak gamma frequency and fMRI amplitude in response to visual
stimulation in humans. Proc Natl Acad Sci 106:8356–8361.

Muthukumaraswamy SD, Liley DTJ (2018) 1/f electrophysiological spectra in resting and drug-
induced states can be explained by the dynamics of multiple oscillatory relaxation processes.
NeuroImage 179:582–595.

Muthukumaraswamy SD, Singh KD, Swettenham JB, Jones DK (2010) Visual gamma oscillations and
evoked responses: Variability, repeatability and structural MRI correlates. NeuroImage
49:3349–3357.

Owsley C (2011) Aging and vision. Vision Res 51:1610–1622.

Palop JJ, Mucke L (2016) Network abnormalities and interneuron dysfunction in Alzheimer disease.
Nat Rev Neurosci 17:777–792.

Pantazis D, Fang M, Qin S, Mohsenzadeh Y, Li Q, Cichy RM (2018) Decoding the orientation of
contrast edges from MEG evoked and induced responses. NeuroImage 180:267–279.

Pesaran B, Pezaris JS, Sahani M, Mitra PP, Andersen RA (2002) Temporal structure in neuronal
activity during working memory in macaque parietal cortex. Nat Neurosci 5:805–811.

Podvalny E, Noy N, Harel M, Bickel S, Chechik G, Schroeder CE, Mehta AD, Tsodyks M, Malach R
(2015) A unifying principle underlying the extracellular field potential spectral responses in
the human cortex. J Neurophysiol 114:505–519.

Ray S, Maunsell JHR (2015) Do gamma oscillations play a role in cerebral cortex? Trends Cogn Sci
19:78–85.

Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RSR, Busa E, Morris JC, Dale AM, Fischl B (2004)
746   Thinning of the Cerebral Cortex in Aging. Cereb Cortex 14:721–730.

747   Sheehan TC, Sreekumar V, Inati SK, Zaghloul KA (2018) Signal Complexity of Human Intracranial EEG
748   Tracks Successful Associative-Memory Formation across Individuals. J Neurosci 38:1744–
749   1755.

750   Shirhatti V, Borthakur A, Ray S (2016) Effect of Reference Scheme on Power and Phase of the Local
751   Field Potential. Neural Comput 28:882–913.

752   Sohal VS, Zhang F, Yizhar O, Deisseroth K (2009) Parvalbumin neurons and gamma rhythms enhance
753   cortical circuit performance. Nature 459:698–702.

754   Tada M, Nagai T, Kirihara K, Koike S, Suga M, Araki T, Kobayashi T, Kasai K (2014) Differential
755   Alterations of Auditory Gamma Oscillatory Responses Between Pre-onset High-risk
756   Individuals and First-episode Schizophrenia. Cereb Cortex:bhu278.

757   Uhlhaas PJ, Singer W (2007) What Do Disturbances in Neural Synchrony Tell Us About Autism? Biol
758   Psychiatry 62:190–191.

759   Vaden RJ, Hutcheson NL, McCollum LA, Kentros J, Visscher KM (2012) Older adults, unlike younger
760   adults, do not modulate alpha power to suppress irrelevant information. NeuroImage
761   63:1127–1133.

762   Veit J, Hakim R, Jadi MP, Sejnowski TJ, Adesnik H (2017) Cortical gamma band synchronization
763   through somatostatin interneurons. Nat Neurosci 20:951–959.

764   Verret L, Mann EO, Hang GB, Barth AMI, Cobos I, Ho K, Devidze N, Masliah E, Kreitzer AC, Mody I,
765   Mucke L, Palop JJ (2012) Inhibitory Interneuron Deficit Links Altered Network Activity and
766   Cognitive Dysfunction in Alzheimer Model. Cell 149:708–721.

767   Voytek B, Kramer MA, Case J, Lepage KQ, Tempesta ZR, Knight RT, Gazzaley A (2015) Age-Related
768   Changes in 1/f Neural Electrophysiological Noise. J Neurosci 35:13257–13265.

769   Wilson TW, Rojas DC, Reite ML, Teale PD, Rogers SJ (2007) Children and Adolescents with Autism
770   Exhibit Reduced MEG Steady-State Gamma Responses. Biol Psychiatry 62:192–197.

771   Yuval-Greenberg S, Tomer O, Keren AS, Nelken I, Deouell LY (2008) Transient Induced Gamma-Band
772   Response in EEG as a Manifestation of Miniature Saccades. Neuron 58:429–441.

 Figure legends

Figure 1. Slow and fast gamma in an example elderly subject. a) Trial-averaged EEG trace

(1st row, blue); time-frequency spectrograms of raw power (2nd row) and change in power from

baseline (3rd row); and change in power with time (4th row) in alpha (8-12 Hz, violet), slow

(20-34 Hz, pink) and fast gamma (36-66 Hz, orange) bands averaged across 10 unipolar (left

column) and 9 bipolar (right column) electrodes. Vertical dashed lines represent actual stimulus

duration (0-0.8 s, black) and period used for analysis within stimulus duration (0.25-0.75 s,

red). Horizontal lines represent baseline (-0.5-0 s, black) and stimulus (0.25-0.75 s, red)

analysis periods. White lines in spectrograms represent slow (solid) and fast (dashed) gamma

frequency ranges. b) Right ordinate shows raw power spectral densities (PSDs, black traces)

vs frequency in baseline (dotted) and stimulus (solid) periods averaged across 10 unipolar

electrodes (left column) and 9 bipolar (right column) electrodes; left ordinate shows the same

for change in PSD (in dB, solid blue trace) in stimulus period from baseline. Solid pink lines

and dashed orange lines represent slow and fast gamma bands respectively. c) Scalp maps

showing 112 bipolar electrodes (represented as disks). Colour of each disk represents change

in slow (left) and fast (right) gamma power. 9 electrodes used in 1a and 1b (right column) are

marked with dots.

Figure 2. Baseline PSDs, slopes and alpha power. Baseline PSDs (averaged across 10

unipolar or 9 bipolar electrodes) for three age-groups on a log-log scale for unipolar (left) and

bipolar (right) reference, plotted for males (2a) and females (2b). Thickness of traces indicate

SEM across subjects. Age-group limits and the number of subjects in the respective age-groups

are indicated on the left plot. c) Same as in 2a and 2b, but for males and females, pooled across

all age-groups. d) Mean baseline PSDs for three ranges of baseline alpha power (8-12 Hz,

power ranges for respective traces indicated on the plots) pooled across all age-groups.

Thickness of traces and numbers indicate SEM across subjects and number of subjects in

respective alpha power ranges. Coloured bars on the abscissa indicate alpha (8-12 Hz, violet),

slow (20-34 Hz, pink) and fast gamma (36-66 Hz, orange) frequency bands.

Figure 3. Slow and fast gamma in younger and elderly subjects. a) Scatter plot showing

change in slow (abscissa) and fast (ordinate) gamma power. Dotted lines represent 0.5 dB

threshold. Points represent subjects with no gamma (dark blue), only slow gamma (light blue),

only fast gamma (green) and both gamma rhythms (yellow) with change in power above 0.5

dB threshold. b) Change in PSDs vs frequency averaged across subjects (numbers denoted by

n) as categorized in 3a. Thickness of traces indicate SEM. Solid pink and dashed lines represent

slow and fast gamma ranges respectively. c) Bar plot showing percentage of subjects in three

age-groups (marked by respective colours) categorized as in 3a. d) Schematic showing

placements of left and right anterolateral and posteromedial group of electrodes used for

analysis on the scalp, as well as ground and reference electrodes. e) Average scalp maps of 112

bipolar electrodes (disks) for three age-groups for slow (top row) and fast (bottom row) gamma.

Colour of disks represents change in respective gamma power. Electrode groups represented

as in 3d.

Figure 4. Change in gamma power vs age for anterolateral group of electrodes. Mean

time-frequency change in power spectrograms (4a) and change in power spectra vs frequency

(4b) for three age-groups separately for males (top row) and females (bottom row). Thickness

of traces and numbers in 4b indicate SEM and number of subjects respectively. Solid and

dashed lines indicate slow and fast gamma frequency ranges respectively. c) Left column: bar

plots showing mean change in slow gamma power for three age-groups separately for males and females. Number of subjects for respective age-groups are indicated on top. Error bars indicate SEM. Right column: scatter plot for change in slow gamma power vs age for all elderly subjects (>49 years age-group, n=227), plotted separately for males (in orange) and females (in yellow). Orange, yellow and black solid lines indicate regression fits for males, females and data pooled across gender respectively. p-values of the regression fits are indicated in respective colours. d) Same as in 4c but for fast gamma.

Figure 5. Centre frequency of slow and fast gamma vs age for elderly subjects. a) Change in power vs frequency for 227 elderly subjects arranged in ascending order of age. Age for each subject is indicated on ordinate. Colour represents change in power in dB for each frequency. Circles and triangles represent centre frequency of slow and fast gamma rhythms respectively, indicated only for those subjects who have change in power in respective gamma range above 0.5 dB. b) Scatter plots showing centre frequency vs age for slow (top row) and fast (bottom row) gamma for those subjects who have change in power in respective gamma range above 0.5 dB (numbers indicated on the plots). Solid lines in both 5a and 5b are regression fits for centre frequency vs age. p-values for these fits are indicated on plots in 5b. Left and right columns show analysis for anterolateral and posteromedial group of electrodes respectively.

Figure 6. Change in alpha power vs age. a) Left column: bar plots showing mean change in alpha power across anterolateral group of electrodes for three age-groups separately for males and females. Number of subjects for respective age-groups are indicated at bottom. Right column: scatter plot for change in alpha power vs age for all elderly subjects (>49 years age-

group, n=227), plotted separately for males (in orange) and females (in yellow). Same format

as in Figure 4c. b) Scalp maps for 112 electrodes (disks) averaged across all subjects separately

for three age-groups. Colour indicates change in alpha power for each electrode, same format

as in Figure 3e.

Figure 7. Eye position, microsaccades and pupillary reactivity across age for elderly

subjects. a) Eye-position in horizontal (top row) and vertical (middle row) directions; and

histogram showing microsaccade rate (bottom row) vs time (-0.5-0.75 s of stimulus onset) for

elderly subjects (n=226). Number of subjects in each age-group is indicated on top. Thickness

indicates SEM. b) Main sequence showing peak velocity and maximum displacement of all

microsaccades (number indicated by n) extracted for both elderly age-groups. Average

microsaccade rate (mean±SEM) across all subjects for each elderly age-group is also indicated.

c) Scatter plot showing change in power vs age for slow (top row) and fast (bottom row) gamma

for all elderly subjects with analysable data after removal of trials containing microsaccades.

Solid lines indicate regression fits. Numbers of subjects with analysable data in each age-group

is indicated on top. d) Scatter plots for coefficient of variation (CV) of pupil diameter vs age

(top row), change in slow (middle row) and fast (bottom row) gamma power. Pearson

correlation coefficients (r) and p-values are also indicated.

Figure 8. Change in SSVEP power vs age for posteromedial group of electrodes. Time-

frequency change in power spectrograms (8a) and change in power spectra vs frequency (8b)

for three age-groups separately for males (top row) and females (bottom row). Thickness of

traces in 8b indicates SEM. Insets in 8b display zoomed-in images of respective main plots,

showing clear SSVEP peaks at 32 Hz. c) Top row: bar plots showing mean change in SSVEP

power for three age-groups separately for males and females; Numbers of subjects in each age-group is indicated on top. Error bars indicate SEM. Bottom row: scatter plot for change in SSVEP power vs age for all elderly subjects (>49 years age-group, n=197), plotted separately for males (in orange) and females (in yellow). Orange, yellow and black solid lines indicate regression fits for males, females and data pooled across gender respectively. p-values of the regression fits are indicated in respective colours. d) Scalp maps for 112 electrodes (disks) averaged across all subjects separately for three age-groups. Colour indicates change in SSVEP power at 32 Hz for each electrode.

Supplementary Figure 1. PSD slopes (16-44 Hz) across age-groups, gender and baseline alpha power

a) Bar plots showing mean slopes of baseline PSDs for three age-groups separately for males and females in 16-44 Hz range for unipolar (left) and bipolar (right) reference. Error bars indicate SEM. Numbers on top indicate number of subjects. b) Scatter plot showing baseline slopes in 16-44 Hz range vs alpha power, for unipolar (left) and bipolar (right) reference for all 274 subjects used for analysis (227 elderly and 47 younger). Lines indicate regression fits. Pearson correlation coefficients (r) and p-values are also indicated.

Supplementary Figure 2. Change in gamma power vs age for posteromedial group of electrodes. Same format as in Figure 4.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

b

Figure 6

Figure 7

Figure 8

Supplementary Figure 1

a

b

Supplementary Figure 2