Genomic signature of shifts in selection in a sub-alpine ant and its physiological adaptations

Cicconardi Francesco[1], Krapf Patrick[2], D'Annessa Ilda[3], Gamisch Alexander[2,4], Wagner Herbert C[2], Nguyen Andrew D[5], Economo Evan P[6], Mikheyev Alexander S[7], Guénard Benoit[8], Grabherr Reingard[9], Arthofer Wolfgang[2], di Marino Daniele[10], Steiner Florian M[2], Schlick-Steiner Birgit C[2]

[1] Department of Zoology, University of Cambridge, Cambridge, UK
[2] Department of Ecology, University of Innsbruck, Innsbruck Austria
[3] Istituto di Chimica del Riconoscimento Molecolare, CNR (ICRM-CNR), Milan, Italy
[4] Department of Biosciences, University of Salzburg, Salzburg, Austria
[5] Department of Entomology and Nematology, University of Florida, USA
[6] Biodiversity & Biocomplexity Unit, Okinawa Institute of Science & Technology, Onna, Japan
[7] Ecology and Evolution Unit, Okinawa Institute of Science & Technology, Onna, Japan
[8] School of Biological Sciences, The University of Hong Kong, Hong Kong, China
[9] Institute of Biotechnology, University of Natural Resources and Life Sciences, Vienna, Austria
[10] Polytechnic University of Marche, Department of Life and Environmental Sciences, Ancona, Italy
Corresponding author: francicco@gmail.com

Abstract

Understanding how organisms adapt to extreme environments is fundamental and can provide insightful case studies for both evolutionary biology and climate-change biology. Here, we take advantage of the vast diversity of lifestyles in ants to identify genomic signatures of adaptation to extreme habitats such as high altitude. We hypothesised two parallel patterns would occur in a genome adapting to an extreme habitat: *i*) strong positive selection on genes related to adaptation and, *ii*) a relaxation of previous purifying selection. We tested this hypothesis by sequencing the high-elevation specialist *Tetramorium alpestre* and four related species. In support of our hypothesis, we recorded a strong shift of selective forces in *T. alpestre*. We further disentangled candidate molecular adaptations in both gene expression and protein-coding sequence that were identified by our genome wide analyses. In particular, we demonstrate that *T. alpestre* has *i*) a derived level of expression for *stv* and other heat-shock proteins in chill shock tests, and *ii*) enzymatic enhancement of Hex-T1, a rate-limiting regulatory enzyme that controls the entry of glucose into the glycolytic pathway. Together, our analyses highlight the adaptive molecular changes that support colonisation of high altitude environments.

Introduction

Adaptation of organisms to climate drives variation in diversification rates and species richness among clades. Their climatic niche affects a species' distribution in space and time [1], critically affecting both speciation and extinction. Therefore, understanding how organisms manage thermal adaptation is important in terms of both evolutionary biology and climate-change biology, considering the potential world-wide loss of ecological niches [2,3]. In this context, high elevations, characterized by a short growing season and low annual minimum and mean temperatures, with high daily fluctuation temperatures [4], are an important open-air laboratory to study speciation and adaptation to cold habitats; and compared to the great effort in gene-based studies on cold tolerance of model organisms, a lesser effort has been put towards understanding truly cold-tolerant animals at the genomic level [5,6]. This is even more evident when considering that among the approximately 600 sequenced insect genomes available today only four belong to species ecologically restricted to high altitudes or Antarctic habitats [7–10].

As yet, potential patterns of genomic signatures in organisms adapting to more extreme habitats, such as high elevations, have not been identified and, more generally, there is no theory that predicts the rates of genomic change for extreme habitats. Here, we hypothesise two parallel patterns to occur in a genome adapting to an extreme habitat. (1) Strong positive selection on genes related to adaptation, such as genes involved in metabolic pathways; and (2) Relaxation of previous selecting forces, as the conditions of the previous niche are lacking in the new niche. The latter should lead to a reduced number and/or a different set of genes under purifying selection. Specifically, in the case of high-elevation habitats, some heat-

shock proteins (HSPs), necessary for coping with extreme heat, should be under

relaxation, because heat resistance is not strongly selected for in the average alpine

species. A problem with the hypothesis of these two parallel patterns is that there is

no direct test available for the functional consequences of the loss or gain of a

specific gene. There are often many pleiotropic effects for genes that make

interpretations difficult, and HSPs, expressed upon exposure to stress or during

development and growth, are likely to fall into this category [11]. Therefore, we

expect strong positive selection in genes involved in energetic metabolism, and an

increase in the relaxation rates in other genes, without being able to specify which

these might be. Identifying such genes will help to set up hypotheses to test in the

future. A comparative approach offers the strongest method for testing our

hypothesis, by comparing genomes of species closely related to each other, but

divergent in terms of adaptation to different environments. However, this approach

can be limited by insufficient niche divergence within a group or the number of

genomes sequenced. Ants are emerging as a leading system for comparative

genomics due to the constantly increasing number of available genomes. This

resource offers the opportunity to improve the accuracy of orthology detection [12],

to scan for specific mutations, candidate genes, and patterns of acceleration and

relaxation in the ant genomes associated with adaptation to cold habitats.

Ants are key species in the Earth's terrestrial ecosystems. They comprise

more than 15,442 described species and display an impressive diversity of lifestyles

[13–15]. Ants are especially notable among insects for their ecological dominance as

predators, scavengers, and indirect herbivores. They compose at least one third of

the entire insect biomass [16] and colonise all kinds of habitats, including

thermobiologically challenging environments. The formicine ant *Melophorus bagoti*,

for instance, is active during the hottest periods of the summer day, when air

temperatures at ant height exceed 50 ˚C [17]. On the other hand, for example, the

myrmicine ant *Tetramorium alpestre* inhabits the montane and subalpine belt of the

Central and South European mountain systems, with the Alps as its main distribution

area [18,19]. This species lives mainly between 1300 and 2300 m above sea level

(a.s.l.), forages below the ground, and nests are established in cool grassland under

stones, in moss, rootage, and dead wood, especially subalpine and alpine grass mats

[15]. Since *T. alpestre* belongs to a species complex in which no relation between the

ecological niche and worker morphology has been detected, searching for adaptive

phenotypical differences outside of morphology, for example physiology, has already

been suggested [18]. *Tetramorium alpestre* is also relevant because, among the 79

Palearctic species of the worldwide distributed genus *Tetramorium* (Bolton 2016), it

is one of the only two species to form colonies with multiple queens (polygynous

colonies), showing intermediate states of aggression and a transition from

multicoloniality to supercoloniality [15,20]. *Tetramorium alpestre* is thus a system

well suited for studying both genomic adaptation to high elevation and socio-

behavioural evolution.

Here, we tested our hypothesis of both increased diversifying and reduced

purifying (relaxing) selection resulting from adaptation to an extreme niche using *T.*

alpestre. We newly sequenced its genome and those of four related *Tetramorium*

species with diverging ecological niches: *T. immigrans*, a species of the *T. caespitum*

complex to which *T. alpestre* also belongs to, in sympatry with *T. alpestre*, but

distinct in its altitudinal and ecological habitat; *T. parvispinum*, restricted to

mountain forest habitats of the Austral-Asian and Indo-Malayan subregions [21]; and

T. bicarinatum and *T. simillimum*, invasive generalist species which occur in warmer

habitats [22,23]. Our goals were to define the ecological niche of these ant species

based on environmental data, and subsequently to perform gene family

expansion/contraction and protein-coding scans for signatures of diversifying and

relaxing selection. We also annotated the five HSP subfamilies for 19 ant species to

test for possible shifts in selection acting on these gene families. Among the

signatures of diversifying selection detected, we performed two experiments in *T.*

alpestre and its relative, *T. immigrans*: we assessed (i) chill-shock triggered gene-

expression patterns of *starvin* (*stv*), a modulator the activity of Hsp70 chaperone

machinery during recovery from cold stress, and certain HSPs involved in recovering

from extreme cold, and (ii) the temperature dependence of the enzyme activity of

Hexokinase type 1 (Hex-T1), a rate-limiting and regulatory enzyme that controls the

entry of glucose into the glycolytic pathway, one of the most conserved and

essential hexokinase isoenzymes [24]. The genomic data presented here contribute

to a foundation for studying insect genome evolution, particularly also in the light of

climate change.

Figure 1. *a*) Venn diagram displaying overlap in orthologous genes in the five *Tetramorium* spp. + *V. emeryi* (Crematogastrini). Coloured circles with number represent phylogenetic splits in the nuDNA (N^{nu}) and mitochondrial phylogenies in *b*; *b*) The dated Crematogastrini nuclear phylogeny (above). Branch colour are based on the concordance factors of the two datasets (nuDNA and mtDNA). On each branch, the numbers of expanded (green arrows) or contracted (purple arrows) genes is shown, inferred from observed OGs sizes at terminal branches. In the bottom, the global $\delta^{18}O$ (‰) derived from analyses of two common and long-lived benthic taxa are given, *Cibicidoides* and *Nuttallides,* which reflect the global deep-sea oxygen and carbon isotope and thus the temperature (from Zachos et al., 2001). Bars represent the 95% confidence interval for the BI analysis, circles with numbers the phylogenetic splits in the nuDNA (N^{nu}) and mtDNA (N^{mt}) phylogenies. *c*) Boxplots of the five bioclimatic variables significantly different for *T. alpestre* and the other ant species.

Results

Climatic niches

To define the environment of *T. alpestre* and the other four *Tetramorium* species, a dataset contains 1835 localities representing 17 species was assembled, using georeferenced occurrence data compiled from the Global Ant Biodiversity Informatics database (see method section), with the number of localities per species ranging from 9 to 591 (Table S1). Of the 19 bioclimatic variables, four (bio1: Annual Mean Temperature, bio5: Max Temperature of Warmest Month, bio8: Mean Temperature of Wettest Quarter, bio10: Mean Temperature of Warmest Quarter)

differentiated between *T. alpestre* and other ants (adjusted *P*-values < 0.001) (Figure

1a). In detail, *T. immigrans* and *T. alpestre* occupy habitats that are colder in the

growing season than those of the remaining species, with *T. alpestre* revealing even

colder habitats compared with *T. immigrans*.

Sequencing, assembly, and annotation of the *Tetramorium* genomes

The combination of overlapping paired-end libraries with very high coverage,

standard short paired-end libraries, and mate pair libraries resulted in an overall

complete assembly of the five *de novo Tetramorium* genomes, at the level of both

contiguity and scaffolding. In particular, the *T. alpestre* genome contiguity (contig

N50) is the 3^rd^ best among all ant genomes available (Table S2). The analysis of the *k*-

mer frequency distribution generated was unimodal (Figure S1; Supplementary

Material online), and an analysis of the distribution of GC content across the five *de*

novo Tetramorium genomes revealed similar distributions with a mean of 38%, that

is, slightly less than that found in *V. emeryi* (42%) (Figure S2). No significant bias was

recovered considering GC content and coverage (Figure S3). All five *de novo* draft

assemblies were ~241 Mb in length, comparable in size with the average of all ant

genomes available (~278 Mb) (Table S2). The total *T. alpestre* genome size was

245.72 Mb, including 18 Mb of gaps and unknown characters (N/X), slightly smaller

than the estimated genome size by flow-cytometry of 291.84 Mb +/- 1.76 Mb (n=12).

Repetitive elements made up similar proportions of each sequenced genome (*T.*

alpestre⬜=⬜18%, *T. immigrans*⬜= 20%, *T. bicarinatum* = 21%, *T. simillimum*⬜=⬜20%,

T. parvispinum = 23%; Table S2), indicating a strong correlation between genome

size and total interspersed repeat content (Pearson ρ = 0.976, *P*-value < 0.005; Figure

S5). For the gene annotations in *T. alpestre*, a combinatorial approach of

unsupervised RNA-seq-based, homology-based, *ab initio*, and finally *de novo*

methods were used. By doing so, we approached the number of predicted protein-

coding genes in the phylogenetically closest annotated species (15,085 genes

predicted vs. 14,872 in the *V. emeryi* genome, Table S2). This improved the

homology-based annotation of the remaining four *Tetramorium* genome assemblies

that did not differ much in terms of estimated gene content, ranging from 16k to 15k

annotated loci (Table S2). BUSCO analyses estimated overall good representation,

with recovered genes in between the 97.6% and 99.9% (Table S2; Figure S6), and all

statistics on mRNA, coding regions (CDS), exon, and intron length gave highly similar

and overlapping distributions among all *Tetramorium* species and *D. melanogaster*

as a reference (Figure S7). Also, the transcript completeness gave good results with a

comparable distribution of percentage of alignment across species (Figure S8).

The orthology search analysis produced a total of 7195 OGs between *D.*

melanogaster and Hymenoptera; 3261 of these represented scOGs present in all 22

species analysed. Restricting the orthology analysis to *Tetramorium* spp. and *V.*

emeryi resulted in 6494 OGs; among *Tetramorium*, the highest fraction was

recovered between *T. alpestre* and *T. immigrans* with 729 OGs, followed by *T.*

simillimum and *T. parvispinum* (540 OGs) (Figure 1a). For all *Tetramorium* species.,

the fraction of orthologous genes with at least a match with one of the three

outgroups (*A. mellifera*, *N. vitripennis*, *D. melanogaster*) was on average 7286, with

the highest in *T. alpestre* (7879) and the lowest in *T. parvispinum* (6977), in line with

the average value found in ants (7831).

Assembly and annotation of ant mitogenomes and their gene translocations

We assembled complete mtDNA genome sequences from 24 ant species. Because of the differences among SRA datasets, we used different numbers of reads to assemble the complete sequences. The new mtDNA genomes were slightly longer (on average, 17k vs. 16k; Table S2), probably due to the reconstruction of the D-loop and ribosomal genes, which are sometimes harder to assemble. We slightly mis-assembled a coding gene (*nad2*) in the *T. bicarinatum* genome and a few other tRNA loci in other genomes (Table S2). The average GC-content for coding genes was 23.9 ± 3.0% and varied slightly among taxa with the lowest in *Formica fusca* (19.0%) and the highest in *Leptomyrmex pallens* (32.8%) (Table S2). While almost all species had the typical gene order and orientation, some rearrangements were observed. All myrmicine ants were characterized by a translocation of *trnV* after *rrnS*; *Solenopsis* spp. had a swapped position of *rrnS* and *trnN*; *Monomorium pharaonis* had at least three rearrangements: the inversion of *nad6*, *cob*, and *trnS2*, the translocation of *trnE* upstream to *trnA*, and the translocation of *trnR* upstream to *trnE*; *Tetramorium* spp. underwent a swap of *trnR* and *trnN* (Figure S9).

Phylogenetic analyses of *Tetramorium* spp., their diversification dating, and the Miocene-Pliocene origin of the lineage leading to *T. alpestre*

The estimated divergence ages from both datasets (mtDNA, nuDNA) were congruent for most nodes, except for recent nodes within the Attini (Figure 1b; Figure S10-13). In this case, the penalized likelihood approach estimates were always more recent than the estimations made with BI using BEAST. The dating of basal nodes like the MRCAs (e.g., Crematogastrini, Myrmicinae, and Formicinae) were congruent with a

previous analysis based on more taxa but fewer genes [26]. The two methods also

converged regarding the date of the most recent common ancestor (MRCA) between

T. alpestre and *T. immigrans*. In that specific case, PL dated that split at 6.8 million

years ago (mya) within the confidence interval given by BI (Median: 5.15 mya; 95%

HPD: 2.66, 8.23).

The phylogenetic relationships were generally highly congruent and with

predominantly strong support (Figure 1b and Figures S10-13). In the ML phylogeny

of scOGs (nuDNA), all nodes were supported with a bootstrap value of 1. Across the

whole phylogeny, only a few nodes had weak signals, resulting in very few

topological disagreements and low coalescent units (CU). Within the clade of

Crematogastrini (*Tetramorium* spp. + *V. emeryi*), the relation of *T. bicarinatum* and *T.*

simillimum was uncertain: nuDNA phylogeny recovered them as sister species with

low CU (0.11), while mtDNA phylogeny placed *T. bicarinatum* as sister to the *T.*

alpestre-*T. immigrans* cluster, with a bootstrap value of 0.77 and a posterior

probability of 0.99.

Contractions of OGs in the *T. immigrans* and *T. alpestre* lineages and expansion of

retrotransposon-related genes in *T. alpestre*

We examined the evolutionary dynamics of OGs looking at more dynamic ranges

corresponding with terminal branches. We observed an increased number of OG

deaths in the *T. parvispinum* lineage (− 642) and in the lineage leading to *T. alpestre*

and *T. immigrans* (− 516). Ortholog groups showed very little expansion in internal

branches of the *Tetramorium* radiation. Expanded OGs were common in terminal

branches, except for *T. parvispinum* with very few expanded OGs (+ 14), suggesting a

possible lack of annotation. *Tetramorium alpestre* showed the highest number of

expanded OGs (+ 364) (Figure 1b), with more than 370 of them (~ 40%) belonging to

genes with reverse transcriptase (PFAM: rve, rve_1, rve_2) or transposase and

retrotransposon activities (PFAM: Retrotrans_gag, Retrotran_gag_2). Among other

PFAM, haemolymph juvenile hormone-binding proteins (PFAM: JHBP; 20 genes), zinc

fingers (PFAM: zf-CCHC, zf-H2C2, zf-met, 32 genes), and sugar transporters (PFAM:

Sugar_tr, 4 genes) were found.

257 Dual signature of evolutionary pressures in single-copy OGs in the *T. alpestre*

258 genome

We investigated the impact of evolutionary pressures on one-to-one orthology in the

five *Tetramorium* spp. and 14 other ant species and three outgroups (*Apis mellifera*,

Nasonia vitripennis, and *Drosophila melanogaster*) by computing both mean ω

(d_N/d_S) for terminal branches and relaxing selection (k) within the Crematogastrini.

The measure of the mean branch ω showed an overall high level of purifying

selection along the whole ant phylogeny, including for short branches [27,28], with

median values of 0.10. However, this distribution was significantly shifted in *T.*

alpestre, which showed a median value of 0.38 (Wilcoxon rank-sum test '*greater*', P-

value < $2.2e^{-16}$), whilst keeping the proportion of $\omega > 1$ in each branch unchanged

(Figure 2a, S13). The evaluation of k in the five *Tetramorium* spp. and *V. emeryi* had a

mean of median values of 0.99, that is, twice the value in *T. alpestre* (median k =

0.47; Wilcoxon rank-sum test '*less*', P-value < $2.2e^{-16}$) (Figure 2b). We investigated

the nature of the increased ω (episodic diversifying selection) and decreased k

(relaxed selection) in *T. alpestre* by combining the two measures. Two evolutionary

trajectories appeared to act simultaneously (two tails of the distribution, Figure 2c):

One trajectory seems to have led to episodic diversifying selection,

Figure 2. *a*) Boxplots showing the distribution and median (vertical line) of mean ω (d_N/d_S) rates (square shapes)
in terminal branches of *Tetramorium* spp. and *V. emeryi* (Crematogastrini) in scOGs, and the proportion of genes
for which ω is higher than one. *b*) Boxplots overlapped by violin plots showing the distribution of k in
Crematogastrini. The distribution in *T. alpestre* is bimodal as in *T. immigrans*, but with a much more skewed
distribution towards zero values. In both sections, asterisks indicate the degree of significance between *T.*
alpestre and the other species (Wilcoxon rank-sum tests). *c*) Scatter plots for each Crematogastrini species
included, showing for each scOG branch both values of ω and k. The green dots are scOGs with uncorrected *P*-
values associated with diversifying selection less than 0.05. The distribution in *T. alpestre* shows the longest tail

corresponding to very low values of *k* and high values of *ω*. It is also notable a much wider and scattered
distribution of scOGs with high *k* and *ω* compared with all other species.

promoting the fixation of nonsynonymous mutations with presumably advantageous

fitness effects for genes with increased values of *ω* and *k*, also present in the other

species. The other trajectory showed a relaxation of the overall purifying selection in

genes with increased values of *ω* and decreased values of *k*, only mildly present in *T.*

immigrans.

Figure 3. For each enriched Gene Ontology (GO) term category *a*) the median values of *ω*; and *b*) *k* for genes
enriching the specific category. *Tetramorium* spp. are colour coded, the other 14 species, merged to obtain a
background signal, are in gray. Blue circles are scOGs tested for each GO term, red circles are scOGs returned as
significant. Circle sizes are proportional to the number of genes. To be noted are the overall extremely higher

values of ω and lower values of k in *T. alpestre* branches compared to the other species. Seldomly also *T.*
immigrans shows few GO terms with high ω and low k values, in particular the Apical junction assembly
(GO:0043297) shows and inverted trend between *T. alpestre* and *T. immigrans*.
299

After a stringent correction for multiple testing, we found 175 scOGs with a

putative signature of diversifying selection (adjusted *P*-values < 0.005; Table S5),

with 50 genes enriching 13 biological process terms (*P*-values < 0.005; Figure 3a,

Table S6), nine related to cell development (e.g., cell migration involved in

gastrulation, photoreceptor cell development, cell fate determination) and

organization (e.g., regulation of organelle organization), two related to signalling

pathways (enzyme linked receptor protein signalling pathway and torso signalling

pathway), one to heart development, and one to organic hydroxy-compound

metabolic processes. The most representative genes were *Pten* and *Gbeta13F*,

involved in six biological processes, *DCTN1-p150* and *dsh*, involved in five of them. All

key genes are involved in the proliferation and division of cells, especially

neurological stem cells (*Gbeta13F*), respiratory system development (*Pten* and *dsh*),

and the serine/threonine-protein kinase *polo*, present in four enriched Gene

Ontology (GO) terms. In the catalytic domain of this protein (255 aa), we found five

amino acid substitutions compared with *T. immigrans*, three of them corresponding

to active sites (*TalpPolo$_{G29V}$*, *TalpPolo$_{G30R}$*, *TalpPolo$_{S104R}$*). We also tested for possible

enrichment in KEGG pathways and found three significant pathways (*P*-values < 0.02;

Table S7), all involved in the metabolism of sugars, specifically: glycolysis, pentose

phosphate pathway and galactose metabolism. Within these pathways, five enzymes

were found under diversifying selection: a glucose-6-phosphate (*CG9008*), the

hexokinase type 1 (*Hex-T1*), a phosphofructokinase (*Pfk*), a transketolase (*CG8036*),

and a phosphoglucosemutase (*pgm*).

132 scOGs had a signature of putative relaxation (*P*-values < 0.005; Table S8).

While the absolute number of scOGs under relaxation was slightly lower than the

number of genes under diversifying selection, the gene-set enrichment tests

revealed many more biological processes significantly enriched: 70 terms (*P*-values <

0.005; Figure 3b, Table S9) enriched by 118 genes. Fifteen terms were related to the

regulation of cellular development, signal transduction, and cell communication

(e.g., regulation of response to stimuli, regulation of cell communication, regulation

of Ras signal transduction), 12 terms related to neurogenesis, axon development,

and synaptic transmission (e.g., neuron differentiation, trans-synaptic signalling,

axon guidance), seven terms were related to cell morphogenesis and junction

organization, and many other related to response to stimuli (e.g., chemotaxis,

response to light) or related to adult development and morphogenesis (e.g.,

anatomical structure morphogenesis, molting cycle, chitin-based cuticle). The

enrichment of KEGG pathways revealed three pathways: mRNA surveillance

pathway, spliceosome, and protein processing in endoplasmic reticulum (*P*-values <

0.05; Table S10).

The Hex-T1 activity, its model structure, and molecular dynamics simulations in *T.

alpestre and *T. immigrans***

As we describe in the previous section, the enrichment for KEGG pathways led to the

identification of three metabolic pathways, all involved in the metabolism of sugars,

with five genes under selection. In order to experimental validate these *in silico*

results, we cloned and expressed the Hex-T1 forms for both *T. alpestre* and *T.*

immigrans into *Drosophila melanogaster* cells lines, in order to compare their

activities. The purified enzyme of both ants was assayed in at 6, 16, 26, and 36 °C to

establish their efficiency under different thermic conditions and identify possible

metabolic adaptations to cold-environments. The assays showed that the Km

(substrate concentration necessary to reach maximum reaction speed) of Hex-T1 of

T. alpestre was lower than the enzyme of *T. immigrans* at all temperatures.

351
Figure 4. *a*) Scatter plot of Km score across the temperature gradient in the two species, *T. immigrans* on top and
T. alpestre on the bottom. Lines represent average values with confidence intervals. Significances: *** < 0.001;
** < 0.001; * < 0.05. *b*) Overview of the homology-modeled structure of Hex-T1 in *T. alpestre* showing the two
Hexokinase domains, PFAM00349 (green) and PFAM03727 (pink). The lower panels show the location of TalpHex-
T1$_{G74}$ (left) and TimmHex-T1$_{E70}$ (right) as well as the arginine residue that forms a salt bridge with E70 (dashed
line). *c*) Per-Residue Root Mean Square Fluctuation (RMSF) calculated along 500 ns trajectories carried out at 6 °C
(upper panel) and 26 °C (lower panel) for TalpHex-T1 (left panel, blue line) and TimmHex-T1 (right panel, light

blue line). The location of the residues found mutated between the two sequences are indicated. *d*) Time
evolution of the atomic distance between the centre of mass of glucose and the lateral chain nitrogen atom of
TalpHex-T1$_{K168}$ (blue line) and TimmHex-T1$_{K164}$ (light blue line) in the binding pocket calculated along the 500 ns
trajectories carried out at 6 °C (upper panel) and 26 °C (lower panel). *e*) Representative configurations extracted
from the simulations at 26 °C showing the different orientation of TalpHex-T1$_{K168}$ and TimmHex-T1$_{K164}$, with the
first one being the only one able to bind the glucose (dashed line).

(Wilcoxon rank-sum test 'less', *P*-value < 0.00004, Figure 4a), with values ranging

from 0.08 mM to 0.01 mM in *T. alpestre* versus 0.15 mM to 0.05 mM in *T.*

immigrans, from the highest to the lowest temperature. These results demonstrate

that Hex-T1 of *T. alpestre* was always more efficient than *T. immigrans*, requiring less

substrate to reach maximum activity. Although the number of replicates per

condition was small, we performed statistical tests to attempt obtaining insight in

the enzymatic activity differences within species at different temperatures and in

their variance. Interestingly, it seems that for both species there is an improvement

in the Km values shifting towards the lowest temperatures (6 °C *vs* 16 °C; two-way

ANOVA followed by a Tukey's multiple comparison; adjusted *P*-values < 0.05), and a

possible deterioration at higher temperatures in *T. alpestre*, from 0.04±0.01 mM (26

°C) to 0.08±0.01 mM (36 °C) (two-way ANOVA followed by a Tukey's multiple

comparison; adjusted *P*-value < 0.0002). An F test (VAR.TEST() implemented in R) to

compare the variances of the enzymatic activities (Km values) between the two

species – by pooling the replicates at 16 and 26 °C – suggested that there is a

possible significant difference in their variance (*P*-value = 0.015), with the confidence

interval of Hex-T1 activity in *T. immigrans* twice wider as in *T. alpestre*.

The Hex-T1 enzyme is made of two structurally similar domains, Hexokinase 1

and 2 (PFAM00349 and PFAM03727), and an N-terminus that is highly variable both in

terms of length and amino acidic composition (Figure 4b). The two domains are

relatively well conserved across all ants, other Hymenoptera and *D. melanogaster*.

Excluding the hypervariable N-terminus, and considering the two functional

domains, the amino acid sequences of *T. alpestre* and *T. immigrans* only differ by

two amino-acidic substitutions: a glutamic acid (E) replaced by a glycine (G) in

position 74 (TalpHex-T1$_{E74G}$) (Figure 4b inserts) and an asparagine (N) replaced by a

lysine (K) in position 344 (TalpHex-T1$_{N344K}$). While the second change probably does

not entail a significant functionality effect, the *TalpHex-T1$_{E74G}$* instead may be

significant. Because of the lack of the long chain and the negative charge of glutamic

acid, this substitution could give the protein a higher flexibility and therefore an

overall higher kinetics. We tested this hypothesis by simulating the protein dynamics

in the presence of the glucose substrate at different temperatures, i.e. 6 and 26 °C,

in the two species, with the intent of mirroring the condition in the in vitro assay. In

the simulations we can see a dual effect. For TalpHex-T1$_{N344K}$, located at the C-

terminal of a α-helix forming a helix-turn-helix motif (VSETE**K**DPKG), resulted in a

lower degree of Root Mean Square Fluctuation (RMSF) revealed a lower degree of

fluctuation in TalpHex-T1 (Figure 4c). This is possibly due to the alternation of

positive and negative charges creates a salt-bridges network that stabilizes the

whole motif (data not shown). In particular TalpHex-T1$_{K344}$ is involved in a stable

interaction with TalpHex-T1$_{E341}$ (data not shown), while TimmHex-T1$_{N340}$ is not

available for the formation of salt bridges and mainly establishes hydrogen bonds

with TimmHex-T1$_{S336}$ or TimmHex-T1$_{E337}$ at 6 or 26 °C, respectively. Although the *T.*

alpestre salt bridge brings a higher flexibility of the helix-turn-helix motive, it does

not interfere with the catalytic site, given that this motif is far away in the C-terminal

region of the protein. In contrast, at the other region (TalpHex-T1$_{E74G}$), the mutation-

caused amino acid change allows the loop to explore a broader conformational

space equipping it with a higher degree of flexibility already at 6 °C (Figure 4c).

Nevertheless, by increasing the temperature, TalpHex-T1 was subjected to an even

higher degree of fluctuation, once again confirming the intrinsic flexibility led by the

glycine. Notably, the increased flexibility is also transferred to residues relatively far

from the mutation, in particular in regions near the active site (168-169), causing a

different pattern of interaction with the substrate (Figure 4d, e). Indeed, TalpHex-T1

established a stronger interaction with glucose at both temperatures with respect to

TimmHex-T1 (Figure 4e).

The heat-shock proteins in ants and their evolution in *T. alpestre*

Members of the five subfamilies of the heat shock proteins (HSPs: Hsp90s, Hsp70s,

Hsp60s, Hsp40s, and sHsps) were fully recovered and characterized in 19 ant species.

Overall, a strong purifying selection was observed within subfamilies (Figure 5a,

Table S11). In the Hsp90 gene family (*trap1*, *gp93*, and *Hsp83*) *Hsp83* was found

duplicated within the lineage of *W. auropunctata* (inparalog), as well as the *T.*

parvispinum gp93; duplication that probably occurred before emergence of the

lineage (outparalog) (Figure S14). Members of the Hsp70 family were recovered for

all species with 11 ortholog groups, with few duplications in *Hsc70-3* and *Hsc70-5*

(two inparalogs and two outparalogs, respectively). We confirmed the lack of Hsp70s

in ants [29] and a multiple duplication of *Hsc70-4* in Hymenoptera. One paralog

seemed to be lost in many species while the other two copies were still present in

almost all species, with the presence of other outparalogs in various species, mostly

in Attini (Figure S15). The chaperonin Cpn60/TCP-1 family (Hsp60) had nine OGs; six

inparalogs were found in *CCT7, CCT3* and four outparalogs in *Tcp1a, CCT8*. The heat

shock protein 40 kDa/Chaperone DnaJ was the largest subfamily, with 37 well-

defined OGs. Its turnover of duplication was relatively small, and only six OGs

showed some level of duplication. The small HSPs (sHsps) showed the expansion of

the lethal (2) essential for life (*l(2)efl*) gene paralog of *D. melanogaster*, with five

distinct OGs in all Hymenoptera. *L(2)efl3* seemed to be retained by only in 11

species, and with *l(2)efl4*, they are the two most divergent OGs, with ω equal to 0.23

and 0.33, respectively. The less divergent were *Hspb1* and *l(2)efl5*, for which ω was

0.11 and 0.07, respectively. Overall, the sHsps showed a significantly higher

distribution of ω (median = 0.22, *P*-value = 0.013), followed by Hsp40s, which also

showed a wide distribution of ω. In contrast, Hsp90s, Hsp70s, and Hsp60s seemed to

be very stable and under a very high purifying selection.

To understand other possible mechanisms of adaptation of *T. alpestre* to the

cold, we computed the mean values of ω and k for all branches leading to

Tetramorium spp. and *V. emeryi*, integrating 13 ant species, and scanned for

diversifying and relaxed selection specifically in *T. alpestre*. The first part of the

analysis showed a similar landscape as observed in scOGs. Among the five

Tetramorium species and *V. emeryi*, five species display a distribution of the mean-

branch ω ranging between 0.07 (*T. immigrans*) and 0.13 (*T. parvispinum*); *T. alpestre*

instead showed significant 2.7-fold higher value of the median mean-branch ω (ω =

0.36, Wilcoxon rank-sum test '*greater*', adjusted *P*-values < 0.004, Figure 5b). The

distribution of k among HSPs in the six species showed median values of k all close to

the expected (k = 1), between 0.90 in *V. emeryi* and 1.05 in *T. simillimum*, again with

the exception of *T. alpestre*, where the median k = 0.63, 1.4-fold lower than the

lowest median *k* (Wilcoxon rank-sum test '*less*', adjusted *P*-values < 0.05, Figure 5c).

The scan for diversifying positive and relaxed selection in branches of the HSPs of *T.*

alpestre

Figure 5. *a*) Boxplots showing the distribution and median (horizontal black line) of mean ω (d_N/d_S) rates for each
of the HSP subfamily. *b*) Violin plot showing the distribution of the mean ω values; and *c*) *k* in the terminal
branches of the Crematogastrini (from up to bottom: *T. alpestre, T. immigrans, T. bicarinatum, T. simillimum, T.*
parvispinum, V. emeryi). *d-f*) Scatter plot of target gene (*stv, Hsc70-4d, Hsc70-4e*) vs. housekeeping gene (HK)
concentrations; concentrations given as total number of cycles minus cycle threshold (Ct) values, that is, higher
values represent higher concentrations. Straight lines are linear regressions for each target gene against HK using
the different recovery times separately; based on one-way analysis of covariance followed by correction for
multiple comparison, just lines for those treatment / recovery-time combinations are shown for which
significances for just a single species arose, that is, *T. alpestre* coma vs. non-coma after 0.5 h (*d, e*; no difference
in *T. immigrans*) or for which the two species differed, that is, *T. alpestre* coma vs *T. immigrans* coma after 4.0 h
(*f*).

showed no gene under putative diversifying positive selection, while six loci were

identified to be under putative relaxing selection: one Hsp90 (*Trap1*), two Hsp60

(*CCT4*, *CCT5*), and three Hsp40s (*Sec63*, *DnajC8*, *DnajC11*) (k < 0.14, adjusted *P*-values

< 0.005). Using these genes in prediction interaction networks, we found that the

two Hsp60s interacted with other 21 genes, giving 17 significant enriched functions

(FDR < 0.05), mainly related to protein folding and microtubule, spindle organization;

the three Hsp40s interacted with 21 genes, giving two enriched functions (FDR <

0.05), related to response to heat and response to temperature stimulus (Figure

S19).

Chill shock effect, recovery times, and expression patterns of *stv* and HSPs in *T.*

alpestre* and *T. immigrans

Among the genes with a putative signature of diversifying selection, we also found

starvin (*stv*), a BAG-family member which seems to be implicated in Hsp70 ATPase

activity during cold recovery in *D. melanogaster* [30]. Therefore, we explored if the

modulation of its expression may play a role in how *T. alpestre* differently withstands

the cold in comparison to the related species *T. immigrans*. We thus performed a

pilot test measuring the different expression patterns of *stv*, two paralogs of *D.*

melanogaster Hsc70-4 (*Hsc70-4d* and *Hsc70-4e*), a member of the Hsp90 subfamily

(*Hsp83*), and sHsp *l(2)efl4* in the two species.

494   We measured the transcription activity of three recovery times (0.5 h, 2.0 h,

and 4.0 h) and compared coma and non-coma individuals, within and between

species, and found a significant difference in the expression patterns of *stv*, *Hsc70-*

4d, *Hsc70-4e*, and *Hsp83* were observed, but not of *l(2)efl4* (one-way ANCOVA

followed by Bonferroni-Holm correction). In particular, looking at the different

recovery times, *stv* and *Hsc70-4d* responded quickly in *T. alpestre* but not in *T.*

immigrans, being significantly expressed only in workers of *T. alpestre* after 0.5 h

recovery, but not in *T. immigrans* (Figure 5d-e). Comparing the expression levels

between species, we found higher mRNA levels for *Hsc70-4e* in *T. alpestre* after 4 h

recovery (Figure 5f).

Discussion

Tetramorium alpestre represents a critical example of evolutionary adaptation to

cold environments such as alpine habitats (Figure 1c). By comparing its genome with

other 18 ant species, we shed new light on how *T. aspestre* evolved to colonize sub-

alpine habitats and proposed it as a model for studies on arthropods adaptation to

cold. We found significant concordance between the nuclear and the mitochondrial

dated phylogenies, concerning the dating of the speciation of *T. alpestre*, which

seems to have diverged from its closest relative, *T. immigrans*, between 2 and 9 mya.

This period overlaps with the beginning of a new climatic zonation of the European

continent during the Middle and earliest Late Miocene [25]. At the end of this

period, a rapid uplift (~5 mya) fundamentally changed the paleogeographic and

topographic setting of central and southern Europe and triggered Alpine glaciation

[31]. Interestingly, both paleoclimatic and paleogeographic changes co-occurred

with the origin of the lineage leading to *T. alpestre* (Figure 1b). In fact, while these

glaciations dramatically reshaped global biodiversity patterns, eliminating terrestrial

biota from many mid- to high-latitude areas, they may have been the main driver of

diversification on highlands by reducing gene flow among populations [32].

The present results support our prediction that two parallel patterns will

occur in the genome of species adapting to extreme habitats. The genome of *T.*

alpestre appears to be under the influence of two strong evolutionary trajectories:

on the one hand, a diversifying selective force, two-fold higher compared with all

other ants, and on the other hand, a relaxation of the overall purifying selection

present in other ants, identified by the skewed distribution of k towards zero.

Specifically, the diversifying selective force affects more than a hundred genes in

many key biological processes, such as those related to development, cell migration,

and gastrulation. Genes under the category of biological processes are also under

selection in the genome of the Antarctic midge *Belgica antarctica* [8]. In fact, early

stages are the most fragile ones, having the highest rate of lethal phenotypes during

heat shock and UV irradiation, particularly around gastrulation [33]. The same may

pertain to genes related to cellular-level organization (e.g., spindle midzone

assembly, supramolecular fiber organization, actin cytoskeleton organization), where

environmental stress may jeopardise a well working cell division machinery, as SNPs

associated to long-term cold-adaptation plasticity were found to be present in genes

related to cytoskeletal and membrane structural components in *D. melanogaster*

[34]. Among 25 genes involved in these biological processes, *polo*, a polo-like kinase

which plays a central role as regulator of cell division and is required for several

events of mitosis and cytokinesis [35], shows three amino acid substitutions in

positions corresponding to active binding sites, and thus could be a good candidate

for further studies.

The two evolutionary trajectories were predicted in our hypothesis, and while

diversifying selection is somewhat expected in species adapting to a new

environmental niche, the magnitude of relaxed selection found is more surprising

and never recorded. Although both ω and k distributions in *T. alpestre* are highly

skewed, the absolute number of genes under diversifying and relaxed selection are

not markedly greater than other species in similar studies [27,28,36]. This cannot be

directly ascribed to the type *I* error rate because the same rate should be equally

randomly present in all the other short or long branches of the phylogeny.

Nevertheless, *T. alpestre*, as part of a species complex [18,19], may have more

recently diverged from more closely related species not yet adapted to the alpine

habitat. Magnitude of this relaxed selection is evidenced by the 70 enriched

biological processes, and can be seen as the consequence of a shift and/or

decreased magnitude in the purifying selection. The direct implication of this

relaxation is not clear, especially its effect on genes and their biological processes,

such as the regulation of cellular development, signal transduction, cell

communication, neurogenesis, axon development, and many others. If the

physiology of cold adaptation utilises different metabolic and developmental

strategies to minimise and optimise energy consumption, a new balance needs to be

reached. The two patterns here observed (diversifying/relaxing selection) might

represent this new shift. Interestingly, *T. immigrans*, which inhabit intermediate

environmental conditions (Figure 1c), shows also an intermediate pattern of *k*, with

an intriguing bimodal distribution (Figure 2b). Therefore, we can speculate that the

highly skewed distributions point to ongoing adaptation to a colder climate, and a

relaxation of selective forces present in warmer habitats. To our knowledge, this is

the first time that this effect is found in an organism, most likely due to the

underrepresentation of genomic studies on alpine and cold adapted species, and,

importantly, the lack of application of the RELAXED test in a genome-wise manner.

Examining the effect of those evolutionary trajectories, energetic metabolism

seems to be significantly involved in the sub-alpine ant's adaptation. Several clues

are pointing in that direction, such as the more specialized diet underlined by the

presence of commensal underground aphids in *T. alpestre* nests but not in those of

T. immigrans (unpublished data), the expansion of OGs related to sugar transporters,

and five enzymes under diversifying selection related to metabolism of sugars. We

deeply explored and validated one of these enzymes, Hex-T1, a key regulator and

rate-limiting enzyme for energy (sugars) metabolism and reactive oxygen species

(ROS) activity in insects [37], by contrasting its enzymatic activity between *T. alpestre*

and *T. immigrans.* We showed that the mutated form in *T. alpestre* is indeed more

efficient, with less substrate needed to reach the highest enzymatic activity, with the

protein modeling simulations demonstrating that the single amino acid mutation

enhances the catalytic efficiency of the enzyme due to the formation of a more

functional network of hydrogen bonds with its substrate. In fact, the path of the

hydrogen bonds found in TalpHex-T1 resembled the one depicted by the X-ray

solved structure of Hex-T1 from [38], used as a template to obtain the three-

dimensional coordinates of both TimmHex-T1 and TalpHex-T1 (see Methods).

Mulichak and colleagues (2000) identified their model residues Lys621 and Asp657

as crucial for the catalytic activity of the protein; the corresponding residues

TalpHex-T1$_{K168}$ and TalpHex-T1$_{D204}$ are indeed interacting, together with other

residues, with glucose in TalpHex-T1 at both 6 and 26 °C. On the other hand, while

the interaction with TalpHex-T1$_{D204}$ is conserved, the one with the lysine is lost in

TimmHex-T1 and a looser protein-sugar network is formed, explaining the higher

catalytic efficiency of TalpHex-T1 versus that of TimmHex-T1. We can indeed infer

that the higher flexibility conferred by TalpHex-T1$_{G74}$ increases the shift of the

residue TalpHex-T1$_{K168}$, belonging to the catalytic site (Figure 4d, e), allowing it to

move closer and establish a stronger interaction with glucose, possibly contributing

to lower the Km of the catalytic reaction. In eco-evolutionary terms all these findings

hint at a more intimate relation between *T. alpestre* and the aphids. This relationship

between ants and aphids is one of the most studied examples of mutualistic

relationships in the animal kingdom. Aphids produce honeydew, and in return ants

offer protection from predators, parasitoids, fungal infection and adverse conditions.

Honeydew contains monosaccharides (glucose and fructose), disaccharides (maltose,

sucrose), and Melezitose, a trisaccharide, that is one of the main sugar synthesized

by aphids, and which has been found to attract ants and maintain the ant-aphid

mutualism [39]. Small amounts of amino-acids, proteins, and lipids are also present

[40]. Usually, this interaction is above ground level, on plant leaves and branches,

but there are cases in which ants are specialized to farm subterranean aphids, just

like in *Lasius flavus* [41]. *Tetramorium alpestre* may have evolved the same behavior

through a coevolution of physiological factors such as palatability, fluid intake rate

and digestibility of sugar molecules, improving the ants' survival by enhancing

energy intake, a limiting factor in high-elevation ecosystems.

The complete annotation and characterization of the five HSP subfamilies in

19 ant species show good overall conservation, and strong purifying selection was

observed in most subfamilies. The exception are sHsps, which seem to be the most

diversified HSP subfamily, with the highest and the widest range of ω. It is also the

only subfamily that took a different evolutionary path compared with Diptera,

having very few orthologs in common with them. Given that sHsps are virtually

ubiquitous molecular chaperones that can prevent the irreversible aggregation of

denaturing proteins [42], they may have played a central role in the adaptation of

ants to strongly different climates and environments. Although we observe a higher

distribution of ω in *T. alpestre*, no HSPs show a significant sign of diversifying

selection. Rather, six loci show relaxed purifying selection (Hsp90: *Trap1*;

Hsp60:*CCT4*, *CCT5*; Hsp40: *Sec63*, *DnajC8*, *DnajC11*). The impact of the TNF receptor-

associated protein (Trap1) on cellular bioenergetics of *T. alpestre* may contribute to

its adaptation to cold environments by differently modulating cellular metabolism.

Given that it functions as a negative regulator of mitochondrial respiration, able to

modulate the balance between oxidative phosphorylation and aerobic glycolysis,

possibly another form of metabolic adaptation. It has been shown that reduced or

absent *Trap1* expression deregulated mitochondrial respiration leading to a high

energy state. This state is characterized by accumulation of ATP, tricarboxylic acid

cycle intermediates and reactive oxygen species, as well as an increase in fatty acid

oxidation [43]. *Trap1* overexpression is associated with increased expression of

genes associated with cell proliferation [44]. Since the two Hsp60s are associated to

protein folding and cytoplasmic microtubule organization, biological processes also

enriched by genes under diversifying selection, this suggests profound changes in

the cellular physiology of *T. alpestre*. It is also plausible that adapting to a cold

environment will depress selection on genes directly related to severe heat shock. As

we have shown, *T. alpestre*'s habitat is characterised by temperatures that rarely

exceed 20 °C (median = 16.5 °C; 99% confidence interval = 17.8 °C), although rocky

habitats may produce warmer microhabitats. This could mean that this species

experienced relaxed selection for HSPs associated with the denaturation to protein

folding by heat shock, as suggested by the relaxation of purifying selection on *Cct5*

and the three Hsp40s associated with responses to heat and temperature stimulus.

Although we provide a range of evidence that adaptation to cold

environments can be related to changes in protein-coding sequences, it has long

been postulated that phenotypic divergence between closely related species, such as

T. alpestre and *T. immigrans*, is primarily driven by quantitative and spatiotemporal

changes in gene expression, mediated by alterations in regulatory elements (e.g.:

Danko et al., 2018; Prescott et al., 2015). We tested this hypothesis by looking at the

expression patterns of *stv* and four other HSPs (*Hsc70-4d*, *Hsc70-4e*, *Hsp83*, *l(2)efl4*),

between *T. alpestre* and *T. immigrans*. *Stv* is a member of Bcl-2-associated

athanogene (BAG)-family proteins, which interact with Hsc70s and Hsp70s, and can

modulate, either positively or negatively, the functions of these chaperones [47]. It is

implicated in the recovery from chill shock in *D. melanogaster* during cold recovery

[30,48]. The concerted synthesis of *D. melanogaster stv* and Hsp70s suggests

cooperation to offset cold injury, possibly by preserving the folding/degradation and

by regulating apoptosis [30]. In our experiment, an acute cold stress (-6 °C) was used

to knock down individuals. By looking at the differential expression of these genes in

coma and non-coma individuals within and between species, we were not only able

to show that *stv* may interact with Hsc70s, but also that *T. alpestre* has a quicker

response in these genes compared with, *T. immigrans,* the related species not cold

adapted. Given these results, which are in line with the selection signature we

identified, it is probable that *stv* may play a similar role in the physiology of ants. As

we have shown in this study, ants completely lack Hsp70s (Figure S14) and retained

only the heat shock cognates (Hsc70s) with mainly two conserved paralogs (*Hsc70-*

4d and *Hsc70-4e*), which may have evolved to function as the Dipteran Hsp70s.

Although we found an interesting and promising correlation between chill shock, *stv*,

and Hsc70s, it will be important to continue exploring other expression patterns in

ant physiology to assess whether *stv* has tissue-specific expression patterns, as was

found in *D. melanogaster* [49], and to study its expression across different species

with different ecological adaptations.

One of the most enduring problems of evolutionary biology is explaining how

complex adaptive traits originate [50]. Although it is widely assumed that new traits

arise solely from selection on genetic variation, many researchers ask whether

phenotypic plasticity precedes and facilitates adaptation [51]. This alternative

scenario, the plasticity-first hypothesis, states that under novel conditions,

environmentally induced variation can be refined by selection and, depending on

whether plasticity is favoured, become developmentally canalized through genetic

assimilation. Under this scenario, environmentally induced phenotypic change can

precede and promote the evolutionary origins of a complex adaptive trait.

Four criteria have been proposed for verifying this hypothesis [51]: 1) the

focal trait needs to be environmentally induced in the ancestral-proxy lineages; 2)

cryptic genetic variation will be uncovered when ancestral-proxy lineages experience

the derived environment; 3) the focal trait will exhibit evidence of having undergone

an evolutionary change in its regulation, in its form, or in both in the derived

lineages; and 4) the focal trait will exhibit evidence of having undergone adaptive

refinement in the derived lineages. Though it is methodically difficult to prove each

point, our findings on the evolution and adaptation of *T. alpestre* to cold appear

consistent with plasticity-first evolution. In more detail, *T. immigrans*, the ancestral-

proxy lineage, shows considerable plasticity under variable temperature conditions

compared to other ant species (Criterion 1, Figure 1c), showing an ability to recover

from severe chill-shocks almost as good as the derived lineage (*T. alpestre*), and,

unexpectedly, better efficiency of Hex-t1 at lower rather than higher temperatures

(Figure 4a). We suggest that the derived lineage, *T. alpestre*, underwent genetic

assimilation of the cold response (Criterion 3), followed by evolutionary refinement

via changes in regulation, resulting, for example, in *stv* differential expression

(Criterion 4). Consistent with evolutionary refinement, the derived lineage has

evolved more extreme traits such as an overall improved activity of Hex-T1 with

reduced variability.

Our comparative genomic analysis shows how natural selection could trigger

complicated patterns of change in genomes, especially protein-coding sequences,

both in terms of diversifying selection and relaxation of purifying selection. Many of

these changes are in genes that may be associated with aspects of development,

either directly or through the associated complex changes in ecology and natural

history. Furthermore, this work characterized some candidate genes activities, both

in vitro and *in silico*, confirming our hypothesis and opening for the identification of

other genes possibly playing important role in adaptation to new ecological niches.

Overall, this study represents the first systematic attempt to provide a framework

for the genomic analysis of adaptation to extreme environments and underlines the

importance of studying organisms with different ecological niches in understanding

the genetic basis of ecological adaptation.

Methods

Species-locality and climatic data

To define the environment of *T. alpestre* and the other four *Tetramorium* species,

georeferenced occurrence data were compiled and complemented with those for

the native ranges of other ant species with available genomic data. Occurrence

records for 17 species (Table S2) were compiled from the Global Ant Biodiversity

Informatics database (GABI, Guénard et al., 2017, accessible via www.antmaps.org,

Janicki et al., 2016). The species determinations in the records were evaluated by

specialists in ant taxonomy, and erroneous data were excluded. Slightly incorrect

coordinates (i.e. in the ocean within 8.5 km of land) were assigned to the nearest

land point. Additionally, georeferenced occurrence data of two species (*T. alpestre,*

T. immigrans) were compiled from the latest taxonomic revision [18]. The

occurrence data of each species were spatially rarefied at a 5 km level using the R

package SPTHIN v. 0.1.0 [53], to counter spatial autocorrelation and to provide some

protection against sampling bias issues [54]. The Bioclim variables bio1-bio19 from

the WORLDCLIM database v. 1.4 [55] were extracted for each unique species locality at

a 2.5 arc min resolution (4.65 x 4.65 = 21.6225 km^2 at the equator) using ARCGIS v.

10.4 (ESRI, Redland, CA). A Wilcoxon-Mann-Whitney rank sum test as implemented

in the R package WILCOX.TEST (http://www.r-project.org) was adopted to test for

differences between *T. alpestre* and the other species individually for each of the

Bioclim variables. The Bonferroni-Holm sequential rejection procedure (Abdi 2010)

was used to control for multiple testing.

Biological sample processing, genome size estimation, and genome sequencing

Six colonies of *T. alpestre* ants were sampled, two for flow cytometry (nest 18811:

Penser Joch, Italy, 46.83379° N, 11.44652° E, 21 June 2016; nest 18813: Kühtai,

Austria, 47.27923, 11.07590° E, 21 June 2016), three for RNA extraction (nest 18586:

Zirmbachalm, Austria, 47.21920° N, 11.07530° E, 30 July 2015; nest 18590:

Zirmbachalm, 47.21910° N, 11.07620° E, 30 July 2015; nest 18594: Penser Joch,

46.82940°N, 11.43830° E, 2 August 2015), and one for DNA extraction (nest 18592:

Penser Joch, 46.83380° N, 11.44610° E, 2 August 2015). The ants were brought alive

to the laboratory (for flow cytometry) or immediately submerged in RNAlater (for

RNA extraction) or 96% ethanol (for DNA extraction) and stored at -70 °C (RNA) or -

20 °C (DNA).

Flow cytometry was used to estimate relative genome size from single *T.*

alpestre worker heads, using strain ISO-1 of *Drosophila melanogaster* as internal

standard (obtained from Bloomington Drosophila Stock Center, Indiana University,

Bloomington, USA; nuclear DNA content: 0.35 pg, Gregory 2018). Six individuals each

were measured from *T. alpestre* nests 18811 and 18813. In more detail, single *T.*

alpestre heads and, separately, heads of 10 ISO-1 females were chopped in 500 µL of

ice-cold Otto 1 Buffer (0.1 mol/L citric acid, 0.5% Tween20 (Merck KGaA, Darmstadt,

Germany)). The suspensions were filtered using 42-µm nylon meshes and incubated

for 5 min. Then, 1 mL of staining solution (DAPI (4 µL/mL) and 2-mercaptoethanol (2

µL/mL) in Otto 2 buffer (0.4 mol/L $Na_2HPO_4 \cdot 12 H2O$)) was added to each suspension.

For each measurement, 200 µL suspension of a single *T. alpestre* worker and 10 µL of

the ISO-1 standard were used. Fluorescence intensity of 3000 cell nuclei per sample

was measured with a Partec CyFlow space flow cytometer (Partec GmbH, Münster,

Germany). Gating and peak analysis were done automatically using the software

Partec Flo Max.

For the DNA and RNA extractions, the individuals were thawed on ice and the

gasters removed. DNA was extracted using the QIAamp DNA Micro Kit (Qiagen). The

extraction followed the protocol of the manufacturer, except that 20 µl RNase A

solution (Sigma-Aldrich) was added to the vials and incubated for 5 minutes at room

temperature. DNA concentration and quality were checked via Nanodrop (Thermo

Scientific) and gel electrophoresis. Two adult males were used to generate the

overlapping-read and the paired-end (PE) libraries, while for the mate-pair (MP)

libraries, four adult females were extracted of nest 18592. RNA was extracted using

the RNeasy Mini Kit (Qiagen) following the instructions of the manufacturer. Six

developmental stages were extracted as pools of individuals (nest 18586: worker

larvae – 30 individuals, worker pupae – 40, worker adult – 50; nest 18594: gyne

pupae – 2; nest 18590: gyne adult – 7, male adult – 7). RNA concentration and

quality were checked via Nanodrop (Thermo Scientific) and gel electrophoresis.

All *T. alpestre* DNA-seq and RNA-seq Illumina libraries were prepared by

IGATech (http://igatechnology.com/). Specifically, four libraries were used for

genome sequencing of *T. alpestre*: one overlapping-reads library (insert size ~400 bp;

paired-end (PE) sequencing with 2×250 bp read length), one regular PE library (insert

size ~400 bp, 2×100 bp read length), and two mate pair libraries (insert size ~4 kb,

2×100 bp read length) (see Table S2). For RNA-seq, six strand-specific libraries (insert

size ~500 bp, 2×200 bp read length) were constructed from three stages of workers

(larvae, pupae, adult), two stages of gynes (larvae and adult), and adult males. All

libraries were sequenced on an Illumina HiSeq 2500 platform in standard (DNA) and

RAPID (RNA) mode. For each *T. immigrans*, *T. bicarinatum*, *T. simillimum*, and *T.*

parvispinum, one TrueSeq library (insert size ~500 bp, 2×100 bp read length) was

prepared and sequenced on an Illumina HiSeq 4000 machine at OIST sequencing

center.

Genomic and mitogenomic assembly and gene prediction

Genomic sequencing reads of *T. alpestre* were filtered to remove low-quality reads

and PCR duplicates, corrected, and assembled using ALLPATHS-LG v. 52488 [56].

Contigs were first constructed based on overlapping reads and then scaffolded using

paired-end and mate-pair information from all DNA libraries. Given the high

coverage of the overlapping reads, based on the estimated genome size, multiple

trials were performed using different coverages to establish the best combination of

read coverage. The genomes of the other four related species were assembled using

SPADES v. 3.10.0 [57] with enabled read error correction algorithm, a *k*mer range of

21, 33, 55, 77, 99, and mismatch correction. Scaffolds belonging to contaminants

were identified and removed using BLOBOLOGY v. 2013-09-12 [58].

A bioinformatic pipeline that included combinatorial approaches of

homology, *ab initio*, and *de novo* methods were implemented to predict gene

models. REPEATMASKER v. 4.0.6 [59] was used to generate repeat hints. For the

homology-based exon hints, protein data sets of *D. melanogaster*, *Apis mellifera*,

Nasonia vitripennis, and 12 ant species (see Table S1) were aligned to the *T. alpestre*

genome using TBLASTN (*e*-value cut-off: 1e^{-5}) and EXONERATE v. 2.2.0 [60]. As an *ab*

initio procedure, the merged RNA-seq libraries were mapped using STAR v. 020201

[61] (settings: ALIGNENDSTYPE Local; ALIGNINTRONMIN 30; ALIGNINTRONMAX 450,000;

OUTSJFILTERINTRONMAXVSREADN 80 100 500 1000 2000 5000 20,000;

ALIGNSJOVERHANGMIN 10; ALIGNTRANSCRIPTSPERREADNMAX 100,000) to generate the

splice-site position-specific scoring matrix (PSSM) and CUFFLINKS v. v2.2.1 [62] to

generate UTR and intron hints. This information was then integrated with GENEMARK-

ES v. 4 [63] to train the analysis and use the outcome under the BRAKER1 procedure

[63] and AUGUSTUS v. 3.2.1. [64]. To avoid missing genes due to unassembled

regions of the genome, previously unmapped reads were *de novo* assembled using

TRINITY v. 2.2.0 [65] and annotated with FRAMA [66]. The three results where then

merged, and Benchmarking Universal Single Copy Orthologs (BUSCO v. 1.2) [67] was

adopted to evaluate the annotation completeness together with integrity and

completeness evaluation using DELTABLAST [68]. A visual inspection of the annotation

and gene integrity was also performed on more than 250 loci.

All available complete ant mitochondrial genomes (mtDNA) were

downloaded from GENBANK [69] (Table S1). Illumina raw SRA data of other ant

species were downloaded and processed following Cicconardi et al., (2017b). In

brief, reads were quality filtered and assembled using IDBA-UD v. 1.1.1 [70] and

SPADES v. 3.10.0 [57], and annotated using MITOS v. 2 web server [71].

Phylogenetic analysis

Nuclear single-copy ortholog genes (scOGs) and complete mitochondrial genomes

were used to compute the ant species tree. Once scOGs were determined (see

below), single-locus trees and a species tree of concatenated scOGs were estimated

using Maximum Likelihood (ML) search as implemented in FASTTREE v. 2.1.8 SSE3

[72]. Gene trees were summarized with MP-EST [73] using the STRAW web-server

[74].

For the mtDNA phylogeny, only protein-coding genes were used. Each gene

was aligned using MACSE v. 1.01b [75], and all alignments were concatenated. The

best partitions and models of evolution were identified with PARTITIONFINDER v.

1.1.1_Mac [76]. MtDNA trees were searched with ML and Bayesian Inference (BI)

algorithms. For ML, GARLI v. 2.01.1067 (http://code.google.com/p/garli) was used

performing 20 + 5 runs from random starting trees. The runs were continued until no

further improvement in log-likelihood was found, followed by 1000 bootstrap

pseudoreplicates. The results were summarized using SUMTREES v. 3.3.1

(http://bit.ly/DendroPy), using *A. mellifera*, *N. vitripennis*, and *D. melanogaster* as

outgroups. For BI, BEAST v. 2 [77] was adopted to obtain the dated mtDNA tree and

to estimate the divergence of the most recent common ancestors (MRCAs)

(template: BEAST [78]). Fossil data were taken mainly from Moreau & Bell (2013)

and PALEOBIO DB (paleobiodb.org) (see Table S4 for a detailed list of references) and

used as minimum calibration points constraining nodes in the topology as listed in

Table S4. Lognormal prior distribution was implemented with an offset

corresponding to the minimum fossil age, log(mean) of 1.0, and log(SD) of 1.0. Ten

independent runs of 8 x 10^7 generations were sampled every 100,000th generation.

Each partition was modelled with an uncorrelated relaxed clock with a Yule process,

species tree prior, and its best substitution model. Evolutionary models not

implemented in BEAUTI v2 [77] were manually edited in the xml file. TRACER v. 1.6

(http://beast.bio.ed.ac.uk/Tracer) was used to evaluate convergence and the

parameters' effective sampling size (ESS) and a burn-in was manually implemented

and roughly 10% from each run were removed. LOGCOMBINER [77] and TREEANNOTATOR

[77] were used to summarize the results in a single consensus tree. The non-ant

species were excluded from the analysis, and no outgroup was adopted. Divergence

dates for the maximum-likelihood nuclear DNA (nuDNA) tree (only ants) were also

inferred using the penalized likelihood approach [79] implemented in the R package

APE v. 5.1 [80] by calibrating four nodes: the root, the *Tetramorium*, the

Solenopsis+Monomorium, and the Attini branches (Table S4).

Functional annotation and orthologous-group dynamic evolution

Genes from 19 ant and three outgroup species (Table S3) were clustered into

ortholog groups (OGs) with HIERANOID v. 2 [81], which implements INPARANOID v. 8

[82] using a guide tree topology based on Moreau & Bell (2013) and BLASTP (*e*-value

cutoff of 10^{-5}) to improve the reciprocal hit accuracy [83], while homology

relationships were searched with DELTABLAST [68]. The functional annotation of each

putative protein coding sequence was performed by identifying both the protein

domain architecture and Gene Ontology (GO) terms. For each sequence, first

HMMER v. 3.1b2 (HMMSCAN) [84] was used to predict PFAM-A v. 31.0 domains, then

Domain Annotation by a Multi-objective Approach (DAMA) v. 2 [85] was applied to

identify architectures combining scores of domain matches, previously observed

multi-domain co-occurrence, and domain overlapping. Annotation of GO terms was

performed as implemented in the CATH assignments for large sequence datasets

[86]. Briefly, each input sequence was scanned against the library of CATH functional

families (FᴜɴFᴀᴍs) HMMs [87] using HMMER3 [84] to assign FᴜɴFᴀᴍs to regions on

the query sequence (with conditional E-value < 0.005). Then the GO annotations for

a matching FᴜɴFᴀᴍ were transferred to the query sequence with its confidence

scores, calculated by considering the GO term frequency among the annotated

sequences. Finally, a non-redundant set of GO annotations was retained, making up

the GO annotations for the query protein sequence [87].

To identify orthologous groups specifically expanded in the *T. alpestre*

genome, BᴀᴅɪRᴀᴛᴇ v. 1.35 [88] was performed twice, once for the nuDNA and one for

the mtDNA ultrametric tree topologies, reconstructing ancestral family sizes, gain,

death, and innovation (GDI), applying stochastic models and allowing estimation of

the family turnover rate (λ) by ML. Using this approach, we observed a correlation

between the turnover rate (λ) and branch lengths (Spearman correlation: ρ = -0.52,

P-value = 0), and a bimodal distribution of λ. We interpreted the distribution closer

to 0 as background noise and therefore considered only expanded and contracted

OGs with λ within the second distribution. The boundary between the two

distributions was defined by the valley values between the two, which were

computed using ᴏᴘᴛɪᴍɪᴢᴇ and ᴀᴘᴘʀᴏxFᴜɴ functions implemented in the sᴛᴀᴛs v. 3.4.0

package in R.

Heat-shock protein family analysis

From the functional annotation analyses of all Hymenoptera, all sequences bearing a

valid protein domain matching HSPs were extracted and aligned using the CʟᴜsᴛᴀʟW

v.1.2.1 EBI web server [89] (settings: ᴍʙᴇᴅ true; ᴍʙᴇᴅɪᴛᴇʀᴀᴛɪᴏɴ true; ɪᴛᴇʀᴀᴛɪᴏɴs 5;

ɢᴛɪᴛᴇʀᴀᴛɪᴏɴs 5; ʜᴍᴍɪᴛᴇʀᴀᴛɪᴏɴs 5). A phylogenetic tree was computed using ML as

implemented in FASTTREE v. 2.1.8 SSE3 [72] and manually checked to identify and

provisionally annotate the five main subfamily proteins (Hsp90, Hsp70, Hsp60,

Hsp40, and sHsp) and their OG. Sequences falling outside clusters and with low

bootstrap values were manually checked with the DELTABLAST [68] web-server and

with the CONSERVED DOMAIN DATABASE (CDD) web-server [90] to identify and remove

bacterial contaminants and spurious/incomplete sequences. Then, nucleotide

sequences belonging to each subfamily were revers translated into amino acids,

aligned using CLUSTALW v.1.2.1, and the obtained protein alignment was used to

derive the nucleotide one. FASTTREE was used again to compute phylogenetic trees

for the five HSP subfamilies.

Selection on single-copy orthologous groups and gene families

All selected scOGs and HSP-OGs were scanned to evaluate selection signature on

coding regions of *T. alpestre*. This was done by computing the mean ω and the

relaxation of each branch of the phylogeny of *Tetramorium* spp. + *Vollenhovia*

emeryi. More specifically, adopting a pipeline similar to that in Cicconardi et al.

(2017a, 2017b), the signatures of diversifying selection were searched in codon-

based aligning groups of one-to-one orthologous ant genes with MACSE v. 1.01b

[75], filtering with GBLOCKS v. 0.91b (Castresana 2000) under a "relaxed" condition

[10,28,91] and using the aBSREL algorithm as implemented in the HYPHY batch

language [92] using a batch script (BRANCHSITEREL) in HYPHY

(http://github.com/veg/hyphy).

In parallel, the same scOGs and HSP-OGs were scanned using the RELAX test

[93] to search for putative signals of relaxation. In brief, RELAX tests the hypothesis

of evolutionary-rate relaxation in selected branches of a phylogenetic tree compared

with reference branches. A k value is computed to evaluate whether the selective

strength ω shifts towards neutrality. The rate of d_N/d_S (ω) can relax ($k < 1$), stay

stable ($k = 1$), or intensify ($k > 1$). For both scOGs and HSP-OGs, and for both the

ABSREL and RELAX tests, the Bonferroni-Holm sequential rejection procedure (Abdi

2010) was used to control the false discovery rate with a very stringent cutoff value

of 0.005 for the adjusted P-values applied [94]. Hsp60s and Hsp40s under relaxed

selection were analyzed using the GENEMANIA prediction server [95] to predict their

functions by creating an interaction network with genes by including protein and

genetic interactions, pathways, co-expression, co-localization, and protein domain

similarity.

Finally, the GOSTATS package for R [96] (settings: ANNOTATION org.Dm.eg.db;

CONDITIONAL TRUE; TESTDIRECTION over) was used to check for GO terms of biological

processes and KEGG pathway enrichments of scOGs under selection and relaxation

using the whole set of tested scOGs as background. P-values < 0.005 and < 0.05 were

implemented for GO terms and KEGG pathways, respectively.

Synthesis of Hex-t1 gene and its enzyme activity assays

Following identification as under diversifying selection in *T. alpestre* (see results

section "Dual signature of evolutionary pressures in single-copy OGs in the *T.*

alpestre genome"), synthesis of genes and enzymes was attempted for

phosphofructokinase, transketolase, phosphoglucosemutase, and Hex-t1 was

successful just for the Hex-t1. In more detail, the Hex-t1 genes of *Tetramorium*

alpestre (1398 bp) and *T. immigrans* (1386 bp) were synthesized by ThermoFisher

lifetechnologies (Carlsbad, CA, USA) and Eurofins Genomics (Ebersberg, Germany),

respectively. Both constructs carried a 5' EcoRI and a 3' HindIII recognition site for

subsequent cloning as well as a C-terminal His-tag. The genes were excised from

their carrier backbones by enzymatic digestion. The fragments were purified by

agarose gel excision and sub-cloned into the pACEBac1 expression vector

(MULTIBACTM, Geneva Biotech, Geneva, Switzerland). Correct insertion was verified

by Sanger sequencing, and transformation into *D. melanogaster* cells and protein

expression followed the standard protocols of the expression vector manufacturer.

After cell harvesting, the presence of the recombinant protein in the cell lysates was

verified by a western blot targeting the His-tag. The relative amounts of protein

were quantified from the western blot using ImageJ (https://imagej.net/Welcome).

The cell lysates were centrifuged at 4000 rpm at 4 °C for 5 min, and the supernatant

was directly used as protein source.

Enzyme activity assays followed a modified protocol of Crabtree &

Newsholme (1972). Briefly, 1000 µl assay medium containing 75 mM Tris pH=7.5

(Merck Millipore, Burlington, MA, USA), 7.5 mM $MgCl_2$ (Merck), 0.8 mM EDTA

(Sigma-Aldrich, St. Louis, MO, USA), 1.5 mM KCl (Merck), 4.0 mM mercaptoethanol

(Sigma), 0.4 mM NADP+ (Abcam, Cambridge, UK), 2.5 mM ATP (Abcam), and 10 mM

creatine phosphate (Abcam) were placed in a disposable cuvette (Brand, Germany)

in a SPECORD 210 PLUS UV/VIS spectrophotometer (Analytic Jena AG, Germany)

with an attached 1157P programmable thermostate (VWR, Radnor, PA, USA).

Variable amounts of D-glucose (VWR) and synthetic protein were added, and the

cuvette was allowed to reach the assay temperature. Then, the reaction was started

by adding 15 U creatine phosphokinase (Sigma) and 5 U glucose 6-phospahte

dehydrogenase (Sigma).

The Hex-t1 activity was measured in triplicate as the rate of reduction of

NADP$^+$ causing an extinction at 340 nm. Each measurement lasted for 60 s, and every

second, an extinction value was recorded. For the determination of the Michaelis-

Menten constant (Km-value) of both enzymes, assays were conducted along a

substrate gradient from 0.00167 to 24.30000 mM D-glucose, achieving saturated

extinction curves. Assay temperatures ranged from 6 °C to 36 °C with 10 K

increments. The software PRISM v. 8 (GraphPad Software, San Diego, CA, USA) was

used to calculate the Km-values.

Hex-t1 structure modelling and molecular dynamics simulations

The three-dimensional structures of *T. alpestre* and *T. immigrans* Hex-t1 proteins are

not available. To study the effect of the amino acid changes on the proteins'

function/dynamics, we modeled the structures of these two proteins via homology

modelling. The X-ray structure of the Hex-t1 isoform from *Schistosoma mansoni* [38],

solved at 2.6 Å resolution was used as a template and the models were generated

using the software MODELLER v. 9.21 [97]. Once we obtained the models, we built the

simulative systems to run molecular dynamics simulations. The two proteins

TalpHex-t1 and TimmHex-t1, in complex with the glucose substrate, were immersed

in a triclinic box filled with TIP3P water molecules [98] and rendered electroneutral

by the addition of cloride counterions. The topology of the two systems, consisting

of ~ 80.000 atoms each, was built using the AMBER14 force field [99], further

converted in GROMACS v. 4.6 [100] format using ACPYPE [101]. Simulations were run

on a GPU cluster using GROMACS v. 4.6 with the following protocol: 1) 25000 steps of

steepest descent followed by 25000 steps of coniugate gradient minimization; 2)

5*100 ps of equilibration runs in the NVT environment starting at 50 K and

performed by increasing the temperature of 50 K after each run untill a final value of

250 K; 3) 5*100 ps of equilibration runs in the NPT environment starting at 50 K and

performed by increasing the temperature of 50 K after each run untill a final value of

250 K; 4) the equilibrated systems were simulated for 500 ns at 279 or 300 K, i.e. at 6

or 26 °C, for a total of four simulations and 2 μs of sampling. Electrostatic

interactions were considered by means of the Particle Mesh Ewald method (PME)

with a cut off of 1.2 nm for the real space and Van der Waals interactions [102].

Bond lengths and angles were constrained via the LINCS algorithm [103]. The

temperatures were kept constant at 279 or 300 °K by using the velocity rescale with

a coupling constant of 0.1 ps, and the pressure was kept at 1 bar using the

Parrinello-Rahman barostat with a coupling constant of 1.0 ps during sampling [104].

The four trajectories were collected and comparative analyses performed with the

GROMACS suite or with in house written code.

1015

Chill coma assay and quantitative qRT-PCR assay

The cold hardiness and expression of selected genes were compared between *T.*

alpestre and *T. immigrans* using chill coma assays. From four colonies additional to

those used for other experiments in this study, two per species, about 500 workers

per colony were collected in August 2017; *T. alpestre*: Jaufenpass (18980: 46.83791°

N, 11.29768° E) and Penser Joch (18978: 46.81402° N, 11.44198° E); *T. immigrans*:

Vienna (18977: 48.12448° N, 16.43522° E.; 18983: 48.31341° N, 16.42529° E).

Workers were collected alive using aspirators, transported to the laboratory in

Innsbruck, and kept in polypropylene boxes (18.5 cm x 11 cm) with chambers of

various sizes. The walls of the boxes were Fluon-coated (GP1, De Monchy

International BV, Rotterdam, Netherlands) to prevent workers from escaping. Food

(sugar-honey-water and deep-frozen *Drosophila* flies) and tap water were provided

ad libitum three times a week. Workers were kept in a climate chamber (MIR-254,

Panasonic, Etten Leur, Netherlands) at a 12L:12D photoperiod at constant 18 °C. The

temperature of 18 °C was the average of the monthly mean temperatures

experienced at 200 m a.s.l. (*T. immigrans*, approx. 23.7 °C) and at 2000 m a.s.l. (*T.*

alpestre, approx. 12.3 °C) in July 2017. Workers were acclimatized to 18 °C for at

least three weeks before the chill coma assays.

After the acclimation period, from each of the four colonies, 90 workers were

randomly chosen and divided into 18 equivalent pools, each were used for the chill

coma and for the control assays. For the chill coma assay, worker pools were

transferred into empty 5-ml glass vials, sealed, and immersed for 6.5 hours in a

water:ethane-1,2-diol mix (1:1) bath set at -6 °C; in pilot experiments, this

temperature had been determined as the highest temperature at which the ants had

fallen into chill coma after 6.5 hours exposition; the temperature was identical for

both species. Temperature was monitored using an electronic thermometer (TFX

430, ebro Electronic GmbH; Ingolstadt, Germany) with an accuracy of 0.05 °C

inserted into an additional, empty vial treated in the same way as the vials with ants.

After chill coma, the vials were transferred to a climate chamber at 18 °C, and the

workers were allowed to recover for 0.5, 2.0, and 4.0 hours. For each recovery time,

three replicate pools were used. After recovery, the workers were transferred to

RNAse-free reaction tubes and killed in liquid nitrogen. Tubes were stored at -70 °C.

For the control assay, worker pools were transferred to empty glass vials and placed

in the climate chamber at 18 °C. After 6.5 hours plus respective recovery time, the

workers were killed and stored as described above.

For each recovery time and replicate, complete worker pools were used for

the molecular analyses. RNA was extracted using the NUCLEOSPIN® RNA Kit

(Macherey-Nagel, Düren, Germany) following the instructions of the manufacturer.

First-strand cDNA was synthesized using 200 U REVERTAID reverse transcriptase (all

reagents by Thermo Fisher Scientific, Waltham, USA), 40 U RIBOLOCK RNASE inhibitor,

5 µM random hexamer primers, 100 µM dNTPs, and 2 µl RNA extract in a total

volume of 40 µl. The mixture was incubated for 5 min at 25 °C, 60 min at 42 °C, and 5

min at 70 °C on a UnoCycler 1200 (VWR, Radnor, USA). Quantitative PCR was

conducted on a Rotorgene Q (Qiagen) PCR system. Each reaction contained 1×

Rotor-Gene SYBR Green PCR Mastermix (Qiagen), 0.2 µM of primers, specifically

designed for target genes (Table S12), and 1 µl cDNA in a total volume of 10 µl. All

qRT-PCR reactions were performed as triplicates. Cycling conditions were 95 °C for 5

min, followed by 40 cycles of 94 °C for 15 s, 58 °C for 10 s, and 72 °C for 15 s.

Fluorescence was acquired at the end of each elongation step. PCR was followed by

a melting curve analysis from 60 to 95 °C with 0.1 °C increments held for 5 s before

fluorescence acquisition. Cycle threshold (Ct) values were subtracted from total

number of cycles. On these values, linear-regression analyses were based separately

for each target gene, treatment, recovery time, and species (in each regression

analysis, two populations and three replicate values were used, i.e., n = 6) using the

housekeeping gene as independent variable and the target gene as dependent

variable. For each target gene / recovery-time combination one-way analysis of

covariance (ANCOVA) followed by Bonferroni-Holm correction was then used to

identify significant differences between treatments within species and between

species within treatment.

1075

1076

Data access

The sequence data from this study have been submitted to the National Center for

Biotechnology Information (NCBI) Sequence Read Archive (SRA) under BioProject

numbers PRJNA532334, PRJNA533534, PRJNA533535, PRJNA533536 and

PRJNA533537 (https://www.ncbi.nlm.nih.gov/bioproject/). The mtDNA genome

assemblies have been submitted to the NCBI (GenBank accessions: MK861047 -

MK861070). All accession codes of deposited and retrieved data are provided in

Table S1.

1085

1086

Acknowledgements

Philipp Andesner and Stefan Gross for valuable work in the laboratory. Research was

supported by the Austrian Science Fund (FWF, P23409, P30861). The computational

results presented here have been achieved in part by using the HPC infrastructure of

the University of Innsbruck (LEO), the Vienna Scientific Cluster (VSC), the

collaborative system of the Universities Innsbruck and Linz (MACH), and the Swiss

National Supercomputing Centre (CSCS) hosted in Lugano.

1094

1095

References

1.   Soberón J. Grinnellian and Eltonian niches and geographic distributions of

species. Ecology Letters. 2007. pp. 1115–1123. doi:10.1111/j.1461-

0248.2007.01107.x

2.   Lamprecht A, Semenchuk PR, Steinbauer K, Winkler M, Pauli H. Climate

change leads to accelerated transformation of high-elevation vegetation in

the central Alps. New Phytol. 2018; 447–459. doi:10.1111/nph.15290

3.   Rogora M, Frate L, Carranza ML, Freppaz M, Stanisci A, Bertani I, et al.

Assessment of climate change effects on mountain ecosystems through a

cross-site analysis in the Alps and Apennines. Sci Total Environ. Elsevier B.V.;

2018;624: 1429–1442. doi:10.1016/j.scitotenv.2017.12.155

4.   Körner C, Paulsen J, Spehn EM. A definition of mountains and their bioclimatic

belts for global comparisons of biodiversity data. Alp Bot. 2011;121: 73–78.

doi:10.1007/s00035-011-0094-4

5.   Clark MS, Worland MR. How insects survive the cold: molecular

mechanisms—a review. J Comp Physiol B. 2008;178: 917–933.

doi:10.1007/s00360-008-0286-4

6.   Parker DJ, Wiberg RAW, Trivedi U, Tyukmaeva VI, Gharbi K, Butlin RK, et al.

Inter and Intraspecific Genomic Divergence in Drosophila montana Shows

Evidence for Cold Adaptation. 2018;10: 2086–2101. doi:10.1093/gbe/evy147

7.   Keeling CI, Yuen MM, Liao NY, Roderick Docking T, Chan SK, Taylor G a, et al.

Draft genome of the mountain pine beetle, Dendroctonus ponderosae

Hopkins, a major forest pest. Genome Biol. BioMed Central Ltd; 2013;14: R27.

doi:10.1186/gb-2013-14-3-r27

8. Kelley JL, Peyton JT, Fiston-Lavier A-S, Teets NM, Yee M-C, Johnston JS, et al.

Compact genome of the Antarctic midge is likely an adaptation to an extreme

environment. Nat Commun. Nature Publishing Group; 2014;5: 4611.

doi:10.1038/ncomms5611

9. Macdonald HC, Cunha L, Bruford MW, Florestas E, Macdonald HC, Sir T, et al.

Development of genomic resources for four potential environmental

bioindicator species: *Isoperla grammatica*, *Amphinemura sulcicollis*, *Oniscus*

asellus and *Baetis rhodani*. bioRxivorg. 2016; doi:10.1101/046227

10. Cicconardi F, Di Marino D, Olimpieri PP, Arthofer W, Schlick-Steiner BC, Steiner

FM. Chemosensory adaptations of the mountain fly Drosophila nigrosparsa

(Insecta: Diptera) through genomics' and structural biology's lenses. Sci Rep.

Nature Publishing Group; 2017;7: 43770. doi:10.1038/srep43770

11. King AM, MacRae TH. Insect Heat Shock Proteins During Stress and Diapause.

Annu Rev Entomol. 2015;60: 59–75. doi:10.1146/annurev-ento-011613-

162107

12. Nygaard S, Hu H, Li C, Schiøtt M, Chen Z, Yang Z, et al. Reciprocal genomic

evolution in the ant-fungus agricultural symbiosis. Nat Commun. 2016; 1–9.

doi:10.1038/ncomms12233

13. Hölldobler B, Wilson EO. The Ants. Harvard University Press. 1990.

14. Bolton B. An online catalog of the ants of the world. Available from

http://antcat.org [Internet]. 2018.

15. Seifert B. The ants of Central and North Europe. Tauer, editor. Lutra Verlags-

und Vertriebsgesellschaft; 2018.

16.   Wilson EO. Success and Dominance in Ecosystems: The Case of Social Insects.

Excellence in ecology. 1990.

17.   Christian KA, Morton SR. Extreme thermophilia in a central Australian ant,

Melophorus bagoti. Physiol Zool. 1992;65: 885–905.

18.   Wagner HC, Arthofer W, Seifert B, Muster C, Steiner FM, Schlick-Steiner BC.

Light at the end of the tunnel: Integrative taxonomy delimits cryptic species in

the Tetramorium caespitum complex (Hymenoptera: Formicidae).

Myrmecological News. 2017;25: 95–129.

19.   Steiner FM, Seifert B, Moder K, Schlick-Steiner BC. A multisource solution for a

complex problem in biodiversity research: Description of the cryptic ant

species Tetramorium alpestre sp.n. (Hymenoptera: Formicidae). Zool Anzeiger

- A J Comp Zool. 2010;249: 223–254. doi:10.1016/j.jcz.2010.09.003

20.   Krapf P, Russo L, Arthofer W, Möst M, Steiner FM, Schlick-Steiner BC. An

Alpine ant's behavioural polymorphism: monogyny with and without internest

aggression in Tetramorium alpestre. Ethol Ecol Evol. 2018;30: 220–234.

doi:10.1080/03949370.2017.1343868

21.   Liu C, Guénard B, Garcia FH, Yamane S, Blanchard B, Yang DR, et al. New

records of ant species from Yunnan, China. Zookeys. 2015;78: 17–78.

doi:10.3897/zookeys.477.8775

22.   Bertelsmeier C, Ollier S, Liebhold A, Keller L. Recent human history governs

global ant invasion dynamics. Nat Ecol Evol. 2017;1. doi:10.1038/s41559-017-

0184

23.   Guénard B, Weiser MD, Gómez K, Narula N, Economo EP. The Global Ant

Biodiversity Informatics (GABI) database: Synthesizing data on the geographic

distribution of ant species (Hymenoptera: Formicidae). Myrmecological News.

2017. pp. 83–89.

24.   Jayakumar PC, Shouche YS, Patole MS. Functional analysis of *Drosophila*

melanogaster hexokinase Hex-A locus⬚: multiple Initiator-like elements

enhance DPE containing promoter activity. 2007;16: 3–13.

doi:10.1111/j.1365-2583.2006.00664.x

25.   Zachos J, Pagani M, Sloan L, Thomas E, Billups K. Trends, Rythms, and

Aberration in Global Climate 65 Ma to Present. Science (80-). 2001;292: 686–

693. doi:10.1126/science.1059412

26.   Moreau CS, Bell CD. Testing The Museum Versus Cradle Tropical Biological

Diversity Hypothesis: Phylogeny, Diversification, And Ancestral Biogeographic

Range Evolution Of The Ants. Evolution (N Y). 2013;67: 2240–2257.

doi:10.1111/evo.12105

27.   Roux J, Privman E, Moretti S, Daub JT, Robinson-Rechavi M, Keller L. Patterns

of positive selection in seven ant genomes. Mol Biol Evol. 2014;31: 1661–

1685. doi:10.1093/molbev/msu141

28.   Cicconardi F, Marcatili P, Arthofer W, Schlick-Steiner BC, Steiner FM. Positive

diversifying selection is a pervasive adaptive force throughout the Drosophila

radiation. Mol Phylogenet Evol. 2017;112: 230–243.

doi:10.1016/j.ympev.2017.04.023

29.   Nguyen AD, Gotelli NJ, Cahan SH. The evolution of heat shock protein

sequences, cis-regulatory elements, and expression profiles in the eusocial

Hymenoptera. BMC Evol Biol. BMC Evolutionary Biology; 2016;16: 15.

doi:10.1186/s12862-015-0573-0

30.   Colinet H, Hoffmann A. Gene and protein expression of *Drosophila* Starvin

during cold stress and recovery from chill coma. Insect Biochem Mol Biol.

Elsevier Ltd; 2010;40: 425–428. doi:10.1016/j.ibmb.2010.03.002

31.   Kuhlemann J. Paleogeographic and paleotopographic evolution of the Swiss

and Eastern Alps since the Oligocene. Glob Planet Change. 2007;58: 224–236.

doi:10.1016/j.gloplacha.2007.03.007

32.   Wallis GP, Waters JM, Upton P, Craw D. Transverse Alpine Speciation Driven

by Glaciation. Trends Ecol Evol. Elsevier Ltd; 2016;31: 916–926.

doi:10.1016/j.tree.2016.08.009

33.   Uchida Y, Uesaka M, Yamamoto T, Takeda H, Irie N. Embryonic lethality is not

sufficient to explain hourglass-like conservation of vertebrate embryos.

Evodevo. BioMed Central; 2018;9: 1–11. doi:10.1186/s13227-018-0095-0

34.   Gerken AR, Eller OC, Hahn D a, Morgan TJ. Constraints, independence, and

evolution of thermal plasticity: Probing genetic architecture of long- and

short-term thermal acclimation. Proc Natl Acad Sci U S A. 2015;112: 4399–

404. doi:10.1073/pnas.1503456112

35.   Archambault V, Lépine G, Kachaner D. Understanding the Polo Kinase

machine. Oncogene. 2015;34: 4799–4807. doi:10.1038/onc.2014.451

36.   Harpur BA, Kent CF, Molodtsova D, Lebon JMD, Alqarni AS, Owayss AA, et al.

Population genomics of the honey bee reveals strong signatures of positive

selection on worker traits. Proc Natl Acad Sci. 2014;111: 2614–2619.

doi:10.1073/pnas.1315506111

37.   Xian-Wu L, Wei-Hua X. Hexokinase is a key regulator of energy metabolism

and ROS activity in insect lifespan extension. Aging (Albany NY). 2016;8: 245–

258. Available:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4789580/pdf/aging-08-

245.pdf

38.   Mulichak AM, Wilson JE, Padmanabhan K, Garavito RM. The structure of

mammalian hexokinase-1. Nat Struct Biol. 2002; doi:10.1038/811

39.   Fischer MK, Shingleton AW. Host plant and ants influence the honeydew

sugar. Funct Ecol. 2001;15: 544–550.

40.   Pringle EG, Novo A, Ableson I, Barbehenn R V, Vannette RL. Plant-derived

differences in the composition of aphid honeydew and their effects on

colonies of aphid-tending ants. 2014; doi:10.1002/ece3.1277

41.   Ivens ABF, Kronauer DJC, Pen I, Weissing FJ, Boomsma JJ. Ants farm

subterranean aphids mostly in single clone groups - An example of prudent

husbandry for carbohydrates and proteins? BMC Evol Biol. 2012;12.

doi:10.1186/1471-2148-12-106

42.   Haslbeck M, Vierling E. A first line of stress defense: Small heat shock proteins

and their function in protein homeostasis. J Mol Biol. Elsevier Ltd; 2015;427:

1537–1548. doi:10.1016/j.jmb.2015.02.002

43.   Yoshida S, Tsutsumi S, Muhlebach G, Sourbier C, Lee M-J, Lee S, et al.

Molecular chaperone TRAP1 regulates a metabolic switch between

mitochondrial respiration and aerobic glycolysis. Proc Natl Acad Sci. 2013;110:

E1604–E1612. doi:10.1073/pnas.1220659110

44.   Liu D, Hu J, Agorreta J, Cesario A, Zhang Y, Harris AL, et al. Tumor necrosis

factor receptor-associated protein 1(TRAP1) regulates genes involved in cell

cycle and metastases. Cancer Lett. 2010;296: 194–205.

doi:10.1016/j.canlet.2010.04.017

45.   Prescott SL, Srinivasan R, Marchetto MC, Grishina I, Narvaiza I, Selleri L, et al.

Enhancer Divergence and cis-Regulatory Evolution in the Human and Chimp

Neural Crest. Cell. 2015;163: 68–84. doi:10.1016/j.cell.2015.08.036

46.   Danko CG, Choate LA, Marks BA, Rice EJ, Wang Z, Chu T, et al. Dynamic

evolution of regulatory element ensembles in primate CD4+ T cells. Nat Ecol

Evol. Springer US; 2018;2. doi:10.1038/s41559-017-0447-5

47.   Doong H, Vrailas A, Kohn EC. What's in the "BAG"? - A functional domain

analysis of the BAG-family proteins. Cancer Lett. 2002;188: 25–32.

doi:10.1016/S0304-3835(02)00456-1

48.   Colinet H, Hoffmann AA. Comparing phenotypic effects and molecular

correlates of developmental, gradual and rapid cold acclimation responses in

Drosophila melanogaster. Funct Ecol. 2012;26: 84–93. doi:10.1111/j.1365-

2435.2011.01898.x

49.   Coulson M, Robert S, Saint R. Drosophila starvin encodes a tissue-specific

BAG-domain protein required for larval food uptake. Genetics. 2005;171:

1799–1812. doi:10.1534/genetics.105.043265

50.   Wagner GP, Lynch VJ. Evolutionary novelties. Curr Biol. 2010;20: 48–52.

Available: https://pdf.sciencedirectassets.com/272099/1-s2.0-

S0960982210X00021/1-s2.0-S0960982209019459/main.pdf?x-amz-security-

token=AgoJb3JpZ2luX2VjECIaCXVzLWVhc3QtMSJHMEUCIDsUh2jRKJL1EeOiMr

NglWa%2BljWFLis7tEu%2FF65JtK%2FrAiEA3Ge%2FEhYUyff41n0NLuT%2F6f3O

MYi9qUIAiE

51.   Levis NA, Pfennig DW. Evaluating 'Plasticity-First' Evolution in Nature: Key

1263   Criteria and Empirical Approaches. Trends Ecol Evol. Elsevier Ltd; 2016;31:

1264   563–574. doi:10.1016/j.tree.2016.03.012

1265   52.   Janicki J, Narula N, Ziegler M, Guénard B, Economo EP. Visualizing and

1266   interacting with large-volume biodiversity data using client-server web-

1267   mapping applications: The design and implementation of antmaps.org. Ecol

1268   Inform. 2016;32: 185–193. doi:10.1016/j.ecoinf.2016.02.006

1269   53.   Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP. spThin:

1270   An R package for spatial thinning of species occurrence records for use in

1271   ecological niche models. Ecography (Cop). 2015;38: 541–545.

1272   doi:10.1111/ecog.01132

1273   54.   Merow C, Smith MJ, Silander JA. A practical guide to MaxEnt for modeling

1274   species' distributions: What it does, and why inputs and settings matter.

1275   Ecography (Cop). 2013;36: 1058–1069. doi:10.1111/j.1600-0587.2013.07872.x

1276   55.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution

1277   interpolated climate surfaces for global land areas. Int J Climatol. 2005;25:

1278   1965–1978. doi:10.1002/joc.1276

1279   56.   Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, et al.

1280   High-quality draft assemblies of mammalian genomes from massively parallel

1281   sequence data. Proc Natl Acad Sci U S A. 2011;108: 1513–1518.

1282   doi:10.1073/pnas.1017351108

1283   57.   Bankevich A, Nurk S, Antipov D, Gurevich A a., Dvorkin M, Kulikov AS, et al.

1284   SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell

1285   Sequencing. J Comput Biol. 2012;19: 455–477. doi:10.1089/cmb.2012.0021

1286 58. Kumar S, Jones M, Koutsovoulos G, Clarke M, Blaxter M. Blobology: exploring

1287   raw genome data for contaminants, symbionts and parasites using taxon-

1288   annotated GC-coverage plots. Front Genet. 2013;4: 237.

1289   doi:10.3389/fgene.2013.00237

1290 59. Smit A, Hubley R, Green P. RepeatMasker Open-4.0. 2013-2015 . [Internet].

1291   http://www.repeatmasker.org. 2013. Available: http://repeatmasker.org

1292 60. Slater GSC, Birney E. Automated generation of heuristics for biological

1293   sequence comparison. BMC Bioinformatics. 2005;6: 1–11. doi:10.1186/1471-

1294   2105-6-31

1295 61. Dobin A, Davis C a, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

1296   ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.

1297   doi:10.1093/bioinformatics/bts635

1298 62. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential

1299   gene and transcript expression analysis of RNA-seq experiments with TopHat

1300   and Cufflinks. Nat Protoc. Nature Publishing Group; 2012;7: 562–578.

1301   doi:10.1038/nprot.2012.016

1302 63. Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. BRAKER1:

1303   Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and

1304   AUGUSTUS. Bioinformatics. 2015;32: 767–769.

1305   doi:10.1093/bioinformatics/btv661

1306 64. Stanke M, Schöffmann O, Morgenstern B, Waack S. Gene prediction in

1307   eukaryotes with a generalized hidden Markov model that uses hints from

1308   external sources. BMC Bioinformatics. 2006;7: 62. doi:10.1186/1471-2105-7-

1309   62

65.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson D a, Amit I, et al. Full-

length transcriptome assembly from RNA-Seq data without a reference

genome. Nat Biotechnol. 2011;29: 644–52. doi:10.1038/nbt.1883

66.   Bens M, Sahm A, Groth M, Jahn N, Morhart M, Holtze S, et al. FRAMA⬚: from

RNA-seq data to annotated mRNA assemblies. BMC Genomics. BMC

Genomics; 2016; 1–12. doi:10.1186/s12864-015-2349-8

67.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM.

BUSCO: Assessing genome assembly and annotation completeness with

single-copy orthologs. Bioinformatics. 2015;31: 3210–3212.

doi:10.1093/bioinformatics/btv351

68.   Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL.

Domain enhanced lookup time accelerated BLAST. Biol Direct. 2012;7: 12.

doi:10.1186/1745-6150-7-12

69.   Benson DA, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW.

GenBank. Nucleic Acids Res. 2014;43: D30–D35. doi:10.1093/nar/gku1216

70.   Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a *de novo* assembler for

single-cell and metagenomic sequencing data with highly uneven depth.

Bioinformatics. 2012;28: 1420–1428. doi:10.1093/bioinformatics/bts174

71.   Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, et al.

MITOS: Improved de novo metazoan mitochondrial genome annotation. Mol

Phylogenet Evol. Elsevier Inc.; 2013;69: 313–319.

doi:10.1016/j.ympev.2012.08.023

72.   Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood

trees for large alignments. PLoS One. 2010;5: e9490.

doi:10.1371/journal.pone.0009490

73.   Liu L, Yu L, Edwards S V. A maximum pseudo-likelihood approach for

estimating species trees under the coalescent model. BMC Evol Biol. 2010;10:

302. doi:10.1186/1471-2148-10-302

74.   Shaw TI, Ruan Z, Glenn TC, Liu L. STRAW: Species TRee Analysis Web server.

Nucleic Acids Res. 2013;41: 238–241. doi:10.1093/nar/gkt377

75.   Ranwez V, Harispe S, Delsuc F, Douzery EJP. MACSE: Multiple alignment of

coding SEquences accounting for frameshifts and stop codons. PLoS One.

2011;6: e22594. doi:10.1371/journal.pone.0022594

76.   Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A. Selecting optimal

partitioning schemes for phylogenomic datasets. BMC Evol Biol. 2014;14: 82.

doi:10.1186/1471-2148-14-82

77.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: a

software platform for Bayesian evolutionary analysis. PLoS Comput Biol.

2014;10: e1003537. doi:10.1371/journal.pcbi.1003537

78.   Heled J, Drummond AJ. Bayesian inference of species trees from multilocus

data. Mol Biol Evol. 2010;27: 570–80. doi:10.1093/molbev/msp274

79.   Sanderson MJ. Estimating absolute rates of molecular evolution and

divergence times: a penalized likelihood approach. Mol Biol Evol. 2002;19:

101–109. doi:10.1093/oxfordjournals.molbev.a003974

80.   Paradis E, Claude J, Strimmer K. APE: Analyses of phylogenetics and evolution

in R language. Bioinformatics. 2004;20: 289–290.

doi:10.1093/bioinformatics/btg412

81.   Kaduk M, Sonnhammer E. Improved orthology inference with Hieranoid 2.

Bioinformatics. 2017;5: btw774. doi:10.1093/bioinformatics/btw774

82.   Sonnhammer ELL, Östlund G. InParanoid 8: Orthology analysis between 273

proteomes, mostly eukaryotic. Nucleic Acids Res. 2015;43: D234–D239.

doi:10.1093/nar/gku1203

83.   Edgar RC. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics. 2010;26: 2460–2461. doi:10.1093/bioinformatics/btq461

84.   Eddy SR. Accelerated profile HMM searches. PLoS Comput Biol. 2011;7.

doi:10.1371/journal.pcbi.1002195

85.   Bernardes JS, Vieira FRJ, Zaverucha G, Carbone A. A multi-objective

optimization approach accurately resolves protein domain architectures.

Bioinformatics. 2015;32: 345–353. doi:10.1093/bioinformatics/btv582

86.   Das S, Sillitoe I, Lee D, Lees JG, Dawson NL, Ward J, et al. CATH FunFHMMer

web server: Protein functional annotations using functional family

assignments. Nucleic Acids Res. 2015;43: W148–W153.

doi:10.1093/nar/gkv488

87.   Sillitoe I, Lewis TE, Cuff A, Das S, Ashford P, Dawson NL, et al. CATH:

Comprehensive structural and functional annotations for genome sequences.

Nucleic Acids Res. 2015;43: D376–D381. doi:10.1093/nar/gku947

88.   Librado P, Vieira FG, Rozas J. BadiRate: Estimating family turnover rates by

likelihood-based methods. Bioinformatics. 2012;28: 279–281.

doi:10.1093/bioinformatics/btr623

89.   Larkin M a., Blackshields G, Brown NP, Chenna R, Mcgettigan P a., McWilliam

H, et al. Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23: 2947–

2948. doi:10.1093/bioinformatics/btm404

90.   Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE:

1383   Functional classification of proteins via subfamily domain architectures.

1384   Nucleic Acids Res. 2017;45: D200–D203. doi:10.1093/nar/gkw1129

1385   91.   Parker J, Tsagkogeorga G, Cotton J a, Liu Y, Provero P, Stupka E, et al. Genome-

1386   wide signatures of convergent evolution in echolocating mammals. Nature.

1387   Nature Publishing Group; 2013;502: 1–9. doi:10.1038/nature12511

1388   92.   Kosakovsky Pond SL, Frost SDW, Muse S V. HyPhy: Hypothesis testing using

1389   phylogenies. Bioinformatics. 2005;21: 676–679.

1390   doi:10.1093/bioinformatics/bti079

1391   93.   Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. RELAX:

1392   Detecting relaxed selection in a phylogenetic framework. Mol Biol Evol.

1393   2014;32: 1–13. doi:10.1093/molbev/msu400

1394   94.   Benjamin DJ, Berger JO, Johannesson M, Nosek BA, Wagenmakers E-J, Berk R,

1395   et al. Redefine statistical significance. Nat Hum Behav. 2017;

1396   doi:10.1038/s41562-017-0189-z

1397   95.   Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al.

1398   The GeneMANIA prediction server: biological network integration for gene

1399   prioritization and predicting gene function. Weaver J, Anija K, Daspit T,

1400   editors. Nucleic Acids Res. Oxford University Press; 2010;38: W214–W220.

1401   doi:10.1093/nar/gkq537

1402   96.   Falcon S, Gentleman R. Using GOstats to test gene lists for GO term

1403   association. Bioinformatics. 2007;23: 257–258.

1404   doi:10.1093/bioinformatics/btl567

97.   Webb B, Sali A. Protein structure modeling with MODELLER. Methods Mol

Biol. 2014;1137: 1–15. doi:10.1007/978-1-4939-0366-5_1

98.   Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. Comparison

of simple potential functions for simulating liquid water. J Chem Phys. 1983;

doi:10.1063/1.445869

99.   Case DAA, Babin V, Berryman JTT, Betz RMM, Cai Q, Cerutti DSS, et al.

AMBER14. University of California, San Francisco. 2014. doi:10.1007/s13398-

014-0173-7.2

100.   Hess B, Kutzner C, Van Der Spoel D, Lindahl E. GRGMACS 4: Algorithms for

highly efficient, load-balanced, and scalable molecular simulation. J Chem

Theory Comput. 2008; doi:10.1021/ct700301q

101.   Sousa Da Silva AW, Vranken WF. ACPYPE - AnteChamber PYthon Parser

interfacE. BMC Res Notes. 2012; doi:10.1186/1756-0500-5-367

102.   Darden T, York D, Pedersen L. Particle mesh Ewald: An N·log(N) method for

Ewald sums in large systems. J Chem Phys. 1993; doi:10.1063/1.464397

103.   Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. LINCS: A Linear Constraint

Solver for molecular simulations. J Comput Chem. 1997;

doi:10.1002/(SICI)1096-987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-H

104.   Parrinello M, Rahman A. Polymorphic transitions in single crystals: A new

molecular dynamics method. J Appl Phys. 1981; doi:10.1063/1.328693

1425