Genomic signature of shifts in selection in a sub-alpine ant and its physiological adaptations

Cicconardi Francesco[1], Krapf Patrick[2], D'Annessa Ilda[3], Gamisch Alexander[2,4], Wagner Herbert C[2], Nguyen Andrew D[5], Economo Evan P[6], Mikheyev Alexander S[7], Guénard Benoit[8], Grabherr Reingard[9], Arthofer Wolfgang[2], di Marino Daniele[10], Steiner Florian M[2], Schlick-Steiner Birgit C[2]

[1] Department of Zoology, University of Cambridge, Cambridge, UK
[2] Department of Ecology, University of Innsbruck, Innsbruck Austria
[3] Istituto di Chimica del Riconoscimento Molecolare, CNR (ICRM-CNR), Milan, Italy
[4] Department of Biosciences, University of Salzburg, Salzburg, Austria
[5] Department of Entomology and Nematology, University of Florida, USA
[6] Biodiversity & Biocomplexity Unit, Okinawa Institute of Science & Technology, Onna, Japan
[7] Ecology and Evolution Unit, Okinawa Institute of Science & Technology, Onna, Japan
[8] School of Biological Sciences, The University of Hong Kong, Hong Kong, China
[9] Institute of Biotechnology, University of Natural Resources and Life Sciences, Vienna, Austria
[10] Polytechnic University of Marche, Department of Life and Environmental Sciences, Ancona, Italy
Corresponding author: francicco@gmail.com

Abstract

Adaptation to climate can drive variation in diversification rates and species richness. Understanding how organisms adapt to extreme environments can therefore provide insightful case studies for both evolutionary biology and climate-change biology. Here, we take advantage of the vast diversity of lifestyles in ants to identify genomic signatures of adaptation to extreme habitats such as high altitude. We hypothesised two parallel patterns would occur in a genome adapting to an extreme habitat: *i)* strong positive selection on genes related to adaptation and, *ii)* a relaxation of previous purifying selection. We tested this hypothesis by sequencing the high-elevation specialist *Tetramorium alpestre* and four related species. In support of our hypothesis, we recorded a strong shift of selective forces in *T. alpestre*. We further disentangled candidate molecular adaptations in both gene expression and protein-coding sequence that were identified by our genome wide analyses. In particular, we demonstrate that *T. alpestre* has *i)* a derived level of expression for *stv* and other heat-shock proteins in chill shock tests, and *ii)* enzymatic enhancement of Hex-T1, a rate-limiting regulatory enzyme that controls the entry of glucose into the glycolytic pathway. Together, our analyses highlight the adaptive molecular changes that support colonisation of high altitude environments.

Introduction

Adaptation of organisms to climate drives variation in diversification rates and species richness among clades. Their climatic niche influences a species' occurrence in space and time (Soberón 2007) and thus is critically important for both speciation (e.g., ecological speciation through climatic niche divergence) and extinction (e.g., due to climate change). Therefore, understanding how organisms manage thermal adaptation is important in terms of both evolutionary biology and climate-change biology, considering the potential world-wide loss of ecological niches (Lamprecht et al. 2018; Rogora et al. 2018). In this context, high elevations, characterized by a short growing season and low annual minimum and mean temperatures, with high daily fluctuation temperatures (Körner et al. 2011), are an important open-air laboratory to study speciation and adaptation to cold habitats; and compared to the great effort in gene-based studies on cold tolerance of model organisms, a lesser effort has been put towards understanding truly cold-tolerant animals at the genomic level (Clark and Worland 2008; Parker et al. 2018). This is even more evident when considering that among the approximately 600 sequenced insect genomes available today only four belong to species ecologically restricted to high altitudes or Antarctic habitats (Keeling et al. 2013; Kelley et al. 2014; Macdonald et al. 2016; Cicconardi et al. 2017a).

As yet, potential patterns of genomic signatures in organisms adapting to more extreme habitats, such as high elevations, have not been identified and, more generally, there is no theory that predicts the rates of genomic change for extreme habitats. Here, we hypothesise two parallel patterns to occur in a genome adapting to an extreme habitat. On the one hand, strong positive selection on genes related to adaptation, such as genes involved in metabolic pathways, and on the other hand, a

relaxation of previous selecting forces, as the conditions of the previous niche are

lacking in the new niche. The latter should lead to a reduced number and/or a

different set of genes under purifying selection. Specifically, in the case of high-

elevation habitats, some heat-shock proteins (HSPs), necessary for coping with

extreme heat, should be under relaxation, because heat resistance is not strongly

selected for in the average alpine species. A problem with the hypothesis of these two

parallel patterns is that there is no test available for the functional consequences of

the loss or gain of a specific gene. There are often many pleiotropic effects for genes

that make interpretations difficult, and HSPs, expressed upon exposure to stress or

during development and growth, are likely to fall into this category. Therefore, we

expect signatures of strong positive selection in metabolic genes, and an increase in

the relaxation rates in other genes, without being able to specify which these might

be. Identifying such genes will help to set up hypotheses to test in the future. A

comparative approach offers the strongest method for testing our hypothesis, by

comparing genomes of species closely related to each other, but divergent in terms of

adaptation to different environments. However, this approach can be limited by

insufficient niche divergence within a group or the number of genomes sequenced.

Ants are emerging as a leading system for comparative genomics due to the constantly

increasing number of available genomes. This resource offers the opportunity to

improve the accuracy of orthology detection (Nygaard et al. 2016), to scan for specific

mutations, candidate genes, and patterns of acceleration and relaxation in the ant

genomes associated with adaptation to cold habitats.

Ants are key species in the Earth's terrestrial ecosystems. They comprise more

than 15,442 described species and display an impressive diversity of lifestyles

(Hölldobler and Wilson 1990; Bolton 2018; Seifert 2018). Ants are especially notable

among insects for their ecological dominance as predators, scavengers, and indirect

herbivores. They compose at least one third of the entire insect biomass (Wilson 1990)

and colonise all kinds of habitats, including thermobiologically challenging

environments. The formicine ant *Melophorus bagoti*, for instance, is active during the

hottest periods of the summer day, when air temperatures at ant height exceed 50 ˚C

(Christian and Morton 1992). On the other hand, for example, the myrmicine ant

Tetramorium alpestre inhabits the montane and subalpine belt of the Central and

South European mountain systems, with the Alps as its main distribution area (Wagner

et al. 2017; Steiner et al. 2010). This species lives mainly between 1300 and 2300 m

above sea level (a.s.l.), forages below the ground, and nests are established in cool

grassland under stones, in moss, rootage, and dead wood, especially subalpine and

alpine grass mats (Seifert 2018). Since *T. alpestre* belongs to a species complex in

which no relation between the ecological niche and worker morphology has been

detected, searching for adaptive phenotypical differences outside of morphology, for

example physiology, has already been suggested (Wagner et al. 2017). This species is

also relevant because, among the 79 Palearctic species of the worldwide distributed

genus *Tetramorium* (Bolton 2016), it is one of only two species to form colonies with

multiple queens (polygynous colonies), showing intermediate states of aggression and

a transition from multicoloniality to supercoloniality (Krapf et al. 2018; Seifert 2018).

Tetramorium alpestre is thus a system well suited for studying both genomic

adaptation to high elevation and socio-behavioural evolution.

Here, we tested our hypothesis of both increased diversifying and reduced

purifying (relaxing) selection resulting from adaptation to an extreme niche using *T.*

alpestre. We newly sequenced its genome and those of four related *Tetramorium*

species with diverging ecological niches: *T. immigrans*, a species of the *T. caespitum*

complex to which *T. alpestre* also belongs to, in sympatry with *T. alpestre*, but distinct

in its altitudinal and ecological habitat; *T. parvispinum*, restricted to mountain forest

habitats of the Austral-Asian and Indo-Malayan subregions (Liu et al. 2015); and *T.*

bicarinatum and *T. simillimum*, invasive generalist species which occur in warmer

habitats (Bertelsmeier et al. 2017; Guénard et al. 2017). Our goals were to define the

ecological niche of these ant species based on environmental data, and subsequently

to perform gene family expansion/contraction and protein-coding scans for signatures

of diversifying and relaxing selection. We also annotated the five HSP subfamilies for

19 ant species to test for possible shifts in selection acting on these gene families.

Among the signatures of diversifying selection detected, we performed two

experiments in *T. alpestre* and its relative, *T. immigrans*: we assessed (i) chill-shock

triggered gene-expression patterns of *starvin* (*stv*), a Hsp70 modulator, and certain

HSPs involved in recovering from extreme cold, and (ii) the temperature dependence

of the enzyme activity of Hexokinase type 1 (Hex-T1), a rate-limiting and regulatory

enzyme that controls the entry of glucose into the glycolytic pathway, one of the most

conserved and essential hexokinase isoenzymes (Jayakumar et al. 2007). The genomic

data presented here contribute to a foundation for studying insect genome evolution,

particularly also in the light of climate change.

Figure 1. *a*) Venn diagram displaying overlap in orthologous genes in the five *Tetramorium* spp. + *V. emeryi* (Crematogastrini). Coloured circles with number represent phylogenetic splits in the nuDNA (N^{nu}) and mitochondrial phylogenies in *b*; *b*) The dated Crematogastrini nuclear phylogeny (above). Branch colour are based on the concordance factors of the two datasets (nuDNA and mtDNA). On each branch, the numbers of expanded (green arrows) or contracted (purple arrows) genes is shown, inferred from observed OGs sizes at terminal branches. In the bottom, the global δ18O (‰) derived from analyses of two common and long-lived benthic taxa are given, *Cibicidoides* and *Nuttallides*, which reflect the global deep-sea oxygen and carbon isotope and thus the temperature (from Zachos et al., 2001). Bars represent the 95% confidence interval for the BI analysis, circles with numbers the phylogenetic splits in the nuDNA (N^{nu}) and mtDNA (N^{mt}) phylogenies. *c*) Boxplots of the five bioclimatic variables significantly different for *T. alpestre* and the other ant species.

Results

Climatic niches

A dataset contains 1835 localities representing 17 species was assembled, with the number of localities per species ranging from 9 to 591 (Table S1). Of the 19 bioclimatic variables, four (bio1: Annual Mean Temperature, bio5: Max Temperature of Warmest Month, bio8: Mean Temperature of Wettest Quarter, bio10: Mean Temperature of Warmest Quarter) differentiated between *T. alpestre* and other ants (adjusted *P*-values < 0.001) (Figure 1a). In detail, *T. immigrans* and *T. alpestre* occupy habitats that

are colder in the growing season than those of the remaining species, with *T. alpestre*

revealing even colder habitats compared with *T. immigrans*.

Sequencing, assembly, and annotation of the *Tetramorium* genomes

The combination of overlapping paired-end libraries with very high coverage, standard

short paired-end libraries, and mate pair libraries resulted in an overall complete

assembly of the five *de novo Tetramorium* genomes, at the level of both contiguity

and scaffolding. In particular, the *T. alpestre* genome contiguity (contig N50) is the 3[rd]

best among all ant genomes available (Table S2). The analysis of the *k*-mer frequency

distribution generated was unimodal (Figure S1; Supplementary Material online), and

an analysis of the distribution of GC content across the five *de novo Tetramorium*

genomes revealed similar distributions with a mean of 38%, that is, slightly less than

that found in *V. emeryi* (42%) (Figure S2). No significant bias was recovered

considering GC content and coverage (Figure S3). All five *de novo* draft assemblies

were ~241 Mb in length, comparable in size with the average of all ant genomes

available (~278 Mb) (Table S2). The total *T. alpestre* genome size was 245.72 Mb,

including 18 Mb of gaps and unknown characters (N/X), slightly smaller than the

estimated genome size by flow-cytometry of 291.84 Mb +/- 1.76 Mb (n=12). Repetitive

elements made up similar proportions of each sequenced genome (*T. alpestre* = 18%,

T. immigrans = 20%, *T. bicarinatum* = 21%, *T. simillimum* = 20%, *T. parvispinum* = 23%;

Table S2), indicating a strong correlation between genome size and total interspersed

repeat content (Pearson ρ = 0.976, *P*-value < 0.005; Figure S5). For the gene

annotations in *T. alpestre*, a combinatorial approach of unsupervised RNA-seq-based,

homology-based, *ab initio*, and finally *de novo* methods were used. By doing so, we

approached the number of predicted protein-coding genes in the phylogenetically

closest annotated species (15,085 genes predicted vs. 14,872 in the *V. emeryi* genome,

Table S2). This improved the homology-based annotation of the remaining four

Tetramorium genome assemblies that did not differ much in terms of estimated gene

content, ranging from 16k to 15k annotated loci (Table S2). BUSCO analyses estimated

overall good representation, with recovered genes in between the 97.6% and 99.9%

(Table S2; Figure S6), and all statistics on mRNA, coding regions (CDS), exon, and intron

length gave highly similar and overlapping distributions among all *Tetramorium*

species and *D. melanogaster* as a reference (Figure S7). Also, the transcript

completeness gave good results with a comparable distribution of percentage of

alignment across species (Figure S8).

The orthology search analysis produced a total of 7195 OGs between *D.*

melanogaster and Hymenoptera; 3261 of these represented scOGs present in all 22

species analysed. Restricting the orthology analysis to *Tetramorium* spp. and *V. emeryi*

resulted in 6494 OGs; among *Tetramorium*, the highest fraction was recovered

between *T. alpestre* and *T. immigrans* with 729 OGs, followed by *T. simillimum* and *T.*

parvispinum (540 OGs) (Figure 1a). For all *Tetramorium* species., the fraction of

orthologous genes with at least a match with one of the three outgroups (*A. mellifera*,

N. vitripennis, *D. melanogaster*) was on average 7286, with the highest in *T.*

alpestre (7879) and the lowest in *T. parvispinum* (6977), in line with the average value

found in ants (7831).

Assembly and annotation of ant mitogenomes and their gene translocations

We assembled complete mtDNA genome sequences from 24 ant species. Because of

the differences among SRA datasets, we used different numbers of reads to assemble

the complete sequences. The new mtDNA genomes were slightly longer (on average,

17k vs. 16k; Table S2), probably due to the reconstruction of the D-loop and ribosomal

genes, which are sometimes harder to assemble. We slightly mis-assembled a coding

gene (*nad2*) in the *T. bicarinatum* genome and a few other tRNA loci in other genomes

(Table S2). The average GC-content for coding genes was 23.9 ± 3.0% and varied

slightly among taxa with the lowest in *Formica fusca* (19.0%) and the highest in

Leptomyrmex pallens (32.8%) (Table S2). While almost all species had the typical gene

order and orientation, some rearrangements were observed. All myrmicine ants were

characterized by a translocation of *trnV* after *rrnS*; *Solenopsis* spp. had a swapped

position of *rrnS* and *trnN*; *Monomorium pharaonis* had at least three rearrangements:

the inversion of *nad6*, *cob*, and *trnS2*, the translocation of *trnE* upstream to *trnA*, and

the translocation of *trnR* upstream to *trnE*; *Tetramorium* spp. underwent a swap of

trnR and *trnN* (Figure S9).

Phylogenetic analyses of *Tetramorium* spp*., their diversification dating, and the

Miocene-Pliocene origin of the lineage leading to *T. alpestre*

The estimated divergence ages from both datasets (mtDNA, nuDNA) were congruent

for most nodes, except for recent nodes within the Attini (Figure 1b; Figure S10-13).

In this case, the penalized likelihood approach estimates were always more recent

than the estimations made with BI using BEAST. The dating of basal nodes like the

MRCAs (e.g., Crematogastrini, Myrmicinae, and Formicinae) were congruent with a

previous analysis based on more taxa but fewer genes (Moreau and Bell 2013). The

two methods also converged regarding the date of the most recent common ancestor

(MRCA) between *T. alpestre* and *T. immigrans*. In that specific case, PL dated that split

at 6.8 million years ago (mya) within the confidence interval given by BI (Median: 5.15

mya; 95% HPD: 2.66, 8.23).

The phylogenetic relationships were generally highly congruent and with

predominantly strong support (Figure 1b and Figures S10-13). In the ML phylogeny of

scOGs (nuDNA), all nodes were supported with a bootstrap value of 1. Across the

whole phylogeny, only a few nodes had weak signals, resulting in very few topological

disagreements and low coalescent units (CU). Within the clade of Crematogastrini

(*Tetramorium* spp. + *V. emeryi*), the relation of *T. bicarinatum* and *T. simillimum* was

uncertain: nuDNA phylogeny recovered them as sister species with low CU (0.11),

while mtDNA phylogeny placed *T. bicarinatum* as sister to the *T. alpestre-T. immigrans*

cluster, with a bootstrap value of 0.77 and a posterior probability of 0.99.

Contractions of OGs in the *T. immigrans* and *T. alpestre* lineages and expansion of

retrotransposon-related genes in *T. alpestre*

We examined the evolutionary dynamics of OGs looking at more dynamic ranges

corresponding with terminal branches. We observed an increased number of OG

deaths in the *T. parvispinum* lineage (– 642) and in the lineage leading to *T. alpestre*

and *T. immigrans* (– 516). Ortholog groups showed very little expansion in internal

branches of the *Tetramorium* radiation. Expanded OGs were common in terminal

branches, except for *T. parvispinum* with very few expanded OGs (+ 14), suggesting a

possible lack of annotation. *Tetramorium alpestre* showed the highest number of

expanded OGs (+ 364) (Figure 1b), with more than 370 of them (~ 40%) belonging to

genes with reverse transcriptase (PFAM: rve, rve_1, rve_2) or transposase and

retrotransposon activities (PFAM: Retrotrans_gag, Retrotran_gag_2). Among other

PFAM, haemolymph juvenile hormone-binding proteins (PFAM: JHBP; 20 genes), zinc

fingers (PFAM: zf-CCHC, zf-H2C2, zf-met, 32 genes), and sugar transporters (PFAM:

Sugar_tr, 4 genes) were found.

Dual signature of evolutionary pressures in single-copy OGs in the *T. alpestre*

genome

We investigated the impact of evolutionary pressures on one-to-one orthology in the

five *Tetramorium* spp. and 14 other ant species and three outgroups (*Apis mellifera*,

Nasonia vitripennis, and *Drosophila melanogaster*) by computing both mean ω (d_N/d_S)

for terminal branches and relaxing selection (k) within the Crematogastrini.

The measure of the mean branch ω showed an overall high level of purifying

selection along the whole ant phylogeny, including for short branches (Roux et al.

2014; Cicconardi et al. 2017b), with median values of 0.10. However, this distribution

was significantly shifted in *T. alpestre*, which showed a median value of 0.38 (Wilcoxon

rank-sum test '*greater*', P-value $< 2.2e^{-16}$), whilst keeping the proportion of $\omega > 1$ in

each branch unchanged (Figure 2a, S13). The evaluation of k in the five *Tetramorium*

spp. and *V. emeryi* had a mean of median values of 0.99, that is, twice the value in *T.*

alpestre (median $k = 0.47$; Wilcoxon rank-sum test '*less*', P-value $< 2.2e^{-16}$) (Figure 2b).

We investigated the nature of the increased ω (episodic diversifying selection) and

decreased k (relaxed selection) in *T. alpestre* by combining the two measures. Two

evolutionary trajectories appeared to act simultaneously (two tails of the distribution,

Figure 2c): One trajectory seems to have led to episodic diversifying selection,

Figure2. *a*) Boxplots showing the distribution and median (vertical line) of mean ω (d_N/d_S) rates (square shapes) in
terminal branches of *Tetramorium* spp. and *V. emeryi* (Crematogastrini) in scOGs, and the proportion of genes for
which ω is higher than one. *b*) Boxplots overlapped by violin plots showing the distribution of *k* in Crematogastrini.
The distribution in *T. alpestre* is clearly bimodal as in *T. immigrans*, but with a much more skewed distribution
towards zero values. In both sections, asterisks indicate the degree of significance between *T. alpestre* and the
other species (Wilcoxon rank-sum tests). *c*) Scatter plots for each Crematogastrini species included, showing for
each scOG branch both values of ω and *k*. The green dots are scOGs with uncorrected *P*-values associated with
diversifying selection less than 0.05. The distribution in *T. alpestre* shows the longest tail corresponding to very low

values of *k* and high values of *ω*. It is also notable a much wider and scattered distribution of scOGs with high *k* and
ω compared with all other species.
285

promoting the fixation of nonsynonymous mutations with presumably advantageous

fitness effects for genes with increased values of *ω* and *k*, also present in the other

species. The other trajectory showed a relaxation of the overall purifying selection in

genes with increased values of *ω* and decreased values of *k*, only mildly present in *T.*

immigrans.

Figure 3. For each enriched Gene Ontology (GO) term category *a*) the median values of *ω*; and *b*) *k* for genes
enriching the specific category. *Tetramorium* spp. are colour coded, the other 14 species, merged to obtain a
background signal, are in gray. Blue circles are scOGs tested for each GO term, red circles are scOGs returned as
significant. Circle sizes are proportional to the number of genes. To be noted are the overall extremely higher

values of ω and lower values of k in *T. alpestre* branches compared to the other species. Seldomly also *T. immigrans*
shows few GO terms with high ω and low k values, in particular the Apical junction assembly (GO:0043297) shows
and inverted trend between *T. alpestre* and *T. immigrans*.
298

After a stringent correction for multiple testing, we found 175 scOGs with a

putative signature of diversifying selection (adjusted *P*-values < 0.005; Table S5), with

50 genes enriching 13 biological process terms (*P*-values < 0.005; Figure 3a, Table S6),

nine related to cell development (e.g., cell migration involved in gastrulation,

photoreceptor cell development, cell fate determination) and organization (e.g.,

regulation of organelle organization), two related to signalling pathways (enzyme

linked receptor protein signalling pathway and torso signalling pathway), one to heart

development, and one to organic hydroxy-compound metabolic processes. The most

representative genes were *Pten* and *Gbeta13F*, involved in six biological processes,

DCTN1-p150 and *dsh*, involved in five of them. All key genes are involved in the

proliferation and division of cells, especially neurological stem cells (*Gbeta13F*),

respiratory system development (*Pten* and *dsh*), and the serine/threonine-protein

kinase *polo*, present in four enriched Gene Ontology (GO) terms. In the catalytic

domain of this protein (255 aa), we found five amino acid substitutions compared with

T. immigrans, three of them corresponding to active sites (*TalpPolo$_{G29V}$*, *TalpPolo$_{G30R}$*,

TalpPolo$_{S104R}$). We also tested for possible enrichment in KEGG pathways and found

three significant pathways (*P*-values < 0.02; Table S7), all involved in the metabolic

processes of galactose metabolism and the glycolysis pathway, the pentose

phosphate pathway, and galactose metabolism, with five enzymes under diversifying

selection: a glucose-6-phosphate (*CG9008*), the hexokinase type 1 (*Hex-T1*), a

phosphofructokinase (*Pfk*), a transketolase (*CG8036*), and a phosphoglucosemutase

(*pgm*).

132 scOGs had a signature of putative relaxation (*P*-values < 0.005; Table S8).

While the absolute number of scOGs under relaxation was slightly lower than the

number of genes under diversifying selection, the gene-set enrichment tests revealed

many more biological processes significantly enriched: 70 terms (*P*-values < 0.005;

Figure 3b, Table S9) enriched by 118 genes. Fifteen terms were related to the

regulation of cellular development, signal transduction, and cell communication (e.g.,

regulation of response to stimuli, regulation of cell communication, regulation of Ras

signal transduction), 12 terms related to neurogenesis, axon development, and

synaptic transmission (e.g., neuron differentiation, trans-synaptic signalling, axon

guidance), seven terms were related to cell morphogenesis and junction organization,

and many other related to response to stimuli (e.g., chemotaxis, response to light) or

related to adult development and morphogenesis (e.g., anatomical structure

morphogenesis, molting cycle, chitin-based cuticle). The enrichment of KEGG

pathways revealed three pathways: mRNA surveillance pathway, spliceosome, and

protein processing in endoplasmic reticulum (*P*-values < 0.05; Table S10).

The Hex-T1 activity, its model structure, and molecular dynamics simulations in *T.

alpestre and *T. immigrans***

As we describe in the previous section, the enrichment for KEGG pathways led to the

identification of three metabolic pathways, all involved in the metabolism of sugars,

with five genes under selection. We therefore aimed at the experimental validation of

their functionalities comparing their activity in *T. alpestre* and the most closely related

species in this dataset, *T. immigrans*. We tried to assay the four enzymes, but

unfortunately, we were able to correctly synthesise only Hex-T1. The assay was

designed to measure the efficiency of Hex-T1 in both species and in a temperature

gradient, at 6, 16, 26, and 36 °C. The assays showed that the Km value was always

347
Figure 4. a) Scatter plot of Km score across the temperature gradient in the two species, *T. immigrans* on top and
T. alpestre on the bottom. Lines represent average values with confidence intervals. Significances: *** < 0.001; **
< 0.001; * < 0.05. *b*) Overview of the homology-modeled structure of Hex-T1 in *T. alpestre* showing the two
Hexokinase domains, PFAM00349 (green) and PFAM03727 (pink). The lower panels show the location of TalpHex-
T1$_{G74}$ (left) and TimmHex-T1$_{E70}$ (right) as well as the arginine residue that forms a salt bridge with E70 (dashed line).
c) Per-Residue Root Mean Square Fluctuation (RMSF) calculated along 500 ns trajectories carried out at 6 °C (upper
panel) and 26 °C (lower panel) for TalpHex-T1 (left panel, blue line) and TimmHex-T1 (right panel, light blue line).
The location of the residues found mutated between the two sequences are indicated. *d*) Time evolution of the
atomic distance between the centre of mass of glucose and the lateral chain nitrogen atom of TalpHex-T1$_{K168}$ (blue
line) and TimmHex-T1$_{K164}$ (light blue line) in the binding pocket calculated along the 500 ns trajectories carried out
at 6 °C (upper panel) and 26 °C (lower panel). *e*) Representative configurations extracted from the simulations at
26 °C showing the different orientation of TalpHex-T1$_{K168}$ and TimmHex-T1$_{K164}$, with the first one being the only
one able to bind the glucose (dashed line).

lower in *T. alpestre* than in *T. immigrans* (Wilcoxon rank-sum test 'less', *P*-value <

0.00004, Figure 4a), with values ranging from 0.08 mM to 0.01 mM in *T. alpestre*

versus 0.15 mM to 0.05 mM in *T. immigrans*, from the highest to the lowest

temperature; reflecting that less substrate concentration was needed for the *T.*

alpestre enzyme to reach half of its maximal reaction speed, that is, Hex-T1 of *T.*

alpestre was always more efficient than of *T. immigrans*. Although the number of

replicates per condition was small, we performed statistical tests to attempt obtaining

insight in the enzymatic activity differences within species at different temperatures

and in their variance. Interestingly, it seems that for both species there is an

improvement in the Km values shifting towards the lowest temperatures (6 °C *vs* 16

°C; two-way ANOVA followed by a Tukey's multiple comparison; adjusted *P*-values <

0.05), and a possible deterioration at higher temperatures in *T. alpestre*, from

0.04±0.01 mM (26 °C) to 0.08±0.01 mM (36 °C) (two-way ANOVA followed by a Tukey's

multiple comparison; adjusted *P*-value < 0.0002). An F test (VAR.TEST() implemented in

R) to compare the variances of the enzymatic activities (Km values) between the two

species – by pooling the replicates at 16 and 26 °C – suggested that there is a possible

significant difference in their variance (*P*-value = 0.015), with the confidence interval

of Hex-T1 activity in *T. immigrans* twice wider as in *T. alpestre*.

The Hex-T1 enzyme is made of two structurally similar domains, Hexokinase 1

and 2 (PFAM00349 and PFAM03727), and an N-terminus that is highly variable both in

terms of length and amino acidic composition (Figure 4b). The two domains are

relatively well conserved across all ants, other Hymenoptera and *D. melanogaster*.

Excluding the hypervariable N-terminus, and considering the two functional domains,

the amino acid sequences of *T. alpestre* and *T. immigrans* only differ by two amino-

acidic changes: a glutamic acid (E) replaced by a glycine (G) in position 74 (TalpHex-

T1$_{E74G}$) (Figure 4b inserts) and an asparagine (N) replaced by a lysine (K) in position

344 (TalpHex-T1$_{N344K}$). While the second change probably does not entail a significant

functionality effect, the *TalpHex-T1$_{E74G}$* instead may be significant. Because of the lack

of the long chain and the negative charge, this modification could give the protein a

higher flexibility and therefore an overall higher kinetics. We tested this hypothesis by

simulating the protein dynamics in the presence of the glucose substrate at different

temperatures, i.e. 6 and 26 °C, with the intent of mirroring the condition in the in vitro

assay. In the simulations we can see a dual effect. For TalpHex-T1$_{N344K}$, located at the

C-terminal of an α-helix forming a helix-turn-helix motif (VSETE**K**DPKG), the Root

Mean Square Fluctuation (RMSF) revealed a lower degree of fluctuation in TalpHex-

T1 (Figure 4c). This is possibly due to the alternation of positive and negative charges

creates a salt-bridges network that stabilizes the whole motif (data not shown). In

particular TalpHex-T1$_{K344}$ is involved in a stable interaction with TalpHex-T1$_{E341}$ (data

not shown), while TimmHex-T1$_{N340}$ is not available for the formation of salt bridges

and mainly establishes hydrogen bonds with TimmHex-T1$_{S336}$ or TimmHex-T1$_{E337}$ at 6

or 26 °C, respectively. Although the *T. alpestre* salt bridge brings a higher flexibility of

the helix-turn-helix motive, it does not interfere with the catalytic site, given that this

motif is far away in the C-terminal region of the protein.

In contrast, at the other region (TalpHex-T1$_{E74G}$), the mutation-caused amino

acid change allows the loop to explore a broader conformational space equipping it

with a higher degree of flexibility already at 6 °C (Figure 4c). Nevertheless, by

increasing the temperature, TalpHex-T1 was subjected to an even higher degree of

fluctuation, once again confirming the intrinsic flexibility led by the glycine. Notably,

the increased flexibility is also transferred to residues relatively far from the mutation,

in particular in regions near the active site (168-169), causing a different pattern of

interaction with the substrate (Figure 4d, e). Indeed, TalpHex-T1 established a

stronger interaction with glucose at both temperatures with respect to TimmHex-T1

(Figure 4e).

The heat-shock proteins in ants and their evolution in *T. alpestre*

Members of the five subfamilies of the heat shock proteins (HSPs: Hsp90s, Hsp70s,

Hsp60s, Hsp40s, and sHsps) were fully recovered and characterized in 19 ant species.

Overall, a strong purifying selection was observed within subfamilies (Figure 5a, Table

S11). In the Hsp90 gene family (*trap1*, *gp93*, and *Hsp83*) *Hsp83* was found duplicated

within the lineage of *W. auropunctata* (inparalog), as well as the *T. parvispinum gp93*;

duplication that probably occurred before emergence of the lineage (outparalog)

(Figure S14). Members of the Hsp70 family were recovered for all species with 11

ortholog groups, with few duplications in *Hsc70-3* and *Hsc70-5* (two inparalogs and

two outparalogs, respectively). We confirmed the lack of Hsp70s in ants (Nguyen et

al. 2016) and a multiple duplication of *Hsc70-4* in Hymenoptera. One paralog seemed

to be lost in many species while the other two copies were still present in almost all

species, with the presence of other outparalogs in various species, mostly in Attini

(Figure S15). The chaperonin Cpn60/TCP-1 family (Hsp60) had nine OGs; six inparalogs

were found in *CCT7, CCT3* and four outparalogs in *Tcp1a, CCT8*. The heat shock protein

40 kDa/Chaperone DnaJ was the largest subfamily, with 37 well-defined OGs. Its

turnover of duplication was relatively small, and only six OGs showed some level of

duplication. The small HSPs (sHsps) showed the expansion of the lethal (2) essential

for life (*l(2)efl*) gene paralog of *D. melanogaster*, with five distinct OGs in all

Hymenoptera. *L(2)efl3* seemed to be retained by only in 11 species, and with *l(2)efl4*,

they are the two most divergent OGs, with ω equal to 0.23 and 0.33, respectively. The

less divergent were *Hspb1* and *l(2)efl5*, for which ω was 0.11 and 0.07, respectively.

Overall, the sHsps showed a significantly higher distribution of ω (median = 0.22, *P*-

value = 0.013), followed by Hsp40s, which also showed a wide distribution of ω. In

contrast, Hsp90s, Hsp70s, and Hsp60s seemed to be very stable and under a very high

purifying selection.

To understand other possible mechanisms of adaptation of *T. alpestre* to the

cold, we computed the mean values of ω and k for all branches leading to

Tetramorium spp. and *V. emeryi*, integrating 13 ant species, and scanned for

diversifying and relaxed selection specifically in *T. alpestre*. The first part of the

analysis showed a similar landscape as observed in scOGs. Among the five

Tetramorium species and *V. emeryi*, five species display a distribution of the mean-

branch ω ranging between 0.07 (*T. immigrans*) and 0.13 (*T. parvispinum*); *T. alpestre*

instead showed significant 2.7-fold higher value of the median mean-branch ω (ω =

0.36, Wilcoxon rank-sum test '*greater*', adjusted *P*-values < 0.004, Figure 5b). The

distribution of k among HSPs in the six species showed median values of k all close to

the expected (k = 1), between 0.90 in *V. emeryi* and 1.05 in *T. simillimum*, again with

the exception of *T. alpestre*, where the median k = 0.63, 1.4-fold lower than the lowest

median k (Wilcoxon rank-sum test '*less*', adjusted *P*-values < 0.05, Figure 5c). The scan

for diversifying positive and relaxed selection in branches of the HSPs of *T. alpestre*

Figure 5. *a*) Boxplots showing the distribution and median (horizontal black line) of mean ω (d_N/d_S) rates for each of the HSP subfamily. *b*) Violin plot showing the distribution of the mean ω values; and *c*) *k* in the terminal branches of the Crematogastrini (from up to bottom: *T. alpestre*, *T. immigrans*, *T. bicarinatum*, *T. simillimum*, *T. parvispinum*, *V. emeryi*). *d-f*) Scatter plot of target gene (*stv*, *Hsc70-4d*, *Hsc70-4e*) vs. housekeeping gene (HK) concentrations; concentrations given as total number of cycles minus cycle threshold (Ct) values, that is, higher values represent higher concentrations. Straight lines are linear regressions for each target gene against HK using the different recovery times separately; based on one-way analysis of covariance followed by correction for multiple comparison, just lines for those treatment / recovery-time combinations are shown for which significances for just a single species arose, that is, *T. alpestre* coma vs. non-coma after 0.5 h (*d*, *e*; no difference in *T. immigrans*) or for which the two species differed, that is, *T. alpestre* coma vs *T. immigrans* coma after 4.0 h (*f*).

showed no gene under putative diversifying positive selection, while six loci were

identified to be under putative relaxing selection: one Hsp90 (*Trap1*), two Hsp60

(*CCT4, CCT5*), and three Hsp40s (*Sec63, DnajC8, DnajC11*) (*k* < 0.14, adjusted *P*-values

< 0.005). Using these genes in prediction interaction networks, we found that the two

Hsp60s interacted with other 21 genes, giving 17 significant enriched functions (FDR

< 0.05), mainly related to protein folding and microtubule, spindle organization; the

three Hsp40s interacted with 21 genes, giving two enriched functions (FDR < 0.05),

related to response to heat and response to temperature stimulus (Figure S19).

Chill shock effect, recovery times, and expression patterns of *stv* and HSPs in *T.*

alpestre* and *T. immigrans

Among the genes with a putative signature of diversifying selection, we also found

starvin (*stv*), a BAG-family member which seems to be implicated in Hsp70 ATPase

activity during cold recovery in *D. melanogaster* (Colinet and Hoffmann 2010).

Therefore, we explored if the modulation of its expression may play a role in how *T.*

alpestre differently withstands the cold compared to the related species *T. immigrans.*

We thus performed a pilot test measuring the different expression patterns of *stv*, two

paralogs of *D. melanogaster Hsc70-4* (*Hsc70-4d* and *Hsc70-4e*), a member of the

Hsp90 subfamily (*Hsp83*), and sHsp *l(2)efl4* in the two species.

We measured the transcription activity of three recovery times (0.5 h, 2.0 h,

and 4.0 h) and compared coma and non-coma individuals and found a significant

difference in the expression patterns of *stv*, *Hsc70-4d*, *Hsc70-4e*, and *Hsp83*, but not

of *l(2)efl4* (one-way ANCOVA followed by Bonferroni-Holm correction). In particular,

looking at the different recovery times, *stv* and *Hsc70-4d* responded quickly in *T.*

alpestre but not in *T. immigrans,* being significantly expressed only in workers of *T.*

alpestre after 0.5 h recovery, but not in *T. immigrans* (Figure 5d-e)*.* Comparing the

expression levels between species, we found higher mRNA levels for *Hsc70-4e* in *T.*

alpestre after 4 h recovery (Figure 5f)*.*

Discussion

Tetramorium alpestre represents a critical example of evolutionary adaptation to cold

environments such as alpine habitats, characterized by low temperatures during the

growing season (Figure 1c). By analyzing its genome in the context of 18 other ant

species, we shed new light on the evolution and adaptation of this ant, which can

serve as a potential model for arthropod adaptation to cold habitats. We found good

concordance between the nuclear and the mitochondrial dated phylogenies,

concerning the dating of the speciation of *T. alpestre*, which seems to have diverged

from its closest relative, *T. immigrans*, between 2 and 9 mya. This period overlaps with

the beginning of a new climatic zonation of the European continent during the Middle

and earliest Late Miocene (Zachos et al. 2001). At the end of this period, a rapid uplift

(∼5 mya) fundamentally changed the paleogeographic and topographic setting of

central and southern Europe and triggered Alpine glaciation (Kuhlemann 2007).

Interestingly, both paleoclimatic and paleogeographic changes co-occurred with the

origin of the lineage leading to *T. alpestre* (Figure 1b). In fact, while these glaciations

dramatically reshaped global biodiversity patterns, eliminating terrestrial biota from

many mid- to high-latitude areas, they may have been the main driver of

diversification on highlands by reducing gene flow among populations (Wallis et al.

2016).

The present results support our prediction that two parallel patterns will occur

in the genome of species adapting to extreme habitats. The genome of *T. alpestre*

appears to be under the influence of two strong evolutionary trajectories: on the one

hand, a diversifying selective force, two-fold higher compared with all other ants, and on the other hand, a relaxation of the overall purifying selection present in other ants, identified by the skewed distribution of k towards zero. Specifically, the diversifying selective force affects more than a hundred genes in many key biological processes, such as those related to development, cell migration, and gastrulation. Genes under the category of biological processes are also under selection in the genome of the Antarctic midge *Belgica antarctica* (Kelley et al. 2014). In fact, early stages are the most fragile ones, having the highest rate of lethal phenotypes during heat shock and UV irradiation, particularly around gastrulation (Uchida et al. 2018). The same may pertain to genes related to cellular-level organization (e.g., spindle midzone assembly, supramolecular fiber organization, actin cytoskeleton organization), where environmental stress may jeopardise a well working cell division machinery, as SNPs associated to long-term cold-adaptation plasticity were found to be present in genes related to cytoskeletal and membrane structural components in *D. melanogaster* (Gerken et al. 2015). Among 25 genes involved in these biological processes, *polo*, a polo-like kinase which plays a central role as regulator of cell division and is required for several events of mitosis and cytokinesis (Archambault et al. 2015), shows three amino acid substitutions in positions corresponding to active binding sites, and thus could be a good candidate for further studies.

The two evolutionary trajectories were predicted in our hypothesis, and while diversifying selection is somewhat expected in species adapting to a new environmental niche, the magnitude of relaxed selection found is more surprising and never recorded. Although both ω and k distributions in *T. alpestre* are highly skewed, the absolute number of genes under diversifying and relaxed selection are not

markedly greater than other species in similar studies (Cicconardi et al. 2017b; Roux

et al. 2014; Harpur et al. 2014). This cannot be directly ascribed to the type *I* error rate

because the same rate should be equally randomly present in all the other short or

long branches of the phylogeny. Nevertheless, *T. alpestre*, as part of a species complex

(Steiner et al. 2010; Wagner et al. 2017), may have more recently diverged from more

closely related species not yet adapted to the alpine habitat. Magnitude of this relaxed

selection is evidenced by the 70 enriched biological processes, and can be possibly

seen as the consequence of a shift and/or decreased magnitude in the purifying

selection. The direct implication of this relaxation is not clear, especially its effect on

genes and their biological processes, such as the regulation of cellular development,

signal transduction, cell communication, neurogenesis, axon development, and many

others. If the physiology of cold adaptation utilises different metabolic and

developmental strategies to minimise and optimise energy consumption, a new

balance needs to be reached. The two patterns observed in present study

(diversifying/relaxing selection) might represent this new shift. Interestingly, *T.*

immigrans, which inhabit intermediate environmental conditions (Figure 1c), shows

also an intermediate pattern of *k*, with an intriguing bimodal distribution (Figure 2b).

Therefore, we can speculate that the highly skewed distributions point to ongoing

adaptation to a colder climate, and a relaxation of selective forces present in warmer

habitats. To our knowledge, this is the first time that this effect is found in an

organism, most likely due to the underrepresentation of genomic studies on alpine

and cold adapted species, and, importantly, the lack of application of the RELAXED

test in a genome-wise manner.

Examining the effect of those evolutionary trajectories, metabolism seems to

be significantly involved in the sub-alpine ant`s adaptation. Several clues are pointing

in that direction, such as the presence of underground aphids in *T. alpestre* nests but

not in those of *T. immigrans* (unpublished data), the expansion of OGs related to sugar

transporters, and five enzymes under diversifying selection related to metabolism and

phosphorylation of sugars. We deeply explored and validated one of these enzymes,

Hex-T1, a key regulator and rate-limiting enzyme for energy (sugars) metabolism and

reactive oxygen species (ROS) activity in insects (Xian-Wu and Wei-Hua 2016), by

contrasting its enzymatic activity between *T. alpestre* and *T. immigrans.* We showed

that the mutated form in *T. alpestre* is indeed overall more efficient, with less

substrate needed to reach the highest reaction velocity of the enzyme, possibly due

to a single amino acid mutation that enhance effect on the catalytic efficiency of the

enzyme due to the formation of a more functional network of hydrogen bonds with

its substrate*.* In fact, the path of the hydrogen bonds found in TalpHex-T1 resembled

the one depicted by the X-ray solved structure of Hex-T1 from (Mulichak et al. 2002)*,*

used as a template to obtain the three-dimensional coordinates of both TimmHex-T1

and TalpHex-T1 (see Methods). Mulichak and colleagues identified their model

residues Lys621 and Asp657 as crucial for the catalytic activity of the protein; the

corresponding residues TalpHex-T1_{K168} and TalpHex-T1_{D204} are indeed interacting,

together with other residues, with glucose in TalpHex-T1 at both 6 and 26 °C. On the

other hand, while the interaction with TalpHex-T1_{D204} is conserved, the one with the

lysine is lost in TimmHex-T1 and a looser protein-sugar network is formed, explaining

the higher catalytic efficiency of TalpHex-T1 versus that of TimmHex-T1. We can

indeed infer that the higher flexibility conferred by TalpHex-T1_{G74} increases the shift

of the residue TalpHex-T1$_{K168}$, belonging to the catalytic site (Figure 4d, e), allowing it

to move closer and establish a stronger interaction with glucose, possibly contributing

to lower the Km of the catalytic reaction. In eco-evolutionary terms all these findings

hint at a more intimate relation between *T. alpestre* and the aphids. This relationship

of ants and aphids is one of the most studied mutualistic relationships in the animal

kingdom. Aphids produce honeydew, and in return ants offer protection from

predators, parasitoids, fungal infection and adverse conditions. Honeydew contains

monosaccharides (glucose and fructose), disaccharides (maltose, sucrose), and

Melezitose, a trisaccharide, that is one of the main sugar synthesized by aphids, and

which has been found to attract ants and maintain the ant-aphid mutualism (Fischer

and Shingleton 2001). Small amounts of amino-acids, proteins, and lipids are also

present (Pringle et al. 2014). Usually, this interaction is above ground level, on plant

leaves and branches, but there are cases in which ants are specialized to farm

subterranean aphids, just like in *Lasius flavus* (Ivens et al. 2012). *Tetramorium alpestre*

may have evolved the same behavior through a coevolution of physiological factors

such as palatability, fluid intake rate and digestibility of sugar molecules, improving

the ants' survival by enhancing energy intake, a limiting factor in high-elevation

ecosystems.

The complete annotation and characterization of the five HSP subfamilies in 19

ant species show good overall conservation, and strong purifying selection was

observed in most subfamilies. The exception are sHsps, which seem to be the most

diversified HSP subfamily, with the highest and the widest range of ω. It is also the

only subfamily that took a different evolutionary path compared with Diptera, having

very few orthologs in common with them. Given that sHsps are virtually ubiquitous

molecular chaperones that can prevent the irreversible aggregation of denaturing

proteins (Haslbeck and Vierling 2015), they may have played a central role in the

adaptation of ants to strongly different climates and environments. Although we

observe a higher distribution of ω in *T. alpestre*, no HSPs show a significant sign of

diversifying selection. Rather, six loci show relaxed purifying selection (Hsp90: *Trap1*;

Hsp60:*CCT4*, *CCT5*; Hsp40: *Sec63*, *DnajC8*, *DnajC11*). The impact of the TNF receptor-

associated protein (Trap1) on cellular bioenergetics of *T. alpestre* may contribute to

its adaptation to cold environments by differently modulating cellular metabolism.

Given that it functions as a negative regulator of mitochondrial respiration, able to

modulate the balance between oxidative phosphorylation and aerobic glycolysis,

possibly another form of metabolic adaptation. It has been shown that reduced or

absent *Trap1* expression leads to deregulated mitochondrial respiration, that is, a high

energy state characterized by elevated ATP and reactive oxygen species, an increase

in mitochondrial respiration and fatty acid oxidation, and a cellular accumulation of

tricarboxylic acid cycle intermediates (Yoshida et al. 2013). *Trap1* overexpression is

associated with increased expression of genes associated with cell proliferation (Liu et

al. 2010a). Since the two Hsp60s are associated to protein folding and cytoplasmic

microtubule organization, biological processes also enriched by genes under

diversifying selection, this suggests profound changes in the cellular physiology of *T.

alpestre*. It is also plausible that adapting to a cold environment will depress selection

on genes directly related to severe heat shock. As we have shown, *T. alpestre*'s habitat

is characterised by temperatures that rarely exceed 20 °C (median = 16.5 °C; 99%

confidence interval = 17.8 °C), although rocky habitats may produce warmer

microhabitats. This could mean that this species experienced relaxed selection for

HSPs associated with the denaturation to protein folding by heat shock, as suggested

by the relaxation of purifying selection on *Cct5* and the three Hsp40s associated with

responses to heat and temperature stimulus.

Although we provide a range of evidence that adaptation to cold environments

can be related to changes in protein-coding sequences, it has long been postulated

that phenotypic divergence between closely related species, such as *T. alpestre* and *T.*

immigrans, is primarily driven by quantitative and spatiotemporal changes in gene

expression, mediated by alterations in regulatory elements (e.g.: Danko et al., 2018;

Prescott et al., 2015). We tested this hypothesis by looking at the expression patterns

of *stv* and four other HSPs (*Hsc70-4d*, *Hsc70-4e*, *Hsp83*, *l(2)efl4*), between *T. alpestre*

and *T. immigrans*. *Stv* is a member of Bcl-2-associated athanogene (BAG)-family

proteins, which interact with Hsc70s and Hsp70s, and can modulate, either positively

or negatively, the functions of these chaperones (Doong et al. 2002). It is implicated

in the recovery from chill shock in *D. melanogaster* during cold recovery (Colinet and

Hoffmann 2010, 2012). The concerted synthesis of *D. melanogaster stv* and Hsp70s

suggests cooperation to offset cold injury, possibly by preserving the

folding/degradation and by regulating apoptosis (Colinet and Hoffmann 2010). In our

experiment, an acute cold stress (-6 °C) was used to knock down individuals. By looking

at the differential expression of these genes in coma and non-coma individuals within

and between species, we were not only able to show that *stv* may interact with

Hsc70s, but also that *T. alpestre* has a quicker response in these genes compared with,

T. immigrans, the related species not cold adapted. Given these results, which are in

line with the selection signature we identified, it is probable that *stv* may play a similar

role in the physiology of ants. As we have shown in this study, ants completely lack

Hsp70s (Figure S14) and retained only the heat shock cognates (Hsc70s) with mainly

two conserved paralogs (*Hsc70-4d* and *Hsc70-4e*), which may have evolved to function

as the Dipteran Hsp70s. Although we found an interesting and promising correlation

between chill shock, *stv*, and Hsc70s, it will be important to continue exploring other

expression patterns in ant physiology to assess whether *stv* has tissue-specific

expression patterns, as was found in *D. melanogaster* (Coulson et al. 2005), and to

study its expression across different species with different ecological adaptations.

One of the most enduring problems of evolutionary biology is explaining how

complex adaptive traits originate (Wagner and Lynch 2010). Although it is widely

assumed that new traits arise solely from selection on genetic variation, many

researchers ask whether phenotypic plasticity precedes and facilitates adaptation

(Levis and Pfennig 2016). This alternative scenario, the plasticity-first hypothesis,

states that under novel conditions, environmentally induced variation can be refined

by selection and, depending on whether plasticity is favoured, become

developmentally canalized through genetic assimilation. Under this scenario,

environmentally induced phenotypic change can precede and promote the

evolutionary origins of a complex adaptive trait.

Four criteria have been proposed for verifying this hypothesis (Levis and

Pfennig 2016): 1) the focal trait needs to be environmentally induced in the ancestral-

proxy lineages; 2) cryptic genetic variation will be uncovered when ancestral-proxy

lineages experience the derived environment; 3) the focal trait will exhibit evidence

of having undergone an evolutionary change in its regulation, in its form, or in both in

the derived lineages; and 4) the focal trait will exhibit evidence of having undergone

adaptive refinement in the derived lineages. Although it is methodically difficult to

prove each point, our findings on the evolution and adaptation of *T. alpestre* appear

consistent with plasticity-first evolution. In more detail, *T. immigrans*, the ancestral-

proxy lineage, shows considerable plasticity under variable temperature conditions

compared to other ant species (Criterion 1, Figure 1c), showing an ability to recover

from severe chill-shocks that is almost as good as the derived lineage (*T. alpestre*),

and, unexpectedly, better efficiency of Hex-t1 at lower rather than higher

temperatures (Figure 4a). We suggest that the derived lineage, *T. alpestre*, underwent

genetic assimilation of the cold response (Criterion 3), followed by evolutionary

refinement via changes in regulation, resulting, for example, in *stv* differential

expression (Criterion 4). Consistent with evolutionary refinement, the derived lineage

has evolved more extreme traits such as an overall improved activity of Hex-T1 with

reduced variability.

Our comparative genomic analysis shows how natural selection could trigger

complicated patterns of change in genomes, especially protein-coding sequences,

both in terms of diversifying selection and relaxation of purifying selection. Many of

these changes are in genes that may be associated with aspects of development,

either directly or through the associated complex changes in ecology and natural

history, and all of these hypotheses are now open for in-depth scrutiny. Furthermore,

this work identifies promising gene expression patterns that are possibly playing an

important role in adaptation to new ecological niches. Overall, this study represents

the first systematic attempt to provide a framework for the genomic analysis of

adaptation to extreme environments and underlines the importance of studying

organisms with different ecological niches in understanding the genetic basis of

ecological adaptation.

Methods

Species-locality and climatic data

To define the environment of *T. alpestre* and the other four *Tetramorium* species, georeferenced occurrence data were compiled and complemented with those for the native ranges of other ant species with available genomic data. Occurrence records for 17 species (Table S2) were compiled from the Global Ant Biodiversity Informatics database (GABI, Guénard et al., 2017, accessible via www.antmaps.org, Janicki et al., 2016). The species determinations in the records were evaluated by specialists in ant taxonomy, and erroneous data were excluded. Slightly incorrect coordinates (i.e. in the ocean within 8.5 km of land) were assigned to the nearest land point. Additionally, georeferenced occurrence data of two species (*T. alpestre, T. immigrans*) were compiled from the latest taxonomic revision (Wagner et al. 2017). The occurrence data of each species were spatially rarefied at a 5 km level using the R package SPTHIN v. 0.1.0 (Aiello-Lammens et al. 2015), to counter spatial autocorrelation and to provide some protection against sampling bias issues (Merow et al. 2013). The Bioclim variables bio1-bio19 from the WORLDCLIM database v. 1.4 (Hijmans et al. 2005) were extracted for each unique species locality at a 2.5 arc min resolution (4.65 x 4.65 = 21.6225 km^2 at the equator) using ARCGIS v. 10.4 (ESRI, Redland, CA). A Wilcoxon-Mann-Whitney rank sum test as implemented in the R package WILCOX.TEST (http://www.r-project.org) was adopted to test for differences between *T. alpestre* and the other species individually for each of the Bioclim variables. The Bonferroni-

Holm sequential rejection procedure (Abdi 2010) was used to control for multiple

testing.

Biological sample processing, genome size estimation, and genome sequencing

Six colonies of *T. alpestre* ants were sampled, two for flow cytometry (nest 18811:

Penser Joch, Italy, 46.83379° N, 11.44652° E, 21 June 2016; nest 18813: Kühtai,

Austria, 47.27923, 11.07590° E, 21 June 2016), three for RNA extraction (nest 18586:

Zirmbachalm, Austria, 47.21920° N, 11.07530° E, 30 July 2015; nest 18590:

Zirmbachalm, 47.21910° N, 11.07620° E, 30 July 2015; nest 18594: Penser Joch,

46.82940°N, 11.43830° E, 2 August 2015), and one for DNA extraction (nest 18592:

Penser Joch, 46.83380° N, 11.44610° E, 2 August 2015). The ants were brought alive

to the laboratory (for flow cytometry) or immediately submerged in RNAlater (for RNA

extraction) or 96% ethanol (for DNA extraction) and stored at -70 °C (RNA) or -20 °C

(DNA).

Flow cytometry was used to estimate relative genome size from single *T.*

alpestre worker heads, using strain ISO-1 of *Drosophila melanogaster* as internal

standard (obtained from Bloomington Drosophila Stock Center, Indiana University,

Bloomington, USA; nuclear DNA content: 0.35 pg, Gregory 2018). Six individuals each

were measured from *T. alpestre* nests 18811 and 18813. In more detail, single *T.*

alpestre heads and, separately, heads of 10 ISO-1 females were chopped in 500 μL of

ice-cold Otto 1 Buffer (0.1 mol/L citric acid, 0.5% Tween20 (Merck KGaA, Darmstadt,

Germany)). The suspensions were filtered using 42-μm nylon meshes and incubated

for 5 min. Then, 1 mL of staining solution (DAPI (4 μL/mL) and 2-mercaptoethanol (2

μL/mL) in Otto 2 buffer (0.4 mol/L $Na_2HPO_4 \cdot 12\ H2O$)) was added to each suspension.

For each measurement, 200 µL suspension of a single *T. alpestre* worker and 10 µL of

the ISO-1 standard were used. Fluorescence intensity of 3000 cell nuclei per sample

was measured with a Partec CyFlow space flow cytometer (Partec GmbH, Münster,

Germany). Gating and peak analysis were done automatically using the software

Partec Flo Max.

For the DNA and RNA extractions, the individuals were thawed on ice and the

gasters removed. DNA was extracted using the QIAamp DNA Micro Kit (Qiagen). The

extraction followed the protocol of the manufacturer, except that 20 µl RNase A

solution (Sigma-Aldrich) was added to the vials and incubated for 5 minutes at room

temperature. DNA concentration and quality were checked via Nanodrop (Thermo

Scientific) and gel electrophoresis. Two adult males were used to generate the

overlapping-read and the paired-end (PE) libraries, while for the mate-pair (MP)

libraries, four adult females were extracted of nest 18592. RNA was extracted using

the RNeasy Mini Kit (Qiagen) following the instructions of the manufacturer. Six

developmental stages were extracted as pools of individuals (nest 18586: worker

larvae – 30 individuals, worker pupae – 40, worker adult – 50; nest 18594: gyne pupae

– 2; nest 18590: gyne adult – 7, male adult – 7). RNA concentration and quality were

checked via Nanodrop (Thermo Scientific) and gel electrophoresis.

All *T. alpestre* DNA-seq and RNA-seq Illumina libraries were prepared by

IGATech (http://igatechnology.com/). Specifically, four libraries were used for

genome sequencing of *T. alpestre*: one overlapping-reads library (insert size ~400 bp;

paired-end (PE) sequencing with 2×250 bp read length), one regular PE library (insert

size ~400 bp, 2×100 bp read length), and two mate pair libraries (insert size ~4 kb,

2×100 bp read length) (see Table S2). For RNA-seq, six strand-specific libraries (insert

size ~500 bp, 2×200 bp read length) were constructed from three stages of workers

(larvae, pupae, adult), two stages of gynes (larvae and adult), and adult males. All

libraries were sequenced on an Illumina HiSeq 2500 platform in standard (DNA) and

RAPID (RNA) mode. For each *T. immigrans*, *T. bicarinatum*, *T. simillimum*, and *T.*

parvispinum, one TrueSeq library (insert size ~500 bp, 2×100 bp read length) was

prepared and sequenced on an Illumina HiSeq 4000 machine at OIST sequencing

center.

Genomic and mitogenomic assembly and gene prediction

Genomic sequencing reads of *T. alpestre* were filtered to remove low-quality reads

and PCR duplicates, corrected, and assembled using ALLPATHS-LG v. 52488 (Gnerre et

al. 2011). Contigs were first constructed based on overlapping reads and then

scaffolded using paired-end and mate-pair information from all DNA libraries. Given

the high coverage of the overlapping reads, based on the estimated genome size,

multiple trials were performed using different coverages to establish the best

combination of read coverage. The genomes of the other four related species were

assembled using SPADES v. 3.10.0 (Bankevich et al. 2012) with enabled read error

correction algorithm, a *k*mer range of 21, 33, 55, 77, 99, and mismatch correction.

Scaffolds belonging to contaminants were identified and removed using BLOBOLOGY v.

2013-09-12 (Kumar et al. 2013).

A bioinformatic pipeline that included combinatorial approaches of homology,

ab initio, and *de novo* methods were implemented to predict gene models.

REPEATMASKER v. 4.0.6 (Smit et al. 2013) was used to generate repeat hints. For the

homology-based exon hints, protein data sets of *D. melanogaster*, *Apis mellifera*,

Nasonia vitripennis, and 12 ant species (see Table S1) were aligned to the *T. alpestre*

genome using TBLASTN (*e*-value cut-off: 1e^{-5}) and EXONERATE v. 2.2.0 (Slater and Birney

2005). As an *ab initio* procedure, the merged RNA-seq libraries were mapped using

STAR v. 020201 (Dobin et al. 2013) (settings: ALIGNENDSTYPE Local; ALIGNINTRONMIN 30;

ALIGNINTRONMAX 450,000; OUTSJFILTERINTRONMAXVSREADN 80 100 500 1000 2000 5000

20,000; ALIGNSJOVERHANGMIN 10; ALIGNTRANSCRIPTSPERREADNMAX 100,000) to generate

the splice-site position-specific scoring matrix (PSSM) and CUFFLINKS v. v2.2.1 (Trapnell

et al. 2012) to generate UTR and intron hints. This information was then integrated

with GENEMARK-ES v. 4 (Hoff et al. 2015) to train the analysis and use the outcome

under the BRAKER1 procedure (Hoff et al. 2015) and AUGUSTUS v. 3.2.1. (Stanke et al.

2006). To avoid missing genes due to unassembled regions of the genome, previously

unmapped reads were *de novo* assembled using TRINITY v. 2.2.0 (Grabherr et al. 2011)

and annotated with FRAMA (Bens et al. 2016). The three results where then merged,

and Benchmarking Universal Single Copy Orthologs (BUSCO v. 1.2) (Simão et al. 2015)

was adopted to evaluate the annotation completeness together with integrity and

completeness evaluation using DELTABLAST (Boratyn et al. 2012). A visual inspection of

the annotation and gene integrity was also performed on more than 250 loci.

All available complete ant mitochondrial genomes (mtDNA) were downloaded

from GENBANK (Benson et al. 2014) (Table S1). Illumina raw SRA data of other ant

species were downloaded and processed following Cicconardi et al., (2017b). In brief,

reads were quality filtered and assembled using IDBA-UD v. 1.1.1 (Peng et al. 2012)

and SPADES v. 3.10.0 (Bankevich et al. 2012), and annotated using MITOS v. 2 web

server (Bernt et al. 2013).

Phylogenetic analysis

Nuclear single-copy ortholog genes (scOGs) and complete mitochondrial genomes were used to compute the ant species tree. Once scOGs were determined (see below), single-locus trees and a species tree of concatenated scOGs were estimated using Maximum Likelihood (ML) search as implemented in FASTTREE v. 2.1.8 SSE3 (Price et al. 2010). Gene trees were summarized with MP-EST (Liu et al. 2010b) using the STRAW web-server (Shaw et al. 2013).

For the mtDNA phylogeny, only protein-coding genes were used. Each gene was aligned using MACSE v. 1.01b (Ranwez et al. 2011), and all alignments were concatenated. The best partitions and models of evolution were identified with PARTITIONFINDER v. 1.1.1_Mac (Lanfear et al. 2014). MtDNA trees were searched with ML and Bayesian Inference (BI) algorithms. For ML, GARLI v. 2.01.1067 (http://code.google.com/p/garli) was used performing 20 + 5 runs from random starting trees. The runs were continued until no further improvement in log-likelihood was found, followed by 1000 bootstrap pseudoreplicates. The results were summarized using SUMTREES v. 3.3.1 (http://bit.ly/DendroPy), using *A. mellifera*, *N. vitripennis*, and *D. melanogaster* as outgroups. For BI, BEAST v. 2 (Bouckaert et al. 2014) was adopted to obtain the dated mtDNA tree and to estimate the divergence of the most recent common ancestors (MRCAs) (template: BEAST (Heled and Drummond 2010)). Fossil data were taken mainly from Moreau & Bell (2013) and PALEOBIO DB (paleobiodb.org) (see Table S4 for a detailed list of references) and used as minimum calibration points constraining nodes in the topology as listed in Table S4. Lognormal prior distribution was implemented with an offset corresponding to the minimum fossil age, log(mean) of 1.0, and log(SD) of 1.0. Ten independent runs of 8 x

10^7 generations were sampled every 100,000th generation. Each partition was

modelled with an uncorrelated relaxed clock with a Yule process, species tree prior,

and its best substitution model. Evolutionary models not implemented in BEAUTi v2

(Bouckaert et al. 2014) were manually edited in the xml file. TRACER v. 1.6

(http://beast.bio.ed.ac.uk/Tracer) was used to evaluate convergence and the

parameters' effective sampling size (ESS) and a burn-in was manually implemented

and roughly 10% from each run were removed. LOGCOMBINER (Bouckaert et al. 2014)

and TREEANNOTATOR (Bouckaert et al. 2014) were used to summarize the results in a

single consensus tree. The non-ant species were excluded from the analysis, and no

outgroup was adopted. Divergence dates for the maximum-likelihood nuclear DNA

(nuDNA) tree (only ants) were also inferred using the penalized likelihood approach

(Sanderson 2002) implemented in the R package APE v. 5.1 (Paradis et al. 2004) by

calibrating four nodes: the root, the *Tetramorium*, the *Solenopsis*+*Monomorium*, and

the Attini branches (Table S4).

Functional annotation and orthologous-group dynamic evolution

Genes from 19 ant and three outgroup species (Table S3) were clustered into ortholog

groups (OGs) with HIERANOID v. 2 (Kaduk and Sonnhammer 2017), which implements

INPARANOID v. 8 (Sonnhammer and Östlund 2015) using a guide tree topology based on

Moreau & Bell (2013) and BLASTP (*e*-value cutoff of 10^{-5}) to improve the reciprocal hit

accuracy (Edgar 2010), while homology relationships were searched with DELTABLAST

(Boratyn et al. 2012). The functional annotation of each putative protein coding

sequence was performed by identifying both the protein domain architecture and

Gene Ontology (GO) terms. For each sequence, first HMMER v. 3.1b2 (HMMSCAN) (Eddy

2011) was used to predict PFAM-A v. 31.0 domains, then Domain Annotation by a

Multi-objective Approach (DAMA) v. 2 (Bernardes et al. 2015) was applied to identify

architectures combining scores of domain matches, previously observed multi-

domain co-occurrence, and domain overlapping. Annotation of GO terms was

performed as implemented in the CATH assignments for large sequence datasets (Das

et al. 2015). Briefly, each input sequence was scanned against the library of CATH

functional families (FUNFAMS) HMMs (Sillitoe et al. 2015) using HMMER3 (Eddy 2011)

to assign FUNFAMS to regions on the query sequence (with conditional E-value < 0.005).

Then the GO annotations for a matching FUNFAM were transferred to the query

sequence with its confidence scores, calculated by considering the GO term frequency

among the annotated sequences. Finally, a non-redundant set of GO annotations was

retained, making up the GO annotations for the query protein sequence (Sillitoe et al.

2015).

To identify orthologous groups specifically expanded in the *T. alpestre*

genome, BADIRATE v. 1.35 (Librado et al. 2012) was performed twice, once for the

nuDNA and one for the mtDNA ultrametric tree topologies, reconstructing ancestral

family sizes, gain, death, and innovation (GDI), applying stochastic models and

allowing estimation of the family turnover rate (λ) by ML. Using this approach, we

observed a correlation between the turnover rate (λ) and branch lengths (Spearman

correlation: ρ = -0.52, P-value = 0), and a bimodal distribution of λ. We interpreted the

distribution closer to 0 as background noise and therefore considered only expanded

and contracted OGs with λ within the second distribution. The boundary between the

two distributions was defined by the valley values between the two, which were

computed using OPTIMIZE and APPROXFUN functions implemented in the STATS v. 3.4.0

package in R.

Heat-shock protein family analysis

From the functional annotation analyses of all Hymenoptera, all sequences bearing a

valid protein domain matching HSPs were extracted and aligned using the CLUSTALW

v.1.2.1 EBI web server (Larkin et al. 2007) (settings: MBED true; MBEDITERATION true;

ITERATIONS 5; GTITERATIONS 5; HMMITERATIONS 5). A phylogenetic tree was computed using

ML as implemented in FASTTREE v. 2.1.8 SSE3 (Price et al. 2010) and manually checked

to identify and provisionally annotate the five main subfamily proteins (Hsp90, Hsp70,

Hsp60, Hsp40, and sHsp) and their OG. Sequences falling outside clusters and with low

bootstrap values were manually checked with the DELTABLAST (Boratyn et al. 2012)

web-server and with the CONSERVED DOMAIN DATABASE (CDD) web-server (Marchler-

Bauer et al. 2017) to identify and remove bacterial contaminants and

spurious/incomplete sequences. Then, nucleotide sequences belonging to each

subfamily were revers translated into amino acids, aligned using CLUSTALW v.1.2.1, and

the obtained protein alignment was used to derive the nucleotide one. FASTTREE was

used again to compute phylogenetic trees for the five HSP subfamilies.

Selection on single-copy orthologous groups and gene families

All selected scOGs and HSP-OGs were scanned to evaluate selection signature on

coding regions of *T. alpestre*. This was done by computing the mean ω and the

relaxation of each branch of the phylogeny of *Tetramorium* spp. + *Vollenhovia emeryi*.

More specifically, adopting a pipeline similar to that in Cicconardi et al. (2017a,

2017b), the signatures of diversifying selection were searched in codon-based aligning groups of one-to-one orthologous ant genes with MACSE v. 1.01b (Ranwez et al. 2011), filtering with GBLOCKS v. 0.91b (Castresana 2000) under a "relaxed" condition (Parker et al. 2013; Cicconardi et al. 2017a, 2017b) and using the aBSREL algorithm as implemented in the HYPHY batch language (Kosakovsky Pond et al. 2005) using a batch script (BRANCHSITEREL) in HYPHY (http://github.com/veg/hyphy).

In parallel, the same scOGs and HSP-OGs were scanned using the RELAX test (Wertheim et al. 2014) to search for putative signals of relaxation. In brief, RELAX tests the hypothesis of evolutionary-rate relaxation in selected branches of a phylogenetic tree compared with reference branches. A k value is computed to evaluate whether the selective strength ω shifts towards neutrality. The rate of d_N/d_S (ω) can relax ($k <$ 1), stay stable ($k = 1$), or intensify ($k > 1$). For both scOGs and HSP-OGs, and for both the ABSREL and RELAX tests, the Bonferroni-Holm sequential rejection procedure (Abdi 2010) was used to control the false discovery rate with a very stringent cutoff value of 0.005 for the adjusted P-values applied (Benjamin et al. 2017). Hsp60s and Hsp40s under relaxed selection were analyzed using the GENEMANIA prediction server (Warde-Farley et al. 2010) to predict their functions by creating an interaction network with genes by including protein and genetic interactions, pathways, co-expression, co-localization, and protein domain similarity.

Finally, the GOSTATS package for R (Falcon and Gentleman 2007) (settings: ANNOTATION org.Dm.eg.db; CONDITIONAL TRUE; TESTDIRECTION over) was used to check for GO terms of biological processes and KEGG pathway enrichments of scOGs under selection and relaxation using the whole set of tested scOGs as background. P-values < 0.005 and < 0.05 were implemented for GO terms and KEGG pathways, respectively.

Synthesis of Hex-t1 gene and its enzyme activity assays

Following identification as under diversifying selection in *T. alpestre* (see results

section "Dual signature of evolutionary pressures in single-copy OGs in the *T. alpestre*

genome"), synthesis of genes and enzymes was attempted for phosphofructokinase,

transketolase, phosphoglucosemutase, and Hex-t1 was successful just for the Hex-t1.

In more detail, the Hex-t1 genes of *Tetramorium alpestre* (1398 bp) and *T. immigrans*

(1386 bp) were synthesized by ThermoFisher lifetechnologies (Carlsbad, CA, USA) and

Eurofins Genomics (Ebersberg, Germany), respectively. Both constructs carried a 5'

EcoRI and a 3' HindIII recognition site for subsequent cloning as well as a C-terminal

His-tag. The genes were excised from their carrier backbones by enzymatic digestion.

The fragments were purified by agarose gel excision and sub-cloned into the

pACEBac1 expression vector (MULTIBACTM, Geneva Biotech, Geneva, Switzerland).

Correct insertion was verified by Sanger sequencing, and transformation into

Drosophila melanogaster cells and protein expression followed the standard protocols

of the expression vector manufacturer. After cell harvesting, the presence of the

recombinant protein in the cell lysates was verified by a western blot targeting the

His-tag. The relative amounts of protein were quantified from the western blot using

ImageJ (https://imagej.net/Welcome). The cell lysates were centrifuged at 4000 rpm

at 4 °C for 5 min, and the supernatant was directly used as protein source.

Enzyme activity assays followed a modified protocol of Crabtree & Newsholme

(1972). Briefly, 1000 µl assay medium containing 75 mM Tris pH=7.5 (Merck Millipore,

Burlington, MA, USA), 7.5 mM $MgCl_2$ (Merck), 0.8 mM EDTA (Sigma-Aldrich, St. Louis,

MO, USA), 1.5 mM KCl (Merck), 4.0 mM mercaptoethanol (Sigma), 0.4 mM NADP+

(Abcam, Cambridge, UK), 2.5 mM ATP (Abcam), and 10 mM creatine phosphate

(Abcam) were placed in a disposable cuvette (Brand, Germany) in a SPECORD 210

PLUS UV/VIS spectrophotometer (Analytic Jena AG, Germany) with an attached 1157P

programmable thermostate (VWR, Radnor, PA, USA). Variable amounts of D-glucose

(VWR) and synthetic protein were added, and the cuvette was allowed to reach the

assay temperature. Then, the reaction was started by adding 15 U creatine

phosphokinase (Sigma) and 5 U glucose 6-phospahte dehydrogenase (Sigma).

The Hex-t1 activity was measured in triplicate as the rate of reduction of NADP$^+$

causing an extinction at 340 nm. Each measurement lasted for 60 s, and every second,

an extinction value was recorded. For the determination of the Michaelis-Menten

constant (Km-value) of both enzymes, assays were conducted along a substrate

gradient from 0.00167 to 24.30000 mM D-glucose, achieving saturated extinction

curves. Assay temperatures ranged from 6 °C to 36 °C with 10 K increments. The

software PRISM v. 8 (GraphPad Software, San Diego, CA, USA) was used to calculate the

Km-values.

Hex-t1 structure modelling and molecular dynamics simulations

The three-dimensional structures of *T. alpestre* and *T. immigrans* Hex-t1 proteins are

not available. To study the effect of the amino acid changes on the proteins'

function/dynamics, we modeled the structures of these two proteins via homology

modelling. The X-ray structure of the Hex-t1 isoform from *Schistosoma mansoni*

(Mulichak et al. 2002), solved at 2.6 Å resolution was used as a template and the

models were generated using the software MODELLER v. 9.21 (Webb and Sali 2014).

Once we obtained the models, we built the simulative systems to run molecular

dynamics simulations. The two proteins TalpHex-t1 and TimmHex-t1, in complex with

the glucose substrate, were immersed in a triclinic box filled with TIP3P water

molecules (Jorgensen et al. 1983) and rendered electroneutral by the addition of

cloride counterions. The topology of the two systems, consisting of ~ 80.000 atoms

each, was built using the AMBER14 force field (Case et al. 2014), further converted in

GROMACS v. 4.6 (Hess et al. 2008) format using ACPYPE (Sousa Da Silva and Vranken

2012). Simulations were run on a GPU cluster using GROMACS v. 4.6 with the following

protocol: 1) 25000 steps of steepest descent followed by 25000 steps of coniugate

gradient minimization; 2) 5*100 ps of equilibration runs in the NVT environment

starting at 50 K and performed by increasing the temperature of 50 K after each run

untill a final value of 250 K; 3) 5*100 ps of equilibration runs in the NPT environment

starting at 50 K and performed by increasing the temperature of 50 K after each run

untill a final value of 250 K; 4) the equilibrated systems were simulated for 500 ns at

279 or 300 K, i.e. at 6 or 26 °C, for a total of four simulations and 2 μs of sampling.

Electrostatic interactions were considered by means of the Particle Mesh Ewald

method (PME) with a cut off of 1.2 nm for the real space and Van der Waals

interactions (Darden et al. 1993). Bond lengths and angles were constrained via the

LINCS algorithm (Hess et al. 1997). The temperatures were kept constant at 279 or

300 °K by using the velocity rescale with a coupling constant of 0.1 ps, and the pressure

was kept at 1 bar using the Parrinello-Rahman barostat with a coupling constant of

1.0 ps during sampling (Parrinello and Rahman 1981). The four trajectories were

collected and comparative analyses performed with the GROMACS suite or with in

house written code.

1020

Chill coma assay and quantitative qRT-PCR assay

The cold hardiness and expression of selected genes were compared between *T. alpestre* and *T. immigrans* using chill coma assays. From four colonies additional to those used for other experiments in this study, two per species, about 500 workers per colony were collected in August 2017; *T. alpestre*: Jaufenpass (18980: 46.83791° N, 11.29768° E) and Penser Joch (18978: 46.81402° N, 11.44198° E); *T. immigrans*: Vienna (18977: 48.12448° N, 16.43522° E.; 18983: 48.31341° N, 16.42529° E). Workers were collected alive using aspirators, transported to the laboratory in Innsbruck, and kept in polypropylene boxes (18.5 cm x 11 cm) with chambers of various sizes. The walls of the boxes were Fluon-coated (GP1, De Monchy International BV, Rotterdam, Netherlands) to prevent workers from escaping. Food (sugar-honey-water and deep-frozen *Drosophila* flies) and tap water were provided *ad libitum* three times a week. Workers were kept in a climate chamber (MIR-254, Panasonic, Etten Leur, Netherlands) at a 12L:12D photoperiod at constant 18 °C. The temperature of 18 °C was the average of the monthly mean temperatures experienced at 200 m a.s.l. (*T. immigrans*, approx. 23.7 °C) and at 2000 m a.s.l. (*T. alpestre*, approx. 12.3 °C) in July 2017. Workers were acclimatized to 18 °C for at least three weeks before the chill coma assays.

After the acclimation period, from each of the four colonies, 90 workers were randomly chosen and divided into 18 equivalent pools, each were used for the chill coma and for the control assays. For the chill coma assay, worker pools were transferred into empty 5-ml glass vials, sealed, and immersed for 6.5 hours in a water:ethane-1,2-diol mix (1:1) bath set at -6 °C; in pilot experiments, this temperature had been determined as the highest temperature at which the ants had

fallen into chill coma after 6.5 hours exposition; the temperature was identical for

both species. Temperature was monitored using an electronic thermometer (TFX 430,

ebro Electronic GmbH; Ingolstadt, Germany) with an accuracy of 0.05 °C inserted into

an additional, empty vial treated in the same way as the vials with ants. After chill

coma, the vials were transferred to a climate chamber at 18 °C, and the workers were

allowed to recover for 0.5, 2.0, and 4.0 hours. For each recovery time, three replicate

pools were used. After recovery, the workers were transferred to RNAse-free reaction

tubes and killed in liquid nitrogen. Tubes were stored at -70 °C. For the control assay,

worker pools were transferred to empty glass vials and placed in the climate chamber

at 18 °C. After 6.5 hours plus respective recovery time, the workers were killed and

stored as described above.

For each recovery time and replicate, complete worker pools were used for the

molecular analyses. RNA was extracted using the NUCLEOSPIN® RNA Kit (Macherey-

Nagel, Düren, Germany) following the instructions of the manufacturer. First-strand

cDNA was synthesized using 200 U REVERTAID reverse transcriptase (all reagents by

Thermo Fisher Scientific, Waltham, USA), 40 U RIBOLOCK RNASE inhibitor, 5 µM random

hexamer primers, 100 µM dNTPs, and 2 µl RNA extract in a total volume of 40 µl. The

mixture was incubated for 5 min at 25 °C, 60 min at 42 °C, and 5 min at 70 °C on a

UnoCycler 1200 (VWR, Radnor, USA). Quantitative PCR was conducted on a Rotorgene

Q (Qiagen) PCR system. Each reaction contained 1× Rotor-Gene SYBR Green PCR

Mastermix (Qiagen), 0.2 µM of primers, specifically designed for target genes (Table

S12), and 1 µl cDNA in a total volume of 10 µl. All qRT-PCR reactions were performed

as triplicates. Cycling conditions were 95 °C for 5 min, followed by 40 cycles of 94 °C

for 15 s, 58 °C for 10 s, and 72 °C for 15 s. Fluorescence was acquired at the end of

each elongation step. PCR was followed by a melting curve analysis from 60 to 95 °C

with 0.1 °C increments held for 5 s before fluorescence acquisition. Cycle threshold

(Ct) values were subtracted from total number of cycles. On these values, linear-

regression analyses were based separately for each target gene, treatment, recovery

time, and species (in each regression analysis, two populations and three replicate

values were used, i.e., n = 6) using the housekeeping gene as independent variable

and the target gene as dependent variable. For each target gene / recovery-time

combination one-way analysis of covariance (ANCOVA) followed by Bonferroni-Holm

correction was then used to identify significant differences between treatments

within species and between species within treatment.

1079

1080

Data access

The sequence data from this study have been submitted to the National Center for

Biotechnology Information (NCBI) Sequence Read Archive (SRA) under BioProject

numbers PRJNA532334, PRJNA533534, PRJNA533535, PRJNA533536 and

PRJNA533537 (https://www.ncbi.nlm.nih.gov/bioproject/). The mtDNA genome

assemblies have been submitted to the NCBI (GenBank accessions: MK861047 -

MK861070). All accession codes of deposited and retrieved data are provided in Table

S1.

1089

1090

Acknowledgements

Philipp Andesner and Stefan Gross for valuable work in the laboratory. Research was

supported by the Austrian Science Fund (FWF, P23409, P30861). The computational

results presented here have been achieved in part by using the HPC infrastructure of

the University of Innsbruck (LEO), the Vienna Scientific Cluster (VSC), the

collaborative system of the Universities Innsbruck and Linz (MACH), and the Swiss

National Supercomputing Centre (CSCS) hosted in Lugano.

1098

1099

References

Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP. 2015. spThin:

An R package for spatial thinning of species occurrence records for use in

ecological niche models. *Ecography (Cop)* **38**: 541–545.

Archambault V, Lépine G, Kachaner D. 2015. Understanding the Polo Kinase

machine. *Oncogene* **34**: 4799–4807.

Bankevich A, Nurk S, Antipov D, Gurevich A a., Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: A New Genome

Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J Comput*

Biol **19**: 455–477.

Benjamin DJ, Berger JO, Johannesson M, Nosek BA, Wagenmakers E-J, Berk R, Bollen

KA, Brembs B, Brown L, Camerer C, et al. 2017. Redefine statistical significance.

Nat Hum Behav. http://www.nature.com/articles/s41562-017-0189-z.

Bens M, Sahm A, Groth M, Jahn N, Morhart M, Holtze S, Hildebrandt TB, Platzer M,

Szafranski K. 2016. FRAMA : from RNA-seq data to annotated mRNA assemblies.

BMC Genomics 1–12. http://dx.doi.org/10.1186/s12864-015-2349-8.

Benson DA, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2014.

GenBank. *Nucleic Acids Res* **43**: D30–D35.

http://nar.oxfordjournals.org/lookup/doi/10.1093/nar/gku1216.

Bernardes JS, Vieira FRJ, Zaverucha G, Carbone A. 2015. A multi-objective

optimization approach accurately resolves protein domain architectures.

Bioinformatics **32**: 345–353.

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J,

Middendorf M, Stadler PF. 2013. MITOS: Improved de novo metazoan

mitochondrial genome annotation. *Mol Phylogenet Evol* **69**: 313–319.

http://dx.doi.org/10.1016/j.ympev.2012.08.023.

Bertelsmeier C, Ollier S, Liebhold A, Keller L. 2017. Recent human history governs

global ant invasion dynamics. *Nat Ecol Evol* **1**.

https://www.nature.com/articles/s41559-017-0184.pdf.

Bolton B. 2018. An online catalog of the ants of the world. Available from

http://antcat.org.

Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. 2012.

Domain enhanced lookup time accelerated BLAST. *Biol Direct* **7**: 12.

http://www.ncbi.nlm.nih.gov/pubmed/22510480%5Cnhttp://www.pubmedcen

tral.nih.gov/articlerender.fcgi?artid=PMC3438057.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard M a, Rambaut A,

Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary

analysis. *PLoS Comput Biol* **10**: e1003537.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3985171&tool=pm

centrez&rendertype=abstract (Accessed May 27, 2014).

Case DAA, Babin V, Berryman JTT, Betz RMM, Cai Q, Cerutti DSS, Cheatham TE, III,

Darden TAA, Duke REE, et al. 2014. AMBER14. *Univ California, San Fr*.

Christian KA, Morton SR. 1992. Extreme thermophilia in a central Australian ant,

Melophorus bagoti. *Physiol Zool* **65**: 885–905.

Cicconardi F, Di Marino D, Olimpieri PP, Arthofer W, Schlick-Steiner BC, Steiner FM.

2017a. Chemosensory adaptations of the mountain fly Drosophila nigrosparsa

(Insecta: Diptera) through genomics' and structural biology's lenses. *Sci Rep* **7**:

43770. http://dx.doi.org/10.1038/srep43770.

Cicconardi F, Marcatili P, Arthofer W, Schlick-Steiner BC, Steiner FM. 2017b. Positive

diversifying selection is a pervasive adaptive force throughout the Drosophila

radiation. *Mol Phylogenet Evol* **112**: 230–243.

Clark MS, Worland MR. 2008. How insects survive the cold: molecular mechanisms—

a review. *J Comp Physiol B* **178**: 917–933.

http://link.springer.com/10.1007/s00360-008-0286-4.

Colinet H, Hoffmann A. 2010. Gene and protein expression of *Drosophila* Starvin

during cold stress and recovery from chill coma. *Insect Biochem Mol Biol* **40**:

425–428. http://dx.doi.org/10.1016/j.ibmb.2010.03.002.

Colinet H, Hoffmann AA. 2012. Comparing phenotypic effects and molecular

correlates of developmental, gradual and rapid cold acclimation responses in

Drosophila melanogaster. *Funct Ecol* **26**: 84–93.

Coulson M, Robert S, Saint R. 2005. Drosophila starvin encodes a tissue-specific BAG-

domain protein required for larval food uptake. *Genetics* **171**: 1799–1812.

Danko CG, Choate LA, Marks BA, Rice EJ, Wang Z, Chu T, Martins AL, Dukler N,

Coonrod SA, Tait Wojno ED, et al. 2018. Dynamic evolution of regulatory

element ensembles in primate CD4+ T cells. *Nat Ecol Evol* **2**.

http://www.nature.com/articles/s41559-017-0447-5.

Darden T, York D, Pedersen L. 1993. Particle mesh Ewald: An N·log(N) method for

Ewald sums in large systems. *J Chem Phys*.

Das S, Sillitoe I, Lee D, Lees JG, Dawson NL, Ward J, Orengo CA. 2015. CATH

FunFHMMer web server: Protein functional annotations using functional family

assignments. *Nucleic Acids Res* **43**: W148–W153.

Dobin A, Davis C a, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:

15–21.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3530905&tool=pm

centrez&rendertype=abstract.

Doong H, Vrailas A, Kohn EC. 2002. What's in the "BAG"? - A functional domain

analysis of the BAG-family proteins. *Cancer Lett* **188**: 25–32.

Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput Biol* **7**.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics **26**: 2460–2461.

Falcon S, Gentleman R. 2007. Using GOstats to test gene lists for GO term

association. *Bioinformatics* **23**: 257–258.

Fischer MK, Shingleton AW. 2001. Host plant and ants influence the honeydew

sugar. *Funct Ecol* **15**: 544–550.

Gerken AR, Eller OC, Hahn D a, Morgan TJ. 2015. Constraints, independence, and

evolution of thermal plasticity: Probing genetic architecture of long- and short-

term thermal acclimation. *Proc Natl Acad Sci U S A* **112**: 4399–404.

http://www.ncbi.nlm.nih.gov/pubmed/25805817.

Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall

G, Shea TP, Sykes S, et al. 2011. High-quality draft assemblies of mammalian

genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A* **108**:

1513–1518.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson D a, Amit I, Adiconis X, Fan L,

Raychowdhury R, Zeng Q, et al. 2011. Full-length transcriptome assembly from

RNA-Seq data without a reference genome. *Nat Biotechnol* **29**: 644–52.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3571712&tool=pm

centrez&rendertype=abstract (Accessed March 19, 2014).

Guénard B, Weiser MD, Gómez K, Narula N, Economo EP. 2017. The Global Ant

Biodiversity Informatics (GABI) database: Synthesizing data on the geographic

distribution of ant species (Hymenoptera: Formicidae). *Myrmecological News*

24: 83–89.

Harpur BA, Kent CF, Molodtsova D, Lebon JMD, Alqarni AS, Owayss AA, Zayed A.

2014. Population genomics of the honey bee reveals strong signatures of

positive selection on worker traits. *Proc Natl Acad Sci* **111**: 2614–2619.

http://www.pnas.org/cgi/doi/10.1073/pnas.1315506111.

Haslbeck M, Vierling E. 2015. A first line of stress defense: Small heat shock proteins

and their function in protein homeostasis. *J Mol Biol* **427**: 1537–1548.

http://dx.doi.org/10.1016/j.jmb.2015.02.002.

Heled J, Drummond AJ. 2010. Bayesian inference of species trees from multilocus

data. *Mol Biol Evol* **27**: 570–80.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2822290&tool=pm

centrez&rendertype=abstract (Accessed February 29, 2012).

Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. 1997. LINCS: A Linear Constraint

Solver for molecular simulations. *J Comput Chem*.

Hess B, Kutzner C, Van Der Spoel D, Lindahl E. 2008. GRGMACS 4: Algorithms for

highly efficient, load-balanced, and scalable molecular simulation. *J Chem*

Theory Comput.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution

interpolated climate surfaces for global land areas. *Int J Climatol* **25**: 1965–

1978.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2015. BRAKER1:

Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and

AUGUSTUS. *Bioinformatics* **32**: 767–769.

Hölldobler B, Wilson EO. 1990. *The Ants*.

Holt RD. 2009. Bringing the Hutchinsonian niche into the 21st century: Ecological and

evolutionary perspectives. *Proc Natl Acad Sci* **106**: 19659–19665.

http://www.pnas.org/cgi/doi/10.1073/pnas.0905137106.

Ivens ABF, Kronauer DJC, Pen I, Weissing FJ, Boomsma JJ. 2012. Ants farm

subterranean aphids mostly in single clone groups - An example of prudent

husbandry for carbohydrates and proteins? *BMC Evol Biol* **12**.

Janicki J, Narula N, Ziegler M, Guénard B, Economo EP. 2016. Visualizing and

interacting with large-volume biodiversity data using client-server web-mapping

applications: The design and implementation of antmaps.org. *Ecol Inform* **32**:

185–193.

Jayakumar PC, Shouche YS, Patole MS. 2007. Functional analysis of *Drosophila*

melanogaster hexokinase Hex-A locus : multiple Initiator-like elements enhance

DPE containing promoter activity. **16**: 3–13.

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison

of simple potential functions for simulating liquid water. *J Chem Phys*.

Kaduk M, Sonnhammer E. 2017. Improved orthology inference with Hieranoid 2.

Bioinformatics **5**: btw774.

http://bioinformatics.oxfordjournals.org/lookup/doi/10.1093/bioinformatics/bt

w774.

Keeling CI, Yuen MM, Liao NY, Roderick Docking T, Chan SK, Taylor G a, Palmquist DL,

Jackman SD, Nguyen A, Li M, et al. 2013. Draft genome of the mountain pine

beetle, Dendroctonus ponderosae Hopkins, a major forest pest. *Genome Biol*

14: R27.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4053930&tool=pm

centrez&rendertype=abstract.

Kelley JL, Peyton JT, Fiston-Lavier A-S, Teets NM, Yee M-C, Johnston JS, Bustamante

CD, Lee RE, Denlinger DL. 2014. Compact genome of the Antarctic midge is likely

an adaptation to an extreme environment. *Nat Commun* **5**: 4611.

http://www.nature.com/ncomms/2014/140812/ncomms5611/full/ncomms561

1.html.

Körner C, Paulsen J, Spehn EM. 2011. A definition of mountains and their bioclimatic

belts for global comparisons of biodiversity data. *Alp Bot* **121**: 73–78.

Kosakovsky Pond SL, Frost SDW, Muse S V. 2005. HyPhy: Hypothesis testing using

phylogenies. *Bioinformatics* **21**: 676–679.

Krapf P, Russo L, Arthofer W, Möst M, Steiner FM, Schlick-Steiner BC. 2018. An

Alpine ant's behavioural polymorphism: monogyny with and without internest

aggression in Tetramorium alpestre. *Ethol Ecol Evol* **30**: 220–234.

https://www.tandfonline.com/doi/full/10.1080/03949370.2017.1343868.

Kuhlemann J. 2007. Paleogeographic and paleotopographic evolution of the Swiss

and Eastern Alps since the Oligocene. *Glob Planet Change* **58**: 224–236.

Kumar S, Jones M, Koutsovoulos G, Clarke M, Blaxter M. 2013. Blobology: exploring

raw genome data for contaminants, symbionts and parasites using taxon-

annotated GC-coverage plots. *Front Genet* **4**: 237.

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3843372&tool=pm

centrez&rendertype=abstract.

Lamprecht A, Semenchuk PR, Steinbauer K, Winkler M, Pauli H. 2018. Climate change

leads to accelerated transformation of high-elevation vegetation in the central

Alps. *New Phytol* 447–459. http://doi.wiley.com/10.1111/nph.15290.

Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A. 2014. Selecting optimal

partitioning schemes for phylogenomic datasets. *BMC Evol Biol* **14**: 82.

http://www.biomedcentral.com/1471-2148/14/82.

Larkin M a., Blackshields G, Brown NP, Chenna R, Mcgettigan P a., McWilliam H,

Valentin F, Wallace IM, Wilm a., Lopez R, et al. 2007. Clustal W and Clustal X

version 2.0. *Bioinformatics* **23**: 2947–2948.

Levis NA, Pfennig DW. 2016. Evaluating ' Plasticity-First ' Evolution in Nature : Key

Criteria and Empirical Approaches. *Trends Ecol Evol* **31**: 563–574.

http://dx.doi.org/10.1016/j.tree.2016.03.012.

Librado P, Vieira FG, Rozas J. 2012. BadiRate: Estimating family turnover rates by

likelihood-based methods. *Bioinformatics* **28**: 279–281.

Liu C, Guénard B, Garcia FH, Yamane S, Blanchard B, Yang DR, Economo E. 2015. New

records of ant species from Yunnan, China. *Zookeys* **78**: 17–78.

Liu D, Hu J, Agorreta J, Cesario A, Zhang Y, Harris AL, Gatter K, Pezzella F. 2010a.

Tumor necrosis factor receptor-associated protein 1(TRAP1) regulates genes

involved in cell cycle and metastases. *Cancer Lett* **296**: 194–205.

Liu L, Yu L, Edwards S V. 2010b. A maximum pseudo-likelihood approach for

estimating species trees under the coalescent model. *BMC Evol Biol* **10**: 302.

http://www.biomedcentral.com/1471-

2148/10/302%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=

2976751&tool=pmcentrez&rendertype=abstract.

Macdonald HC, Cunha L, Bruford MW, Florestas E, Macdonald HC, Sir T, Evans M,

Avenue M. 2016. Development of genomic resources for four potential

environmental bioindicator species: *Isoperla grammatica*, *Amphinemura*

sulcicollis, *Oniscus asellus* and *Baetis rhodani*. *bioRxivorg*.

Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK,

Geer RC, Gonzales NR, et al. 2017. CDD/SPARCLE: Functional classification of

proteins via subfamily domain architectures. *Nucleic Acids Res* **45**: D200–D203.

Merow C, Smith MJ, Silander JA. 2013. A practical guide to MaxEnt for modeling

species' distributions: What it does, and why inputs and settings matter.

Ecography (Cop) **36**: 1058–1069.

Moreau CS, Bell CD. 2013. Testing The Museum Versus Cradle Tropical Biological

Diversity Hypothesis: Phylogeny, Diversification, And Ancestral Biogeographic

Range Evolution Of The Ants. *Evolution (N Y)* **67**: 2240–2257.

Mulichak AM, Wilson JE, Padmanabhan K, Garavito RM. 2002. The structure of

mammalian hexokinase-1. *Nat Struct Biol*.

Nguyen AD, Gotelli NJ, Cahan SH. 2016. The evolution of heat shock protein

sequences, cis-regulatory elements, and expression profiles in the eusocial

Hymenoptera. *BMC Evol Biol* **16**: 15. http://www.biomedcentral.com/1471-

2148/16/15.

Nygaard S, Hu H, Li C, Schiøtt M, Chen Z, Yang Z, Xie Q, Ma C, Deng Y, Dikow R, et al.

2016. Reciprocal genomic evolution in the ant-fungus agricultural symbiosis.

Nat Commun 1–9.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of phylogenetics and evolution

in R language. *Bioinformatics* **20**: 289–290.

Parker DJ, Wiberg RAW, Trivedi U, Tyukmaeva VI, Gharbi K, Butlin RK, Hoikkala A,

Kankare M, Ritchie MG. 2018. Inter and Intraspecific Genomic Divergence in

Drosophila montana Shows Evidence for Cold Adaptation. **10**: 2086–2101.

Parker J, Tsagkogeorga G, Cotton J a, Liu Y, Provero P, Stupka E, Rossiter SJ. 2013.

Genome-wide signatures of convergent evolution in echolocating mammals.

Nature **502**: 1–9.

http://dx.doi.org/10.1038/nature12511%5Cnpapers2://publication/doi/10.103

8/nature12511.

Parrinello M, Rahman A. 1981. Polymorphic transitions in single crystals: A new

molecular dynamics method. *J Appl Phys*.

Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a *de novo* assembler for

single-cell and metagenomic sequencing data with highly uneven depth.

Bioinformatics **28**: 1420–1428.

http://www.ncbi.nlm.nih.gov/pubmed/22495754 (Accessed July 24, 2014).

Prescott SL, Srinivasan R, Marchetto MC, Grishina I, Narvaiza I, Selleri L, Gage FH,

Swigut T, Wysocka J. 2015. Enhancer Divergence and cis-Regulatory Evolution in

the Human and Chimp Neural Crest. *Cell* **163**: 68–84.

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 - Approximately maximum-likelihood

trees for large alignments. *PLoS One* **5**: e9490.

Pringle EG, Novo A, Ableson I, Barbehenn R V, Vannette RL. 2014. Plant-derived

differences in the composition of aphid honeydew and their effects on colonies

of aphid-tending ants.

Ranwez V, Harispe S, Delsuc F, Douzery EJP. 2011. MACSE: Multiple alignment of

coding SEquences accounting for frameshifts and stop codons. *PLoS One* **6**:

e22594.

Rogora M, Frate L, Carranza ML, Freppaz M, Stanisci A, Bertani I, Bottarin R,

Brambilla A, Canullo R, Carbognani M, et al. 2018. Assessment of climate

change effects on mountain ecosystems through a cross-site analysis in the Alps

and Apennines. *Sci Total Environ* **624**: 1429–1442.

https://doi.org/10.1016/j.scitotenv.2017.12.155.

Roux J, Privman E, Moretti S, Daub JT, Robinson-Rechavi M, Keller L. 2014. Patterns

of positive selection in seven ant genomes. *Mol Biol Evol* **31**: 1661–1685.

Sanderson MJ. 2002. Estimating absolute rates of molecular evolution and

divergence times: a penalized likelihood approach. *Mol Biol Evol* **19**: 101–109.

http://www.ncbi.nlm.nih.gov/pubmed/11752195.

Seifert B. 2018. *The ants of Central and North Europe*. ed. Tauer. Lutra Verlags- und

Vertriebsgesellschaft.

Shaw TI, Ruan Z, Glenn TC, Liu L. 2013. STRAW: Species TRee Analysis Web server.

Nucleic Acids Res **41**: 238–241.

Sillitoe I, Lewis TE, Cuff A, Das S, Ashford P, Dawson NL, Furnham N, Laskowski RA,

Lee D, Lees JG, et al. 2015. CATH: Comprehensive structural and functional

annotations for genome sequences. *Nucleic Acids Res* **43**: D376–D381.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. 2015. BUSCO:

Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**: 3210–3212.

Slater GSC, Birney E. 2005. Automated generation of heuristics for biological

sequence comparison. *BMC Bioinformatics* **6**: 1–11.

Smit A, Hubley R, Green P. 2013. RepeatMasker Open-4.0. 2013-2015 .

http://www.repeatmasker.org. http://repeatmasker.org.

Soberón J. 2007. Grinnellian and Eltonian niches and geographic distributions of

species. *Ecol Lett* **10**: 1115–1123.

Sonnhammer ELL, Östlund G. 2015. InParanoid 8: Orthology analysis between 273

proteomes, mostly eukaryotic. *Nucleic Acids Res* **43**: D234–D239.

Sousa Da Silva AW, Vranken WF. 2012. ACPYPE - AnteChamber PYthon Parser

interfacE. *BMC Res Notes*.

Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in

eukaryotes with a generalized hidden Markov model that uses hints from

external sources. *BMC Bioinformatics* **7**: 62.

Steiner FM, Seifert B, Moder K, Schlick-Steiner BC. 2010. A multisource solution for a

complex problem in biodiversity research: Description of the cryptic ant species

Tetramorium alpestre sp.n. (Hymenoptera: Formicidae). *Zool Anzeiger - A J*

Comp Zool **249**: 223–254.

http://linkinghub.elsevier.com/retrieve/pii/S0044523110000537 (Accessed

April 9, 2014).

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,

Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of

RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**: 562–578.

http://www.nature.com/doifinder/10.1038/nprot.2012.016 (Accessed March 1,

2012).

Uchida Y, Uesaka M, Yamamoto T, Takeda H, Irie N. 2018. Embryonic lethality is not

sufficient to explain hourglass-like conservation of vertebrate embryos. *Evodevo*

9: 1–11. https://doi.org/10.1186/s13227-018-0095-0.

Wagner GP, Lynch VJ. 2010. Evolutionary novelties. *Curr Biol* **20**: 48–52.

https://pdf.sciencedirectassets.com/272099/1-s2.0-S0960982210X00021/1-

s2.0-S0960982209019459/main.pdf?x-amz-security-

token=AgoJb3JpZ2luX2VjECIaCXVzLWVhc3QtMSJHMEUCIDsUh2jRKJL1EeOiMrN

glWa%2BljWFLis7tEu%2FF65JtK%2FrAiEA3Ge%2FEhYUyff41n0NLuT%2F6f3OMY

i9qUIAiE.

Wagner HC, Arthofer W, Seifert B, Muster C, Steiner FM, Schlick-Steiner BC. 2017.

Light at the end of the tunnel: Integrative taxonomy delimits cryptic species in

the Tetramorium caespitum complex (Hymenoptera: Formicidae).

Myrmecological News **25**: 95–129.

Wallis GP, Waters JM, Upton P, Craw D. 2016. Transverse Alpine Speciation Driven by

Glaciation. *Trends Ecol Evol* **31**: 916–926.

http://dx.doi.org/10.1016/j.tree.2016.08.009.

Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz M,

Grouios C, Kazi F, Lopes CT, et al. 2010. The GeneMANIA prediction server:

biological network integration for gene prioritization and predicting gene

function eds. J. Weaver, K. Anija, and T. Daspit. *Nucleic Acids Res* **38**: W214–

W220. http://nar.oxfordjournals.org/content/38/suppl_2/W214.short

(Accessed March 1, 2013).

Webb B, Sali A. 2014. Protein structure modeling with MODELLER. *Methods Mol Biol*

1137: 1–15.

http://www.springerlink.com/openurl.asp?genre=book&id=doi:10.1385/15925

93682%5Cnhttp://link.springer.com/10.1007/978-1-4939-0366-

5%5Cnhttp://link.springer.com/protocol/10.1007/978-1-60327-058-

8_8%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/24573470.

Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2014. RELAX:

Detecting relaxed selection in a phylogenetic framework. *Mol Biol Evol* **32**: 1–

13. http://www.ncbi.nlm.nih.gov/pubmed/25540451.

Wilson EO. 1990. *Success and Dominance in Ecosystems: The Case of Social Insects*.

Xian-Wu L, Wei-Hua X. 2016. Hexokinase is a key regulator of energy metabolism and

ROS activity in insect lifespan extension. *Aging (Albany NY)* **8**: 245–258.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4789580/pdf/aging-08-

245.pdf.

Yoshida S, Tsutsumi S, Muhlebach G, Sourbier C, Lee M-J, Lee S, Vartholomaiou E,

Tatokoro M, Beebe K, Miyajima N, et al. 2013. Molecular chaperone TRAP1

regulates a metabolic switch between mitochondrial respiration and aerobic

glycolysis. *Proc Natl Acad Sci* **110**: E1604–E1612.

http://www.pnas.org/cgi/doi/10.1073/pnas.1220659110.

Zachos J, Pagani M, Sloan L, Thomas E, Billups K. 2001. Trends, Rythms, and

Aberration in Global Climate 65 Ma to Present. *Science (80-)* **292**: 686–693.

1430