Machine-learning classification suggests that many alphaproteobacterial prophages may instead be gene transfer agents

Roman Kogay[1], Taylor B. Neely[1,2], Daniel P. Birnbaum[1,3], Camille R. Hankel[1,4], Migun Shakya[1,5], and Olga Zhaxybayeva[1,6,*]

[1] Department of Biological Sciences, Dartmouth College, Hanover, NH, USA

[2] Present address: Amazon.com Inc., Seattle, WA, USA

[3] Present address: School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

[4] Present address: Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA

[5] Present address: Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM, USA

[6] Department of Computer Science, Dartmouth College, Hanover, NH, USA

* Author for Correspondence: Olga Zhaxybayeva, Department of Biological Sciences, Dartmouth College, Hanover, NH, USA, Tel: 603-646-8616, E-mail: olga.zhaxybayeva@dartmouth.edu

Data deposition

Sequence alignments and phylogenetic trees are available in a **FigShare** repository at DOI 10.6084/m9.figshare.8796419. The Python source code of the described classifier and additional scripts used in the analyses are available via a **GitHub** repository at https://github.com/ecg-lab/GTA-Hunter-v1

Abstract

Many of the sequenced bacterial and archaeal genomes encode regions of viral provenance. Yet, not all of these regions encode *bona fide* viruses. Gene transfer agents (GTAs) are thought to be former viruses that are now maintained in genomes of some bacteria and archaea and are hypothesized to enable exchange of DNA within bacterial populations. In Alphaproteobacteria, genes homologous to the 'head-tail' gene cluster that encodes structural components of the *Rhodobacter capsulatus* GTA (RcGTA) are found in many taxa, even if they are only distantly related to *Rhodobacter capsulatus*. Yet, in most genomes available in GenBank RcGTA-like genes have annotations of typical viral proteins, and therefore are not easily distinguished from their viral homologs without additional analyses. Here, we report a 'support vector machine' classifier that quickly and accurately distinguishes RcGTA-like genes from their viral homologs by capturing the differences in the amino acid composition of the encoded proteins. Our open-source classifier is implemented in Python and can be used to scan homologs of the RcGTA genes in newly sequenced genomes. The classifier can also be trained to identify other types of GTAs, or even to detect other elements of viral ancestry. Using the classifier trained on a manually curated set of homologous viruses and GTAs, we detected RcGTA-like 'head-tail' gene clusters in 57.5% of the 1,423 examined alphaproteobacterial genomes. We also demonstrated that more than half of the *in silico* prophage predictions are instead likely to be GTAs, suggesting that in many alphaproteobacterial genomes the RcGTA-like elements remain unrecognized.

Keywords

Virus exaptation, GTA, *Rhodobacter capsulatus*, support vector machine, binary classification, carbon depletion

Introduction

Viruses that infect bacteria (phages) are extremely abundant in biosphere (Keen 2015). Some of the phages integrate their genomes into bacterial chromosomes as part of their infection cycle and survival strategy. Such integrated regions, known as prophages, are very commonly observed in sequenced bacterial genomes. For example, Touchon et al. (2016) report that 46% of the examined bacterial genomes contain at least one prophage. Yet, not all of the prophage-like regions represent *bona fide* viral genomes (Koonin and Krupovic 2018). One such exception is a Gene Transfer Agent, or GTA for short (reviewed most recently in Lang et al. [2017] and Grull et al. [2018]). Many of genes that encode GTAs have significant sequence similarity to phage genes, but the produced tailed phage-like particles generally package pieces of the host genome unrelated to the "GTA genome" (Hynes et al. 2012; Tomasch et al. 2018). Moreover, the particles are too small to package complete GTA genome (Lang et al. 2017). Hence, GTAs are different from lysogenic viruses, as they do not use the produced phage-like particles for the purpose of their propagation.

Currently, five genetically unrelated GTAs are known to exist in Bacteria and Archaea (Lang et al. 2017). The best studied GTA is produced by the alphaproteobacterium *Rhodobacter capsulatus* and is referred hereafter as the RcGTA. Since RcGTA's discovery 45 years ago (Marrs 1974), the genes for the related, or RcGTA-like, elements have been found in many of the alphaproteobacterial genomes (Shakya et al. 2017). For a number of *Rhodobacterales* isolates that carry RcGTA-like genes, there is an experimental evidence of GTA particle production (Fu et al. 2010; Nagao et al. 2015; Tomasch et al. 2018). Seventeen of the genes of the RcGTA "genome" are found clustered in one locus and encode proteins that are involved in DNA packaging and head-tail morphogenesis (**Figure 1** and **Supplementary Table S1**). This locus is referred to as a 'head-tail cluster'. The remaining seven genes of the RcGTA genome are distributed across four loci and are involved in maturation, release and regulation of RcGTA production (Hynes et al. 2016). Since the head-tail cluster resembles a typical phage genome with genes organized in modules similar to those of a λ phage genome (Lang et al. 2017), and since many of its

69   genes have homologs in *bona fide* viruses and conserved phage gene families (Shakya et al. 2017), the

70   cluster is usually designated as a prophage by algorithms designed to detect prophage regions in a genome

71   (Shakya et al. 2017). The RcGTA's classification as a prophage raises a possibility that some of the '*in*

72   *silico*'-predicted prophages may instead represent genomic regions encoding RcGTA-like elements.

73   Presently, to distinguish RcGTA-like genes from the truly viral homologs one needs to examine

evolutionary histories of the RcGTA-like and viral homologs and to compare gene content of a putative

RcGTA-like element to the RcGTA "genome". These analyses can be laborious and often require

subjective decision making in interpretations of phylogenetic trees. An automated method that could

quickly scan thousands of genomes is needed. Notably, the RcGTA-like genes and their viral homologs

have different amino acid composition (**Figure 1** and **Supplementary Figure S1**). Due to the purifying

selection acting on the RcGTA-like genes at least in the *Rhodobacterales* order (Lang et al. 2012) and of

their overall significantly lower substitution rates when compared to viruses (Shakya et al. 2017), we

hypothesize that the distinct amino acid composition of the RcGTA-like genes is preserved across large

evolutionary distances, and therefore the RcGTA-like genes can be distinguished from their *bona fide*

viral homologs by their amino acid composition.

84   Support vector machine (SVM) is a machine learning algorithm that can quickly and accurately

separate data into two classes from the differences in specific features within each class (Cortes and

Vapnik 1995). The SVM-based classifications have been successfully used to delineate protein families

(e.g., DNA binding proteins (Bhardwaj et al. 2005), G-protein coupled receptors (Karchin et al. 2002),

and herbicide resistance proteins (Meher et al. 2019)), to distinguish plastid and eukaryotic host genes

(Kaundal et al. 2013), and to predict influenza host from DNA and amino acid oligomers found in the

sequences of the flu virus (Xu et al. 2017). During the training step, the SVM constructs a hyperplane that

best separates the two classes. During the classification step, data points that fall on one side of the

hyperplane are assigned to one class, while those on the other side are assigned to the other class. In our

case, the two classes of elements in need of separation are phages and GTAs, while their distinguishing

features are several metrics that capture the amino acid composition of the encoding genes.

In this study, we developed, implemented, and cross-validated an SVM classifier that

distinguishes RcGTA-like head-tail cluster genes from their phage homologs with high accuracy. We then

applied the classifier to 1,423 alphaproteobacterial genomes to examine prevalence of putative RcGTA-

like elements in this diverse taxonomic group and to assess how many of the RcGTA-like elements are

mistaken for prophages in the *in silico* predictions.

Materials and Methods

The Support Vector Machine (SVM) classifier and its implementation

Let's denote as u a homolog of an RcGTA-like gene g that needs to be assigned to a class y,

"GTA" ($y = -1$) or "virus" ($y = 1$). The assignment is carried out using a weighted soft-margin SVM

classifier, which is trained on a dataset of m sequences $T^g = \{T_1^g, ..., T_m^g\}$ that are homologous to u (see

"SVM training data" section below). The basis of the classification is the *n*-dimensional vector of

features $\boldsymbol{x}$ associated with sequences u and T_i^g (see **"Generation of sequence features"** section below).

Each sequence T_i^g is known to belong to a class y_i.

Using the training dataset T^g, we identify hyperplane that separates two classes as an optimal

solution to the objective function:

111
$$min\left(\frac{1}{2}||\boldsymbol{w}||^2 + \boldsymbol{C}\sum_{i=1}^{m}\xi_i\right) (eq. 1)$$

subject to:

$$\forall_i: y_i(\boldsymbol{w}\boldsymbol{x_i} + b) \geq 1 - \xi_i, where\ \xi_i \geq 0, i = 1, \dots, m \ (eq.\ 2)$$

where $\boldsymbol{w}$ and b define the hyperplane $f(\boldsymbol{x}) = \boldsymbol{w}\boldsymbol{x_i} + b$ that divides the two classes, ξ_i is the slack variable

that allows some training data points not to meet the separation requirement, and $\boldsymbol{C}$ is a regularization

parameter, which is represented as an $m \times m$ diagonal matrix. The $\boldsymbol{C}$ matrix determines how lenient the

soft-margin SVM is in allowing for genes to be misclassified: larger values "harden" the margin, while

smaller values "soften" the margin by allowing more classification errors. The product $\boldsymbol{C}\xi$ represents the

cost of misclassification. The most suitable values for the $\boldsymbol{C}$ matrix were determined empirically during

cross-validation, as described in the **"Model training, cross validation, and assessment"** section below.

To solve equation 1, we represented this minimization problem in the Lagrangian dual form $L(\alpha)$:

$$\begin{matrix} max \\ \alpha_i \end{matrix} \quad L(\alpha) = \sum_{i=1}^{m} \alpha_i - \frac{1}{2}\sum_{i=1}^{m}\sum_{i=j}^{m} \alpha_i \alpha_j y_i y_j K(\boldsymbol{x_i}\boldsymbol{x_j}) \ (eq.\ 3)$$

subject to:

$$\forall_i: \sum_{i=1}^{m} \alpha_i y_i = 0 \ and \ 0 \leq \alpha_i \leq C, \qquad i = 1, \dots, m$$

where K represents a kernel function. The minimization problem was solved using the convex

optimization (CVXOPT) quadratic programming solver (Andersen et al. 2012). The pseudocode of the

algorithm for the weighted soft-margin SVM classifier training and prediction is shown in **Figure 2**.

SVM training data

To train the classifier, sets of "true viruses" (class $y = 1$) and "true GTAs" (class $y = -1$) were

constructed separately for each RcGTA-like gene g. To identify the representatives of "true viruses",

amino acid sequences of 17 genes from the RcGTA head-tail cluster were used as queries in BLASTP (E-

value < 0.001; query and subject overlap by at least 60% of their length) and PSI-BLASTP searches (E-

value < 0.001; query and subject overlap by at least 40% of their length; maximum of six iterations) of

the viral RefSeq database release 90 (last accessed in November 2018; accession numbers of the viral

entries are provided in **Supplementary Table S2**). BLASTP and PSI-BLAST executables were from the

BLAST v. 2.6.0+ package (Altschul et al. 1997) . The obtained homologs are listed in **Supplementary**

Table S3. Due to few or no viral homologs for some of the queries, the final training sets T^g were limited

to 11 out of 17 RcGTA-like head-tail cluster genes (*g2, g3, g4, g5, g6, g8, g9, g12, g13, g14, g15*; see

Supplementary Table S1 for functional annotations of these genes).

As the representatives of the "true GTAs", we used the RcGTA-like regions that were designated

as such via phylogenetic and genome neighborhood analyses by Shakya et al. (2017). To make sure that

our "true GTAs" do not contain any other regions, we created a database of the 235 complete

alphaprotebacterial genomes that were available in the RefSeq database prior to January 2014

(**Supplementary Table S4**). To identify the representatives of "true GTAs" in this database, amino acid

sequences of 17 genes from the RcGTA head-tail cluster (Lang et al., 2017) were used as queries in

BLASTP (E-value < 0.001; query and subject overlap by at least 60% of their length) and PSI-BLAST

searches (E-value < 0.001; query and subject overlap by at least 40% of their length; maximum of six

iterations) of the database. For each genome, the retrieved homologs were designated as an RcGTA-like

head-tail cluster if at least 9 of the homologs had no more than 5,000 base pairs between any two adjacent

genes. The maximum distance cutoff was based on the observed distances between the neighboring

RcGTA head-tail cluster genes. This assignment was determined by clustering of the obtained homologs

with the DBSCAN algorithm (Ester et al. 1996) using an in-house Python script (available in a **GitHub**

repository; see **"Software Implementation"** section below). The resulting set of 88 "true GTAs" is

provided in **Supplementary Table S5** and was verified to represent a subset of RcGTA-like elements that

were identified by Shakya et al. (2017).

Since GTA functionality has been extensively studied only in *Rhodobacter capsulatus* SB1003

(Lang et al. 2017) and horizontal gene transfer likely occurred multiple times between the putative GTAs

and bacterial viruses (Hynes et al. 2016; Zhan et al. 2016), the bacterial homologs that were both too

divergent from other bacterial RcGTA-like homologs and more closely related to the viral homologs were

eliminated from the training sets to reduce possible noise in classification. To do so, for each of the 11

trainings sets T^g, all detected viral and bacterial homologs were aligned using MUSCLE v3.8.31 (Edgar

2004) and then pairwise phylogenetic distances were estimated under PROTGAMMAJTT substitution

model using RAxML version 8.2.11 (Stamatakis 2014). For each bacterial homolog in a set T^g, the

pairwise phylogenetic distances between it and all other bacterial homologs were averaged. This average

distance was defined as an outlier distance (o) if it satisfied the inequality:

166
$$o > Q_3 + 1.5 * (Q_3 - Q_1) \ (eq.\ 4)$$

where Q_1 and Q_3 are the first and third quartiles, respectively, of the distribution of the average distances

for all bacterial homologs in the training set T^g. If an outlier distance was greater than the shortest

distance from it to a viral homolog in the set T^g, the bacterial homolog was removed from the dataset.

The alignments, list of removed sequences and the associated calculations are available in the **FigShare**

repository.

Additionally, for each gene g, the sequences that had the same RefSeq ID (and therefore 100%

amino acid identity) were removed from the training data sets. The final number of sequences in each

training dataset are listed in **Table 1**.

Assignment of weights to the training set sequences

Highly similar training sequences can have an increased influence on the position of the

hyperplane, as misclassification of two or more similar sequences can be considered less optimal than

misclassification of only one sequence. This could be a problem for our classifier, since there is generally

a highly unequal representation of taxonomic groups in the RefSeq database. To correct for this

taxonomic bias, a weighting scheme was introduced into the soft-margin of the SVM classifier during

training. To do so, sequences in each training set $T^g = \{T_1, \dots, T_m\}$ were aligned in MUSCLE v3.8.31

(Edgar 2004) (The alignments are available in the **FigShare** repository). For each pair of sequences in a

training set T^g, phylogenetic distances were calculated in RAxML version 8.2.11 (Stamatakis 2014)

under the best substitution model (PROTGAMMAAUTO; the selected substitution matrices are listed in

the **Supplementary Table S6**). The farthest-neighbor hierarchical clustering method was used to group

sequences with distances below a specified threshold t. Weight d_i of each sequence in a group was

defined as a reciprocal of the number of genes in the group. These weights are used to adjust the cost of

misclassification by multiplying C_{ii} for each sequence T_i by d_i. The most suitable value of t was

determined empirically during cross-validation, as described in the **"Model training, cross validation,**

and assessment" section below.

Generation of sequence features

To use amino acid sequences in the SVM classifier, each sequence was transformed to an *n*-

dimensional vector of compositional features. Three metrics that capture different aspects of sequence

composition were implemented: frequencies of "words" of size k (*k*-mers), pseudo amino-acid

composition (PseAAC), and physicochemical properties of amino acids.

In the first feature type, amino acid sequence of a gene is broken into a set of overlapping

subsequences of size k, and frequencies of these *n* unique *k*-mers form a feature vector $\boldsymbol{x}$. Values of k

equal to 1, 2, 3, 4, 5 and 6 were evaluated for prediction accuracy (see the **"Model training, cross**

validation, and assessment" section below).

The second feature type, pseAAC, has n=(20+λ) dimensions and takes into account frequencies

of 20 amino acids, as well as correlations of hydrophobicity, hydrophilicity and side-chain mass of amino

acids that are λ positions apart in the sequence of the gene (after Chou [2001]), More precisely, PseAAC

feature set x of a sequence of length L consisting of amino acids $R_1R_2…R_L$ is defined as follows:

$$
x_i = \begin{cases} \dfrac{r_i}{\sum_{i=1}^{20} r_i + \omega \sum_{k=1}^{\lambda} s_k}, & if\ 1 \le i \le 20, \\[3mm] \dfrac{\omega s_{j-20}}{\sum_{i=1}^{20} r_i + \omega \sum_{k=1}^{\lambda} s_k}, & if\ 21 \le j \le 20 + \lambda \end{cases} \qquad (eq.5)
$$

205   where r_i is the frequency of the i-th amino acid (out of 20 possible), ω is a weight constant for the order

206   effect that was set to 0.05, and s_k $(k = 1, …, \lambda)$ are sequence order-correlation factors. These factors are

207   defined as

$$
s_k = \frac{1}{L-k} \sum_{i=1}^{L-k} J_{i,i+k} \quad (eq.6)
$$

209   where

$$
J_{i,j} = \frac{1}{3} \left[\big(H1(R_j) - H1(R_i)\big)^2 + \big(H2(R_j) - H2(R_i)\big)^2 + \big(M(R_j) - M(R_i)\big)^2 \right] \quad (eq.7)
$$

211   and $H_1(R_i)$, $H_2(R_i)$, and $M(R_i)$ denote the hydrophobicity, hydrophilicity, and side-chain mass of amino

212   acid R_i, respectively. The $H_1(R_i)$, $H_2(R_i)$, and $M(R_i)$ scores were subjected to a conversion as described

213   in the following equation:

$$\begin{cases} H_1(i) = \dfrac{H_1^0(i) - \sum_{i=1}^{20}\dfrac{H_1^0(i)}{20}}{\sqrt{\dfrac{\sum_{i=1}^{20}\left[H_1^0(i) - \sum_{i=1}^{20}\dfrac{H_1^0(i)}{20}\right]^2}{20}}} \\[2em] H_2(i) = \dfrac{H_2^0(i) - \sum_{i=1}^{20}\dfrac{H_2^0(i)}{20}}{\sqrt{\dfrac{\sum_{i=1}^{20}\left[H_2^0(i) - \sum_{i=1}^{20}\dfrac{H_2^0(i)}{20}\right]^2}{20}}} \\[2em] M(i) = \dfrac{M^0(i) - \sum_{i=1}^{20}\dfrac{M^0(i)}{20}}{\sqrt{\dfrac{\sum_{i=1}^{20}\left[M^0(i) - \sum_{i=1}^{20}\dfrac{M^0(i)}{20}\right]^2}{20}}} \end{cases} \quad (eq.\,8),$$

where $H_1^0(i)$ is the original hydrophobicity value of the i-th amino acid, $H_2^0(i)$ is hydrophilicity value,

and $M^0(i)$ is the mass of its side chain. Values of λ equal to 3 and 6 were evaluated for prediction

accuracy (see the **"Model training, cross validation, and assessment"** section below).

The third feature type relies on classification of amino acids into 19 overlapping classes of

physicochemical properties (**Supplementary Table S7**; after Kaundal et al. [2013]). For a given

sequence, each of its encoded amino acids was counted towards one of the 19 classes, and the overall

scores for each class were normalized by the length of the sequence to form $n = 19$-dimensional feature

vector $\boldsymbol{x}$.

Model training, cross validation, and assessment

For each GTA gene, parameter, and feature type, the accuracy of the classifier was evaluated

using a five-fold cross-validation scheme, in which a dataset was randomly divided into five different

sub-samples. Four parts were combined to form the training set, while the fifth part was used as the

validation set and its SVM-assigned classifications compared to the known classes. This step was

repeated five times, so that every set was tested as a known class at least once.

For each class y ("GTA" and "Virus"), the results were evaluated by their accuracy scores,

defined as the number of correctly classified homologs divided by the total number of homologs that were

tested. The cross-validation procedure was repeated ten times to reduce the partitioning bias, and the

generated results were averaged, resulting in an Average Accuracy Score (AAS) for each gene and each

class. To ensure that "GTA" and "Virus" classes had equal impact on the accuracy assessment, each class

was assigned a weight of 0.5. The final, Weighted Accuracy Score (WAS) was calculated as:

$$WAS^g = 100 * (AAS^g_{GTA} * 0.5 + AAS^g_{Virus} * 0.5) \quad (eq.\,9)$$

The most suitable "softness" of the SVM margin was determined by trying all possible

combinations of several raw diagonal values of the matrix $\boldsymbol{C}$ $(0.01, 0.1, 1, 100, 10000)$ and the threshold t

$(0, 0.01, 0.02, 0.03, 0.04, 0.05, 0.1)$. The set of parameters and features that resulted in the highest WAS

was defined as the optimal set for a gene g. If multiple parameter and feature sets resulted in the equally

highest WAS, we applied the following parameter selection criteria, in the priority order listed, until only

one parameter set was left: first, we selected parameter set(s) with k-mer size that on average performed

better than other k-mer sizes; second, we avoided parameter set(s) that included PseAAC and

physicochemical composition features; third, we selected parameter set(s) with the value of $\boldsymbol{C}$ that gives

the highest average accuracy across the remaining parameter sets; and finally, we opted for the parameter

set with the value of t that also gives the highest WAS across the remaining parameter sets. Additionally,

we evaluated classifier accuracy using the Matthews correlation coefficient (MCC) (Matthews 1975).

Selection of alphaproteobacterial genomes for testing the presence of RcGTA-like genes

From the alphaproteobacterial genomes deposited to the RefSeq database between January 2014

and January 2019, we selected 636 complete and 789 high-quality draft genomes, with the latter defined

as genome assemblies with N50 length >400 kbp. The taxonomy of each genome was assigned using the

GTDB-Tk toolkit (Parks et al. 2018). The GTDB assignment is based on the combination of Average

Nucleotide Identity (Jain et al. 2018) and phylogenetic placement on the reference tree (as implemented in

the *pplacer* program (Matsen et al. 2010)). Three of the 1,425 genomes could not be reliably placed into a

known alphaproteobacterial order, and hence were left unclassified. Two of the 1,425 genomes were

removed from further analyses due to their classification outside of the Alphaproteobacteria class,

resulting in 635 complete and 788 high-quality genomes in our dataset (**Supplementary Table S8**).

Detection of RcGTA-like genes and head-tail clusters in Alphaproteobacteria

The compiled training datasets of the RcGTA-like genes (see the **"SVM training data"** section)

were used as queries in BLASTP (E-value < 0.001; query and subject overlap by at least 60% of their

length) searches of amino acid sequences of all annotated genes from the 1,423 alphaproteobacterial

genomes. Acquired homologs of unknown affiliation (sequences *u*) were then assigned to either "GTA" or

"virus" category by running the SVM classifier with the identified optimal parameters for each gene *g*

(**Table 2**).

The proximity of the individually predicted RcGTA-like genes in each genome was evaluated by

running the DBSCAN algorithm (Ester et al. 1996) implemented in an in-house Python script (available

in a **GitHub** repository; see **"Software Implementation"** section below). The retrieved homologs were

designated as an RcGTA-like head-tail cluster only if at least 6 of the RcGTA-like genes had no more

than 8,000 base pairs between any two adjacent genes. The maximum distance cutoff was increased from

the 5,000 base pairs used for the clustering of homologs in the training datasets (see **"SVM Training**

Data" section) because the SVM classifier evaluates only 11 of the 17 RcGTA-like head-tail cluster

homologs and therefore the distances between some of the identified RcGTA-like genes can be larger.

To reduce the bias arising from the overrepresentation of particular taxa in the estimation of the

RcGTA-like cluster abundance in Alphaproteobacteria, the 1,423 genomes were grouped into Operational

Taxonomic Units (OTUs) by computing pairwise Average Nucleotide Identity (ANI) using the FastANI

v1.1 program (Jain et al. 2018) and defining boundaries between OTUs at the 95% threshold. Since not all

OTUs consist uniformly of genomes that were either all with or all without the RcGTA-like clusters, each

RcGTA-like cluster in an OTU was assigned a weight of "1/[number of genomes in an OTU]". The

abundance of the RcGTA-like clusters in different alphaproteobacterial orders was corrected by summing

up the weighted numbers of RcGTA-like clusters.

Software Implementation

The above described SVM classifier, generation of sequence features, and preparation and

weighting of training data are implemented in a Python program called "GTA-Hunter". The source code

of the program is available via GitHub at https://github.com/ecg-lab/GTA-Hunter-v1. The repository also

contains training data for the detection of the RcGTA-like heat-tail cluster genes, examples of how to run

the program, and the script for clustering of the detected RcGTA-like genes using the DBSCAN

algorithm.

Assessment of prevalence of the RcGTA-like clusters among putative prophages

Putative prophages in the 1,423 alphaproteobacterial genomes were predicted using the

PHASTER web server (Arndt et al. [2016]; accessed in January, 2019). The PHASTER program was

chosen due to its solid performance in benchmarking studies (Sousa et al. 2018) and its useful scoring

system that ranks predictions based on a prophage region completeness (Song et al. 2019). To restrict our

evaluation to likely functional prophages, only predicted prophages with the PHASTER score >90 (i.e.,

classified as "intact" prophages) were retained for further analyses. The proportion of these predicted

intact prophages classified by the GTA-Hunter as "GTA"s was calculated by comparing the overlap

between the genomic locations of the predicted intact prophages and the putative RcGTA-like regions.

Construction of the alphaproteobacterial reference phylogeny

From the set of 120 phylogenetically informative proteins (Parks et al. 2017), 83 protein families

that are present in a single copy in >95% of 1,423 alphaproteobacterial genomes were extracted using

hmmsearch (E-value < 10^{-7}) via modified AMPHORA2 scripts (Wu and Scott 2012) (**Supplementary**

Table S9). For each protein family, homologs from *Escherichia coli* str. K12 substr. DH10B and

Pseudomonas aeruginosa PAO1 genomes (also retrieved using *hmmsearch*, as described above) were

added to be used as an outgroup in the reconstructed phylogeny. The amino acid sequences of each

protein family were aligned using MUSCLE v3.8.31 (Edgar 2004). Individual alignments were

concatenated, keeping each alignment as a separate partition in further phylogenetic analyses (Chernomor

et al. 2016). The most suitable substitution model for each partition was selected using

ProteinModelSelection.pl script downloaded from https://github.com/stamatak/standard-

RAxML/tree/master/usefulScripts. Gamma distribution with 4 categories was used to account for rate

heterogeneity among sites (Yang 1994). The maximum likelihood phylogenetic tree was reconstructed

with IQ-TREE v 1.6.7 (Nguyen et al. 2014). One thousand ultrafast bootstrap replicates were used to get

support values for each branch (Hoang et al. 2017; Minh et al. 2013). The concatenated sequence

alignment in PHYLIP format and the reconstructed phylogenetic tree in Newick format are available in

the **FigShare** repository.

Examination of conditions associated with the decreased fitness of the knock-out mutants

of the RcGTA-like head-tail cluster genes

From the three genomes that are known to contain RcGTA-like clusters (*Caulobacter crescentus*

NA100, *Dinoroseobacter shibae* DFL-12, and *Phaeobacter inhibens* BS107), fitness experiments data

associated with the knock-out mutants of the RcGTA-like head-tail cluster genes were retrieved from the

Fitness Browser (Price et al. [2018]; accessed in May, 2019 via http://fit.genomics.lbl.gov/cgi-

bin/myFrontPage.cgi). Price et al. (2018) defined gene fitness as the log2 change in abundance of knock-

out mutants in that gene during the experiment. For our analyses, the significantly decreased fitness of

each mutant was defined as a deviation from the fitness of 0 with a $|t - score| \geq 4$. The conditions

associated with the significantly decreased fitness were compared across the RcGTA-like head-tail cluster

genes in all three genomes.

Results

GTA-Hunter is an effective way to distinguish RcGTA-like genes from their viral homologs

The performance of the developed SVM classifier depends on values of parameters that

determine type and composition of sequence features, specify acceptable levels of misclassification, and

account for biases in taxonomic representation of the sequences in the training sets. To find the most

effective set of parameters, for each of the 11 RcGTA-like head-tail genes with the sufficient number of

homologs available (**Figure 1**; also, see **Materials and Methods** for details) we evaluated the

performance of 1,435 different combinations of the parameters using a cross-validation technique

(**Supplementary Table S10**).

Generally, the classifiers that only use k-mers as the feature have higher median WAS values than

the classifiers that solely rely either on physicochemical properties of amino acids or on pseudo amino

acid composition (PseAAC) (**Supplementary Figure S2** and **Supplementary Table S10**), indicating that

the conservation of specific amino acids blocks is important in delineation of RcGTA-like genes from

their viral counterparts. However, the WAS values are lower for the large k-mer sizes (**Supplementary**

Figure S2), likely due to the feature vectors becoming too sparse. Consequently, parameter combinations

with values of k above 6 were not tested. The WAS values are also lower for $k=1$, likely due to the low

informativeness of the feature. The lowest observed WAS values involve usage of physicochemical

properties of proteins as a feature (**Supplementary Figure S2** and **Supplementary Table S10**),

suggesting the conservation of physicochemical properties of amino acids among proteins of similar

function in viruses and RcGTA-like regions despite their differences in the amino acid composition. The

more sophisticated re-coding of physicochemical properties of amino acids as the PseAAC feature

performs better, but for all genes its performance is worse than the best-performing *k*-mer

(**Supplementary Figure S2** and **Supplementary Table S10**).

For several genes, the highest value of WAS was obtained with multiple combinations of features

and parameter values (**Supplementary Table S10**). Based on the above-described observations of the

performance of individual features, we preferred parameter sets that did not include PseAAC and

physicochemical composition features, and selected *k*-mer size that on average performed better than

other *k*-mer sizes (see **Materials and Methods** for the full description of the parameter selection

procedure).

For individual genes, the WAS of the selected parameter set ranges from 95.6 to 100% (**Table 2**),

with 5 out of 11 genes reaching the WAS of 100%. The two genes with the highest WAS below 99% (*g6*

and *g12*) have the smallest number of viral homologs available for training (**Table 2**). Additionally,

several viral homologs in the training datasets for *g6* and *g12* genes have smaller phylogenetic distances

to "true GTA" homologs than to other "true virus" homologs (**Supplementary Table S11**). As a result,

the SVM classifier erroneously categorizes some of the RcGTA-like *g6* and *g12* genes as "viral",

resulting in the reduced classifier efficacy (**Supplementary Table S10**).

Assessment of accuracy using the Matthews correlation coefficient (MCC) generally agrees with

the results based on WAS (**Table 2** and **Supplementary Table 10**). For 10 out of 11 genes, the set of

parameters with the highest WAS also has the highest MCC. For gene *g6*, there are sets of parameters

with higher MCC than the MCC for set of parameters with the highest WAS, but the differences among

the MCC values are small (**Supplementary Table S10**). Therefore, the combinations of features and

parameters chosen using the WAS scheme (**Table 2**) were selected to classify homologs of the RcGTA

genes in the 1,423 alphaproteobacterial genomes (**Supplementary Table S8**).

GTA-Hunter predicts abundance of RcGTA-like head-tail clusters in Alphaproteobacteria

The 1,423 examined alphaproteobacterial genomes contain 7,717 homologs of the 11 RcGTA genes. The GTA-Hunter classified 6,045 of these homologs as "GTA" genes (**Supplementary Table S12**). However, many genomes are known to contain regions of decaying viruses that may be too divergent to be recognizably "viral" and there is at least one known case of horizontal gene transfer of several GTA genes into a viral genome (Zhan et al. 2016), raising a possibility that some of the predicted "GTA" genes may not be part of "true GTA" genomic regions. To minimize such false positives, we imposed an extra requirement of multiple predicted RcGTA-like genes to be in proximity on a chromosome. Specifically, we called a genomic region the putative RcGTA-like cluster only if it consisted of at least 6 genes classified as "GTA". We found that the RcGTA-like clusters defined that way are present in one (and only one) copy in 818 of the 1,423 (~57.5%) examined alphaproteobacterial genomes (**Supplementary Table S13** and **Table 3**). Uneven taxonomic representation of Alphaproteobacteria among the analyzed genomes may inflate this estimation of the abundance of the GTA-harboring genomes within the class. To correct for this potential bias, 1,423 genomes were grouped into 797 Operational Taxonomic Units (OTUs) based on the average nucleotide identity (ANI) of their genomes (**Supplementary Table S14**). Although indeed some taxonomic groups are overrepresented in the set of 1,423 genomes, in 450 of the 797 OTUs (56.4%) all OTU members contain the putative RcGTA-like clusters (**Supplementary Table S14**).

RcGTA-like clusters are widely distributed within a large sub-clade of Alphaproteobacteria

The 818 genomes with the RcGTA-like gene clusters detected in this study are not evenly distributed across the class (**Table 3**), but are found only in a clade that includes seven orders (clade 4 in **Figure 3**). Overall, 66% of the examined OTUs within the clade 4 are predicted to have an RcGTA-like cluster (**Table 3**). RcGTA-like clusters are most abundant in clade 6 (**Figure 3**), a group that consists of the orders *Rhodobacterales* and *Caulobacterales* (**Table 3**).

Although the two unclassified orders that contain RcGTA-like clusters are represented by only

two genomes (clades 2 and 3 in **Figure 3**), their position on the phylogenetic tree of Alphaproteobacteria

suggests that the RcGTA-like element may have originated earlier than was proposed by Shakya et al.

(2017) (clade 5 on **Figure 3**). Given that RcGTA-like head-tail cluster genes are readily detectable in viral

genomes, it is unlikely that the RcGTA-like clusters remained completely undetectable in the examined

genomes outside of the clade 4 due to the sequence divergence. Therefore, an RcGTA-like element was

unlikely to be present in the last common ancestor of all Alphaproteobacteria (clade 7 on **Figure 3**),

which was suggested when only a limited number of genomic data was available (Lang and Beatty 2007).

Most of the detected RcGTA-like clusters can be mistaken for prophages

Among the 818 detected RcGTA-like clusters, the functional annotations of the 11 examined

genes were similar to the prophages and none of them refer to a "gene transfer agent" (data not shown).

Since at least 11 of the 17 RcGTA head-tail cluster genes have detectable sequence similarity to viral

genes (**Supplementary Table S3**), it is likely that, if not recognized as GTAs, many of the putative

RcGTA-like clusters will be designated as "prophages" in genome-wide searches of prophage-like

regions. To evaluate this hypothesis, we predicted prophages in the set of 1,423 alphaproteobacterial

genomes, and limited our analyses to the predicted prophage regions that are more likely to be functional

integrated viruses ('intact' prophages; see **Materials and Methods** for the criteria). Indeed, of the 1,235

'intact' prophage regions predicted in the clade 4 genomes, 664 (54%) coincide with the RcGTA-like

clusters (**Figure 4**). Conversely, 664 out of 818 of the predicted RcGTA-like clusters (81%) are classified

as intact prophages. Of the 351 RcGTA-like clusters that contain *all* 11 examined genes, 323 (92%) are

classified as intact prophages.

Interestingly, within 818 genomes that contain RcGTA-like clusters, the average number of

predicted intact prophages is 1.23 per genome (**Figure 5**), which is significantly higher than 0.51

prophages per genome in genomes not predicted to contain RcGTA-like clusters (p-value $< 0.22 * 10^{-17}$;

Mann-Whitney U test). If the 664 RcGTA-like regions classified as intact prophages are removed from

the genomes that contain them, the average number of predicted 'intact' prophages per genome drops to

0.42 (**Figure 5**) and the difference becomes insignificant (p-value = 0.1492; Mann-Whitney U test). This

analysis suggests that an elevated number of the observed predicted prophage-like regions in some

alphaproteobacterial genomes may be due to the presence of unrecognized RcGTA-like elements.

Discussion

Our study demonstrates that RcGTA-like and *bona fide* viral homologs can be clearly separated

from each other using a machine learning approach. The highest accuracy of the classifier is achieved

when it primarily relies on short amino acid *k*-mers present in the examined genes. This suggests that the

distinct primary amino acid composition of the RcGTA-like and truly viral proteins is what allows the

separation of the two classes of elements (**Figure 1**). However, the cause of the amino acid preferences of

the RcGTA-like genes, and especially enrichment of the encoded proteins in alanine and glycine amino

acids (**Figure 1**), remains unknown. Given the structure of the genetic code, the skewed amino acid

composition may be the driving force behind the earlier described significantly higher %G+C of the

genomic region encoding the RcGTA-like head-tail cluster than the average %G+C in the host genome

(Shakya et al. 2017). Regardless of the cause of the skewed amino acid composition, the successful

identification of the putative RcGTA-like elements in alphaproteobacterial taxa only distantly related to

Rhodobacter capsulatus (clade 4 in **Figure 3**) suggests that the selection to maintain these elements likely

extends beyond the *Rhodobacterales* order. Nevertheless, whether these putative elements indeed encode

GTAs, as we currently understand them, remains to be experimentally validated.

The benefits associated with the GTA production that would underlie the selection to maintain

them remain unknown. In a recently published high-throughput screen for phenotypes associated with

specific genes (Price et al. 2018), knockout of the RcGTA-like genes in the three genomes that encode the

RcGTA-like elements resulted in decreased fitness of the mutants (in comparison to the wild type) under

some of the tested conditions (**Supplementary Table S15**). Interestingly, the conditions associated with

the most statistically significant decreases in fitness correspond to the growth on non-glucose sugars, such

as D-Raffinose, β-Lactose, D-Xylose and m-Inositol. Overall, carbon source utilization is the most

common condition that elicits statistically significant fitness decreases in the mutants. The RcGTA

production was also experimentally demonstrated to be stimulated by carbon depletion (Westbye et al.

2017). Further experimental work is needed to identify the link between the RcGTA-like genes expression

and carbon utilization. Conversely, absence of the RcGTA-like elements in some of the clade 4 genomes

(**Figure 3**) indicates that in some ecological settings RcGTA-like elements are either deleterious or

"useless" and thus their genes were either purged from the host genomes (if RcGTA-like element

evolution is dominated by vertical inheritance) or not acquired (if horizontal gene transfer plays a role in

the RcGTA-like element dissemination).

Previous analyses inferred that RcGTA-like elements had evolved primarily vertically, with few

horizontal gene exchanges between closely related taxa (Hynes et al. 2016; Lang and Beatty 2007;

Shakya et al. 2017). Under this hypothesis, the distribution of the RcGTA-like head-tail clusters in

alphaproteobacterial genomes suggests that RcGTA-like element originated prior to the last common

ancestor of the taxa in clade 4 (**Figure 3**). This places the origin of the RcGTA-like element to even

earlier timepoint than the one proposed in Shakya et al. (2017). However, it should be noted that our

inference is sensitive to the correctness of the inferred relationships of taxa within the

alphaproteobacterial class, which remain to be disputed due to compositional biases and unequal rates of

evolution of some alphaproteobacterial lineages (Munoz-Gomez et al. 2019). The most recent

phylogenetic inference that takes into account these heterogeneities (Munoz-Gomez et al. 2019) is

different from the reference phylogeny shown in **Figure 3**. Relevant to the evolution of RcGTA-like

elements, on the phylogeny in Munoz-Gomez et al. (2019) the order Pelagibacterales is located within the

clade 4 instead of being one of the early-branching alphaproteobacterial orders (**Figure 3**). No RcGTA-

like clusters were detected in Pelagibacterales, although in our analyses the order is represented by only

five genomes. Better sampling of genomes within this order would be needed either to show a loss of the

RcGTA-like element in this order or to re-assess the hypothesis about origin and transmission of the

RcGTA-like elements within Alphaproteobacteria.

Genes in the detected RcGTA-like head-tail clusters remain mainly unannotated as "gene transfer

agents" in GenBank records, and therefore they can be easily confused with prophages. For example,

recently described "conserved prophage" in *Sphingomonadales* (Viswanathan et al. 2017) is predicted to

be an RcGTA-like element by GTA-Hunter. Incorporation of a GTA-Hunter-like machine learning

classification into an automated genome annotation pipeline will help improve quality of the gene

annotations in GenBank records and facilitate discovery of GTA-like elements in other taxa. Moreover,

application of the presented GTA-Hunter program is not limited to the detection of the RcGTA-like

elements. With appropriate training datasets, the program can be applied to the detection of GTAs that do

not share evolutionary history with the RcGTA (Lang et al. 2017) and of other elements that are

homologous to viruses or viral sub-structures, such as type VI secretion system (Leiman et al. 2009) and

encapsulins (Giessen and Silver 2017).

Acknowledgements

481 This work was supported by the National Science Foundation [NSF-DEB 1551674 to O.Z.]; the Simons

482 Foundation Investigator in Mathematical Modeling of Living Systems [327936 to O.Z.]; Dartmouth Dean

483 of Faculty start-up funds to O.Z.; and Dartmouth James O. Freedman Presidential Scholarship to T.N.

References

Altschul SF, et al. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25: 3389-3402.

Andersen M, Dahl J, Liu Z, Vandenberghe L. 2012. Interior-point methods for large-scale cone programming. In: Sra S, Nowozin S, Wright SJ, editors. Optimization for Machine Learning: MIT Press. p. 55–83.

Arndt D, et al. 2016. PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids Res 44: W16-W21.

Bhardwaj N, Langlois RE, Zhao G, Lu H. 2005. Kernel-based machine learning protocol for predicting DNA-binding proteins. Nucleic Acids Res 33: 6486-6493.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. Syst Biol 65: 997-1008.

Chou KC. 2001. Prediction of protein cellular attributes using pseudo-amino acid composition. Proteins 43: 246-255.

Cortes C, Vapnik V. 1995. Support-vector networks. Mach Learn 20: 273-297.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32: 1792-1797.

Ester M, Kriegel H-P, Sander J, Xu X. 1996. A density-based algorithm for discovering clusters a density-based algorithm for discovering clusters in large spatial databases with noise. In Simoudis E, Han J, Fayyad U editors. *Proceedings of the Second International Conference on Knowledge Discovery and Data Mining*: 3001507: AAAI Press. p. 226-231.

Fu Y, et al. 2010. High diversity of *Rhodobacterales* in the subarctic North Atlantic Ocean and gene transfer agent protein expression in isolated strains. Aquat Microb Ecol 59: 283-293.

Giessen TW, Silver PA. 2017. Widespread distribution of encapsulin nanocompartments reveals functional diversity. Nat Microbiol 2: 17029.

Grull MP, Mulligan ME, Lang AS. 2018. Small extracellular particles with big potential for horizontal gene transfer: membrane vesicles and gene transfer agents. FEMS Microbiol Lett 365: fny192.

Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2017. UFBoot2: improving the ultrafast bootstrap approximation. Mol Biol Evol 35: 518-522.

Hynes AP, Mercer RG, Watton DE, Buckley CB, Lang AS. 2012. DNA packaging bias and differential expression of gene transfer agent genes within a population during production and release of the *Rhodobacter capsulatus* gene transfer agent, RcGTA. Mol Microbiol 85: 314-325.

Hynes AP, et al. 2016. Functional and evolutionary characterization of a gene transfer agent's multilocus "genome". Mol Biol Evol 33: 2530-2543.

Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun 9: 5114.

Karchin R, Karplus K, Haussler D. 2002. Classifying G-protein coupled receptors with support vector machines. Bioinformatics 18: 147-159.

Kaundal R, Sahu SS, Verma R, Weirick T. 2013. Identification and characterization of plastid-type proteins from sequence-attributed features using machine learning. BMC Bioinformatics 14: S7.

Keen EC. 2015. A century of phage research: bacteriophages and the shaping of modern biology. Bioessays 37: 6-9.

Koonin EV, Krupovic M. 2018. The depths of virus exaptation. Curr Opin Virol 31: 1-8.

Lang AS, Beatty JT. 2007. Importance of widespread gene transfer agent genes in alpha-proteobacteria. Trends Microbiol 15: 54-62.

Lang AS, Westbye AB, Beatty JT. 2017. The distribution, evolution, and roles of gene transfer agents in prokaryotic genetic exchange. Ann Rev Virol 4: 87-104.

Lang AS, Zhaxybayeva O, Beatty JT. 2012. Gene transfer agents: phage-like elements of genetic
exchange. Nat Rev Microbiol 10: 472-482.
Leiman PG, et al. 2009. Type VI secretion apparatus and phage tail-associated protein complexes share a
common evolutionary origin. Proc Natl Acad Sci U S A 106: 4154-4159.
Marrs B. 1974. Genetic recombination in *Rhodopseudomonas capsulata*. Proc Natl Acad Sci USA 71:
971-973.
Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-likelihood and Bayesian
phylogenetic placement of sequences onto a fixed reference tree. BMC Bioinformatics 11: 538.
Matthews BW. 1975. Comparison of the predicted and observed secondary structure of T4 phage
lysozyme. Biochimica et Biophysica Acta (BBA) - Protein Structure 405: 442–451.
Meher PK et al. 2019. HRGPred: Prediction of herbicide resistant genes with k -mer nucleotide
compositional features and support vector machine. Sci Rep 9: 1–16.
Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast approximation for phylogenetic bootstrap. Mol
Biol Evol 30: 1188-1195.
Munoz-Gomez SA, et al. 2019. An updated phylogeny of the *Alphaproteobacteria* reveals that the
parasitic *Rickettsiales* and *Holosporales* have independent origins. Elife 8: e42535.
Nagao N, et al. 2015. The gene transfer agent-like particle of the marine phototrophic bacterium
Rhodovulum sulfidophilum. Biochem Biophys Rep 4: 369-374.
Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32: 268-274.
Parks DH, et al. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. Nat Biotechnol 36: 996-1004.
Parks DH, et al. 2017. Recovery of nearly 8,000 metagenome-assembled genomes substantially expands
the tree of life. Nat Microbiol 2: 1533-1542.
Price MN, et al. 2018. Mutant phenotypes for thousands of bacterial genes of unknown function. Nature
557: 503-509.
Shakya M, Soucy SM, Zhaxybayeva O. 2017. Insights into origin and evolution of α-proteobacterial gene
transfer agents. Virus Evol 3: vex036.
Song W et al. 2019. Prophage Hunter: an integrative hunting tool for active prophages. Nucleic Acids
Res. 47: W74–W80.
Sousa AL de et al. 2018. PhageWeb – Web Interface for Rapid Identification and Characterization of
Prophages in Bacterial Genomes. Front Genet 9: 644.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30: 1312-1313.
Tomasch J, et al. 2018. Packaging of *Dinoroseobacter shibae* DNA into gene transfer agent particles is
not random. Genome Biol Evol 10: 359-369.
Touchon M, Bernheim A, Rocha EP. 2016. Genetic and life-history traits associated with the distribution
of prophages in bacteria. ISME J 10: 2744-2754.
Viswanathan V, Narjala A, Ravichandran A, Jayaprasad S, Siddaramappa S. 2017. Evolutionary
Genomics of an Ancient Prophage of the Order *Sphingomonadales*. Genome Biol Evol 9: 646-
658.
Westbye AB, O'Neill Z, Schellenberg-Beaver T, Beatty JT. 2017. The *Rhodobacter capsulatus* gene
transfer agent is induced by nutrient depletion and the RNAP omega subunit. Microbiol 163:
1355-1363.
Wu M, Scott AJ. 2012. Phylogenomic analysis of bacterial and archaeal sequences with AMPHORA2.
Bioinformatics 28: 1033-1034.
Xu B, Tan Z, Li K, Jiang T, Peng Y. 2017. Predicting the host of influenza viruses based on the word
vector. PeerJ 5: e3579.
Yang Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates
over sites: approximate methods. J Mol Evol 39: 306-314.

Zhan Y, Huang S, Voget S, Simon M, Chen F. 2016. A novel roseobacter phage possesses features of
584   podoviruses, siphoviruses, prophages and gene transfer agents. Sci Rep 6: 30372.
585
586

587 **Tables**

588 **Table 1. Number of the RcGTA homologs in the "true GTA" and "true virus" training datasets.**

Gene	"true GTAs"	"true viruses"
g2	69	1646
g3	65	769
g4	62	465
g5	67	627
g6	61	19
g8	62	96
g9	66	61
g12	63	12
g13	73	57
g14	67	124
g15	67	155

590 **Table 2. The combinations of features and parameters that showed the highest weighted accuracy**

591 **score (WAS) in cross-validation.** The listed parameter sets were used in predictions of the RcGTA-like

592 genes in 1,423 alphaproteobacterial genomes. See **Materials and Methods** for the procedure on selecting

593 one parameter set in the cases where multiple parameter sets had the identical highest WAS.

Gene	Weighted Accuracy Score, WAS (%)	Matthews Correlation Coefficient, MCC	k-mer (size)	PseAAC (value of λ)	Grouping based on physicochemical properties of amino acids	C	t
g2	100	1	2	-[1]	-	10000	0.02
g3	100	1	3	-	-	10000	0.02
g4	100	1	3	3	-	10000	0.02
g5	100	1	3	-	-	100	0.02
g6	95.9	0.88	4	-	+	0.1	0.02
g8	99.4	0.98	2	3	-	0.1	0.03
g9	100	1	2	-	-	100	0.1
g12	95.6	0.90	5	-	-	10000	0.05
g13	99.1	0.98	2	-	-	100	0
g14	99.6	0.99	6	6	-	0.01	0.03
g15	99.7	0.99	2	-	-	10000	0.02

595 [1] throughout the table, "-" denotes that the feature type was not used

Table 3. Distribution of prophages and RcGTA-like elements across different orders within class Alphaproteobacteria.

Order	Number of genomes	Number of prophages	Number of RcGTA-like clusters	Number of OTUs	Corrected abundance of RcGTA-like clusters[1]	Percentage of OTUs that have RcGTA-like clusters
Acetobacterales	62	34	0	34	0	0
Azospirillales	13	10	0	12	0	0
Caedibacterales	1	0	0	1	0	0
Caulobacterales	50	30	39	45	35	78
Elsterales	1	0	0	1	0	0
Kiloniellales	5	1	0	3	0	0
Oceanibaculales	2	1	0	2	0	0
Paracaedibacterales	1	2	0	1	0	0
Parvibaculales	5	5	2	5	2	40
Pelagibacterales	5	0	0	5	0	0
Rhizobiales	730	763	435	300	155	52
Rhodobacterales	241	318	208	174	150	86
Rhodospirillales	24	10	0	15	0	0
Rickettsiales	70	18	0	24	0	0
Sneathiellales	2	1	0	2	0	0
Sphingomonadales	207	115	132	169	110	65
Thalassobaculales	1	0	0	1	0	0
Unclassified order 1	1	0	0	1	0	0
Unclassified order 2	1	2	1	1	1	100
Unclassified order 3	1	2	1	1	1	100

[1] See "**Detection of RcGTA-like genes and head-tail clusters in Alphaproteobacteria**" subsection of the **Materials and Methods** for explanation about the correction.

Figure Legends

Figure 1. The 'head-tail' cluster of the *Rhodobacter capsulatus* GTA "genome" and the amino acid composition of viral and alphaproteobacterial homologs for some of its genes. Genes that are used in the machine learning classification are highlighted in grey. For those genes, the heatmap below a gene shows the relative abundance of each amino acid (rows) averaged across the RcGTA-like and viral homologs that were used in the classifier training (columns). The amino acids are sorted by the absolute difference in the average relative abundance between RcGTA-like and viral homologs, which was additionally averaged across 11 genes. The heatmaps of the amino acid composition in the individual homologs are shown in **Supplementary Figure S1**.

Figure 2. The pseudocode of the SVM classifier algorithm that distinguishes RcGTA-like genes from the 'true' viruses. The algorithm is implemented in the GTA-Hunter software package (see **"Software Implementation"** section in **Materials and Methods**).

Figure 3. Distribution of the detected RcGTA-like clusters across the class Alphaproteobacteria. The presence of RcGTA-like clusters is mapped to a reference phylogenetic tree that was reconstructed from a concatenated alignment of 83 marker genes (See **Materials and Methods** and **Supplementary Table S9**). The branches of the reference tree are collapsed at the taxonomic rank of "order", and the number of OTUs within the collapsed clade is shown in parentheses next to the order name. Orange and brown bars depict the proportion of OTUs with and without the predicted RcGTA-like clusters, respectively. The orders that contain at least one OTU with an RcGTA-like cluster are colored in green. Nodes 1, 2 and 3 mark the last common ancestors of the unclassified orders. Node 4 marks the lineage where, based on this study, the RcGTA-like element should have already been present. Nodes 5 and 7

mark the lineages that were previously inferred to represent last common ancestor of the RcGTA-like

element by Shakya et al. (2017) and Lang and Beatty (2007), respectively. Node 6 marks the clade where

RcGTA-like elements are the most abundant. The tree is rooted using homologs from *Escherichia coli* str.

K12 substr. DH10B and *Pseudomonas aeruginosa* PAO1 genomes. Branches with ultrafast bootstrap

values >= 95% are marked with black circles. The scale bar shows the number of substitutions per site.

The full reference tree is provided in the **FigShare** repository.

Figure 4. An overlap between prophage and GTA predictions. The "predicted RcGTA-like clusters"

set refers to the GTA-Hunter predictions, while the "predicted intact prophages" set denotes predictions

made by the PHASTER program (Arndt et al. 2016) on the subset of the genomes that are found within

clade 4 (**Figure 3**).

Figure 5. The number of predicted 'intact' prophages in alphaproteobacterial genomes. The 1,423

genomes were divided into two groups: those without GTA-Hunter-predicted RcGTA-like clusters (in

brown) and those with these RcGTA-like clusters (in dark orange). For the latter group, the number of

prophages was re-calculated after the RcGTA-like clusters that were designated as prophages were

removed (in light orange). The distribution of the number of predicted intact prophages within each

dataset is shown as a violin plot with the black point denoting the average value. The datasets with

significantly different average values are denoted by asterisks ($p < 0.001$; Mann-Whitney U test).

Supplementary Figure Legends and Table Captions

Supplementary Figure S1. The amino acid composition of viral and alphaproteobacterial homologs of the 11 RcGTA genes. These homologs were used in the training and cross-validation of the SVM classifier. Each heatmap corresponds to one of the 11 genes (see **Supplementary Table S1** for the functional annotations of the genes). Each row in a heatmap corresponds to an individual homolog of the RcGTA gene. The homologs from viruses and alphaproteobacterial are separated by the black line and labeled as "True Virus" and "True GTA", respectively. The heatmap shows the relative abundance of each amino acid within a homolog. For each gene, the amino acids on the X axis are sorted by the absolute difference in the average relative abundance between "true viruses" and "true GTAs" (from highest on the left to the lowest on the right).

Supplementary Figure S2. The weighted accuracy scores for different types of features. The boxplots for the three feature types are color coded. The data for six examined k-mer sizes $(1, 2, 3, 4, 5, 6)$ are shown from the left to the right on the graphs. Each boxplot shows a median value bounded by the first and third quartiles, and the whiskers depict a deviation that was calculated using the 1.5*InterQuartile Range rule. Outliers are shown as dots.

Supplementary Table S1. Functional annotations of the 'head-tail' cluster genes of the *Rhodobacter capsulatus* gene transfer agent.

Supplementary Table S2. List of the 7,995 viral assemblies used to find RcGTA homologs for the training datasets.

Supplementary Table S3. List of 1,939 viruses with at least one detected RcGTA homolog. The data

in the columns show the accession numbers of these homologs.

Supplementary Table S4. List of 235 alphaproteobacterial genomes used to find RcGTA-like

clusters for the training datasets.

Supplementary Table S5. List of 88 alphaproteobacterial RcGTA-like clusters detected in 85

genomes. The data in the columns show the accession numbers of these homologs.

Supplementary Table S6. Substitution matrices that were used to generate pairwise phylogenetic

distances within training datasets.

Supplementary Table S7. Grouping of amino acids into classes based on their physicochemical

properties (after Kaundal et al., 2013).

Supplementary Table S8. List of 1,423 alphaproteobacterial genomes used for testing the presence

of RcGTA-like genes and regions.

Supplementary Table S9. Information about 83 marker genes that were used to reconstruct

reference phylogeny of Alphaproteobacteria.

Supplementary Table S10. Summary of the classifier cross-validation. Results for each gene are

shown in separate tabs. Each row represents one of the 1,435 tested combinations of the parameters

(columns A-E), number and fraction of correctly classified homologs averaged across 10 replicates

(columns F-I), and the overall weighted accuracy score (column J) and the Matthews Correlation

Coefficient (Column K) of the parameter combination. When a feature was not used, the value of the

parameter shown in columns A-C is set to 0.

Supplementary Table S11. Phylogenetic distances of the "truly viral" homologs of the genes *g6* and

***g12* to "true GTAs" and to other "true viruses" in the training datasets.** Data for the *g6* and *g12*

homologs are shown in separate tabs. Viral homologs that are more closely related to "true GTAs" than to

other "true viruses" are highlighted in yellow.

Supplementary Table S12. Summary of the alphaproteobacterial RcGTA homologs' classification.

Supplementary Table S13. Information about the 818 detected RcGTA-like clusters. Data in

columns D-N correspond to the RefSeq accession numbers of the encoded proteins.

Supplementary Table S14. Presence of the RcGTA-like clusters in the reconstructed

alphaproteobacterial Operational Taxonomic Units (OTUs).

Supplementary Table S15. Results of the fitness experiments with the knock-out mutants of the

RcGTA-like head-tail cluster genes in three alphaproteobacterial genomes. The data was retrieved

from the Fitness Browser (Price et al 2018). Each row corresponds to a separate experiment, in which the

specified gene was knocked out (column B) and decreased fitness (columns E and F) was associated with

a specific condition (column D). The conditions are classified into groups (column C). Rows

corresponding to the "carbon source" group are highlighted in yellow. This group is the most common

among the listed experiments and is found in experiments associated with each of the three genomes. For

description of conditions, refer to the Fitness Browser (Price et al 2018).

Figure 1. The 'head-tail' cluster of the Rhodobacter capsulatus GTA "genome" and the amino acid composition of viral and alphaproteobacterial homologs for some of its genes. Genes that are used in the machine learning classification are highlighted in grey. For those genes, the heatmap below a gene shows the relative abundance of each amino acid (rows) averaged across the RcGTA-like and viral homologs that were used in the classifier training (columns). The amino acids are sorted by the absolute difference in the average relative abundance between RcGTA-like and viral homologs, which was additionally averaged across 11 genes. The heatmaps of the amino acid composition in the individual homologs are shown in **Supplementary Figure S1**.

1: Let $T = (T_1, \ldots, T_m)$ be an array of training sequences T_i, $1 \leq i \leq m$

2: Let $X = (x_i)$ be the feature sets for genes $T_i \in T$

3: Let $Y = (y_i)$ be the classes for genes $T_i \in T$

4: Let $W = (d_i)$ be the weights for genes $T_i \in T$

5: Let $y_i = -1$ if T_i is a GTA and $y_i = 1$ if it is a virus

6: Let $QUADPROG$ be a quadratic programming solver

7: **procedure** $SVMTrain(T, C)$

8:      $Compute\ Lagrange - multipliers = QUADPROG(X, Y, C * W)$

9:      Let $alphas = \{\alpha_i \in Lagrange - multipliers : \alpha_i > 10^{-5}\}$

10:      Let $support\ vectors = \{T_i \in Lagrange - multipliers : \alpha_i > 10^{-5}\}$

11:      **return** $alphas$, $support\ vectors$

12: **end procedure**

13:

14: Let u be an unclassified gene, where x_u is the feature set of u

15: **procedure** $SVMPredict(alphas, supportvectors, x_u)$

16:      Let $score = 0$

17:      for $\alpha_i \in alphas$ and $T_i \in support\ vectors$ **do**

18:          $score = score + (\alpha_i * y_i * K(x_i * x_u))$

19:      **end for**

20:      if $score < 0$ then

21:          **return "**GTA"

22:      else

23:          **return** "virus"

24:      **end if**

25: **end procedure**

Figure 2. The pseudocode of the SVM classifier algorithm that distinguishes RcGTA-like genes from the 'true' viruses. The algorithm is implemented in the GTA-Hunter software package.

Figure 3. Distribution of the detected RcGTA-like clusters across the class Alphaproteobacteria. The presence of RcGTA-like clusters is mapped to a reference phylogenetic tree that was reconstructed from a concatenated alignment of 83 marker genes (See Materials and Methods and Supplementary Table S9). The branches of the reference tree are collapsed at the taxonomic rank of "order", and the number of OTUs within the collapsed clade is shown in parentheses next to the order name. Orange and brown bars depict the proportion of OTUs with and without the predicted RcGTA-like clusters, respectively. The orders that contain at least one OTU with an RcGTA-like cluster are colored in green. Nodes 1, 2 and 3 mark the last common ancestors of the unclassified orders. Node 4 marks the lineage where, based on this study, the RcGTA-like element should have already been present. Nodes 5 and 7 mark the lineages that were previously inferred to represent last common ancestor of the RcGTA-like element by (Shakya et al. 2017) and (Lang and Beatty 2007), respectively. Node 6 marks the clade where RcGTA-like elements are the most abundant. The tree is rooted using homologs from Escherichia coli str. K12 substr. DH10B and Pseudomonas aeruginosa PAO1 genomes. Branches with ultrafast bootstrap values >= 95% are marked with black circles. The scale bar shows the number of substitutions per site. The full reference tree is provided in the FigShare repository (see Materials and Methods).

Figure 4. An overlap between prophage and GTA predictions. The "predicted RcGTA-like clusters" set refers to the GTA-Hunter predictions, while the "predicted intact prophages" set denotes predictions made by the PHASTER program (Arndt et al. 2016) on the subset of the genomes that are found within Clade 4 (Figure 3).

Figure 5. The number of predicted 'intact' prophages in alphaproteobacterial genomes.
The 1,423 genomes were divided into two groups: those without GTA-Hunter-predicted RcGTA-like clusters (in brown) and those with these RcGTA-like clusters (in dark orange). For the latter group, the number of prophages was re-calculated after the RcGTA-like clusters that were designated as prophages were removed (in light orange). The distribution of the number of predicted intact prophages within each dataset is shown as a violin plot with the black point denoting the average value. The datasets with significantly different average values are denoted by asterisks (p < 0.001; Mann-Whitney U test).

Supplementary Figure S1. The amino acid composition of viral and alphaproteobacterial homologs of the 11 RcGTA genes. These homologs were used in the training and cross-validation of the SVM classifier. Each heatmap corresponds to one of the 11 genes (see Supplementary Table S1 for the functional annotations of the genes). Each row in a heatmap corresponds to an individual homolog of the RcGTA gene. The homologs from viruses and alphaproteobacterial are separated by the black line and labeled as "True Virus" and "True GTA", respectively. The heatmap shows the relative abundance of each amino acid within a homolog. For each gene, the amino acids on the X axis are sorted by the absolute difference in the average relative abundance between "true viruses" and "true GTAs" (from highest on the left to the lowest on the right).

Supplementary Figure S2. The weighted accuracy scores for different types of features. The boxplots for the three feature types are color coded. The data for six examined k-mer sizes (1, 2, 3, 4, 5, 6) are shown from the left to the right on the graphs. Each boxplot shows a median value bounded by the first and third quartiles, and the whiskers depict a deviation that was calculated using the 1.5*InterQuartile Range rule. Outliers are shown as dots.