Single Cell Analysis Reveals Partial Reactivation of X-chromosome Instead of chromosome wide dampening in Naïve Human Pluripotent Stem Cells

Mandal S, Chandel D, Kaur H, Majumdar S, Arava M, and Gayen S*

Department of Molecular Reproduction, Development and Genetics, Indian Institute of Science, Bangalore-560012, India.

 *Correspondence: srimonta@iisc.ac.in

Abstract

To balance the dosage of X-linked gene expression between the sexes, eutherian female mammals inactivate one of the two X-chromosomes present in a diploid cell. Once enacted in individual cells of the early female embryo, X-inactivation is stably transmitted such that all descendant cells maintain silencing of that X-chromosome. Recently, it has been demonstrated that in human however, X-chromosome dosage compensation initiates through the dampening of X-linked gene expression from both X-chromosomes in the preimplantation embryos and which later on switches to the X-chromosome inactivation. Subsequently, Sahakyan *et al.* found that conversion of primed human pluripotent stem cells (hPSCs) to naïve state is associated with X-chromosome dampening. However, both in embryos and hPSCs, the existence of dampened X-chromosomes remains controversial. Specially, in preimplantation embryos it has been shown that there is actually inactivation of X-chromosome instead of dampening. Here, we have performed allelic analysis of X-linked genes at single cell level in hPSCs and found that there is partial reactivation of the inactive X-chromosome instead of chromosome wide dampening upon conversion from primed to naïve state. In addition, our analysis suggests that the reduced X-linked gene expression in naïve hPSCs might be due to the erasure of X-chromosome upregulation.

Keywords: X-chromosome dampening; X-chromosome inactivation; X-chromosome upregulation, Human pluripotent stem cells; Naive & primed pluripotency; Human embryos; *Xist*

Introduction

In therian mammals, presence of two X-chromosomes in female and one X and one Y chromosome in male leads to the dosage imbalance of X-chromosome between the sexes. Therefore, to balance the X-chromosome dosage between male and female, one X-chromosome becomes inactive in female cells (Lyon, 1961). On the other hand, dosage imbalance between single active X-chromosome and two copies of autosomes (AA) is compensated through the upregulation of the active-X chromosome in both male and female (Deng et al., 2011; Larsson et al., 2019; Ohno, 1967). Recently, a unique form of X-chromosome dosage compensation has been inferred in human preimplantation embryos, which happens through the dampening of X-linked gene expression from both X-chromosomes (Petropoulos et al., 2016). However, X-chromosome dampening phenomenon in human embryos remains controversial (De Mello et al., 2017; Saiba et al., 2018). De Mello *et al.* found inactivation of X-chromosome instead of dampening when they reanalyzed the same scRNA-seq dataset of Petropoulos *et al.* with more stringency. Subsequently, Sahakyan *et al.* have found that naïve human pluripotent stem cells (hPSCs) also reflect the X-chromosome dampening of embryos (Sahakyan et al., 2017a). However, similar to the human embryos, X-chromosome states in hPSCs remains elusive (Kaur et al., 2019; De Mello et al., 2017). Therefore, here we have taken single cell approach with allelic resolution to get more insight about X-chromosome states in hPSCs using the single-cell RNA-sequencing (scRNA-seq) dataset from Sahakyan *et al.*

Results

Increased *Xist* expression and reduction in X-linked gene expression upon conversion of early to late naïve state

In rodents and human, two transient pluripotent state have been described: naïve and primed (De Los Angeles et al., 2015; Zhang et al., 2016). Pluripotency first arises in inner cell mass of preimplantation blastocysts in a naïve state and then transformed to a primed state in post implantation epiblast. Conventional hPSCs derived from blastocysts represent a primed state instead of naïve state and therefore, unable to recapitulate the preimplantation X-chromosome states (Nichols and Smith, 2009; Sahakyan et al., 2017b). Therefore, to model the preimplantation X-chromosome state, Sahakyan *et al.* converted primed hPSCs to a naïve state

(Fig. 1A). The primed cell line UCLA1, they used for their study, harbored one active-X
chromosome and one inactive X chromosome (X_aX_i). The transition of primed to naïve state
happened through an intermediate early naïve state (Fig. 1A). It was suggested that there was
reactivation of the inactive-X chromosome upon transition from primed to the early naïve state
and followed by X-chromosome dampening in late naïve cells (Fig. 1A). Moreover, X-
reactivation was associated with the transformation of cells from *Xist* positive to negative state,
whereas X-chromosome dampening was associated with the transformation of cells from *Xist*
negative to positive state. In this study, we have analyzed scRNA-seq dataset of early naïve
cells and late naïve cells (Fig. 1A). First, to test if the transition of early to late naïve state is
associated with the conversion of cells from *Xist* negative to positive state, we analyzed the
expression of *Xist* in early and late naïve cells. We found that indeed majority of the early naïve
cells had very low levels of *Xist* expression, whereas late naïve cells mostly showed
significantly higher levels of *Xist* expression as it was inferred by the Sahakyan *et al* (Fig. S1;
Supplementary file1). Next, we proceeded for analysis of X-linked gene expression. We sorted
out top 60 out of 96 early naïve cells based on low levels of *Xist* expression and top 59 out of
96 late naïve cells having highest levels of *Xist* expression. Finally, from this list we selected
out top 28 early naïve cells and 30 late naïve cells based on total gene expression (RPKM sum
and mean) (Fig. 1B; Supplementary file1) for comparison of X-linked gene expression
analysis. Comparison of *Xist* expression in these cells, showed significant increase in *Xist*
expression in late naïve cells (Fig. 1B). From the same cells (28+30), we selected 57 X-linked
genes having mean RPKM ($5 < \mu_{RPKM} < 200$) for our analysis. We found that there was a
significant reduction in X-linked gene expression in late naïve cells compared to the early naïve
cells (Fig. 1C; Supplementary file1). Taken together, our analysis of scRNA-Seq data showed
increased *Xist* expression and reduction in X-linked gene expression during transition of early
to late naïve state just like the RNA-Seq analysis of whole cell population as it was described
in Sahakyan *et al.*

Reduction in X-linked gene expression in late naïve cells is independent of *Xist*

X-chromosome dampening in preimplantation embryos is associated with the expression of
Xist from both X-chromosomes. In fact, X-dampening initiates concomitantly with the
initiation of *Xist* expression and therefore it is thought that *Xist* might have important role in
dampening process. Therefore, we wanted to examine what fraction of cells show *Xist*
expression from both X-chromosomes in late naïve cells. In fact, Sahakyan *et al.* tested allelic
expression of *Xist* at single cell level using RNA-Seq data and found ~28% of cells having *Xist*

expression from both X-chromosomes. Consistent with the Sahakyan's analysis we also found ~32 % cells (8 of 25) having the *Xist* expression from both X-chromosomes (Fig. 1D; Supplementary file 2). Next, we tested if the reduction in X-linked gene expression in late naïve cells was restricted to the *Xist* biallelic cell population only as in embryos X-dampening is mostly associated with the biallelic *Xist* expression. To test this, we compared the global X-linked gene expression of *Xist* biallelic cells with the *Xist* monoallelic cells. However, we did not find any significant difference in X-linked gene expression between *Xist* biallelic and monoallelic cells (Fig. 1E; Supplementary file 2). Interestingly, we observed that there is no significant difference also when we compared with the *Xist* negative cells even (Fig. 1E; Supplementary file 2). Altogether, we concluded that there is no significant correlation between the reduction in X-linked gene expression and *Xist* expression in late naïve cells.

No evidence of X-chromosome inactivation or dampening

We then looked further about the mechanisms behind the reduction in X-linked gene expression in late naïve hPSCs. At first, we considered it could be due to the either X-chromosome inactivation or dampening. To examine if it is due to the X-inactivation, we analyzed allelic expression pattern of multiple X-linked genes distributed across the X-chromosome based on single nucleotide polymorphisms (SNPs) in both early and late naïve cells (n=12 SNPs covering 10 genes). Interestingly, we found allelic gene expression pattern between the cells varied dramatically both in early and late naïve cells (Fig. 2A; Fig. S2; Supplementary file 2). Importantly, there was incomplete reactivation of X-linked genes in early naïve state as there many cells showed monoallelic expression along with biallelically expressed X-linked genes (Fig. 2A; Fig. S2). Almost similar pattern of allelic expression of X-linked genes was observed in late naïve cells. In addition, we found that there is no difference in allelic expression pattern between *Xist* monoallelic and biallelic cells (Fig. 2A). To validate the accuracy of our allele specific expression analysis, we also analyzed allelic expression of genes of an autosome (Chr17). As expected, we found mostly biallelic expression of SNPs, thus validating our allelic expression analysis (Fig. 2A). Next, we calculated percent of SNPs showing biallelic and monoallelic expression in each cells of early and late naïve cells (Fig. 2B). We found that there was huge variation in percent of biallelic vs monoallelic expression between cell to cell and there was overall little increase in fraction of SNPs showing biallelic expression in late naïve cells compared to the early naïve cells. Overall, we found that both early and late naïve cells have partially reactivated X-chromosome. Moreover, frequency of reactivation is increased in late naïve cells compared to the early naïve cells, however reactivation remained incomplete.

We concluded that late naïve cells do not harbor inactive X-chromosome rather they have
partially reactivated X-chromosome. If the late naïve cells underwent X-inactivation, then we
should have expected monoallelic expression of most of the X-linked genes, which we did not
observe. Next, we looked if X-dampening is leading the reduction in X-linked gene expression
in late naïve cells. In fact, from the allelic analysis of X-linked gene expression it is clear that
late naïve cells harbor partially reactivated X-chromosome and if they harbored dampened X-
chromosomes, we should have expected biallelic expression of most of the genes, which we
did not observe. In addition, we compared median expression of biallelically expressed genes
of early cells with that of the late naïve cells. If there are X-dampening, we should have seen
significant decrease in median expression. However, we did not find any significant differences
(Fig. 2C). Therefore, we concluded that there is neither X-inactivation or dampening upon
conversion of early to late naïve cells. Next, to find out the mechanism behind the reduction in
X-linked gene expression in late naïve cells we checked X-chromosome ploidy of the cells. We
thought, it may be possible that upon conversion of early to late naïve state, cells are losing one
X-chromosome. Though, from the allelic expression pattern in late naïve cells, it is clear that
cells harbored two X-chromosomes as evident by the biallelic expression of some X-linked
genes (Fig. 2A). However, it is also possible that they may lose some part of the X-chromosome
in late naïve cells and which is leading to the reduction in X-linked gene expression. To test
this, we analyzed gene expression ratio of X-linked genes across the X-chromosome and we
did not find significant differences between early vs. late naïve cells (Fig. 2D; Supplementary
file 3). Therefore, we confirmed that loss of X-chromosome is not the cause of reduction in X-
linked gene expression in late naïve cells, some other mechanism might be existing for this.

Erasure of X-upregulation might be the cause of reduced X-linked gene expression in late
naïve cells

We investigated if the erasure of X-chromosome upregulation might be leading the reduction
in X-linked gene expression in late naïve cells. Sahakyan *et al.* in their analysis of X-linked
gene expression did not consider the possibility of having upregulated X-chromosome in
primed cells. Recent years, existence of X-chromosome upregulation has been increasingly
documented in mammals (Chen and Zhang, 2016; Lin et al., 2011; Sangrithi et al., 2017; Wang
et al., 2016). Moreover, De Mello *et al.* recently have shown that indeed primed hPSCs harbor
upregulated active-X chromosome. Based on this, we first wanted to confirm if UCLA1 primed
cells truly harbor upregulated active-X chromosome. To test this, we calculated X to autosomal
(X:A) expression ratio. For this, we used whole cell population RNA-Seq dataset as the

scRNA-Seq dataset was not available for UCLA1 primed cells. If primed cells harbor one

upregulated active-X chromosome and one inactive-X chromosome then we should expect X:A

ratio more than 0.5 and near to 1. Indeed, we found X:A ratio ~0.8 in primed cells, which

strongly suggests that UCLA1 primed cells harbor upregulated active-X chromosome (Fig. 2E;

Supplementary file 3). Interestingly, we found significant increase in X:A ratio upon transition

of primed to early naïve state (Fig. 2E). As there was X-reactivation upon transition of primed

to early naïve cells, it is obvious that X:A ratio will increase provided there was not complete

erasure of X-upregulation as well. On the other hand, we observed decrease in X:A ratio from

early to late naïve state, which indicated that it might be possible that the X-upregulation is

getting erased upon transition of early to late naïve cells and thereby leading to the overall X-

linked gene expression. Albeit, the decrease in X:A ratio from early to late naïve state is not

significant as expected. We think increase in biallelic X-liked gene expression from early to

late naïve state is masking this to some extent. To make sure that changes in X:A ratio is not

contributed by the autosomal gene expression we compared autosomal gene expression in

primed early naïve and late naïve cells, which showed no significant differences (Fig. 2F).

Altogether, we think that X-dampening is not the causing component for the reduction in X-

linked gene expression in late naïve cells and erasure of X-upregulation might be leading to

the decrease of X-linked gene expression (Fig. 2G).

Discussion

The initiation of X-chromosome dosage compensation in human remains elusive till date.

Although it has been proposed recently that X-chromosome dosage compensation is initiated

in female human preimplantation embryos through the dampening of X-linked gene expression

from both X-chromosomes, other studies have provided evidence of inactivation of one of the

X-chromosome. The major reason behind why till date this issue remains unresolved is the lack

of availability of surplus number of human embryos for the experiment. Therefore, hPSCs

derived from the human embryos serve as the alternative system to resolve this issue. However,

conventional hPSCs represent the primed state instead of the naïve state (Davidson et al., 2015;

Sahakyan et al., 2017b). Recently, Sahakyan *et al*. converted primed hPSCs to naïve state to

model the X-chromosome states of preimplantation embryos. They suggested that naïve hPSCs

have dampened X-chromosomes like preimplantation blastocysts, which strengthened the

dampening observation of human embryos a step further. In this study, using scRNA-seq

dataset, we have demonstrated that upon conversion of primed hPSCs to naïve state, there is
partial reactivation of the inactive-X chromosome instead of the X-chromosome dampening.

Sahakyan *et al.* demonstrated that conversion of primed to naïve state happened through one
intermediate early naïve state; i.e. Primed to early naïve and then to late naïve (Fig1A). They
studied X-chromosome dynamics during the conversion of primed to naïve state using bulk
RNA-sequencing using cell population. Here, we have analyzed the single cell RNA-Seq
dataset generated from the early naïve state and late naïve state. Single cell RNA-Seq provides
better clarity on cellular state and gene expression to distinguish the heterogeneity among the
cells in a population. We found that early naïve cells were mostly *Xist* negative and becomes
Xist positive in late naïve state along with the reduction in X-linked gene expression, which is
consistent with the Sahakyan *et al.* Moreover, we found in late naïve state 32% cells showed
Xist expression from both X-chromosomes, similar to Sahakyan *et al.* In contrary, in
preimplantation embryos ~85% cells express *Xist* from both X-chromosomes (Okamoto et al.,
2011). It is believed that *Xist* might have important role in X-dampening. However, we found
that reduction in X-linked gene expression was *Xist* independent as *Xist* biallelic, monoallelic
and negative cells showed almost similar level of gene expression in late naïve cells (Fig. 1E).
Next, our allelic analysis of X-linked gene expression revealed that early and late naïve cells
harbored partially reactivated X-chromosome (Fig. 2A), which is dissimilar to the observation
of Sahakyan *et al.* where they found almost all genes they analyzed showed biallelic
expression. Importantly, we found huge variation in allelic pattern of gene expression in
different cells, such as same gene in some cells is monoallelic, whereas in others biallelic.
Therefore, in that scenario, whole cell population analysis will always show biallelic
expression of the genes, indeed which was the case in Sahakyan's study. In addition, we
observed that the frequency of biallelic is increasing in late naïve cells compared to the early
naïve cells, which indicated that the cells are still undergoing through the reactivation process.
Besides this, we did not find significant difference of the median expression of biallelically
expressed genes between early and late naïve cells, which indicated that there is no X-
dampening. Altogether, we concluded that the conversion of primed to naïve state is associated
with the incomplete reactivation of X-chromosome instead of X-inactivation or X-dampening.
Importantly, some observations by Sahakyan *et al.* also indicated that there is an incomplete
erasure of epigenetic memory on the inactive X in the naïve state. They found, upon
differentiation, naïve hPSCs undergo non-random X-inactivation dissimilar to the normal
development, and the same X-chromosome that was inactive in the primed hPSCs got selected

for inactivation. Even, the naïve cells showed an accumulation of H3K27me3 repressive marks

on one of the X-chromosomes, that has not been seen in preimplantation blastocyst (Okamoto

et al., 2011). We think presence of H3k27me3 is the result of incomplete erasure of inactive-X

marks, which support our observation of partially reactivated X-chromosome. Taking these

facts together, we strongly suggest that the naïve cells is still in the process of removal of

inactive-X epigenetic marks. In contrary, Sahakyan *et al.* found that RNA-fluorescent in situ

hybridization (FISH) analysis of three X-linked genes (*HUWE1, ATRX, and THOC2*) showed

biallelic expression in naïve cells. We think this small number of genes assayed is not enough

to conclude the X-chromosome wide gene expression status.

On the other hand, we found presence of upregulated X-chromosome in primed UCLA1.

Moreover, our analysis indicates that there might be erasure of X-upregulation upon conversion

of primed to naïve state. In fact, Theunissen *et al.* found a reduction in X-linked gene

expression in male naïve hPSCs compared to that in primed hPSCs, which may be due to the

erasure of X-chromosome upregulation (Theunissen et al., 2016). Our data also indicate that

the erasure of upregulation is might be incomplete in early naïve state and gets completed in

late naïve state (Fig. 2E; 2G). In that scenario, we believe that erasure of X-upregulation is

leading to the reduction in X-linked gene expression upon transition of early to late naïve state.

Therefore, in future, thorough analysis of X-chromosome dynamics in light of erasure of X-

upregulation is important.

In conclusion, our study provides compelling evidence that X-chromosome states in naïve

hPSCs do not reflects the X-chromosome states of human preimplantation embryos. We

suggest that the conversion of primed cells to the naïve state leads to the partial reactivation of

the inactive X-chromosome instead of chromosome wide dampening. Importantly, our data

indicates that erasure of X-upregulation might leading the reduction in X-linked gene

expression in naïve hPSCs. Finally, we think more improvement is necessary to establish a

suitable culture condition to establish naïve hPSCs recapitulating the X-chromosome states of

preimplantation embryos.

Methods

Key resources table

Softwares/Tools	Source/ Reference	Identifiers (RRID)
FASTQC	http://www.bioinformatics.babraham.ac.uk/projects/fastqc/	SCR_014583
STAR	https://github.com/alexdobin/STAR	SCR_015899
SAMtools	https://github.com/samtools/samtools	SCR_002105
HTSeq-count	https://htseq.readthedocs.io/en/release_0.9.1/	SCR_005514
Picard	http://broadinstitute.github.io/picard/	SCR_006525
GATK	https://software.broadinstitute.org/gatk/	SCR_001876
R	https://www.r-project.org/	SCR_001905
CGH Explorer	http://www.softgenetics.com/CGHExplorer.html	SCR_010920

Experimental model and subject details NA

Method details

Data acquisition: RNA-Seq datasets used for the analysis were acquired from Gene Expression Omnibus (GEO) under the accession number GSE872309 (Sahakyan et al., 2017a). We analyzed 192 single cell RNA-Seq data consisting of 96 early naive (Exp2) and 96 late naive (Exp1) cells. We also analyzed whole cell population RNA-Seq data from UCLA1 primed, UCLA1 Clone4 *Xist* negative, UCLA1 Clone12 *Xist* positive, UCLA1 Clone9 *Xist* positive, each having two replicates.

Reads mapping and counting: Reads were mapped to the human genome (hg38) using STAR with default settings (Dobin et al., 2013). Mapped reads were then processed using SAM tools (Li et al., 2009) and the number of reads mapping to each gene was counted using HTSeq-count (Anders et al., 2015).

Expression analyses: For both single cell data and whole cell population dataset, the counted reads from HTSeq were used to calculate RPKM (Reads Per Kilobase per Million) using rpkmforgenes (Ramsköld et al., 2009). For single cell RNA-Seq analysis, we checked for *Xist* expression in all 192 single cells. For early naive cells we selected 60 cells out of 96

showing very low *Xist* expression (0 to 0.5 rpkm) and for late naive cells, 59 out of 96 showing

very high *Xist* expression (3 to 39 rpkm). For quality control of the selected (59+60) cells, we

considered RPKM sum (> 500) and mean (> 0.2) and we reduced to 28 early naive cells and

30 late naive cells. Outlier cells were removed from the further analysis. To check the

expression of X-linked genes, we considered only those genes which were having $5 < \mu_{RPKM} <$

200 (n=57 genes) and these genes were only considered for further downstream analyses.

Variant calling and SNP annotation: Aligned reads from the 58 (28 early naive + 30 late

naive) single cells were marked for duplicate reads using Picard tools

v2.18.11(https://broadinstitute.github.io/picard/). To retrieve allelic read counts for SNPs,

GATK (v3.8) (Van der Auwera et al., 2013; Depristo et al., 2011) "HaplotypeCaller" was used.

Only validated SNPs specific to UCLA1 cell line (GSM2420529) were taken into consideration

for downstream analysis. SNPs were then annotated using dbSNP Build 152 (GRCh38.p12).

We considered a SNP as informative, which showed biallelic expression in at least one cell.

Allele expression analyses: The allelic expression of two different alleles was calculated by

directly counting the allele-specific reads covering a SNP position mapped to the reference or

the alternative allele and then dividing it by the total number of reads covering that

position. Low coverage SNPs (≤ 5 reads) were excluded from the analysis, and SNPs that

occurred at least in ten different cells were only retained. Biallelic expression was considered

when an allele expression ratio ranged from 0.1 to 0.9 and monoallelic expression otherwise.

Also, we only considered allelic ratios of SNPs (n=12, covering 10 X-liked genes) that were

having RPKM ≥ 4 or else deemed as not determined.

Ploidy: To analyze the X-chromosomal ploidy, we considered 73 genes having RPKM mean

≥ 3 distributed across the X-chromosome for 56 cells (28 early naive cells + 28 late naive cells).

We calculated gene expression ratio through dividing the individual gene's RPKM by its

median value across all the 56 cells. To visualize the profile of X-chromosome in all these

cells, the normalised gene expression values (gene expression ratio) were used to build a

moving average plot (neighbourhood size, k=9) using methods described by Mayshar et al.

(Mayshar et al., 2010).

X chromosome to autosomes expression ratio: For calculation of X to autosomal gene

expression ratio (X:A) we used whole cell population RNA-Seq dataset. For each cell line,

normalized gene expressions for all the expressed autosomal and X-linked genes (RPKM>0)

excluding escapees and genes in the pseudo autosomal regions of X-chromosome were

considered for X:A ratio. X:A ratios were calculated by dividing the median RPKM values of

the X chromosome by those of the autosomes. Also, for comparison of autosomal gene

expression analysis in these cell lines, we considered the expressed genes with RPKM>0.

Statistical tests and Box-plots: All statistical tests and boxplots were performed using R

version 3.5.1(R Development Core Team, 2016).

Figure legends

Figure 1: Transition of early naïve to late naïve state is associated with increased *Xist*

expression and reduction in X-linked gene expression. (A) Schematic representation of

different stages of conversion of primed hPSCs to naïve state as described in Sahakyan *et al*.

We performed analysis of single cell RNA-Seq dataset acquired from early and late naïve state.

(B) Comparison of *Xist* expression (RPKM) between early naïve (n=28 cells) and late naïve

cells (n=30 cells). $p<0.00001$ (Mann Whitney U test) (C) Comparison of X-linked gene

expression (n=57) between early and late naïve state. $p<0.00001$ (Mann Whitney U test) (D)

Allelic expression of *Xist* in late naïve cells (n=25 cells). (E) Comparison of X-linked gene

expression among Xist biallelic, Xist monoallelic and Xist negative cells. Non-significant at

$p<0.05$ (Kruskal-Wallis test)

Figure 2: Reduction in X-linked gene expression in naïve hPSCs not because of X-

dampening and might be due to the erasure of X-upregulation. (A) Allelic expression

analysis of X-linked genes in early and late naïve cells at single cell levels. Right, validation

of our allelic gene expression analysis by analyzing allelic expression of an autosome (Chr17).

(B) Histogram showing the percent of SNPs showing monoallelic and biallelic in each cells of

early and late naïve state. (C) Comparison of median of expression level of biallelically

expressed genes between early and late naïve cells. Non-significant at $p<0.05$ (Kruskal-Wallis

test) (D) Ploidy analysis of the cells of early and late naïve state through the analysis of gene

expression ratio across the X-chromosome. (E) Analysis of ratio of X-linked gene expression

and autosomal gene expression in primed, early naïve and late naïve cells. $P<0.05$ (Student's

t-test) (F) Comparison of autosomal gene expression among primed, early naïve and late naïve

cells. Non-significant at $p<0.05$ (Kruskal-Wallis test) (G) Proposed model representing the X-

chromosome states during the conversion of primed hPSCs to the naïve state.

Author's Contribution

SG, SM (Susmita Mandal), DC, and HK conceptualized the study. SG supervised the study. Bioinformatic analyses was done by SM (Susmita Mandal) and DC. SG, SM (Susmita Mandal), DC, MA, and SM wrote, edited and proofread the manuscript. Final manuscript was edited and approved by all the authors.

Acknowledgments

We thank Pavithra RV and Karunyaa M for their help in art work and Amitesh Panda for taking part during the discussion. The study is supported by IISC-MHRD Start up grant, Ramalingaswamy-DBT fellowship to SG, and IISC-DBT partnership program.

References:

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq-A Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166–169.

Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., et al. (2013). From fastQ data to high-confidence variant calls: The genome analysis toolkit best practices pipeline. Curr. Protoc. Bioinforma.

Chen, X., and Zhang, J. (2016). The X to autosome expression ratio in haploid and diploid human embryonic stem cells. Mol. Biol. Evol. *33*, 3104–3107.

Davidson, K.C., Mason, E.A., and Pera, M.F. (2015). The pluripotent state in mouse and human. Development *142*, 3090–3099.

Deng, X., Hiatt, J.B., Nguyen, D.K., Ercan, S., Sturgill, D., Hillier, L.W., Schlesinger, F., Davis, C.A., Reinke, V.J., Gingeras, T.R., et al. (2011). Evidence for compensatory upregulation of expressed X-linked genes in mammals, Caenorhabditis elegans and Drosophila melanogaster. Nat. Genet. *43*, 1179–1185.

Depristo, M.A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J.R., Hartl, C., Philippakis, A.A., Del Angel, G., Rivas, M.A., Hanna, M., et al. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat. Genet. *43*, 491–

501.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., and Gingeras, T.R. (2013). STAR: Ultrafast universal RNA-seq aligner. Bioinformatics

29, 15–21.

Kaur, H., Rv, P., and Gayen, S. (2019). Dampened X-chromosomes in human pluripotent

stem cells: dampening or erasure of X-upregulation ? Chromosoma *1–3*.

Larsson, A.J.M., Coucoravas, C., Sandberg, R., and Reinius, B. (2019). Transcriptional

kinetics of X-chromosome upregulation. 1–6.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools.

Bioinformatics *25*, 2078–2079.

Lin, H., Halsall, J.A., Antczak, P., O'Neill, L.P., Falciani, F., and Turner, B.M. (2011).

Relative overexpression of X-linked genes in mouse embryonic stem cells is consistent with

Ohno's hypothesis. Nat. Genet. *43*, 1169–1170.

De Los Angeles, A., Ferrari, F., Xi, R., Fujiwara, Y., Benvenisty, N., Deng, H.,

Hochedlinger, K., Jaenisch, R., Lee, S., Leitch, H.G., et al. (2015). Hallmarks of

pluripotency. Nature *525*, 469–478.

Lyon, M.F. (1961). Gene action in the X-chromosome of the mouse (mus musculus L.).

Nature *190*, 372–373.

Mayshar, Y., Ben-David, U., Lavon, N., Biancotti, J.C., Yakir, B., Clark, A.T., Plath, K.,

Lowry, W.E., and Benvenisty, N. (2010). Identification and classification of chromosomal

aberrations in human induced pluripotent stem cells. Cell Stem Cell *7*, 521–531.

De Mello, J.C.M., Fernandes, G.R., Vibranovski, M.D., and Pereira, L. V. (2017). Early X

chromosome inactivation during human preimplantation development revealed by single-cell

RNA-sequencing. Sci. Rep. *7*, 1–12.

Nichols, J., and Smith, A. (2009). Perspective Naive and Primed Pluripotent States. Stem Cell

4, 487–492.

Ohno, S. (1967). Sex chromosomes and sex-linked genes.

Okamoto, I., Patrat, C., Thépot, D., Peynot, N., Fauque, P., Daniel, N., Diabangouaya, P.,

Wolf, J.P., Renard, J.P., Duranthon, V., et al. (2011). Eutherian mammals use diverse
strategies to initiate X-chromosome inactivation during development. Nature *472*, 370–374.

Petropoulos, S., Deng, Q., Panula, S.P., Codeluppi, S., Reyes, A.P., Linnarsson, S., Sandberg,
R., and Lanner, F. (2016). Single-cell RNA-seq reveals lineage and X-chromosome dynamics
in human preimplantation embryos. Cell.

R Development Core Team (2016). R: A language and environment for statistical computing.
R Found. Stat. Comput.

Ramsköld, D., Wang, E.T., Burge, C.B., and Sandberg, R. (2009). An abundance of
ubiquitously expressed genes revealed by tissue transcriptome sequence data. PLoS Comput.
Biol. *5*, e1000598.

Sahakyan, A., Kim, R., Chronis, C., Sabri, S., Bonora, G., Theunissen, T.W., Kuoy, E.,
Langerman, J., Clark, A.T., Jaenisch, R., et al. (2017a). Human Naive Pluripotent Stem Cells
Model X Chromosome Dampening and X Inactivation. Cell Stem Cell *20*, 87–101.

Sahakyan, A., Plath, K., and Rougeulle, C. (2017b). Regulation of X-chromosome dosage
compensation in human: Mechanisms and model systems. Philos. Trans. R. Soc. B Biol. Sci.
372.

Saiba, R., Arava, M., and Gayen, S. (2018). Dosage compensation in human pre-implantation
embryos: X-chromosome inactivation or dampening? EMBO Rep. *19*, e46294.

Sangrithi, M.N., Royo, H., Mahadevaiah, S.K., Ojarikre, O., Bhaw, L., Sesay, A., Peters,
A.H.F.M., Stadler, M., and Turner, J.M.A. (2017). Non-Canonical and Sexually Dimorphic X
Dosage Compensation States in the Mouse and Human Germline. Dev. Cell *40*, 289–301.e3.

Theunissen, T.W., Friedli, M., He, Y., Planet, E., O'Neil, R.C., Markoulaki, S., Pontis, J.,
Wang, H., Iouranova, A., Imbeault, M., et al. (2016). Molecular Criteria for Defining the
Naive Human Pluripotent State. Cell Stem Cell *19*, 502–515.

Wang, F., Shin, J., Shea, J.M., Yu, J., Bošković, A., Byron, M., Zhu, X., Shalek, A.K.,
Regev, A., Lawrence, J.B., et al. (2016). Regulation of X-linked gene expression during early
mouse development by Rlim. Elife *5*, 1–19.

Zhang, H., Gayen, S., Xiong, J., Zhou, B., Shanmugam, A.K., Sun, Y., Karatas, H., Liu, L.,
Rao, R.C., Wang, S., et al. (2016). MLL1 Inhibition Reprograms Epiblast Stem Cells to

Naive Pluripotency. Cell Stem Cell *18*, 481–494.

Fig.1

Fig.2