Temporal changes in genetic admixture are linked to heterozygosity and health diagnoses in
humans

Brian S. Mautz,[1,2] Jacklyn N. Hellwege,[1,2] Chun Li,[4] Yaomin Xu,[5] Siwei Zhang,[5] Joshua C. Denny,[2,7]
Dan M. Roden,[1,2,6,7] Tracy L. McGregor,[8,9] Digna R. Velez Edwards,[1,3,7,9] and Todd L. Edwards[1,2,9*]

[1] Vanderbilt Epidemiology Center and Vanderbilt Genetics Institute, Vanderbilt University
School of Medicine, Nashville, TN, USA

[2] Department of Medicine, Vanderbilt University School of Medicine, Nashville, TN, USA

[3] Division of Quantitative Sciences, Department of Obstetrics and Gynecology, Vanderbilt
University School of Medicine, Nashville, TN, USA

[4] Department of Population and Quantitative Health Sciences, Case Western Reserve
University, Cleveland, OH, USA

[5] Department of Biostatistics, Center for Quantitative Sciences, Vanderbilt University School of
Medicine, Nashville, TN, USA

[6] Department of Pharmacology and Biomedical Informatics, Vanderbilt University School of
Medicine, Nashville, TN, USA

[7] Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville,
TN, USA

[8] Department of Pediatrics, Vanderbilt University School of Medicine, Nashville, TN, USA

[9]Senior author

[*]Correspondence: todd.l.edwards@vanderbilt.edu

Abstract

Reproduction between individuals from different ancestral populations creates genetically admixed offspring. Admixture can have positive and negative impacts on individual health, feeding back to population health. Historical and forced migrations, and recent mobility, have brought formerly disparate populations of humans together. Here we sought to better understand how temporal changes in genetic admixture influence levels of heterozygosity and health outcomes. We evaluated variation in ancestry over 100 birth years in 35,842 individuals from a genetic database linked to health records in a population in the Southeastern United States. Analysis of 2,678 ancestrally informative markers revealed increased admixture and heterozygosity for all clinically-defined race groups since 1990. Most groups also exhibited increasing long-range linkage disequilibrium over time. A phenome-wide association study of clinical outcomes detected protective associations with female reproductive disorders and increased risk for diseases with links to autoimmunity dysfunction. These mixed effects have important ramifications for epidemiology of human health.

Keywords: electronic health records, genetic admixture, heterozygosity fitness correlation, health outcomes, long-range linkage disequilibrium

Introduction

Understanding temporal changes in heterozygosity has important implications for individual- and population-level health in humans that remain unexplored. The heterozygosity-fitness correlation (HFC) is a measure used to quantify the extent to which proxies of fitness are positively correlated with the level of heterozygosity across sets of genetic loci or the whole genome (David, 1998; Head et al., 2017; Szulkin et al., 2010). Evidence linking fitness components to levels of heterozygosity has been found in animals (Keller and Waller, 2002), but results are equivocal and controversial (Chapman et al., 2009; Szulkin et al., 2010). HFCs arise through some level of inbreeding depression or can be generated through immigration and population admixture (reviewed in Szulkin et al., 2010). Both positive and negative effects on individual and population health can occur.

Three factors make human population genetic data ideal to study the connection between heterozygosity and health. First, the human genome is sequenced and genetic databases with large numbers of genetic loci and individuals are readily available. Second, there are substantial numbers of subjects with genetic data linked to electronic health records (EHR) (Roden et al., 2008; Sudlow et al., 2015). Third, the diseases and their etiologies used in health records are known in much more detail and well categorized than most other species, facilitating the estimation the relationship between heterozygosity and changes in (disease) phenotypes used as fitness proxies.

Additionally, we have detailed information on the movements of many human populations. Geographically disparate human populations have come into contact through

migration, as well as more recent mobility and urbanization, increasing the opportunity for

genetic admixture and HFCs to occur. Genetic admixture has previously been used to identify

migration patterns across several humans populations (Baharian et al., 2016; Bryc et al., 2010,

2015; Han et al., 2017; Moreno-Estrada et al., 2014; Tishkoff et al., 2009; Wang et al., 2013) and

the genetic basis of diseases (Franceschini et al., 2013; Kato et al., 2015; Reich and Patterson,

2005). Two studies have shown temporal increases in heterozygosity due to urbanization, one

in a Croatian population (Rudan et al., 2008) and one in a US population of European ancestry

(Nalls et al., 2009). However, these studies of admixture have not connected migratory or

urbanization patterns to health outcomes.

Finally, accruing evidence in humans suggests HFCs occur. For example, several studies

have linked levels of heterozygosity in non-admixed populations to specific disease states

(Bittles and Black, 2010; Campbell et al., 2007, 2009; Pemberton and Szpiech, 2018).

Additionally, a meta-analysis of populations of European American and African American

ancestry found a positive association between levels of heterozygosity and mortality in humans

(Bihlmeyer et al., 2014). However, these studies are limited in scope, investigating single

outcomes and ignoring the impact of temporal trends in admixture and heterozygosity on

epidemiological outcomes.

Here, we link temporal changes in genetic admixture to patterns of heterozygosity and

epidemiological outcomes across a broad array of human diseases. We used BioVU, a

biorepository of genetic data linked to EHR (Roden et al., 2008) to investigate temporal changes

in genetic admixture and heterozygosity in a population in the Southeastern United States.

Though race has little biological meaning, we used race as instrumental groupings (Maglo et al.,

2016) to reflect corresponding socio-cultural dynamics in this geographic area and investigate

differential changes to admixture and heterozygosity within these groupings. In addition, we

used a phenome-wide association study (PheWAS, Denny et al., 2010), to connect genetic data

with the clinical phenome representing an entire collection of disease outcomes associated

with EHRs in BioVU.

Results

We identified 2,678 ancestry-informative markers (AIMs) from genetic data in BioVU.

These AIMs were from ExomeChip data in a cohort of 28,723 Whites, 4,129 Blacks, 550

Hispanic/Latinos, and 270 Asians, based on the EHR third-party race designation (see methods).

Sample population demographics by EHR race and birth decade are in presented in

Supplementary Table S1-2 and Figure S1. After combining our data with the 1000 Genomes as a

reference group (Durbin et al., 2010), we calculated principal components to identify patterns

of ancestry in each individual. We used the ancestral classifications to test for temporal trends

in mean and variance in admixture proportion.

Analysis of temporal trends in genetic admixture showed an increase in ancestral

diversity with time (Figure 1; see Table S2). The admixture proportion consistently increased

with younger ages in Asian and Hispanic/Latino groups. In Whites, the level of admixture

remained stable until approximately 1990 when it began to increase. Blacks had a decrease in

admixture proportion in those born the early- to mid-20th century, but an increase after the

1980's. In addition to the distinct clusters in the principal component plots representing

individuals who are identified as either White and Black and who contain both European and

African ancestries, the overall proportions of ancestry across all recorded race groups exhibit

increasing admixture in younger individuals (Figure S3-S7). Additionally, a significant increase in

the variance of admixture proportions over time was observed (variance coefficient = 0.0193 ±

0.0009 [SE], p-value < 1.44x10^{-100}), indicating that there is a linear increase in variance of the

admixture proportion of 0.0193 with every birth year. This is consistent with recent and

ongoing admixture (Verdu and Rosenberg, 2011).

Figure 1. Admixture proportion plotted against birth year. Admixture proportion is defined as (1
- maximum ancestry proportion). The smoothing curves are obtained using the generalized
additive model method with a cubic spline basis implemented in R package mgcv and plotted
using R package ggplot2. Cohorts are grouped by race and 95% confidence intervals are
included.

We detect similar patterns in three additional sources. First, we calculated heterozygosity

across AIMs and tested for temporal trends to compare with patterns of genetic admixture.

There was strong support for increasing heterozygosity in younger individuals (Figure 2; Table

S3 and Figure S8). The timing of inflection for increased standardized heterozygosity varied

between the race groups, but the data indicated that ancestry diversity has accelerated rapidly

in Asian, Black, and White cohorts since about 1980, while Hispanics/Latinos have exhibited a

relatively steady rate of increasing diversity since the 1940s. This reflects the increasing number

of children born to couples of predominantly different ancestral backgrounds over the past few

decades.

Figure 2. Standardized heterozygosity plotted against birth year. The y-axis depicts the
standardized heterozygosity of individuals with birth year on the x-axis. Cohorts are grouped by
stated race and 95% confidence intervals are indicated.

Second, we verified patterns by estimating pairwise long-range linkage disequilibrium in our

dataset. Using common single nucleotide polymorphisms (SNPs) (minor allele frequency [MAF]

> 5%) from the genotyping array, we calculated pairwise D' using Haploview (Barrett et al.,

2005) and plotted D' against physical distance for all pairs of SNPs between 9–10 megabases of

each other (Figure 3; Table S4). Results show a small drop in long range disequilibrium (LRLD) in

individuals born before the 1950s, followed by significant steady increases in the 1950s through

the 1980s and subtle fluctuation at higher levels in the 1990s to 2010s.

Figure 3. Pairwise D' for SNPs between 9-10 Mb for all common SNPs on the exome array in all
study individuals stratified by intervals of birth decade. Pairwise D' for all common SNPs
between 9 - 10 megabases apart for individuals are presented on the y-axis and physical
distance in Mb is presented on the x-axis. The plots are paneled by decades of birth to depict
relative changes over time.

Third, we estimated changes in the number of ethnicities in self-reported ancestry in the

2013 American Communities Survey. Respondents were instructed to select

Hispanic/Latino/Spanish status and all applicable races for each individual in the household

(Ruggles, 2014). We analyzed the average number of race categories selected for each

individual by birth year and stratified these results within race categories for Tennessee (Figure

S9), the East South-Central Region which includes Tennessee, Alabama, Kentucky, and

Mississippi (Figure S10), and the entire United States (Figure S11). The results from all regions

are concordant, indicating that younger individuals are more likely to indicate multiple races.

The inflection point appears to have been earlier in the Asian race category but is demonstrated

in all race groups by the mid-1960s in the entire US sample.

Changes in health diagnoses with admixture

We evaluated the association between individual heterozygosity and disease diagnoses

using a phenome-wide association study approach (PheWAS, Denny et al., 2010). Increasing

genetic admixture resulted in fewer diagnoses of female reproductive traits across all data

(Table 1). These results remain statistically significant when considering adults only (Table S5)

and after correction for multiple tests. Phenotype codes for "disorders of menstruation and

other abnormal bleeding from female genital tract" and "irregular menstrual cycle/bleeding"

were significantly associated with protection by increasing heterozygosity (p-value = 7.21×10^{-6}

and 4.37×10^{-5}, respectively; Table 1, Figure S12). Other protective findings were also

gynecological in nature, including cervical cancer/dysplasia and abnormal Papanicolaou smear

results. Significant phenotypes in adults were predominantly detected for females. Outside

genitourinary findings, other nominally significant associations (Bonferroni significant < p ≤

0.05) show increased risk with genetic admixture and include atopic dermatitis, AV Block,

obstructive asthma, Sicca syndrome.

Table 1. Results from the phenome-wide association study of heterozygosity and clinical
outcomes for full sample.

PheCode	Phenotype	P-value	OR (95% Confidence Interval)
626	Disorders of menstruation and other abnormal bleeding from female genital tract	7.21×10^{-6}	0.37 (0.24 – 0.57)
626.1	Irregular menstrual cycle/bleeding	4.37×10^{-5}	0.37 (0.23 – 0.60)
939	Atopic/contact dermatitis due to other or unspecified	2.86×10^{-4}	1.82 (1.32 – 2.52)
180	Cervical cancer and dysplasia	4.02×10^{-4}	0.19 (0.08 – 0.48)
792.1	Papanicolaou smear of cervix or vagina with atypical squamous cells	5.36×10^{-4}	0.21 (0.09 – 0.51)

180.3	Cervical intraepithelial neoplasia [CIN] [Cervical dysplasia]	6.24×10^{-4}	$0.15 \ (0.05 - 0.45)$
426.2	Atrioventricular [AV] block	7.62×10^{-4}	$2.97 \ (1.58 - 5.61)$
495.11	Chronic obstructive asthma with exacerbation	8.13×10^{-4}	$4.63 \ (1.89 - 11.34)$
785	Abdominal pain	1.04×10^{-3}	$0.70 \ (0.57 - 0.87)$
792	Abnormal Papanicolaou smear of cervix and cervical Human Papilloma Virus (HPV)	1.04×10^{-3}	$0.33 \ (0.17 - 0.64)$
709.2	Sicca syndrome	1.66×10^{-3}	$4.38 \ (1.75 - 10.99)$
318	Tobacco use disorder	1.75×10^{-3}	$0.48 \ (0.30 - 0.76)$
626.12	Excessive or frequent menstruation	1.80×10^{-3}	$0.28 \ (0.12 - 0.62)$
628	Ovarian cyst	2.83×10^{-3}	$0.23 \ (0.09 - 0.60)$
426.24	Atrioventricular block, complete	3.57×10^{-3}	$3.30 \ (1.48 - 7.38)$

Discussion

We found increasing genetic admixture and heterozygosity through time in a population of the

Southeastern United States. Comparing these changes to socio-cultural shifts in this geographic

area puts these results in context. In the Southern United States from the Civil War until the

mid-1960's, laws and policies enforced segregation of populations of European and African

ancestry. Consistent with these socio-cultural boundaries, there is little change in admixture

through the 1960's. Additionally, despite legal rulings and perceived socio-cultural

transformation, there remained a very slow increase in admixture and heterozygosity for an

additional 20-30 years. For individuals with an EHR designation of White, the mean proportion

of European ancestry was greater than 98% until the 1990's but decreased to 92% in the

youngest cohort as the proportion of African ancestry increased to 6% (Figure S4). In individuals

designated as Black in the EHR, the mean African proportion was approximately 80% in those

born before the 1990's but decreased to 77% in those born more recently (Figure S5). The

largest change of any group was in those identified as Hispanic/Latino. The European ancestry

proportion in these individuals decreased by 15%, from approximately 75% in those born

before the 1950's to under 60% after the 1980's. This decrease in European ancestry was

associated with an increase in the East Asian ancestry proportion, likely a proxy for Native

American ancestry, while the African proportion remained under 10% (Figure S6). The

continental ancestry proportions in EHR-designated Asians were more difficult to discern

because of the small cohort size, although we did observe a slight decrease in the East Asian

proportion over time (Figure S7). These trends are consistent with previous reported trends in

decreases in homozygosity in other populations (Nalls et al., 2009; Rudan et al., 2008).

Long-range linkage disequilibrium occurs when subdivided populations mix (Koch et al.,

2013; Nei and Li, 1973). Consistent with this and with patterns we detected in admixture and

heterozygosity, we find that LRLD changes through time (Figure 3). There was small drop in

LRLD for older individuals born before 1950. This was followed by a consistent increase in LRLD

in individuals born up to 1980. Subtle fluctuations at higher LRDL levels occurs in those

individuals born between 1990 and 2010. It should be noted, however, that we cannot

necessarily rule out other possible sources of LRLD (e.g. drift, epistatic selection, hitchhiking as

in Koch et al., 2013), though they seem unlikely given the recent nature of admixture and

formation of LRLD.

Our results are qualitatively echoed in the 2013 American Communities Survey. Younger

individuals, across multiple geographic areas (Tennessee, East South-Central Region, United

States) are more likely to indicate multiple races when asked about their ancestral origins

(Figure S9-11). Across race group the inflection point occurs by the mid-1960's of the entire US

sample, though appears earlier for the Asian race category.

Changes in health diagnoses with admixture

We show that changes in population genetic parameters have important consequences

for population health (Table 1; Table S5 and Figure S12), which have not been previously

documented. There were several statistically significant associations of genetic admixture with

adult female genitourinary diagnosis codes (e.g. irregular menstrual cycle/bleeding; cervical

cancer/dysplasia). These findings linking admixture to protection from menstruation and

gynecological abnormalities suggests that ancestral diversity is related to risk of disorders that

could affect reproduction. This is consistent with long-standing models that propose ancestral

diversity and heterozygosity is related to fertility in plants and animals, as well as other

beneficial characteristics in offspring (Dickerson, 1973; Keller and Waller, 2002; Lamkey and

Edwards, 1999; Shull, 1948; Szulkin et al., 2010).

The changes in reproductive diagnoses were detected predominantly for females,

suggesting a largely sex-specific response to changes in admixture and heterozygosity. In some

animals, there are sex-specific responses in reproduction to inbreeding (Keller and Waller,

2002), providing some circumstantial support for our findings. However, the sex-specific

response we detected could also be a results of gender differences in other factors. For

example, men are less likely to seek medical attention and report medical issues (Galdas et al.,

2005; Mansfield et al., 2003). Alternatively, male reproductive traits, such as sperm quality,

might not be routinely checked and reported as with female reproductive parameters. More

research is needed to disentangle biological and sociological factors driving the sex-specific

responses to determine the source of these trends.

Other interesting patterns emerge when considering all data and nominally significant

PheWAS results (i.e. $p < 0.05$). Several of these disease phenotypes, show positive associations

with increasing genetic admixture. Importantly, each of these diseases have at least suggestive

links to autoimmunity issues (*atopic dermatitis*: Mittermann et al., 2004, *AV block*: Buyon et al.,

1998; Villuendas et al., 2014, *asthma*: Barnes, 2008; Tedeschi and Asero, 2008, *Sicca/Sjögren*

syndrome: Brito-Zerón et al., 2016). The nominally significant diseases detected here also have

reported connections with each other in the literature. For example, atopic dermatitis and

asthma have a similar mechanistic basis via allergen exposure through the epidermis (Spergel,

2010). Similarly, autoimmune congenital heart block is linked to Ro and La autoantigens,

Sjögren-syndrome-related antigen A and B respectively, which cross the placenta from the

mother to the fetus to cause cardiac abnormalities and cutaneous rashes (Brito-Zerón et al.,

2015). These patterns suggest increasing heterozygosity are linked to increases in activity in the

immune system. Our results show continued increases in genetic admixture with time. If

heterozygosity is positively linked to autoimmunity-related disease, then we would predict

continued increases in prevalence of these types of disease with time. Future research should

address these intriguing immunity-disease relationships to determine the validity and

consistency of these patterns within and across human populations.

These results have important ramifications in several areas. First, this work has

implications for genetic association study designs, disease prevalence and disparities, and

classification of individuals for their medical care. Frequently genetic analyses are stratified by

ancestry or race, and our data show that simple categorical race classifiers, whether by self-

report or derived from other data, are becoming less accurate over time. When conducting

genetic studies in children, researchers may need to collect the race(s) of the parents or even

grandparents to allow for appropriate classification, or consider AIMs to estimate ancestry

proportions (Kodaman et al., 2013; Ruiz-Narvá Ez et al., 2011). This increase in ancestral

diversity of the younger population may also enable admixture mapping studies in populations

not traditionally considered for these types of studies.

Second, understanding race-associated factors in patients with complex ancestries may

be increasingly important for effectively delivering precision medical care, which is a major

National Institutes of Health (NIH) goal and one of the key promises of the genetics research

enterprise in the United States. Genetic information has been used to investigate the roots of

human evolution and subsequent diversity. These same approaches can be applied to analyze

ongoing trends in regional and national demographics which have significant implications for

scientific, societal, and governmental policies.

Finally, these results have important implications for HFCs and their use in biological

research (Keller and Waller, 2002; Chapman et al., 2009; Szulkin et al. 2010). We detected

positive changes in reproductive traits in females, providing evidence that genetic admixture

and heterozygosity are linked to potential fitness-related components. Though there were

protective effects for reproductive diagnoses, our results provide additional details showing

increased risk associated with other disease states that might be linked with the immune

system as genetic admixture increases and that HFCs might be sex-specific. The effect of

changes to levels on health and fitness related parameters are complex, so HFCs should be

interpreted with caution and more detail is needed for proper interpretation.

Materials and methods

Individuals were selected from the BioVU DNA Repository which links clinical data from

de-identified electronic medical records to DNA specimens obtained from patients at Vanderbilt

University Medical Center (VUMC) (Roden et al., 2008). Each individual is assigned a single race

in the Electronic Heath Record (EHR) of White, Black, Asian, Pacific Islander, American

Indian/Alaska Native, or declined/unknown, and an ethnicity of Hispanic/Latino, Not

Hispanic/Latino, or declined/unknown. BioVU also contains third-party-designated race, which

is a good predictor of self-reported ethnicity/ancestry in this database (Dumitrescu et al., 2010).

This study of de-identified data was determined to be non-human subject research by the

institutional review board (IRB) of Vanderbilt University, Nashville, TN.

DNA Extraction and Genotyping

All DNA samples were isolated from whole blood using the Autopure LS system (QIAGEN

Inc., Valencia, CA). Genomic DNA was quantitated via an ND-8000 spectrophotometer and DNA

quality was evaluated via gel electrophoresis. Individuals were genotyped using the Illumina

Infinium HumanExome Array [12v1-1] (Illumina Inc., San Diego, CA). The data were processed

for genotype calling using Illumina's Genome Studio (Illumina Inc., San Diego, CA).

Genotyping Quality Control

Data on 240,117 SNPs and 35,842 individuals (16,289 males, 19,552 females) were

available prior to implementation of quality control (QC) measures. No individuals were

excluded for low genotyping efficiency (<98%). 6,599 SNPs were excluded for low genotyping

efficiency (<98%) and 71,667 SNPs were monomorphic. Twenty-six individuals (14 EHR males,

12 EHR females) were excluded for inconsistent genetic and database sex. After QC, 163,135

SNPs remained for analyses in 35,456 individuals. No SNPs were removed for deviations from

Hardy-Weinberg equilibrium.

Quantification and Statistical Analyses

Descriptive statistics of demographic and clinical characteristics were expressed as

mean with standard deviation or median with interquartile range for continuous covariates and

as frequencies or proportions for categorical data using SPSS statistical software (IBM

Corporation, Armonk, NY).

A subset of 2,678 ancestry-informative markers (AIMs) were selected for subsequent

analysis. We selected AIMs from the ExomeChip content, which were selected to have strong

differences between African and European ancestry populations as well as Native American and

European ancestry populations. Average differences in allele frequency, Δ_p, between 1000

Genomes (Price et al., 2006) continental populations (European, Asian, Americas, African) for

the AIMs are presented in Table S6. AIMs were used rather than using pruned SNP data due to

the genotype data coming from an ExomeChip platform, which was designed with a panel of

AIMs to enable evaluation of ancestry.

EIGENSTRAT v6.0.1 software was used to conduct principal components analysis (PCA)

to estimate continuous axes of ancestry from AIMs running all populations together (Price et

al., 2006).

STRUCTURE software v2.3.3 (Falush et al.; Pritchard et al., 2000) was used to quantify

ancestry in combined study and 1000 Genomes Project (Price et al., 2006) individuals using the

AIMs. We estimated proportions of ancestry assuming ancestral clusters (K) ranging from one

to 16, where 16 is the number of sub-populations in the 1000 Genomes data plus two. We

assumed unlinked SNPs and used 5,000 iterations of burn-in and 10,000 iterations for analysis

without providing population information to the software. We observed that the –log-

likelihood of the data given K did not vary significantly for K's greater than three and observed

that K's greater than three primarily subdivided the European populations (data not shown).

The three STRUCTURE clusters corresponded to African, European, and Asian ancestry based on

comparisons to the 1000 Genomes reference data.

Each of the three derived proportions of continental ancestry from STRUCTURE were

regressed onto birth year using generalized additive models with integrated smoothness

estimation (GAM) (Wood, 2010) implemented in the R package mgcv in all study individuals,

Non-Hispanic Whites , Non-Hispanic Blacks, Hispanics/Latinos, and Non-Hispanic Asians. We

derived admixture proportion, α, for the ith individual using the formula $\alpha_i = 1 -$ maximum

(%European, %African, %East Asian). We regressed α_i onto birth year using generalized additive

models with integrated smoothness estimation for all study individuals, Non-Hispanic Whites,

Non-Hispanic Blacks, Non-Hispanic Asians, and Hispanics or Latinos. Proportions of continental

ancestry in the Non-Hispanic Whites, Non-Hispanic Blacks, Non-Hispanic Asians, and Hispanics

or Latinos by birth decade are presented.

It has been previously shown by Verdu and Rosenberg (2011) that when parental

populations stop contributing to admixture, that the variance of admixture proportions

decreases rapidly, and when they continue to contribute to the admixture, that the variance of

admixture proportions increases over time. To test null hypothesis that the observed levels of

admixture in our data were not due to ongoing and increasing rates of admixture, we analyzed

the association between the variance of α and birth year. We used the software package

MVtest and modeled the admixture proportion variance as a log-linear function of birth year

and five principal components of ancestry using estimating equations.

Analysis of admixture proportion variance over time

We model admixture proportion mean and variance simultaneously as functions of birth

year and covariates. Specifically, let m_i be the mean and σ_i^2 be the variance of the trait Y_i for

the i-th individual. We model them as

$$m_i = \beta_0 + \beta_g G_i + \sum_{j=1}^p \beta_j X_{ij},$$

and

$$\ln(\sigma_i^2) = \gamma_0 + \gamma_g G_i + \sum_{j=1}^p \gamma_j X_{ij}.$$

where G_i is the birth year or variable of interest for the i-th individual, and $X_{i1}, ..., X_{ip}$ are p

covariates. In this model the variance is monotonic with respect to birth year, an assumption

that holds in most circumstances. The parameters are estimated simultaneously. This

framework allows for testing of the null hypothesis of no effect on mean, variance, or both for

any term in the model. These correspond to a mean test with null $H_0: \beta_g = 0$, a variance test

with $H_0: \gamma_g = 0$, both having one degree of freedom (DF), and a 2-DF test with $H_0: \beta_g = 0, \gamma_g =$

0.

Model fitting with estimating equations

The parameters are estimated through the estimating equations approach, which does

not require a full specification of the outcome distribution, but only a few constraints for the

parameters of interest. These constraints are often written as equations, and the parameter

estimates can be obtained by solving the equations. The asymptotic distribution for the

parameter estimates can be derived (Stefanski and Boos, 2002). Specifically, suppose the

random variable Y_i has mean $m_i = \beta_0 + \beta_g G_i + \sum_{j=1}^{p} \beta_j X_{ij}$, and log-variance $\ln(\sigma_i^2) = \gamma_0 +$

$\gamma_g G_i + \sum_{j=1}^{p} \gamma_j X_{ij}$. There are $k = 2(p + 2)$ parameters, which can be written as a vector

$\boldsymbol{\theta} = (\boldsymbol{\beta}, \boldsymbol{\gamma})$, where $\boldsymbol{\beta} = (\beta_0, \beta_g, \beta_1, \dots, \beta_p)$ and $\boldsymbol{\gamma} = (\gamma_0, \gamma_g, \gamma_1, \dots, \gamma_p)$. Let y_i and $\boldsymbol{x}_i =$

$(1, g_i, x_{i1}, \dots, x_{ip})^T$ be the observed values for subject i. If we had assumed normality for the

outcome, the log-likelihood for observation i would have been

$$l_i(\boldsymbol{\theta}) = -\frac{1}{2}\log(2\pi) - \frac{1}{2}\boldsymbol{\gamma}'\boldsymbol{x}_i - \frac{(y_i - \boldsymbol{\beta}'\boldsymbol{x}_i)^2}{2\exp(\boldsymbol{\gamma}'\boldsymbol{x}_i)},$$

378   for which the partial derivatives with respect to the parameters $\boldsymbol{\theta}$ is a k-vector,

$$\Psi_i(\boldsymbol{\theta}) = \begin{pmatrix} \frac{\partial l_i}{\partial \boldsymbol{\beta}} \\ \frac{\partial l_i}{\partial \boldsymbol{\gamma}} \end{pmatrix} = \begin{pmatrix} \frac{y_i - \boldsymbol{\beta}'\boldsymbol{x}_i}{\exp(\boldsymbol{\gamma}'\boldsymbol{x}_i)}\boldsymbol{x}_i \\ \frac{1}{2}\left[\frac{(y_i - \boldsymbol{\beta}'\boldsymbol{x}_i)^2}{\exp(\boldsymbol{\gamma}'\boldsymbol{x}_i)} - 1\right]\boldsymbol{x}_i \end{pmatrix},$$

379   and the maximum likelihood estimates of the parameters could have been obtained by solving

380   the k equations $\sum_{i=1}^{n} \Psi_i(\boldsymbol{\theta}) = 0$. This motivated us to use these k equations,

$$\sum_{i=1}^{n} \Psi_i(\boldsymbol{\theta}) = 0,$$

as the starting point for our estimating equations approach to obtain parameter estimates

$\widehat{\boldsymbol{\theta}} = (\widehat{\boldsymbol{\beta}}, \widehat{\boldsymbol{\gamma}})$. If normality holds, then $\widehat{\boldsymbol{\theta}}$ are the maximum likelihood estimates. Note that

although the estimating equations were motivated by the Gaussian likelihood, one can always

start from these equations to obtain $\widehat{\boldsymbol{\beta}}$ and $\widehat{\boldsymbol{\gamma}}$, whether normality holds or not, and proceed

with statistical inference using the M-estimation theory (Stefanski and Boos, 2002). This is a

major advantage for using estimating equations. The partial derivative of $\Psi_i(\boldsymbol{\theta})$ is a $k \times k$

matrix, denoted as $\psi_i(\boldsymbol{\theta})$. Using the M-estimation theory, we have

$$\sqrt{n}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{d} N(0, V),$$

388   where the $k \times k$ covariance matrix V can be estimated as $A^{-1}B(A^{-1})^T$ with

389    $A = -\frac{1}{n}\sum_{i=1}^{n}\psi_i(\widehat{\boldsymbol{\theta}})$ and $B = \frac{1}{n}\sum_{i=1}^{n}\Psi_i(\widehat{\boldsymbol{\theta}})\Psi_i(\widehat{\boldsymbol{\theta}})^T$.

390   If our interest is on the effect of G, the asymptotic result for the joint distribution for the

391   parameter estimates $\widehat{\beta}_g$ and $\widehat{\gamma}_g$ is

$$\sqrt{n}\left[\begin{pmatrix}\widehat{\beta}_g \\ \widehat{\gamma}_g\end{pmatrix} - \begin{pmatrix}\beta_g \\ \gamma_g\end{pmatrix}\right] \xrightarrow{d} N(0, V_2),$$

392   where V_2 is the corresponding 2×2 submatrix of V, with diagonal values denoted as $\widehat{\sigma}_{\beta_g}^2$ and

$\widehat{\sigma}_{\gamma_g}^2$, respectively. A mean test (H$_0$: $\beta_g = 0$) can be performed by comparing $\sqrt{n}\widehat{\beta}_g$ with

$N\left(0, \widehat{\sigma}_{\beta_g}^2\right)$, and similarly, a variance test (H$_0$: $\gamma_g = 0$) by comparing $\sqrt{n}\widehat{\gamma}_g$ with $N\left(0, \widehat{\sigma}_{\gamma_g}^2\right)$. A 2-

DF joint test (H$_0$: $\beta_g = 0, \gamma_g = 0$) can be performed by comparing $n(\widehat{\beta}_g, \widehat{\gamma}_g)V_2^{-1}\begin{pmatrix}\widehat{\beta}_g \\ \widehat{\gamma}_g\end{pmatrix}$ with a chi-

squared distribution with two degrees of freedom. MVtest software for genetic analysis of SNP

data or general analysis of variables is freely available at https://github.com/edwards-

lab/MVtest.

Heterozygosity Analysis

Standardized measures of heterozygosity among the AIMs were calculated in order to

evaluate trends in heterozygosity over time relative to expectations. We first estimated the

expected number of heterozygous genotypes in an individual in the k-th subpopulation as

(1) $H_k = \sum_i 2p_{ik}q_{ik}$,

where the sum is over all SNPs in our analysis, and p_{ik} and q_{ik} = 1-p_{ik} are the allele frequencies

for the i-th SNP. Hardy-Weinberg equilibrium was assumed. Then for every individual j in the k-

th subpopulation, we standardized the observed number of heterozygous genotypes, O_{kj}, by

comparing it with the expected number H_k:

(2) $\left(O_{kj} - H_k\right)/H_k$

Standardized heterozygosity was regressed onto birth year using GAM in all study individuals,

and the results were plotted for Non-Hispanic White, Non-Hispanic Black, Non-Hispanic Asian,

and Hispanic or Latino.

Pairwise linkage disequilibrium (LD) D' statistics were calculated for all pairs of common

(MAF > 0.05) SNPs within 10 megabases (Mb) using Haploview software (Wood, 2011). D'

statistics were regressed onto physical distance between SNPs using GAM for distances in the

interval from 9-10 Mb for each birth decade.

We downloaded a 1% representative sample of individual-level responses to the

American Community Survey (ACS) from the Integrated Public Use Microdata Series (IPUMS)

(Ruggles, 2014). Race and ethnicity was self-reported in the ACS IPUMS. We regressed the

number of major race groups claimed by individuals onto their reported birth year using GAM

and frequency weights provided by IPUMS for TN, the South East Central census region, and the

entire US. For individual groups, such as "White" and "Black or African American", we plotted

all individuals who responded affirmatively to those items; thereby, the samples for the

individual race group plots are not independent and overlap at observations where participants

claim two or more race groups.

Phenotype Classification

Each individual was classified according to 1,645 phenotypes based on the International

Classification of Disease, Ninth Revision, Clinical Modification (ICD9) Codes (Denny et al., 2013).

Our classification strategy includes all ICD9 features except for procedures. Additionally, the

system is hierarchical such that disease subtypes are also classified, such as cardiac arrhythmias

are the parent to atrial fibrillation and atrial flutter. Additional phenotypes that are not

represented directly in the ICD9 hierarchy are also included, such as inflammatory bowel

disease as the parent for Crohn's disease and ulcerative colitis. Diagnoses that were not

possible for an individual were set to missing, such as pregnancy for males, or prostate diseases

for females. Detailed features of all phenotype algorithms used are available from:

http://phewascatalog.org.

Clinical Outcomes

For each phenotype, we regressed the binary outcome onto the standardized

heterozygosity from equation 2 above, adjusted for birth year and the top 5 principal

components of ancestry using logistic regression. We limited analysis to outcomes with 40 or

more cases and individuals with at least 2 ICD9 codes. We determined the threshold for

statistical significance by Bonferroni correction for the number of analyses where the model

converged.

Data and Software Availability

The data that support the findings of this study are available from the BioVU resource,

but restrictions apply to the availability of these data, which were used under license for the

current study, and so are not publicly available. Data are however available upon reasonable

request with permission from BioVU.

Acknowledgements

The authors would like to thank J. Henshaw and W. Kunce for useful feedback on the

manuscript. The dataset used in the analyses described were obtained from Vanderbilt

University Medical Centers BioVU which is supported by institutional funding and by the

Vanderbilt CTSA grant UL1 TR000445 from NCATS/NIH. This work was supported by the

National Institutes of Health (grant numbers RC2GM092618, U01HG004603, K23HD000001,

R25/T32CA160056, R01LM010685).

Declaration of Interests

The authors declare no competing interests.

References

Baharian, S., Barakatt, M., Gignoux, C.R., Shringarpure, S., Errington, J., Blot, W.J., Bustamante,

C.D., Kenny, E.E., Williams, S.M., Aldrich, M.C., et al. (2016). The great migration and African-

American genomic diversity. PLOS Genet. *12*, e1006059.

Balloux, F., Amos, W., and Coulson, T. (2004). Does heterozygosity estimate inbreeding in real

populations? Mol. Ecol. *13*, 3021–3031.

Barnes, P.J. (2008). Immunology of asthma and chronic obstructive pulmonary disease. Nat.

Rev. Immunol. *8*, 183–192.

Barrett, J.C., Fry, B., Maller, J., and Daly, M.J. (2005). Haploview: analysis and visualization of LD

and haplotype maps. Bioinformatics *21*, 263–265.

Bihlmeyer, N.A., Brody, J.A., Smith, A.V., Lunetta, K.L., Nalls, M., Smith, J.A., Tanaka, T., Davies,

G., Yu, L., Mirza, S.S., et al. (2014). Genetic diversity is a predictor of mortality in humans. BMC

Genet. *15*, 159.

Bittles, A.H., and Black, M.L. (2010). Consanguinity, human evolution, and complex diseases.

Proc. Natl. Acad. Sci. *107*, 1779–1786.

Brito-Zerón, P., Izmirly, P.M., Ramos-Casals, M., Buyon, J.P., and Khamashta, M.A. (2015). The

clinical spectrum of autoimmune congenital heart block. Nat. Rev. Rheumatol. *11*, 301–312.

Brito-Zerón, P., Baldini, C., Bootsma, H., Bowman, S.J., Jonsson, R., Mariette, X., Sivils, K.,

Theander, E., Tzioufas, A., and Ramos-Casals, M. (2016). Sjögren syndrome. Nat. Rev. Dis. Prim.

2, 16047.

Bryc, K., Velez, C., Karafet, T., Moreno-Estrada, A., Reynolds, A., Auton, A., Hammer, M.,

Bustamante, C.D., and Ostrer, H. (2010). Colloquium paper: genome-wide patterns of

population structure and admixture among Hispanic/Latino populations. Proc. Natl. Acad. Sci.

U. S. A. *107 Suppl 2*, 8954–8961.

Bryc, K., Durand, E.Y., Macpherson, J.M., Reich, D., and Mountain, J.L. (2015). The genetic

ancestry of African Americans, Latinos, and European Americans across the United States. Am.

J. Hum. Genet. *96*, 37–53.

Buyon, J.P., Hiebert, R., Copel, J., Craft, J., Friedman, D., Katholi, M., Lee, L.A., Provost, T.T.,

Reichlin, M., Rider, L., et al. (1998). Autoimmune-associated congenital heart block:

demographics, mortality, morbidity and recurrence rates obtained from a national neonatal

lupus registry. J. Am. Coll. Cardiol. *31*, 1658–1666.

Campbell, H., Carothers, A.D., Rudan, I., Hayward, C., Biloglav, Z., Barac, L., Pericic, M.,

Janicijevic, B., Smolej-Narancic, N., Polasek, O., et al. (2007). Effects of genome-wide

heterozygosity on a range of biomedically relevant human quantitative traits. Hum. Mol. Genet.

16, 233–241.

Campbell, H., Rudan, I., Bittles, A.H., and Wright, A.F. (2009). Human population structure,

genome autozygosity and human health. Genome Med. *1*, 91.

Chapman, J.R., Nakagawa, S., Coltman, D.W., Slate, J., and Sheldon, B.C. (2009). A quantitative

review of heterozygosity-fitness correlations in animal populations. Mol. Ecol. *18*, 2746–2765.

David, P. (1998). Heterozygosity–fitness correlations: new perspectives on old problems.

Heredity. *80*, 531–537.

Denny, J.C., Ritchie, M.D., Basford, M.A., Pulley, J.M., Bastarache, L., Brown-Gentry, K., Wang,

D., Masys, D.R., Roden, D.M., and Crawford, D.C. (2010). PheWAS: demonstrating the feasibility

of a phenome-wide scan to discover gene-disease associations. Bioinformatics *26*, 1205–1210.

Denny, J.C., Bastarache, L., Ritchie, M.D., Carroll, R.J., Zink, R., Mosley, J.D., Field, J.R., Pulley,

J.M., Ramirez, A.H., Bowton, E., et al. (2013). Systematic comparison of phenome-wide

association study of electronic medical record data and genome-wide association study data.

Nat. Biotechnol. *31*, 1102–1111.

Dickerson, G.E. (1973). Inbreeding and heterosis in animals. J. Anim. Sci. *1973*, 54–77.

Dumitrescu, L., Ritchie, M.D., Brown-Gentry, K., Pulley, J.M., Basford, M., Denny, J.C.,

Oksenberg, J.R., Roden, D.M., Haines, J.L., and Crawford, D.C. (2010). Assessing the accuracy of

observer-reported ancestry in a biorepository linked to electronic medical records. Genet. Med.

12, 648–650.

Durbin, R.M., Altshuler, D.L., Durbin, R.M., Abecasis, G.R., Bentley, D.R., Chakravarti, A., Clark,

A.G., Collins, F.S., De La Vega, F.M., Donnelly, P., et al. (2010). A map of human genome

variation from population-scale sequencing. Nature *467*, 1061–1073.

Falush, D., Stephens, M., and Pritchard, J.K. Inference of Population Structure Using Multilocus

Genotype Data: Linked Loci and Correlated Allele Frequencies.

Franceschini, N., Fox, E., Zhang, Z., Edwards, T.L., Nalls, M.A., Sung, Y.J., Tayo, B.O., Sun, Y. V.,

Gottesman, O., Adeyemo, A., et al. (2013). Genome-wide association Analysis of blood-pressure

traits in African-Ancestry Individuals reveals common associated genes in African and Non-

African populations. Am. J. Hum. Genet. *93*, 545–554.

Frankham, R., Ballou, J.D., Eldridge, M.D.B., Lacy, R.C., Ralls, K., Dudash, M.R., and Fenster, C.B.

(2011). Predicting the probability of outbreeding depression. Conserv. Biol. *25*, 465–475.

Galdas, P.M., Cheater, F., and Marshall, P. (2005). Men and health help-seeking behaviour:

literature review. J. Adv. Nurs. *49*, 616–623.

Han, E., Carbonetto, P., Curtis, R.E., Wang, Y., Granka, J.M., Byrnes, J., Noto, K., Kermany, A.R.,

Myres, N.M., Barber, M.J., et al. (2017). Clustering of 770,000 genomes reveals post-colonial

population structure of North America. Nat. Commun. *8*, 14238.

Head, M.L., Kahn, A.T., Henshaw, J.M., Keogh, J.S., and Jennions, M.D. (2017). Sexual selection

on male body size, genital length and heterozygosity: Consistency across habitats and social

settings. J. Anim. Ecol. *86*, 1458–1468.

Kato, N., Loh, M., Takeuchi, F., Verweij, N., Wang, X., Zhang, W., Kelly, T.N., Saleheen, D., Lehne,

B., Leach, I.M., et al. (2015). Trans-ancestry genome-wide association study identifies 12 genetic

loci influencing blood pressure and implicates a role for DNA methylation. Nat. Genet. *47*,

1282–1293.

Keller, L.F., and Waller, D.M. (2002). Inbreeding effects in wild populations. Trends Ecol. Evol.

17, 230–241.

Koch, E., Ristroph, M., and Kirkpatrick, M. (2013). Long range linkage disequilibrium across the

human genome. PLoS One *8*, e80754.

Kodaman, N., Aldrich, M.C., Smith, J.R., Signorello, L.B., Bradley, K., Breyer, J., Cohen, S.S., Long,

J., Cai, Q., Giles, J., et al. (2013). A small number of candidate gene SNPs reveal continental

ancestry in african americans. Ann. Hum. Genet. *77*, 56–66.

Lamkey, and Edwards, J. (1999). Quantitative genetics of heterosis. Genet. Explor. Heterosis

Crop. Am. Soc. Agron. Madison.

Lofquist, D., Lugaila, T., O'Connell, M., and Feliz, S. (2012). Household and families: 2010. 2010

census briefs.

Lynch, M. (1991). The genetic interpretation of inbreeding depression and outbreeding

depression. Evolution (N. Y). *45*, 622–629.

Maglo, K.N., Mersha, T.B., and Martin, L.J. (2016). Population genomics and the statistical

values of race: an interdisciplinary perspective on the biological classification of human

populations and implications for clinical genetic epidemiological research. Front. Genet. *7*, 22.

Mansfield, A., Addis, M., Health, J.M.-J. of M., and 2003, U. (2003). "Why won't he go to the

doctor?": The psychology of men's help seeking. Int. J. Mens. Health *2*, 93–109.

Mittermann, I., Aichberger, K.J., Bünder, R., Mothes, N., Renz, H., and Valenta, R. (2004).

Autoimmunity and atopic dermatitis. Curr. Opin. Allergy Clin. Immunol. *4*, 367–371.

Moreno-Estrada, A., Gignoux, C.R., Fernandez-Lopez, J.C., Zakharia, F., Sikora, M., Contreras, A.

V., Acuna-Alonzo, V., Sandoval, K., Eng, C., Romero-Hidalgo, S., et al. (2014). The genetics of

Mexico recapitulates Native American substructure and affects biomedical traits. Science (80-.

). *344*, 1280–1285.

Nalls, M.A., Simon-Sanchez, J., Gibbs, J.R., Paisan-Ruiz, C., Bras, J.T., Tanaka, T., Matarin, M.,

Scholz, S., Weitz, C., Harris, T.B., et al. (2009). Measures of autozygosity in decline:

globalization, urbanization, and its implications for medical genetics. PLoS Genet. *5*, e1000415.

Nei, M., and Li, W.-H. (1973). Linkage disequilibrium in subdivided populations. Genetics *75*,

213–219.

Pemberton, T.J., and Szpiech, Z.A. (2018). Relationship between deleterious variation, genomic

autozygosity, and disease risk: insights from The 1000 Genomes Project. Am. J. Hum. Genet.

102, 658–675.

Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., and Reich, D. (2006).

Principal components analysis corrects for stratification in genome-wide association studies.

Nat. Genet. *38*, 904–909.

Pritchard, J.K., Stephens, M., and Donnelly, P. (2000). Inference of Population Structure Using

Multilocus Genotype Data.

Reich, D., and Patterson, N. (2005). Will admixture mapping work to find disease genes? Philos.

Trans. R. Soc. Lond. B. Biol. Sci. *360*, 1605–1607.

Roden, D., Pulley, J., Basford, M., Bernard, G., Clayton, E., Balser, J., and Masys, D. (2008).

Development of a large-scale de-identified DNA biobank to enable personalized medicine. Clin.

Pharmacol. Ther. *84*, 362–369.

Rosenberg, N.A. (2011). A population-genetic perspective on the similarities and differences

among worldwide human populations. Hum. Biol. *83*, 659–684.

Rudan, I., Carothers, A.D., Polasek, O., Hayward, C., Vitart, V., Biloglav, Z., Kolcic, I., Zgaga, L.,

Ivankovic, D., Vorko-Jovic, A., et al. (2008). Quantifying the increase in average human

heterozygosity due to urbanisation. Eur. J. Hum. Genet. *16*, 1097–1102.

Ruggles, S. (2014). Big microdata for population research. Demography *51*, 287–297.

Ruiz-Narvá Ez, E.A., Rosenberg, L., Wise, L.A., Reich, D., and Palmer, J.R. (2011). Practice of

epidemiology validation of a small set of ancestral informative markers for control of

population admixture in african americans. Am. J. Epidemiol. *173*.

Shull, G.H. (1948). What Is "Heterosis"? Genetics *33*, 439–446.

Spergel, J.M. (2010). From atopic dermatitis to asthma: the atopic march. Ann. Allergy, Asthma

Immunol. *105*, 99–106.

Stefanski, L.A., and Boos, D.D. (2002). The Calculus of M-Estimation. The American Statistician.

56, 29-38.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green,

J., Landray, M., et al. (2015). UK Biobank: an open access resource for identifying the causes of

a wide range of complex diseases of middle and old age. PLOS Med. *12*, e1001779.

Szulkin, M., Bierne, N., and David, P. (2010). Heterozygosity-fitness correlations: a time for

reappraisal. Evolution (N. Y). *64*, 1202–1217.

Tedeschi, A., and Asero, R. (2008). Asthma and autoimmunity: a complex but intriguing relation.

Expert Rev. Clin. Immunol. *4*, 767–776.

Tishkoff, S.A., Reed, F.A., Friedlaender, F.R., Ehret, C., Ranciaro, A., Froment, A., Hirbo, J.B.,

Awomoyi, A.A., Bodo, J.-M., Doumbo, O., et al. (2009). The genetic structure and history of

Africans and African Americans. Science *324*, 1035–1044.

Verdu, P., and Rosenberg, N.A. (2011). A general mechanistic model for admixture histories of

hybrid populations. Genetics *189*, 1413–1426.

Villuendas, R., Olivé, A., Juncà, G., Salvador, I., Martínez-Morillo, M., Santos-Pardo, I.,

Pereferrer, D., Zamora, E., and Bayes-Genis, A. (2014). Autoimmunity and Atrioventricular Block

of Unknown Etiology in Adults. J. Am. Coll. Cardiol. *63*, 1335–1336.

Wang, C., Xu, Z., Jin, G., Hu, Z., Dai, J., Ma, H., Jiang, Y., Hu, L., Chu, M., Cao, S., et al. (2013).

Genome-wide analysis of runs of homozygosity identifies new susceptibility regions of lung

cancer in Han Chinese. J. Biomed. Res. *27*, 208–214.

Wood, S.N. (2010). Fast stable REML and ML estimation of semiparametric GLMs. J. R. Stat. Soc.

Ser. B Statistical Methodol. *73*, 3–36.