Gestational pertussis vaccination and the infant's cellular immune response against

whole-cell pertussis vaccine in the first year of life

Carolina Argondizo-Correia[a,b,*]; Lourdes Rehder de Andrade Vaz-de-Lima[b], Elaine

Uchima Uehara[c], Eder Gatti Fernandes[d], Helena Keico Sato[d], Ana Paula Sayuri Sato[e],

Expedito José de Albuquerque Luna[a], Euclides Ayres de Castilho[b], Cyro Alves de

Brito[a,b,#]

[a]Institute of Tropical Medicine, University of São Paulo, São Paulo, Brazil

[b]Immunology Centre, Adolfo Lutz Institute, São Paulo, SP, Brazil

[c]Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil

[d]CVE/CCD/SES, São Paulo, Brazil

[e]School of Public Health, University of São Paulo, São Paulo, Brazil

Running Head: Infant's pertussis cellular immune response

#Address correspondesse to Cyro Alves de Brito, cbrito@usp.br

*Present address: Carolina Argondizo Correia, Laboratory of Medical Investigation 60,

Medical School, University of São Paulo.

Word count (abstract): 249 words

Word count (text): 2735 words

Abstract

Pertussis resurgence worldwide calls for new prevention strategies, as the recently incorporated vaccine booster dose during pregnancy, whose aim is to protect newborns from infection. In Brazil, maternal Tdap vaccination is recommended since 2014, and we reported that this strategy promotes high transplacental transfer of anti-PT IgG and it is effective in protecting infants early in life. Young children are the most susceptible group and with higher mortality rates, however, it is not well known whether the elicited anti-pertussis maternal antibodies could influence in the children's immune responses further in life, especially after their own vaccination against pertussis. Considering this scenario, we conducted a study with children born to mothers who either received or not the booster dose during pregnancy, after their primary pertussis vaccination, in order to investigate the first impact of maternal immunisation on the response to infant immunisation regarding the cellular immune response, while comparing with data from the literature. As transfer of maternal antibodies could result in attenuation of the immune response to vaccination in infants, this study performed to determine whether higher levels of maternal antibodies could influence in the immune response of infants to the whole-pertussis vaccination series. Results showed no difference in cytokine production between groups, a first suggestion that maternal vaccination may not interfere with recognition and cellular response generation to vaccination. This data, together with humoral immunity and epidemiological studies, is important for the implementation of maternal immunisation strategies nationwide and will contribute to assure public policies regarding vaccination schemes.

Importance: Pertussis, or whopping cough, is a respiratory infectious disease caused by a bacterial agent, resulting in violent coughs and possibly death in vulnerable groups,

such as young children and neonates. It is known that pregnant mothers transfer antibodies to their developing foetuses for protection in early life, however anti-pertussis antibodies are not highly detected in young children. Thus, a pertussis maternal vaccination was implemented to increase maternal anti-pertussis antibodies levels in pregnant women and therefore the transference to the foetus. However, maternal antibodies can also interfere in the child immune response in the first months of life. The significance of our research is in analysing the cellular immune response of children born from pertussis-vaccinated mothers, which will give a first glimpse on how maternal antibodies could modulate the child's response to pertussis in early life.

Introduction

Pertussis is a contagious respiratory disease caused by the *Bordetella pertussis* bacteria [1]. Despite high vaccination coverage, it remains an important public health problem, re-emerging in several countries every several years [2,3]. It has a high rate of morbimortality in young children, and estimates from the World Health Organization suggest that ~50 million pertussis cases and 300,000 deaths occurred annually, mostly in children under five years of age [4]. In Brazil, between 2010 and 2014, over 22,000 people were infected [5], most cases in children younger than 1 year of age [5,6].

The current childhood vaccination schedule includes three doses of the diphtheria, tetanus and whole cell pertussis vaccine, combined with *Haemophilus influenza* b and hepatitis B (DTwP-Hib-HBV) in the first year of life, at 8, 16 and 24 weeks of age, leaving neonates without specific protection. So, a dose of tetanus, reduced diphtheria and acellular pertussis vaccine (Tdap) during pregnancy was proposed to address this immunity gap in young infants and promote a higher transplacental transfer of maternal

antipertussis antibodies (MatAb) to the foetus, resulting in improved protection during the neonatal period and until their own vaccination scheme is completed [7].

Since 2012, this has been recommended in the United States, United Kingdom and Australia [8], and since November 2014 in Brazil [9]. Recently, we have shown Tdap maternal vaccination promotes high titers of anti-PT IgG in newborn [10], also this strategy showed a vaccine effectiveness of 82.6% in protecting infants aged <8 weeks from pertussis in Brazil [11]. However, until now, little is known whether it may affect the subsequent childhood vaccination, which uses whole-cell pertussis (wP) [12], while other countries recommend the acellular pertussis (aP) version [13]. There is also a concern that in addition to promoting protection to young infants, high MatAb titres could interfere or attenuate the immune response to the primary childhood pertussis vaccination series [14–16].

The humoral response is not the only responsible for the protection against infection, as well as there are no correlates of protection for serological levels of antipertussis antibodies [17]. Several studies show that the cellular immune response is required for effective clearance of infection from the respiratory tract and disease prevention, through effector mechanisms mediated by IFN-γ and IL-2, which are mostly produced by the T helper lymphocyte 1 (Th1). This cell type is described as having a more inflammatory profile, opposed to Th2 cells, characterized by the production of IL-4, IL-5 and IL-13. Some authors also relate the presence of IL-17 in the mechanism against *B.pertussis in vivo*, a cytokine produced by Th17 cells [18–23].

In light of this information, we sought to analyse the production of effector cytokines that could imply the activation of the infants' cellular immune response against *B. pertussis* antigen *in vitro*, in the context of the presence of maternal anti-pertussis antibodies.

Material and Methods

Study Design and Participants

This study included 43 children around 7 months of age, born either to mothers vaccinated

during pregnancy with a Tdap boost vaccine during the third trimester of pregnancy

(n=33) or mothers who did not get vaccinated (n=10). This cohort is derived from a larger

cohort described elsewhere [10]. Mothers were followed up after parturition until their

children received the primary pertussis vaccination series, composed by three DTwP-Hib-

HBV doses (produced by Bio-Manguinhos/Fiocruz), at approximately 2, 4 and 6 months

of age. The protocol was approved by the Ethics Committee from the Adolfo Lutz

Institute (CAAE: 37581114.0.0000.0059).

Sample Processing and In Vitro Stimulation

5 mL of heparinized venous peripheral blood samples were collected by nurses from

young children around one month after the third DTwP dose.

Samples were diluted in culture medium (RPMI 1640 [Gibco, Massachusetts/USA]

supplemented with 2 g of $NaHCO_3$, 10 mL of nonessential amino acids [Gibco], 40 mg

of gentamycin and 10 mL of 200 mM L-glutamine [Sigma, Missouri/USA]). Peripheral

blood mononuclear cells (PBMC) were isolated using a Ficoll-Paque Plus 1440 gradient

(GE Healthcare, Uppsala/Sweden), according to the manufacturer's instructions. Cells

were washed and resuspended in culture medium, counted and rested overnight (~16 h,

$1x10^6$ cells/mL) in 48-well flat-bottom plates at 37ºC in a 5% CO_2 incubator. On the

following day, cells were stimulated with 2 µg/mL phytohemagglutinin (PHA) for 48 h

[Sigma, Missouri/USA] or 5 µg/mL of inactivated pertussis toxoid (PT) for 120 h [kindly

donated by GSK]. After culture, cells and supernatant were collected and stored at -80ºC

for later assays. Cell pellets were stabilized using RNAlater (Sigma) and phosphate

buffered saline (21.02 g of Na_2HPO_4, 7.16 g of $NaH_2PO_4.H_2O$, 164.64 g of NaCl and

reverse-osmosis H_2O for 1 L) until total RNA extraction.

Cytokine Detection

Frozen supernatants were thawed at room temperature, and cytokines were measured

using both *Cytometric Bead Array* Th1/Th2/Th17 kit (BD Biosciences, New Jersey/USA)

and LEGENDplex *Human Th Cytokine Panel* (13-*plex*) kit (BioLegend, California/USA),

according to the manufacturers' instructions, in order to compare results and analyse a

larger range of cytokines. Both curve and samples were read in a BD LSRFortessa flow

cytometer (BD Biosciences). Minimum detection limits were 4.88 pg/mL (BD) and 2.4

pg/mL (BioLegend). Corresponding analyses were performed with FCAP Array software

v3.0 (BD Biosciences) and LEGENDplex Data Analysis software (BioLegend).

RNA Extraction and Quantitative Polymerase Chain Reaction (qPCR)

RNA extraction was performed using Qiagen's RNeasy Mini kit (Qiagen,

Hilden/Germany), following the manufacturer's instructions. To remove any traces of

DNA, a DNA digestion step was performed using an RNAqueous Mini Kit (Ambion,

Massachusetts/USA). RNA purity and quantity were evaluated by spectrophotometry in

a Nanodrop (Thermo Scientific, Massachusetts/USA). RNA was reverse transcribed to

cDNA using an iScript cDNA Synthesis Kit (Bio-Rad, California/USA) according to the

manufacturer's instructions.

In an attempt to identify different cell types, the chosen genes were the transcription factor

RORC2 and the cytokine gene *IL17A*, representing Th17 cells, the transcription factor

GATA3 and the cytokine gene *IL4* representing Th2 cells, and the cytokines *IFNG* for

Th1 and *IL10* for regulatory response [24]. qPCR assays were run for each sample

containing cDNA as a template, specific forward (5'-3') and reverse (5'-3') primers,

respectively:

RORC2: TGGAAGTGGTGCTGGTTAGGA/AAGGCTCGGAACAGCTCCAT;

GATA3: AGATGGCACGGGACACTACCT/TAATTCGGGTTCGCTTCCG;

IFNG:

GTTTTGGGTTCTCTTGGCTGTTA/AAAAGAGTTCCATTATGCGCTACATC;

IL17A: GACTCCTGGGAAGACCTCATTGG/CTTGTCCTCAGAATTTGGGCATCC;

IL4: ACAGCCTCACAGAGCAGAAGACT/TGTTCTTGGAGGCAGCAAAGA;

IL10: CAGGGCACCCAGTCTGAGAA/CACATGCGCCTTGATGTCTG;

GAPDH: GAAGGTGAAGGTCGGAGT/GAAGATGGTGATGGGATTTCCA;

and SYBR Green PCR Master Mix (Life Technologies, Massachusetts/USA) on an

Applied Biosystems 7500 Real-Time PCR System (Applied Biosystems,

California/USA). *GAPDH* gene expression was used as a control to normalize the data.

Results were represented as fold change given by the equation described by Livak and

Schmittgen [25].

Statistical Analysis

Descriptive and inferential statistical analyses were performed with GraphPad Prism

software (version 5.0). Nonparametric Mann-Whitney, Kruskal-Wallis, Friedman, Dunn

and Wilcoxon tests were used for inter- and intragroup analyses, and Spearman's

correlation coefficient was used for demographic and experimental data. We used $\alpha=5\%$,

95% confidence interval, level of significance=5% and 2-sided tests.

Results

The study population consisted of young children born from women who received (VC)

or not (NVC) a Tdap boost dose during pregnancy (Table 1). Samples were randomly

chosen from a larger cohort described by Vaz-de-Lima et al. [10]. The groups presented

a similar and representative demographic profile when compared to the larger cohort.

In order to evaluate whether Tdap vaccination during pregnancy could induce

modification in the cellular immune response of children, we evaluated a few related

genes and cytokines that could represent the main cells in the cellular immune response,

herein described as Th1, Th2 and Th17.

Transcriptional profile (Figure 1) showed no difference between groups, though the

median levels of *GATA3*, *IL10* and *IFNG* were higher in children born to nonvaccinated

mothers. When comparing the normalized expression of both basal and stimulated

condition between groups, NVCs had lower levels for the basal condition concerning

GATA3 (p=0.0279) and *IFNG* (p=0.0290) expression. In the stimulated condition, the

differences disappear, so the higher median level in the fold change analysis can indicate

that in order to achieve the same expression upon PT stimulation, NVCs must have a

greater increase in the expression of *GATA3* and *IFNG* due to the lower basal expression.

To verify whether the results found in gene expression reflected the cytokine production,

cytokines from Th1, Th2 and Th17 profiles were quantified via flow cytometry (Table

2). PHA stimulation was used as a positive control in a few samples for the main cytokines

for the 3 Th profiles, due to samples and kit availability.

In Table 2 it shown the total quantification from detectable all cytokines but IL-9 and IL-

17F, as those values were undetectable.

Overall, TNF, IFN-γ, IL-13 and IL-6 were the most produced cytokines in both groups,

with IL-6 being more produced in PT-stimulated cells derived from children born to

vaccinated mothers (VC). In contrast, IL-2 was higher is children whose mothers were

not vaccinated (NVC).

As both groups showed simultaneous IFN-γ and IL-13 production, which could indicate

a mixed Th1/Th2 response or a good response by individuals to all cytokines, we

evaluated the correlation between cytokines from different profiles, looking for a

predominant response in all individuals or whether some individuals were indistinctly

good responders to all T helper cell profiles (data not shown). However, no correlation

was found between groups or cytokines. Individual responses were different from each

other, making it challenging to find a correlation in a small group.

Regarding infants with higher cytokine levels, we analysed whether factors such as Tdap

dose, gestational age, route of birth, infant's sex, period of DTwP vaccination and

exclusive breastfeeding could influence the response deviation (data not shown), but no

significant correlations were found.

Discussion

This study was the first to show the cellular immune response in older children, after the

primary pertussis vaccination series, born to Tdap-vaccinated mothers in Brazil. Despite

the use of Tdap during pregnancy was implemented in Brazil and other countries in the

last few years and substantial data regarding antibody transfer and its influence on

children's humoral response to vaccination [10,26–28], little is known whether this

strategy could affect the infant's cellular immune response.

A few studies have been published recently about the influence of maternal antibodies in

the neonatal humoral response, as the study made by Ibrahim et al. [29]. Women that did

not receive Tdap during pregnancy but had detectable pertussis antibody levels showed

that maternal antibody titres at delivery did not blunt infant vaccine response among

children that receive at least 2 wP doses, and the authors observed that infant antibody

titres increased with increasing maternal antibody levels against all pertussis antigens. A

Brazilian study found that Tdap maternal immunization yielded significantly higher anti-

PT IgG levels in vaccinated mothers and their infants compared to their unvaccinated

counterparts and a strong positive correlation between the anti-PT IgG levels in maternal

and cord blood at delivery, but did not look for antibodies against other pertussis antigens

[10].

Lima et al. [28] showed the same correlation regarding antibodies against other bacterial

antigens, and showed for the first time the analysis of the cellular immune response in

whole cord blood cells, showing IFN-γ, IL-6, IL-10 and TNF production by neonatal cells

upon stimulation with the whole inactivated *B. pertussis*, and low levels of IL-2, IL-4 and

IL-17A.

In our study, we complemented the neonatal data by evaluating the cytokine productions

in older children, with the complete primary series of vaccination against pertussis.

The cytokines analysed were TNF-α and IL-2, which belong to a Th1 profile, in

association with IFN-γ; IL-9 by Th9 cells, IL-4, IL-5 and IL-13, which contribute to the

Th2 response profile; IL-10 is secreted by T regulatory cells and IL-17A, IL-17F, IL-21

and Il-22 secreted by Th17 cells. IL-6 is an inflammatory cytokine and is important in the

differentiation of Th17 cells [24].

We used purified PBMC stimulated with PT, for five days, as described in the literature

[30]. By using PBMCs it easier to access lymphocytes present in the samples and PT is

the only specific antigen of *B. pertussis* [31], so our group decided to first run the assays

with this antigen to have a first glimpse of what could have been happening, as well as,

with PT, we analysed the adaptive response directly related to this stimulation. We could

not determine the purity of PT used in the assays, but even with other bacterial antigens

present we could see cytokine production upon stimulus.

Despite using samples from older children, our study agrees with Lima et al. [28] in the

production of IFN-γ, IL-6 and TNF, which we found elevated in infants' samples, but not

IL-10. We also found low levels of IL-4 and IL-17A, but IL-13, which is an indicative of

the Th2 response and was not available in the assay the authors performed, was found

elevated in our samples. In all cytokines evaluated, there was one or more individuals that

we did not detect any production, indicating a strong variability amongst human samples.

As our both groups presented similar characteristics, the slight difference in gene

expression could suggest modulation by vaccination. Even so, samples were limited, and

the mRNA analysed was just to have a glimpse of what cells populations may have been

present upon stimulus. Even though cytokine mRNA are produced earlier under these

experimental conditions, the presence of the genes can show a tendency to a certain

profile. This is seen in both the cytokine measurement and gene expression and could be

complemented by the analysis of *TBX21*, the gene responsible for Th1 cell polarization

[32].

When attempting to find correlation between different cytokines production, we found

that in both groups every individual produced high levels of at least one cytokine. In the

NVC group, a correlation between IFN-γ and IL-10 could point to a balance between

activation and regulation, while in the VC group the same correlation was found, as well

as between and IFN-γ and IL-13 (data not shown). This could indicate a mixed Th1/Th2

profile, with IL-10 possibly balancing the production of IFN-γ and the activation of Th1

cells. In both groups, cytokine levels were variable between individuals, and thus, it is

not possible to determine a predominant response pattern.

According to Li *et al.* [33], Th2 cells do not prevent Th1 polarization but induce their

apoptosis. In addition, they also observed that IL-10 could promote Th1 cell anergy

through negative regulation of accessory molecules or IL-12 in antigen-presenting cells.

Even inducing protection, maternal antibodies can also interfere with the child's vaccine

response, reducing the vaccine efficiency [34], leading to a decrease in the child's

antibody production via immune complex formation, antigen elimination via

phagocytosis or epitope masking [35]. However, immune cellular response appears to be

unaffected by these mechanisms [26]. As in Brazil children are vaccinated with DTwP,

unlike many countries, we aimed to analyse the cellular immune response of vaccinated

children born to either vaccinated or non vaccinated women, to confirm whether the

cellular response would be intact. No difference was found between the two groups.

Regardless of the different sample sizes and individual variabilities, both common factors

in human vaccine studies [36–38], we could see IFN-γ and TNF-α production, indicative

of the Th1 response, while only IL-13 from the Th2 profile was detected.

Overall, this work is the first step towards a complete scenario of maternal antibodies

interference in cellular immune responses. Most pertussis studies describe humoral

responses, before [39] and after the implementation of maternal vaccination [40,41],

evaluating immunoglobulin levels produced by mothers and transferred to newborns, in

countries using aP vaccines in children, unlike Brazil. A paper in Argentina describes the

use of DTwP in children in the country; however, the study shows only the influence of

maternal antibodies on the humoral responses of children before DTwP vaccination

[41].Despite the limitations, we evidenced cellular immune response in the context of

pertussis vaccination, which is shown to be important for protection [42]. Our results

show no difference in cytokine production against a bacterial antigen between groups,

regardless of maternal vaccination, and this could set the path to other works that will

support public health policies regarding vaccination in pregnancy. This knowledge is

essential to the enhancement of vaccine protocols in pregnant women and as a basis for

physicians and healthcare managers to recommend this strategy.

Conclusion

This work shows cytokine production against a pertussis antigen in the context of

maternal antibodies, and data could be strengthened by verifying these observations with

more than one pertussis antigens. Also, there are ongoing studies that measured antibody

responses to infant immunisation, in order to complement data and promote discussion

about a potential difference in the impact of maternal immunisation of infant cellular and

humoral responses.

Acknowledgments: The authors would like to thank the teams at Hospital Maternidade

Interlagos and Hospital Maternidade Leonor Mendes de Barros for recruitment and their

help with blood specimen collection, Laboratory of Dermatology and

Immunodeficiencies (LIM-56, Medical School of the University of São Paulo) and

Immunology Centre from the Adolfo Lutz Institute for the infrastructure and support, the

funding sources and all participants of the study. This study was supported by funding

provided by the Special Health Fund for Mass Immunization and Disease Control

(FESIMA) and São Paulo Research Foundation (FAPESP), grants #2015/19726-7 and

#2015/16157-1. The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication. The authors also declare

no conflicts of interest to disclose.

References

[1]   World Health Organization. Pertussis vaccines: WHO position paper - August
2015. Weely Epidemiol Rec 2015;90:433–60. doi:10.1371/jour.

[2]   Pan American Health Organization. Epidemiological alert. Pertussis (Whooping
Cough). PAHO Bull 2012:1–2.

[3]   Luz PM, Codeço CT, Werneck GL. A reemergência da coqueluche em países
desenvolvidos: um problema também para o Brasil? Cad Saúde Pública
2003;19:1209–13.

[4]   World Health Organization. WHO–recommended standards for surveillance of
selected vaccine-preventable diseases. Geneva: 2003.

[5]   Secretaria de Vigilância em Saúde, Ministério da Saúde. Coqueluche no Brasil:
análise da situação epidemiológica de 2010 a 2014. Bol Epidemiológico
2015;46:1–8.

[6]   Rocha EL, Leite D, Camargo CH, Martins LM, Silva RSN, Martins VP, et al. The
characterization of Bordetella pertussis strains isolated in the Central-Western
region of Brazil suggests the selection of a specific genetic profile during 2012–
2014   outbreaks.   Epidemiol   Infect   2017;145:1392–7.
doi:10.1017/S0950268816003332.

[7]   Locht C, Mielcarek N. New pertussis vaccination approaches: En route to protect
newborns? FEMS Immunol Med Microbiol 2012;66:121–33. doi:10.1111/j.1574-
695X.2012.00988.x.

[8]   Centers for Disease Control and Prevention. Updated Recommendations for Use
of Tetanus Toxoid, Reduced Diphtheria Toxoid, and Acellular Pertussis Vaccine
(Tdap) in Pregnant Women — Advisory Committee on Immunization Practices
(ACIP), 2012. Morb Mortal Wkly Rep 2013.

[9]   Ministério da Saúde. Nota informativa sobre mudanças no calendário nacional de

345  vacinação para o ano de 2017. Brasília: 2016.

346 [10] Vaz-de-Lima LRA, Sato HK, Fernandes EG, Sato APS, Pawloski LC, Tondella

347  ML, et al. Association between the timing of maternal vaccination and newborns'

348  anti-pertussis toxin antibody levels. Vaccine 2019.

349  doi:10.1016/j.vaccine.2019.04.079.

350 [11] Fernandes EG, Sato APS, Vaz-de-Lima LRA, Rodrigues M, Leite D, de Brito CA,

351  et al. The effectiveness of maternal pertussis vaccination in protecting newborn

352  infants in Brazil: A case-control study. Vaccine 2019.

353  doi:10.1016/j.vaccine.2019.03.049.

354 [12] Domingues CMAS, Teixeira AM da S. Coberturas vacinais e doenças

355  imunopreveníveis no Brasil no período 1982-2012: avanços e desafios do

356  Programa Nacional de Imunizações. Epidemiol e Serviços Saúde 2013;22:9–27.

357  doi:10.5123/S1679-49742013000100002.

358 [13] Noble GRGR, Bernier RHRH, Esber ECEC, Hardegree MCC, Hinman ARAR,

359  Klein D, et al. Acellular and whole-cell pertussis vaccines in Japan. Report of a

360  visit by US scientists. JAMA 1987;257:1351–6.

361  doi:10.1001/jama.1987.03390100089032.

362 [14] Osborn JJ, Dancis J, Julia JF. Studies of the immunology of the newborn infant.

363  Pediatrics 1952;10:339–45.

364 [15] Premenko-Lanier M, Hodge G, Rota P, Tamin A, Bellini W, McChesney M.

365  Maternal antibody inhibits both cellular and humoral immunity in response to

366  measles vaccination at birth. Virology 2006;350:429–32.

367  doi:10.1016/j.virol.2006.02.029.

368 [16] Gall SA, Myers J, Pichichero M. Maternal immunization with

369  tetanusdiphtheriapertussis vaccine: Effect on maternal and neonatal serum

antibody levels. Am J Obstet Gynecol 2011;204:334.e1-334.e5.
doi:10.1016/j.ajog.2010.11.024.

[17] Cherry JD, Gornbein J, Heininger U, Stehr K. A search for serologic correlates of
immunity to Bordetella pertussis cough illnesses. Vaccine 1998;16:1901–6.
doi:10.1016/S0264-410X(98)00226-6.

[18] Mahon BBP, Sheahan BJ, Griffin F, Murphy G, Mills KHG. of Mice with Targeted
Disruptions of Interferon- □ Receptor 1997;186.

[19] Ross PJ, Sutton CE, Higgins S, Allen AC, Walsh K, Misiak A, et al. Relative
contribution of Th1 and Th17 cells in adaptive immunity to Bordetella pertussis:
towards the rational design of an improved acellular pertussis vaccine. PLoS
Pathog 2013;9:e1003264. doi:10.1371/journal.ppat.1003264.

[20] Barbic J, Leef MF, Burns DL, Shahin RD. Role of gamma interferon in natural
clearance of Bordetella pertussis infection. Infect Immun 1997;65:4904–8.
doi:10.1371/journal.ppat.1003264.

[21] Ryan M, Murphy G, Gothefors L, Nilsson L, Storsaeter J, Mills KH. Bordetella
pertussis respiratory infection in children is associated with preferential activation
of type 1 T helper cells. J Infect Dis 1997;175:1246–50. doi:10.1086/593682.

[22] Higgs R, Higgins SC, Ross PJ, Mills KHG. Immunity to the respiratory pathogen
Bordetella pertussis. Mucosal Immunol 2012;5:485–500. doi:10.1038/mi.2012.54.

[23] Mills KHG. Immunity to Bordetella pertussis. Microbes Infect 2001;3:655–77.

[24] Raphael I, Nalawade S, Eagar TN, Forsthuber TG. T cell subsets and their
signature cytokines in autoimmune and inflammatory diseases. Cytokine
2015;74:5–17. doi:10.1016/j.cyto.2014.09.011.

[25] Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. Methods 2001;25:402–8.

doi:10.1006/meth.2001.1262.

[26]   Siegrist C-A, Barrios C, Martinez X, Brandt C, Berney M, Córdova M, et al.

397   Influence of maternal antibodies on vaccine responses: inhibition of antibody but

398   not T cell responses allows successfull early prime-boost strategies in mice. Eur J

399   Immunol 1998;28:4138–48. doi:10.1002/(SICI)1521-4141(199812)28.

400   [27]   Siegrist CA, Córdova M, Brandt C, Barrios C, Berney M, Tougne C, et al.

401   Determinants of infant responses to vaccines in presence of maternal antibodies.

402   Vaccine 1998;16:1409–14. doi:10.1016/S0264-410X(98)00100-5.

403   [28]   Lima L, Molina M da GF, Pereira BS, Nadaf MLA, Nadaf MIV, Takano OA, et

404   al. Acquisition of specific antibodies and their influence on cell-mediated immune

405   response in neonatal cord blood after maternal pertussis vaccination during

406   pregnancy. Vaccine 2019;37:2569–79. doi:10.1016/j.vaccine.2019.03.070.

407   [29]   Ibrahim R, Ali SA, Kazi AM, Rizvi A, Guterman LB, Bednarczyk RA, et al.

408   Impact of maternally derived pertussis antibody titers on infant whole-cell

409   pertussis vaccine response in a low income setting. Vaccine 2018;36:7048–53.

410   doi:10.1016/j.vaccine.2018.09.045.

411   [30]   Palazzo R, Carollo M, Bianco M, Fedele G, Schiavoni I, Pandolfi E, et al.

412   Persistence of T-cell immune response induced by two acellular pertussis vaccines

413   in children five years after primary vaccination. New Microbiol 2016;39:35–47.

414   [31]   Mattoo S, Cherry JD. Molecular pathogenesis, epidemiology, and clinical

415   manifestations of respiratory infections due to Bordetella pertussis and other

416   Bordetella   subspecies.   Clin   Microbiol   Rev   2005;18:326–82.

417   doi:10.1128/CMR.18.2.326-382.2005.

418   [32]   Brummelman J, Raeven RHM, Helm K, Pennings JLA, Metz B, van Eden W, et

419   al. Transcriptome signature for dampened Th2 dominance in acellular pertussis

vaccine-induced CD4+ T cell responses through TLR4 ligation. Sci Rep

2016;6:25064. doi:10.1038/srep25064.

[33] Li L, Lee H-HH, Bell JJ, Gregg RK, Ellis JS, Gessner A, et al. IL-4 utilizes an

alternative receptor to drive apoptosis of Th1 cells and skews neonatal immunity

toward Th2. Immunity 2004;20:429–40. doi:10.1016/S1074-7613(04)00072-X.

[34] Siegrist CA. The Challenges of Vaccine Responses in Early Life: Selected

Examples. J Comp Pathol 2007;137:S4–9. doi:10.1016/j.jcpa.2007.04.004.

[35] Siegrist CA. Mechanisms by which maternal antibodies influence infant vaccine

responses: Review of hypotheses and definition of main determinants. Vaccine

2003;21:3406–12. doi:10.1016/S0264-410X(03)00342-6.

[36] Lee YC, Newport MJ, Goetghebuer T, Siegrist CA, Weiss HA, Pollard AJ, et al.

Influence of genetic and environmental factors on the immunogenicity of Hib

vaccine in Gambian twins. Vaccine 2006;24:5335–40.

doi:10.1016/j.vaccine.2006.04.021.

[37] Newport MJ, Goetghebuer T, Weiss HA, Whittle H, Siegrist C-A, Marchant A, et

al. Genetic regulation of immune responses to vaccines in early life. Genes Immun

2004;5:122–9. doi:10.1038/sj.gene.6364051.

[38] Tan PL, Jacobson RM, Poland GA, Jacobsen SJ, Pankratz VS. Twin studies of

immunogenicity-determining the genetic contribution to vaccine failure. Vaccine

2001;19:2434–9.

[39] Marcellini V, Piano Mortari E, Fedele G, Gesualdo F, Pandolfi E, Midulla F, et al.

Protection against Pertussis in Humans Correlates to Elevated Serum Antibodies

and Memory B Cells. Front Immunol 2017;8:1–11.

doi:10.3389/fimmu.2017.01158.

[40] Maertens K, Caboré RN, Huygen K, Vermeiren S, Hens N, Van Damme P, et al.

Pertussis vaccination during pregnancy in Belgium: Follow-up of infants until 1

month after the fourth infant pertussis vaccination at 15 months of age. Vaccine

2016;34:3613–9. doi:10.1016/j.vaccine.2016.04.066.

[41] Fallo AA, Neyro SE, Manonelles G V., Lara C, Hozbor D, Zintgraff J, et al.

Prevalence of Pertussis Antibodies in Maternal Blood, Cord Serum, and Infants

From Mothers With and Those Without Tdap Booster Vaccination During

Pregnancy in Argentina. J Pediatric Infect Dis Soc 2016:1–7.

doi:10.1093/jpids/piw069.

[42] van Twillert I. Cellular and humoral immunity after infection with B. pertussis The

role of age, antigen and vaccination history. 2017.

470   Table 1. Information about the study groups

		With Tdap	Without Tdap
N (infants)		33	10
Sex (%)	Female	16 (49%)	7 (70%)
	Male	17 (52%)	3 (30%)
Route of birth (%)	Vaginal	14 (42%)	6 (60%)
	C-section	20 (58%)	4 (40%)
Exclusive breastfeeding, median (range), months		6.2 (1.8)	5.7 (2.3)
Mother's age at birth, mean (SD), years		29.9 (7.8)	24.3 (5.2)
Mother's gestational age at vaccination, mean (SD), weeks		29.6 (2.8)	-
Mother's gestational age at birth, mean (SD), weeks		39.1 (1.7)	39.6 (1.2)
Time gap between vaccination and birth, mean (SD), weeks		9.5 (2.8)	-
1st DTwP dose age (minimum-maximum, days)		21-92	62-70
2nd DTwP dose age (minimum-maximum, days)		99-155	122-194
3rd DTwP dose age (minimum-maximum, days)		182-232	183-267
Time gap between 3rd dose and sample collection (minimum-maximum, days)		10-127	7-129

DTwP: diphtheria, tetanus and whole-cell pertussis vaccine; SD: standard deviation;

Tdap: tetanus, diphtheria and acellular pertussis vaccine.

Figure 1: Cytokine expression in PT-stimulated PBMCs from infants born either from

vaccinated (VC) or nonvaccinated (NVC) mothers. (A) Relative gene expression is

similar in children born to Tdap-vaccinated (VC) or nonvaccinated (NVC) mothers when

culturing PMBCs with PT for 120 h. Box represents variation between 25th and 75th

percentiles, and whiskers represent minimum and maximum values. (B) VC presented

higher basal expression of *GATA3* and *IFNG* than NVC, though differences disappear

with PT stimulation for 120 h. Box represents variation between the 25th and 75th

percentiles, and whiskers represent minimum and maximum values. Statistically

significant differences are indicated.

494   Table 2. Cytokine concentration in culture supernatant of peripheral blood mononuclear cells (PBMC) from infants.

Cytokines (pg/ml)	NV				VC			
	n=2	n=2	n=10	n=10	n=4	n=4	n=33	n=28/33
	(-) basal 48h	(+) PHA	(-) basal 120h	PT	(-) basal 48hl	(+) PHA	(-) basal 120h	PT
IL-5	*	*	0(0-0)	0(0-0)	*	*	0(0-0)	3,45(0-96,54)
IL-13	*	*	0(0-0)	775,03(0-3360)	*	*	0(0-0)	725,41(0-5966)
IL-2	0(0-0)	102,61(15,53-189,68)	1,3(0-13,0)	430,25(0-2795,84)	0,81(0-3,24)	5,45(0-12,91)	5,57(0-94,46)	206,27-0-4174,33)
IL-6	29,59(14,95-44,22)	*	182,03(0-1798)	245,24(0-1167)	74,02(49,56-113,11)	*	1.969,41(0-20351)	2098,78(0-17292)

IL-10	1,02(0,63-1,41)	278,64(274,75-282,54)	0,01(0-0,1)	14,29(0-71,73)	0,35(0-0,86)	35,42(10,83-57,28)	2,03(0-38,31)	21,57(0-174,53)
IFN	0,05(0,02-0,08)	2605,14(2058,35-3151,94)	0,01(0-0,1)	799,12(0-2632,72)	0,04(0-0,08)	62,94(0,10-248,88)	4,34(0-127,68)	638,10(0-3633,75)
TNF	7,42(6,58-8,25)	2877,63(1809,63-3945,63)	0(0-0)	303,32(0-2292)	5,28(1,69-8,25) N=3	370,02(105,96-948,70)	0(0-0)	226,90(0-2603)
IL-17A	3,73(0-7,45)	19,80(4,18-35,41)	0,67(0-6,68)	3,85(0-13,86)	0(0-0)	5,94(0-23,76)	1,40(0-8,58)	7,47(0-32,02)
IL-4	0,85(0,58-1,12)	3,6(1,52-5,68)	0,55(0-1,19)	1,1(0-2,48)	0,63(0-1,31)	1,04(0-1,9)	0,79(0-1,45)	1,48(0-5,5)
IL-21	*	*	0(0-0)	0(0-0)	*	*	0,66(0-6,55)	1,045(0-6,99)
IL-22	*	*	0(0-0)	0,954(0-9,54)	*	*	0,18(0-5,91)	0(0-0)

495   PBMC were stimulated *in vitro* with 2 μL/mL PHA (+) or 5 μg/mL pertussis toxin (PT), or left unstimulated (-) for 48 h and 120 h respectively,

496   and the supernatants collected to measure cytokine concentrations by flow cytometry. Values are presented as mean (minimum-maximum). *

497   Samples were not available for IL-5, IL-13, IL-21 and IL-22 quantification after PHA stimulation. NVC: children born from unvaccinated women;

498   VC: children born from Tdap-vaccinated women; pg/mL, picogram/milliliter.