2 **Adintoviruses: An Animal-Tropic Family of Midsize Eukaryotic Linear dsDNA**
3 **(MELD) Viruses**

5 Gabriel J. Starrett[†], Michael J. Tisza[†], Nicole L. Welch, Anna K. Belford, Alberto Peretti,

6 Diana V. Pastrana, and Christopher B. Buck*

7
8 Lab of Cellular Oncology, NCI, NIH, Bethesda, MD, 20892 USA
9
10 [†]These authors contributed equally
11
12 *Corresponding author: buckc@mail.nih.gov
13

Abstract

Polintons (also known as Mavericks) were initially identified as a widespread class of eukaryotic transposons named for their hallmark type B DNA *pol*ymerase and retrovirus-like *int*egrase genes. It has since been recognized that many polintons encode possible capsid proteins and viral genome-packaging ATPases similar to those of a diverse range of double-stranded DNA (dsDNA) viruses. This supports the inference that at least some polintons are viruses that remain capable of cell-to-cell spread. At present, there are no polinton-associated capsid protein genes annotated in public sequence databases. To rectify this deficiency, we used a data-mining approach to investigate the distribution and gene content of polinton-like elements and related DNA viruses in animal genomic and metagenomic sequence datasets. The results define a discrete family-like clade of animal-specific viruses with two genus-level divisions. We suggest the family name *Adintoviridae,* connoting similarities to *ad*enovirus virion proteins and the presence of a retrovirus-like *int*egrase gene. Although adintovirus-class PolB sequences were detected in datasets for fungi and various unicellular eukaryotes, sequences resembling adintovirus virion proteins and accessory genes appear to be restricted to animals. Degraded adintovirus sequences are endogenized into the germlines of a wide range of animals, including humans.

Introduction

Analyses based on conserved protein structural features have increasingly revealed commonalities between families of eukaryotic viruses with double-stranded DNA (dsDNA) genomes. A current model places a loosely defined group known as polinton-like viruses at the center of a network of evolutionary relationships (Koonin, Dolja et al. 2015, Koonin, Krupovic et al. 2015). Polintons (also known as Mavericks) are defined by the presence of a type B DNA *pol*ymerase (PolB) and a retrovirus-like *int*egrase gene. Although polintons were first recognized as transposons, the observation that many of them encode predicted virion proteins supports the proposal that most elements initially designated as polinton transposons are actually integrated proviruses that may remain capable of infectious cell-to-cell spread (Krupovic, Bamford et al. 2014, Krupovic and Koonin 2015).

Adenoviruses, poxviruses, and baculoviruses are familiar groups of animal-tropic viruses that encode genes distantly similar to polinton PolB and virion proteins (Koonin, Dolja et al. 2015). An emerging group of viruses known as virophages, which are named for their ability to parasitize megaviruses that infect unicellular eukaryotes, also encode polinton-like PolB and virion protein genes as well as, in some cases, retrovirus-like integrase genes. (Duponchel and Fischer 2019).

Although polintons have been widely recognized in animal genomics and transcriptomics datasets (Krupovic, Bamford et al. 2014), the proposed capsid genes of these elements are not currently annotated in public sequence databases. This has led to confusion. For instance, a recent study detected two "Maverick transposons" in insect cell cultures but failed to annotate the capsid genes that identify them as likely viruses (Geisler 2018). In another example, a set of classic polinton PolB gene fragments detected in mouse fecal samples appear in GenBank with annotations incorrectly indicating that they are parvovirus structural proteins (Williams, Che et al. 2018). A primary goal of this study is to develop a coherent classification system for animal-tropic viruses with polinton-like genes and to facilitate further discovery by rendering annotated examples of these viruses searchable in public databases.

Results

Classification of animal-associated contigs with polinton-like PolB genes

TBLASTN searches using the inferred virion maturational protease (Adenain) of an arbitrarily chosen *Parasteatoda* spider contig (AOMJ02256338) identified hundreds of >10kb contigs of interest in NCBI's whole genome shotgun (WGS) and transcriptome shotgun assembly (TSA) databases, as well as in *de novo* assemblies of various datasets of interest from the Sequence Read Archive (SRA). In animal datasets, a great majority of the larger adenain-bearing elements were found to encode either an archetypal polinton-like PolB (pfam03175) or a divergent PolB <30% identical to the pfam03175 type. Both PolB types encode a distinctive N-terminal domain with predicted structural similarity to the ovarian tumor superfamily of ubiquitin-specific proteases (OTU). Adenovirus PolB sequences lack the OTU domain. In this study, we refer to

the OTU-pfam03175 PolB class as Alpha and the second OTU-PolB class as Beta. In RepBase
https://www.girinst.org/, polinton groups 1, 2, 3, 4, and 9 each contain both Alpha and Beta PolB
genes. Alpha PolB genes have previously been binned with hybrid virophages, ungrouped
polinton-like viruses, and Polintons group 2, while Beta PolB genes have been binned with
ungrouped polinton-like viruses, plant and fungal mitochondrial plasmids, and Polintons group 1
(Moriyama, Terasawa et al. 2008, Yutin, Raoult et al. 2013, Yutin, Kapitonov et al. 2015, Yutin,
Shevchenko et al. 2015).

In BLASTP searches, Alpha PolB sequences give strong hits (E-values ~1e-60) for an emerging
group of bipartite parvovirus-like viruses called bidnaviruses or bidensoviruses (Krupovic and
Koonin 2014). Use of the DELTA-BLAST algorithm (Boratyn, Schaffer et al. 2012) yields
stronger hits (E-values <1e-100) for adenoviruses. Beta PolB sequences typically do not yield
bidnavirus hits in BLASTP searches and instead give moderate hits (E-value ~1e-15) for the
PolB proteins of megaviruses (e.g., Faustovirus and Klosneuvirus) as well as various
bacteriophages (Figure 1). Neither of the two PolB classes detects known virophage PolB
sequences in BLASTP or DELTA-BLAST searches.

94
95
Figure 1: PolB BLASTP relationships. PolB protein sequences were subjected to all-
against-all sequence similarity network analysis with a BLASTP E-value cutoff of 1e-06.
Figure supplement 1: an interactive version of Figure 1 that can be viewed using
Cytoscape software https://cytoscape.org
Figure supplement 2: sequence compilations for PolB and other proteins (fasta format, zip
compressed)
Figure supplement 3: network analysis of Hexon and Penton proteins

In addition to Adenain and PolB, nearly all >10 kb contigs from the WGS and TSA surveys
encode a retrovirus-like integrase (protein family rve) as well as a protein similar to a group of

FtsK/HerA-type nucleoside triphosphatases (FtsK) that are thought to mediate the packaging of
viral genomes into virions (Iyer, Makarova et al. 2004).
108
Alignments of selected contigs back to parent read datasets showed that coverage depth fell to
zero near the ends of some contigs. An example is shown graphically in Figure 2 Figure
supplement 2. In some cases, such as a *Mayetiola destructor* (barley midge) read dataset, a single
predominant apparently free-ended sequence could be assembled but the dataset also contained a
range of lower-coverage variant reads near the termini, some of which extended into inverted
terminal repeats (ITRs) and host genomic DNA sequences. The observation suggests that the
integrase gene is functional and mediates integration events akin to those observed in virophages
that encode rve integrases (Fischer and Hackl 2016).
117
118   Based on the similarities to *ad*enoviruses and the presence of *int*egrase and *virus* genome-
119   packaging genes we suggest that this group of animal-associated elements could be referred to as
120   "adintoviruses." Maps of reference adintoviruses are shown in Figure 2.
121

122
Figure 2: Genome maps of two representative adintoviruses.
Figure supplement 1: accession numbers and full Linnaean designations of animal hosts (MS
Excel table).
Figure supplement 2: graphical examples of the gene-annotation process.
Figure supplement 3: graphical maps of additional adintoviruses.
Figure supplement 4: annotated nucleotide maps of adintoviruses and related viruses
(GenBank-formatted text file).
130
HHpred searches confirmed the presence of ORFs with high-probability predicted structural
similarity to the double-jellyroll major capsid proteins (Hexons) and single-jellyroll vertex minor
capsid proteins (Pentons) of adenoviruses, virophages, megaviruses, or poxviruses (see Figure 2
Figure supplement 2 for illustrated examples of annotation methods). As expected, contigs with
Beta PolB genes encode Hexon and Penton proteins that occupy discrete clusters that encompass
the *Terrapene* Beta adintovirus cognates (Figure 1 Figure supplement 3). Although most contigs
with Alpha PolB genes encode Hexon and Penton proteins that cluster with the *Mayetiola*
cognates, some Alpha PolB contigs unexpectedly encode virion proteins that are interspersed
within the *Terrapene* cluster. Similar results were observed in analyses using traditional
phylogenetic trees. The results suggest the existence of distinct Alpha and Beta adintovirus
lineages, but with some examples reflecting horizontal transfer of virion protein operons from
the Beta PolB lineage into the Alpha PolB lineage. We have previously proposed a similar intra-
family horizontal gene transfer scenario for some species of polyomaviruses (Buck, Van
Doorslaer et al. 2016).
145
146

Other adintovirus genes

Adintoviruses encode three classes of proteins with predicted structures resembling known membrane-active proteins. A previously noted class (Yutin, Raoult et al. 2013) is similar to the phospholipase A2 (PLA2) domain of parvovirus VP1 virion proteins (Figure 2 Figure supplement 2). In parvoviruses, the domain is thought to be involved in membrane disruption during the infectious entry process. The PLA2-like genes, which are characteristic of *Mayetiola*-class (Alpha) virion protein operons, include a C-terminal domain similar to adenovirus virion core protein ten (pX). We suggest the gene name PLA2X.

Beta adintoviruses, as well as Alpha PolB adintoviruses with *Terrapene*-class (Beta) virion protein operons, encode homologs of the C-terminal regulatory domain of gasdermins, a group of pore-forming proteins that serve as executioners in pyroptosis (a form of inflammatory programmed cell death)(Dubois, Sorgeloos et al. 2019). Like PLA2X, adintovirus gasdermin homologs typically encode a pX-like domain near the C-terminus. Apparent homologs of a membrane-active spider venom protein known as cupiennin were also observed in Beta-class virion protein operons. The pairing of hallmark Beta-class virion accessory genes (GasderminX, Cupiennin) with a subset of Alpha PolB adintoviruses (Figure 2 Figure supplement 3) supports the hypothesis that some adintovirus species arose through chimerization between the Alpha and Beta adintovirus lineages.

Some classes of predicted protein sequences were conserved among adintoviruses but did not show clear hits for known proteins in BLASTP or HHpred searches. We assigned these groups of adintovirus-conserved proteins of unknown function numbered "Adintoc" names.

Small DNA tumor viruses (adenoviruses, polyomaviruses, and papillomaviruses (Pipas 2019)) encode proteins harboring conserved LXCXE motifs that that are known to engage cellular retinoblastoma (Rb) and related tumor suppressor proteins (de Souza, Iyer et al. 2010). Adenovirus E1A, papillomavirus E7, polyomavirus LT, and parvovirus NS3 oncoproteins typically encode the Rb-binding motif just upstream of a consensus casein kinase 2 acceptor motif ((ST)XX(DE)). Some oncogenes, such as E1A, encode an additional conserved region ((DEN)(LIMV)XX(LM)(FY)), referred to as CR1, that binds the groove containing the A and B cyclin folds within the Rb pocket domain (Pipas 1992, Gouw, Michael et al. 2018). In general, these predicted Rb-interacting motifs are adjacent to potential zinc- or iron-sulfur-binding motifs (typically, paired CXXC). Open reading frames encoding combinations of these short linear motifs were observed in adintovirus contigs. We refer to these predicted proteins, which typically occupy a region upstream of the PolB gene, as "Oncoid" genes, conjnoting their similarities to the known oncogenes of small DNA tumor viruses. Adintovirus homologs of anti-apoptotic proteins, such as Bcl2 and IAP, were also observed (Figure 2 Figure supplement 3).

Distribution of adintovirus-like PolB sequences in eukaryotic WGS datasets

The conserved catalytic core PolB sequences of either the *Mayetiola* barley midge Alpha adintovirus or *Terrapene* box turtle Beta adintovirus were used separately as baits in TBLASTN searches of WGS databases for eukaryotes. Retrieved protein sequences were trimmed to 80% similarity and subjected to clustering with an alignment score threshold of 60 (Shannon, Markiel

192 et al. 2003, Li and Godzik 2006, Huang, Niu et al. 2010, Fu, Niu et al. 2012, Zallot, Oberg et al.
193 2018). The clustering segregated away Beta adintovirus-like PolB sequences encoded by plant
194 and fungal mitochondria (e.g., EU365401, AF061244). The filtered sequences were subjected to
195 phylogenetic analyses (Figure 3).
196

197
198
Figure 3: Phylogenetic trees comprised of WGS hits for Alpha or Beta adintovirus PolB
sequences (left and right panels, respectively). Hits from insect datasets are colored
yellow, hits from tetrapod datasets are red and fungus-associated hits are blue. All other
types of eukaryotes are represented by black lines. Annotated branches show two Alpha
adintovirus sequences associated with bovine (*Bos*) lung samples clustering with
adintovirus sequences from insect datasets, suggesting an environmental insect source. In
contrast, exemplar *Mayetiola* and *Terrapene* adintoviruses cluster with sequences found
in other insect or terrestrial vertebrate datasets, respectively. Similarly, adintovirus PolB-
like sequences from *Powellomyces* and *Rhizophagus* fungi cluster with sequences from
other types of fungi.
Figure supplements 1 and 2: interactive Nexus-format tree files that can be viewed using
FigTree software http://tree.bio.ed.ac.uk/software/figtree/
211
212
Two complete Alpha adintovirus-like contigs (NKLS02000104, NKLS02001728) were observed
in assemblies of a PacBio-based WGS survey of bovine lung tissue. Sequences outside the
inferred proviral inverted terminal repeats (ITRs) in the two sequences were highly diverse and
mostly unidentifiable, but in a few reads the extra-proviral host sequences showed BLASTN
similarity to genomic DNA sequences of various beetles, including *Tribolium castaneum* (a flour
beetle that commonly infests cattle feed). Furthermore, the *Bos* lung-associated PolB sequences
occupy phylogenetic clades comprised of insect-associated PolB sequences (Figure 3). These
observations suggest that the two Alpha adintovirus sequences in the bovine datasets are insect-
derived environmental contaminants, rather than mammal-tropic viruses. Similarly, several Beta
adintovirus-like contigs (e.g., AANG04004209) found in a housecat oral swab sample show
close phylogenetic affinity for adintovirus sequences observed in salmon WGS datasets. In
another example, integrated adintoviruses found in a genomic dataset for olive trees (*Olea*
europaea) showed insect-like sequences outside the inferred ITRs and showed phylogenetic

affinity with PolB sequences from insect WGS datasets. Other adintovirus-like sequences found
in plant datasets resembled adintovirus PolB sequences associated with nematode datasets. It
thus appears that adintovirus sequences in some datasets are derived from environmental
sources, as opposed to a productive infection of the organism that was the target of the
sequencing effort.
231
Although there are examples of apparent environmental contamination, most adintovirus
sequences form discrete clades that recapitulate the phylogeny of the host organisms that were
the subjects of the WGS surveys. For example, a distinct clade of Alpha adintovirus PolB
sequences was observed in datasets for multiple related species of venomous snakes. Several
distinct clades of Beta adintoviruses were observed in datasets for amphibians and reptiles,
including the well-populated clade that houses the exemplar *Terrapene* adintovirus. The
exemplar *Mayetiola* adintovirus likewise occupies a clade exclusively populated by sequences
found in insect WGS datasets.
240
TBLASTN searches against *Terrapene* box turtle Beta adintovirus PolB and Hexon protein
sequences both yielded weak hits (E-value ~1e-05) for a locus on human chromosome 7. An
adintovirus GasderminX sequence was also detected at the locus. Alignments to *Terrapene*
adintovirus protein sequences were used to assign pseudogene annotations (Figure 4). The
detected element is a highly disrupted endogenized Beta adintovirus. Homologous nucleotide
sequences were detected in the genomes of primates, rodents, shrews, afrotherians, and
xenarthrans but not in datasets for ungulates, carnivores, bats, marsupials, or prototherians.
Endogenized adintovirus sequences observed in amphibian and reptile genomes do not share
recognizable nucleotide similarity with placental mammal-endogenized adintovirus sequences. It
is unclear whether a single adintovirus endogenization event affected an early placental mammal
and the endogenized virus was then lost in non-shrew Laurasiatherians or whether multiple
distinct endogenization events occurred in separate placental mammal lineages. Identification of
extant examples of placental mammal adintoviruses could help resolve this question.
254

255
Figure 4: An endogenized Beta adintovirus relic found on human chromosome 7. Degraded
pseudogenes interrupted by nonsense and frameshift mutations were reconstructed based on
alignments to the protein sequences of *Terrapene* box turtle adintovirus. Question marks
indicate that the reconstructed gene does not yield hits in BLAST searches of GenBank's
viruses taxon. Tentative gene assignments are based on synteny with the *Terrapene*
adintovirus. The reconstructed Hexon, GasderminX, and PolB protein sequences yield
DELTA-BLAST hits with E-values of 1e-21, 4e-05, and 3e-25, respectively.
Figure supplement 1: annotated GenBank-format nucleotide map of the human chromosome
7 endogenized adintovirus depicted graphically in Figure 4.
265
266 **Viruses with adinto-like genes in non-animal eukaryote datasets**
267

Eukaryotic viruses with midsize (10-50 kb) linear dsDNA genomes show a remarkable degree of
genomic modularity (Koonin, Dolja et al. 2015, Yutin, Kapitonov et al. 2015, Yutin, Shevchenko
et al. 2015). The apparently promiscuous horizontal gene transfer and lack of any single defining
gene for these viruses makes the group taxonomically challenging. We propose the collective
acronym MELD (midsize eukaryotic linear dsDNA) virus for the dizzyingly polyphyletic
category. The name, which would encompass adenoviruses and adintoviruses, is intended to fill a
gap between other operationally defined umbrella groups, such as CRESS viruses, small DNA
tumor viruses, nucleocytoplasmic large DNA viruses, and megaviruses.
276
Datasets for *Blastocystis hominis* (a diatom-related unicellular eukaryote that commonly inhabits
the human gut) contain MELD virus sequences that unite Alpha adintovirus-like PolB and
integrase genes with inferred virion proteins whose primary sequences are not recognizably
similar to known virion proteins (Figure 5). Gene identities for the *Blastocystis* virus were
inferred based on HHpred results. Comparable MELD viruses were confirmed in rumen
metagenomic datasets for sheep (Yutin, Kapitonov et al. 2015) and cattle, as well as in WGS
datasets for green algae and fungi. In phylogenetic analyses, the PolB sequences of these viruses
occupy long branches that are distant from animal-associated PolB clades (Figure 3).
285

286
287
Figure 5: MELD viruses (and related elements) with adintovirus-like PolB genes. Greek
letters indicate genes similar to adintoviruses in BLASTP or DELTA-BLAST searches.
"Ad-" indicates similarity to adenovirus sequences. Abbreviations: Fiber, predicted
structural or primary sequence similarity to bacteriophage tail fibers or coiled-coil
proteins; Lectin, predicted structural similarity to galactose-binding domains; Dcm,
predicted structural similarity to cytosine DNA methyltransferases; TBP, similar to

TATA binding proteins; Matrix/Capsid/NC/Pro, similarities to retroviral Gag and
retropepsin; S3H, poxvirus D5-like superfamily 3 helicase.
296
297
Contigs encoding Alpha adintovirus-like PolB and integrase genes were found in metagenomics
datasets for bioreactor-cultured human feces, human urine samples, and human oral swab
samples (Santiago-Rodriguez, Ly et al. 2015). This group of closely related sequences was only
detected in datasets from a single laboratory and not in other human metagenomics surveys.
Divergent variants of predicted proteins from the feces-associated virus were found in contigs
from datasets for *Cyanophora paradoxa*, a species of glaucophyte algae (e.g., QPMI01000557),
suggesting that the human feces-associated adintovirus-like sequences were derived from an
environmental source.
306
Six MELD virus genomes assembled from a single *Powellomyces* SRA dataset unite sequences
resembling adenovirus vertex fiber proteins with either a Beta adintovirus-like PolB (Figure 5) or
a surprising variety of non-PolB DNA replicases (Figure 6). MELD virus genomes encoding
genes similar to Alpha adintovirus virion proteins (E-values ~1e-7 to 1e-21) were assembled
from datasets for *Capsaspora owczarzaki, Monosiga brevicollis,* and *Trichoplax H2* (unicellular
eukaryotes that are thought to be closely related to animals). Aside from the abovementioned
insect- and nematode-associated adintovirus sequences found in datasets for plants, adintovirus-
like virion protein sequences were not detected in datasets for other non-animal eukaryotes.
Capsaspora MELD virus 1 and the *Trichoplax* MELD virus both encode superfamily 1 helicase
(S1H) genes instead of a PolB gene. Various megaviruses and bacteriophages encode similar
S1H genes, as does a MELD virus observed in *Physarum polycephalum* slime mold and in
Powellomyces MELD virus 4. Full-length S1H replicase genes of this class were not detected in
animal WGS datasets, with the exception of seemingly endogenized degraded virus-like contigs
in datasets for several coral and jellyfish species and a helitron-like element found in
Branchiostoma lancelets (e.g., RDEB01009762, ABEP02037959).
322
323

Figure 6: MELD viruses with other replicases. Greek letters indicate genes with sequences similar to Alpha or Beta adintoviruses in BLASTP searches. Sequences similar to adenoviruses are marked with "Ad-." Yellow flags represent predicted tRNA genes. Abbreviations: dUTPase, similar to poxvirus deoxy-UTP diphosphatases; Fiber, similarity to bacteriophage tail fibers or other coiled-coil proteins; Tapemeasure, similarity to phage tail tape measure proteins; S1H, RecD/Pif1-like superfamily 1 helicase; YRec, homolog of phage tyrosine recombinases; TRAF, predicted structural similarity to TNF receptor associated factor 3; UCH, predicted structural similarity to ubiquitin C-terminal hydrolase cysteine proteases; IVa2, sequence similarity to adenovirus pIVa2 viral genome-packaging ATPases; DnaA-like, sequence distantly similar to DnaA and DnaB-like helicases; VI, similar to adenovirus virion core protein six; PolA, DNA polymerase family A (Pfam:00476); S3H superfamily 3 helicase similar to those observed in virophages and megaviruses; S2H, superfamily 2 helicase similar to DEAD-box helicase of Yellowstone Lake virophage 7 (YP_009177696); TRAF UCH TBP Fbox, homologs of host proteins with these gene symbols; SLATT, homolog of host SMODS and SLOG-associating 2TM effector domain proteins; VLTF, homolog of mimivirus VLTF3-like transcription factor. See main text for information about other gene names.

In WGS searches for sequences resembling human adenovirus type 5 PolB, we did not detect any
contigs resembling full-length viruses in non-animal datasets. The searches did reveal the
complete ITR-bounded genome of a typical mastadenovirus in a dataset for *Dipodomys ordii* (a
type of kangaroo rat) as well as apparently complete MELD viruses in datasets for *Hydra*
oligactis (brown hydra) and *Anelosimus studiosus* (a type of tangle-web spider). Like known
adenoviruses, the *Hydra* and *Anelosimus* MELD viruses do not encode integrase genes and their
PolB genes do not encode detectable OTU domains.
352
353 **PolB⁺ parvoviruses**
354
BLASTP searches using Alpha adintovirus PolB sequences return high-likelihood matches (E-
values <1e-80) for the PolB genes of an emerging group of bipartite parvoviruses referred to as
bidnaviruses (Krupovic and Koonin 2014)(Figure 1). Like adintoviruses, bidnavirus PolB genes
encode an N-terminal OTU domain. We searched assemblies of SRA datasets of interest for
additional examples of bidnavirus genomes. PolB⁺ contigs were detected in datasets for the gut
contents of African termites (*Cubitermes ugandensis*), dog (*Canis lupus familiaris*) feces, the silk
glands of a false wolf spider (*Tengella perfuga*), and Tasmanian devil (*Sarcophilus harrisii*)
feces (Figure 7). The dog feces PolB sequence is 53% similar to the "structural protein" of Fresh
Meadows "densovirus" 3 previously detected in mouse (*Mus musculus*) feces
(AWB14611)(Williams, Che et al. 2018).
365

366
367 Figure 7: Bidnaparvovirus and non-animal parvovirus genome maps. Abbreviations: GIIM,
368 similarity to group II intron maturases; tailspike, similarity to bacteriophage short tail
369 fibers.
370
In previously reported bidnaviruses, the termini of each of the two genome segments have
matching nucleotide sequences. To search for second segments, we probed the assemblies for
examples of other contigs with termini similar to the ITRs of the initially observed bidnavirus
contigs. The datasets were also searched for contigs with sequences similar to previously
reported bidnavirus proteins. Second segments were not detected, suggesting that the five new
bidnaviruses may be monopartite. We suggest that the apparently monopartite viruses could still
be referred to as bidnaviruses (or, more specifically, bidnaparvoviruses) but with the "bidna"
moniker connoting the presence of two types of DNA replicase genes, as opposed to the original
connotation of a virus with two genomic DNA segments.
380

381   Searches for examples of parvovirus NS1-like sequences did not reveal clear examples outside of
382   multicellular animal datasets. A marginal exception was a group of sequences found in datasets
383   for *Abeoforma whisleri* and *Ichthyophonus hoferi*, two unicellular eukaryotes that are thought to
384   be closely related to multicellular animals. The observations suggest an early-animal origin for
385   parvoviruses that involved acquisition of genes from Alpha adintoviruses.
386
387
388   **Discussion**
389
We have identified a coherent family-like grouping of animal viruses that we call adintoviruses,
connoting their hallmark adenovirus-like virion protein genes and retrovirus-like integrase genes.
Adintovirus sequences are detectable either as apparently free linear DNA molecules or as
endogenized integrants in WGS datasets representing all eumetazoan phyla. Although sequences
resembling the PolB proteins of Alpha and Beta adintoviruses can also be found in datasets for
non-animal eukaryotes, the sequences of adintovirus virion proteins appear to be restricted to
animals.
397
We imagine that the related Alpha and Beta adintovirus-like lineages might have infected early
eukaryotes and the two lineages gradually co-evolved with major divisions of eukaryotes,
including multicellular animals. In this model, the sequences of the virion protein genes
presumably evolved more rapidly than the conserved catalytic core of PolB, resulting in
distinctive mutually unrecognizable virion protein sequences specific to each major division of
eukaryotes. The model suggests that adenoviruses could be thought of as a related sister lineage
that also arose in or before the first animals. Although adenoviruses are currently only known to
infect vertebrates, the idea that the lineage long predates the emergence of vertebrates is
consistent with our identification of a distantly adenovirus-like sequence in a spider dataset
(Figure 6).
408
In non-animal eukaryote datasets, adintovirus-like PolB sequences can be found in a wide range
of sequence contexts, ranging from elements with no obvious virion proteins to the genomes of
megaviruses. Conversely, it appears that the adintovirus-like PolB can readily be replaced with
other types of DNA replicase genes (Figure 6). This presumably reflects the previously proposed
rampant horizontal gene transfer among virus lineages that infect unicellular eukaryotes.
Although similar horizontal gene transfer events appear to have occurred between various
animal-tropic virus families, including adintoviruses and parvoviruses (Figure 7), adomaviruses
and polyomaviruses (Mizutani, Sayama et al. 2011, Dill, Camus et al. 2018)
https://www.biorxiv.org/content/10.1101/341131v2 and papillomaviruses and polyomaviruses
(Woolford, Rector et al. 2007), each of these cases appears to represent single ancient event. It
may be that the evolution of distinct tissues and organs or the development of cell-mediated
immunity in multicellular animals placed limits on the likelihood that different virus lineages can
co-infect a single cell and productively recombine. From this view, the distinctive gene
combinations seen in adintoviruses and adenoviruses might simply be bottlenecked examples of
the much larger range of gene combinations observed in MELD viruses of unicellular eukaryotes
(Yutin, Kapitonov et al. 2015, Yutin, Shevchenko et al. 2015).
425

It has generally been assumed that the functionally similar oncogenes found in adenoviruses,
papillomaviruses, parvoviruses, and polyomaviruses arose through convergent evolution or
through horizontal gene transfer between virus families (de Souza, Iyer et al. 2010). Although
small DNA tumor virus oncogenes show low overall sequence similarity, they can be roughly
defined based on the presence of short linear motifs. Many adintoviruses encode candidate
"Oncoid" proteins with these motifs. Bombyx silkworm bidnaparvovirus NS3, which we have
designated as a candidate Oncoid (Figure 7) has previously been shown to be similar to a
baculovirus protein of unknown function (Krupovic and Koonin 2014). We note that many of the
proposed baculovirus homologs (e.g., YP_009506034) share potential zinc-coordinating cysteine
residues as well as a C-terminal LXCXE/CK2 site, qualifying the baculovirus proteins as
candidate Oncoids as well. Surprisingly, candidate Oncoids were also observed in MELD viruses
of unicellular eukaryotes (Figure 6). The predicted Oncoid2 gene of *Capsaspora* protist MELD
virus 2 detects polyomavirus Large T oncogenes in DELTA-BLAST searches (E-value 4e-16). It
is interesting to imagine that oncogenes in a broad range of animal DNA viruses might share an
ancestry that pre-dates the emergence of multicellular animals.
441
Adintoviruses encode a number of accessory genes that appear to be homologs of membrane-
active proteins found in animal venom. These include bee and snake venom PLA2 and melittin,
as well as a spider venom protein called cupiennin. Interestingly, venom PLA2 and melittin
(which shows similarity to adenovirus pX in HHpred searches) act in concert (Vogt, Patzer et al.
1970), suggesting the speculative hypothesis that these venom genes might have arisen from a
captured viral PLA2X-like gene.
448
In unpublished work, our group used a standard baculovirus-based expression system
(ThermoFisher) to generate a virus-like particle (VLP) vaccine against BK polyomavirus
(BKV)(Peretti, Geoghegan et al. 2018). The project provided an inadvertent natural experiment.
Recombinant baculoviruses were generated in Sf9 cells and bulk protein expression was
performed using the *Trichoplusia ni* cell line High Five. BKV VLPs were purified according to
previously reported methods (Cardone, Moyer et al. 2014) involving ultracentrifugation through
density gradients, nuclease digestion, and size exclusion chromatography. Deep sequencing of
DNA extracted from the purified VLP preparation shows high-depth coverage of *Spodoptera*
adintovirus genomes alongside incomplete patchy coverage of endogenized *Trichoplusia*-
specific homologs of the two *Spodoptera* viruses (Supplemental File 1). It appears that Sf9-
derived adintoviruses infected the High Five cells and this led to the production of adintovirus
virions that co-purified with the recombinant BKV VLPs. The results suggest that standard insect
cell cultures could serve as a laboratory model for productive adintovirus infection.
462
A Beta adintovirus was detected in transcriptomic and WGS datasets for Mexican blind tetra
cavefish (*Astyanax mexicanus*). Adintovirus transcripts were most abundant in head, kidney, and
intestine samples and least abundant in muscle and whole embryo samples (Supplemental Table
1). Analysis of the WGS dataset showed that adintovirus DNA reads outnumbered reads for a
single copy host gene (gamma tubulin, NW_019172896) by a factor of 25. At both an RNA and
DNA level the *Astyanax* sequence showed a high degree of uniformity, suggesting a clonal
infection. In contrast, pet store samples of a different species of tetra, *Gymnocorymbus ternetzi*
(SRR2040422), showed such a complex range of adintovirus sequence variants that assembly of

contigs representing complete viral genomes was challenging. These observations suggest that
tetras might serve as a tractable laboratory model for adintovirus infection.
473
Adintoviruses have a number of features that could make them useful as recombinant gene
transfer vectors. Their genome size is substantially larger than commonly used retroviral and
parvoviral vectors. In contrast to adenovirus- and baculovirus-based vector systems, adintovirus
genomes are small enough to be manipulated entirely in the setting of standard plasmids. An
intriguing feature of the adintovirus integrase gene is the presence of a predicted chromodomain
that, in LTR retrotransposons, is believed to influence integration site specificity (Kordis 2005).
This could theoretically offer an advantage over retroviral vectors, which show little integration
site specificity. Another potential practical use for adintoviruses might be as biocontrol agents
for pest organisms, such as *Mayetiola destructor* barley midges or chytrid fungi that parasitize
amphibians.
484
An important implication of this study is that there may be additional unappreciated families of
animal viruses hiding in plain sight in sequence databases. Adintoviruses may have been
relatively easy quarry because they are able to integrate into host genomes, such that they are
detectable in WGS datasets of randomly sampled animals that did not happen to be suffering
from an active infection. In contrast to the hundreds of adintovirus-like contigs detected in our
initial WGS survey, focused searches for adenoviruses (which do not encode integrases) detected
only a single complete adenovirus genome. For future discovery efforts, it will be important to
develop higher throughput methods using sensitive structure-guided searches to identify
divergent new examples of viral hallmark genes in sequence datasets representing many
individuals, including subjects suffering from disease. The key goal will be to understand which
combinations of genes tend to co-occupy single contigs. Recently reported bioinformatics
pipelines, such as Cenote-Taker (Tisza, Pastrana et al. 2019) and Mash Screen (Ondov, Starrett
et al. 2019), should be useful for these purposes. Deposition of annotated viral genome
sequences into publicly searchable databases will be critical for further expanding our
understanding of the eukaryotic virome.
500

Materials and Methods

Detection and analysis of viral sequences

Adomavirus LO8 (Adenain) sequences were initially used for TBLASTN searches of the NCBI
TSA and WGS databases. The relationship between adomavirus and adintovirus virion proteins
is the subject of a separate manuscript (BioRxiv 341131v2). The Adenain sequences of *Nephila*
orb-weaver spider contig (GFKT014647032) or a *Parasteatoda* spider contig (AOMJ02256338)
were arbitrarily chosen for further TBLASTN searches of eukaryotic datasets in TSA and WGS
databases. Adenain-bearing contigs 4-50kb in length were further searched (using CLC
Genomics Workbench) for BLASTP-detectable PolB homologs. Contigs were inspected for the
presence of nearly overlapping arrays of large (>100 AA) open reading frames. Contigs with
inverted repeats flanking the ORF cluster were favored, but this was not a strict sorting criterion.

Selected contigs of interest were initially annotated using DELTA-BLAST searches of GenBank
nr or HHpred analyses of single or aligned protein sequences against PDB_mmCIF70,
COG_KOG, Pfam-A, and NCBI_CD databases (Altschul, Madden et al. 1997, Altschul,
Wootton et al. 2005, Soding 2005, Hildebrand, Remmert et al. 2009, Gerlt, Bouvier et al. 2015,
Meier and Soding 2015, Zimmermann, Stephens et al. 2017). Protein sequences were extracted
from the contigs using getORF http://bioinfo.nhri.org.tw/cgi-bin/emboss/getorf (Rice, Longden
et al. 2000). Extracted protein sequences were clustered using EFI-EST
https://efi.igb.illinois.edu/efi-est/ (Gerlt, Bouvier et al. 2015, Zallot, Oberg et al. 2018) and
displayed using Cytoscape v3.7.1 (Shannon, Markiel et al. 2003). Multiple sequence alignments
were constructed using MAFFT https://toolkit.tuebingen.mpg.de/#/tools/mafft. Contigs were
annotated using Cenote-Taker (Tisza, Pastrana et al. 2019) with an iteratively refined library of
conserved adintovirus protein sequences. Compiled protein sequences are provided as a zipped
set of fasta-format text files in Figure 1 Figure supplement 2. Maps were drawn using MacVector
17 software. Phylogenetic analyses were performed using MAFFT 7
https://mafft.cbrc.jp/alignment/server/ (Kuraku, Zmasek et al. 2013, Katoh, Rozewicki et al.
2019) and displayed using FigTree 1.4.4 http://tree.bio.ed.ac.uk/software/figtree/.

Selected contigs for which SRA datasets were available were subjected to reference-guided re-
assembly using Megahit 1.2.9 (Li, Liu et al. 2015, Li, Luo et al. 2016) and/or the map reads to
reference function of CLC Genomics Workbench. Annotated maps were submitted to GenBank
as third party annotation assemblies (TPA_asm). Graphical examples of the annotation process
are depicted in Figure 2 Figure supplement 2.
533

Data Availability

GenBank accession numbers for sequences deposited in association with this study are:
BK010888 BK010889 BK010890 BK010893 BK010894 BK010998 BK010999 BK011000
BK011001 BK011002 BK011003 BK011004 BK011005 BK011006 BK011007 BK011008
BK011009 BK011010 BK011011 BK011022 BK011023 BK011024 BK011025 BK011026
BK012042 BK012043 BK012044 BK012045 BK012046 BK012047 BK012048 BK012049
BK012050 BK012051 BK012052 BK012053 BK012054 BK012055 BK012056 BK012057
BK012058 BK012059 BK012060 BK012061 BK012062 BK012063 BK012064 BK012084
BK012085 BK012086.

Acknowledgments

545 The authors are indebted to Eugene Koonin and Natalya Yutin for their generous guidance and
546 for the spirited discussions that inspired us to pursue this study. We thank Karl Münger and Jim
547 Pipas for their extensive advice about oncogene sequence motifs and Chris Bellas for useful
548 discussions and for critical evaluation of the manuscript. We are also grateful to Patrick
549 McTamney for guiding the production of BK polyomavirus VLP stocks.

550
551

References

Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller and D. J. Lipman (1997). "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs." Nucleic Acids Res **25**(17): 3389-3402.

Altschul, S. F., J. C. Wootton, E. M. Gertz, R. Agarwala, A. Morgulis, A. A. Schaffer and Y. K. Yu (2005). "Protein database searches using compositionally adjusted substitution matrices." FEBS J **272**(20): 5101-5109.

Boratyn, G. M., A. A. Schaffer, R. Agarwala, S. F. Altschul, D. J. Lipman and T. L. Madden (2012). "Domain enhanced lookup time accelerated BLAST." Biol Direct **7**: 12.

Buck, C. B., K. Van Doorslaer, A. Peretti, E. M. Geoghegan, M. J. Tisza, P. An, J. P. Katz, J. M. Pipas, A. A. McBride, A. C. Camus, A. J. McDermott, J. A. Dill, E. Delwart, T. F. Ng, K. Farkas, C. Austin, S. Kraberger, W. Davison, D. V. Pastrana and A. Varsani (2016). "The Ancient Evolutionary History of Polyomaviruses." PLoS Pathog **12**(4): e1005574.

Cardone, G., A. L. Moyer, N. Cheng, C. D. Thompson, I. Dvoretzky, D. R. Lowy, J. T. Schiller, A. C. Steven, C. B. Buck and B. L. Trus (2014). "Maturation of the human papillomavirus 16 capsid." MBio **5**(4): e01104-01114.

de Souza, R. F., L. M. Iyer and L. Aravind (2010). "Diversity and evolution of chromatin proteins encoded by DNA viruses." Biochim Biophys Acta **1799**(3-4): 302-318.

Dill, J. A., A. C. Camus, J. H. Leary and T. F. F. Ng (2018). "Microscopic and Molecular Evidence of the First Elasmobranch Adomavirus, the Cause of Skin Disease in a Giant Guitarfish, Rhynchobatus djiddensis." MBio **9**(3).

Dubois, H., F. Sorgeloos, S. T. Sarvestani, L. Martens, Y. Saeys, J. M. Mackenzie, M. Lamkanfi, G. van Loo, I. Goodfellow and A. Wullaert (2019). "Nlrp3 inflammasome activation and Gasdermin D-driven pyroptosis are immunopathogenic upon gastrointestinal norovirus infection." PLoS Pathog **15**(4): e1007709.

Duponchel, S. and M. G. Fischer (2019). "Viva lavidaviruses! Five features of virophages that parasitize giant DNA viruses." PLoS Pathog **15**(3): e1007592.

Fischer, M. G. and T. Hackl (2016). "Host genome integration and giant virus-induced reactivation of the virophage mavirus." Nature **540**(7632): 288-291.

Fu, L., B. Niu, Z. Zhu, S. Wu and W. Li (2012). "CD-HIT: accelerated for clustering the next-generation sequencing data." Bioinformatics **28**(23): 3150-3152.

Geisler, C. (2018). "A new approach for detecting adventitious viruses shows Sf-rhabdovirus-negative Sf-RVN cells are suitable for safe biologicals production." BMC Biotechnol **18**(1): 8.

Gerlt, J. A., J. T. Bouvier, D. B. Davidson, H. J. Imker, B. Sadkhin, D. R. Slater and K. L. Whalen (2015). "Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST): A web tool for generating protein sequence similarity networks." Biochim Biophys Acta **1854**(8): 1019-1037.

Gouw, M., S. Michael, H. Samano-Sanchez, M. Kumar, A. Zeke, B. Lang, B. Bely, L. B. Chemes, N. E. Davey, Z. Deng, F. Diella, C. M. Gurth, A. K. Huber, S. Kleinsorg, L. S. Schlegel, N. Palopoli, K. V. Roey, B. Altenberg, A. Remenyi, H. Dinkel and T. J. Gibson (2018). "The eukaryotic linear motif resource - 2018 update." Nucleic Acids Res **46**(D1): D428-D434.

Hildebrand, A., M. Remmert, A. Biegert and J. Soding (2009). "Fast and accurate automatic structure prediction with HHpred." Proteins **77 Suppl 9**: 128-132.

Huang, Y., B. Niu, Y. Gao, L. Fu and W. Li (2010). "CD-HIT Suite: a web server for clustering and comparing biological sequences." Bioinformatics **26**(5): 680-682.

Iyer, L. M., K. S. Makarova, E. V. Koonin and L. Aravind (2004). "Comparative genomics of the
FtsK-HerA superfamily of pumping ATPases: implications for the origins of chromosome
segregation, cell division and viral capsid packaging." Nucleic Acids Res **32**(17): 5260-5279.
Katoh, K., J. Rozewicki and K. D. Yamada (2019). "MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization." Brief Bioinform **20**(4): 1160-1166.
Koonin, E. V., V. V. Dolja and M. Krupovic (2015). "Origins and evolution of viruses of
eukaryotes: The ultimate modularity." Virology **479-480**: 2-25.
Koonin, E. V., M. Krupovic and N. Yutin (2015). "Evolution of double-stranded DNA viruses of
eukaryotes: from bacteriophages to transposons to giant viruses." Ann N Y Acad Sci **1341**: 10-
24.
Kordis, D. (2005). "A genomic perspective on the chromodomain-containing retrotransposons:
Chromoviruses." Gene **347**(2): 161-173.
Krupovic, M., D. H. Bamford and E. V. Koonin (2014). "Conservation of major and minor jelly-
roll capsid proteins in Polinton (Maverick) transposons suggests that they are bona fide viruses."
Biol Direct **9**: 6.
Krupovic, M. and E. V. Koonin (2014). "Evolution of eukaryotic single-stranded DNA viruses of
the Bidnaviridae family from genes of four other groups of widely different viruses." Sci Rep **4**:
5347.
Krupovic, M. and E. V. Koonin (2015). "Polintons: a hotbed of eukaryotic virus, transposon and
plasmid evolution." Nat Rev Microbiol **13**(2): 105-115.
Kuraku, S., C. M. Zmasek, O. Nishimura and K. Katoh (2013). "aLeaves facilitates on-demand
exploration of metazoan gene family trees on MAFFT sequence alignment server with enhanced
interactivity." Nucleic Acids Res **41**(Web Server issue): W22-28.
Li, D., C. M. Liu, R. Luo, K. Sadakane and T. W. Lam (2015). "MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn graph."
Bioinformatics **31**(10): 1674-1676.
Li, D., R. Luo, C. M. Liu, C. M. Leung, H. F. Ting, K. Sadakane, H. Yamashita and T. W. Lam
(2016). "MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced
methodologies and community practices." Methods **102**: 3-11.
Li, W. and A. Godzik (2006). "Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences." Bioinformatics **22**(13): 1658-1659.
Meier, A. and J. Soding (2015). "Automatic Prediction of Protein 3D Structures by Probabilistic
Multi-template Homology Modeling." PLoS Comput Biol **11**(10): e1004343.
Mizutani, T., Y. Sayama, A. Nakanishi, H. Ochiai, K. Sakai, K. Wakabayashi, N. Tanaka, E.
Miura, M. Oba, I. Kurane, M. Saijo, S. Morikawa and S. Ono (2011). "Novel DNA virus isolated
from samples showing endothelial cell necrosis in the Japanese eel, Anguilla japonica." Virology
412(1): 179-187.
Moriyama, T., K. Terasawa, M. Fujiwara and N. Sato (2008). "Purification and characterization
of organellar DNA polymerases in the red alga Cyanidioschyzon merolae." FEBS J **275**(11):
2899-2918.
Ondov, B. D., G. J. Starrett, A. Sappington, A. Kostic, S. Koren, C. B. Buck and A. M. Phillippy
(2019). "Mash Screen: High-throughput sequence containment estimation for genome
discovery." bioRxiv: 557314.
Peretti, A., E. M. Geoghegan, D. V. Pastrana, S. Smola, P. Feld, M. Sauter, S. Lohse, M.
Ramesh, E. S. Lim, D. Wang, C. Borgogna, P. C. FitzGerald, V. Bliskovsky, G. J. Starrett, E. K.
Law, R. S. Harris, J. K. Killian, J. Zhu, M. Pineda, P. S. Meltzer, R. Boldorini, M. Gariglio and

C. B. Buck (2018). "Characterization of BK Polyomaviruses from Kidney Transplant Recipients Suggests a Role for APOBEC3 in Driving In-Host Virus Evolution." <u>Cell Host Microbe</u> **23**(5): 628-635 e627.

Pipas, J. M. (1992). "Common and unique features of T antigens encoded by the polyomavirus group." <u>J Virol</u> **66**(7): 3979-3985.

Pipas, J. M. (2019). "DNA Tumor Viruses and Their Contributions to Molecular Biology." <u>J Virol</u> **93**(9).

Rice, P., I. Longden and A. Bleasby (2000). "EMBOSS: the European Molecular Biology Open Software Suite." <u>Trends Genet</u> **16**(6): 276-277.

Santiago-Rodriguez, T. M., M. Ly, M. C. Daigneault, I. H. Brown, J. A. McDonald, N. Bonilla, E. A. Vercoe and D. T. Pride (2015). "Chemostat culture systems support diverse bacteriophage communities from human feces." <u>Microbiome</u> **3**: 58.

Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B. Schwikowski and T. Ideker (2003). "Cytoscape: a software environment for integrated models of biomolecular interaction networks." <u>Genome Res</u> **13**(11): 2498-2504.

Soding, J. (2005). "Protein homology detection by HMM-HMM comparison." <u>Bioinformatics</u> **21**(7): 951-960.

Tisza, M. J., D. V. Pastrana, N. L. Welch, B. Stewart, A. Peretti, G. J. Starrett, Y.-Y. S. Pang, A. Varsani, S. R. Krishnamurthy, P. A. Pesavento, D. H. McDermott, P. M. Murphy, J. L. Whited, B. Miller, J. M. Brenchley, S. P. Rosshart, B. Rehermann, J. Doorbar, B. A. Ta'ala, O. Pletnikova, J. Troncoso, S. M. Resnick, A. M. Segall and C. B. Buck (2019). "Discovery of several thousand highly diverse circular DNA viruses." 555375.

Vogt, W., P. Patzer, L. Lege, H. D. Oldigs and G. Wille (1970). "Synergism between phospholipase A and various peptides and SH-reagents in causing haemolysis." <u>Naunyn Schmiedebergs Arch Pharmakol</u> **265**(5): 442-454.

Williams, S. H., X. Che, J. A. Garcia, J. D. Klena, B. Lee, D. Muller, W. Ulrich, R. M. Corrigan, S. Nichol, K. Jain and W. I. Lipkin (2018). "Viral Diversity of House Mice in New York City." <u>MBio</u> **9**(2).

Woolford, L., A. Rector, M. Van Ranst, A. Ducki, M. D. Bennett, P. K. Nicholls, K. S. Warren, R. A. Swan, G. E. Wilcox and A. J. O'Hara (2007). "A novel virus detected in papillomas and carcinomas of the endangered western barred bandicoot (Perameles bougainville) exhibits genomic features of both the Papillomaviridae and Polyomaviridae." <u>J Virol</u> **81**(24): 13280-13290.

Yutin, N., V. V. Kapitonov and E. V. Koonin (2015). "A new family of hybrid virophages from an animal gut metagenome." <u>Biol Direct</u> **10**: 19.

Yutin, N., D. Raoult and E. V. Koonin (2013). "Virophages, polintons, and transpovirons: a complex evolutionary network of diverse selfish genetic elements with different reproduction strategies." <u>Virol J</u> **10**: 158.

Yutin, N., S. Shevchenko, V. Kapitonov, M. Krupovic and E. V. Koonin (2015). "A novel group of diverse Polinton-like viruses discovered by metagenome analysis." <u>BMC Biol</u> **13**: 95.

Zallot, R., N. O. Oberg and J. A. Gerlt (2018). "'Democratized' genomic enzymology web tools for functional assignment." <u>Curr Opin Chem Biol</u> **47**: 77-85.

Zimmermann, L., A. Stephens, S. Z. Nam, D. Rau, J. Kübler, M. Lozajic, F. Gabler, J. Söding, A. N. Lupas and V. Alva (2017). "A Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core." <u>J Mol Biol</u> **S0022-2836**(17): 30587-30589.