Metacontrast masking as a measure of change detection: Children with poor reading fluency show impaired change detection for both letters and shapes.

David Crewther[1], Jacqueline Rutkowski[1], Sheila Crewther[2]

[1]Brain Sciences Institute, Swinburne University of Technology, Victoria 3122, Australia

[2]School of Psychological Science, La Trobe University, Victoria 3086, Australia

Corresponding author:
David P Crewther
Centre for Human Psychopharmacology
Swinburne University of Technology
Melbourne, VIC 3122
Australia
Tel: +61 3 9214 5877
Fax: +61 3 9214 5525
e-mail: dcrewther@swin.edu.au

Abstract

Developmental dyslexia, a specific learning difficulty in reading, manifests as trouble in decoding words and as such is commonly associated with reduced phonemic awreness. However, its underlying cause remains elusive, with magnocellular visual processing, temporal auditory processing, visual attentional deficits and cerebellar dysfunction all gaining some traction, with integrative attempts identifying perceptual speed as a common factor. Reading in a crowded environment requires the suppression of distraction. More recent theories have concerned the visual span of attention, measuring the parallel attentive capacity of the sensory visual system. However the task as implemented used reports, both conscious recall and recognition of letters, that activate many cortical areas beyond sensory visual cortex. Change detection, on the other hand, does not require the conscious recognition of items, but simply awareness that the stimulus has changed, or not, again testing attention in a parallel fashion, but avoiding the complications of higher order cognitive processes. Thus, we investigated change detection in 33 good and poor readers with ages of around 10 yr, using a gap paradigm. Change detection performance for groups of letters or shapes, each item surrounded by an annular frame filled with dynamic random noise of variable contrast was manipulated by varying the contrast of these meta-contrast mask frames. Performance was measured as the threshold contrast of the frames at which participants could just detect change. In two separate experiments, letters and rectangular shapes were used as target items respectively, in order to test whether previous findings of superior change detection in good compared with poor readers was a result of greater automaticity in letter recognition of the good readers. The results indicate that the good readers were able to detect change at higher levels of masking distraction for both the letter and shape targets, indicating that this difference is not specifically related to to the training of graphemic or lexical information but more likely reflects a difference in alerting or pre-recognition stages of visual processing. Together, the results provide further support of the notion that there is a low level attentional performance difference between dyslexic and normal reading children. Thus, the results further bring transient spatial attention directly into the spotlight as an ability critical for learning to read.

Introduction

Developmental Dyslexia (DD) is an impairment in the acquisition of literacy skills despite normal intelligence, socioeconomic and educational circumstances, as defined under DSM-IV-TR (American Psychiatric Association, 2000), but positioned with specific learning disorder in DSM-5 (American Psychiatric Association, 2013). It is estimated to affect approximately 5-10% of school-aged children (Habib, 2000). These deficits tend to persist through to adulthood. Children with DD are reported to show deficits in many areas of perception, motor skills and learning but the primary deficit has long been considered to involve phonological processing (reviewed Bishop and Snowling (2004), Snowling and Hulme (2012)). Deficits in rapid temporal auditory processing have also been hypothesised as the basis of dyslexia by Merzenich et al. (1996), as dyslexics have difficulty in rapidly processing sequential information. Indeed, rapid temporal dot counting has also been shown to be more difficult for children with dyslexia than spatial dot counting (Eden et al., 1995). Other multisensory processing deficits have been demonstrated with both auditory and visual gap detection impaired in children with dyslexia compared with normal reading children of the same age (Van Ingelghem et al., 2001) indicate the possibility of a multisensory temporal processing deficit. More recent experiments employing stimuli designed to probe magnocellular activity have also identified perceptual speed as a common factor accounting for variance in reading ability (McLean et al., 2011).

Competing visual hypotheses also exist for the underlying mechanism of dyslexia. For example, dysfunction in the magnocellular visual pathway (Stein and Walsh, 1997;Habib, 2000;Gori et al., 2016) has been proposed on the basis of reduced motion and motion coherence sensitivity (Cornelissen et al., 1995;Talcott et al., 2000), and reduced brain activation in Area V5/MT+ (Eden et al., 1996) to moving stimuli with Demb et al. (1998) showing a 3-way interaction between brain activiation in MT+, thresholds for speed discrimination and reading speed.

Interest has also grown in visual processing of stimuli and abnormalities in transient attention. Indeed deficiencies in visual attention as the basis for dyslexia are supported by reports of slower serial search speeds and lowered search performance (Eskenazi and Diamond, 1983;Casco and Prunetti, 1996). Early work investigating the effect of backward masking (Grosser and Trzeciak, 1981) on the recognition of briefly presented single letters, demonstrated poorer performance in a group of dyslexic children compared with normals. Facoetti and Molteni (2001) showed that dyslexia ias associated with an alteration in the gradient of attention with consequent abnormalities in engagement and disengagament (Facoetti et al., 2008). Visual attention span (VA) was introduced as a means of measuring the proportion of distinct visual elements that can be correctly processed in parallel in a briefly presented multi-element array (Bosse et al., 2007). Bosse et al. proposed that a deficit in VA

span might contribute to developmental dyslexia, independently of abnormalities in phonological processing. This was supported through studies indicating a visual rather than verbal modality (Lobier et al., 2012). Also, fMRI studies using categorization of alphanumeric and non-alphanumeric character strings showed differential fMRI activation in the superior parietal lobule between participants with and without dyslexia (Lobier et al., 2014).

Other aspects of attentional processing, such as ability to detect change in the environment rapidly (change detection), have received relatively less attention, despite strong intergroup differences. Rutkowski et al. (2003) using a gap-paradigm change detection task, involving a stimulus array of 4 letters in circular place holders, showed that a much longer presentation time (an inspection time) for the initial group of letters was needed in order to just detect change, comparing dyslexic children with normal readers. There are some similarities between change detection and visual attention span, centred around the fact that for both tasks, several objects have to be processed in parallel. A key difference is that with change detection, the target objects do not have to be brought fully to conscious awareness to report change at better than chance level. VA span, on the other hand, requires identification and hence conscious perception of the objects. While change detection relies on the integration of the initial items in terms of depth of recognition, it is possible that the ability to detect change depends on a low level attentional function related to the sudden reappearance of the second group of items following the gap. If misdirection of attention provided by the placeholders around each object provides a masking signal, then this can potentially be used to measure change detection ability, independent of integration times for recognition of the initial display in the gap paradigm stmuli but rather as a function of ability to filter salient information by inhibiting distracting visual stimuli.

Thus we proposed to further investigate masking as a distracter in change detection, providing a novel thresholding mechanism using dynamic random noise of variable contrast in annular place holders as a variable meta-contrast masking of the second presentation stimulus of the gap change detection paradigm. This method of measuring change detection sensitivity (Rutkowski et al., 2003) was then used across letter and shape targets to establish whether previously reported change detection differences could be more attributable to better training in letter recognition in good readers rather than impairment in visual processing in those with dyslexia. The shape stimuli were used to test whether good readers, when presented with novel, untrained stimuli under similar task conditions, would continue to perform better than poor readers.

Methods

Participants

Thirty-three children were recruited from two non-selective state primary schools and from a summer camp for children with learning difficulties. Reading ability was assessed by trained experimenters using the The Neale Analysis of Reading Test (Neale, 1999). A reading age was established for each participant on the basis of the published norms. Assignment of participants to good and poor reading groups was on the basis of chronological age minus reading age

Two similar change detection tasks were carried out – one using detection of changes in letter identity (Expt 1a) and the other using change in orientation or colour of rectangular shapes (Expt 1b). Delay in the implementation of testing for Expt 1b resulted in overlapping but not identical populations for the two experiments.

Table 1. Participant Age and Reading Profiles for Experiments 1a and 1b

Stimulus	Reading Group	Chronological Age	Reading Age	N
Letter	Poor	10.9±0.6	7.7± 0.5	14
	Good	9.2±0.3	11.3±0.5	17
Shape	Poor	9.0±0.9	7.4±0.7	7
	Good	9.2±0.4	11.4±0.5	16

Change Detection for Letter Stimuli

In this experiment, a fixation cross appeared for a period of 1 sec, prior to the appearance of the first stimulus, which consisted of four upper case letters , each subtending a visual angle of 1° high x 0.8° wide and presented at an eccentricity of 2° for a participant seated 57 cm from the computer monitor (see Figure 1a). The presentation time for the first stimulus was fixed at 0.7 sec, followed by a fixed gap of 0.25 sec (with only the fixation cross showing). The second display was as for the first display (with a probablity of 50% that one item was changed), except that each letter was surrounded by an annulus of width 0.2 subtending 2° at the eye filled with dynamic random noise (2 pixel), the contrast of which could be varied from 0 to 1. Stimuli were created and presented using VPixx (www.vpixx.com) on an iMac computer (see Fig 1a). Participants were required (in a two alternative forced choice) to press the keyboard right arrow if they thought change had occurred and keyboard left arrow if they thought the two displays were identical. Convergence to threshold for the dynamic random dot noise was under the control of PEST model. Participants performed 40 trials.

Figure 1: Experimental stimuli used in experimentas 1a and 1b. Both employed annular place holders around the 4 stimuli of the second presentation, with variable contrast random dynamic noise (shown at high contrast for the letter stimuli in Fig 1a, and low contrast in the shape stimuli for Fig 1b).

Change Detection for Shape Stimuli

T\he shape experiment was identical to the letter experiment, except that shapes were substituted for letters. The shapes were rectangles oriented either horizontally or vertically and coloured either red or blue (see Fig. 1b). With 50% probability one of the shapes could change its orientation, its colour, or both, with conditions randomised. The shapes subtended 1° x 0.5° while the circular place holders subtended 2° at the eye of the participant seated 57 cm from the computer monitor. Stimuli were created and presented using VPixx on an iMac computer. Background luminance for both experiments was 50 cd/m^2.

Results

Letters

Good readers performed signficantly better than poor readers on the change detection for letters task. This can be seen in Figure 2 which shows the mean threshold contrast (with standard errors) for the

masking annuli. As the contrast of the flickering mask increases it becomes harder to detect change, and hence the contrast threshold repesents a measure of change detection ability. Analysis of variance (ANOVA) was carried out on the threshold masking contrast compared between reading groups, finding a significant main effect of Reading Group ($F(1, 29) = 8.35$, $p = 0.007$).

Figure 2: Threshold masking contrasts for letter and shape stimuli, for which groups of good and poor readers (GR, PR) could just detect change. The data are presented as means (with standard errors). Contrast thresholds for both letter and shape stimuli were higher for Good *cf* Poor reader groups.

Experiment 1b: Shapes

Good readers performed better than Poor readers on the change detection for shapes as well (see Figure 2). Again, higher contrast indicates a better ability to withstand the distraction caused by the appearance of the rings. Despite the smaller population size of Poor readers in the Shapes experiment, the effect of reading group was again significant $F(1, 22) = 5.75$, $p = 0.02$.

The similarity in threshold results for the two tasks suggests the degree of correlation be investigated. Taken across all readers, the correlation between threholds for letters and shapes was significant ($F(1,20) = 4.77$, *p*=.04, *Rsq*=.19).

Discussion

The results of these psychophysical experiments validate the use of variable masking as a measure of change detection, as the results replicate those established using different methods of measurement (titrating the initial display presentation time to estimate detection threshold) (Rutkowski et al., 2003). The metacontrast masking method has the advantage that it opens up other possibilities for investigating the role of attention in change detection, particularly as it applies to the mechanisms of dyslexia. It has the benefit that a wide dynamic range of distraction is available – especially using the

random dot noise which is qualitatively much more effective at high contrast than a static noise ring of the same geometrical dimensions and contrast. The fact that mask contrast can be used in such a controllable fashion to modify detection rate also suggests that the sudden appearance of the second stimulus (see Fig.1) may interfere with change detection precisely because attention is distributed across the field of possible targets. The results of this experiment reinforce the idea (Rutkowski et al., 2003) that the sudden appearance of the second stimulus is likely to be the the factor that controls change blindness, for this type (display – gap – display) of change experiment.

A common criticism of studies in dyslexia that use letters as stimuli is that letter processing by poor readers may be deficient because they have generally read less and practised letter recognition less. This criticism provides the rationale for examining change detection using shapes. The second experiment, despite being less statistically powered, due to a smaller PR population, showed that when geometrical shapes were used as targets, change detection performance was still inferior for poor readers compared with good readers.

The significance of the difference in change detection between good fluent and poor nonfluent readers suggests that rapidity of object processing might play an important role in the development of fluent reading (see (Grosser and Trzeciak, 1981), irrespective of need for phonological awareness of various letters and words. This is borne out by recent experiments associating perceptual speed with reading ability (Hecht et al., 2004;Thomson et al., 2006;McLean et al., 2011). The current findings are hard to reconcile with Goswami's opinion piece (2015) because while letters may have been learned better by those with good reading skills, the spatial arrangement of the change detection task is something that has not been trained, and yet the change detection performance on the shapes task was similarly impaired in those with poor reading skills.

Acknowledgements

The authors would like to acknowledge support from the philanthropic Andrew Fildes Foundation for sponsoring summer camps, the Department of Education and Training Victoria, and an Australian Research Council Discovery Project.

References

American psychiatric association (2000). *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR* Washington, USA: American Psychiatric Association.

American psychiatric association (2013). *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Arlington, VA, USA: American Psychiatric Association.

Bishop, D.V., and Snowling, M.J. (2004). Developmental dyslexia and specific language impairment: same or different? *Psychol Bull* 130, 858-886.

Bosse, M.L., Tainturier, M.J., and Valdois, S. (2007). Developmental dyslexia: the visual attention span deficit hypothesis. *Cognition* 104, 198-230.

Casco, C., and Prunetti, E. (1996). Visual search of good and poor readers: effects with targets having single and combined features. *Percept Mot Skills* 82, 1155-1167.

Cornelissen, P., Richardson, A., Mason, A., Fowler, S., and Stein, J. (1995). Contrast sensitivity and coherent motion detection measured at photopic luminance levels in dyslexics and controls. *Vision Res* 35, 1483-1494.

Demb, J.B., Boynton, G.M., and Heeger, D.J. (1998). Functional magnetic resonance imaging of early visual pathways in dyslexia. *J Neurosci* 18, 6939-6951.

Eden, G.F., Stein, J.F., Wood, H.M., and Wood, F.B. (1995). Temporal and spatial processing in reading disabled and normal children. *Cortex* 31, 451-468.

Eden, G.F., Vanmeter, J.W., Rumsey, J.M., Maisog, J.M., Woods, R.P., and Zeffiro, T.A. (1996). Abnormal processing of visual motion in dyslexia revealed by functional brain imaging. *Nature* 382, 66-69.

Eskenazi, B., and Diamond, S.P. (1983). Visual exploration of non-verbal material by dyslexic children. *Cortex* 19, 353-370.

Facoetti, A., and Molteni, M. (2001). The gradient of visual attention in developmental dyslexia. *Neuropsychologia* 39, 352-357.

Facoetti, A., Ruffino, M., Peru, A., Paganoni, P., and Chelazzi, L. (2008). Sluggish engagement and disengagement of non-spatial attention in dyslexic children. *Cortex* 44, 1221-1233.

Gori, S., Seitz, A.R., Ronconi, L., Franceschini, S., and Facoetti, A. (2016). Multiple Causal Links Between Magnocellular-Dorsal Pathway Deficit and Developmental Dyslexia. *Cereb Cortex* 26, 4356-4369.

Goswami, U. (2015). Visual attention span deficits and assessing causality in developmental dyslexia. *Nat Rev Neurosci* 16, 225-226.

Grosser, G.S., and Trzeciak, G.M. (1981). Durations of recognition for single letters, with and without visual masking, by dyslexics and normal readers. *Percept Mot Skills* 53, 991-995.

Habib, M. (2000). The neurological basis of developmental dyslexia: an overview and working hypothesis. *Brain* 123 Pt 12, 2373-2399.

Hecht, R., Crewther, D., and Crewther, S. (2004). Rate of learning and asymptotic performance in an automatization task and the relation to reading. *Perceptual and motor skills* 99, 1103-1121.

Lobier, M., Zoubrinetzky, R., and Valdois, S. (2012). The visual attention span deficit in dyslexia is visual and not verbal. *Cortex* 48, 768-773.

Lobier, M.A., Peyrin, C., Pichat, C., Le Bas, J.F., and Valdois, S. (2014). Visual processing of multiple elements in the dyslexic brain: evidence for a superior parietal dysfunction. *Front Hum Neurosci* 8, 479.

Mclean, G.M., Stuart, G.W., Coltheart, V., and Castles, A. (2011). Visual temporal processing in dyslexia and the magnocellular deficit theory: the need for speed? *J Exp Psychol Hum Percept Perform* 37, 1957-1975.

Merzenich, M.M., Jenkins, W.M., Johnston, P., Schreiner, C., Miller, S.L., and Tallal, P. (1996). Temporal processing deficits of language-learning impaired children ameliorated by training. *Science* 271**,** 77-81.

Neale, M.D. (1999). *Neale analysis of reading ability: manual:* . Melbourne.: ACER Press.

Rutkowski, J., Crewther, D., and Crewther, S. (2003). Change detection is impaired in children with dyslexia. *Journal of Vision* 3**,** 95-105.

Snowling, M.J., and Hulme, C. (2012). Annual research review: the nature and classification of reading disorders--a commentary on proposals for DSM-5. *J Child Psychol Psychiatry* 53**,** 593-607.

Stein, J., and Walsh, V. (1997). To see but not to read; the magnocellular theory of dyslexia. *Trends Neurosci* 20**,** 147-152.

Talcott, J.B., Hansen, P.C., Assoku, E.L., and Stein, J.F. (2000). Visual motion sensitivity in dyslexia: evidence for temporal and energy integration deficits. *Neuropsychologia* 38**,** 935-943.

Thomson, B., Crewther, D.P., and Crewther, S.G. (2006). Wots that werd? Pseudowords (non-words) may be a misleading measure of phonological skills in young learner readers. *Dyslexia* 12**,** 289-299.

Van Ingelghem, M., Van Wieringen, A., Wouters, J., Vandenbussche, E., Onghena, P., and Ghesquiere, P. (2001). Psychophysical evidence for a general temporal processing deficit in children with dyslexia. *Neuroreport* 12**,** 3603-3607.