Critical Role for the Unique N-Terminus of Chlamydial MreB in Directing Its Membrane Association and Interaction with Elements of the Divisome

Junghoon Lee[1], John V. Cox[2], and Scot P. Ouellette[1]

[1] Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE, USA 68198

[2] Department of Microbiology, Immunology, and Biochemistry, University of Tennessee Health Science Center, Memphis, TN, USA 38163

Abstract

Chlamydiae lack the conserved central coordinator protein of cell division FtsZ, a tubulin-like homolog. Current evidence indicates *Chlamydia* uses the actin-like homolog, MreB, to substitute for the role of FtsZ. Interestingly, we observed MreB as a ring at the septum in dividing cells of *Chlamydia*. We hypothesize that, for MreB to substitute for FtsZ in *Chlamydia*, it must possess unique properties compared to canonical MreB homologs. Sequence differences between chlamydial MreB and homologs in other bacteria revealed that chlamydial MreB possesses an extended N-terminal region as well as a conserved amphipathic helix found in other bacterial MreBs. The extended N-terminal region is sufficient to restore the localization of a truncated MreB mutant of *E.coli* lacking its amphipathic helix to the membrane and is crucial for interactions with cell division components RodZ and FtsK, though the region is not required for homotypic interactions. In addition, the N-terminal region was sufficient to direct GFP to the membrane when expressed in *Chlamydia*. From these data, the extended N-terminal region of chlamydial MreB is critical for correct localization and interactions of this protein. Our data provide mechanistic support for the ability of chlamydial MreB to serve as a substitute for FtsZ.

Word Count: 197/200

Importance

Chlamydia is an obligate intracellular bacterial pathogen, which causes sexually transmitted diseases and trachoma. A better understanding of basic chlamydial physiology is important for developing novel therapeutic strategies for these diseases. Chlamydiae divide by a unique polarized cell division process not requiring the canonical cell division coordinator FtsZ. We previously demonstrated a role for MreB in directing chlamydial division. We hypothesize chlamydial MreB encodes unique features allowing it to substitute for FtsZ in this organism. Here, we performed a detailed characterization of chlamydial MreB as a cell division coordinator. Chlamydial MreB forms a ring at the septum, like FtsZ in *E.coli*. It possesses an extended N-terminus that is conserved amongst *Chlamydiaeceae*. This region displays amphipathicity that is sufficient to drive membrane localization of distinct proteins. In addition, this region is crucial for interactions with other division proteins. We propose that this N-terminal region confers unique properties to chlamydial MreB.

Word count: 149/150

Introduction

Chlamydia is an obligate intracellular pathogen that causes diverse diseases. These Gram-negative cocci differentiate between two morphologically and functionally distinct cellular forms during their developmental cycle: the elementary body (EB) and the reticulate body (RB)(1). The EB is the infectious but non-dividing form whereas the RB is the dividing but non-infectious form. After entering the host cell, *Chlamydia* remains within a membrane-bound parasitic organelle, termed an inclusion(2). RBs undergo multiple rounds of cell division until such point they engage a secondary differentiation program and convert to EBs, which are subsequently released from the host cell to propagate the infection.

In evolving to obligate intracellular dependence, *Chlamydia* has significantly reduced its genome and eliminated *ftsZ*(3), encoding the conserved tubulin-like cell division coordinator of binary fission(4). However, *Chlamydia* encodes rod-shape determining proteins associated with peptidoglycan synthesis in the lateral cell wall of bacilli. That *Chlamydia* has retained these genes is unusual since these are coccoid bacteria and only synthesize peptidoglycan during division(5). The absence of FtsZ in chlamydiae suggest they might not divide by binary fission. Recent work from our labs supports this. We employed both live-cell and fixed-cell imaging to localize various markers, including a cell division protein (FtsQ(6)), during EB-to-RB differentiation and the first division of the RB. Interestingly, we observed that chlamydiae are highly polarized throughout this time and that division itself occurs via a polarized process(7). This polarized budding process remains the primary mode of division even at later times during the developmental cycle (Cox et al., *in prep*).

In 2012, we hypothesized and presented evidence that *Chlamydia* co-opted the rod-shape determining protein MreB to function as the central coordinator of division(8). To perform this function, we hypothesize that chlamydial MreB possesses unique properties.

Many MreB homologs encode an N-terminal amphipathic helix that facilitates MreB

association with the inner membrane(9). At the membrane periphery it engages PG synthesis

along short moving filaments(10-12). Here, we investigated the unique structural features of

chlamydial MreB that allow it to substitute for FtsZ. Chlamydial MreB forms rings at the

division site, and bioinformatics analysis indicates that chlamydial MreB encodes an extended

N-terminal region (amino acids 1-23: aa1-23) in addition to the conserved amphipathic helix.

The aa1-23 residues also encode predicted amphipathicity. We confirmed that the conserved

amphipathic helix, but not aa1-23, is important for directing membrane-localization of GFP.

Nevertheless, when aa1-23 of chlamydial MreB was fused to a cytosolic *E. coli* MreB lacking

its endogenous amphipathic helix, *E. coli* MreB was redirected to the membrane. Importantly,

aa1-23 were sufficient to direct GFP to the membrane in *Chlamydia* directly. We determined

the unique N-terminal region of chlamydial MreB is crucial for its interactions with proteins

related to cell division but not its ability to form homo-oligomers. Our data indicate the unique

N-terminal region of chlamydial MreB is important for membrane localization and interaction

with the divisome. These properties may allow it to function as a central organizer of the cell

division machinery in *Chlamydia* and in a manner distinct from other MreB homologs.

Results

Chlamydial MreB_6xH localizes as a ring at the division site. Given our hypothesis for the

role of chlamydial MreB in directing polarized division, we examined its localization during

the first division of an RB. We were unable to detect endogenous MreB using the antibody

previously reported to image the distribution of MreB in *Chlamydia*(13). We then attempted to

image wild-type and various truncation mutants of MreB using a GFP sandwich (SW) fusion

strategy that, in *E. coli*, was reported to be functional(14) (N- or C-terminal fluorescent protein

fusions with MreB display an artifactual localization(15)). However, when we inducibly

expressed MreB_GFP$_{SW}$ in *C. trachomatis* L2, bacterial growth was significantly inhibited,

even with low levels of expression, as observed by the presence of a small number of enlarged

bacteria (Suppl. Fig. 1). We, therefore, opted to construct an MreB fusion with a small C-

terminal hexahistidine tag (MreB_6xH). The human epithelial cell line, HeLa, was infected

with a transformant of *C. trachomatis* L2 carrying a plasmid to inducibly express MreB_6xH.

At 6 hours post-infection (hpi), expression of MreB_6xH was induced with

anhydrotetracycline (aTc), and cells were fixed at 10.5 hpi, a time when the nascent RB has

begun its first division(7). As seen in Figure 1, we observed using super-resolution structured

illumination microscopy (SIM) the polar localization of the major outer membrane protein

(MOMP) in the budding daughter cell as previously reported(7). Consistent with its proposed

role in division(8), MreB_6xH was localized in a band-like structure across the septum and not

in puncta as previously reported(13, 16). Interestingly, when 3D reconstructions were

assembled from the SIM images, we observed that MreB_6xH formed clear ring-like structures

at the septum.

Chlamydial MreB encodes an amphipathic helix and an extended N-terminal region

conserved in *Chlamydiae*. In many Gram-negative bacteria, there is an amphipathic helix at

the N-terminus of MreB, and the helix is important for the membrane localization of MreB(9).

This localization is crucial for the role of MreB in directing PG synthesis during lateral cell

wall growth(9, 17). To confirm whether chlamydial MreB also has these features, we predicted

its secondary structure and amphipathic regions using bioinformatics tools (Fig. 2). The

alignment data show that chlamydial MreB encodes an extended N-terminal region (amino

acids 1-23: aa1-23), which other Gram-negative bacteria lack (Fig. 2A). Importantly, this

extended N-terminal region is conserved amongst *Chlamydiae* (Suppl. Fig. 2). Following this

region, there is a predicted amphipathic helix, from R24 to F29 (Fig. 2B), that aligns with the

amphipathic helices of the other bacteria (Fig. 2A). A high amphipathic score (A-score) extends

to residue F32 in chlamydial MreB. Interestingly, aa1-23 also has a high A-score though the

region was not predicted to form an amphipathic helix (Fig. 2B). Nevertheless, performing a

helical wheel analysis with 8 amino acid windows revealed that each window shows

amphipathicity (Fig. 2C). Based on these observations, we conclude that the extended N-

terminal region of chlamydial MreB may possess membrane-associating properties.

Chlamydial MreB is unable to complement an *mreB*-deficient mutant of *E.coli*. The

bioinformatics data suggested that chlamydial MreB encodes unique structural features absent

in other bacterial MreBs (Fig. 2A). To assess the activities associated with these unique

structural elements, we introduced chlamydial MreB into an *mreB*-deficient mutant of *E.coli*.

As *mreB* is an essential gene in *E.coli*, Bendezu et al. made the strain P2733, which has a

deletion in the *mreB* gene and is conditionally viable by the overexpression of the *ftsQAZ*

operon(18). This strain has been used for complementation assays and grows as coccoid

bacteria (since MreB is necessary for rod-shape)(18). To test whether chlamydial MreB

(CtrMreB) is capable of complementing the *mreB* deficient mutant, P2733 was transformed

with an arabinose-inducible vector encoding chlamydial *mreB*. Cell shape was compared after

inducing MreB expression in comparison to an empty vector negative control and an *E. coli*

MreB (EcMreB) positive control. When EcMreB is expressed from an arabinose-inducible

promoter in the P2733 strain, the *E. coli* cells begin to adopt a rod shape (Fig. 3B), consistent

with other reports(14, 18, 19). When chlamydial MreB was induced, the morphology of P2733

was unchanged compared to uninduced or empty vector controls, indicating that chlamydial

MreB does not complement the *mreB*-deficient mutant of *E.coli* (Fig. 3C). To determine

whether the inability of chlamydial MreB to complement was due to its extended N-terminal

region, we performed the same experiment with ΔN22 chlamydial MreB, which more closely

resembles *E. coli* MreB in size and characteristics at the N-terminus. We observed that ΔN22

chlamydial MreB also does not complement the *mreB* deficient *E.coli*, suggesting that the

extended N-terminal region is not the only reason for the failure to complement (Fig. 3D – see

also Discussion). Chlamydial MreB expression was confirmed in these experiments by western

blotting (Suppl. Fig. 3).

The amphipathic helix, but not the extended N-terminal region, of chlamydial MreB is

sufficient to direct GFP to the membrane in *E. coli*. We hypothesized that, because of their

amphipathicity, the unique sequence elements at the N-terminus of chlamydial MreB could

direct the membrane localization of MreB (Fig. 1). To test this, we performed a series of

localization studies in *E. coli* using the N-terminal regions of chlamydial MreB (CtrMreB)

fused to GFP. A similar strategy was used to demonstrate that two copies of the amphipathic

helix of *E. coli* MreB (EcMreB) were sufficient to direct GFP to the membrane(9). We made

various fusions in which portions of the N-terminal region of chlamydial MreB were fused to

GFP either in one copy or two (illustrated in Fig. 4A). We then observed the localization of

these proteins in both wild-type (MG1655) and MreB-deficient (P2733) *E.coli* (Fig. 4B and

Suppl. Fig. 4). We observed that a single copy of aa1-32 ($MreB_{1-32}$), encoding all high A-score

residues, was sufficient to direct GFP to the membrane. Interestingly, aa1-28 ($MreB_{1-28}$),

encoding both the predicted amphipathic helix and the extended N-terminus, was also sufficient

to direct GFP to the membrane, but, in wild-type cells, its localization was primarily restricted

to the poles. Consistent with *E. coli*, two copies of aa23-32 ($MreB_{23-32}$), encoding the predicted

amphipathic helix, directed GFP to the membrane. Contrary to our prediction, one or two copies

of aa1-23-GFP ($MreB_{1-23}$-GFP) was not able to direct GFP to the membrane even though the

region has a high A-score (Fig. 2B).

The extended N-terminus of chlamydial MreB is sufficient to direct an *E. coli* MreB

lacking its amphipathic helix to the membrane. Although aa1-23 of CtrMreB were not able

to direct GFP to the membrane, we speculated that its membrane-associating properties may

be revealed when fused directly to an EcMreB truncation mutant lacking its amphipathic helix

(ΔN7aa EcMreB), which fails to localize at the membrane(9). To test this, we fused the N-

terminal region of chlamydial MreB to ΔN7aa EcMreB_GFP (illustrated in Fig. 4D) and

observed the localization of these constructs in *E. coli* (Fig. 4E). Wild-type EcMreB_GFP is

localized to the membrane whereas ΔN7aa EcMreB_GFP appears as aggregates in the

cytosol(9). Both chlamydial aa1-23 and aa1-28 clearly redirected ΔN7aa EcMreB_GFP to the

membrane, suggesting that these sequence elements possess amphipathic features sufficient to

direct the membrane localization of an MreB homolog (Fig. 4E). From these data, we conclude

that the amphipathic helix is important for the membrane localization of MreB and aa1-23 may

confer additional membrane binding affinities and/or properties to chlamydial MreB.

The extended N-terminal region of chlamydial MreB is sufficient to direct GFP to the

membrane in *C. trachomatis* L2. Given the ability of aa1-23 of chlamydial MreB to direct

ΔN7aa EcMreB_GFP to the membrane in *E. coli*, we next asked the question whether this

region, alone or in combination with the amphipathic helix (aa24-28), could direct GFP to the

membrane directly in *Chlamydia*. The MreB$_{1-23aa}$_GFP or MreB$_{1-28aa}$_GFP fusions were moved

to an inducible chlamydial expression plasmid and transformed into *C. trachomatis* L2. These

transformants were then used to infect HeLa cells, and construct expression was induced at

either 6 or 16 hpi. Infected cells were fixed and imaged at 10.5 or 20 hpi, respectively (Fig. 5

with additional images in Suppl. Fig. 5). GFP alone is a cytosolic protein when expressed in

chlamydiae (Fig. 5A). For both fusion proteins, GFP fluorescence was observed at membrane

sites (Fig. 5B&C with arrowheads in 5B indicating individual bacteria with membrane

localization of GFP) at both time points examined. We conclude from these data that the

extended N-terminal region of chlamydial MreB is sufficient to direct GFP to the membrane in

Chlamydia.

An N-terminal truncation of MreB shows reduced membrane-localization in *Chlamydia*.

To further investigate the role of the N-terminal region of MreB in *Chlamydia*, we constructed three truncations each fused to a C-terminal hexahistidine tag: one lacking the additional N-terminal region (ΔN22), one lacking both the N-terminal region and the amphipathic helix (ΔN28), and one lacking all high A-score residues as described in Figure 2 (illustrated in Suppl. Fig. 6A). Each MreB_6xH was inducibly expressed directly in *C. trachomatis* L2. Only the ΔN32 MreB_6xH truncation showed reduced membrane binding with more accumulation in the cytosol compared to the other constructs (Suppl. Fig. 6B). An important caveat to these studies is that the endogenous chromosomally encoded MreB is still present as it is not currently possible to generate a reliable conditional knockout in *Chlamydia* (see Discussion).

The extended N-terminal region is dispensable for MreB homo-oligomerization but essential for interactions with other division proteins. Given the caveat noted above, we next asked whether the N-terminal region is necessary for MreB homo-oligomerization. In *E.coli*, MreB interacts with itself to form short filaments(10). In addition, since MreB participates in both cell shape determination and cell division, it interacts with diverse proteins related to cell division and peptidoglycan synthesis(14, 19, 20). These features are shared with chlamydial MreB, which interacts with itself and with several proteins such as the cytoskeletal protein RodZ and the cell division components FtsK and FtsQ(6, 8, 21). Based on these previous reports, we hypothesized that the extended N-terminal region is important for these interactions. To test this, we used the bacterial adenylate cyclase-based two hybrid (BACTH) assay, which is based on the reconstitution of enzyme activity by two interacting proteins that bring catalytic adenylate cyclase fragments (T25 and T18) into close proximity. We designed BACTH constructs encoding truncated MreB mutants and then performed BACTH assays (Fig. 6). The N-terminal truncations of MreB interacted with full-length MreB (Fig. 6A and summarized in B), providing a potential explanation for the localization data of MreB_6xH

truncations in *Chlamydia* (Suppl. Fig. 6). However, these truncated MreBs did not interact with

RodZ, full-length FtsK, or the N-terminal region of FtsK lacking its ATPase domain whereas

full-length MreB did interact with these constructs (Fig. 6C and summarized in D)(8, 21). From

these data, we conclude that the extended N-terminal region is dispensable for MreB homo-

oligomerization but is critical for interacting with other proteins related to cell division.

Discussion

MreB is a well characterized rod-shape determining protein, which is conserved in most

bacilli(18). For example, when the *mreB* gene is deleted, the cell shape is changed from bacillus

to coccoid in *E.coli*(18). MreB participates primarily in organizing PG synthesis by interacting

with other proteins, such as MreC and RodZ, to direct Pbp2 and RodA activity at the

membrane(19, 22). Recently, MreB has been observed as a dynamic cytoskeletal protein

because it forms short filaments and rotates around the membrane perpendicularly to the

longitudinal axis of the cell(11). This motion is driven by PG synthesis and is critical for cell

elongation and maintenance of rod shape(23). Given these characteristics and properties of

MreB, it is surprising that *Chlamydia*, a Gram-negative coccoid bacterium that lacks PG in its

cell wall, encodes MreB. This suggests that the maintenance of cell morphology may not be

the primary function of chlamydial MreB(5, 24). Rather, given the function of chlamydial PG

strictly in cell division, chlamydial MreB may function in this process.

Since *Chlamydia* does not have the conserved cell division organizing protein FtsZ, we have

hypothesized and presented evidence that MreB is a functional substitute for FtsZ(6-8, 21).

However, this raises the intriguing question of how MreB, which is highly conserved between

Chlamydia and phylogenetically unrelated bacteria, can serve as the divisome organizer.

Interestingly, when MreB is reintroduced into MreB-depleted cells in *B. subtilis*, a budding

shape is observed that is enriched in MreB(25). This eventually leads to recruitment of cell

wall machinery and formation of a rod-shape. However, in *Chlamydia*, which also undergoes

a budding-like polarized division, this process is spatially restricted, thus what prevents

Chlamydia from continuing to synthesize PG to produce a rod shape? One hypothesis is that

chlamydial MreB exhibits a restricted distribution that is at least in part dependent upon its

unique structural features, and we have shown here that the unique N-terminal region of

chlamydial MreB can facilitate the membrane association of this cytoskeletal protein.

To initiate this study, we attempted to localize endogenous chlamydial MreB using an antibody

previously described(13). Prior reports suggested that MreB is localized as puncta at sites of

division(13, 16). Our goal was to capture MreB localization at high resolution and at early

stages of division since it becomes more difficult to resolve individual organisms within the

inclusion as RBs multiply. However, we were unable to detect endogenous MreB using the

previously reported antibody. We, therefore, took advantage of genetic tools that have been

recently developed for *Chlamydia* to transform this bacterium with a plasmid construct

encoding an inducible MreB_6xH. Using this strategy, we detected MreB_6xH localized at the

division septum during the polarized division of the first RB. More interestingly, MreB formed

a ring at the septum in the dividing cell that resembled the FtsZ-ring at the septum of *E. coli*

(Fig.1). The septal MreB ring is very similar to the PG ring previously observed in

Chlamydia(5). We speculate that the MreB puncta previously observed may be due to the low

affinity of the MreB antibody as we were also unable to detect MreB at later stages during

infection using this reagent. Given the effects of inhibiting MreB activity on chlamydial

division and PG synthesis(7, 8, 16), these data further suggest that chlamydial MreB likely has

a scaffolding function for cell divisome proteins to direct PG synthesis in a manner similar to

the function of FtsZ in *E.coli*(26).

To further investigate how chlamydial MreB may substitute for FtsZ in *Chlamydia*, we

performed a simple alignment of MreB homologs from *Chlamydia* species and other rod-shape

bacteria. Many Gram-negative MreB homologs possess an amphipathic helix at their N-

terminus that allows MreB to associate with the inner membrane(9). *Chlamydia* species have

conserved this feature. The most striking difference between the chlamydial species and other

"canonical" MreB homologs is the presence of an extended N-terminus of 23 amino acids (aa1-

23). Amphipathic prediction algorithms suggested this region possesses amphipathicity.

Interestingly, canonical MreB filaments are excluded from areas of high membrane curvature

in bacilli that are typically enriched in anionic phospholipids such as cardiolipin (i.e. the cell

poles(27)), but *Chlamydia* are coccoid with its membrane displaying curvature similar to the

poles of bacilli. Therefore, we hypothesized that the additional residues at the N-terminus of

chlamydial MreB may allow it to associate more efficiently with membranes that exhibit

curvature. In addition, these residues may be critical for establishing polarity in this organism

as we previously observed that inhibiting MreB activity with A22 or MP265 not only blocks

cell division but also depolarizes the RB(7, 8).

The amphipathic helix of canonical Gram-negative MreB homologs is critical for the

association of MreB with membranes. This was demonstrated by two means: expressing two

copies of the helix in tandem fused to the N-terminus of GFP and deleting the helix from MreB

and assessing its localization(9). In the former, GFP was directed to the inner membrane(9). In

the latter, MreB is no longer localized as membrane patches but rather as cytosolic aggregates

in *E. coli*(9). Using the former approach, we observed that two copies, but not one copy, of the

chlamydial amphipathic helix were sufficient to direct MreB to the membrane, consistent with

what has been observed in *E. coli* (Fig. 4B). However, the additional N-terminal region of

chlamydial MreB (aa1-23) together with a single copy of the helix (aa24-28) directed GFP to

the membrane. Intriguingly, the membrane localization was restricted to the cell polar region

but did not appear as an inclusion body. These data suggest that the additional N-terminal

region (aa1-23) can direct membrane localization and may possess affinity for regions of high

membrane curvature. Yet, this region alone or in two copies was not sufficient to direct GFP to

the membrane. This may indicate that the membrane affinity of this N-terminal region is

context specific. Therefore, we tested if it was capable of redirecting an *E. coli* truncation

mutant of MreB, lacking its amphipathic helix, to the membrane. This was the case. Therefore,

we conclude that the additional N-terminal region of chlamydial MreB encodes a membrane-

targeting function. Indeed, this was supported by the ability of this region to direct GFP to the

membrane when expressed in *Chlamydia* directly. We propose that chlamydial MreB, with both

its extended N-terminus and amphipathic helix, may be more tightly associated with

membranes and is critical for it to form ring structures associated with regions of high

membrane curvature. We are currently working to prepare recombinant protein of full-length

and truncated chlamydial MreB to test this in liposomes.

As *Chlamydia* is an obligate intracellular bacterium, it is time-consuming and difficult to

genetically modify its genes in a targeted manner, and there is a high failure rate in doing so.

Ideally, we would create a conditional knockout such that we could express truncated mutants

of chlamydial MreB in the absence of the chromosomal full-length copy. We did attempt to

localize truncation mutants of MreB in *Chlamydia*, but only the mutant lacking all predicted

amphipathicity showed a different profile. The truncations lacking the extended N-terminus

(aa1-23) and both that region and the amphipathic helix (aa1-28) still showed membrane

localization (Suppl. Fig. 6). While the mutant lacking all predicted amphipathicity (aa1-32)

appeared more cytosolic than the other constructs, some signal was detected at the membrane.

Not surprisingly, we detected interactions between the MreB truncations and the full-length

MreB, thus one interpretation is that the truncations, when expressed in *Chlamydia*, interacted

with full-length MreB (Fig. 6), which is itself at the membrane. In *E.coli*, the residues, V121

and S284, participate in forming the protofilament of MreB, and chlamydial MreB possesses

similar residues, V139 and A302, respectively(28). Furthermore, the N-terminal region is

important for interactions with other proteins related to cell division and PG synthesis. When

the region was deleted, chlamydial MreB no longer interacted with RodZ and FtsK by bacterial

two-hybrid analysis. One possible explanation of these data is that the truncated MreB isoforms

do not associate with the membrane efficiently enough to interact with the membrane proteins

RodZ and FtsK. Nevertheless, these data suggest the extended N-terminal region may function

as a scaffolding platform for MreB to interact with other proteins related to cell division and

may be crucial for the cell divisome formation at the septum, like FtsZ.

The sequence similarity between chlamydial MreB and *E.coli* MreB is 71%, and we previously

observed that these homologs will interact by two-hybrid analysis(6, 21). Therefore, we tested

whether chlamydial MreB could functionally complement an *E. coli mreB* deletion strain.

Given the differences at the N-terminus and the role of MreB as part of a multicomponent

machinery that directs cell wall synthesis, it is perhaps not surprising that chlamydial MreB

failed to complement an *E. coli mreB* mutant strain even when the additional N-terminal region

was removed (Fig. 3). To date, we have been unable to complement any structural components

of the *E. coli* divisome (e.g. FtsQ(6)) or Mre system (MreB or RodZ(21)), which reinforces the

importance of recapitulating all necessary interactions to efficiently complement. In contrast,

cytosolic components of the PG synthesis pathway possessing enzymatic activity have been

shown to complement *E. coli* conditional mutants(29-31). These data are consistent with the

notion that the PG synthesis machinery is conserved in these organisms but the mechanisms by

which PG is used and directed are not.

Outstanding questions remain: how is polarity established in *Chlamydia*? Is it MreB-dependent

or are other factors involved? Is MreB actively excluded from other parts of the membrane by

an unknown system similar to the function of the MinCDE system in restricting FtsZ to division

septa during binary fission? These are active areas of investigation in our labs and are of great

interest to the community. In this study, we performed the first systematic investigation of

chlamydial MreB and its function as a cell division coordinator. Our data suggest that

chlamydial MreB has multiple membrane-associating domains that may promote its assembly

at the septum. This in turn may allow it to functionally substitute for the role of FtsZ in

organizing the division septum.

Materials and Methods

Organisms and Cell Culture

McCoy (kind gift of Dr. Harlan Caldwell) and HeLa (ATCC, Manassas, VA) cells were cultured

at 37°C with 5% CO_2 in Dulbecco's Modified Eagle Medium (DMEM; Invitrogen, Waltham,

MA) containing 10% fetal bovine serum (FBS; Hyclone, Logan, UT) and 10 μg/mL gentamicin

(Gibco, Waltham, MA). *Chlamydia trachomatis* serovar L2 (strain 434/Bu) lacking the

endogenous plasmid (-pL2) was infected and propagated in McCoy cells for use in

transformations. HeLa cells were infected with chlamydial transformants in DMEM containing

10% FBS, 10 μg/mL gentamicin, 1 U/mL penicillin G, and 1 μg/mL cycloheximide. All cell

cultures and chlamydial stocks were routinely tested for *Mycoplasma* contamination using the

Mycoplasma PCR detection kit (Sigma, St. Louis, MO). For *E.coli*, wild-type MG1655 and the

mreB-deficient strain P2733 (kind gift of Dr. Piet de Boer) were cultured at 37°C with 225 rpm

shaking in Lysogeny broth (LB) media containing no or several antibiotics as indicated: 50

μg/mL spectinomycin, 34 μg/mL chloramphenicol, or 25 μg/mL tetracycline. All chemicals

and antibiotics were obtained from Sigma unless otherwise noted.

Cloning

The plasmids and primers used in this study are described in Supplemental Table 1. The wild-type and truncated *Chlamydia trachomatis mreB* genes were amplified by PCR with Phusion DNA polymerase (NEB, Ipswich, MA) using 100 ng *C. trachomatis* L2 genomic DNA as a template. Some gene segments were directly synthesized as a gBlock fragment (Integrated DNA Technologies, Coralville, IA). The PCR products were purified using a PCR purification kit (Qiagen, Hilden, Germany). The HiFi Assembly reaction master mix (NEB) was used according to the manufacturer's instructions in conjunction with plasmids pASK-GFP-mKate2-L2 (kind gift of Dr. P. Scott Hefty) cut with FastDigest AgeI and EagI (Thermofisher, Waltham, MA), pBOMB4-Tet (kind gift of Dr. Ted Hackstadt) cut with EagI and KpnI, pKT25 or pUT18C cut with BamHI and EcoRI, or pBAD33 cut with XbaI and SalI depending on the construct being prepared. All plasmids were also dephosphorylated with FastAP (ThermoFsher). The products of the HiFi reaction were transformed into either NEB-10beta (for chlamydial transformation plasmids) or DH5alpha (Iq) competent cells (NEB), plated on appropriate antibiotics, and plasmids were subsequently isolated from individual colonies grown overnight in LB broth by using a mini-prep kit (Qiagen). For chlamydial transformation, the constructs were transformed into *dam⁻ dcm⁻* competent cells (NEB) and purified as de-methylated constructs.

Bioinformatics Analysis

Sequences for *Chlamydia trachomatis* serovar L2/434, *Pseudomonas aeruginosa* (PAO1), and *Vibrio cholerae* (O395) were obtained from the NCBI database (https://www.ncbi.nlm.nih.gov/) and for *E.coli* MG1655 from Ecocyc database (https://ecocyc.org/)(32). Protein sequence alignment was performed using Clustal Omega website (https://www.ebi.ac.uk/Tools/msa/clustalo/)(33) and the ESPript3 program (http://espript.ibcp.fr)(34). Helical wheels were made by using Helixator

385   (http://www.tcdb.org/progs/helical_wheel.php)(35). Amphipathic helixes were predicted by

386   using   Amphipaseek   program   (https://npsa-prabi.ibcp.fr/cgi-

387   bin/npsa_automat.pl?page=/NPSA/npsa_amphipaseek.html)(36).

Transformation of *Chlamydia trachomatis*

McCoy cells were plated in a six-well plate the day before beginning the transformation

procedure. *Chlamydia trachomatis* serovar L2 without plasmid (-pL2) (kind gift of Dr. Ian

Clarke) was incubated with 2 µg demethylated plasmid in Tris-CaCl$_2$ buffer (10 mM Tris-Cl,

50 mM CaCl$_2$) for 30 min at room temperature. During this step, the McCoy cells were washed

with 2 mL Hank's Balanced Salt Solution (HBSS) media containing Ca^{2+} and Mg^{2+} (Gibco).

After that, McCoy cells were infected with the transformants in 2 mL HBSS per well. The plate

was centrifuged at 400 *x g* for 15 min at room temperature and incubated at 37°C for 15 min.

The inoculum was aspirated, and DMEM containing 10% FBS and 10 µg/mL gentamicin was

added per well. At 8 h post infection (hpi), the media was changed to media containing 1 µg/mL

cycloheximide and 1 or 2 U/mL penicillin G, and the plate was incubated at 37°C until 48 hpi.

At 48 hpi, the transformants were harvested and used to infect a new McCoy cell monolayer.

These harvest and infection steps were repeated every 48 hpi until mature inclusions were

observed.

Complementation assay

The arabinose-inducible vectors encoding nothing, *E. coli* MreB, chlamydial MreB, or ΔN22

chlamydial MreB were transformed in both wild-type (MG1655) and *mreB* mutant strain

(P2733; kind gift of Dr. Piet de Boer). Chemically competent cells of each strain were prepared

using standard techniques with CaCl$_2$. The strains were cultured overnight, diluted 1/50 in LB

media containing chloramphenicol (34 µg/mL), tetracycline (25 µg/mL) or spectinomycin (50

µg/mL) and cultured for 2 hours when 0.01% arabinose was added as an inducer or cells were

left uninduced as a control. After 2 hours and 6 hours of induction, 4 µL of the cells were

mounted on 1% LB agar pad and covered with a cover slip. The morphologies of the strains

were observed by a Zeiss Imager.Z2 equipped with an Apotome2 using a 100X objective.

The localization of chlamydial MreB fusion proteins in *E. coli*

The *E.coli* MG1655 wild-type and Δ*mreB* mutant (P2733) transformed with the arabinose-

inducible vectors encoding GFP with various MreB N-terminal peptides were cultured at 37℃

with 255 rpm shaking overnight. Overnight cultures were diluted 1:100 (MG1655) or 1:50

(P2733) in LB media containing appropriate antibiotics and cultured at 37℃ with 255 rpm

shaking. After 2 hours of shaking, the cells were induced, or not, with 0.05% (w/v) arabinose

and cultured for more 2 hours. Before mounting the samples, SynaptoRed-C2 (FM4-64;

Cayman Chemical, Ann Arbor, MI) was added in the samples to a final concentration 1.5 µM

to stain the membrane. After 15 min, 4 µL of each culture were placed on 1% LB agar pad and

covered with coverslip. The samples were observed with a Zeiss Imager.Z2 equipped with an

Apotome2 using a 100X objective.

Indirect Immunofluorescence (IFA) Microscopy

HeLa cells were seeded in 24-well plates on coverslips at a density of 10^5 cells per well the day

before infection. Chlamydial strains expressing wild-type or truncated MreBs with a six-

histidine tag at the C-terminus were used to infect HeLa cells in DMEM media containing

penicillin G and cycloheximide. At 6 hpi, 10 nM anhydrotetracycline (aTc) was added. At 10.5

hpi, the coverslips of infected cells were washed with DPBS and fixed with fixing solution

(3.2% formaldehyde and 0.022% glutaraldehyde in DPBS) for 2 min. The samples were then

washed three times with DPBS and permeabilized with ice-cold 90% methanol for 1 min.

Afterwards, the fixed cells were labeled with primary antibodies including goat anti-major outer-membrane protein (MOMP; Meridian, Memphis, TN), rabbit anti-MreB$_{CT}$ antibody (custom anti-peptide antibody directed against the C-terminus of *C. trachomatis* serovar L2 MreB; ThermoFisher), rabbit anti-Hsp60 (kind gift of Dr. Rick Morrison), mouse and rabbit anti-six histidine tag (Genscript, Nanjing, China and Abcam, Cambridge, UK, respectively). Secondary antibodies donkey anti-goat antibody (594), donkey anti-rabbit antibody (647), and donkey anti-mouse (488) were used to visualize the primary antibodies. The secondary antibodies were obtained from Invitrogen. Coverslips were observed by using either a Zeiss LSM 800 confocal microscope or a super-resolution SIM scope (Zeiss ELYRA PS.1).

BACTH Assay

Competent DHT1 *E. coli*, an adenylate cyclase-deficient strain, were co-transformed with pKT25 and pUT18C vectors encoding the genes of interest or empty vectors and spread on M63 minimal media plates containing 50 μg/mL ampicillin, 25 μg/mL kanamycin, 0.5 mM IPTG, 40 μg/mL X-gal, 0.04% casamino acid, and 0.2% maltose. The plates were incubated at 30℃ for 5-7 days. Blue colonies indicate positive interactions whereas no growth or small white colonies indicate no interactions.

Acknowledgments

The authors would like to thank Dr. Lisa Rucks for critical review of the manuscript. The authors would like to thank the following individuals for providing reagents used in this study: Dr. Harlan Caldwell (National Institute of Health), Dr. Ian Clarke (University of Southampton), Dr. Piet de Boer (Case Western University), Dr. Ted Hackstadt (Rocky Mountain Labs/NIH), Dr. P. Scott Hefty (University of Kansas), and Dr. Rick Morrison (University of Arkansas for Medical Sciences). This work was supported by a grant from the National Institute for General

Medical Science (R35GM124798-01) within the National Institutes of Health to SPO. The

University of Nebraska Medical Center Advanced Microscopy Core Facility receives partial

support from the National Institute for General Medical Science INBRE - P20 GM103427 and

COBRE - P30 GM106397 grants, as well as support from the National Cancer Institute (NCI)

for The Fred & Pamela Buffett Cancer Center Support Grant- P30 CA036727 and the Nebraska

Research Initiative.

References

1.   Abdelrahman YM, Belland RJ. 2005. The chlamydial developmental cycle. FEMS
Microbiol Rev 29:949-59.

2.   Moore ER, Ouellette SP. 2014. Reconceptualizing the chlamydial inclusion as a
pathogen-specified parasitic organelle: an expanded role for Inc proteins. Front Cell
Infect Microbiol 4:157.

3.   Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger
L, Tatusov RL, Zhao Q, Koonin EV, Davis RW. 1998. Genome sequence of an obligate
intracellular pathogen of humans: Chlamydia trachomatis. Science 282:754-9.

4.   Bi EF, Lutkenhaus J. 1991. FtsZ ring structure associated with division in Escherichia
coli. Nature 354:161-4.

5.   Liechti GW, Kuru E, Hall E, Kalinda A, Brun YV, VanNieuwenhze M, Maurelli AT.
2014. A new metabolic cell-wall labelling method reveals peptidoglycan in Chlamydia
trachomatis. Nature 506:507-10.

6.   Ouellette SP, Rueden KJ, AbdelRahman YM, Cox JV, Belland RJ. 2015. Identification
and Partial Characterization of Potential FtsL and FtsQ Homologs of Chlamydia. Front
Microbiol 6:1264.

7.   Abdelrahman Y, Ouellette SP, Belland RJ, Cox JV. 2016. Polarized Cell Division of
Chlamydia trachomatis. PLoS Pathog 12:e1005822.

8.   Ouellette SP, Karimova G, Subtil A, Ladant D. 2012. Chlamydia co-opts the rod shape-
determining proteins MreB and Pbp2 for cell division. Mol Microbiol 85:164-78.

9.   Salje J, van den Ent F, de Boer P, Lowe J. 2011. Direct membrane binding by bacterial
actin MreB. Mol Cell 43:478-87.

10.   Srinivasan R, Mishra M, Murata-Hori M, Balasubramanian MK. 2007. Filament

formation of the Escherichia coli actin-related protein, MreB, in fission yeast. Curr Biol

17:266-72.

11.   van Teeffelen S, Wang S, Furchtgott L, Huang KC, Wingreen NS, Shaevitz JW, Gitai

Z. 2011. The bacterial actin MreB rotates, and rotation depends on cell-wall assembly.

Proc Natl Acad Sci U S A 108:15822-7.

12.   Strahl H, Burmann F, Hamoen LW. 2014. The actin homologue MreB organizes the

bacterial cell membrane. Nat Commun 5:3442.

13.   Kemege KE, Hickey JM, Barta ML, Wickstrum J, Balwalli N, Lovell S, Battaile KP,

Hefty PS. 2015. Chlamydia trachomatis protein CT009 is a structural and functional

homolog to the key morphogenesis component RodZ and interacts with division septal

plane localized MreB. Mol Microbiol 95:365-82.

14.   Bendezu FO, Hale CA, Bernhardt TG, de Boer PA. 2009. RodZ (YfgA) is required for

proper assembly of the MreB actin cytoskeleton and cell shape in E. coli. EMBO J

28:193-204.

15.   Swulius MT, Jensen GJ. 2012. The helical MreB cytoskeleton in Escherichia coli

MC1000/pLE7 is an artifact of the N-Terminal yellow fluorescent protein tag. J

Bacteriol 194:6382-6.

16.   Liechti G, Kuru E, Packiam M, Hsu YP, Tekkam S, Hall E, Rittichier JT,

VanNieuwenhze M, Brun YV, Maurelli AT. 2016. Pathogenic Chlamydia Lack a

Classical Sacculus but Synthesize a Narrow, Mid-cell Peptidoglycan Ring, Regulated

by MreB, for Cell Division. PLoS Pathog 12:e1005590.

17.   Wang S, Arellano-Santoyo H, Combs PA, Shaevitz JW. 2010. Actin-like cytoskeleton

filaments contribute to cell mechanics in bacteria. Proc Natl Acad Sci U S A 107:9182-

5.

18.   Bendezu FO, de Boer PA. 2008. Conditional lethality, division defects, membrane

involution, and endocytosis in mre and mrd shape mutants of Escherichia coli. J

Bacteriol 190:1792-811.

19.   Fenton AK, Gerdes K. 2013. Direct interaction of FtsZ and MreB is required for septum

synthesis and cell division in Escherichia coli. EMBO J 32:1953-65.

20.   Kruse T, Bork-Jensen J, Gerdes K. 2005. The morphogenetic MreBCD proteins of

Escherichia coli form an essential membrane-bound complex. Mol Microbiol 55:78-89.

21.   Ouellette SP, Rueden KJ, Gauliard E, Persons L, de Boer PA, Ladant D. 2014. Analysis

of MreB interactors in Chlamydia reveals a RodZ homolog but fails to detect an

517   interaction with MraY. Front Microbiol 5:279.

518 22. van den Ent F, Johnson CM, Persons L, de Boer P, Lowe J. 2010. Bacterial actin MreB
519   assembles in complex with cell shape protein RodZ. EMBO J 29:1081-90.

520 23. Morgenstein RM, Bratton BP, Nguyen JP, Ouzounov N, Shaevitz JW, Gitai Z. 2015.
521   RodZ links MreB to cell wall synthesis to mediate MreB rotation and robust
522   morphogenesis. Proc Natl Acad Sci U S A 112:12510-5.

523 24. Fox A, Rogers JC, Gilbart J, Morgan S, Davis CH, Knight S, Wyrick PB. 1990.
524   Muramic acid is not detectable in Chlamydia psittaci or Chlamydia trachomatis by gas
525   chromatography-mass spectrometry. Infect Immun 58:835-7.

526 25. Hussain S, Wivagg CN, Szwedziak P, Wong F, Schaefer K, Izore T, Renner LD, Holmes
527   MJ, Sun Y, Bisson-Filho AW, Walker S, Amir A, Lowe J, Garner EC. 2018. MreB
528   filaments align along greatest principal membrane curvature to orient cell wall synthesis.
529   Elife 7.

530 26. Hale CA, de Boer PA. 1997. Direct binding of FtsZ to ZipA, an essential component of
531   the septal ring structure that mediates cell division in E. coli. Cell 88:175-85.

532 27. Kawazura T, Matsumoto K, Kojima K, Kato F, Kanai T, Niki H, Shiomi D. 2017.
533   Exclusion of assembled MreB by anionic phospholipids at cell poles confers cell
534   polarity for bidirectional growth. Mol Microbiol 104:472-486.

535 28. van den Ent F, Izore T, Bharat TA, Johnson CM, Lowe J. 2014. Bacterial actin MreB
536   forms antiparallel double filaments. Elife 3:e02634.

537 29. Hesse L, Bostock J, Dementin S, Blanot D, Mengin-Lecreulx D, Chopra I. 2003.
538   Functional and Biochemical Analysis of Chlamydia trachomatis MurC, an Enzyme
539   Displaying UDP-N-Acetylmuramate:Amino Acid Ligase Activity. Journal of
540   Bacteriology 185:6507-6512.

541 30. McCoy AJ, Maurelli AT. 2005. Characterization of Chlamydia MurC-Ddl, a fusion
542   protein exhibiting D-alanyl-D-alanine ligase activity involved in peptidoglycan
543   synthesis and D-cycloserine sensitivity. Mol Microbiol 57:41-52.

544 31. Patin D, Bostock J, Blanot D, Mengin-Lecreulx D, Chopra I. 2009. Functional and
545   biochemical analysis of the Chlamydia trachomatis ligase MurE. J Bacteriol 191:7430-
546   5.

547 32. Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martinez C, Caspi
548   R, Fulcher C, Gama-Castro S, Kothari A, Krummenacker M, Latendresse M, Muniz-
549   Rascado L, Ong Q, Paley S, Peralta-Gil M, Subhraveti P, Velazquez-Ramirez DA,

Weaver D, Collado-Vides J, Paulsen I, Karp PD. 2017. The EcoCyc database: reflecting
new knowledge about Escherichia coli K-12. Nucleic Acids Res 45:D543-D550.

33.   Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,
Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools
APIs in 2019. Nucleic Acids Res doi:10.1093/nar/gkz268.

34.   Robert X, Gouet P. 2014. Deciphering key features in protein structures with the new
ENDscript server. Nucleic Acids Res 42:W320-4.

35.   Saier MH, Jr., Reddy VS, Tsu BV, Ahmed MS, Li C, Moreno-Hagelsieb G. 2016. The
Transporter Classification Database (TCDB): recent advances. Nucleic Acids Res
44:D372-9.

36.   Sapay N, Guermeur Y, Deleage G. 2006. Prediction of amphipathic in-plane membrane
anchors in monotopic proteins using a SVM classifier. BMC Bioinformatics 7:255.

Figure 1. Localization of chlamydial MreB_6xH in *C. trachomatis* using structured illumination microscope (SIM). *C. trachomatis* without plasmid (-pL2) was transformed with anhydrotetracycline (aTc)-inducible vector encoding chlamydial MreB with six histidine tag at the C-terminus. HeLa cells were infected with this strain and chlamydial MreB_6xH expression was induced with 10 nM aTc at 6 hpi. At 10.5 hpi, the infected cells were fixed (3.2% Formaldehyde, 0.022% Glutaraldehyde in PBS) for 2 min and permeabilized with 90% methanol (MeOH) for 1 min. This sample was stained for major outer membrane protein (MOMP; red) and chlamydial MreB (green). Structured illumination microscope (SIM) images were acquired on a Zeiss ELYRA PS.1 super-resolution microscope. The scale bar = 0.5 μm.

(B)

	10	20	30
Amino acid	MSPYRSLYKIKHLSNRLYNKALG**RFDRVF**NFFSGNVGIDL...		
2nd structure	ccccchhhhhhhhhhhhhhhhhhhhhhhhhhhhcccccceec...		
Amphipathic(A) score	2334344343444434344555554544443321122222...		

Extra N-terminal region in *C.trachomatis*
Predicted Amphipathic helix residues

Figure 2. Bioinformatics analysis of chlamydial MreB. Chlamydial MreB has unique

structural features based on the bioinformatics analysis. (A) Protein sequence alignment of

chlamydial and other bacterial MreBs. The blue box represents the extended N-terminus and

predicted amphipathic helix of chlamydial MreB. The alignment was performed with Clustal

Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) and represented with ESPript 3.0

(http://espript.ibcp.fr). (B) The amphipathic score and the predicted secondary structure of the

N-terminus of chlamydial MreB. The prediction of amphipathicity is performed by using

AMPHIPASEEK. The blue and red residues represent the extended N-terminus and predicted

amphipathic helix, respectively. (C) The amphipathicity of the fragments of the N-terminal

region of chlamydial MreB. These amphipathic structures were predicted by online helical

wheel program, named Helixator (http://www.tcdb.org/progs/helical_wheel.php). The circles

represent residues. (yellow circle: non-polar residue; green circle: uncharged polar residue; red

circle: basic polar residue; blue circle: acid polar residue).

Figure 3. Test of complementation of chlamydial MreB in _E.coli_. An _E.coli_ the _mreB_-

deficient mutant (P2733) strain was transformed with an empty arabinose-inducible vector (A)

and the vectors encoding _E.coli_ MreB (B), chlamydial MreB (C) and truncated chlamydial

MreB lacking the extended N-terminal region (D). Stationary phase cultures were diluted to

1:50 in LB media containing 50 μg/mL spectinomycin, 25 μg/mL tetracycline, and 34 μg/mL

chloramphenicol and cultured at 37°C with 225 rpm shaking for 2 h. The cells were then

induced or not with 0.01% (w/v) arabinose. After induction, 4 μL of each culture at 2 h and 6

h were spotted under a 1% LB agar pad and covered with a coverslip. Images were acquired

on a Zeiss Imager.Z2 equipped with an Apotome2 using a 100X objective. The arrows indicate

the cells complemented by the induction of _E.coli_ MreB. Scale bar = 2 μm.

(A)

```
1         10          20        30 32
MSPYRSLYKIKHLSNRLYNKALGRFDRVFNFF    GFP
```

CtrMreB₁₋₃₂ₐₐ-GFP
CtrMreB₁₋₂₈ₐₐ-GFP
CtrMreB₁₋₂₃ₐₐ-GFP
x2 CtrMreB₁₋₂₃ₐₐD-GFP
CtrMreB₂₃₋₃₂ₐₐ-GFP
x2 CtrMreB₂₃₋₃₂ₐₐD-GFP

(C)

(B)

MG1655

| CtrMreB₁₋₃₂ₐₐ-GFP | CtrMreB₁₋₂₈ₐₐ-GFP | CtrMreB₁₋₂₃ₐₐ-GFP | CtrMreB₁₋₂₃ₐₐD-GFP | CtrMreB₂₃₋₃₂ₐₐ-GFP | CtrMreB₂₃₋₃₂ₐₐD-GFP |

GFP / FM 4-64 / Merge

(D)

EcMreB - GFP
ΔN7aa EcMreB - GFP

```
1         10         20      28
MSPYRSLYKIKHLSNRLYNKALGRFDRV    ΔN7aa EcMreB    GFP
```

CtrMreB₁₋₂₈ₐₐ-ΔNEc-GFP
CtrMreB₁₋₂₃ₐₐ-ΔNEc-GFP

(E)

| EcMreB-GFP | ΔN7aa EcMreB(ΔNEc)-GFP | CtrMreB₁₋₂₃ₐₐ-ΔNEc-GFP | CtrMreB₁₋₂₈ₐₐ-ΔNEc-GFP |

GFP / FM 4-64 / Merge

Figure 4. Localization in *E.coli* of various N-terminal regions of chlamydial MreB fused

to GFP. Wild-type MG1655 *E. coli* were transformed with the arabinose-inducible vectors

encoding GFP fusions with various N-terminal regions of chlamydial MreB (A,B) or truncated

E.coli MreB lacking the amphipathic helix (D,E). Stationary phase cultures strains were diluted

1:100 in LB media containing 34 µg/ml chloramphenicol and cultured for 2 h. The cells were

then induced or not with 0.05% (w/v) arabinose and cultured for 2 h more. To stain the

membrane, SynaptoRed-C2 (FM4-64) was added to the samples at a final concentration of 1.5

µM. After 15 min, 4 µL of each culture was spotted under a 1% LB agar pad and covered with

a coverslip. The images were acquired on a Zeiss Imager.Z2 equipped with an Apotome2 using

a 100X objective. (A) The structure of the various N-terminal MreB-GFP fusion peptides. In

the boxed region containing the N-terminus of chlamydial MreB; red residues represent the

predicted amphipathic helix residues. Red constructs show the membrane localization and

black constructs show cytosolic localization. (B) The localization of various N-terminal regions

of chlamydial MreB fused to GFP in MG1655. Note the polar membrane localization of the

MreB$_{1-28aa}$-GFP. Scale bar = 2 µm. (C) A zoomed image of MreB$_{1-28aa}$-GFP localization

represented by the box in (B). The arrowheads represent the polar membrane localization of

the MreB$_{1-28aa}$-GFP. Scale bar = 0.5 µm. (D) The structure of the various N-terminal chlamydial

MreB regions fused with truncated *E.coli* MreB lacking the amphipathic helix. The orange

region represents the amphipathic helix of *E. coli* MreB, and the red residues are predicted

amphipathic helix. (E) The localization of the various N-terminal regions of chlamydial MreB

fused to the truncated *E.coli* MreB in MG1655. Scale bar = 2 µm.

Figure 5. Localization in *C. trachomatis* of various N-terminal regions of chlamydial MreB

fused to GFP. *C. trachomatis* serovar L2 without plasmid (-pL2) was transformed with

anhydrotetracycline (aTc)-inducible vectors encoding GFP (A), chlamydial MreB$_{1-23aa}$-GFP

fusion peptide (B), or chlamydial MreB$_{1-28aa}$-GFP fusion peptide (C). HeLa cells were infected

with the indicated strains, and expression of the GFP fusions was induced at 6 hpi or 16 hpi

with 10 nM aTc. At 10.5 hpi or 20 hpi, the samples were fixed (3.2% Formaldehyde, 0.022%

Glutaraldehyde in 1X PBS) for 2 min and permeabilized with 90% methanol (MeOH) for 1

min. These samples were stained for major outer membrane protein (MOMP; red) with GFP

imaged in green. The arrowheads represent the MreB$_{1-23aa}$-GFP localized at the membrane.

Images were acquired on a Zeiss LSM 800 confocal microscope with 63X objective. Scale bar

= 0.5 μm (10.5 hpi) or 1 μm (20 hpi).

Figure 6. BACTH assay to test protein-protein interactions of full-length and truncated chlamydial MreB. *E.coli* DHT1 (Δ*cya*) was co-transformed with vectors encoding the indicated genes fused to the T25 and T18 catalytic domains of the *B. pertussis* adenylate cyclase. Transformants were plated on M63 minimal medium plates containing 50 μg/mL ampicillin, 25 μg/mL kanamycin, 0.5 mM IPTG, 40 μg/mL X-gal, 0.04% casamino acid, and 0.2% maltose. The plates were incubated at 30℃ for 5-7 days. (A) BACTH assay of the interaction between wild-type and truncated chlamydial MreBs. (B) BACTH assay of the interaction between wild-type or truncated chlamydial MreBs and previously described chlamydial cell division components FtsK, the N-terminal domain of FtsK (FtsKN), and RodZ. A positive (+) control is the interaction between T25-Zip and T18-Zip, the GCN4 leucine zipper motif, and a negative (-) control is the lack of interaction between T25 and the mixture of T18-MreB and T18-ΔN22 MreB. These tests were performed a minimum of two times. Slash box: Not tested; N/D: Non detected.

**Supplementary figure 1. Localization of chlamydial MreB_GFPsw proteins in *C.*
*trachomatis.*** HeLa cells were infected with *C. trachomatis* transformants containing aTc-
inducible vectors encoding MreB_GFPsw proteins. At 12 hpi, expression of the GFP sandwich
fusions was induced with 10 nM aTc, and the samples were fixed (3.2% Formaldehyde, 0.022%
Glutaraldehyde in 1X PBS) at 16 hpi for 2 min and permeabilized with 90% methanol. The
samples were stained for major outer membrane protein (MOMP; red) and GFP (green). Images
were acquired on a Zeiss LSM 800 confocal microscope. Scale bar = 1 μm

660 **Supplementary figure 2. Protein sequence alignment of the N-terminus of MreB from**

661 **diverse *Chlamydia* genera.** The extended N-terminus of chlamydial MreB is conserved across

662 *Chlamydia*. The alignment was performed with Clustal Omega

663 (https://www.ebi.ac.uk/Tools/msa/clustalo/) and represented with ESPript 3.0

664 (http://espript.ibcp.fr).

Supplementary figure 3. Interaction between chlamydial and *E. coli* MreBs and the expression of chlamydial MreB in *E. coli.* (A) BACTH assays were carried out to test interactions between chlamydial MreB and *E. coli* MreB. DHT1 *E. coli* were co-transformed with plasmids encoding the indicated fusion proteins and plated on M63 minimal medium containing 50 μg/mL ampicillin, 25 μg/mL kanamycin, 0.5 mM IPTG, 40 μg/mL X-gal, 0.04% casamino acid, and 0.2% maltose. The plates were incubated at 30°C for 5-7 days. A positive control is the interaction between T25-zip and T18-zip. A negative control is the lack of interaction between T25 and T18-chlamydial MreB and T18-*E. coli* MreB. These tests were performed a minimum of two times. (B) Western blotting was performed to test the expression of chlamydial MreB in strains used in the complementation assay depicted in Figure 3. Whole cell lysates from cultures tested in the complementation assay were separated by SDS-PAGE and transferred to a PVDF membrane. The chlamydial MreB was detected with rabbit anti-MreB primary antibody and IRDye goat anti-rabbit 800CW (LI-COR, Lincoln, NE).

Supplementary figure 4. Localization of N-terminus of chlamydial MreB-GFP fusion proteins in Δ*mreB* mutant (P2733). The Δ*mreB* mutant (P2733) was transformed with the arabinose-inducible vectors encoding GFP fused with diverse N-terminal regions of chlamydial MreB. Samples were prepared as described in the legend to Figure 4 with the membrane labeled with FM4-64. Images were acquired on a Zeiss Imager.Z2 equipped with an Apotome2 using a 100X objective. Scale bar = 2 μm.

Supplementary figure 5. Localization in *C. trachomatis* of various N-terminal regions of chlamydial MreB fused to GFP. *C. trachomatis* without plasmid (-pL2) was transformed with anhydrotetracycline (aTc)-inducible vector encoding GFP (A), chlamydial MreB$_{1-23aa}$-GFP fusion peptide (B) or chlamydial MreB$_{1-28aa}$-GFP fusion peptide (C) as described in the legend to Figure 5. These samples were stained for major outer membrane protein (MOMP; red). Images were acquired on a Zeiss LSM 800 confocal microscope with 63X objective. Scale bar = 0.5 μm (10.5 hpi) or 1 μm (20 hpi).

(A)

```
                     10        20        30
Amino acid   MSPYRSLYKIKHLSNRLYNKALGRFDRVFNFFSGNVGIDL...
2ⁿᵈ structure ccccchhhhhhhhhhhhhhhhhhhhhhhhcccccceec...
Amphipathic(A) score 233443443434344443434455555454444321122222...
```

MreB

ΔN22 MreB

ΔN28 MreB

ΔN32 MreB

Extra N-terminal region in *C.trachomatis*
Predicted Amphipathic helix residues

(B)

MreB-6xH	MOMP	Merge
ΔN22 MreB-6xH	MOMP	Merge
ΔN28 MreB-6xH	MOMP	Merge
ΔN32 MreB-6xH	MOMP	Merge

Supplementary figure 6. The localization of various truncated chlamydial MreBs in *C.*

***trachomatis* L2.** (A) Representation of the various truncated chlamydial MreBs tested. The

blue and red residues represent the extended N-terminus and predicted amphipathic helix,

respectively. (B) *C. trachomatis* serovar L2 transformants containing aTc-inducible vectors

encoding the truncated MreBs were used to infect HeLa cells. At 16 hpi, expression of the

MreB constructs was induced with 10 nM aTc, and these samples were fixed (3.2%

Formaldehyde, 0.022% Glutaraldehyde in 1X DPBS) at 20 hpi for 2 min and permeabilized

with 90% methanol. The samples were stained for major outer membrane protein (MOMP; red)

and six histidine tag (green). Images were acquired on a Zeiss Imager.Z2 equipped with an

Apotome2 using a 100X objective. The white box represents the cells which are zoomed in at

the upper right. Scale bar = 2 μm

 Supplementary table 1. The list of Strains, Plasmids and Primers

Construct Plasmid	Relevant genotype	Ori	Source of Reference
pASK-GFP-mKate-L2 (pTLR2)	*bla* P*tet*	ColE1	(1)
pTLR2-*mreB*_6xH	*bla* P*tet*::*Ctr_mreB*_6xH	ColE1	This study
pBAD33	*cat* P*ara*	p15A	(2)
pBAD33-*Ec_mreB*	*cat* P*ara*::*Ec_mreB*	p15A	This study
pBAD33-*mreB*	*cat* P*ara*::*Ctr_mreB*	p15A	This study
pBAD33-*mreB*_67	*cat* P*ara*::ΔN66nt *Ctr_mreB*	p15A	This study
pBAD33G-*mreB*1-96nt	*cat* P*ara*::*Ctr_mreB*1-96nt-*gfp*	p15A	This study
pBAD33G-*mreB*1-84nt	*cat* P*ara*::*Ctr_mreB*1-84nt-*gfp*	p15A	This study
pBAD33G-*mreB*1-69nt	*cat* P*ara*::*Ctr_mreB*1-69nt-*gfp*	p15A	This study
pBAD33G-*mreB*1-69nt duplicate	*cat* P*ara*::*Ctr_mreB*1-69nt *Ctr_mreB*1-69nt -*gfp*	p15A	This study
pBAD33G-*mreB*67-96nt	*cat* P*ara*::*Ctr_mreB*67-96nt-*gfp*	p15A	This study
pBAD33G-*mreB*67-96nt duplicate	*cat* P*ara*::*Ctr_mreB*67-96nt *Ctr_mreB*67-96nt-*gfp*	p15A	This study
pBAD33G-*Ec_mreB*	*cat* P*ara*::*Ec_mreB-gfp*	p15A	This study
pBAD33G-*Ec_mreB*_22	*cat* P*ara*::ΔN21nt *Ec_mreB-gfp*	p15A	This study
pBAD33G-*mreB*1-84nt-*Ec_mreB*_22	*cat* P*ara*::*Ctr_mreB*1-84nt-ΔN21nt *Ec_mreB-gfp*	p15A	This study
pBAD33G-*mreB*1-69nt-*Ec_mreB*_22	*cat* P*ara*::*Ctr_mreB*1-69nt-ΔN21nt *Ec_mreB-gfp*	p15A	This study
pBOMB4-Tet	*bla* P*tet*	pUC19	(3)
pBOMB4-Tet (no GFP)-*MreB*1-69nt-*gfp*	*bla* P*tet*::*Ctr_mreB*1-69nt-*gfp*	pUC19	This study
pBOMB4-Tet (no GFP)-*MreB*1-84nt-*gfp*	*bla* P*tet*::*Ctr_mreB*1-84nt-*gfp*	pUC19	This study
pKT25	*aph* P*lac*::*t25*	p15A	(4)
pUT18C	*bla*P*lac*::*t18*	ColE1	(4)
pKT25-*mreB*	*aph* P*lac*::*t25-Ctr_mreB*	p15A	This study
pKT25-*mreB*_67	*aph* P*lac*::*t25-*ΔN66nt *Ctr_mreB*	p15A	This study
pKT25-*mreB*_85	*aph* P*lac*::*t25-*ΔN84nt *Ctr_mreB*	p15A	This study
pKT25-*mreB*_97	*aph* P*lac*::*t25-*ΔN96nt *Ctr_mreB*	p15A	This study
pUT18C-*mreB*	*bla*P*lac*::*t18-Ctr_mreB*	ColE1	This study

pUT18C-*mreB_67*	*bla*P*lac*::*t18*-ΔN66nt *Ctr_mreB*	ColE1	This study
pUT18C-*mreB_85*	*bla*P*lac*::*t18*-ΔN84nt *Ctr_mreB*	ColE1	This study
pUT18C-*mreB_97*	*bla*P*lac*::*t18*-ΔN96nt *Ctr_mreB*	ColE1	This study
pUT18C-*rodZ*	*bla*P*lac*::*t18*-Ctr_*rodZ*	ColE1	(5)
pUT18C-*ftsK*	*bla*P*lac*::*t18*-Ctr_*ftsK*	ColE1	(6)
pUT18C-*ftsK*N	*bla*P*lac*::*t18*-N-terminus of *Ctr_ftsK*	ColE1	(6)
pKT25-zip	*aph* P*lac*::*t25-zip*	p15A	(4)
pUT18C-zip	*bla*P*lac*::*t18-zip*	ColE1	(4)
pTLR2-*mreB*-*gfp*_sw	*bla* P*tet*::*Ctr_mreB-gfp_sw*	ColE1	This study
pTLR2-*mreB_67*-*gfp*_sw	*bla* P*tet*::ΔN66nt *Ctr_mreB-gfp_sw*	ColE1	This study
pTLR2-*mreB_85*-*gfp*_sw	*bla* P*tet*::ΔN84nt *Ctr_mreB-gfp_sw*	ColE1	This study
pTLR2-*mreB_97*-*gfp*_sw	*bla* P*tet*::ΔN96nt *Ctr_mreB-gfp_sw*	ColE1	This study
pKT25-*Ec_mreB*	*aph* P*lac*::*t25-Ec_mreB*	p15A	This study
pUT18C-*Ec_mreB*	*bla*P*lac*::*t18-Ec_mreB*	ColE1	This study
pTLR2-*mreB_67*_6xH	*bla* P*tet*::ΔN66nt *Ctr_mreB_6xH*	ColE1	This study
pTLR2-*mreB_85*_6xH	*bla* P*tet*::ΔN84nt *Ctr_mreB_6xH*	ColE1	This study
pTLR2-*mreB_97*_6xH	*bla* P*tet*::ΔN96nt *Ctr_mreB_6xH*	ColE1	This study
pFB112	*tet sdiA*	ColE1	(7)
pFB124	*aadA* cI857(Ts) PλR::*mreC mreD*-LE	pSC101	(7)

Strain	Relevant genotype	Source of Reference
MG1655	F⁻, lambda⁻, *rph-1*	
FB17	*dadR trpE trpA tna mreBCD<> frt*	(7)
DHT1	F⁻ *glnV44* (AS) *recA1 endA1 gyrA96* (Nal^R) *thi-1 hsdR17 spoT1 rfbD1 cya-854 ilv-691 ::Tn10 (TetR)*	(8)
P2733	FB17 pFB112 pFB124	(7)

Primer name	Sequence	Features	Usage
mreB 5'/pTLR2/LIC_F	tttgtttaactttaagaaggagata ATGAGCCCATACCGCAGC	lower case for plasmid overlap construction	for amplification of *mreB*
mreB6XH/pTLR2/R	ccatttttcacttcacaggtcaacc ttaatggtgatggtgatggtg TACTAAACTCTCTTTTCGTTTCTTCAAT TG	lower case for plasmid overlap construction, adds 6xH to *mreB* sequence	for amplification of *mreB*
mreB_1_5'/pBAD 33/F	ttcgagctcggtacccggggatcct tttgtttaactttaagaaggagatataca ATGAGCCCATACCGCAGC	lower case for plasmid overlap construction, adds	for amplification of *mreB*

		Shine-Dalgano (SD) sequence	
mreB_67_5'/pBAD33/F	ttcgagctcggtacccggggatcct tttgtttaactttaagaaggagatatacaatg GGTCGTTTCGATCGTGTATTTAATTTTT TTTC	lower case for plasmid overlap construction, adds Shine-Dalgano (SD) sequence	for amplification of ΔN66nt *mreB*
mreB 3'/pBAD33/LIC_R	ctagaggatccccgggtaccgagct TCATACTAAACTCTCTTTTCGTTTC	lower case for plasmid overlap construction	for amplification of *mreB*s
EcmreB/pBAD33/F	gctcggtacccggggatcct tttgtttaactttaagaaggagatataca ATGTTGAAAAAATTCGTG	lower case for plasmid overlap construction, adds Shine-Dalgano (SD) sequence	for amplification of *Ec_mreB*
EcmreB/pBAD33/R	caagcttgcatgcctgcagg TTACTCTTCGCTGAACAGG	lower case for plasmid overlap construction	for amplification of *Ec_mreB*
mreB/pBAD33G/F	tgggctagcgaattcgagct tttgtttaactttaagaaggagatataca ATGAGCCCATACCGCAGC	lower case for plasmid overlap construction, adds Shine-Dalgano (SD) sequence	for amplification of *mreB_1-96nt*
mreB1-96nt/pBAD33G/R	gtcgactctagaggatcccc c AAAAAAATTAAATACACGATCGAAAC G	lower case for plasmid overlap construction	for amplification of *mreB_1-96nt*
Ctr L2/mreB/pBAD/5'	TATAT<u>GGTACC</u>TGATTAACTTTATAAGG AGGAAAAACAT**A**TGAGCCCATACCGC AGCTTATAT	Underline for KpnI site	for amplification of *mreB*_1-69nt and 1-84nt
Ctr L2/mreB_69/pBAD/3'	CGCCC<u>TCTAGA**A**</u>CCCAACGCCTTGTT ATACAGACGGTTAGA	Underline for XbaI site	for amplification of *mreB*_1-69nt
Ctr L2/mreB_84/pBAD/3'	CGCCC<u>TCTAGA**T**</u>ACACGATCGAAACG ACCCAACGCCTTGTTATAC	Underline for XbaI site	for amplification of *mreB*_1-84nt
mreB_67-84D/pBAD33/F	gctcggtacccggggatcct tttgtttaactttaagaaggagatataca ATGGGTCGTTTCGATCGTG	lower case for plasmid overlap construction, adds Shine-Dalgano (SD) sequence	for amplification of *mreB_67-84nt-gfp* from gBlock
GFP/pBAD33/R	caagcttgcatgcctgcagg TTATTTGTATAGTTCATCCATGCCATG	lower case for plasmid overlap construction	for amplification of *mreB_67-84nt-gfp* from gBlock
Ec mreB/pBAD/5'	CCCCC<u>GAGCTC</u>TGATTAACTTTATAAG GAGG<u>AAAAACAT**A**TGCTGAAAAAATT</u> TCGTGGCATGT	Underline for SacI site	for amplification of Ec_*mreB*

Ec mreB_22/pBAD/ 5'	CCCCCGAGCTCTGATTAACTTTATAAG GAGGAAAAACATATGTTCTCCAATGA CTTGTCCATTGA	Underline for SacI site	for amplification of ΔN21nt Ec_*mreB*
Ec mreB/pBADGFP/ 3'	ATAATGTCGACCTCTTCGCTGAACAG GTCGCCGCCGTGCATGTCGATC	Underline for SalI site	for amplification of Ec_*mreB*
mreB/pBOMB4/F	gatctaaagaggagaaaggatctgc ATGAGCCCATACCGCAG	lower case for plasmid overlap construction	for amplification of *mreB*_1-69nt and 1-84nt fused with *gfp*
GFP/pBOMB4/R	tttgaatggtcgaccggtac TTATTTGTATAGTTCATCCATGCCATG	lower case for plasmid overlap construction	for amplification of *mreB*_1-69nt and 1-84nt fused with *gfp*
mreB_4/T25/F	ctgcagggtcgactctagag AGCCCATACCGCAGCTTATATAAG	lower case for plasmid overlap construction	for amplification of *mreB*
mreB_67/T25/F	ctgcagggtcgactctagag GGTCGTTTCGATCGTGTATTTAATTTTT TTTC	lower case for plasmid overlap construction	for amplification of ΔN66nt *mreB*
mreB_85/T25/F	ctgcagggtcgactctagag TTTAATTTTTTTTCCGGG	lower case for plasmid overlap construction	for amplification of ΔN84nt *mreB*
mreB_97/T25/F	ctgcagggtcgactctagag TCCGGGAATGTTGGTATC	lower case for plasmid overlap construction	for amplification of ΔN96nt *mreB*
mreB_1101/(pKT 25)/R	tcacgacgttgtaaaacgacggccg TCATACTAAACTCTCTTTTCGTTTC	lower case for plasmid overlap construction	for amplification of *mreBs*
mreB_4/T18/F	actgcaggtcgactctagag AGCCCATACCGCAGCTTATATAAG	lower case for plasmid overlap construction	for amplification of *mreB*
mreB_67/T18/F	actgcaggtcgactctagag GGTCGTTTCGATCGTGTATTTAATTTTT TTTC	lower case for plasmid overlap construction	for amplification of ΔN66nt *mreB*
mreB_85/T18/F	actgcaggtcgactctagag TTTAATTTTTTTTCCGGG	lower case for plasmid overlap construction	for amplification of ΔN84nt *mreB*
mreB_97/T18/F	actgcaggtcgactctagag TCCGGGAATGTTGGTATC	lower case for plasmid overlap construction	for amplification of ΔN96nt *mreB*
mreB_1101/(pUT	accatattacttagttatatcgatg	lower case for	for

18C)/R	TCATACTAAACTCTCTTTTCGTTTC	plasmid overlap construction	amplification of *mreBs*
Ct009/rodZ/BACTH/5'BamHI	ATATA<u>GGATCC</u>A**GG**TAGAGCAAACCAGGGGAATTAC	Underline for BamHI site	for amplification of *rodZ*
Ct009/rodZ/BACTH/3'KpnI	AGGGT<u>GGTACC</u>**T**TA**G**AAAAGGTTGAATAGATTCCCTAG	Underline for KpnI site	for amplification of *rodZ*
Ct739/ftsK/BACTH/5'XbaI	ACTAGCTGGT<u>TCTAGA</u>T**G**GAAAAGAACGGAAGAAA	Underline for XbaI site	for amplification of *ct739(Ctr_ftsK)*
Ct739/ftsK/BACTH/3'KpnI	CACAGCTAGA<u>GGTACC</u>AT**T**AATCGTCCTGATTTGA	Underline for KpnI site	for amplification of *ct739(Ctr_ftsK)*
Ct739/ftsKN/BACTH/5'XbaI	ACTAG<u>TCTAGA</u>T**G**GAAAAGAACGGAAGAAA	Underline for XbaI site	for amplification of N-terminus of *ct739(Ctr_ftsK)*
Ct739/ftsKN/BACTH/3'KpnI	CTAGA<u>GGTACC</u>AT**T**TGAGAAATTTTAGGAGA	Underline for KpnI site	for amplification of N-terminus of *ct739(Ctr_ftsK)*
mreB67 5'/pTLR2/LIC_F	tttgtttaactttaagaaggagata ATGGGTCGTTTCGATCGTG	lower case for plasmid overlap construction	for amplification of ΔN66nt *mreB*
mreB85 5'/pTLR2/LIC_F	tttgtttaactttaagaaggagata ATGTTTAATTTTTTTTCCGGGAATG	lower case for plasmid overlap construction	for amplification of ΔN84nt *mreB*
mreB_97_5'/pTLR2/F	tttgtttaactttaagaaggagata atgTCCGGGAATGTTGGTATC	lower case for plasmid overlap construction	for amplification of ΔN96nt *mreB*
gfp/pTLR2-mreB/LIC F	ttatccgttaggtggaggtagt AGTAAAGGAGAAGCACTTTC	lower case for plasmid overlap construction	for amplification of *gfp*
gfp/pTLR2-mreB/LIC R	cctgatcactacctcc GAGTCCGGACTTGTATAGTTC	lower case for *mreB* 3' insert DNA overlap construction	for amplification of *gfp*
mreB 5'/pTLR2/LIC_R 2	cctttactactacctcc ACCTAACGGATAAGCAGAACCTATAG	lower case for *gfp* insert DNA overlap construction	for amplification of N-terminus of *mreB*

		lower case for *gfp* insert DNA overlap construction, adds link residues, GGS	for amplification of C-terminus of *mreB*
mreB 3'/pTLR2/LIC_F2	caagtccggactc *ggaggtagt* GATCAGGAATTGGAGATG		
Ec mreB/BACTH/5' Bam	CCTTG<u>GGATCC</u>**T**CTGAAAAAATTTCGT GGCATGT	Underline for BamHI site	for amplification of Ec_*mreB*
Ec mreB/BACTH/3' Sacstp	ATATT<u>GAGCTC</u>**T**TA**C**TCTTCGCTGAAC AGGTCGCC	Underline for SacI site	for amplification of Ec_*mreB*
Ec mreB/BACTH/3' Sac	TATAT<u>GAGCTC</u>CC**C**TCTTCGCTGAACA GGTCGCC	Underline for SacI site	for amplification of Ec_*mreB*

1. Wickstrum J, Sammons LR, Restivo KN, Hefty PS. 2013. Conditional gene expression in Chlamydia trachomatis using the tet system. PLoS One 8:e76743.

2. Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. J Bacteriol 177:4121-30.

3. Bauler LD, Hackstadt T. 2014. Expression and targeting of secreted proteins from Chlamydia trachomatis. J Bacteriol 196:1325-34.

4. Karimova G, Ullmann A, Ladant D. 2001. Protein-protein interaction between Bacillus stearothermophilus tyrosyl-tRNA synthetase subdomains revealed by a bacterial two-hybrid system. J Mol Microbiol Biotechnol 3:73-82.

5. Ouellette SP, Rueden KJ, Gauliard E, Persons L, de Boer PA, Ladant D. 2014. Analysis of MreB interactors in Chlamydia reveals a RodZ homolog but fails to detect an interaction with MraY. Front Microbiol 5:279.

6. Ouellette SP, Karimova G, Subtil A, Ladant D. 2012. Chlamydia co-opts the rod shape-determining proteins MreB and Pbp2 for cell division. Mol Microbiol 85:164-78.

7. Bendezu FO, de Boer PA. 2008. Conditional lethality, division defects, membrane involution, and endocytosis in mre and mrd shape mutants of Escherichia coli. J Bacteriol 190:1792-811.

8. Dautin N, Karimova G, Ullmann A, Ladant D. 2000. Sensitive genetic screen for protease activity based on a cyclic AMP signaling cascade in Escherichia coli. J Bacteriol 182:7060-6.