Mast seeding promotes evolution of scatter-hoarding

Running head: Masting and the evolution of seed caching

Rafał Zwolak[1], Dale Clements[2], Andrew Sih[2], and Sebastian J. Schreiber[2]

[1]Department of Systematic Zoology
Institute of Environmental Biology
Adam Mickiewicz University
Umultowska 89
61-614 Poznań, Poland
10
11   [2]Center of Population Biology
12   University of California
13   One Shields Avenue
14   Davis, California, 95616 USA
15

KEYWORDS (4-6): Evolutionary stable strategies, masting, scatterhoarding, pilferage, seed caching,

synzoochory.

Number of words: 7127

ABSTRACT

Many plant species worldwide are dispersed by scatterhoarding granivores: animals that hide seeds in

numerous, small caches for future consumption. Yet, the evolution of scatterhoarding is difficult to

explain because undefended caches are at high risk of pilferage. Previous models have attempted to

solve this problem by giving cache owners unrealistically large advantages in cache recovery, by kin

selection (but individuals that cache and those that pilfer are usually unrelated), or by introducing

reciprocal pilferage of "shared" seed resources. However, the role of environmental variability has

been so far overlooked in this context. One important form of such variability is masting, which is

displayed by many plant species dispersed by scatterhoarders. We use a mathematical model to

investigate the influence of masting on the evolution of scatter-hoarding. The model accounts for

periodically varying annual seed fall, caching and pilfering behavior, and the demography of

scatterhoarders. Masting, through its effects on population density, reduces cache pilferage and

lowers the reproductive cost of caching (i.e. the cost of caching for the future rather than using seeds

for current reproduction). These reductions promote the evolution of scatter-hoarding behavior

especially when interannual variation in seed fall and the period between masting events are high.

                                                            Introduction

Many species of woody plants, particularly those that produce nuts and nut-like dry fruits, rely for

seed dispersal on scatterhoarders: animals such as corvids, rodents, and certain marsupials that hide

seeds for future consumption in numerous, small, widely-dispersed caches (Forget and Vander Wall

2001; Peserdorfer et al. 2016; Vander Wall et al. 2017). However, explaining the evolutionary

advantage of this behavior has been difficult. The caches are undefended and often suffer very high

rates of pilferage by con- and heterospecifics (Schmidt and Ostfeld 2008; Jansen et al. 2012; Zwolak

et al. 2016; Dittel et al. 2017), thus scatter-hoarding appears vulnerable to cheating by non-caching

pilferers (Anderson and Krebs 1978; Smulders 1998; Vander Wall and Jenkins 2003). How can scatter-

hoarding be advantageous under such circumstances?

This problem can be solved by invoking kin selection (but cache owners and pilferers are

usually unrelated) or by giving cache owners a high advantage in cache recovery over naïve

individuals (Anderson and Krebs 1978). Empirical estimates of the owner advantage are widely

varied, but in most studies cache owners are 2-4 times more likely to recover their caches when

compared with naïve individuals (Jacobs and Liman 1991; Jacobs 1992; Briggs and Vander Wall 2003;

Thayer and Vander Wall 2005; Vander Wall et al. 2006; 2008; Hirsch et al. 2013; Gu et al. 2017; see

also Steele et al. 2011). Yet, some of these estimates may be too high. First, many studies estimated

the *per capita* advantage of owner versus pilferer, but under natural conditions there typically are

many potential pilferers per each cache owner. Thus, caches are more likely to be emptied by a

pilferer than by the owner even when *per capita* advantage of the latter is substantial. Second, most

empirical studies have been short-term, with cache recovery examined over several days at most.

Long-term (e.g. overwinter) removal rates might be very similar for cache owners and pilferers

because cache locations are forgotten over time (Balda and Kamil 1992; Bednekoff et al. 1997;

Barkley and Jacobs 1998). Over the long-term, cache owners will increasingly rely on olfaction or

random search, just like naïve individuals do. As a consequence, the owners advantage appears to be

insufficient to prevent cache loss to pilferage (2-30% lost per day, according to a review by Vander

Wall and Jenkins 2003).

Another putative solution to this puzzle is the reciprocal pilferage hypothesis (Vander Wall

and Jenkins 2003; see also Smulders 1998). In principle, caching can represent an adaptive, stable

strategy when all caches are reciprocally pilfered by scatter-hoarding animals with overlapping home

ranges. Under this scenario, caches represent a collective resource used by selfish individuals

(Vander Wall and Jenkins 2003). The reciprocal pilferage hypothesis predicts that animals are unlikely

to avoid pilferage, but can compensate for it by pilfering caches of other individuals. As a corollary,

individuals should invest in their pilfering tactics rather than in theft-reducing strategies (Vander Wall

and Jenkins 2003).

Yet, many species of scatter-hoarding animals put considerable effort in behaviors apparently

aimed to reduce pilferage (Dally et al. 2006a). This phenomenon has been particularly well-studied in

corvids (e.g. Dally et al. 2006b; Shaw and Clayton 2013), but scatter-hoarding rodents also use a

diverse array of theft-reduction strategies, including some that are costly in time and energy (e.g.

transporting seeds further to reduce the density of caches: Galvez et al. 2009; Hirsch et al. 2012),

some that are potentially dangerous (e.g. caching in open places where predation risk is higher, but

the risk of pilferage is lower: Muñoz and Bonal 2011; Steele et al. 2013, 2015) and some that appear

quite sophisticated (e.g. deceiving observers by making "fake" caches: Steele et al. 2008). This level

of investment in pilferage prevention rather than pilferage tolerance is unexpected under the

reciprocal pilferage hypothesis, and indeed has been interpreted as evidence against it (Leaver et al.

2007; Pravosudov et al. 2010).

However, existing models of scatter-hoarding have largely overlooked the role of

environmental variability in the evolution of this behavior. Such variability is pervasive in ecosystems

dominated by plants that produce dry, scatterhoarder-dispersed fruits because such plants usually

show pronounced masting (Herrera et al. 1998; Vander Wall 2001). Examples of scatterhoarder-

dispersed masting plants can be found in the tropics (Norden et al. 2007; Mendoza et al. 2018),

deserts (Meyer and Pendleton 2015; Auger et al. 2016), and in temperate zones (Koenig and Knops

2000; Schauber et al. 2002; Shibata et al. 2002). While studies of masting have often emphasized the

benefit of masting to plants in terms of reduced *per capita* seed predation, masting also has

important effects on consumer population dynamics that can feedback to affect the evolution of

both masting and caching. In particular, masting has long been known to cause cycles of feast and

famine that induce striking fluctuations in consumer population size (Ostfeld and Keesing 2000; Yang

et al. 2010; Bogdziewicz et al. 2016). Typically, masting triggers a temporary increase in consumer

population size followed by a pronounced crash. Thus, when the next mast year comes, seed to

consumer ratios are particularly high (Kelly 1994; Ostfeld and Keesing 2000).

We use a mathematical model to investigate the influence of mast-related fluctuations in

scatterhoarder population size on the evolution of scatter-hoarding. The model accounts for periodic

seed fall and demography of the scatter-hoarding species. The scatterhoarders consume or cache

harvested seeds and pilfer or recover their own caches over years that differ in the magnitude of

seed fall. Previous models demonstrated that caching is influenced by the owner's advantage in

cache recovery and the probability that scatterhoarders survive long enough to use the caches

(Anderson and Krebs 1978; Smulders 1998; Vander Wall and Jenkins 2003). Thus, we include these

factors in our investigation, but expand on previous models by examining the effects of

environmental variability resulting from mast seeding. We examine how the proportion of seeds

scatter-hoarded versus immediately consumed varies with (1) masting intensity, (2) the frequency of

mast years, (3) the owner's advantage in cache recovery, and (4) the survival of scatterhoarders. Our

results demonstrate that mast-related fluctuations in scatterhoarder population size reduce the risk

of cache loss to pilferers and the reproductive cost of caching (i.e. the cost of caching seeds for future

use rather than using seeds for current reproduction), thus promoting the evolution of scatter-

hoarding.

                                                Methods

                                         Model Description

   We consider a population of scatterhoarders that experience three distinctive periods of seed

   availability in each year. During the first period (fall), there is seed fall of $S(t)$ from the primary seed

   source. Seeds are gathered at a rate proportional to the density $n_1(t)$ of scatterhoarders during this

   period. The proportionality constant a_1 corresponds to the per-capita seed harvest rate. Seeds are

   also lost to other sources (e.g. competitors, germination, decay, etc.) at a per-capita rate of L_1. If all

   seeds are gathered or lost to other sources by the end of the first period, then the amount of seed

   gathered per individual equals:

$$G(t) = \frac{a_1 S(t)}{L_1 + a_1 n_1(t)}$$

   All seeds above a threshold, T, are cached by individuals for later in the year. The seeds which are not

   cached, min{G(t),T}, are used for survival and reproduction. The number of offspring produced by

   individual, $R_1(t)$, at the end of this first period is a decelerating function with a maximal number of

   offspring *b* and a half saturation constant *h* (i.e. *h* is the amount of resources required to produce *b*/2

   offspring). The fraction of adults surviving from the first period (fall) to the second period of the year

   (winter/spring) equals s_1. Thus, the total density $n_2(t)$ of individuals entering the second period

   equals

$$n_2(t) = [R_1(t) + s_1]n_1(t) \text{ where } R_1(t) = \frac{b \, min\{G(t), T\}}{h + min\{G(t), T\}}$$

   The main resource available to individuals during this second period is the total cache of seeds

   max{G(t)-T,0}$n_1(t)$. Owners of the cached seed who survived gather their cached seed at a rate

   proportional to the size of their cache. This proportionality constant a_2 corresponds to the per-capita

   rediscovery and use rate of their cache. All other individuals are assumed to pilfer seed from others'

143   caches at a per-capita rate a_{pil}. Seeds are lost to other sources at a per-capita rate of L_2. If all cached

144   seeds are gathered or lost by the end of the second period, then the fraction of cache recovered by

145   an individual that survived from period 1 to period 2 is

146
$$M(t) = \frac{a_2}{L_2 + a_2 + a_{pil}(n_2(t) - 1)}$$

147   while the fraction of this individual's cache that was pilfered by another individual is

148
$$O(t) = \frac{a_{pil}}{L_2 + a_2 + a_{pil}(n_2(t) - 1)}$$

149   For seed cache whose owner died, the fraction that was recovered by a living non-owner is

150
$$D(t) = \frac{a_{pil}}{L_2 + a_{pil}n_2(t)}$$

151   It follows that the amount of cached seed gathered by a surviving individual from period 1 is

152
$$C_{survivor}(t) = \max\{G(t) - T, 0\}[M(t) + O(t)(s_1 n_1(t) - 1) + D(t)(1 - s_1)n_1(t)]$$

153   and the amount of cached seed gathered by a new individual from period 2 is

154
$$C_{new}(t) = \max\{G(t) - T, 0\}[O(t)s_1 n_1(t) + D(t)(1 - s_1)n_1(t)]$$

155   If s_2 is the fraction of individuals surviving to period 3 (summer), then the density of individuals

156   entering period 3 equals

157
$$n_3(t) = [R_2(t) + s_2]n_2(t)$$

158   where $R_2(t)$ is the population-level per-capita fecundity corresponding to the weighted combination

159   of reproductive contributions of individuals surviving from period 1 and new born individuals from

160   period 1 (for simplicity, we assume the same value of h for surviving and new born individuals):

161
$$R_2(t) = \frac{b\,C_{survivor}(t)}{h + C_{survivor}(t)}\frac{s_1}{R_1(t) + s_1} + \frac{b\,C_{new}(t)}{h + C_{new}(t)}\frac{R_1(t)}{R_1(t) + s_1}$$

162   During this final period of the year (summer), individuals rely on other resources with availability A to

163   reproduce and survive with probability s_3. Thus, the density of individuals entering the Fall of the

164   next year equals

165
$$n_1(t + 1) = [R_3(t) + s_3]n_3(t) \text{ where } R_3(t) = \frac{b\,A}{h + n_3(t)}$$

By composing the equations across the three periods of the year, the yearly update rule for

population densities at the beginning of Fall is

$$n_1(t+1) = [R_3(t) + s_3][R_2(t) + s_2][R_1(t) + s_1]n_1(t)$$

We modeled seedfall S(t) in the fall as a periodic function of time where the period P corresponds to

the time between masting years. In the masting years, S(t)=S_{high} and otherwise S_{low}. Our analysis

assumes that the total seed output, S(1)+S(2)+…+S(P), is fixed and varies the proportion of total seed

output in the masting year. Higher intensity of masting means more seeds during the masting year,

but concomitantly fewer seeds in other years (as opposed to just increasing seed output in masting

years with no effect on seed production in other years). Similarly, when we vary the number of years

between masting events, the total seed output remains the same (i.e., longer intermast interval

corresponds with higher seed production in mast years).

Figure 1 illustrates typical dynamics of the model for the baseline parameters described

below. For all our simulations, the dynamics asymptotically exhibit stable cyclic behavior with a

period corresponding to the period of the masting events. For low caching thresholds (dotted line in

Fig. 1b), individuals cache most of their seeds every fall. For high caching thresholds (dashed line in

Fig. 1b), individuals either only cache on masting years as shown or not at all (not shown). Population

densities (Fig. 1c) exhibit seasonal as well as yearly variation. Moreover, for any given season,

population densities are maximal following a masting event and decrease year to year until the next

masting event.

Fig. 1. *The dynamics of fall seed masting (a), per-cent seed cached by an individual for different caching thresholds T (b), and population densities (c) for the different caching thresholds T (see model description). In (a), there is a 4 year interval between fall masting events. In (b), the percent seeds cached by individuals for populations with a low (dotted), intermediate (solid), and high (dashed) caching threshold. In (c), the intra- and interannual variation in population densities for populations with a low (dotted), intermediate (solid), and high (dashed) caching threshold. The dynamics in (b) and (c) correspond to a stable 4-year periodic orbit of the model.*

Model Parameters

The parameter values mimic an interaction between a masting tree and a scatterhoarding rodent and are based mostly on research on European beech (*Fagus sylvatica*) and yellow-necked mice (*Apodemus flavicollis*). Empirical results were used to generate order of magnitude parameter estimates, which were then adjusted, by no more than a factor of 10, so that the equilibrium population sizes were roughly in line with those given in the literature. While we found it useful to base our parameter estimates on a specific, reasonably well-studied system, moderate variation in these parameters does not change the qualitative results.

The parameters a_1 and L_1 may be reduced to the single parameter L_1/a_1. Rearranging the equation for G yields $L_1/a_1 = \frac{(1-G/S_0)n_1}{G/S_0}$. The parameters n_1 and n_2 were taken to be the average

species-wide density for A*podemus flavicollis*: 17.7 individuals/ha (Jones et al. 2009 [PanTHERIA]).

Estimates of the proportion of seeds removed from the forest floor (G/S_0) tend to be variable with Zwolak et al. (2016) reporting 78% seed removal during mast years and 91% seed removal during non-mast years for *A. flavicollis*, and Lelouarn and Schmitt (1972) reporting 41% and 74% seed removal by *Apodemus sylvaticus* during two different years. We selected the intermediate value of 74% as tentative estimate of seed removal. Thus, $L_1/a_1 = 6.22$. In all cases where L_1 and a_1 were treated as separate parameters, we used $a_1 = 1$ and set L_1 equal to our choice for L_1/a_1.

The parameters a_{pil} and L_2 may similarly be reduced to L_2/a_{pil}. Zwolak et al. (2016) estimated the recovery of seeds from artificial caches to be 54% during nonmast years and 5% during mast years, which we assumed to be roughly equivalent to the proportion of seeds recovered from an abandoned cache ($D/\max\{G(t) - T, 0\}$). We then estimated L_2/a_{pil} to be between 15.1 and 336.3. In our analysis, we set this value to the upper end of this range (300) as this leads to more conservative estimates of when caching evolves. In all cases where L_2 and a_{pil} were treated as separate parameters, we used $a_{pil} = 1$ and set L_2 equal to our choice for L_2/a_{pil}.

We let $a_2 = 3$ (when $a_{pil} = 1$). This value approximates the results of several studies on scatterhoarding rodents (Vander Wall et al. 2006, 2008; Thayer and Vander Wall 2005; Hirsch et al. 2013) that documented seed removal rates by cache owners and naïve foragers. Although some research (Gu et al. 2017) suggests that the owner advantage might be even higher, it also might be lower, particularly later in the season, as we explained in the Introduction.

We assumed a maximum litter size of 11 individuals (Macdonald and Tattersall 2001), with one breeding event per period (2-3 litters per year: Pucek 1984). Assuming that half the population are female and half of the individuals born are female, this yields $b = 5.5$.

We used Jensen's (1982) monthly winter survival estimate of 77.5% as the yearlong monthly survival rate (see also Pucek 1992 for similar values). We assumed that each period lasts four months, yielding 36.1% as the survival rate for each period ($s_1, s_2,$ and s_3).

Half-saturation constant for mid-year reproduction (*h*) was set as 124 seeds/offspring

multiplied by half the maximum number of offspring (*b/2*). This value was calculated on the basis of

energy contents of beech seeds (Grodziński and Sawicka-Kapusta 1970), energy requirements of

yellow-necked mice (0.60 kcal/g/day: Jensen 1982; average body mass of yellow-necked mice is 28.3

g: AnAge), and typical costs of reproduction-related energy expenditure in small mammals (25%

increase in energy expenditure during gestation and 200% increase during lactation: Millar 1978;

1979; Gittleman and Thompson 1988; Sikes 1995; Zhu et al. 2015), given the length of gestation and

lactation in yellow-necked mice (26 and 22 days, respectively: AnAge).

239   *Numerical Methods*

To identify the evolutionary stable caching strategies, we examined whether a small mutant

subpopulation using the germination threshold T_m can invade a resident population using the

germination threshold *T*. When the mutant subpopulation densities $m_i(t)$ in each of the periods

i=1,2,3 are sufficiently small, the density feedback of the mutant population on the resident

population and itself is negligible. Hence, the dynamics of the mutant in the initial phase of invasion

can be approximated by a linearization of its dynamics. We now describe these linearized dynamics.

As the mutant and resident individuals only differ in their caching strategy, the amount of seeds

gathered in year *t* by a mutant individual equals the amount of seeds gathered G(t) by a resident

individual. As for the resident dynamics, yearly update of the mutant's fall density is of the form

250                $$m_1(t+1) = [Q_3(t) + s_3][Q_2(t) + s_2][Q_1(t) + s_1]m_1(t)$$

251    $Q_1(t)$ corresponds to the number of offspring produced by a mutant individual during the fall and

252   only differs from the resident in its threshold T_m

253                        $$Q_1(t) = \frac{b \, min\{G(t), T_m\}}{h + min\{G(t), T_m\}}$$

$Q_2(t)$ corresponds to the number of offspring produced by a mutant individual during the

winter/spring given by a weighted combination due to the fraction of individuals that survived from

the fall and individuals born in the fall:

$$Q_2(t) = \frac{b \, C_{m,survivor}(t)}{h + C_{m,survivor}(t)} \frac{s_1}{Q_1(t) + s_1} + \frac{b \, C_{new}(t)}{h + C_{new}(t)} \frac{Q_1(t)}{Q_1(t) + s_1}$$

where $Q_2(t)$ differs from $R_2(t)$ only in its first term due to surviving individuals with the mutant

caching strategy: $C_{m,survivor}(t) = \max\{G(t) - T_m, 0\}M(t) + \max\{G(t) - T, 0\}[O(t)(s_1 n_1(t) -$

$1) + D(t)(1 - s_1)n_1(t)]$. Finally, the number of offspring produced by a mutant over the summer is

the same as the resident i.e. $Q_3(t) = R_3(t)$.

Whether the mutants playing strategy T_m are able to invade the residents playing the strategy T or

not depends on their long-term per-capita growth rate

$$s(T, T_m) = \lim_{t \to \infty} \tfrac{1}{t} \log[Q_3(t) + s_3][Q_2(t) + s_2][Q_1(t) + s_1]$$

provided the limit exists. Over the parameter space (see previous section) that we simulated, the

population dynamics always converged to a periodic solution whose period kP is a multiple k of the

seed masting period P. Typically, this multiple was 1 or 2 or 4, the latter two corresponding to period-

doubling bifurcations. We developed R code to efficiently approximate these periodic solutions. For

these periodic solutions of the resident dynamics, the long-term per-capita growth rate of mutant

strategy T_m against resident strategy T equals

$$s(T, T_m) = \frac{1}{kP} \sum_{t=1}^{kP} \log[Q_3(t) + s_3][Q_2(t) + s_2][Q_1(t) + s_1]$$

A strategy T is an ESS for caching if $s(T, T_m) < 0$ for all strategies $T_m \neq T$. Using the method of adaptive

dynamics (Geritz et al. 1997), we identified potential ESSs by solving for roots of the fitness gradient

$\frac{\partial s}{\partial T_m}(T, T)$. In the Appendix, we derive an explicit expression of this fitness gradient

Fig. 2. *Period 1 (fall) population density in individuals/ha (a), pilferage risk, defined as the probability that cached seed would be pilfered (b), marginal reproductive costs of caching (c), and proportion of seeds cached rather than eaten (d) as a function of masting intensity (expressed as the percentage of total seed production that occurs during mast years), with mast occurring every 4th year. Dashed, solid, and dotted lines represent the magnitude of owner's advantage in cache recovery (owners 1.5, 3, and 6 times more likely to discover their own caches relatively to naïve foragers). Plotted variables represent values in the mast year. Vertical gray lines correspond to the minimal masting intensity at which caching is selected for.*

Results

Increasing intensity of masting generally results in decreased period 1 (fall) scatterhoarder population density (i.e., the density that enters masting events; Fig. 2a). Intuitively, because reproduction is a saturating function of seeds gathered, the reproductive gains of higher seed availability during masting years were outweighed by the reproductive losses due to lower seed

availability during non-mast years. As seed caching generally increases fall population densities by

preventing population crashes in winter/spring, this indirect positive effect of masting intensity

appears when masting intensities is just sufficient to select for caching and results in intermediate

positive trends in fall density (Fig. 2a).

Increasing masting intensity also reduces the risk that a cached seed would be pilfered (Fig.

2b), particularly when 80% or more seeds are produced during mast years. Notice that the responses

of pilferage risk and fall density are tightly correlated (see, also, Figs. 3 and 4). Furthermore,

increasing masting intensity is associated with a decline in marginal reproductive costs of caching

(Fig. 2c) and with an accelerating increase in the ESS proportion of seeds cached rather than eaten

(Fig. 2d). In the baseline scenario, scatterhoarders start caching when about 50% of seeds are

produced during mast years.

Higher recovery advantage tends to increase scatterhoarder densities (Fig. 2, dotted vs. solid

vs. dashed line), probably because (i) on average individuals are recovering more of the cached

seeds, and (ii) there is selection for increased seed caching (Fig 2d), which tends to be ecologically

advantageous. Increasing the owner's advantage in cache recovery (Fig. 2, dotted line) selects for

more caching despite increases in population densities and cache pilfering; decreasing the owner's

advantage (Fig. 2, dashed line) decreases the proportion of seeds cached even though population

densities and cache pilferage also drop. However, the effect of the owner's advantage in cache

recovery is relatively weak compared to the effect of masting intensity.

Fig. 3. *Period 1 (fall) population density (a), pilferage risk (b), marginal reproductive costs of caching*

(c), and proportion of seeds cached rather than eaten (d) as a function of masting interval. Dashed,

solid, and dotted lines represent masting intensity (50, 70 or 90% of total seed production occurring

during mast years). Plotted variables represent values in the mast year.

Increasing the inter-mast period leads to declines in fall density (Fig. 3a), pilferage risk (Fig. 3b), and

marginal reproductive costs (Fig. 3c), and an increase in the proportion of seeds cached (Fig. 3d).

Intuitively, more years with poor seed crops between masting events leads to lower densities of

individuals entering the fall of a masting year. This reduction in density leads to fewer individuals

entering winter and, thereby, lower risk of seed pilferage. In addition, the low densities of individuals

entering the masting years and the greater amount of seeds available during the masting year

increase the amount of seeds harvested per individual and, due to the effects of diminishing returns,

lower marginal reproductive costs of caching. Collectively, lower reproductive costs and lower

pilferage risk select for more caching. Varying masting intensity (50, 70 or 90% of seeds produced

during mast years: dotted, solid or dashed line on Fig. 3) affects the magnitude of these changes,

with only minor effects on the shape of responses to the masting interval. Scatterhoarders go extinct

when inter-mast periods are long and a high proportion of seeds is produced in mast years.

Fig. 4. *Period 1 (fall) population density (a), pilferage risk (b), marginal reproductive costs of caching*

(c), and proportion of seeds cached rather than eaten (d) as a function of scatterhoarder survival.

Dashed, solid, and dotted lines denote responses to changes in fall, winter, and summer survival,

respectively. Plotted variables represent values in the mast year. Vertical gray lines correspond to the

maximal survivorships at which caching is selected for.

Increasing the survival of scatterhoarders leads to nearly identical increases in period 1 population

density, pilferage risk, and marginal reproductive costs of caching (Fig. 4). These effects are

considerably stronger for changes in winter and summer survival rather than in fall survival.

Intuitively, this is because the greatest concentration of births occurs between the fall and winter,

resulting in the winter population having a higher percentage of new individuals (who are not subject

to mortality during the previous period) than the summer and fall populations. Thus, an increase in

mortality in the fall affects a smaller proportion of the population than an increase in mortality in the

winter or summer. Interestingly, varying summer and winter/spring versus fall survival has

contrasting effects on the percentage of seeds cached. Increasing summer and winter/spring survival

leads to a sharp decline in the predicted proportion of seeds cached most likely due to the large

increase in fall density, and the drastically increased costs associated with increased pilferage risk

and lost reproductive opportunities. When monthly winter/spring and summer survival exceeded

50%, no seeds should be cached. In contrast, increasing fall survival selects for an increase in the

percentage of seeds cached rather than eaten, most likely because the advantage of surviving to

make use of one's cache outweighs the weak increases in costs.

Fig. 5. *Relationships between input parameters (masting interval and intensity, expressed as the proportion of seeds produced during mast years, scatterhoarder survival, and cacher's advantage in cache recovery) and emergent properties of the model (scatterhoarder population density, the proportion of cached seeds that are pilfered, reproductive costs of caching, and the quantity of interest: the proportion of seeds cached by scatterhoarders).*

Discussion

The fact that masting causes strong fluctuations in populations of seed-eating animals has been well-known for a long time (Curran and Leighton 2000; Ostfeld and Keesing 2000; Bogdziewicz et al. 2016), yet the traditional research focus has been on how the satiation-starvation cycle reduces seed losses to pre- and post-dispersal seed predators. More recently, researchers suggested that seed masting is one of the means by which plants manipulate behavior of their dispersers (Vander Wall

2010). According to this reasoning, satiation of current energy needs induces granivores to cache

seeds for future use (Vander Wall 2010). Here we show that the effects of masting on population

dynamics and caching behavior are mutually dependent. By decreasing the degree of pilfering, the

satiation-starvation cycle due to more extreme seed masting events may promote the evolution and

maintenance of seed caching behavior. Thus, the decrease in seed predation, increase in per capita

scatterhorder satiation, and reduction in pilfering pressure may each represent an important

pathway by which the scatterhorder satiation-starvation cycle induced by masting may improve plant

recruitment (Fig. 5). These nuanced interactions between plant and seed predator emphasize the

importance of studying the feedbacks between population dynamics and behavioral evolution.

Results of our simulations suggest that when seed production is highly variable, seed caching

can evolve even when cache owners have little advantage over naive foragers in seed recovery

(compare with Krebs and Anderson 1978). The costs of cache loss to pilferers are reduced in our

model because periods of intense seed production coincide with low densities of scatterhoarders –

and thus few potential pilferers (see Dittel and Vander Wall 2018 for experimental data

demonstrating that the magnitude of cache pilferage is determined by the abundance of

scatterhoarders). When there is pronounced masting with relatively long intervals between masting

events, densities of scatterhoarders entering the start of the next large masting event are low (Figs. 2

& 3). Consequently, individuals are able to collect enough seeds to satiate their reproductive needs.

As the yearly fitness is determined by the geometric mean of their fitness across the seasons and this

geometric mean decreases with seasonal variation in fitness (Lewontin & Cohen 1969; Gillespie 1977;

Schreiber 2015), the benefits of reducing seasonal variation in fitness by increasing winter/spring

reproduction outweigh the diminishing returns of increasing reproduction in the fall.

Our results make a prediction that plants dispersed by scatterhoarders should have high

interannual variation of seed production (typically measured with coefficient of variation, CV)

relatively to plants dispersed by other means. This appears to be the case, at least when plants

dispersed by scatterhoarders (synzoochorously) are compared to plants dispersed by frugivores

(endozoochorously) (Herrera et al. 1998). When explaining this pattern, researchers emphasized

contrasting selective pressures acting on these groups of plants. Avoiding the risk of satiating

frugivores was suggested as a factor that stabilizes seed production in plants dispersed

endozoochorously. On the other hand, variable seed production in synzoochorous plants was

interpreted as an adaptation that enabled reducing seed mortality caused by animals that act as seed

predators and only incidentally disperse seeds (Herrera et al. 1998). However, we suggest that the

high CV of plants dispersed by scatterhoarders can also be linked to the caching behavior of

scatterhoarders.

There are several adaptive hypotheses of masting (e.g. increased pollination efficiency,

predator satiation, and seed dispersal: reviews in Kelly 1994; Kelly and Sork 2002; Pearse et al. 2016)

and these putative fitness benefits of masting can occur simultaneously. Thus, it is difficult to

conclude whether variability in seed production evolved (at least partly) to stimulate more seed

caching, or whether seed caching evolved in granivores that interacted with plants that already had

high CV of seed production. Moreover, these two evolutionary pressures could co-occur, creating a

positive feedback between the variability in plant production and granivore caching behavior.

If, as our simulations suggest, masting intensity and mast interval are important for seed

caching, then changes in plant masting patterns might affect the dynamics of seed caching, and

therefore also the abundance of granivores and recruitment in plant populations. Our model is

loosely based on the European beech – *Apodemus* mice system (Jensen 1982; Zwolak et al. 2016).

Several studies have suggested that the European beech shows more frequent masting, probably due

to global warming (Kantorowicz 2000; Schmidt 2006; Overgaard et al. 2007; Paar et al. 2011; but see

Drobyshev et al. 2014). This could shift the beech-rodent interactions towards antagonism, with

more seed consumed and fewer cached (Fig. 2). On the other hand, a recent meta-analysis of global

data suggests that the interannual variation in seed crops is increasing (Pearse et al. 2017). Such a

change could make seed caching more profitable for granivores. However, extreme interannual

variation in seed crops might lead to a decline and even extinction in granivore populations, due to

the difficulty in tracking resource levels (Fig 3).

Moreover, any environmental change that affects scatterhoarder population dynamics could

alter caching behavior and, thereby, impact seed mortality. For example, we found that increased

scatterhoarder survivorship during the winter or summer may select against caching behavior by

increasing population densities entering the masting years (Fig. 4). Thus, changes in winter or

summer conditions that are favorable for mice could harm seedling recruitment both directly by

increasing seed predation and indirectly by discouraging seed caching. In contrast, the effects of

improved conditions in the autumn on seedling recruitment is more difficult to predict as increased

survivorship of scatterhoarders simultaneously selects for more caching and more seed predation

due to higher population densities.

According to a recent review, at least 1,339 species of plants are dispersed by

scatterhoarders (Gómez et al. 2018). Thus, good understanding of the evolution of caching behavior

and of ecological determinants of seed caching is pivotal when our goal is to comprehend former

selective pressures on synzoochorous plants, disentangle current mechanisms of seed dispersal, or to

predict how global changes might alter seed dispersal patterns. Our study provides a step in this

direction and suggests several promising avenues for future research. For example, future work

should address the evolution of caching reaction norms instead of the simple threshold for caching

considered here. Additionally, evolution of caching strategies could be different when individual

variation in personalities or, more generally, phenotypes of seed dispersing animals (Zwolak 2018) is

taken into account. Finally, comparing systems where reciprocal pilfering is common (e.g. rodents:

Vander Wall and Jenkins 1998) versus those where it is rare (e.g. corvids: Pesendorfer et al. 2016)

could provide further insights on the evolution of caching behavior.

Literature Cited

Andersson, M., and J. Krebs. 1978. On the evolution of hoarding behaviour. Animal Behaviour
26:707-711.

Auger, J., S. E. Meyer, and S. H. Jenkins. 2016. A mast-seeding desert shrub regulates population
dynamics and behavior of its heteromyid dispersers. Ecology and Evolution 6:2275-2296.

Balda, R. P., and A. C. Kamil. 1992. Long-term spatial memory in Clark's nutcracker, *Nucifraga
columbiana*. Animal Behaviour 44: 761-769.

Barkley, C. L., and L. F. Jacobs, L. F. 1998. Visual environment and delay affect cache retrieval
accuracy in a food-storing rodent. Animal Learning & Behavior 26:439-447.

Bednekoff, P. A., R. P. Balda, A. C. Kamil, and A. G. Hile. 1997. Long-term spatial memory in four seed-
caching corvid species. Animal Behaviour 53:335-341.

Bogdziewicz, M., R. Zwolak, and E. E. Crone. 2016. How do vertebrates respond to mast seeding?
Oikos 125:300-307.

Briggs, J. S., S. B. Vander Wall, and S. H. Jenkins. 2009. Forest rodents provide directed dispersal of
Jeffrey pine seeds. Ecology 90:675-687.

Curran, L., and M. Leighton. 2000. Vertebrate responses to spatiotemporal variation in seed
production of mast-fruiting Dipterocarpaceae. Ecological Monographs 70:101–128.

Dally, J. M., N. S. Clayton, and N. J. Emery. 2006a. The behaviour and evolution of cache protection
and pilferage. Animal Behaviour 72:13-23.

Dally, J. M., N. J. Emery, and N. S. Clayton. 2006b. Food-caching western scrub-jays keep track of who
was watching when. Science 312:1662-1665.

Dittel, J. W., R. Perea, and S. B. Vander Wall. 2017. Reciprocal pilfering in a seed-caching rodent
community: implications for species coexistence. Behavioral Ecology and Sociobiology 71:147.

Dittel, J. W., and S. B. Vander Wall. 2018. Effects of rodent abundance and richness on cache
pilfering. Integrative Zoology 13:331-338.

Drobyshev, I., M. Niklasson, M. J. Mazerolle, and Y. Bergeron. 2014. Reconstruction of a 253-year

long mast record of European beech reveals its association with large scale temperature

variability and no long-term trend in mast frequencies. Agricultural and Forest Meteorology

192:9-17.

Forget, P. M., and S. B. Vander Wall. 2001. Scatter-hoarding rodents and marsupials: convergent

evolution on diverging continents. Trends in Ecology & Evolution 16:65-67.

Galvez, D., B. Kranstauber, R. W. Kays, and P. A. Jansen. 2009. Scatter hoarding by the Central

American agouti: a test of optimal cache spacing theory. Animal Behaviour 78:1327-1333.

Gillespie, J. H. 1977. Natural selection for variances in offspring numbers: a new evolutionary

principle. American Naturalist, 111:1010-1014.

Gittleman, J. L., and S. D. Thompson. 1988. Energy allocation in mammalian reproduction. American

Zoologist 28:863-875.

Gu, H., Q. Zhao, and Z. Zhang. 2017. Does scatter-hoarding of seeds benefit cache owners or

pilferers? Integrative Zoology 12:477-488.

Gómez, J. M., E. W. Schupp, and P. Jordano. 2019. Synzoochory: the ecological and evolutionary

relevance of a dual interaction. Biological Reviews 94:874-902.

Grodziński, W., and K. Sawicka-Kapusta. 1970. Energy values of tree-seeds eaten by small mammals.

Oikos 1:52-58.

Herrera, C. M., P. Jordano, J. Guitián, and A. Traveset. 1998. Annual variability in seed production by

woody plants and the masting concept: reassessment of principles and relationship to pollination

and seed dispersal. American Naturalist 152:576-594.

Hirsch, B.T., R. Kays, V. E. Pereira, and P. A. Jansen. 2012. Directed seed dispersal towards areas with

low conspecific tree density by a scatter-hoarding rodent. Ecology Letters 15:1423-1429.

Hirsch, B.T., R. Kays, and P. A. Jansen. 2013. Evidence for cache surveillance by a scatter-hoarding

rodent. Animal Behaviour 85:1511-1516.

Jacobs, L. F., and E. R. Liman. 1991. Grey squirrels remember the locations of buried nuts. Animal

Behaviour 41:103-110.

Jacobs, L. F. 1992. Memory for cache locations in Merriam's kangaroo rats. Animal Behaviour 43:585-

593.

Jansen, P. A., F. Bongers, and L. Hemerik. 2004. Seed mass and mast seeding enhance dispersal by a

neotropical scatter-hoarding rodent. Ecological Monographs 74:569-589.

Jansen, P. A., B. T. Hirsch, W. J. Emsens, V. Zamora-Gutierrez, M. Wikelski, and R. Kays. 2012. Thieving

rodents as substitute dispersers of megafaunal seeds. Proceedings of the National Academy of

Sciences 109:12610-12615.

Jensen, T. S. 1985. Seed-seed predator interactions of European beech, *Fagus silvatica* and forest

rodents, *Clethrionomys glareolus* and *Apodemus flavicollis*. Oikos 44:149–156.

Jones, K. E., J. Bielby, M. Cardillo, S. A. Fritz, J. O'Dell, C. D. L. Orme..., and C. Connolly. 2009.

PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently

extinct mammals: Ecological Archives E090-184. Ecology 90:2648-2648.

Kantorowicz, W. 2000. Half a century of seed years in major tree species of Poland. Silvae Genetica

49:245–248.

Kelly, D. 1994. The evolutionary ecology of mast seeding. Trends in Ecology & Evolution 9:465-470.

Kelly, D., and V. L. Sork. 2002. Mast seeding in perennial plants: why, how, where? Annual Review of

Ecology and Systematics 33:427–447.

Koenig, W. D., and J. M. Knops. 2000. Patterns of annual seed production by northern hemisphere

trees: a global perspective. American Naturalist 155:59-69.

Leaver, L. A., L. Hopewell, C. Caldwell, and L. Mallarky. 2007. Audience effects on food caching in grey

squirrels (*Sciurus carolinensis*): evidence for pilferage avoidance strategies. Animal Cognition

10:23-27.

Lewontin, R. C., and D. Cohen. 1969. On population growth in a randomly varying environment.

Proceedings of the National Academy of Sciences 62:1056-1060.

Macdonald, D. W., and F. T. Tattersall. 2001. Britain's mammals - the challenge for conservation.

Peoples Trust For Endangered Species, London.

Meyer, S. E., and B. K. Pendleton. 2015. Evolutionary drivers of mast-seeding in a long-lived desert

shrub. American Journal of Botany 102:1666-1675.

Mendoza, I., R. S. Condit, S. J. Wright, A. Caubère, P. Châtelet, I. Hardy, and P. M. Forget. 2018. Inter-

annual variability of fruit timing and quantity at Nouragues (French Guiana): insights from

hierarchical Bayesian analyses. Biotropica 50:431-441.

Millar, J. S. 1978. Energetics of reproduction in *Peromyscus leucopus*: the cost of lactation. Ecology

59:1055-1061.

Millar, J. S. 1979. Energetics of lactation in *Peromyscus maniculatus*. Canadian Journal of Zoology

57:1015-1019.

Muñoz, A., and R. Bonal. 2011. Linking seed dispersal to cache protection strategies. Journal of

Ecology 99:1016-1025.

Norden, N., J. Chave, P. Belbenoit, A. Caubère, P. Châtelet, P. M. Forget, and C. Thébaud. 2007. Mast

fruiting is a frequent strategy in woody species of Eastern South America. PLoS One 2:e1079.

Ostfeld, R. S., and F. Keesing. 2000. Pulsed resources and community dynamics of consumers in

terrestrial ecosystems. Trends in Ecology & Evolution 15:232-237.

Övergaard, R., P. Gemmel, and M. Karlsson. 2007. Effects of weather conditions on mast year

frequency in beech (*Fagus sylvatica* L.) in Sweden. Forestry, 80:555-565.

Paar, U., A. Guckland, I. Dammann, M. Albrecht, J. Eichhorn. 2011. Häufigkeit und Intensität der

Fruktifikation der Buche [Frequency and intensity of fruiting in beech]. AFZ-DerWald 6:26-29.

Pearse, I. S., W. D. Koenig, and D. Kelly. 2016. Mechanisms of mast seeding: resources, weather,

cues, and selection. New Phytologist 212:546-562.

Pearse, I. S., J. M. LaMontagne, and W. D. Koenig. 2017. Inter-annual variation in seed production has

increased over time (1900–2014). Proceedings of the Royal Society of London B: Biological

Sciences 284:20171666.

Pesendorfer, M. B., T. S. Sillett, W. D. Koenig, and S. A. Morrison. 2016. Scatter-hoarding corvids as

seed dispersers for oaks and pines: a review of a widely distributed mutualism and its utility to

habitat restoration. Condor 118:215-237.

Pravosudov, V. V., T. C. Roth II, and L. D. LaDage. 2010. Chickadees are selfish group members when

it comes to food caching. Animal Behaviour 80:175-180.

Pucek, Z. 1984. Klucz do oznaczania ssaków Polski. Państwowe Wydawnictwo Naukowe, Warszawa.

Pucek, Z., W. Jędrzejewski, B. Jędrzejewska, and M. Pucek. 1993. Rodent population dynamics in a

primeval deciduous forest (Białowieża National Park) in relation to weather, seed crop, and

predation. Acta Theriologica 38:199-232.

Schauber, E. M., D. Kelly, P. Turchin, C. Simon, W. G. Lee, R. B. Allen, I. J. Payton, P. R. Wilson, P. E.

Cowan, and R. E. Brockie. 2002. Masting by eighteen New Zealand plant species: the role of

temperature as a synchronizing cue. Ecology 83:1214-1225.

Schmidt, K. A., and R. S. Ostfeld. 2008. Eavesdropping squirrels reduce their future value of food

under the perceived presence of cache robbers. American Naturalist 171:386-393.

Schmidt, W. 2006. Temporal variation in beech masting (*Fagus sylvatica* L.) in a limestone beech

forest (1981–2004). Allgemeine Forst und Jagdzeitung 177:9-19.

Schreiber, S. J. 2015. Unifying within-and between-generation bet-hedging theories: an ode to JH

Gillespie. American Naturalist 186:792-796.

Shaw, R. C., and N. S. Clayton. 2013. Careful cachers and prying pilferers: Eurasian jays (*Garrulus*

glandarius) limit auditory information available to competitors. Proceedings of the Royal Society

of London B: Biological Sciences 280:20122238.

Shibata, M., H. Tanaka, S. Iida, S. Abe, T. Masaki, K. Niiyama, and T. Nakashizuka. 2002. Synchronized

annual seed production by 16 principal tree species in a temperate deciduous forest, Japan.

Ecology 83:1727-1742.

Sikes, R. S. 1995. Costs of lactation and optimal litter size in northern grasshopper mice (*Onychomys*

leucogaster). Journal of Mammalogy 76:348-357.

Smulders, T. V. 1998. A game theoretical model of the evolution of food hoarding: applications to the

Paridae. American Naturalist 151:356-366.

Steele, M. A., S. L. Halkin, P. D. Smallwood, T. J. McKenna, K. Mitsopoulos, and M. Beam. 2008. Cache

protection strategies of a scatter-hoarding rodent: do tree squirrels engage in behavioural

deception? Animal Behaviour 75:705-714.

Steele, M. A., M. Bugdal, A. Yuan, A. Bartlow, J. Buzalewski, N. Lichti, and R. Swihart. 2011. Cache

placement, pilfering, and a recovery advantage in a seed-dispersing rodent: could predation of

scatter hoarders contribute to seedling establishment? Acta Oecologica 37:554-560.

Steele, M. A., T. A. Contreras, L. Z. Hadj-Chikh, S. J. Agosta, P. D. Smallwood, and C. N. Tomlinson.

2013. Do scatter hoarders trade off increased predation risks for lower rates of cache pilferage?

Behavioral Ecology 25:206-215.

Steele, M. A., G. Rompre, J. A. Stratford, H. Zhang, M. Suchocki, and S. Marino. 2015. Scatterhoarding

rodents favor higher predation risks for cache sites: the potential for predators to influence the

seed dispersal process. Integrative Zoology 10:257-266.

Thayer, T. C., and S. B. Vander Wall. 2005. Interactions between Steller's jays and yellow pine

chipmunks over scatter-hoarded sugar pine seeds. Journal of Animal Ecology 74:365-374.

Vander Wall, S. B. 2001. The evolutionary ecology of nut dispersal. Botanical Review 67:74-117.

Vander Wall, S. B. 2002. Masting in animal-dispersed pines facilitates seed dispersal. Ecology

83:3508-3516.

Vander Wall, S. B. 2010. How plants manipulate the scatter-hoarding behaviour of seed-dispersing

animals. Philosophical Transactions of the Royal Society B 365:989–997.

Vander Wall, S. B., and S. H. Jenkins. 2003. Reciprocal pilferage and the evolution of food-hoarding

behavior. Behavioral Ecology 14:656-667.

Vander Wall, S. B., J. S. Briggs, S. H. Jenkins, K. M. Kuhn, T. C. Thayer, and M. J. Beck. 2006. Do food-

hoarding animals have a cache recovery advantage? Determining recovery of stored food. Animal

Behaviour 72:189-197.

Vander Wall, S. B., C. J. Downs, M. S. Enders, and B. A. Waitman. 2008. Do yellow-pine chipmunks

prefer to recover their own caches? Western North American Naturalist 68:319-324.

Vander Wall, S. B., S. C. Barga, and A. E. Seaman. 2017. The geographic distribution of seed-dispersal

mutualisms in North America. Evolutionary Ecology 31:725-740.

Yang, L. H., K. F. Edwards, J. E. Byrnes, J. L. Bastow, A. N. Wright, and K. O. Spence. 2010. A meta-

analysis of resource pulse–consumer interactions. Ecological Monographs 80:125-151.

Zhu, W., Y. Mu, J. Liu, and Z. Wang. 2015. Energy requirements during lactation in female *Apodemus*

chevrieri (Mammalia: Rodentia: Muridae) in the Hengduan Mountain region, Italian Journal of

Zoology 82:2.

Zwolak, R., M. Bogdziewicz, A. Wróbel, and E. E. Crone. 2016. Advantages of masting in European

beech: timing of granivore satiation and benefits of seed caching support the predator dispersal

hypothesis. Oecologia 180:749-758.

Appendix. Derivation of the fitness gradient

To derive the fitness gradient, assume that the population dynamics of individuals with the resident threshold strategy T converges to a periodic solution of period kP: $n_i(1), n_i(2), \ldots, n_i(p)$ for $i = 1, 2, 3$. The more general case of aperiodic dynamics is discussed at the end of this appendix. For each $1 \leq t \leq kP$, let $G(t)$ be the amount seeds gathered in Fall by each resident individual in year t, and $Q_1(t), Q_2(t), Q_3(t)$ be the number of offspring produced by an individual playing the mutant strategy T_m during the Fall, Winter/Spring, and Summer, respectively, in year t. Then, the long-term per-capita growth rate of the mutant is

$$r(T, T_m) = \frac{1}{kP} \sum_{t=1}^{kP} \sum_{i=1}^{3} \log(Q_i(t) + s_i).$$

Taking the derivative with respect to T_m and evaluating at $T_m = T$ gives us the fitness gradient:

$$\frac{\partial r}{\partial T_m}(T, T) = \frac{1}{kP} \sum_{t=1}^{kP} \sum_{i=1}^{3} \frac{\left.\frac{\partial Q_i(t)}{\partial T_m}\right|_{T_m=T}}{\left.Q_i(t)\right|_{T_m=T} + s_i} \tag{A1}$$

As $\left.Q_i(t)\right|_{T_m=T}$ corresponds to the fecundity of resident individuals,

$$\left.Q_i(t)\right|_{T_m=T} = R_i(t). \tag{A2}$$

It remains to compute the partial derivatives of the $Q_i(t)$ terms. For $i = 1$, we have (from the main text)

$$Q_1(t) = \frac{b \min\{G(t), T_m\}}{h + \min\{G(t), T_m\}}$$

is piecewise defined depending on whether $G(t) < T$ or $> T$. Specifically, we get

$$\left.\frac{\partial Q_1(t)}{\partial T_m}\right|_{T_m=T} = \begin{cases} 0 & \text{if } T > G(t) \\ \frac{bh}{(h+T)^2} & \text{if } T < G(t). \end{cases} \tag{A3}$$

A-1

For $i = 2$, we have (from the main text)

$$Q_2(t) = \frac{bC_{\text{m,survivor}}(t)}{h + C_{\text{m,survivor}}(t)} \frac{s_1}{Q_1(t) + s_1} + \frac{bC_{\text{new}}(t)}{h + C_{\text{new}}(t)} \frac{Q_1(t)}{Q_1(t) + s_1}$$

where $Q_2(t)$ differs from $R_2(t)$ only in its first term due to surviving individuals with the mutant caching strategy:

$$C_{\text{m,survivor}}(t) = \max\{G(t) - T_m, 0\}M(t) + \max\{G(t) - T, 0\} \left(O(t)(s_1 n_1(t) - 1) + D(t)(1 - s_1)n_1(t)\right).$$

Hence, by chain rule,

$$\frac{\partial Q_2(t)}{\partial T_m}\bigg|_{T_m=T} = \frac{bh}{(h + C_{\text{m,survivor}}(t))^2} \frac{s_1}{Q_1(t) + s_1} \frac{\partial C_{\text{m,survivor}}(t)}{\partial T_m}\bigg|_{T_m=T}.$$

At $T_m = T$, we have $C_{\text{m,survivor}}(t) = C_{\text{survivor}}(t)$ and $Q_1(t) = R_1(t)$. Furthermore,

$$\frac{\partial C_{\text{m,survivor}}(t)}{\partial T_m}\bigg|_{T_m=T} = \begin{cases} 0 & \text{if } T > G(t) \\ -M(t) & \text{if } T < G(t). \end{cases}$$

Thus,

$$\frac{\partial Q_2(t)}{\partial T_m}\bigg|_{T_m=T} = \begin{cases} 0 & \text{if } T > G(t) \\ -\frac{bh}{(h+C_{\text{survivor}}(t))^2} \frac{s_1}{R_1(t)+s_1} M(t) & \text{if } T < G(t). \end{cases} \tag{A4}$$

Finally, as discussed in the main text, $Q_3(t) = R_3(t)$ and, consequently, doesn't depend on T_m. Hence,

$$\frac{\partial Q_3(t)}{\partial T_m}\bigg|_{T_m=T} = 0. \tag{A5}$$

Substituting equations (A2)–(A5) into (A1) provides an explicit expression for the fitness gradient.

These calculations also apply to the more general case when the resident dynamics are asymptotically stationary but not necessarily periodic. By asymptotic stationarity, we mean that there

exists a probability measure μ on the (S, n_1) state space such that

$$\lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} f(S(t), n_1(t)) = \int f(S, n_1)\mu(dS, dn_1)$$

for all continuous functions $f(S, n_1)$ i.e. the Birkhoff averages converge. In this case, the per-capita growth rate of the mutant is well-defined and equals

$$r(T, T_m) = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} \sum_{i=1}^{3} \log(Q_i(t) + s_i).$$

The selection gradient equals

$$\frac{\partial r}{\partial T_m}(T, T) = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} \sum_{i=1}^{3} \frac{\frac{\partial Q_i(t)}{\partial T_m}\big|_{T_m=T}}{Q_i(t)\big|_{T_m=T} + s_i} \tag{A6}$$

and can be explicitly evaluated by using (A2)-(A5).