Multivariate phenotypic divergence along an urbanization gradient

Authors: James S. Santangelo[1, 2, 3*], Carole Advenard[4], and L. Ruth Rivkin[1, 2, 3], Ken A.

Thompson[5]

Affiliations:

1. Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street,

Toronto, Ontario M5S 3B2, Canada.

2. Department of Biology, University of Toronto Mississauga, 3359 Mississauga Road,

Mississauga, Ontario, L5L 1C6, Canada.

3. Centre for Urban Environments, University of Toronto Mississauga, 3359 Mississauga Road,

Mississauga, Ontario, L5L 1C6, Canada.

4. AgroSup Dijon, Dijon 21000, Burgundy, France

5. Department of Zoology & Biodiversity Research Centre, University of British Columbia,

Vancouver, BC, Canada.

*Corresponding author: Email: james.santangelo37@gmail.com

Running title: Multivariate trait evolution in cities

Acknowledgements

We thank M. Hetherington-Rauth for discussions that motivated this work. A. Dao, B. Cohan,

M. Escobar, J. Viliunas, L. Miles, and S. Innes provided valuable assistance in data collection.

We thank M. Kalich and B. Pitton for assistance and space in the greenhouse. Comments from

the EvoEco lab at UofT Mississauga and D. Schluter's group at UBC greatly improved the

manuscript. Authors received funding from NSERC (NSERC PGS-D to JSS and NSERC CGS-D

to LRR and KAT) and the Izaak Walton Killam Foundation (KAT). An NSERC Discovery Grant

to Marc T. J. Johnson provided the supplies, equipment and greenhouse space rental used to

carry out this project.

Author Contributions

32   JSS and KAT planned the study and collected original plant material, and all authors contributed

33   to subsequent study design. JSS wrote the first draft of the paper with sections written by KAT

34   and LRR, and all authors contributed to revisions. CA conducted the common garden experiment

35   and collected data, with assistance from all other authors. JSS oversaw field data collection, with

36   assistance from other authors. JSS analyzed the data with input from KAT and LRR.

Data accessibility

39   All code and data used throughout this manuscript are presently available for reviewers on the

40   GitHub page for J.S.S (https://github.com/James-S-Santangelo/SIC) and will be permanently

41   archived in a data repository following publication.

Conflicts of interest

The authors declare no conflicts of interest.

Abstract

A growing body of evidence suggests that natural populations can evolve to better tolerate the

novel environmental conditions associated with urban areas. Invariably, studies of adaptive

divergence in urban areas examine only one or a few traits at a time from populations residing

only at the most extreme urban and nonurban habitats. Thus, whether urbanization is driving

divergence in many traits simultaneously in a manner that varies with the degree of urbanization

remains unclear. To address this gap, we generated seed families of white clover (*Trifolium*

repens) collected from 27 populations along an urbanization gradient in Toronto, Canada, and

grew them up to measure multiple phenotypic traits in a common garden. Overall, urban

populations had later phenology and germination, larger flowers, thinner stolons, reduced

cyanogenesis, greater biomass, and were more attractive to pollinators. Pollinator observations

revealed near complete turnover between urban and nonurban sites, which may explain some of

the observed divergence in floral traits and phenology. Our results suggest that adaptation to

urban environments involves multiple organismal traits.

Keywords: Anthropocene; cline; urban evolution; rapid evolution; *Trifolium repens*

Introduction

Urbanization is rapidly changing the face of the planet. As cities develop, natural habitats

experience drastic environmental changes, from increased temperatures and pollution to greater

impervious surface and habitat fragmentation [1]. Evidence is accumulating to support the

hypothesis that the environmental features associated with urbanization drive phenotypic

differences between populations in urban and nonurban habitats [2]. For example, urban *Anolis*

lizards have evolved longer limbs and more toe lamellae to improve sprint speed on the smooth

artificial surfaces common in cities [3,4]. Increased impervious surfaces often lead to warmer air

temperatures in cities [5], which has driven increases in thermal tolerance of urban acorn ant [6]

and *Daphnia* populations [7]. These studies, and others (see table S1 in [8]) suggest that many

urban populations are adapting to the novel environments created by humans.

Despite accumulating evidence of evolution in response to urban-driven environmental

change, most studies focusing on phenotypic divergence associated with urbanization have

examined just one or a few traits at a time [4,9–12]. However, theoretical and empirical work in

other systems suggest that selection can drive the evolution of multivariate phenotypic clines

along environmental gradients [13,14]. Multivariate phenotypic analyses incorporate multiple,

often correlated, traits to explain overarching shifts in phenotypes across environments.

Although we can make informed predictions about how particular traits might respond to

urbanization from work in other systems, understanding how cities are driving phenotypic

evolution requires quantifying divergence in many traits that simultaneously influence fitness. It

is thus presently unclear what suites of traits are most often favored as populations become more

urbanized.

Here, we investigate multivariate phenotypic evolution along an urbanization gradient in

Toronto, Canada. Traits involved in plant reproduction are particularly likely to show

multivariate, genetically-based phenotypic associations with urbanization between urban and

nonurban populations due to the direct effect of these traits on fitness. In animal-pollinated

plants, urbanization might impact fitness through changes to their pollinator communities. Both

the diversity and abundance of pollinators are known to change along urbanization gradients

[15,16], with positive [17–19] to negative [20] effects on pollinator visitation. Variation in

pollination along urbanization gradients might drive changes in the extent of pollen limitation

experienced by outcrossing plants [21,22], which can influence the strength and opportunity for

pollinator-mediated selection on plant reproductive traits [23–25]. Consequently, we expect that

plants could compensate for the altered pollination environment in cities by evolving altered

reproductive trait values relative to non-urban plants.

Methods

Detailed methods can be found in the online supplementary materials.

Common garden experiment

We examined multivariate trait divergence along an urbanization gradient using white clover

(*Trifolium repens*) as a model system. We grew 642 F1 generation white clover plants from seed

in a common garden. These plants were distributed among 209 plant families' from 27

populations spanning an urban-rural transect in Toronto, Ontario, Canada (Fig. 1a, [10]). The

garden was located at the University of Toronto Mississauga in summer 2017. We measured

several traits during our experiment, some of which are known to be under selection in this

system (Table 1) [26,27].

Field observations

We quantified variation in pollination among our study populations to identify possible

mechanisms that might underlie phenotypic divergence along the urbanization gradient. We

conducted pollinator observations in all study populations using two 1 m × 1 m quadrats (similar

to [28]) placed in full sun and counting the number of inflorescences visited by pollinators over

20 minutes. We classified pollinators into three morphological groups (termed *morphs* hereafter):

honeybees (*Apis mellifera*), bumblebees (*Bombus spp.*) and sweat bees (Halictidae).

We additionally collected twenty ripe infructescences (i.e., group of fruit) from the same

populations in which we recorded pollinator observations. We counted the number of flowers

and seeds in each infructescence and used the number of seeds per flower as a metric of pollen

limitation (fewer seeds per flower = more pollen limitation). We generated data for the same

metric from plants in the common garden. These data provide information about *in situ* variation

in pollen limitation (field-collected inflorescences) and evolved differences in plants' abilities to

set seed from pollen (common garden data).

Data analysis

We investigated whether urbanization was associated with multivariate phenotypic

divergence using canonical redundancy analysis (i.e., RDA, [29]). The RDA regressed a matrix

of standardized family mean trait values (209 families × 14 traits) as a response variable against

a numeric vector representing the distance (in km) of populations to the urban core as the sole

predictor (see below), which is correlated with % impervious surface cover ($r_{pearson}$ = −0.63,

Fig. S1a) and human population density ($r_{pearson}$ = −0.72, Fig. S1a), has previously been shown to

be correlated with cyanogenesis frequency ($r_{pearson}$= 0.63) in the same species and region [10].

The RDA tests whether distance to the urban center explains more phenotypic variation

than expected by chance. In principle, we could observe a significant RDA even if just one trait

was associated with distance, but our interest lies in understanding whether multiple traits are

evolving together along the transect (i.e., including traits that do not vary significantly on their

own). Therefore, we used the canonical coefficients from the RDA that describe the individual

contribution of phenotypic traits to the first constrained axis of the RDA (RDA1 [i.e., distance])

to calculate a multivariate phenotype score for each individual. This score, referred to as

cline$_{max}$, is the multivariate quantitative trait that shows the strongest association with distance to

the urban center [13]. Regressing **cline$_{max}$** against distance to the urban center enables us to better

visualize the shape of the multivariate cline and to quantify how multivariate phenotypes are

changing along our urbanization gradient.

To examine variation in pollinator visitation among natural populations, we fit a linear

model with the number of visits per inflorescence as a response variable, and distance, pollinator

morph, and their interaction as predictors. To assess how the number of seeds per flower varied

among field-collected inflorescences and plants in the common garden, we fit a model with

population-mean number of seeds per flower as the response and distance, source (field-collected

vs. common garden-collected), and their interaction as predictors. All data (see **Data**

Accessibility) were analyzed in R version 3.6.0 [31] within the RStudio environment [32].

Results

Common garden

Urbanization was associated with multivariate trait divergence and the evolution of a

multivariate phenotypic cline in Toronto, ON (Fig. S2 and Fig. 1b). Distance to the urban core

explained 2.7% of the total variation in the multivariate phenotypic composition of populations

(RDA, $F_{1, 202} = 5.51$, $P < 0.001$, Fig. S2, Fig. S3). The composite trait showing the strongest

association with distance to the urban core (i.e., **cline$_{max}$**) showed a positive cline along the

urbanization gradient ($\beta = 0.01$, $t_{206} = 9.32$, $P < 0.001$, $r^2 = 0.29$, Fig. 1b) that was at least as

strong as any trait when analyzed individually (Fig. S4 and S5, see also standardized beta

coefficients in table S1). The six traits that loaded most strongly (|loading| > 0.3, bolded in Fig.

S2) onto the first axis of the RDA (RDA1)—germination, phenology, flower size, biomass,

HCN, and stolon thickness—and contributed most to the composite trait showing the strongest

association with distance all showed significant univariate clines in the direction predicted based

on their trait loadings (Fig. S2 and Fig. S6a).

Field observations

On average, pollinator visitation per inflorescence was greatest in urban populations (Distance: β

$= -0.032$, $F_{1, 75} = 21.96$, $P < 0.001$; thick black line in Fig. 2a). Visitation rate varied with

pollinator morphs (Morph: $F_{2, 75} = 11.06$, $P < 0.001$): bumblebees had the highest visitation rate

(average $= 0.63 \pm 1.2$ SD visits/inflorescence across all populations), followed by honey bees

(average $= 0.58 \pm 1.05$ visits/inflorescence), and sweat bees (average $= 0.18 \pm 0.41$

visits/inflorescence). Pollinator morphs varied in their response to urbanization (Distance ×

Morph interaction: $F_{2, 75} = 14.11$, $P < 0.001$, Fig. 2a): bumblebee visitation rate was greatest in

urban populations, whereas honeybee visitation was greatest in nonurban populations, and sweat

bees showed little change in visitation across the urbanization gradient (Fig. 2a).

While common garden infructescences were on average more thoroughly pollinated than

those in natural populations (i.e., more seeds per flower; source $F_{1,\,50} = 30.04$, $P < 0.001$,

$mean_{field} = 1.1$, $mean_{garden} = 2.5$, Fig. 2b), the number of seeds per flower was highest in urban

populations (Distance effect: $\beta = -0.013$, $F_{1,\,50} = 5.48$, $P = 0.03$, Fig. 2b). Urban plants were

more pollinated than non-urban plants in both the common garden and in the field (Distance $\times$

source interaction: $F_{1,50} = 0.99$, $P = 0.33$, Fig. 2b).

Discussion

Multivariate phenotypic divergence and clinal variation

Most studies of population differentiation and clinal variation along environmental gradients

focus on only one or a few traits. We know much less about how suites of traits evolve in concert

in response to environmental factors [13,14]. Our study shows clear phenotypic divergence and

clinal variation in multivariate phenotypes along an urbanization gradient. After weighting traits

based on their loadings onto RDA1 [13], distance explained 29% of the variation in the

multivariate phenotype **clinemax**. Urban populations had later germination and flowering (Fig.

S4a and S4b), greater vegetative biomass (Fig. S4c), larger banner petals (Fig. S4d), thinner

stolons (Fig. S4e), and lower HCN frequencies (Fig. S4f). Many of the traits (three of six) most

strongly associated with distance to the urban core were reproductive traits, supporting the

prediction that these traits are likely to show evolutionary divergence between urban and

nonurban habitats, although some notable traits (e.g. reproductive biomass, number of flowers,

and number of inflorescences) showed no association with urbanization.

Urban evolution in white clover

Correlational data suggests that lower cyanogenesis in urban populations has evolved due to

selection for tolerance to freezing damage [10]. Recent data suggest a cost to producing the two

metabolic component of HCN (cyanogenic glucosides and linamarase) under stressful conditions

(e.g., frost) [33], which may partially explain the lower HCN frequencies in urban populations.

Furthermore, small-leaved plants and those with thinner stolons are more frost-tolerant [34,35],

suggesting plants in urban environments should match these phenotypes if they have indeed

evolved to better tolerate frost. Our data do not support the hypothesis that urban populations

have evolved smaller leaves but do support urban plants having thinner stolons (Fig. S4e, table

S1), offering partial support to the hypothesis that urban populations may be more frost tolerant.

Plant evolution in urban environments

Our observed differences in flower size corroborate earlier findings of stronger directional

selection for larger flowers in urban populations [36,37]. In addition, plants from urban

populations were larger than those in nonurban populations, similar to urban and nonurban

Lepidium virginicum plants grown in a common garden [12]. However, increased vegetative

biomass in urban populations is not universal; in ragweed (*Ambrosia artemisiifolia*), there were

no differences in plant size between urban and nonurban populations, although urban populations

flowered earlier than nonurban populations [11]. This contrasts with our experiment in which

urban populations flowered later than nonurban populations. These results suggest that the

effects of urbanization on plant traits vary across species and there currently appears to be no

particular combinations of traits consistently favored in cities.

Urban populations of white clover were less pollen limited and were visited primarily by

bumblebees whereas nonurban populations were visited primarily by honeybees. A single

bumblebee typically visits more white clover flowers per minute than a honeybee [38],

potentially explaining the greater seed set of urban plants, and consistent with other work

showing higher seed set among bumblebee-pollinated urban white clover plants [28]. The

difference in pollinator community between urban and nonurban populations might explain some

of the differences in floral size that have evolved if these pollinators have divergent preferences

or interactions with the flowers [36,39]. In support of this hypothesis, we found that both urban

common garden and field-collected plants set more seed than nonurban plants, suggesting

evolved increases in seed set among urban plants which may be due to increased attractiveness to

the most abundant urban pollinators. The increased pollen limitation of urban plants might have

helped drive the shifts towards—or have resulted from—larger flowers in urban populations.

Conclusion

Our results suggest that natural selection imposed by urbanization is sufficiently strong and

multifarious to have driven the rapid evolution of a multivariate phenotypic cline in white clover

in a large metropolitan area. The data available demonstrate variable responses of different plant

taxa to urbanization, suggesting there is likely no 'one size fits all' solution to life in the city.

Many of the traits showing divergence between urban and nonurban populations are involved in

plant reproduction, and some of this divergence might be due to variation in the pollination

environment. Together, our results suggest that natural selection in urban environments is rapidly

refining whole-organism phenotypes to facilitate adaptation to cities.

References

1. McDonnell MJ, Hahs AK, Breuste JH. 2009 *Ecology of cities and towns: a comparative approach*. Cambridge, UK: Cambridge University Press.
2. Johnson MTJ, Munshi-South J. 2017 Evolution of life in urban environments. *Science.* **8327**. (doi:10.1126/science.aam8327)
3. Winchell KM, Maayan I, Fredette J, Revell LJ. 2018 Linking locomotor performance to morphological shifts in urban lizards. *Proc. R. Soc. Biol. Sci.* **285**, 20180229. (doi:10.1098/rspb.2018.0229)
4. Winchell KM, Reynolds RG, Prado-Irwin SR, Puente-Rolón AR, Revell LJ. 2016 Phenotypic shifts in urban areas in the tropical lizard *Anolis cristatellus*. *Evolution.* **70**, 1009–1022. (doi:10.1111/evo.12925)
5. Oke TR. 1973 City size and the urban heat island. *Atmos. Environ. Pergamon Pres* **7**, 769–779. (doi:10.1016/0004-6981(73)90140-6)
6. Diamond SE, Chick LD, Perez A, Strickler SA, Martin RA. 2018 Evolution of thermal tolerance and its fitness consequences: parallel and non-parallel responses to urban heat islands across three cities. *Proc. R. Soc. B Biol. Sci.* **285**, 20180036.
7. Brans KI, Jansen M, Vanoverbeke J, Tuzun N, Stoks R, De Meester L. 2016 The heat is on: genetic adaptation to urbanization mediated by thermal tolerance and body size. *Glob. Chang. Biol.* **23**, 5218-5227. doi: 10.1111/gcb.13784. (doi:10.1111/ijlh.12426)
8. Rivkin LR *et al.* 2019 A roadmap for urban evolutionary ecology. *Evol. Appl.* **12**, 384–398. (doi:10.1111/eva.12734)
9. Cheptou P-O, Carrue O, Rouifed S, Cantarel A. 2008 Rapid evolution of seed dispersal in an urban environment in the weed *Crepis sancta*. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 3796–3799. (doi:10.1073/pnas.0708446105)
10. Thompson KA, Renaudin M, Johnson MTJ. 2016 Urbanization drives the evolution of parallel clines in plant populations. *Proc. R. Soc. B Biol. Sci.* **283**, 20162180. (doi:10.1098/rspb.2016.2180)
11. Gorton AJ, Moeller DA, Tiffin P. 2018 Little plant, big city: a test of adaptation to urban environments in common ragweed (*Ambrosia artemisiifolia*). *Proc. R. Soc. B Biol. Sci.* **285**, 20180968.
12. Yakub M, Tiffin P. 2017 Living in the city: urban environments shape the evolution of a native annual plant. *Glob. Chang. Biol.* **23**, 2082–2089. (doi:10.1111/gcb.13528)
13. Stock AJ, Campitelli BE, Stinchcombe JR. 2014 Quantitative genetic variance and multivariate clines in the Ivy leaf morning glory, *Ipomoea hederacea*. *Philos. Trans. R. Soc. B Biol. Sci.* **369**, 20130259.
14. Lohman BK, Berner D, Bolnick DI. 2017 Clines arc through multivariate morphospace. *Am. Nat.* **189**, 354–367. (doi:10.1086/690808)
15. Larson JL, Kesheimer AJ, Potter DA. 2014 Pollinator assemblages on dandelions and white clover in urban and suburban lawns. *J. Insect Conserv.* **18**, 863–873. (doi:10.1007/s10841-014-9694-9)
16. Bates AJ, Sadler JP, Fairbrass AJ, Falk SJ, Hale JD, Matthews TJ. 2011 Changing bee and hoverfly pollinator assemblages along an urban-rural gradient. *PLoS One* **6**, e23459. (doi:10.1371/journal.pone.0023459)
17. Theodorou P, Radzevičiūtė R, Settele J, Schweiger O, Murray ET, Paxton RJ. 2016 Pollination services enhanced with urbanisation despite increasing pollinator parasitism.

Proc. R. Soc. B Biol. Sci. **283**, 20160561. (doi:rspb.2016.0561)

18. Desaegher J, Nadot S, Dajoz I, Colas B. 2017 Buzz in Paris: flower production and plant–
pollinator interactions in plants from contrasted urban and rural origins. *Genetica* **145**,
513–523. (doi:10.1007/s10709-017-9993-7)

19. Theodorou P, Albiga K, Radzevičiūtė R, Settele J, Schweiger O, Murray TE, Paxton RJ.
2016 The structure of flower-visitor networks in relation to pollination across an
agricultural to urban gradient. *Funct. Ecol.* **31**, 838–847. (doi:10.1111/1365-2435.12803)

20. Pellissier V, Muratet A, Verfaillie F, Machon N. 2012 Pollination success of *Lotus
corniculatus* (L.) in an urban context. *Acta Oecologica* **39**, 94–100.
(doi:10.1016/j.actao.2012.01.008)

21. Johnson MTJ, Thompson KA, Saini HS. 2015 Plant evolution in the urban jungle. *Am. J.
Bot.* **102**, 1951–1953. (doi:10.3732/ajb.1500386)

22. Eckert CG *et al.* 2009 Plant mating systems in a changing world. *Trends Ecol. Evol.* **25**,
35–43. (doi:10.1016/j.tree.2009.06.013)

23. Trunschke J, Sletvold N, Ågren J. 2017 Interaction intensity and pollinator-mediated
selection. *New Phytol.* **214**, 1381–1389. (doi:10.1111/nph.14479)

24. Caruso CM *et al.* 2017 What are the environmental determinants of phenotypic selection?
A meta-analysis of experimental studies. *Am. Nat.* **190**, 363–376. (doi:10.1086/692760)

25. Sletvold N, Ågren J. 2016 Experimental reduction in interaction intensity strongly affects
biotic selection. *Ecology* **97**, 3091–3098. (doi:10.1002/ecy.1554)

26. Santangelo JS, Thompson KA, Johnson MTJ. 2018 Herbivores and plant defenses affect
selection on plant reproductive traits more strongly than pollinators. *J. Evol. Biol.* **32**, 4–
18. (doi:10.1111/jeb.13392)

27. Thompson KA, Johnson MTJ. 2016 Antiherbivore defenses alter natural selection on plant
reproductive traits. *Evolution.* **70**, 796–810. (doi:10.1017/CBO9781107415324.004)

28. Verboven HAF, Aertsen W, Brys R, Hermy M. 2014 Pollination and seed set of an
obligatory outcrossing plant in an urban-peri-urban gradient. *Perspect. Plant Ecol. Evol.
Syst.* **16**, 121–131. (doi:10.1016/j.ppees.2014.03.002)

29. Légendre P, Légendre L. 2012 *Numerical ecology*. Third. Amsterdam, Netherlands:
Elsevier.

30. Johnson MTJ, Prashad C, Lavoignat M, Saini HS. 2018 Contrasting the effects of natural
selection, genetic drift and gene flow on urban evolution in white clover (*Trifolium
repens*). *Proc. R. Soc. B Biol. Sci.* **285**, 20181019.

31. R Core Team. 2018 R: A language and environment for statistical computing. *R Found.
Stat. Comput.* See https://www.r-project.org/.

32. RStudio Team. 2016 RStudio: Integrated development environment for R.

33. Kooyers NJ, Hartman Bakken B, Ungerer MC, Olsen KM. 2018 Freeze-induced cyanide
toxicity does not maintain the cyanogenesis polymorphism in white clover (*Trifolium
repens*). *Am. J. Bot.* **105**, 1224–1231. (doi:10.1002/ajb2.1134)

34. Caradus JR, Mackay AC, van den Bosch J. 1989 Intraspecific variation for frost hardiness
in white clover. *J. Agric. Sci.* **112**, 151–157.

35. Svenning MM, Røsnes K, Junttila O. 1997 Frost tolerance and biochemical changes
during hardening and dehardening in contrasting white clover populations. *Physiol. Plant.*
101, 31–37.

36. Irwin RE, Warren PS, Adler LS. 2018 Phenotypic selection on floral traits in an urban
landscape. *Proc. R. Soc. B Biol. Sci.* **285**, 20181239.

37. Bode RF, Tong R. 2018 Pollinators exert positive selection on flower size on urban, but
not on rural Scotch broom (*Cytisus scoparius*). *J. Plant Ecol.* **11**, 493–501.
(doi:10.1093/jpe/rtx024)
38. Howlett BG, Lankin-Vega GO, Jesson LK. 2018 Bombus terrestris: a more efficient but
less effective pollinator than Apis mellifera across surveyed white clover seed fields. *New*
Zeal. J. Crop Hortic. Sci. **47**, 1–16. (doi:10.1080/01140671.2018.1466341)
39. Irwin RE, Warren PS, Carper AL, Adler LS. 2014 Plant–animal interactions in suburban
environments: implications for floral evolution. *Oecologia* **174**, 803–815.
(doi:10.1007/s00442-013-2797-2)
40. Center for Internation Earth Science Information (CIESIN). 2016 Gridded population of
the world, version 4 (GPWv4). (doi:http://dx.doi.org/10.7927/H46T0JKB)
348

Tables

Table 1: Traits (with units) measured throughout common garden experiment.

Trait	Unit
Flowing phenology	# days to first flower
Banner petal length	mm
Banner petal width	mmm
# of flowers per inflorescence	Count
# of inflorescences	Count
Defensive phenotype	HCN+ (1) or HCN− (0)
Leaf width	mm
Leaf length	mm
Stolon diameter	mm
Petiole length	mm
Peduncle length	mm
Vegetative biomass	g
Reproductive biomass	g
Relative investment in sexual reproduction	Ratio of reproductive biomass and vegetative biomass

Figures

(a)

Fig. 1: (a) 27 populations from which we collected plants (orange points). Points are scaled by population density (from [40]) within 1 km^2. The city center (yellow star, Yonge-Dundas Square, lat: 43.6561, long: −79.3803) and common garden location (green square, lat: 43.5494, long: −79.6625) are also shown. Inset: Photograph of the common garden. (b) **cline$_{max}$** scores plotted against distance to the urban core (km).

Fig. 2: (a) Mean number of visits per inflorescence against the distance of the population from the urban core (solid black line). We also show a smoothing line (loess line) for bumblebees (red dashed), honeybees (yellow dotted), and sweat bees (blue dots/dashes) separately to illustrate the change in community composition (also see Fig. S7 for point with linear fits). (b) Number of seeds per fruit (a metric of pollen limitation) among field-collected infructescences (grey-filled circles, dashed line) and infructescences from common garden plants (white triangles, dotted line) from these same populations. The thick black line shows the overall decrease in the number of seeds per flower with increasing distance from the urban center.