Title

Synthetic mycobacterial molecular patterns partially complete Freund's adjuvant

Authors/Affiliations

Jean-Yves Dubé [1,2,3*], Fiona McIntosh [2,3], Juan G. Zarruk [4], Samuel David [4], Jérôme Nigou [5], Marcel A. Behr [1,2,3,6*]

1, Department of Microbiology and Immunology, McGill University. 2, Infectious diseases and Immunity in Global Health Program, Research Institute of the McGill University Health Centre. 3, McGill International TB Centre. 4, Centre for Research in Neuroscience, Research Institute of the McGill University Health Centre. 5, Institut de Pharmacologie et de Biologie Structurale, Université de Toulouse, CNRS, Université Paul Sabatier, Toulouse, France. 6, Department of Medicine, McGill University Health Centre, Montréal, Canada.

Contact Info*
jean-yves.dube@mail.mcgill.ca
marcel.behr@mcgill.ca (Lead Contact)
McGill University Health Centre
1001 boul. Décarie
Glen Site Block E, Office #EM3.3212
Montréal, Québec, Canada
H4A 3J1

Abstract

Complete Freund's adjuvant (CFA) has historically been one of the most useful tools of immunologists. Essentially comprised of dead mycobacteria and mineral oil, we asked ourselves what is special about the mycobacterial part of this adjuvant, and could it be recapitulated synthetically? Here, we demonstrate the essentiality of *N*-glycolylated peptidoglycan plus trehalose dimycolate (both unique in mycobacteria) for the complete adjuvant effect using knockouts and chemical complementation. A combination of synthetic *N*-glycolyl muramyl dipeptide and minimal trehalose dimycolate motif GlcC14C18 was able to upregulate dendritic cell effectors, plus induce experimental autoimmunity qualitatively similar but quantitatively milder compared to CFA. This research outlines how to substitute CFA with a consistent, molecularly-defined adjuvant which may inform the design of immunotherapeutic agents and vaccines benefitting from cell-mediated immunity. We also anticipate using synthetic microbe-associated molecular patterns (MAMPs) to study mycobacterial immunity and immunopathogenesis.

Introduction

Infection with *Mycobacterium tuberculosis*, or administration of bacille Calmette-Guérin (BCG), normally leads to cell-meditated immunity (CMI) to the corresponding bacterial antigens. The tuberculin skin test is positive when the immune response, a type IV hypersensitivity reaction or delayed-type hypersensitivity (DTH), occurs to tuberculin (a protein extract of *M. tuberculosis*). This "cutaneous sensitivity" was closely examined in the 1940s using complete Freund's adjuvant (CFA, heat-killed *M. tuberculosis* in mineral oil plus surfactant) [1,2]. These studies provided the first direct evidence for the cellular nature of DTH by transfer from a CFA-immunized guinea pig to a naïve one only through the washed, heat-liable cellular fraction of peritoneal exudates [1,2]. It is now known that DTH is mediated specifically by antigen-sensitive T cells.

Today, CFA is a 'gold standard' adjuvant for eliciting CMI in research models of autoimmune disease. Notable is the experimental autoimmune encephalomyelitis (EAE) model of T-cell meditated destruction of myelin causing ascending paralysis, used most often to model multiple sclerosis [3,4]. CFA is not used in humans because of high reactogenicity [5]. We have asked ourselves: what was the impetus for Jules Freund to develop his eponymous adjuvant with *M. tuberculosis*? For those conducting TB research, it is well appreciated that handling *M. tuberculosis* (a slow growing, clumping, fastidious, and lethally pathogenic organism) is a significant task *per se*. So, why did Freund choose to incorporate these bacteria?

Effort has been made to describe the microbe-associated molecular patterns (MAMPs) in mycobacteria that drive the adjuvant effect. Evidence has pointed to mycobacterial peptidoglycan (PGN), specifically down to the molecular motif muramyl dipeptide (MDP) [6,7]. Mycobacteria are distinct in that they produce the *N*-glycolyl MDP motif in their cell wall [8-10] and this PGN modification has been shown to be more potent compared to the common *N*-acetyl MDP motif possessed by other bacteria [7,11]. MDP is thought to be recognized through the host molecule NOD2 [12], and mutations in NOD2 predispose humans to increased risk of mycobacterial and inflammatory diseases [13-16]. Alternatively, others have pointed to the mycobacterial cell wall lipid trehalose-6,6-dimycolate (TDM) alone or synergistically with purified peptidoglycan [17]. TDM is recognized by the host with the C-type lectin Mincle in concert with FcRγ and MCL [18,19]. TDM has been demonstrated in animal models to alone be sufficient for granuloma formation and immunopathological responses [18,20].

Both *N*-glycolyl MDP and TDM are MAMPs unique to mycobacteria. Their role in mycobacterial immune responses is supported by the literature and their host receptors are well-known. Additionally, *N*-glycolyl MDP is producible synthetically [7,21,22]. Recently a minimal motif of TDM called GlcC14C18 was produced synthetically and shown to retain adjuvancy but with minimal toxicity [23]. As a complex biologic, CFA is subjected to batch inconsistency. We hypothesized that it is possible to create an entirely synthetic CFA using a rational approach to identify essential MAMPs to replace the whole mycobacteria in the adjuvant. In this work, we establish the necessity for *N*-glycolylation of PGN, NOD2 and Mincle for full CFA adjuvancy. We also demonstrate that the mycobacteria in CFA can be replaced with synthetic *N*-glycolyl MDP and GlcC14C18 in combination to partially restore the adjuvant effect in both a murine

92   model of antigen-specific T-cell immunity as well as the EAE model of autoimmune ascending
93   paralysis.
94
95   **Results**
96
97   <u>*N*-glycolylation of PGN by mycobacteria is required for complete adjuvancy</u>
98
Previous work indicated mycobacteria required *N*-glycolylated PGN to elicit a maximal
immune response during live infection [7,11]. To determine if CMI elicited from dead
mycobacteria in the context of Freund's adjuvant similarly required *N*-glycolylated PGN, we
prepared in-house complete Freund's adjuvant with heat-killed *M. tuberculosis* strain H37Rv,
and H37Rv Δ*namH*. NamH is the enzyme responsible for *N*-glycolylation of PGN units as they
are being synthesized in the cytoplasm, and the Δ*namH* mutant was previously shown to be
devoid of *N*-glycolylation [11] (fig. 1A). We next immunized mice against ovalbumin (OVA, an
exemplary antigen) with our CFAs or incomplete Freund's adjuvant (IFA, lacks mycobacteria),
and examined OVA-specific cytokine production by CD4+ T-cells from the draining lymph
nodes seven days post-immunization (fig. 1B; gating in fig. S1A). Both mycobacteria and *namH*
were required to generate the highest proportion of OVA-specific IFN-γ-producing CD4+ T cells
(fig. 1C). OVA-specific IL-17A-producing cells required mycobacteria but did not significantly
depend on *namH* (fig. 1D). Similar results were seen when examining total number of cytokine-
producing CD4+ T cells in the lymph nodes (fig. S2A-B), and a closer analysis of the pooled
data showed that *namH* contributes to about one third of mycobacteria-induced antigen-specific
IFN-γ-producing Th cells (fig. S2C-D). The individual results of experiments are shown
separately in fig. S2E; in all four cases we saw the same trends with IFN-γ. These results are
consistent with the hypothesized role of mycobacteria being a key ingredient in CFA for eliciting
CMI, and that mycobacterial PGN modification by NamH makes a more potent adjuvant.
118
119 <u>Host *Mincle* and *Nod2* are necessary for complete mycobacterial adjuvancy, but do not mediated</u>
120 <u>the entire CFA effect.</u>
121
Recognition of mycobacteria during live infection requires the host molecule NOD2 [24].
Because we hypothesize mycobacterial PGN plays a role in CFA adjuvancy, specifically the
MDP motif where *N*-glycolylation occurs, we addressed whether the PGN/MDP sensor NOD2 is
important for CFA adjuvancy in our OVA model. *Nod2*+/+ and *Nod2*-/- mice were immunized
against OVA in the context of CFA or IFA of commercial provenance. A greater proportion of
OVA-specific IFN-γ-producing CD4+ cells was generated in CFA-immunized mice with *Nod2*
than without (fig. 2A). An independently produced IFN-γ ELISpot of total lymph node cells
supported the same conclusion (fig. S3A), as did another independent experiment using flow
cytometry (fig. S3C). In the absence of *Nod2*, the proportion of OVA-specific IL-17A-
producing cells was also impaired in CFA-immunized mice (fig. 2B and fig. S3C). With IFA as
adjuvant, IFN-γ and IL-17A responses were *Nod2*-independent as expected (fig. 2A-B and S3A-
C). Similar trends were observed with total cell numbers (fig. S3B-C and S4A-B). Together,
these results corroborate the role for NOD2 and accordingly mycobacterial PGN in the ability of
CFA to elicit CMI.
136

The adjuvant effect of mycobacteria declined partially with *Nod2* knock-out, but not
completely. Therefore, there is clearly other mycobacterial MAMP recognition besides that
through NOD2 which is important for the remainder CFA adjuvancy. Others have indicated that
Mincle-mediated recognition of TDM contributes to mycobacterial adjuvancy [17]. We tested the
dependence of our OVA-immunization model with *Mincle*-KO mice [25]. *Mincle*+/+ and *Mincle*-/-
mice were immunized against OVA in the presence of CFA and IFA. Generation of CFA-
elicited OVA-specific IFN-γ- and IL-17A-producing CD4+ T cells was partially dependent on
Mincle (fig. 2C-D), while IFA-immunized mice showed *Mincle*-independent responses, as
expected (fig. 2C-D). Similar trends were again observed with total cell numbers but did not
reach statistical significance (fig. S4C-D). Un-pooled data showed the same trends in both
experiments performed (fig. S3D). These results implicate Mincle and thus its main
mycobacterial ligand, TDM, in CFA adjuvancy.
149
Both *Nod2* and *Mincle* were required for CFA adjuvancy in our model, but neither were
responsible for the entire adjuvant effect. We asked what the sum of the immune responses
elicited by these PRRs is – if it appears these PRRs signal synergistically. We generated *Mincle-
Nod2* double knock-out mice (DKO) and immunized them against OVA with CFA and IFA. The
proportion of OVA-specific IFN-γ- and IL-17A-producing CD4+ T cells in CFA-immunized
mice declined with DKO relative to WT mice, with no change in IFA-immunized mice (fig. 2E-
F). However, the decrease when measuring IFN-γ did not appear obviously larger with DKO
than what was observed with single-KO mice. Likewise, the decrease with IL-17A was
intermediate, and similar trends were seen with total cell numbers (fig. S4E-F). The residual
effect of CFA when both *Nod2* and *Mincle* are disrupted suggests that at least two mycobacterial
MAMPs, *N*-glycolylated PGN as well as TDM, are essential, but insufficient, for the full
adjuvant effect.
162
163   <u>Synthetic minimal structures of mycobacterial PGN and TDM work synergistically to stimulate</u>
164   <u>dendritic cell effector functions.</u>
165
To explore the effect of Mincle and NOD2 signalling on antigen presenting cells
(necessary to generate antigen-specific T-cell immunity), we stimulated bone-marrow derived
dendritic cells (BMDCs) from WT, *Mincle*-/-, *Nod2*-/- and DKO mice with synthetic versions of
the respective PRR ligands. Recently, a rationally determined minimal chemical structure of
TDM called GlcC14C18 was shown to retain activity as a Mincle agonist without the 'toxicity'
associated with purified TDM [23]. Both synthetic *N*-glycolyl and *N*-acetyl MDP are commercially
available. To elicit T-cell immunity, we would expect that mycobacterial MAMPs promote
antigen presentation, co-stimulation and cytokine production by antigen presenting cells, the
classical signals 1, 2 and 3, respectively.
175
After 24 hours of MAMP stimulation *in vitro*, we measured the proportion of cells in
BMDC culture highly expressing MHC-II (fig. 3A-B). With WT cells, MDPs alone had little
effect; GlcC14C18 alone increased the fraction of MHC-II^{hi} cells in WT BMDC cultures (fig.
3A). Combinations of GlcC14C18 with *N*-glycolyl MDP or *N*-acetyl MDP produced the largest
responses. Knock-out of *Mincle* greatly reduced the effect of GlcC14C18 in *Mincle*-/- and DKO
BMDCs, and knock-out of *Nod2* abrogated the effect of MDPs in *Nod2*-/- and DKO BMDCs.
These results suggest these mycobacterial MAMPs can direct myeloid cells towards MHC-II^{hi}

expression as expected for professional antigen presenting cells. Similar results were obtained at
48 hours of stimulation (fig. S5A).

On MHC-II^{hi} cells, we looked at the level of expression of MHC-II, CD40, CD80 and
CD86 by median fluorescence intensity (MFI) after 24 hours of stimulation *in vitro* (fig. S5F-I).
With WT cells and for all markers examined, MDPs alone were unable to substantially enhance
expression over the control (fig. 3C-F). GlcC1418 alone increased marker expression, but
GlcC14C18 together with *N*-glycolyl MDP synergistically produced the greatest responses. *N*-
acetyl MDP was generally less efficacious than *N*-glycolyl MDP. Knockouts behaved as
expected: cells without *Mincle* responded poorly to GlcC14C18; cells without *Nod2* failed to
respond to MDPs (fig. 3C-F). I addition to 24-hour stimulation, cells were also collected after 48
hours of stimulation, yielding similar results (fig. S5B-I).

BMDCs from WT mice produced TNF in a dose-dependent manner in response to
GlcC14C18, but not to MDPs alone (fig. 3G). When combined with limiting or saturating doses
of GlcC14C18, *N*-glycolyl MDP synergistically elevated TNF production in a dose-dependent
manner. *N*-acetyl MDP only significantly elevated TNF production at higher doses of
GlcC14C18 plus MDP and worked less efficaciously than *N*-glycolyl MDP. In the absence of
functional *Mincle*, BMDCs did not respond to GlcC14C18 as expected. *Mincle-/-* BMDCs
stimulated with GlcC14C18 and MDPs behaved like BMDCs exposed to MDP alone (little to no
detectable TNF production). *Nod2-/-* BMDCs responded to GlcC14C18 but not to MDPs as
expected. DKO BMDCs did not respond to either synthetic MAMP as expected. Overall,
GlcC14C18 and *N*-glycolyl MDP combined, through *Mincle* and *Nod2* respectively, are
sufficient to upregulate BMDC antigen presentation, co-stimulation and cytokine production.

Synthetic mycobacterial NOD2 and Mincle ligands can complement IFA to increase antigen-
specific T-cell responses.

As NOD2 and Mincle ligands worked synergistically in BMDCs *in vitro* to elicit effector
functions in these antigen presenting cells, we hypothesized that IFA can be complemented with
NOD2 and/or Mincle ligands to recapitulate at least a portion of the adjuvant effect of whole
mycobacteria. Using minimal synthetic MAMPs for PGN (i.e. *N*-glycolyl MDP) and TDM (i.e.
GlcC14C18), we complemented IFA to test if CMI could be achieved with a completely
synthetic adjuvant containing these essential mycobacterial MAMPs. First, using 10 µg
GlcC14C18 per mouse with varying doses of *N*-glycolyl MDP, we observed a dose dependent
increase in the proportion and numbers of OVA-specific cytokine-producing CD4+ cells which
was significantly greater than IFA alone at the highest dose (fig. 4A). Notably, previous
attempts complementing IFA with only *N*-glycolyl MDP did not induce significant numbers of
IFN-γ producing cells (fig. S6A-B).

Next, using 30 µg of *N*-glycolyl MDP per mouse and varying the dose of GlcC14C18, we
similarly observed a dose dependent increase in the immune response; however this response did
not surpass more than half of the level elicited with CFA by IFN-γ and IL-17A (fig. 4B). An
IFN-γ ELISpot performed in parallel provided a similar conclusion (fig. S6C-D). Together,
these results show that mycobacterial adjuvancy in the context of CFA can be partially

phenocopied synthetically with two mycobacterial MAMPs: namely *N*-glycolyl MDP and
GlcC14C18.
230
231 <u>Synthetic mycobacterial MAMPs increase DC numbers and effectors in lymph nodes.</u>
232
GlcC14C18 and *N*-glycolyl MDP together were sufficient to elicit antigen-specific T-cell
responses. We next looked to see if this MAMP combination was associated with augmentation
of DC effector functions *in vivo* as we had seen *in vitro* with BMDCs. Gating for DC subsets as
well as other leukocytes in the lymph nodes is depicted in fig. S1B (guided by [26]). We looked in
the lymph nodes at 4- and 7-days post-immunization. Both CD11b+ and CD11b- conventional
DC (cDC) numbers were increased in the presence of IFA + 10 μg GlcC14C18 + 30 μg *N*-
glycolyl MDP compared to IFA alone within 7 days, to levels similar to CFA (fig 5A-B). Trends
suggest that early accumulation of cDCs could be driven by MDP alone, but GlcC14C18
together with *N*-glycolyl MDP were required to sustain numbers to day 7.
242
By MFI, expression levels of MHC-I and -II on both CD11b+ and CD11b- cDCs were
not largely altered by MDP nor GlcC14C18 (fig. 5A-B). On CD11b+ cDCs, CD40 expression
was elevated the highest and significantly with IFA + GlcC14C18 MDP at both 4 and 7 days.
CD80 and CD86 expression were also highest with IFA + GlcC14C18 MDP, with statistical
significance at day 4 but not day 7 when expression was possibly waning. On CD11b- cDCs,
CD80 and CD86 expression behaved similarly to that on CD11b+ cDCs. CD40 expression of
CD11b- cDCs was not statistically significantly greater with IFA + GlcC14C18 MDP compared
to IFA alone at both timepoints. Notably, IFA + GlcC14C18 MDP produced equal or greater
costimulatory molecule expression compared to CFA in all cases.
252
In addition to increased numbers of cDCs in the lymph nodes of mice immunized with
GlcC14C18 + *N*-glycolyl MDP, we observed that other leukocytes were substantially elevated in
numbers after immunization with this adjuvant (fig. S7A-F). B cells, CD4+ and CD8+ T cells,
plasmacytoid DCs, monocytes and PMNs were all increased by 7 days post-immunization
relative to IFA, to levels similar to or greater than CFA. By day 7, while MDP alone was
insufficient to elevate cell numbers over the IFA background, MDP in combination with
GlcC14C18 enhanced numbers beyond the levels attained with GlcC14C18 alone. This provides
evidence for synergy between these MAMPs *in vivo*.
261
262 <u>Synthetic mycobacterial MAMPs can induce EAE similar to CFA</u>
263
One of the common uses of CFA is in animal models of autoantigen-specific
autoimmunity. To determine if our synthetic formulation can phenocopy the effect of whole
mycobacteria to produce a more complex biological outcome such as autoimmunity (and thereby
provide a further measure of the 'completeness' of the synthetic formula), we tested IFA +
GlcC14C18 + *N*-glycolyl MDP against CFA in ability to induce relapsing-remitting EAE (RR-
EAE). Briefly, mice were randomly immunized against myelin oligodendrocyte glycoprotein
(MOG) synthetic peptide with CFA or IFA + 10 μg GlcC14C18 + 30 μg *N*-glycolyl MDP, and
onset of RR-EAE was determined by clinically scoring ascending paralysis daily in a blinded
manner (fig. 6A). Both CFA and IFA + GlcC14C18 + MDP produced RR-EAE that was
indistinguishable except quantitatively: over the course of the experiment, the average EAE

score was lower with the synthetic adjuvant compared to the CFA control (fig. 6B), with a
cumulative score suggesting about half the disease burden (58% of CFA cumulative score) (fig.
6C). Expectedly, lower EAE scores also corresponded with less weight-loss (fig. S8A). Of note,
there were mice in the synthetic adjuvant group that reached the same scores as in the CFA
group, but fewer (fig. S8B). Therefore, GlcC14C18 and *N*-glycolyl MDP were sufficient to
recapitulate the adjuvant effect of the whole mycobacterial cell in EAE, albeit quantitatively less
at the tested doses of MAMPs. Additionally, we had attempted RR-EAE with IFA + TDM +
MDP previously, which produced a far less compelling phenocopy of CFA (fig. S8D-G). The
IFA + GlcC14C18 + *N*-glycolyl MDP adjuvant has the advantage of being completely synthetic
and more efficacious than the TDM-containing adjuvant.
284
To verify if there were overt qualitative histopathological differences in EAE between
CFA and the synthetic adjuvant, we examined spinal cords from three mice from each adjuvant
group, having EAE scores 2, 3 and 3.5 at harvest (disease profile of these mice in fig. S8C).
Both cellular infiltrates and demyelination of the white matter looked equivalent between
adjuvants upon visual inspection (fig. 7A). Mice with different EAE scores had correspondingly
different areas of diseased tissue of the spinal cord (fig. 7B), but in comparing adjuvants there
was no detectable difference in the area of diseased spinal cord tissue (with statistical power to
detect as low as +/- 15% CFA levels) (fig. 7C). The main determining variable was EAE score,
not adjuvant. Overall, we observed qualitatively similar autoimmune pathology using IFA +
GlcC14C18 + *N*-glycolyl MDP compared to the gold standard CFA.
295
296 **Discussion**
297
The ability of an adjuvant to elicit CMI can depend on ligand-receptor interactions,
specifically MAMP-PRR interactions. It is well appreciated since the hypothesis of Charles
Janeway Jr. that the host decision to mount an adaptive immune response requires genetically
inborn sensors to detect the presence of microbial products, or microbes by association, and that
this is the foundation of classical adjuvants including CFA [27]. These interactions can be thought
of as an "arms race" between host and microbe, originating through antagonistic evolution in the
case of immune evasion [28]. Additionally, we can imagine the case where a microbe might evolve
to increase a specific PRR interaction that favours a specific active immunological environment
necessary for its lifecycle. When mycobacteria interact with their host, either in the form of *M.*
tuberculosis infection, BCG vaccination or immunization using CFA, CMI normally occurs.
The immune response to mycobacteria has been attributed to its unique cell wall, especially the
MDP motif of PGN [6,7,11,24] and TDM [17,19]. We have shown in the context of both CFA-induced
immunization and autoimmunity with wholly synthetic *N*-glycolyl MDP and TDM (GlcC14C18)
that these MAMP-PRR pathways together contribute partially but significantly to the
mycobacterial adjuvant effect.
313
There are other mycobacterial MAMPs previously identified (and likely more yet
unidentified), plus a greater number of PRRs linked to these microbial products. However, most
are not yet producible synthetically. Purified mannose-capped lipoarabinomannan (ManLAM)
was recently shown to elicit EAE in mice, dependent on the host C-type lectin Dectin-2
(Clec4n), at a dose of 500 µg per animal [29]. Of note, we were not able yet to produce synthetic
mimics of ManLAM that induce Dectin-2 signaling [30]. Similarly, purified TDM has been used

to elicit EAE at 500 µg per mouse, mostly dependent on MCL (Clec4d) and partially on Mincle
(Clec4e) [19]. These findings support a role for mycobacterial MAMPs driving the CFA effect, but
require purification of biologically-sourced MAMPs in high doses. When in combination, our
synthetically pure MAMPs have shown clear activity *in vivo* beginning at 1-10 µg. Of note, with
CFA we inject 50 µg of mycobacterial mass into mice.
325
TLR-2 has been documented to interact with multiple mycobacterial lipids [31]. The
literature is inconsistent on whether or not TLR-2 promotes CFA-induced EAE [32-35].
Mycobacterial DNA was recently shown to signal through the cGAS/STING pathway after
phagosomal disruption via ESX-1 [36,37]. In the context of Freund's adjuvant, it is not clear if heat-
killed mycobacteria could have cytosolic access to activate cGAS/STING. The IFA fraction
perhaps could deliver MAMPs across membranes. TLR-9 also senses DNA, and appears to play
a role during *M. tuberculosis* infection in mice [38], and in humans [39]. TLR-9 was also shown to be
necessary for full induction of EAE [32,34]. Although DNA is not unique to mycobacteria like
TDM and *N*-glycolylated PGN, synthetic DNA is available and pursuing the role of DNA in
mycobacterial adjuvancy interests us, perhaps as a third synthetic MAMP to work with MDP and
GlcC14C18.
337
Mycobacteria are not completely unique in possessing MAMPs that elicit CMI. As an
example, the tuberculosis vaccine candidate M72/AS01E utilizes monophosphoryl lipid A
(MPLA), a TLR-4 agonist based on lipopolysaccharide (LPS), to elicit CMI. LPS is made in
Gram-negative bacteria, not mycobacteria. We concede that many microbes likely contain
MAMPs able to elicit CMI; here we have concerned ourselves primarily with the history of
mycobacterial adjuvancy, CMI and Jules Freund. It is conceivable that a combination of
MAMPs from different bacteria could elicit 'unnatural' immunity that may be beneficial to
control certain infectious agents that otherwise evade natural immune responses.
346
We were able to show that *N*-glycolylation of *M. tuberculosis* PGN altered the adjuvancy
of the mycobacterial cell in the context of CFA. This is consistent with our previous work on the
increased potency of mycobacterial PGN in other contexts [7,11]. Interestingly, the benefit of *N*-
glycolylation of PGN seemed primarily skewed toward IFN-γ production rather than IL-17A,
while *Nod2*-KO showed a greater effect with IL-17A. This may indicate that mycobacteria
retain *N*-glycolylation of PGN specifically to shift immune responses in a Th1 bias and should be
further investigated during live infection. *Nod2* was necessary for full CFA adjuvancy by
measures of IFN-γ and IL-17A, supporting the role for mycobacterial PGN. We were somewhat
surprised that MDP emulsified by itself with IFA and antigen was insufficient to recapitulate the
adjuvant effect of CFA. In a 1975 paper, *N*-acetyl MDP appeared sufficient in a guinea pig
model to elicit DTH using OVA as antigen [6]. In 2009 using a very similar model to our current
work, *N*-glycolyl MDP alone was sufficient to increase ELISpot IFN-γ to near CFA levels, while
N-acetyl MDP failed [7]. One explanation is the possibility of contaminating MAMPs in certain
preparations of OVA. OVA is known to often come with a dose of endotoxin which affects the
immunological outcome of experiments [40]. In the current investigations we have used
endotoxin-free/ultra-pure OVA. Much literature of *in vitro* experiments shows MDP requires
another MAMP to produce measurable outputs like cytokine [7,41-43], although in few cases MDP
was shown to elicit cellular responses by itself [41,42]. Responses measured from MDP alone may
be unrepresentative of a cell's maximum capacity.

366
Other groups have supported a role for TDM in mycobacterial adjuvancy [17,19]. TDM
together with purified PGN was shown to synergistically promote IL-17 production from OVA-
specific (OT-II) CD4+ T cells adoptively transferred into congenic mice [17]. This and other work
directed us to test Mincle-dependence of CFA and MDP synergy with the minimal TDM motif
GlcC14C18. *Mincle*-/- mice allowed us to infer that CFA adjuvancy required TDM for maximal
induction of IFN-γ- and IL-17A-producing Th cells. TDM is thought to be the main Mincle
ligand in mycobacteria; however, there are other ligands. Purified from H37Rv, trehalose-6,6'-
monomycolate, glucose monomycolate, diacyl-trehalose and triacyl-trehalose were shown to
stimulate mouse and human Mincle (plus glycerol monomycolate stimulates human Mincle only)
[23]. It is possible that the phenotype of reduced immunity was because the sensing of these other
MAMPs was decreased. Nevertheless, Mincle signalling was essential for full CFA adjuvancy.
378
Interestingly, simultaneous knock-out of both *Mincle* and *Nod2* resulted in incomplete
abrogation of CFA adjuvancy. DKO mice more closely resembled single-KO from separate
experiments, compared to controls, than IFA. This indicates two things: 1) that much of the
adjuvancy of mycobacteria is not mediated through Mincle and NOD2, and therefore other
mycobacterial MAMPs like those mentioned above may play a similarly important role as PGN
and TDM; 2) that Mincle and NOD2 signalling might work synergistically, where in the absence
of Mincle, NOD2 sensing of PGN is blunted as we have seen clearly *in vitro* with GlcC14C18
and *N*-glycolyl MDP, and therefore DKO resembles single knock-out of either receptor.
387
We demonstrated that GlcC14C18 plus *N*-glycolyl MDP synergistically were sufficient to
promote dendritic cell effector functions *in vitro*. The GlcC14C18 plus *N*-glycolyl MDP
adjuvant was also the most efficacious adjuvant to drive costimulatory molecule upregulation on
cDCs *in vivo*, more than either MAMP alone. While either of these MAMPs by themselves
failed to come close to the whole mycobacteria of CFA in our OVA model of CMI, together we
showed they could produce about half the Th cell IFN-γ/IL-17A response and EAE disease
burden as CFA. Resultingly, we are interested in further understanding the implications for
MAMP signalling (which MAMPs synergize together; what redundancy exists; do different
combinations alter the Th type of CMI?). As the DKO data suggest, other mycobacterial
MAMPs may be necessary to fully recapitulate the adjuvant effect of whole mycobacteria. An
optimal dose of the two MAMPs may not have been reached, although additional dosing
experiments (not shown) failed to find a clearly superior formulation than those presented here.
400
We chose to proceed primarily with the 10 µg GlcC14C18 + 30 µg *N*-glycolyl MDP
doses because these quantities were sufficient for some biological activity on their own and/or in
combination (fig. 4A-B, S6D and not shown). Detailed subset data also showed other biological
synergy at the 10 µg / 30 µg dose (fig. S7). Mycobacteria are an estimated 30% PGN by mass,
of which the muramyl dipeptide motif comprises about half, or 15% total mycobacterial mass [44].
Therefore, when injecting 50 ug of mycobacteria as CFA, we inject about 8 µg of MDP. The 10
µg / 30 µg dose did not quantitatively phenocopy the level of CMI from CFA in the OVA model,
but we did not want to achieve this by using grossly unrepresentative quantities of MAMPs.
Considering the DKO data showing *Mincle* and *Nod2* stimulation do not drive all of the CFA
effect, we proceeded with tempered expectations for EAE. Nonetheless, we have demonstrated
that synthetic TDM and *N*-glycolyl MDP signalling, unique to mycobacteria, enhance the

adjuvancy of CFA. This partly corroborates the natural history of CFA in why it is a uniquely
potent adjuvant. Remaining MAMPs, not necessarily unique to mycobacteria, might fill the
difference.
415
With the wholly synthetic formulation of GlcC14C18 *N*-glycolyl MDP, we were able to
induce RR-EAE in mice that was qualitatively indistinguishable from that induced with CFA, but
milder overall (lower EAE scores on average). We selected spinal cords from mice having
matched EAE scores between CFA and synthetic adjuvant groups; however, extrapolating the
histopathological data to all the mice in our experiment, we would expect that the synthetic
adjuvant produced less pathology in the spinal cord on average since EAE scores were lower on
average. The EAE result correlated with the incomplete complementation in the OVA model.
EAE that failed with TDM plus MDP earlier was also less potent in the OVA model than the
synthetic formulation (not shown), further indicating correlation between these readouts. CFA
often needs to be 'titrated' to account for batch-dependent efficacy in our experience (not
shown), and CFA with higher concentrations of mycobacteria (i.e. 4 mg/ml) is also used when a
more severe, chronic progressive EAE is necessary for studies [45]. These facts point to dosing as
a means of controlling the degree of EAE, which may work with the synthetic adjuvant. Other
experimental uses of CFA include the collagen-induced arthritis (CIA) model and for producing
specific antibodies. CIA takes longer and has a greater role for humoral immunity than EAE
which is T-cell driven, (our interest was in CMI for this study) [46]. Wherever mycobacterial
adjuvancy is useful, we suspect a synthetic adjuvant could be beneficial. Indeed, intravesical
injection of BCG is used to treat bladder cancer (i.e. BCG can provide MAMP-driven protection,
not simply antigen-specific protection). Furthermore, BCG administration to millions of babies
each year can protect not just against childhood tuberculosis, but non-specifically against other
diseases [47].
437
Our work demonstrates that mycobacterial NOD2 and Mincle ligands contribute to the
adjuvant effect of the mycobacterial cell, and that a necessarily combinatorial formulation of
synthetic MAMPs, *N*-glycolyl MDP and GlcC14C18 can partly recapitulate the adjuvant effect
of whole mycobacteria to induce T-cell immunity and EAE. In addition to demonstrating the
first entirely synthetic multi-MAMP mycobacterial adjuvant, we have outlined an approach to
investigate the contribution of other MAMPs to adjuvant design. Moreover, the synthetic
approach may be useful to probe mycobacterial immunity and immunopathogenesis.
445
446 **Methods**
447
Mice
C57BL/6J ('wild-type') as well as *Nod2*-/- mice were obtained from Jackson laboratories and
were bred or used immediately for experiments. *Mincle*-/- mice breeders were provided courtesy
of the laboratory of Christine Wells [25]. Mice were genotyped to confirm absence of the *Dock2*
mutation recently reported in *Nod2*-/- mice [48]. We crossed *Nod2*-/- and *Mincle*-/- mice to obtain
double knock-out mice. All experiments used mice from 6 to 20 weeks of age. All protocols
involving mice followed the guidelines of the Canadian Council on Animal Care (CCAC) and
were approved by the ethics committees of the RI-MUHC.
456
Preparation of adjuvants and ovalbumin immunization

Complete and incomplete Freund's adjuvant were purchased from Sigma or Invivogen. In some
cases, IFA was made by the experimenter from purified mineral oil and mannide monooleate
(Sigma). Adjuvants were prepared on the day of immunization by emulsifying CFA (1 mg/ml
M. tuberculosis) or IFA with a PBS solution containing 1 mg/ml ovalbumin (Endofit brand,
invivogen) in a 1:1 ratio. Emulsification was accomplished using all-plastic syringes and
repeated passage through an 18-G blunt-end needle. Where IFA was complemented with
mycobacterial components: MDP (Invivogen) was diluted in the PBS fraction of the adjuvant
before emulsification; TDM (Sigma) or GlcC14C18 [23] were dissolved in the IFA fraction before
emulsification as described previously [17]; heat-killed mycobacteria in saline were diluted in PBS
and added to the PBS fraction of the adjuvant before emulsification for mice to receive an
equivalent of 10^8 CFU. To make heat-killed mycobacteria, cultures were grown to equivalent
600-nm absorbance at mid-log phase, were pelleted and washed three times with saline, then
heat-killed at 100^oC for 30 minutes and frozen at -80^oC until use. Mice were injected
subcutaneously with 100 µl adjuvant-antigen emulsions in the tail 1-2 cm from the body, towards
the body, for sufficient and consistent drainage to the inguinal lymph nodes. Four or seven days
after injection mice were euthanized and organs were harvested for analysis.

ELISpot and flow cytometry of lymph node cells
Lymph nodes extracted from immunized mice were gently crushed over 70-µm cell strainers to
release the cells therein. Cell concentrations were determined by counting with a
haemocytometer or with a BD Accuri flow cytometer. Lymph node cells (LNCs) were washed
in culture medium, counted and cultured in 100 µl RPMI + 10%FBS (R10) at 250,000 or
500,000 cells per well in IFNγ ELISpot plates (R&D Systems) with or without 100 µg OVA for
~40 hours at 37^oC 5% CO_2 before developing the ELISpot plates. For flow cytometry analysis
of OVA-specific cytokine production, washed LNCs were cultured at 6 million cells per ml in
200 µl R10 with or without 200 µg OVA for ~40 hours at 37^oC 5% CO_2, Brefeldin A
(GolgiPlug[TM], BD) was added for an additional 5 hours, and then cells were stained and fixed
(BD fixation/permeabilization buffer) for flow cytometry on ice. Intracellular staining was
performed on the same or the next day using BD permeabilization buffer. See Table S1 for
antibody probes used. For flow cytometry of leukocyte subsets, 3 million LNCs were taken
immediately after harvest for extracellular staining and fixing. See Table S2 for antibody probes
used. A BD Fortessa X20 was used for cellular phenotyping.

Bone marrow-derived dendritic cells (BMDCs)
Bone marrow was extracted from mice by flushing femora and tibiae with PBS + 2% BSA + 2%
glucose using a 25-G syringe. Red blood cells were lysed and remaining bone marrow cells
were filtered through 70 µm cell strainer. Cells were cultured at 500,000 cells / ml R10 with 20
ng/ml murine rGM-CSF (PeproTech); cells were fed on day 3 with R10 + rGM-CSF, and on day
6 with R10 alone. On day 7, loosely adherent cells were harvested by gentle pipetting and used
in assays. These cells were ~60% CD11c+ MHC-II+ as we obtain routinely. To assess cytokine
production, 100,000 BMDCs were transferred per well to 96-well plates precoated with
GlcC14C18 (by dissolving in isopropanol and drying) and containing MDP in a final volume of
200 µl. BMDCs were incubated with stimulants for 6 hours at 37^oC, 5% CO_2 before supernatant
was removed and stored at -80^oC for later ELISA analysis (mouse TNF ELISA from
ThermoScientific). To asses surface marker expression, 300,000 BMDCs were transferred per
FACS tube precoated with 50 ng GlcC14C18 (by dissolving in isopropanol and drying) and

containing 100 ng/ml MDP in a final volume of 200 µl. BMDCs were incubated with stimulants
for 24 or 48 hours at 37°C, 5% CO_2 before staining and fixing for flow cytometry analysis. See
Table S3 for antibodies used in flow cytometry.
507
Relapsing-remitting experimental autoimmune encephalomyelitis (RR-EAE)
Adjuvants were prepared similarly as above, but with myelin oligodendrocyte glycoprotein
(MOG) used as antigen. Briefly, PBS solution containing 1 mg/ml MOG +/- 600 µg/ml *N*-
glycolyl MDP was emulsified with CFA (1 mg/ml *M. tuberculosis*) / IFA plus 200 µg/ml
GlcC14C18 (or 20 µg/ml TDM) by back-and-forth extrusion through an 18 G two-way needle
with two all-plastic syringes. 8-12 week old female C57BL/6 mice (Charles River) were induced
for RR-EAE by a standard protocol: briefly, on day 0, pertussis toxin (PT) (200 ng) was
administered i.v. in the tail vein, and mice were immunized with 50 µg MOG by bilateral s.c.
injection in the back towards the tail with 100 µl total of an emulsification of CFA or IFA plus
mycobacterial MAMPs (adjuvant group was randomized within and across cages). On day 2,
mice received a second equivalent dose of PT i.v.. After about one week, EAE was scored
blinded (i.e. the scorer did not know the adjuvant received by the mouse) every day according to
these cumulative criteria: 0, no paralysis (normal); 0.5, partial tail weakness observed as < 50%
of tail dragging when mouse walks; 1, tail paralysis observed as >50% of tail dragging as mouse
walks; 2, slow righting reflex (delay < 5 seconds) when mouse is flipped; 3, very slow or absent
righting reflex (> 5 seconds) or inability to bear weight with back legs observed as dragging
hindquarters when walking; 3.5, partial paralysis of one or both hind limbs; 4, complete paralysis
of one or both hindlimbs; 4.5, complete paralysis of one or both hind limbs plus trunk weakness;
5, weakness or paralysis of forelimbs; 6, found dead. Mice were weighed every other day during
scoring. Mice reaching a score of 5 were euthanized within 24 hours. Blinding was
accomplished by having one person inject the mice and another person score/weigh the mice,
unaware of experimental group.
530
Histopathology
At the end of EAE scoring, the experiment was un-blinded and three mice from each group with
matching scores were selected. These six mice were anesthetized with peritoneal ketamine
injection, were perfused with PBS and formalin, and then spinal cords were extracted,
equilibrated in sucrose and then frozen in OTC compound (VWR). Frozen tissue was sectioned
14-nm thick onto slides and then stained (Nissl or Luxol fast blue). Sections of good quality (20
per mouse, 5 cervical, 5 upper thoracic, 5 lower thoracic and 5 lumbar) were randomly selected
and photographed with a Nikon Eclipse microscope. Randomized (blinded) Nissl stain
photographs were used to quantify the area of disease in the spinal cord by subtracting the total
2D area of each tissue section with the area without cellular infiltration in white matter (thus, the
difference of areas is the area with cellular infiltration). This was divided by the total area to
determine the fraction or percent of diseased area.
543
Software, data analysis and statistics
Flow cytometry data was acquired using FACSDiva[TM] software (BD). FCS files were analyzed
using FlowJo V10 (BD). Digital microscopy images were analyzed with ImageJ (NIH). Graphs
were generated with, and routine statistical testing was accomplished with GraphPad Prism v7 or
v8 (GraphPad Software Inc.). Assessment for normal data, then parametric or non-parametric
analyses were applied, as indicated. Sample-size and power calculations were performed

manually using Microsoft Excel. Manuscript figures were assembled with Microsoft
PowerPoint.
552

553 **Acknowledgments**
554
We would like to thank the Containment Level 3 platform of the Research Institute of the
McGill University Health Centre (RI-MUHC) for providing the needed facilities and intellectual
support. We thank the Immunophenotyping platform of the RI-MUHC for cytometers and
advice. We are very grateful to Christine Wells (University of Melbourne, Australia) for
providing *Mincle*-KO mice and to Alexiane Decout for preparing GlcC14C18. For tutelage and
technical assistance, J.Y.D. is thankful to Damien Montamat-Sicotte, Ourania Tsatas, Sarah
Danchuk, Iain Roe, Andréanne Lupien, Nimara Asbah, and Daniel Houle. J.Y.D. was personally
supported by the Canadian Institutes of Health Research (CIHR) Canada Graduate Scholarship –
Master's Program, Fonds de Recherche du Québec – Santé (FRQ-S) Doctoral Training Award,
RI-MUHC studentships (Master's and Ph.D.) and scholarships from the McGill Department of
Microbiology and Immunology. Research activities were funded by CIHR foundation grant held
by M.B. and a CIHR grant held by S.D.

567
568 **Additional Information**
569
570 Competing interests: the authors have none.
571
572 **References**
573

1   Chase, M. W. & Landsteiner, K. Experiments on Transfer of Cutaneous Sensitivity to
Simple Compounds. *Proc. Soc. Exp. Biol. Med.* **49**, 688-690 (1942).
2   Chase, M. W. The Cellular Transfer of Cutaneous Hypersensitivity to Tuberculin. *Proc.*
Soc. Exp. Biol. Med. **59**, 134-135 (1945).
3   Baxter, A. G. The origin and application of experimental autoimmune encephalomyelitis.
Nat. Rev. Immunol. **7**, 904-912 (2007).
4   Constantinescu, C. S., Farooqi, N., O'Brien, K. & Gran, B. Experimental autoimmune
encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *Br. J. Pharmacol.* **164**,
1079-1106, doi:10.1111/bph.2011.164.issue-4 (2011).
5   Vogel, F. R. Improving Vaccine Performance with Adjuvants. *Clin. Infect. Dis.* **30**,
S266–270 (2000).
6   Adam, A., Ellouz, F., Ciorbaru, R., Petit, J. F. & Lederer, E. Peptidoglycan adjuvants:
minimal structure required for activity. *Z. Immunitatsforsch. Exp. Klin. Immunol.* **149**,
341-348 (1975).
7   Coulombe, F. *et al.* Increased NOD2-mediated recognition of *N*-glycolyl muramyl
dipeptide. *J. Exp. Med.* **206**, 1709-1716, doi:10.1084/jem.20081779 (2009).
8   Raymond, J. B., Mahapatra, S., Crick, D. C. & Pavelka, M. S., Jr. Identification of the
namH gene, encoding the hydroxylase responsible for the N-glycolylation of the
mycobacterial peptidoglycan. *J. Biol. Chem.* **280**, 326-333, doi:10.1074/jbc.M411006200
(2005).

9   Mahapatra, S., Crick, D. C., McNeil, M. R. & Brennan, P. J. Unique structural features of
595   the peptidoglycan of Mycobacterium leprae. *J. Bacteriol.* **190**, 655-661,
596   doi:10.1128/JB.00982-07 (2008).

597   10   Essers, L. & Schoop, H. J. Evidence for the incorporation of molecular oxygen, a
598   pathway in biosynthesis of *N*-glycolylmuramic acid in *Mycobacterium phlei*. *Biochim.*
599   *Biophys. Acta* **544**, 180-184 (1978).

600   11   Hansen, J. M. *et al.* *N*-glycolylated peptidoglycan contributes to the immunogenicity but
601   not pathogenicity of Mycobacterium tuberculosis. *J. Infect. Dis.* **209**, 1045-1054,
602   doi:10.1093/infdis/jit622 (2014).

603   12   Girardin, S. E. *et al.* Nod2 is a general sensor of peptidoglycan through muramyl
604   dipeptide (MDP) detection. *J. Biol. Chem.* **278**, 8869-8872, doi:10.1074/jbc.C200651200
605   (2003).

606   13   Miceli-Richard, C. *et al.* CARD15 mutations in Blau syndrome. *Nat. Genet.* **29**, 19-20,
607   doi:10.1038/ng720 (2001).

608   14   Ogura, Y. *et al.* A frameshiftmutation in NOD2 associated with susceptibility to Crohn's
609   disease. *Nat. Lett.* **411**, 603-606 (2001).

610   15   Austin, C. M., Ma, X. & Graviss, E. A. Common nonsynonymous polymorphisms in the
611   NOD2 gene are associated with resistance or susceptibility to tuberculosis disease in
612   African Americans. *J. Infect. Dis.* **197**, 1713-1716, doi:10.1086/588384 (2008).

613   16   Zhang, F. R. *et al.* Genomewide association study of leprosy. *N. Engl. J. Med.* **361**, 2609-
614   2618, doi:10.1056/NEJMoa0903753 (2009).

615   17   Shenderov, K. *et al.* Cord factor and peptidoglycan recapitulate the Th17-promoting
616   adjuvant activity of mycobacteria through Mincle/CARD9 signaling and the
617   inflammasome. *J. Immunol.* **190**, 5722-5730, doi:10.4049/jimmunol.1203343 (2013).

618   18   Ishikawa, E. *et al.* Direct recognition of the mycobacterial glycolipid, trehalose
619   dimycolate, by C-type lectin Mincle. *J. Exp. Med.* **206**, 2879-2888,
620   doi:10.1084/jem.20091750 (2009).

621   19   Miyake, Y. *et al.* C-type lectin MCL is an FcRgamma-coupled receptor that mediates the
622   adjuvanticity of mycobacterial cord factor. *Immunity* **38**, 1050-1062,
623   doi:10.1016/j.immuni.2013.03.010 (2013).

624   20   Bekierkunst, A. Acute Granulomatous Response Produced in Mice by Trehalose-6,6-
625   Dimycolate. *J. Bacteriol.* **96**, 958-961 (1968).

626   21   Shigeru, K. *et al.* Synthesis of Muramyl Dipeptide Analogs with Enhanced Adjuvant
627   Activity. *Bull. Chem. Soc. Japan* **53**, 2570-2577 (1980).

628   22   Xing, S. & Gleason, J. L. A robust synthesis of N-glycolyl muramyl dipeptide via
629   azidonitration/reduction. *Org. Biomol. Chem.* **13**, 1515-1520, doi:10.1039/c4ob02147a
630   (2015).

631   23   Decout, A. *et al.* Rational design of adjuvants targeting the C-type lectin Mincle. *Proc.*
632   *Natl. Acad. Sci. U. S. A.* **114**, 2675-2680, doi:10.1073/pnas.1612421114 (2017).

633   24   Divangahi, M. *et al.* NOD2-Deficient Mice Have Impaired Resistance to Mycobacterium
634   tuberculosis Infection through Defective Innate and Adaptive Immunity. *J. Immunol.* **181**,
635   7157-7165, doi:10.4049/jimmunol.181.10.7157 (2008).

636   25   Wells, C. A. *et al.* The Macrophage-Inducible C-Type Lectin, Mincle, Is an Essential
637   Component of the Innate Immune Response to Candida albicans. *The Journal of*
638   *Immunology* **180**, 7404-7413, doi:10.4049/jimmunol.180.11.7404 (2008).

26   Merad, M., Sathe, P., Helft, J., Miller, J. & Mortha, A. The dendritic cell lineage:
ontogeny and function of dendritic cells and their subsets in the steady state and the
inflamed setting. *Annu. Rev. Immunol.* **31**, 563-604, doi:10.1146/annurev-immunol-
020711-074950 (2013).
27   Janeway, C. A. Approaching the Asymptote? Evolution and Revolution in Immunology.
Cold Spring Harb. Symp. Quant. Biol. **54**, 1-13 (1989).
28   Odendall, C. & Kagan, J. C. Host-Encoded Sensors of Bacteria: Our Windows into the
Microbial World. *MicrobiolSpectrum* **7** (2019).
29   Yonekawa, A. *et al.* Dectin-2 Is a Direct Receptor for Mannose-Capped
Lipoarabinomannan of Mycobacteria. *Immunity* **41**, 402–413 (2014).
30   Decout, A. *et al.* Deciphering the molecular basis of mycobacteria and lipoglycan
recognition by the C-type lectin Dectin-2. *Sci. Rep.* **8**, 16840, doi:10.1038/s41598-018-
35393-5 (2018).
31   Stamm, C. E., Collins, A. C. & Shiloh, M. U. Sensing of Mycobacterium tuberculosis and
consequences to both host and bacillus. *Immunol. Rev.* **264**, 204-219,
doi:10.1111/imr.12263 (2015).
32   Prinz, M. *et al.* Innate immunity mediated by TLR9 modulates pathogenicity in an animal
model of multiple sclerosis. *J. Clin. Invest.* **116**, 456-464, doi:10.1172/JCI26078 (2006).
33   Reynolds, J. M. *et al.* Toll-like receptor 2 signaling in CD4(+) T lymphocytes promotes T
helper 17 responses and regulates the pathogenesis of autoimmune disease. *Immunity* **32**,
692-702, doi:10.1016/j.immuni.2010.04.010 (2010).
34   Miranda-Hernandez, S. *et al.* Role for MyD88, TLR2 and TLR9 but not TLR1, TLR4 or
TLR6 in experimental autoimmune encephalomyelitis. *J. Immunol.* **187**, 791-804,
doi:10.4049/jimmunol.1001992 (2011).
35   Shaw, P. J. *et al.* Signaling via the RIP2 adaptor protein in central nervous system-
infiltrating dendritic cells promotes inflammation and autoimmunity. *Immunity* **34**, 75-84,
doi:10.1016/j.immuni.2010.12.015 (2011).
36   Collins, Angela C. *et al.* Cyclic GMP-AMP Synthase Is an Innate Immune DNA Sensor
for Mycobacterium tuberculosis. *Cell Host & Microbe* **17**, 820-828,
doi:10.1016/j.chom.2015.05.005 (2015).
37   Watson, Robert O. *et al.* The Cytosolic Sensor cGAS Detects Mycobacterium
tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. *Cell Host &*
Microbe **17**, 811-819, doi:10.1016/j.chom.2015.05.004 (2015).
38   Bafica, A. *et al.* TLR9 regulates Th1 responses and cooperates with TLR2 in mediating
optimal resistance to Mycobacterium tuberculosis. *J. Exp. Med.* **202**, 1715-1724,
doi:10.1084/jem.20051782 (2005).
39   Chen, Z. *et al.* Association between toll-like receptors 9 (TLR9) gene polymorphism and
risk of pulmonary tuberculosis: meta-analysis. *BMC Pulm. Med.* **15**, 57,
doi:10.1186/s12890-015-0049-4 (2015).
40   Eisenbarth, S. C. *et al.* Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T
helper cell type 2 responses to inhaled antigen. *J. Exp. Med.* **196**, 1645-1651,
doi:10.1084/jem.20021340 (2002).
41   Kobayashi, K. S. *et al.* Nod2-Dependent Regulation of Innate and Adaptive Immunity in
the Intestinal Tract. *Science* **307**, 731-734 (2004).

42   Khan, N. *et al.* Signaling through NOD-2 and TLR-4 Bolsters the T cell Priming
Capability of Dendritic cells by Inducing Autophagy. *Sci. Rep.* **6**, 19084,
doi:10.1038/srep19084 (2016).
43   Wolfert, M. A., Murray, T. F., Boons, G. J. & Moore, J. N. The origin of the synergistic
effect of muramyl dipeptide with endotoxin and peptidoglycan. *J. Biol. Chem.* **277**,
39179-39186, doi:10.1074/jbc.M204885200 (2002).
44   Allison, D. G. & Lambert, P. A. in *Molecular Medical Microbiology*   Ch. 32, 583-598
(2015).
45   Berard, J. L., Wolak, K., Fournier, S. & David, S. Characterization of relapsing-remitting
and chronic forms of experimental autoimmune encephalomyelitis in C57BL/6 mice.
Glia **58**, 434-445, doi:10.1002/glia.20935 (2010).
46   Brand, D. D., Latham, K. A. & Rosloniec, E. F. Collagen-induced arthritis. *Nat. Protoc.*
2, 1269-1275, doi:10.1038/nprot.2007.173 (2007).
47   Kleinnijenhuis, J. *et al.* Bacille Calmette-Guerin induces NOD2-dependent nonspecific
protection from reinfection via epigenetic reprogramming of monocytes. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 17537-17542, doi:10.1073/pnas.1202870109 (2012).
48   Wong, S. Y. *et al.* B Cell Defects Observed in Nod2 Knockout Mice Are a Consequence
of a Dock2 Mutation Frequently Found in Inbred Strains. *J. Immunol.* **201**, 1442-1451,
doi:10.4049/jimmunol.1800014 (2018).

702
703
704 **Author Contributions**
705
706 Conceptualization, J.Y.D. and M.B.; Methodology, J.Y.D. and J.G.Z.; Investigation, J.Y.D.,
707 F.M. and J.G.Z.; Resources, S.D., J.N. and M.B.; Writing – Original Draft, J.Y.D. and M.B.;
708 Writing – Review & Editing, J.Y.D., F.M., J.G.Z., S.D., J.N. and M.B.; Visualization, J.Y.D.;
709 Supervision, S.D. and M.B.; Funding acquisition, J.Y.D., S.D. and M.B.
710
711 **Legends**
712
Figure 1. CFA-dependent cell-mediated immune responses as a function of mycobacterial
***namH*. A**, PGN of wild-type H37Rv *M. tuberculosis* (left) and PGN of the Δ*namH* mutant
(right). The MDP motif is drawn in red, and the site of *N*-glcolylation is in bold font. With
NamH, *N*-glycolylation was shown on ~70% of muramic acid residues, with *N*-acetylation on the
remaining ~30% [8,11]. **B**, immunization scheme (relevant to figs. 1, 2 and 4): mice were
immunized with adjuvant emulsion containing OVA by s.c. injection at the base of the tail, and
after seven days, inguinal (draining) lymph nodes were harvested. Lymph node cells were
cultured *ex vivo* with or without OVA to examine the OVA-specific cytokine response by flow
cytometry or ELISpot. **C-D**, Proportion of cytokine-producing CD4+CD8- lymph node cells of
mice immunized against OVA with heat-killed *M. tuberculosis* strain H37Rv, H37Rv Δ*namH*, or
IFA alone, seven days prior. Shown are data pooled from four separate experiments with
averages +/- SEM. p-values were calculated with two-tailed student's *t*-tests. *p<0.05. For IFA
+ H37Rv, IFA + H37Rv Δ*namH*, and IFA alone, sample size N = 31, 27 and 16 mice,
respectively. Each plotted point represents the result obtained from an individual mouse.
727

Figure 2. CFA-dependent cell-mediated immune responses as a function of host *Nod2* and *Mincle*. Proportion of cytokine-producing CD4+CD8- lymph node cells of mice immunized against OVA with CFA or IFA seven days prior. **A-B**, *Nod2*+/+ vs. *Nod2*-/- mice, data representative of two independent experiments with averages +/- SEM. p-values were calculated with two-tailed student's *t*-tests. *p<0.05; **p<0.01. For CFA *Nod2*+/+, CFA *Nod2*-/-, IFA *Nod2*+/+ and IFA *Nod2*-/-, sample size N = 12, 13, 9 and 7 mice, respectively. **C-D**, *Mincle*+/+ vs. *Mincle*-/- mice, data pooled from two independent experiments with averages +/- SEM. p-values were calculated with two-tailed student's *t*-tests. *p<0.05; **p<0.01. For CFA *Mincle*+/+, CFA *Mincle*-/-, IFA *Mincle*+/+ and IFA *Mincle*-/-, sample size N = 14, 16, 15 and 11 mice, respectively. **E-F**, WT (*Mincle*+/+*Nod2*+/+) vs. DKO (*Mincle*-/-*Nod2*-/-) mice with averages +/- SEM. p-values were calculated with two-tailed student's *t*-tests. *p<0.05; **p<0.01. For CFA WT, CFA DKO, IFA WT and IFA DKO, sample size N = 10, 7, 7 and 7 mice, respectively.

Figure 3. MHC-II expression, costimulatory molecule upregulation and TNF production and by BMDCs stimulated with GlcC14C18 and MDPs. **A**, percentage of cells expressing MHC-II at high levels (according to gate in panel B) after 24 hours of stimulation with the indicated MAMPs. **B**, gating of MHC-II^{hi} cells amongst live single CD11b+CD11c+ cells. **C-F**, median fluorescence intensity of **C**, MHC-II; **D**, CD40; **E**, CD80; **F**, CD86 on CD11b+CD11c+MHC-II^{hi} cells after 24 hours of stimulation with the indicated MAMPs (use legend of panel A). Shown are averages +/- SD of 3 individually stimulated and assayed cultures. To compare the combination of GlcC14C18+*N*-glycolyl MDP to unstimulated control, GlcC14C18 alone and GlcC14C18+*N*-acetyl MDP with WT cells, p-values were calculated using Dunnett's T3 multiple comparisons test; p<0.05; **p<0.01; ***p<0.001; ****p<0.0001; ns, not significant, p>0.05. **G**, Supernatant TNF after 6-hour stimulation with GlcC14C18 and MDPs. Shown are averages +/- SD of 3-6 individually stimulated and assayed culture wells. To determine which MDP dose elicited significantly more TNF over background dose of GlcC14C18, p-values were calculated with Dunnett's multiple comparisons test. The *, # and & symbols were used for the 17, 50 and 150 ng GlcC14C18 doses respectively, where *p<0.05; **p<0.01; ***p<0.001 (for clarity, non-significant p-values are not noted).

Figure 4. Complementation of IFA with synthetic mycobacterial MAMPs. Proportion of cytokine-producing CD4+CD8- lymph node cells of WT mice immunized against OVA seven days prior. **A**, *N*-glycolyl MDP dose-dependent response with 10 µg GlcC14C18: IFA (N=4), IFA + 10 µg GlcC14C18 (N=7), IFA + 10 µg GlcC14C18 + 1 µg *N*-glycolyl MDP (N=6), and IFA + 10 µg GlcC14C18 + 3 µg *N*-glycolyl MDP (N=7). Shown are data from individual mice, with averages +/- SEM. p-values were calculated with Welch's two-tailed student's *t*-tests. *p<0.05; **p<0.01. **B**, GlcC14C18 dose-dependent response with 30 µg *N*-glycolyl MDP: IFA + 30 µg *N*-glycolyl MDP (N=6), IFA + 30 µg *N*-glycolyl MDP + 10 µg GlcC14C18 (N=7), IFA + 30 µg *N*-glycolyl MDP + 30 µg GlcC14C18 (N=7), and CFA (N=7). Shown are data from individual mice, with averages +/- SEM. In comparing IFA + 30 µg *N*-glycolyl MDP to IFA + *N*-glycolyl MDP + 10 or 30 µg GlcC14C18, p-values were calculated using Dunnett's T3 multiple comparisons test. *p<0.05; **p<0.01; ns, not significant, p>0.05.

Figure 5. Dynamics of cDCs in lymph nodes after immunization with synthetic mycobacterial MAMPs. A, CD11b+ cDC quantities and median fluorescence intensity of

MHC-I, MHC-II, CD40, CD80 and CD86 in mice immunized with the indicated adjuvants after
4 or 7 days. **B**, CD11b- cDC quantities and median fluorescence intensity of MHC-I, MHC-II,
CD40, CD80 and CD86 in mice immunized with the indicated adjuvants after 4 or 7 days. For
all adjuvant groups, N=8 mice (each plotted point represents data from an individual mouse).
Shown are averages +/-SEM. To determine if IFA + GlcC14C18 + *N*-glycolyl MDP altered cDC
parameters compared to IFA alone, p-values were calculated using Dunnett's T3 multiple
comparisons test correcting for multiple timepoints and cDC type (CD11b+/-). *p<0.05;
**p<0.01; ns, not significant, p>0.05.

Figure 6. RR-EAE induced by IFA+GlcC14C18+MDP. A, experimental timeline. **B**,
average EAE score +/- SEM over time of mice induced with CFA (N=15) or IFA + 10 µg
GlcC14C18 + 30 µg *N*-glycolyl MDP (N=15). Mice were euthanized on day 28 post injection.
C, Cumulative EAE score, obtained by adding the EAE score of each mouse over each of the 28
days. Lines represent averages +/- SEM. p-value was calculated with two-tailed student's *t*-test.
**p = 0.0022.

Figure 7. Spinal cord pathology in RR-EAE mice induced by IFA+GlcC14C18+MDP. A,
Nissl and Luxol fast blue (LFB) stains of spinal cord sections from RR-EAE-induced mice at day
28 post injection, having an EAE score of 3.5 upon euthanasia. Red boxes highlight cellular
infiltration and spatially associated demyelination of the white matter seen by Nissl and LFB
staining, respectively. **B**, Quantitative spinal cord pathology per EAE score upon euthanasia.
Statistical significance was determined by Tukey's multiple comparisons test. *p<0.05 and
****p<0.0001. N=40 for each group (20 from CFA and 20 from synthetic adjuvant of
equivalent EAE scores). Lines indicate averages +/- SEM   **C**, Quantitative spinal cord
pathology per adjuvant. Statistical significance was tested with two-tailed unpaired Welch's *t*-
test; a power calculation for given variances and N=60 per group indicated an ability to discern
+/- 15% difference vs. CFA control. Lines indicate averages +/- SEM.

FIG. 1

FIG. 2

FIG. 3

FIG. 4

FIG. 5

FIG. 6

A

Pertussis Toxin i.v.

Adjuvant + MOG s.c.

Monitor for 28 days, EAE signs + weight

Harvest Spinal Cord

B

C

FIG. 7

A

B

y-axis: % area with cell infiltrate

x-axis: EAE score upon euthanasia (2, 3, 3.5)

* ; **** ; ****

C

y-axis: % area with cell infiltrate

x-axis: CFA, IFA + GlcC14C18 + *N*-glycolyl MDP

p = 0.86