Model to link cell shape and polarity with organogenesis

Bjarke Frost Nielsen[1], **Silas Boye Nissen**[1], **Kim Sneppen**[1], **Ala Trusina**[1†], **Joachim Mathiesen**[1†]

[1]Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark

*For correspondence:
trusina@nbi.ku.dk (AT);
mathies@nbi.ku.dk (JM)

[†]Corresponding authors

Abstract   Cell shape change (e.g. wedging), intercalation and proliferation drive epithelial sheet-to-tube transitions. How these complement each other remains an open question. Previously we introduced a particle model of cell sheet dynamics (*Nissen et al., 2018*) which captured features of tubulogenesis but lacked an analog of apical constriction. Here, we model wedging by a cell-intrinsic constraint on apical-basal polarity and can thereby simulate tube budding and extension (in e.g. salivary glands) where *cell rearrangement* and *wedging* act in parallel. We find that either of the two can initiate budding, but isotropic wedging is needed to control tube orientation. Furthermore, we recapitulate the reported role of anisotropic wedging and its coupling to planar cell polarity in neurulation. Simulations predict that the proliferation rate of neural progenitors is a key parameter for successful tube closure. Our work connects changes in cell shape and polarities to macroscopic transitions in organogenesis.

Introduction

Early tubes in embryonic development – gut and neural tubes – form out of epithelial sheets. In mammalian embryos and *Drosophila*, the cell sheet wraps around the tube axis until the edges make contact and fuse. As a result of such *wrapping*, a tube is formed parallel to the sheet. In sea urchin, the gut is formed orthogonal to the epithelial plane by *budding* out of the plane. Budding also appears to be a predominant form of tube formation in organ development (lungs, kidneys, salivary gland and trachea in *Drosophila* (*Andrew and Ewald, 2010*).

Both wrapping and budding sheet-to-tube transitions are driven by the same key mechanisms: changes in cell shape, e.g. cell wedging by changes in apical relative to the basal surfaces – apical constriction (AC) (*Sawyer et al., 2010*) or basal constriction (*Gutzman et al., 2018*; *Visetsouk et al., 2018*); contracting myosin cables spanning across cells and Convergent Extension (CE) by directed cell intercalation (*Chung et al., 2017*; *Andrew and Ewald, 2010*). (In the following, we will refer to apical or basal constriction as *wedging* and directed cell intercalation as CE).

Until recently, the consensus has been that wedging and CE each lead to distinct morphological transformations: wedging bends the sheet and is a primary mechanism for invagination in budding *Paluch and Heisenberg* (*2009*) and CE elongates the sheet and the eventual tube (*Andrew and Ewald, 2010*). However, results by *Chung et al.* (*2017*) and *Sanchez-Corrales et al.* (*2018*) show that invagination in *Drosophila* salivary gland can happen in the absence of wedging and may result from radial CE.

Nishimura et al. (*2012*) suggested that in neurulation CE and wedging are coupled. They argue that Planar Cell Polarity (PCP) may be mediating this coupling: First, the direction of cell intercalations, orthogonal to the tube axis, is set by PCP. Second, wedging has to be anisotropic – have preferred direction parallel to PCP and intercalations – for the sheet to wrap into a tube and not a sphere.

The anisotropy of wedging, however, is rarely considered, possibly because the reported results are
often limited to 2D cross-sectional views. A few examples of studies where anisotropic wedging
is reported are *Nishimura et al.* (*2012*); *Sweeton et al.* (*1991*); *Martin et al.* (*2010*). The coupling
between PCP and wedging – apical *and* basal – is also supported by data at the molecular level (for
neural tube closure (*Ossipova et al., 2014*; *Nishimura et al., 2012*), midbrain-hindbrain boundary in
zebrafish (*Visetsouk et al., 2018*; *Gutzman et al., 2018*), gastrulation in *C. elegans* (*Lee et al., 2006*),
sea urchin (*Croce et al., 2006*), and *Xenopus* (*Choi and Sokol, 2009*)).

This recent development opens for new questions: What are wedging and CE capable of on
their own? Can invagination by CE happen in systems other than salivary glands? Is anisotropy in
wedging required for tubulogenesis and, if so, when?

Theoretical models have been an important part for understanding tissue bending by wedging
and elongation by CE. However, the existing models are often limited to 2D and thus focus on either
wedging or CE (*Collinet et al., 2015*; *Belmonte et al., 2016*; *Spahn and Reuter, 2013*). While there
are several 3D models for budding and neurulation (*Kim et al., 2013*; *Inoue et al., 2016*), models
that are able to either couple CE and wedging, or address the role of PCP-driven anisotropy of AC
are lacking. To probe the above questions theoretically, we expand the recently published model of
polarized cell–cell interactions (*Nissen et al., 2018*). By treating cells as point particles, *Nissen et al.*
(*2018*) focused on directional adhesion mediated by apical-basal (AB) polarity and PCP. This allowed
to successfully capture the coupling between PCP and CE, however, the model did not explicitly
account for changes in cell shapes. Here, we show that the effect of cell wedging can be modeled
within a point-particle representation by modifying cell-cell forces to favor a tilt in AB polarities.

In line with data by *Chung et al.* (*2017*), simulations show that although CE alone can lead to a
budding transition, it is less reliable, with frequent failure of invagination and misorientation. Our
results suggest that the isotropic wedging orients the budding and allows for robust invagination.
When applied to wrapping in neurulation, we find that anisotropic wedging alone was insufficient
for final tube closure. However, both closure and tube separation from the epithelium can be
aided by differential proliferation. Furthermore, we find that anisotropic wedging on its own may
be sufficient for tube elongation. Together, our results support the mutual complementarity of
wedging and CE in bending and elongation.

Results

73 To investigate the role of cell wedging in budding of salivary gland placoids and neurulation we
74 aimed at capturing both isotropic and anisotropic (PCP-driven) wedging with as few parameters as
75 possible.

Modelling wedging of a point particle by favoring tilt in AB

Apical constriction leads to cell wedging and as a consequence the AB axes of neighboring cells
become tilted towards the wedged cell (Figure 1B–C). *Nissen et al.* (*2018*) modeled a flat epithelial
sheet by a cell-cell interaction force favouring parallel AB polarities in neighbor cells (Figure 1A,
Equation 4). To model the effect of wedging we modify the force to favor AB polarity vectors $\mathbf{p}_i$ in
neighbor cells to tilt towards the wedged cell (Figure 1B and 1C). That is, when the force is calculated,
we replace $\mathbf{p}_i$ by $\tilde{\mathbf{p}}_i$ (Equations 1-3).

$$\tilde{\mathbf{p}}_i = \mathbf{p}_i \quad \text{(for no wedging)}, \tag{1}$$

$$\tilde{\mathbf{p}}_i \propto \mathbf{p}_i - \alpha\hat{\mathbf{r}}_{ij} \quad \text{(for isotropic wedging)}, \tag{2}$$

$$\tilde{\mathbf{p}}_i \propto \mathbf{p}_i - \alpha\langle\hat{\mathbf{q}}\rangle_{ij} \quad \text{(for anisotropic wedging)}. \tag{3}$$

77 Here, $\hat{\mathbf{r}}_{ij}$ is the normalized displacement vector between cells i and j while $\langle\hat{\mathbf{q}}\rangle_{ij}$ is the average PCP
78 vector of the two interacting particles.

79   This change required only one parameter, α, setting the extent of the tilt (large α corresponds to
80 pronounced wedging). If the wedging is isotropic, i.e. equally pronounced in all directions (*Sanchez-*
81 *Corrales et al., 2018*), all neighbors to the wedged cell tend to tilt equally. In neurulation, the

Figure 1. Wedging is introduced through a cell-cell interaction that favors tilted AB polarity vectors. α is the extent of the wedging. The blue-red gradient indicates the apical-basal axis.
(**A**) No wedging ($\alpha = 0$), AB polarities (arrows) tend to be parallel. (**B**) With isotropic wedging, the tilt α is the same in all directions. (**C**) With anisotropic wedging, the tilt has a preferred direction. Blue and red signify respectively basal and apical surfaces. p_i and p_j are the AB polarities of cell i and j.

wedging is anisotropic: the wedging happens primarily parallel to the cell's PCP and perpendicular to the axis of the tube (*Nishimura et al., 2012*). To capture this PCP-directed anisotropy, we couple the direction of AB tilting to the orientation of the cell's PCP (Equation 3, Figure 1C). See the Methods section for details of the model and simulations.

Note, that we aim to only capture the effect of wedging-PCP coupling and not the molecular mechanism. Also, in an attempt to generalize our results, we focus on a minimal set of conditions necessary for the final outcome.

To test the validity of our approach, we first consider the complementary roles of CE and wedging in budding.

Complementary and unique roles of CE and wedging in budding

Reflecting the viewpoint that tube budding is a two stage process consisting of wedging-driven invagination and subsequent convergent extension, computational models have generally focused on either of the two stages (*Collinet et al., 2015*; *Belmonte et al., 2016*; *Spahn and Reuter, 2013*). However, to date no computational models have managed to recapitulate the results by *Sanchez-Corrales et al.* (*2018*) and *Chung et al.* (*2017*) suggesting that CE contributes to invagination and may even drive it in the absence of wedging.

To validate our approach, we set to reproduce these experimental observations. We start with a flat sheet of AB polarized cells. Motivated by the possible link between organizing signals (e.g. WNT), PCP and wedging (*Habib et al., 2013*; *Loh et al., 2016*), we define a region of "organizing signals" such that the cells within this region exhibit isotropic wedging and PCP. In salivary glands, the apically constricting cells are distributed on a disk around the future center of the tube. With this configuration, we did not find parameters where both CE and wedging could act in parallel to form the tube. A two-step process, wedging followed by CE, was able to produce invagination and tube extension. However, a ring of basally constricting cells (with or without apically constricting disk) remedied this problem and allowed for wedging and CE to act in parallel. Supporting this, the

data by (*Sanchez-Corrales et al., 2018*) suggests that there are basally constricting cells in the outer
region of the placoid. Furthermore, basal and apical constriction seem to be induced by the same
organizing signal (*Gutzman et al., 2018*) through PCP pathways. Also in sea urchin gastrulation,
both types of wedging seem to be at play (*Kominami and Takata, 2004*). For simplicity, we limit our
simulations to basal wedging, where basally constricting cells are distributed on a ring (Figure 2A
and Figure 2-Figure supplement 5)

Our budding simulations thus show that successful invagination and tube elongation can
proceed if both wedging and PCP (and thus CE) act in parallel (Video 1, Figure 2A-C). We have also
succeeded in simulating sea urchin gastrulation where budding starts from a sphere of cells (Figure
3, Figure 3-Video 1, *Kimberly and Hardin* (*1998*); *Lyons et al.* (*2012*)). This proceeds essentially like
in the planar case. (see Methods for details).

In addition, we find that budding *can* proceed without wedging, however, robustness of the
outcome decreases in two ways. First, the proportion of failed invaginations is higher (Figure
2-Figure supplement 1). Second, the tube can form on either side of the epithelial plane.

With loss of apical constriction, noise is necessary to break the symmetry between the two sides
of the plane and initiate the CE-driven tubulation in one of the two directions orthogonal to the
plane. Thus, it seems that the role of wedging is to aid in the initial invagination and ensure correct
orientation. This is a plausible explanation for the results obtained in *Chung et al.* (*2017*) where the
authors knocked-out wedging in the context of *Drosophila* salivary gland formation and observed
that the budding process could still proceed, but with reduced reliability and orientational stability.
In contrast to our results and the findings by *Sanchez-Corrales et al.* (*2018*), *Chung et al.* (*2017*) do
not consider cell intercalations by CE but propose that supracellular myosin cables drive tissue
bending in the absence of wedging. It will be interesting to extend our approach to include an
analog of myosin cables through e.g. PCP-coupled supracellular forces, however, it is outside of the
scope of the current work.

Cell shape change, intercalation and tissue compression by supracellular myosin cables are also
key players in wrapping (*Nishimura et al., 2012*). The differences that cause some tubes to form
parallel and others orthogonal to the epithelial plane appear to be encoded in the geometrical
arrangement of the cells that participate in these three processes. In budding such cells are
arranged on a ring or a disk (circular symmetry), while in wrapping they are arranged on a band
(axial symmetry).

Anisotropic wedging and differential proliferation are sufficient for wrapping

139 To test if this difference in geometry alone is sufficient for wrapping, we choose a stripe of cells in
140 the middle of the epithelial sheet to represent the neuroepithelium (NE) (shown by gray in Figure
141 2D-E) and the rest of cells to represent ectoderm (E) (colored cells in Figure 2D-E). The NE cells are
142 then assigned anisotropic apical constriction and PCP pointing orthogonal to the future tube axis
143 (Figure 2-Figure supplement 4).

Wrapping requires anisotropy in wedging

In the case of isotropic wedging one would expect a collection of NE cells to eventually form a round
invagination or spherical lumen — the minimum energy state (Video 4). If we impose isotropic
wedging in our neurulation simulations, a bulging, rounded invagination is observed, see Video 3.

Motivated by the results of *Nishimura et al.* (*2012*), showing that wedging is anisotropic (Equa-
tion 3) and cells wedge primarily in the direction orthogonal to the tube axis, we asked if anisotropic
wedging can aid in tube closure. As expected, the tissue bends around the tube axis without capping
at the ends of the tube (Figure 1C, Figure 2-Figure supplement 2).

Interestingly, anisotropic wedging also leads to CE cell intercalation, narrowing and elongating
neuroepithelium (see Figure 2-Figure supplement 3), thus supporting the link between PCP-driven
wedging and cell intercalations. The simple intuitive argument for this comes from how wedged
cells pack in the tube. In the minimum energy state the extent of wedging, α, determines how many

cells can pack around the tube's circumference (Figure 1). To minimize energy, the "extra" cells will be displaced along the tube axis (see Figure 2-Figure supplement 3). CE-driven narrowing of the epithelium was proposed to be important for tube closure (*Wallingford et al., 2002*). In our simulations, wedging and CE alone succeeded in bending the tissue in an axially symmetric fashion (Figure 2-Figure supplement 2), however, we could not obtain successful tube closure even with maximally possible CE and wedging (both tuned by the strength of α in Equation 3). This suggests that additional mechanisms are necessary for final tube closure.

Buckling by proliferation at the NE boundary aids in tube closure

Images of neurulation cross-sections (see e.g. *Galea et al.* (*2018*)) show a strong bending at the neuroepithelium-ectoderm (NE) boundary with the curvature opposite to that inside of neuroepithelium (neural folds) (*Smith and Schoenwolf, 1997*). This is believed to be a result of combined forces from ectoderm due to i) change in cell shape (ectoderm cells become flatter and neuroepithelial cells become taller); ii) adhesion between basal surfaces of NE and E close to the neuroepithelium-ectoderm (NE-E) boundary (*Smith and Schoenwolf, 1997*) and iii) increase in cell density at this boundary either due to cell proliferation or intercalation (*McShane et al., 2015*).

Our goal was to test if the model can capture full tube closure with at least one of the mechanisms, so for simplicity, we focused on differential proliferation. When cells were set to proliferate only at the NE-E interface (*McShane et al., 2015*), we found that the resulting buckling can lead to successful neural tube closure (Figure 2-Video 2). In the simulations, the out-of-equilibrium buckling created by rapid cell proliferation is necessary to create a narrow neck that allows epithelial folds to fuse. We find that tubulation is possible within a rather broad range of cell cycles (3h–16h). Shorter *or* longer cell cycles resulted in open-tube morphologies reminiscent of neural tube defects such as spina bifida (Figure 4). In both cases the folds are too far apart to fuse, but for different reasons. If proliferation is too slow, the folds are far apart because the buckling is too weak. On the other hand, when proliferation is too fast, the sheet does not have time to equilibrate and CE does not catch up in narrowing it. Because of this, some sections of the tube become too wide to fuse. Interestingly this can sometimes lead to tube doubling/splitting (Figure 4-Figure supplement 1).

The effect of slow proliferation in our simulations is in line with the experimental data. *Copp et al.* (*1988*) showed that low proliferation rates could lead to neural tube defects in mice. In humans, mutations of the PAX3 transcription factor are implicated in Waardenburg syndrome (*Tassabehji et al., 1993*; *Baldwin et al., 1994*) characterized by incomplete neural tube closure. The same transcription factor has been shown (*Wu et al., 2015*) to be essential in ensuring sufficient cell proliferation. The effect of increased (compared to wild-type) proliferation has not been addressed experimentally and we hope that our predictions will motivate experiments in this direction.

Discussion

Larger organisms rely on tubes for distributing nutrients across the body as well as for exocrine functions. How these tubes reliably form is an open question, but a few recurrent mechanisms are known, e.g. directed or differential proliferation, changes in cell shapes, supracellular myosin cables, directed adhesion and cell rearrangements. As evolution proceeds by tinkering rather than engineering, it is not surprising that these mechanisms have overlapping functions. Recently quantitative experiments (*Chung et al., 2017*; *Sanchez-Corrales et al., 2018*; *Nishimura et al., 2012*) enabled us to look beyond a "one mechanism, one function" relationship and towards a map of where mechanisms overlap and how they complement each other.

In this work, we have made a step towards charting the functional overlap and complementarity among CE, wedging, and proliferation. A phenomenological point-particle representation allowed us for the first time to combine PCP-driven cell intercalation (CE) and anisotropic wedging in thousands of cells in 3D and with a few free parameters.

This allowed us to arrive at the following key results. First, our simulations recapitulate that CE can drive invagination in the absence of wedging (*Chung et al., 2017*; *Röper, 2012*). Thus suggesting

that this is a general mechanism, does not require forces from surrounding tissues and is possible in invaginating systems, other than the salivary glands in *Drosophila*. The invagination is however unreliable and isotropic wedging plays a complementary role by setting the direction of invagination.

Second, our results predict that anisotropic, PCP-coupled wedging may play a role in tube formation and elongation. Our model predicts that anisotropy in wedging maintains axial symmetry of the tube during wrapping. Remarkably, anisotropic wedging can also effectively result in CE-like cell intercalation and lead to tube elongation. While we have only tested the contribution of anisotropic wedging in wrapping, the same principle may apply in budding, were the reported isotropic wedging (*Röper, 2012*; *Sanchez-Corrales et al., 2018*) seems to become anisotropic along tube axis as the tube is elongating (*Pirraglia et al., 2010*). It will be interesting to explore this hypothesis theoretically and experimentally. Such isotropic to anisotropic transition in wedging has been reported in *Drosophila* furrow formation (*Leptin and Roth, 1994*; *Sweeton et al., 1991*). The visual inspection of tube cross sections in the pancreas and kidneys suggest that cells are wedged, and while by analogy to neurulation it is reasonable to expect wedging to be anisotropic, it remains to be confirmed experimentally by e.g. whole mount 3D imaging of stained tubes.

Third, buckling by differential proliferation (faster at the neuroepithelium/ectoderm boundary than in the remaining tissue) together with anisotropic wedging within neuroepithelium is sufficient for tube closure and separation. Differential proliferation has been proposed by *McShane et al.* (*2015*) as a mechanism for forming DLHP – regions where the tissue curvature has the same sign as at medial hinge points (MHP). We find that modifying the extent of apical constriction or how it is distributed – i.e. throughout entire neuroepithelium, or combinations of DLHPs and MHP, could not result in tube closure. Instead, our results highlight the importance of creating opposite curvature at the boundaries. Our simulations suggest that differential proliferation buckles the boundaries and aids tube closure as it curves the epithelium opposite to the curvature resulting from apical constriction.

Our simulations predict a wide range of proliferation rates capable of producing sufficient buckling for closure. These results call for testing for differential proliferation in systems without DLHP's (by accelerating or reducing proliferation rate in mutants or by molecular inhibitors (*Li et al., 2017*). While not immediately feasible, it is also interesting to consider how to perturb the "opposite" curvature by interfering with differences in cell shapes or basal adhesion (*Smith and Schoenwolf, 1997*) of the neuroepithelium and ectoderm close to the boundary.

Computational models of neurulation date back at least to *Kerszberg and Changeux* (*1998*) who also considered the effects of proliferation in a 2D model. Their model does not consider AB polarity and PCP but rather assumes unbreakable predetermined bonds between cells, allowing cells to move rather like a string of beads. This is unable to capture the interplay of polarities as well as the effects of anisotropy. A 3D vertex model of neurulation focusing on the three mechanisms of cell elongation, apical constriction, and active cell migration was formulated by *Inoue et al.* (*2016*). The model does not include cell proliferation but instead relies on active cell migration to pull the neural cells towards the midline. The model assumes successful neurulation when the folds are brought sufficiently close, but does not cover the separation of the tube from the sheet. Also, a recent 3D model of tube budding in lung epithelium arrived at the conclusion that anisotropic wedging can only result in rounded tubes and is insufficient to drive the entire process (*Kim et al., 2013*).

We have demonstrated that cell wedging can be phenomenologically captured in a point-particle representation. This is not restricted to apical constriction but also covers e.g. basal constriction and can, in a similar spirit, be extended to capture changes in cell height and width. This could allow for modeling a wider range of phenomena where morphological changes are driven by these differences. Furthermore, we are now in a position to address tube branching in e.g. lungs and vascularization, where cells forming the tubes also are the ones that secrete organizing signals that locally re-orient PCP polarities and may induce anisotropic changes in cell shapes.

Model and Methods

Following *Nissen et al.* (*2018*), cells are treated as point particles interacting with neighboring cells through a pair-potential V_{ij}. The potential has a rotationally symmetric repulsive term and a polarity-dependent attractive term. In terms of r_{ij} (the distance between cells i and j), the dimensionless potential can be formulated as

$$V_{ij} = e^{r_{ij}} - \left[\lambda_1 \, S_{ij}(A) + \lambda_2 \, S_{ij}(AP) + \lambda_3 \, S_{ij}(P) \right] e^{-r_{ij}/5}. \tag{4}$$

The parameters λ_i are coupling constants which define the strength of polar interactions in the model. $S_{ij}(A)$ quantifies the coupling between AB polarity and position, whereas $S_{ij}(AP)$ and $S(P)_{ij}$ quantify the coupling of PCP with AB and position, respectively, as described in *Nissen et al.* (*2018*). These couplings are formulated in in terms of AB vectors $\mathbf{p}_i$, PCP vectors $\mathbf{q}_i$ and a unit vector $\hat{\mathbf{r}}_{ij}$ from cell i to j. The coupling $S_{ij}(AP) = (\mathbf{p}_i \times \mathbf{q}_i) \cdot (\mathbf{p}_j \times \mathbf{q}_j)$ dynamically maintains the orthogonality of the PCP unit vectors $\mathbf{q}_i$ and $\mathbf{q}_j$ to their corresponding AB polarity vectors while lateral organization is favored by $S_{ij}(P) = (\hat{\mathbf{r}}_{ij} \times \mathbf{q}_i) \cdot (\hat{\mathbf{r}}_{ij} \times \mathbf{q}_j)$. In the absence of any cell shape effects, the coupling between AB and position is given by $S_{ij}(A) = (\hat{\mathbf{r}}_{ij} \times \mathbf{p}_i) \cdot (\hat{\mathbf{r}}_{ij} \times \mathbf{p}_j)$, which favours a flat cell sheet. Wedging of cells is introduced into our model by a single deformation parameter α, which describes an attractive interaction between the AB polarity unit vectors $\mathbf{p}_i$ and $\mathbf{p}_j$:

$$S_{ij}(A) = (\hat{\mathbf{r}}_{ij} \times \tilde{\mathbf{p}}_i) \cdot (\hat{\mathbf{r}}_{ij} \times \tilde{\mathbf{p}}_j), \tag{5}$$

where $\tilde{\mathbf{p}}_i$ is given by

$$\tilde{\mathbf{p}}_i = \mathbf{p}_i \quad \text{(for no wedging)},$$

$$\tilde{\mathbf{p}}_i = \frac{\mathbf{p}_i - \alpha \hat{\mathbf{r}}_{ij}}{|\mathbf{p}_i - \alpha \hat{\mathbf{r}}_{ij}|} \quad \text{(for isotropic wedging)},$$

$$\tilde{\mathbf{p}}_i = \frac{\mathbf{p}_i - \alpha \langle \hat{\mathbf{q}} \rangle_{ij}}{|\mathbf{p}_i - \alpha \langle \hat{\mathbf{q}} \rangle_{ij}|} \quad \text{(for anisotropic wedging)}. \tag{6}$$

Here, $\langle \hat{\mathbf{q}} \rangle_{ij}$ denotes the mean of PCP vectors $\mathbf{q}_i$ and $\mathbf{q}_j$ belonging to the two interacting cells. $\alpha = 0$ favors a flat sheet (see Figure 1A–B) whereas a non-zero α favours bending of AB polarity vectors towards (or away from) one another and induces curvature in a sheet of cells (Figure 1C–D).

The time development is simulated by overdamped (relaxational) dynamics along the gradient of the above potential, Equation (4). In the case of cell divisions, new daughter cells are placed randomly around the mother cell at a distance of one cell radius.

The source code for the simulations is available on GitHub (*Nielsen, 2019*).

Parameter estimation and robustness

We have tested the robustness of our approach on a number of model cases and find that, for example, *budding* can be reproduced with a broad range of wedging parameters, $\alpha \in [0.1, 0.6]$ and for diverse PCP coupling strengths $\lambda_3 \in [0.8, 0.14]$. For these intervals, the budding is qualitatively similar to that illustrated in Figure 2A.

We further explore our model by re-instating dimensions in the formulation of the potential and the equation of motion and estimating dimensionful quantities. With dimensions reinstated, the pair-potential takes the form

$$V_{ij} = V_0 \left[\exp(-r/\ell) - S \exp(-r/(\beta \ell)) \right]. \tag{7}$$

The overdamped equation of motion (without noise) becomes

$$0 = \gamma \mathbf{v}_i + \frac{\partial V_{ij}}{\partial \mathbf{r}_i}, \tag{8}$$

where $\mathbf{v}_i = \partial \mathbf{r}_i / \partial t$. We now introduce dimensionless (tilded) parameters

$$V_{ij} = V_0 \tilde{V}_{ij}, \quad \mathbf{r}_i = \ell \tilde{\mathbf{r}}_i, \quad v_i = v_0 \tilde{v}_i = \frac{\ell}{t_0} \tilde{v}_i. \tag{9}$$

and insert the dimensionless parameters in our equation of motion

$$\tilde{\mathbf{v}}_i = -\frac{V_0}{\ell \gamma v_0} \frac{\partial \tilde{V}_{ij}}{\partial \tilde{\mathbf{r}}_i}.$$ (10)

We expect $\left|\frac{\partial \tilde{V}_{ij}}{\partial \tilde{\mathbf{r}}_i}\right|$, and thus also $|\tilde{\mathbf{v}}_i|$ to be of order 1 in sheet-orthogonal extrusion. In *Eskandari and Salcudean* (*2008*), a typical value for the dynamical viscosity μ was reported to be on the order of 250Pa s. This should be seen in relation to the coefficient γ in Stokes' Law for viscous drag, $\gamma = 6\pi\mu\ell$. We now compare our model with epithelial cell extrusion and use the typical cell speed reported in *Yamada et al.* (*2017*), $v_0 \approx 1\text{mm h}^{-1}$ and use the typical cell size reported in *Brown and Bron* (*1987*), $2\ell = 13\mu\text{m}$. With these numbers, our model predicts a typical extrusion energy on the order of

$$12V_0 \approx 12 \times 6\pi\mu\ell^2 v_0 \approx 2 \times 10^{-13}\text{J}.$$ (11)

The factor of $12 = 2 \times 6$ is due to the hexagonal structure of the cell sheet. Note, that our estimate of the extrusion energy is consistent with the finding in *Yamada et al.* (*2017*) for epithelial cell extrusion. Here, an actomyosin ring is measured to exhibit a contraction force of the order of 1kPa, which results in an extrusion energy of the order $1\text{kPa} \times \ell^3 \approx 3 \times 10^{-13}\text{J}$.

Modeling neurulation/wrapping

The starting point for our simulation of neurulation is a planar sheet of cells where a line with a width of six cell radii is given non-zero wedging strength $|\alpha| = \alpha_0 > 0$ and all other cells have $\alpha = 0$. The line is centered at $x = 0$ and PCP is initialized orthogonally to this line, along the x direction ($\mathbf{q}|_{t=0} = \hat{\mathbf{x}}$). See Figure 2-Figure supplement 4.

Cell proliferation is simulated as a Poisson process by choosing a rate Γ for *each cell* to divide in each time unit. Only cells at the neuroepithelium-ectoderm interface (defined as cells with $|\alpha| > 0$ who are neighbours of cells with $\alpha = 0$) proliferate (with rate $\Gamma = \Gamma_0 > 0$) while the rest have $\Gamma = 0$. Daughter cells inherit all properties of their mother cell and are initiated randomly in a distance of one cell radius from their mother cell.

It should be noted that the initial width of the strip is not particularly important, since wedging will ensure the correct tube width given sufficient proliferation.

All cells in the simulation have the same coupling constants, typically $\lambda = (0.6, 0.4, 0)$. Typical values for Γ_0 and α_0 are 2.8×10^{-4} and 0.5. respectively.

Modeling gastrulation

In our gastrulation simulation, the assignment of PCP and cell wedging is characterized by two radii, describing an annulus (see Figure 2-Figure supplement 5):

$$r_0 = 7,$$ (12)

$$r_1 = 3r_0 = 21.$$ (13)

PCP is assigned within the disk Ω_1 given by

$$\Omega_1 = \left\{ (x, y, z) \left| \sqrt{x^2 + y^2} < r_1 \right. \right\}.$$ (14)

The PCP coupling strength λ is taken to be

$$\lambda = \begin{cases} (0.5, 0.5 - \lambda_3, \lambda_3) & \text{inside } \Omega_1, \\ (1, 0, 0) & \text{everywhere else.} \end{cases}$$ (15)

where a typical value for λ_3 is between 0.08 and 0.12.

The PCP vector field $\mathbf{q}$ is initially assigned so that it spirals around the axis of tube formation (the z-axis):

$$\mathbf{q}|_{t=0} = \hat{z} \times \mathbf{r},$$ (16)

In the gastrulation simulations, the PCP vector field is fixed on a per-cell basis.

Nonzero apical constriction parameter α is assigned in an annulus Ω_2, which shares its outer radius with the disk Ω_1:

$$\Omega_2 = \left\{ (x, y, z) \,\middle|\, r_0 < \sqrt{x^2 + y^2} < r_1 \right\}. \tag{17}$$

The magnitude of α for the cells in Ω_2 is taken as 0.4:

$$|\alpha| = \begin{cases} 0.4 & \text{inside } \Omega_2, \\ 0 & \text{everywhere else.} \end{cases} \tag{18}$$

296 The regions Ω_1 and Ω_2 are fixed in space and not on a particle basis. The number of particles in our
297 simulation is $N = 4000$.

Modeling budding from plane

298 The budding simulation is, apart from global topology, very similar to the gastrulation simulation.
The relevant length parameters are

$$r_0 = 5, \tag{19}$$

$$r_1 = 3r_0 = 15. \tag{20}$$

Two regions are correspondingly defined – the disk Ω_1 and the annulus Ω_2:

$$\Omega_1 := \left\{ (x, y, z) \,\middle|\, \sqrt{x^2 + y^2} < r_1 \right\}, \tag{21}$$

$$\Omega_2 := \left\{ (x, y, z) \,\middle|\, r_0 < \sqrt{x^2 + y^2} < r_1 \right\}. \tag{22}$$

The PCP coupling strength λ is taken to be

$$\lambda = \begin{cases} (0.5, 0.5 - \lambda_3, \lambda_3) & \text{inside } \Omega_1, \\ (1, 0, 0) & \text{everywhere else.} \end{cases} \tag{23}$$

where a typical value for λ_3 is between 0.08 and 0.12.

The PCP vector field $\mathbf{q}$ is initially assigned so that it spirals around the center of invagination (the origin of coordinates):

$$\mathbf{q}|_{t=0} = \hat{\mathbf{z}} \times \mathbf{r}, \tag{24}$$

In the gastrulation simulations, the PCP vector field is fixed on a per-cell basis.

Nonzero apical constriction parameter α is assigned in the annulus Ω_2 with magnitude 0.5:

$$|\alpha| = \begin{cases} 0.5 & \text{inside } \Omega_2, \\ 0 & \text{everywhere else.} \end{cases} \tag{25}$$

299 The total number of particles in the simulation is 1384.

Acknowledgments

This research has received funding from the Danish National Research Foundation (grant number: DNRF116) and the European Research Council under the European Union's Seventh Framework Programme (FP/2007 2013)/ERC Grant Agreement number 740704.

Declaration of interests

All authors of this article declare no competing interests.

References

Andrew DJ, Ewald AJ. Morphogenesis of epithelial tubes: Insights into tube formation, elongation, and elaboration. Developmental Biology. 2010; 341(1):34–55. doi: https://doi.org/10.1016/j.ydbio.2009.09.024.

Baldwin CT, Lipsky NR, Hoth CF, Cohen T, Mamuya W, Milunsky A. Mutations in PAX3 associated with Waardenburg syndrome type I. Human Mutation. 1994; 3(3):205–211. doi: https://doi.org/10.1002/humu.1380030306.

Belmonte JM, Swat MH, Glazier JA. Filopodial-tension model of convergent-extension of tissues. PLoS Computational Biology. 2016; 12(6):e1004952. doi: https://doi.org/10.1371/journal.pcbi.1004952.

Brown N, Bron AJ. An estimate of the human lens epithelial cell size in vivo. Experimental Eye Research. 1987; 44(6):899–906.

Choi SC, Sokol SY. The involvement of lethal giant larvae and Wnt signaling in bottle cell formation in Xenopus embryos. Developmental Biology. 2009; 336(1):68–75. doi: https://doi.org/10.1016/j.ydbio.2009.09.033.

Chung S, Kim S, Andrew DJ. Uncoupling apical constriction from tissue invagination. eLife. 2017; 6:e22235. doi: https://doi.org/10.7554/eLife.22235.

Collinet C, Rauzi M, Lenne PF, Lecuit T. Local and tissue-scale forces drive oriented junction growth during tissue extension. Nature Cell Biology. 2015; 17(10):1247. doi: https://doi.org/10.1038/ncb3226.

Copp AJ, Brook FA, Roberts HJ. A cell-type-specific abnormality of cell proliferation in mutant (curly tail) mouse embryos developing spinal neural tube defects. Development. 1988; 104(2):285–295.

Croce J, Duloquin L, Lhomond G, McClay DR, Gache C. Frizzled5/8 is required in secondary mesenchyme cells to initiate archenteron invagination during sea urchin development. Development. 2006; 133(3):547–557. doi: https://doi.org/10.1242/dev.02218.

Eskandari H, Salcudean SE. Characterization of the viscosity and elasticity in soft tissue using dynamic finite elements. In: *2008 30th Annual International Conference of the IEEE Engineering in Medicine and Biology Society* IEEE; 2008. p. 5573–5576. doi: https://doi.org/10.1109/IEMBS.2008.4650477.

Galea GL, Nychyk O, Mole MA, Moulding D, Savery D, Nikolopoulou E, Henderson DJ, Greene ND, Copp AJ. Vangl2 disruption alters the biomechanics of late spinal neurulation leading to spina bifida in mouse embryos. Disease models & mechanisms. 2018; 11(3):dmm032219. doi: https://doi.org/10.1242/dmm.032219.

Gutzman JH, Graeden E, Brachmann I, Yamazoe S, Chen JK, Sive H. Basal constriction during midbrain–hindbrain boundary morphogenesis is mediated by Wnt5b and focal adhesion kinase. Biology Open. 2018; 7(11):bio034520. doi: https://doi.org/10.1242/bio.034520.

Habib SJ, Chen BC, Tsai FC, Anastassiadis K, Meyer T, Betzig E, Nusse R. A localized Wnt signal orients asymmetric stem cell division in vitro. Science. 2013; 339(6126):1445–1448. doi: https://doi.org/10.1126/science.1231077.

Inoue Y, Suzuki M, Watanabe T, Yasue N, Tateo I, Adachi T, Ueno N. Mechanical roles of apical constriction, cell elongation, and cell migration during neural tube formation in Xenopus. Biomechanics and Modeling in Mechanobiology. 2016; 15(6):1733–1746. doi: https://doi.org/10.1007/s10237-016-0794-1.

Kerszberg M, Changeux JP. A simple molecular model of neurulation. BioEssays. 1998; 20(9):758–770. doi: https://doi.org/10.1002/(SICI)1521-1878(199809)20:9<758::AID-BIES9>3.0.CO;2-C.

Kim HY, Varner VD, Nelson CM. Apical constriction initiates new bud formation during monopodial branching of the embryonic chicken lung. Development. 2013; 140(15):3146–3155. doi: https://doi.org/10.1242/dev.093682.

Kimberly EL, Hardin J. Bottle cells are required for the initiation of primary invagination in the sea urchin embryo. Developmental Biology. 1998; 204(1):235–250. doi: https://doi.org/10.1006/dbio.1998.9075.

Kominami T, Takata H. Gastrulation in the sea urchin embryo: a model system for analyzing the morpho-
genesis of a monolayered epithelium. Development, Growth & Differentiation. 2004; 46(4):309–326. doi:
https://doi.org/10.1111/j.1440-169x.2004.00755.x.

Lee JY, Marston DJ, Walston T, Hardin J, Halberstadt A, Goldstein B. Wnt/Frizzled signaling controls C. ele-
gans gastrulation by activating actomyosin contractility. Current Biology. 2006; 16(20):1986–1997. doi:
https://doi.org/10.1016/j.cub.2006.08.090.

Leptin M, Roth S. Autonomy and non-autonomy in Drosophila mesoderm determination and morphogenesis.
Development. 1994; 120(4):853–859.

Li Y, Muffat J, Omer A, Bosch I, Lancaster MA, Sur M, Gehrke L, Knoblich JA, Jaenisch R. Induction
of expansion and folding in human cerebral organoids. Cell Stem Cell. 2017; 20(3):385–396. doi:
https://doi.org/10.1016/j.stem.2016.11.017.

Loh KM, van Amerongen R, Nusse R. Generating cellular diversity and spatial form: Wnt signal-
ing and the evolution of multicellular animals. Developmental Cell. 2016; 38(6):643–655. doi:
https://doi.org/10.1016/j.devcel.2016.08.011.

Lyons DC, Kaltenbach SL, McClay DR. Morphogenesis in sea urchin embryos: linking cellular events to gene
regulatory network states. Wiley Interdisciplinary Reviews: Developmental Biology. 2012; 1(2):231–252. doi:
https://doi.org/10.1002/wdev.18.

Martin AC, Gelbart M, Fernandez-Gonzalez R, Kaschube M, Wieschaus EF. Integration of contrac-
tile forces during tissue invagination. The Journal of Cell Biology. 2010; 188(5):735–749. doi:
https://doi.org/10.1083/jcb.200910099.

McShane SG, Molè MA, Savery D, Greene ND, Tam PP, Copp AJ. Cellular basis of neuroepithelial bend-
ing during mouse spinal neural tube closure. Developmental Biology. 2015; 404(2):113–124. doi:
https://doi.org/10.1016/j.ydbio.2015.06.003.

Nielsen BF, OrganogenesisPCP. GitHub; 2019. https://github.com/BjarkeFN/OrganogenesisPCP.

Nishimura T, Honda H, Takeichi M. Planar cell polarity links axes of spatial dynamics in neural-tube closure.
Cell. 2012; 149(5):1084–1097. doi: https://doi.org/10.1016/j.cell.2012.04.021.

Nissen SB, Rønhild S, Trusina A, Sneppen K. Theoretical tool bridging cell polarities with development of robust
morphologies. eLife. 2018; 7:e38407. doi: https://doi.org/10.7554/eLife.38407.

Ossipova O, Kim K, Lake BB, Itoh K, Ioannou A, Sokol SY. Role of Rab11 in planar cell polarity and api-
cal constriction during vertebrate neural tube closure. Nature Communications. 2014; 5:3734. doi:
https://doi.org/10.1038/ncomms4734.

Paluch E, Heisenberg CP. Biology and physics of cell shape changes in development. Current Biology. 2009;
19(17):R790–R799. doi: https://doi.org/10.1016/j.cub.2009.07.029.

Pirraglia C, Walters J, Myat MM. Pak1 control of E-cadherin endocytosis regulates salivary gland lumen size and
shape. Development. 2010; 137(24):4177–4189. doi: https://doi.org/10.1242/dev.048827.

Röper K. Anisotropy of Crumbs and aPKC drives myosin cable assembly during tube formation. Developmental
Cell. 2012; 23(5):939–953. doi: https://doi.org/10.1016/j.devcel.2012.09.013.

Sanchez-Corrales YE, Blanchard GB, Röper K. Radially patterned cell behaviours during tube budding from an
epithelium. eLife. 2018; 7:e35717. doi: https://doi.org/10.7554/eLife.35717.

Sawyer JM, Harrell JR, Shemer G, Sullivan-Brown J, Roh-Johnson M, Goldstein B. Apical constriction: a
cell shape change that can drive morphogenesis. Developmental Biology. 2010; 341(1):5–19. doi:
https://doi.org/10.1016/j.ydbio.2009.09.009.

Smith JL, Schoenwolf GC. Neurulation: coming to closure. Trends in Neurosciences. 1997; 20(11):510–517.

Spahn P, Reuter R. A vertex model of Drosophila ventral furrow formation. PLoS One. 2013; 8(9):e75051. doi:
https://doi.org/10.1371/journal.pone.0075051.

Sweeton D, Parks S, Costa M, Wieschaus E. Gastrulation in Drosophila: the formation of the ventral furrow and
posterior midgut invaginations. Development. 1991; 112(3):775–789.

Tassabehji M, Read AP, Newton VE, Patton M, Gruss P, Harris R, Strachan T. Mutations in the PAX3
gene causing Waardenburg syndrome type 1 and type 2. Nature Genetics. 1993; 3(1):26. doi:
https://doi.org/10.1038/ng0193-26.

Visetsouk MR, Falat EJ, Garde RJ, Wendlick JL, Gutzman JH. Basal epithelial tissue folding is me-
diated by differential regulation of microtubules. Development. 2018; 145(22):dev167031. doi:
https://doi.org/10.1242/dev.167031.

Wallingford JB, Fraser SE, Harland RM. Convergent extension: the molecular control of polarized cell movement
during embryonic development. Developmental Cell. 2002; 2(6):695–706. doi: https://doi.org/10.1016/S1534-
5807(02)00197-1.

Wu TF, Yao YL, Lai IL, Lai CC, Lin PL, Yang WM. Loading of PAX3 to mitotic chromosomes is mediated by
arginine methylation and associated with Waardenburg syndrome. Journal of Biological Chemistry. 2015;
290(33):20556–20564. doi: https://doi.org/10.1074/jbc.M114.607713.

Yamada S, Iino T, Bessho Y, Hosokawa Y, Matsui T. Quantitative analysis of mechanical force re-
quired for cell extrusion in zebrafish embryonic epithelia. Biology Open. 2017; 6(10):1575–1580. doi:
https://doi.org/10.1242/bio.027847.

Figure 2. Isotropic and anisotropic wedging drive budding and wrapping, respectively. Wedging cells are labeled in gray, with a shading that indicates the PCP direction.
(**A-C**) Time evolution of budding simulation (similar to salivary glands). Here, gray cells constrict basally and all cells on and inside the ring intercalate radially. Here, $\lambda_3 = 0.1$, $|\alpha| = 0.5$.
(**D-F**) Time evolution of wrapping simulation (similar to neurulation). Here, gray cells representing neuroepithelium, constrict apically and constriction is anisotropic, follows the direction of PCP (Equation 3). Cells proliferate only at the gray/colored boundary (with 7 hour doubling time), mimicking differential proliferation at the DHLP (*McShane et al., 2015*). Here, $\lambda_3 = 0$, $|\alpha| = 0.5$. Throughout, $\lambda_2 = 0.4$.

Figure 2–video 1. Video of budding simulation.

Figure 2–video 2. Video of neurulation simulation.

Figure 2–video 3. Video of neurulation simulation with isotropic wedging.

Figure 2–video 4. Video of a sheet of isotropically apically constricting cells forming a spherical lumen.

Figure 2–Figure supplement 1. Budding outcomes without wedging at high and low noise as well as in the absence of noise.

Figure 2–Figure supplement 2. The fate of the neural sheet in our simulations in the absence of proliferation.

Figure 2–Figure supplement 3. T1 transition induced by wedging.

Figure 2–Figure supplement 4. The initial configuration of the cell sheet for neurulation.

Figure 2–Figure supplement 5. The initial configuration of the cell sheet for budding.

Figure 3. Isotropic wedging in conjunction with PCP is sufficient to drive sea urchin gastrulation without external forcing. The gray ring shows cells with (isotropic) basal constriction and the shading indicates the direction of planar cell polarity, which curls around the vertical axis in our simulation. See section Modeling Gastrulation for details.
Figure 3–video 1. Video of sea urchin gastrulation simulation.

3.3

4 hours (Observed by McShane et al (2015))

23   Cell cycle/hrs

Figure 4. The cell cycle length at the neuroepithelial-ectoderm boundary affects tube closure. For cell cycle lengths below 3.3h and above 23h the neural tube fails to close in our simulations. It should be noted that this broad interval also contains the cell cycle length of 4h found for cells in the DLHP by *McShane et al.* (*2015*). The insets show outcomes of simulations run at short (2.6h), intermediate (12h) and long (26h) cell cycle lengths.
Figure 4–Figure supplement 1. Tube splitting observed with excessive proliferation rate.

	Normal	Failed	Misoriented
High Noise σ = 0.1 N = 50	40%	0%	60%
Low Noise σ = 0.002 N = 50	20%	50%	30%
No Noise σ = 0 N = 50	0%	100%	0%

Figure 2–Figure supplement 1. Budding outcomes without wedging at high and low noise as well as in the absence of noise. The first column shows the proportion of normal initiations of tubulation, the middle column shows failed invaginations while the last column shows misoriented invaginations.

Figure 2–Figure supplement 2. The fate of the neural sheet in our simulations in the absence of proliferation.

Figure 2–Figure supplement 3. T1 transition induced by wedging.

Figure 2–Figure supplement 4. The initial configurations of the cell sheet for neurulation. Wedging is turned on in a band of width d (gray) with PCP running orthogonal to this band.

Figure 2–Figure supplement 5. The initial configurations of the cell sheet for budding. Wedging is turned on in an annulus (gray) where PCP curls around tangentially.

Figure 4–Figure supplement 1. Tube splitting observed with excessive proliferation rate.

	Normal	Failed	Misoriented
High Noise $\sigma = 0.1$ $N = 50$	40%	0%	60%
Low Noise $\sigma = 0.002$ $N = 50$	20%	50%	30%
No Noise $\sigma = 0$ $N = 50$	0%	100%	0%

3.3

23   Cell cycle/hrs

4 hours (Observed by McShane et al (2015))