Contributions of spatial and temporal control of step length symmetry in the transfer of

locomotor adaptation from a motorized to a non-motorized split-belt treadmill

Daniel L. Gregory1 Frank C. Sup IV[2], Julia T. Choi[1,3]*

[1]Department of Kinesiology, University of Massachusetts Amherst, Amherst, MA, USA

[2]Department of Mechanical and Industrial Engineering, University of Massachusetts Amherst,

Amherst, MA, USA

[3]Neuroscience & Behavior Program, University of Massachusetts Amherst, Amherst MA, USA

Running Head: Motorized and non-motorized split-belt walking

*Corresponding Author:

Julia T. Choi

30 Eastman Lane

University of Massachusetts, Amherst

Amherst, MA 01003

Email: jtchoi@kin.umass.edu

Abstract

The ability to transfer locomotor learning to novel contexts is essential for effective gait rehabilitation. Locomotor adaptation is dependent upon independent processes for spatial and temporal limb control during motorized split-belt walking, where individual belt speeds are fixed with one belt moving faster than the other. Here we investigated the relative contributions of spatial and temporal learning strategies to the transfer of a motorized split-belt adaptation to a non-motorized split-belt treadmill, where each belt is independently user-propelled. In 10 healthy subjects, adaptation of step length symmetry was driven by changes to temporal and spatial motor outputs during motorized split-belt treadmill walking. We found robust after-effects in step length symmetry during transfer to non-motorized split-belt treadmill walking that were driven by changes in temporal motor outputs. We also observed residual after-effects during motorized tied-belt treadmill walking (post-transfer) that were mainly driven by changes in spatial motor outputs. These results suggest independent transfer and de-adaptation of spatial and temporal motor outputs. In addition, our results showed that step length after-effects were mediated by unique patterns of muscle activity during motorized versus non-motorized treadmill walking post-adaptation, suggesting that redundant locomotor outputs are expressed in a context-dependent manner. In sum, this is the first study to show that adaptive learning on a motorized split-belt treadmill is transferable to a non-motorized split-belt treadmill. Future research aims to unravel the independent contributions of spatial and temporal control to gait adaptation and transfer.

Keywords: human gait, interlimb coordination, generalization, electromyography, motor learning

Introduction

Control of human locomotion is highly adaptable. Gait patterns can be altered through

motorized treadmill training, even after central nervous system damage. For example, walking on

a split-belt treadmill which induces asymmetric step lengths by constraining limbs to move at

different speeds is altered over time through an error-based learning process known as motor

adaptation (Reisman et al., 2005, 2007). Training on a split-belt treadmill is promising because

asymmetric gait patterns can be reduced in hemiparetic walkers (Choi et al., 2009; Reisman et al.,

2007). From a clinical perspective, it is desirable to generalize adaptive learning to untrained

walking conditions. However, training on a motorized split-belt treadmill is context-dependent and

limited in the ability to generalize to over-ground walking or to different modes of locomotion

(Choi and Bastian, 2007; Hamzey et al., 2016; Ogawa et al., 2012; Torres-Oviedo and Bastian,

2010, 2012).

Adaptation of symmetric step lengths during motorized split-belt treadmill walking results

from changes in both spatial control of foot placement, and temporal control of step timing (Finley

et al., 2015; Long et al., 2016; Malone et al., 2012). These spatial and temporal motor outputs

reflect distinct neural strategies to overcome the asymmetric belt speeds. Positive asymmetries in

foot placement and step timing achieved during adaptation persist during de-adaptation while limb

excursions (distance the foot travels during stance) immediately return to tied baseline values. The

result is a negative step length asymmetry (i.e. after-effect) which requires time to de-adapt. When

given appropriate feedback about foot placement or step length, participants can block adaptation

of spatial control of locomotion without affecting temporal control (Long et al., 2016; Malone et

al., 2012). What remains to be shown is whether and how spatial and temporal asymmetries used

to reduce step length asymmetry during motorized split-belt treadmill walking transfer to novel

contexts.

Changes in muscle activation patterns during adaptation provides further information about

feedforward (predictive) and feedback (reactive) mechanisms for gait modifications (Iturralde and

Torres-Oviedo, 2019), but have yet to be quantified in a transfer task. Velocity dependent

resistance applied to hip and knee movements with an exoskeleton, or increases swing velocity

induced with an elastic force-field lead to feedforward adaptation of hamstring muscle activity,

evident by post-adaptation after-effects (Blanchette and Bouyer, 2009; Lam et al., 2006).

Furthermore, tibialis anterior muscle activity modulates in time with anterior breaking forces

during split-belt walking, leading to after-effects during the post-adaptation period (Ogawa et al.,

2014). This suggests that transfer of locomotor adaptations would be associated with muscle

responses which indicate a mismatch between current and required muscle control for the novel

environment.

The current study examined whether locomotor adaptation made during motorized split-

belt treadmill walking would transfer to non-motorized treadmill walking, indicated by after-

effects in spatial and temporal control, and muscle activity. Non-motorized treadmills have gained

in popularity for use in athletic settings and in sports science research; e.g., to measure cost of

transport (Smoliga et al., 2015), assess anaerobic performance (Gonzalez et al., 2013), and 5k time

trials (Stevens et al., 2015). Non-motorized treadmills have freely moveable belts which are driven

by participants pushing against an inclined surface. The user-propelled nature of the non-motorized

treadmill allows participants to self-select and express natural variability in gait. However,

research utilizing non-motorized treadmills in a clinical setting or for locomotor adaptation is

limited (Wang et al., 2017). Here, we developed a non-motorized split-belt treadmill that allows

the user to determine their own walking pace while simultaneously allowing for asymmetric

behavior. We hypothesized that due to the contextual similarity between the treadmills, after-

effects would be present during transfer. We further hypothesized that after-effects washed out

during non-motorized treadmill walking should lead to diminished or absent after-effects during

tied motorized treadmill walking. Finally, we hypothesized that leg muscle activity would display

adaptive behavior and therefore after-effects during the transfer condition.

Methods

Participants

Ten healthy volunteers (six female, four male; Age 26.5 ± 5.6 years) with the ability to

walk on a treadmill for 60 minutes and with no neurological or biomechanical impairments were

recruited for this study. All study protocols were approved by the University of Massachusetts,

Amherst Institutional Review Board and all subjects provided informed consent prior to

enrollment. None of the subjects had prior experience walking on a split-belt treadmill.

Experimental setup

Motorized split-belt treadmill. Subjects walked on an instrumented split-belt treadmill

(Bertec Corp, Columbus OH) that has separate left and right belts, each with its own motor and

force transducer. Belt speed acceleration was set at 1 m s^{-2}. During motorized treadmill walking,

subjects wore a non-weight-bearing safety harness suspended from the ceiling. At the beginning

of each trial, subjects were positioned in the middle of the treadmill with one leg on each belt and

were instructed to minimize handrail use and walk with normal arm swing as soon as possible

during all conditions.

Non-motorized spit-belt treadmill. We designed and built a user-propelled non-motorized

split-belt treadmill (**fig. 1a**). The non-motorized treadmill was fabricated from two commercially

available non-motorized treadmills (Inmotion II Manual Treadmill; Stamina, Springfield, MO),

which were designed to share a common support structure and minimize spacing between the belts

(32 mm). To minimize friction with the belt-deck interface, a sheet of polytetrafluoroethylene

(PTFE, 0.30" sheet; ePlastics, San Diego, CA) plastic was secured to the treadmill deck under each

belt independently. A ~1kg mass was added to the flywheel of each belt to increase the inertial

properties of the belt-flywheel system to ensure that the belt continued to move as the limb

transitioned from stance to swing which enabled a smoother transition into the next stance phase.

Each belt flywheel was also fitted with a friction tensioner to increase the resistance of the belt to

moving, however for this study all subjects used the non-motorized treadmill without the

tensioners attached to ensure symmetric resistance on both belts. The treadmill utilizes gravity

(~13° incline) and the users body weight to assist subjects driving the symmetrically resisted but

independently user-propelled belt speed. During non-motorized treadmill walking, subjects lightly

held onto the treadmill handrails throughout baseline and testing conditions and were instructed to

minimize handrail use to avoid supporting body weight. Typically, subjects reported only needing

the support of the handrails for the first two to three strides, however all subjects were encouraged

to maintain minimal contact with the handrail to minimize fall risk.

Split-belt adaptation paradigm

All experimental testing was completed in one session lasting approximately two hours.

The adaptation paradigm (**fig. 1b**) consisted of walking trials that included motorized tied-belt,

motorized split-belt and non-motorized split-belt conditions. Each trial began from a standstill and

subjects were instructed to "walk normally" in all conditions. During all walking conditions,

subjects were instructed to maintain a forward gaze and avoid looking at the feet.

During Baseline trials, subjects walked on the motorized treadmill with belts tied at a slow

(0.67 m/s), medium (1.0 m/s), and fast (1.34 m/s) speeds for two minutes each. The order of the

baseline speeds was randomized across subjects. A Baseline non-motorized treadmill trial was

collected at the subject's preferred walking speed for five minutes. Preferred speed during non-

motorized treadmill walking was determined by instructing subjects to "walk as fast as you would

to a meeting for which you have adequate time to arrive". During *Adaptation* trials, subjects

walked on the motorized split-belt treadmill at a 2:1 speed ratio (0.67 and 1.34 m/s) for ten minutes.

Immediately following adaptation, subjects were instructed to side-step from the motorized

treadmill onto the non-motorized treadmill which was located immediately in front of the

motorized treadmill. Typically, only two side steps were required to move onto the non-motorized

treadmill. Forward stepping was avoided to prevent washout of after-effects. During *Transfer*

trials, subjects walked for five minutes on the non-motorized treadmill to assess transfer of the

adapted locomotor pattern. Then, subjects returned to the motorized treadmill for the *Washout*

period by side-stepping and walked with belts tied at the slow speed (0.67 m/s) for an additional

five minutes. Finally, during *Post-adaptation* subjects walked on the non-motorized treadmill to

assess any residual after-effects. Subjects were randomly assigned left or right limb to the slow

belt during split-belt walking and subsequent references to each respective limb will be as slow or

fast limb regardless of the condition.

Data collection

During motorized treadmill walking, lower limb kinematics were recorded at 100 Hz using

a 4-camera Oqus system (Qualisys, Sweden). Reflective markers were placed bilaterally over the

fifth metatarsal, lateral malleolus, tibial plateau, greater trochanter, and the anterior superior iliac

spines. During non-motorized treadmill walking, lower limb kinematics were relegated to use of

only the ankle markers for identification of spatio-temporal parameters due to limitations in

capture space during data collection.

Electromyographic (EMG) activity was amplified (MA300, Motion Lab Systems, Baton

Rouge, LA) and collected using disposable surface electrodes (Ambu, blue sensor N-00-S) placed

in a bipolar configuration (inter-electrode distance 2 cm) over the tibialis anterior (TA), medial

gastrocnemius (MG), biceps femoris (BF) and the vastus lateralus (VL) according to the Surface

Electromyography for the Non-Invasive Assessment of Muscles project (Hermens et al. 2000).

System gain was set prior to baseline to ensure adequate signal detection and to prevent clipping.

EMG data were sampled at 2000 Hz and synchronized with kinematics data using Qualisys Track

Manager 2.11 (Qualisys, Göteborg, Sweden).

Data analysis

Data analysis was performed using custom software written in MATLAB (Mathworks,

Natick, MA, v. R2017a). Marker data were low pass filtered at 6 Hz with a second order

Butterworth filter. EMG data were high-pass filtered at 45 Hz (3^{rd} order Butterworth), demeaned,

rectified, and low-pass filtered at 35 Hz (3^{rd} order Butterworth).

The same method to determined heel contact time during motorized and non-motorized

treadmill walking (where ground reaction force data is unavailable) was used. We defined heel

contact time as the instant of peak anterior position of the ankle marker for each step and toe-off

as the instant of peak posterior position of the ankle marker for each step.

We used step length difference as our main outcome variable to assess split-belt walking

adaptation (Choi et al., 2009; Reisman et al., 2005, 2007; Roemmich and Bastian, 2015). Step

length was calculated as the anterior-posterior distance between the ankle markers at time of heel

contact. Step length difference was defined as $SL_{fast} - SL_{slow}$, where SL_{fast} is the step length with

fast leg leading at heel strike, and SL_{slow} is step length with slow leg leading (**fig. 2a**). We also

included double support difference as a secondary outcome (see Supplementary materials).

To quantify the spatial and temporal contribution to step length changes, we applied the

following analytical model of step length difference (for derivation, see Finley et al. 2015):

$$SL_{diff} = (\alpha_f - \alpha_s) + \left[\frac{v_s + v_f}{2} * (t_s - t_f) \right] + \left[\frac{t_s + t_f}{2} * (v_s - v_f) \right]$$

where α_f corresponds to where the fast foot is placed relative to the previous slow foot placement,

and α_s corresponds to where the slow foot is placed relative to the previous fast foot placement

(**fig. 2b,c**). v_f and v_s correspond to the speed of the fast and slow foot, respectively. t_f corresponds

to the time between the fast heel strike and the subsequence slow heel strike, and t_s corresponds to

the time between the slow heel strike and the subsequence fast heel strike.

Simplifying the equation to

$$SL_{diff} = \Delta\alpha + \bar{v} \cdot \Delta t + \bar{t} \cdot \Delta v$$

gives us an equation for step length difference where the first term is the step position component

which represents the difference between the relative positions of the feet at the fast and slow heel

strikes , the second term is the step time component which is the difference in slow and fast step

times as a function of average foot speed, and the third term is the step velocity component which

is the difference in slow and fast foot velocities as a function of the average step time. It is

important to note that during motorized treadmill conditions, v_f and v_s are fixed, whereas during

non-motorized treadmill walking, the speed of each belt/limb is strictly user defined and becomes

a third degree of freedom while walking.

To compare EMG across subjects and conditions, EMG amplitude was normalized to the

peak averaged EMG during baseline walking at 1 m/s as this was the average walking speed of

subjects on the non-motorized treadmill during pilot testing. Stride-by-stride changes in EMG were

measured as the mean EMG over a specific phase of the gait cycle for each muscle minus the

baseline speed and condition-matched EMG to form an EMG difference to compare changes

across Adaptation, Transfer and Washout. The following phases of the gait cycle were selected

post-hoc because they showed consistent changes across subjects and represent critical control

phases within the gait cycle for each muscle: a time window from the last 5% of the gait cycle (i.e.

95-100%) to the first 5% (i.e. 0-5%, early weight transfer) was used for the vastus lateralus and

tibialis anterior, referenced as -5:5%; 90-100% of the gait cycle (i.e. terminal swing) was used for

the biceps femoris; 15:25% of the gait cycle (i.e. mid-stance) was used for the medial

gastrocnemius. Note, these were not the only phases of the gait cycle which showed consistent

changes across subjects. However, we limited the analysis to one phase per muscle.

While all muscles were normalized to the same condition amplitudes (i.e. baseline medium

speed), the EMG differences were calculated relative to the condition and speed matched baseline

values across all conditions. During Adaptation, this means that the slow limb differences were

calculated relative to the slow limb baseline and the fast limb differences were calculated relative

to the fast limb baseline. We justified this with the understanding that, if there were no effects of

interlimb coordination, then each limb should individually have the same muscle activation

patterns relative to the baseline condition and speed matched counterparts. This assumes that once

heel contact occurs for a limb, this limb is now in the reference frame of the belt for which that

limb is in contact and therefore references to behavior should be done according to that reference

frame. We did this instead of calculating a difference of each curve from the same baseline value

(e.g. both limbs from medium speed baseline) to avoid differences in amplitude during the adapted

state that would not reflect the adapted state muscle dynamics that are a result of walking on a

split-belt treadmill. We are confident that this is a practical way to analyze this data as several

muscle activation patterns returned to baseline speed-matched values across the gait cycle by the

end of adaptation, specifically the fast VL and the slow MG and TA.

Statistical analysis

For each dependent variable, group means were calculated in time bins across the last 10

strides for each Baseline trial (motorized slow, motorized medium, motorized fast, and non-

motorized preferred) and across the first 5 strides (early) and last 5 strides (late) for Adaptation

(AE – early Adaptation, AL – late Adaptation), Transfer (TE – early Transfer, TL – late Transfer)

and Washout (WE – early Washout, WL – late Washout) trials. Separate repeated measures

ANOVAs were used to test the effects of time on each dependent variable. Post-hoc analyses were

performed using independent sample t-tests to compare each dependent variable at each time point

to the baseline value and for comparisons between early and late time points (i.e. AE vs AL).

Results

To assess the ability to transfer a motorized split-belt treadmill adaptation to non-motorized

split-belt treadmill walking, we exposed subjects to a 2:1 motorized split-belt treadmill condition

which caused large initial differences in step lengths. Subjects gradually adapted to reduce step

length difference to baseline levels over several minutes of walking. We subsequently tested

subjects walking at a preferred pace on the non-motorized split-belt treadmill to assess transfer;

adaptation transfer was evidenced by negative after-effects in step length difference. Finally,

robust after-effects were again present when we assessed for residual differences in a 1:1 motorized

treadmill condition at the slow speed. The group averaged step length data is presented below;

double support data is available in the supplementary materials.

Locomotor adaptation on a motorized split-belt treadmill

Repeated measures ANOVA revealed significant effects of time on step length ($F_{(7,63)}$ =

25.17, $p < 0.001$, **fig. 3a-5a**). Group averaged step-length differences were not significantly

different from zero during baseline tied- slow, medium or fast walking on the motorized split-belt

treadmill (all p-values > 0.05). **Figure 3a** shows gradual changes in step length difference (dark

grey/black) during the first 100 and last 20 strides in Adaptation. During *early Adaptation*, step

length difference showed a large initial change compared to baseline ($p < 0.001$, **fig. 3b,d**). By *late*

Adaptation, step length difference was reduced back to baseline levels ($p = 0.58$, **fig. 3b,d**).

As previously reported (Finley et al., 2015; Long et al., 2016; Malone et al., 2012), step

length difference can be decomposed into independent step velocity (magenta), step position (red)

and step time (blue) contributions (**fig 3c,d**). The step velocity component, which is largely driven

by the difference in belt speeds became increasingly negative over the course of adaptation,

reflecting a large negative velocity induced perturbation which required opposition ($p < 0.001$, **fig**

3d). Both the step position and step time components adapted to oppose the step velocity

component. During *early Adaptation*, the step position component showed a significant initial

asymmetry ($p < 0.001$, **fig. 3d**), however it was not large enough to counter the large negative step

velocity component, as evidenced by a large initial step length difference. The step time component

showed more gradual change, with *early Adaptation* values not significantly different from

baseline (p = 0.26, **fig. 3d**). By *late Adaptation*, the step position and step time components both

increased significantly (p < 0.001, **fig. 3d**) to cancel the negative step velocity component,

resulting in symmetrical step lengths (p = 0.58, **fig. 3d**).

Robust transfer of after-effect to non-motorized split-belt treadmill

We found significant transfer of motorized treadmill adaptations to the non-motorized

split-belt treadmill. **Figure 4a** shows the after-effect in step length difference during the first 100

and last 20 strides in Transfer. During *early Transfer*, step length difference showed a negative

after-effect compared to baseline (p < 0.001, **fig. 4b,d**). By *late Adaptation*, step length difference

returned to baseline levels (p = 0.31).

Figure 4c,d show the contribution of the step velocity, step position and step timing

components to the step length after-effects during Transfer. In *early Transfer*, the step velocity

component displayed a large positive (i.e. an after-effect) component that was significantly

different from baseline (p = 0.03, **fig. 4d**, magenta). By *late Transfer*, the step velocity component

decreased to a negative value that was significantly different from baseline (p = 0.02), indicating

that subjects settled on a step velocity difference in the same direction as the initial motorized split-

belt perturbation (i.e. a negative velocity difference). The step position component was not

significantly different from baseline during *early Transfer* (p = 0.5, **fig. 4d**). There were also no

significant differences between *early* and *late Transfer* for the step position component (p = 0.3).

By *late Adaptation,* the step position component was not different from baseline (p = 0.06, **fig.**

4d), however, upon inspection of individual data, this result may have been driven by a single

outlier (*late Transfer,* red filled circles). These results indicate that the spatial component did not

contribute to the initial after-effects in step length difference. The step time component was

significantly different from baseline during *early Transfer* (p < 0.001, **fig. 4d**). By *late Transfer*,

the step time component decreased back to baseline values (p = 0.58), indicating washout of the

temporal pattern.

Residual after-effects on return to motorized split-belt treadmill

301   Despite washout of step length after-effects during non-motorized treadmill walking in the

302   Transfer period, after-effects in step length reappeared during motorized treadmill walking in *early*

303   *Washout* (p < 0.001). Step length difference was significantly different from baseline during early

304   Washout (p < 0.001 **fig. 5a**) and returned back to baseline by *late Washout* indicated by significant

305   differences from *early* to *late Washout* (p < 0.001) but did not completely return to baseline values

306   (p = 0.03, **fig. 5b,d**).

307   **Figure 5c** shows the contribution of the step velocity, step position and step timing

components to the step length after-effects during the Washout period. The step velocity

components were not different from baseline during *early Washout* (p = 0.72) or *late Washout* (p

= 0.95), due to the symmetric belt speeds (**fig. 5c,d**). The step position component displayed

significant after-effects during *early Washout*, (p < 0.001) that reduced back to baseline by *late*

Washout (p = 0.17, **fig. 5d**). This is unlike the Transfer period, where the step position component

showed no after-effects (see **fig. 4d**, red). The step time component was not significantly different

from baseline during *early Washout* (p = 0.09) nor *late Washout* (p = 0.48, **fig 5d**, blue). Note

however that the step time component increased from low initial values during early Washout

(0.028m ± 0.03) to greater magnitudes around 6-10 strides into the Washout period (SL: ~0.04m)

which later returned to baseline values (**fig. 5c**) indicating a delayed emergence of the temporal

component.

Unlearning during transfer does not interfere with after-effects during washout

We performed a linear regression analysis to determine whether step length after-effects

present during the Transfer condition were the result of unlearning of the motorized treadmill

adaptation. We found that step length error magnitude during *early Transfer* did not predict the

step length error magnitude during *early Washout* ($p = 0.782$, **fig. 6**). In addition, we ran the linear

regression analysis on the individual components to see if there were any interaction effects on the

individual components that may not have been captured by the step length after effect. We found

no interaction between *early Transfer* and *early Washout* for the step position ($p = 0.132$), step

time ($p = 0.451$), nor the velocity components ($p = 0.457$).

EMG activity during motorized split-belt adaptation

Changes in EMG activity during Adaptation (left columns), Transfer (middle, gray

columns) and Washout (right columns) are illustrated, with the number of subjects analyzed for

each muscle indicated, in **figure 7**. Due to issues with EMG acquisition in select subjects, some

muscles were omitted from analysis throughout the experiment. Repeated measures ANOVA

revealed significant effects of time on both fast (VL: $F_{(7,35)} = 5,75$, $p < 0.001$; BF: $F_{(7,42)} = 6.08$,

$p < 0.001$; MG: $F_{(7,49)} = 15.76$, $p < 0.001$; TA: $F_{(7,42)} = 14.59$, $p < 0.001$) and slow limbs (VL:

$F_{(7,35)} = 13.24$, $p < 0.001$; BF: $F_{(7,42)} = 24.51$, $p < 0.001$; MG: $F_{(7,49)} = 9.96$, $p < 0.001$; TA:

$F_{(7,42)} = 13.85$, $p < 0.001$) (**fig. 7**).

During *early Adaptation* (solid line)*, muscle activity of the slow VL, BF, and TA (bottom

panel, cyan) were significantly greater during terminal swing and weight acceptance relative to

slow baseline values ($p < 0.05$, **fig. 7a,b,d**), but not for the MG ($p = 0.65$, **fig. 7c**). By *late*

Adaptation (dashed line), the slow VL and TA activation significantly decreased compared to *early*

Adaptation ($p < 0.05$) indicating significant modulation of muscle activity, yet did not reach

baseline values ($p < 0.05$, **fig. 7a,d**). The slow BF showed a decreasing trend from *early Adaptation*

to *late Adaptation* that was not significant ($p = 0.077$) and was still significantly different from

baseline during the last five strides ($p = 0.005$, **fig. 7b**). The slow MG did not change from *early*

Adaptation to *late Adaptation* ($p = 0.54$) and was not different from baseline during early ($p =$

0.65) or *late Adaptation* ($p = 0.67$, **fig. 7c**). The slow MG during terminal stance (not shown)

initially showed a significant increase in activity ($p < 0.001$) during *early Adaptation* which

remained slightly greater, albeit significantly, relative to baseline slow ($p = 0.008$) during *late*

Adaptation.

In contrast to the slow limb, the VL, BF, and TA of the fast limb (top panel, black) had

significantly lower activation during *early Adaptation* compared to the baseline fast condition (p

< 0.05, **fig. 7a,b,d**). Both the fast VL and TA increased significantly by *late Adaptation* ($p < 0.01$)

with no difference from baseline ($p > 0.05$, **fig. 7a,d**) indicating modulation of each muscle's

activity during terminal swing and early weight acceptance. The fast BF, however, did not change

from *early* to *late Adaptation* during terminal swing ($p = 0.53$, **fig. 7b**). The MG on the fast limb

showed similar behavior to that of the VL, BF, and TA of the slow limb with significant increase

in activation during *early Adaptation* ($p < 0.001$), a significant decrease by *late Adaptation* ($p =$

0.02), and a persistent difference from baseline values during *late Adaptation* ($p < 0.001$, **fig. 7c**),

again, indicating modulation of the muscles activity during this phase of the gait cycle. The fast

MG during terminal stance (not shown) was no different during *early Adaptation* relative to

baseline fast (p = 0.85) and made a non-significant decrease by *late Adaptation* which was

significantly lower than baseline (p < 0.001).

EMG activity during transfer to non-motorized treadmill walking post-adaptation

During *early Transfer*, a significant depression of muscle activity during terminal swing to

weight acceptance was found for the slow VL, BF, and TA indicating the presence of after-effects

(p < 0.003, **fig. 7e,f,h**). The activation of each muscle significantly increased back to baseline

values from *early* to *late Transfer* for the slow VL, BF, and TA (p > 0.05) and were not different

from baseline (p > 0.05 **fig. 7e,f,h**). During *early Transfer*, the slow MG displayed early positive

differences (p = 0.03), which returned to baseline values by *late Transfer* (p = 0.8, **fig. 7g**). The

slow MG during terminal stance in *early Transfer* was no difference from baseline non-motorized

treadmill walking (p = 0.21). By *late Transfer* the slow MG during terminal stance was reduced

relative to baseline (p = 0.005). Of the fast limb muscles, only the BF (p = 0.008, **fig. 7f**) and the

TA (p = 0.001, **fig. 7h**) had a significantly lower activation relative to baseline during *early*

Transfer, and no muscles were different from baseline by *late Transfer* (p ≥ 0.18, **fig. 7e-h**). The

fast MG during terminal stance in *early Transfer* was elevated relative to baseline (p = 0.01), and

like the slow MG during terminal stance showed a minor but significant depression (p < 0.001).

To ensure that suppressed muscle activity during the *early Transfer* period was not just due to

lower walking speeds on the slow limb, we compared the average belt velocity during the same

period to show that the fast limb was significantly slower than the slow limb (0.662 vs 0.755 m/s,

p = 0.03, **fig. 8, middle**).

EMG activity during washout on motorized treadmill post-transfer

Slow limb muscle activity of the VL, BF, and TA were no different from baseline values

($p \geq 0.13$) and did not change from early to late Washout ($p \geq 0.4$, **fig. 7i,j,l**). The slow MG,

however, displayed robust after effects, indicated by a significant increase in muscle activity

relative to baseline ($p < 0.001$, **fig. 7k**) that significantly decreased by late Washout ($p < 0.001$)

and was not different from baseline values ($p = 0.09$), indicating the pattern washed out. The slow

MG during terminal stance showed no difference to baseline throughout the washout period ($p >$

0.05). For the fast limb muscles, only the VL had a minor, yet significant increase in activity during

the early Washout period ($p = 0.01$, **fig. 7i**). However, a small decrease to late Adaptation occurred,

which did not reach significance ($p = 0.05$) and was also not significantly different from baseline

($p = 0.53$).

No remaining after-effects during post-washout non-motorized treadmill walking

No residual step length difference after-effects were present once returned to the non-

motorized treadmill post-washout ($p > 0.05$, not shown). In addition, there were no asymmetries

between any individual component during *early Post-washout* walking ($p > 0.05$). There were

significant decreases in muscle activity between *early Post-washout* walking compared to baseline

non-motorized treadmill values for bilateral VL ($p < 0.05$), BF ($p < 0.001$), and TA ($p < 0.001$),

but not the MG ($p > 0.05$). However, there were no significant differences between limbs for any

muscle (all $p > 0.05$). Additionally, no muscle made a significant change between *early* (first 10

strides) versus *late* (last 10 strides of a 2-minute period) *Post-washout* ($p > 0.05$), indicating that

the amplitude of these muscles was lower throughout the 2-minute *Post-washout* period. Upon

further analysis, this appears to be due to a lower average walking speed between *Baseline* non-

motorized treadmill walking for both slow and fast limbs (1.0 m/s, **fig. 8, left**) and *early Post-*

washout non-motorized treadmill walking (0.91 m/s, p < 0.001, **fig. 8, right**). To ensure that the

lower activation values for the three muscles were not due to residual after-effects, we made sure

there were no significant differences between foot velocities during *early Post-washout* (p = 0.998,

fig. 8, right).

Discussion

Context has been shown to play a major role in modulating the transfer of sensorimotor

adaptations when attempting to apply a motor pattern between environments. Specifically,

locomotor adaptation on a split-belt treadmill has repeatedly demonstrated limited transfer to over-

ground walking in healthy subjects (Reisman et al., 2009; Torres-Oviedo and Bastian, 2010, 2012).

However, patients post-stroke demonstrated greater transfer to over-ground walking, a difference

the authors attributed to a limited ability to utilize context cues as a result of cerebral deficits

(Reisman et al., 2009). Removal of vision either during training, testing, or both has been shown

to substantially improve transfer in healthy walkers, with the greatest transfer present when vision

was occluded in both training and transfer conditions (Torres-Oviedo and Bastian, 2010). The

removal of vision during transfer of locomotor adaptations is thought to prevent the utilization of

visual contextual cues which provide the nervous system with context specific information critical

to successful movement (Gordon et al., 1995; Torres-Oviedo and Bastian, 2010).

Here we demonstrated large after-effects in a new context (i.e. non-motorized split-belt

treadmill walking) after adapting on a motorized split-belt treadmill. In the current study, vision

was never occluded, therefore contextual cues about the walking environment were available

throughout training, transfer, and washout conditions. The only switch in context between

conditions was the treadmill used during walking, which while having several key differences (e.g.

incline, propulsion dynamics), still allowed subjects to maintain normal limb and muscle

coordination. Our results showed robust after-effects during Transfer, which suggest that the

contextual cues available from vision and proprioception were insufficient to block transfer.

Instead, consistency of the dynamics or highly overlapping neural networks between the conditions

may be more relevant to the ability to transfer newly learned locomotor adaptations to a new

context (Choi and Bastian, 2007).

We hypothesized that the washout of after-effects during the Transfer period would lead to

smaller or no after-effects in the subsequent Washout period (Reisman et al., 2009). However, we

found large after-effects in both Transfer and Washout that were driven by asymmetries in step

timing and foot placement differences, respectively. We believe this suggests three potential

mechanisms. First, if the spatial and temporal parameters were uncoupled, this suggests

independent access to the control of spatial and temporal motor outputs, and possibly independent

washout of each parameter in consecutive conditions. Second, the apparent uncoupling of spatial

and temporal after-effects could be a re-organization of spatial, temporal, and velocity

asymmetries resulting from an additional degree of freedom (i.e. belt velocity) when walking on

the non-motorized treadmill. This may effectively save the adapted pattern to be later washed out.

Last, the persistent after-effects in both Transfer and Washout conditions could be an artifact of

walking speed, as this factor has been demonstrated previously to impact after-effect magnitude

following a locomotor adaptation (Vasudevan and Bastian, 2010).

First, previous work has demonstrated the ability to uncouple spatial and temporal control

of locomotor adaptation using feedback during the adaptation period (Long et al., 2016; Malone

et al., 2012). Unlike the previous studies, the proposed uncoupling here occurred during the

Transfer and Washout periods. On the non-motorized treadmill, visual and tactile cues of the

handrail and walking surface may have led to an unintentional restriction of foot placement

asymmetry. In line with this idea, there were no significant changes in foot placement differences

during transfer as step length after-effects were a result of step time and belt velocity asymmetries.

This suggests that by restricting the spatial motor output during Transfer, the temporal motor

output was free to vary and likely washed out, leaving the spatial motor output to be washed out

later during the Washout condition.

Several recent studies indicate that spatial and temporal control of gait may be associated

to functionally distinct neural circuitries. Spatial control has been associated with both state-

estimation (Maeda et al., 2017), a cerebellar process, and pre-planned cognitive control (i.e.

cortical processes) in challenging walking environments (Matthis et al., 2018). In cerebellar lesion

patients, double support difference, but not step length symmetry displays adaptive behavior

(Morton and Bastian, 2006). Hemispherectomy patients have been shown to adapt step length

symmetry, but not double support difference in split-belt walking (Choi et al., 2009). Together,

these studies suggest that decoupling and subsequent independent washout of the temporal and

spatial components of locomotion is possible if constraints limit access to the neural circuits which

are involved in the control of a specific parameter. Our findings here are in line with this idea and

suggest that during split-belt treadmill walking, uncoupling of spatial and temporal components

are dissociable and can be implicitly and independently accessed.

Alternatively, the apparent dissociation between the spatial and temporal components

during the Transfer and Washout conditions may be an artifact of having a third degree of freedom

(i.e. belt speed) while walking on the non-motorized split-belt treadmill. Close inspection of the

individual components during the Transfer condition (**Fig. 4c,d**) might suggest a trade-off in

spatial, temporal, and velocity components, which sum to symmetric step lengths. By the end of

Transfer, belt speeds were asymmetric in the same direction as during Adaptation, while the spatial

component had a positive asymmetry (**Fig. 4d**). Because the context and dynamics of each

treadmill were so similar, this would suggest that subjects persisted with their adapted state from

the motorized split-belt treadmill to trade-off symmetric step times and lengths with asymmetries

in foot placement and step velocity. Consistent with this idea, previous work has demonstrated the

high metabolic cost of locomotion with asymmetric step times (Ellis et al., 2013). Further, step

length asymmetry alone during split-belt walking does not explain the added cost of asymmetry

(Sanchez et al., 2017). This may suggest that subjects preferred temporal symmetry rather than the

foot placement or velocity symmetry as a means of reducing metabolic cost. This further suggests

that subjects adapted a newly learned motor pattern for a novel condition in a way that was more

efficient while meeting the demands of the novel environment without washing out their adapted

motor pattern.

Partial washout or persistent after-effects during Transfer and Washout could also be

caused by after-effects associated with different walking speeds between conditions. The

contribution of speed to after-effect size has been demonstrated previously in both over-ground

and treadmill walking (Hamzey et al., 2016; Vasudevan and Bastian, 2010). Speeds both faster or

slower than the slow speed during adaptation result in diminished after-effects. In the current

study, we show robust after-effect magnitudes during both the Transfer and the Washout periods

independent of speed; i.e. average walking speed during non-motorized treadmill walking in

Transfer was significantly greater than the slow motorized speed (0.8 m/s versus 0.67 m/s, p <

0.05). Walking at a faster speed in Transfer should have led to smaller after-effect magnitudes,

however, this was not the case (p = 0.26). Moreover, we demonstrated no association between

after-effects in Transfer and Washout, suggesting partial washout did not occur. Therefore, we

believe it is unlikely that the persistent after-effects were partial after-effects in each condition as

a function of walking speed.

Finally, flexible control of walking gait is demonstrated here with dissimilar muscle

behavior for similar step length after-effects. Adaptation of muscle activity, but not merely reactive

changes, is demonstrated with robust limb- and phase-dependent after-effects in the Transfer

condition. Specifically, the slow VL, BF, and TA show large suppressions of muscle activity

during late swing and early weight transfer which require substantial time (>25 strides) to recover

to baseline non-motorized treadmill walking values. Further, we also show that the suppression of

muscle activity during this period is not simply a result of slower limb speeds during the early

Transfer period, but instead are suppressed despite faster speeds compared to the fast limb. Tibialis

anterior muscle activity has been demonstrated to display after-effects following split-belt

treadmill walking, behavior which was correlated with anterior breaking forces (Ogawa et al.,

2014). Here we extend these findings to show that the slow limb VL and BF muscles also show

similar predictive control around heel strike and weight acceptance which track similar adaptive

changes found previously (Iturralde and Torres-Oviedo, 2019; Ogawa et al., 2014). This suggests

that muscles of the lower and upper leg participate in predictive control of heel contact and weight

acceptance in split-belt walking and are likely subject to formation of, or modification to, an

internal model which represents the dynamics of lower limb control (Blanchette and Bouyer, 2009;

Lam et al., 2006).

In conclusion, we used a novel non-motorized split-belt treadmill in a locomotor transfer

paradigm to demonstrate that context and locomotor dynamics influence the ability to transfer a

newly learned locomotor adaptation. This study improves on our current understanding of the

transfer of locomotor adaptations and provides further evidence of the adaptability of the human

locomotor system for the control of walking gait in novel challenging environments. We believe

that the low-cost and portable non-motorized split-belt treadmill may provide a unique and useful

tool in gait rehabilitation. Further research is needed to investigate how non-motorized split-belt

treadmills can be used in clinical and home settings, and whether training on this device would

lead to improved over-ground walking.

Acknowledgements

This work is supported by the National Science Foundation grant #1264752 to F.C.S and the

Initiative for Maximizing Student Development (UMass Amherst) to D.L.G. We thank Erin Carey,

Jeffrey Florek, Nicholas Lake, Subhash Vallala, and Eric Murray for designing and building the

non-motorized split-belt treadmill.

References

Blanchette, A., and Bouyer, L.J. (2009). Timing-Specific Transfer of Adapted Muscle Activity After Walking in an Elastic Force Field. Journal of Neurophysiology *102*, 568–577.

Choi, J.T., and Bastian, A.J. (2007). Adaptation reveals independent control networks for human walking. Nature Neuroscience *10*, 1055–1062.

Choi, J.T., Vining, E.P.G., Reisman, D.S., and Bastian, A.J. (2009). Walking flexibility after hemispherectomy: split-belt treadmill adaptation and feedback control. Brain *132*, 722–733.

Ellis, R.G., Howard, K.C., and Kram, R. (2013). The metabolic and mechanical costs of step time asymmetry in walking. Proceedings of the Royal Society B: Biological Sciences *280*, 20122784.

Finley, J.M., Long, A., Bastian, A.J., and Torres-Oviedo, G. (2015). Spatial and Temporal Control Contribute to Step Length Asymmetry During Split-Belt Adaptation and Hemiparetic Gait. Neurorehabilitation and Neural Repair *29*, 786–795.

Gonzalez, A.M., Wells, A.J., Hoffman, J.R., Stout, J.R., Fragala, M.S., Mangine, G.T., McCormack, W.P., Townsend, J.R., Jajtner, A.R., Emerson, N.S., et al. (2013). Reliability of the Woodway Curve(TM) Non-Motorized Treadmill for Assessing Anaerobic Performance. J Sports Sci Med *12*, 104–108.

Gordon, C.R., Fletcher, W.A., Melvill Jones, G., and Block, E.W. (1995). Adaptive plasticity in the control of locomotor trajectory. Exp Brain Res *102*, 540–545.

Hamzey, R.J., Kirk, E.M., and Vasudevan, E.V.L. (2016). Gait speed influences aftereffect size following locomotor adaptation, but only in certain environments. Exp Brain Res *234*, 1479–1490.

Iturralde, P.A., and Torres-Oviedo, G. (2019). Corrective Muscle Activity Reveals Subject-Specific Sensorimotor Recalibration. ENeuro *6*, ENEURO.0358-18.2019.

Lam, T., Anderschitz, M., and Dietz, V. (2006). Contribution of Feedback and Feedforward Strategies to Locomotor Adaptations. Journal of Neurophysiology *95*, 766–773.

Long, A.W., Roemmich, R.T., and Bastian, A.J. (2016). Blocking trial-by-trial error correction does not interfere with motor learning in human walking. Journal of Neurophysiology *115*, 2341–2348.

Maeda, R.S., O'Connor, S.M., Donelan, J.M., and Marigold, D.S. (2017). Foot placement relies on state estimation during visually guided walking., Control of Movement: Foot placement relies on state estimation during visually guided walking. J Neurophysiol *117, 117*, 480, 480–491.

Malone, L.A., Bastian, A.J., and Torres-Oviedo, G. (2012). How does the motor system correct for errors in time and space during locomotor adaptation? Journal of Neurophysiology *108*, 672–683.

Matthis, J.S., Yates, J.L., and Hayhoe, M.M. (2018). Gaze and the Control of Foot Placement When Walking in Natural Terrain. Current Biology *28*, 1224-1233.e5.

Morton, S.M., and Bastian, A.J. (2006). Cerebellar Contributions to Locomotor Adaptations during Splitbelt Treadmill Walking. Journal of Neuroscience *26*, 9107–9116.

Ogawa, T., Kawashima, N., Ogata, T., and Nakazawa, K. (2012). Limited Transfer of Newly Acquired Movement Patterns across Walking and Running in Humans. PLOS ONE *7*, e46349.

Ogawa, T., Kawashima, N., Ogata, T., and Nakazawa, K. (2014). Predictive control of ankle stiffness at heel contact is a key element of locomotor adaptation during split-belt treadmill walking in humans. Journal of Neurophysiology *111*, 722–732.

Reisman, D.S., Block, H.J., and Bastian, A.J. (2005). Interlimb Coordination During Locomotion: What Can be Adapted and Stored? Journal of Neurophysiology *94*, 2403–2415.

Reisman, D.S., Wityk, R., Silver, K., and Bastian, A.J. (2007). Locomotor adaptation on a split-belt treadmill can improve walking symmetry post-stroke. Brain *130*, 1861–1872.

Reisman, D.S., Wityk, R., Silver, K., and Bastian, A.J. (2009). Split-Belt Treadmill Adaptation Transfers to Overground Walking in Persons Poststroke. Neurorehabil Neural Repair *23*, 735–744.

Roemmich, R.T., and Bastian, A.J. (2015). Two ways to save a newly learned motor pattern. J Neurophysiol *113*, 3519–3530.

Sanchez, N., Park, S., and Finley, J.M. (2017). Evidence of Energetic Optimization during Adaptation Differs for Metabolic, Mechanical, and Perceptual Estimates of Energetic Cost. Scientific Reports *7*, 7682.

Smoliga, J.M., Hegedus, E.J., and Ford, K.R. (2015). Increased physiologic intensity during walking and running on a non-motorized, curved treadmill. Physical Therapy in Sport *16*, 262–267.

Stevens, C.J., Hacene, J., Sculley, D.V., Taylor, L., Callister, R., and Dascombe, B. (2015). The Reliability of Running Performance in a 5 km Time Trial on a Non-motorized Treadmill. Int J Sports Med *36*, 705–709.

Torres-Oviedo, G., and Bastian, A.J. (2010). Seeing Is Believing: Effects of Visual Contextual Cues on Learning and Transfer of Locomotor Adaptation. Journal of Neuroscience *30*, 17015–17022.

Torres-Oviedo, G., and Bastian, A.J. (2012). Natural error patterns enable transfer of motor learning to novel contexts. Journal of Neurophysiology *107*, 346–356.

Vasudevan, E.V.L., and Bastian, A.J. (2010). Split-Belt Treadmill Adaptation Shows Different Functional Networks for Fast and Slow Human Walking. Journal of Neurophysiology *103*, 183–191.

602   Wang, J.-C., Sung, W.-H., Chang, Y.-L., Wu, S.-H., and Chuang, T.-Y. (2017). Speed and
603   temporal-distance adaptations during non-motorized treadmill walking in stroke and non-disabled
604   individuals. Eur J Phys Rehabil Med *53*, 863–869.

Figure 1. Non-motorized split-belt treadmill and paradigm. *A.* Non-motorized treadmill dimensions (inches), *left panel*; handrail height from top of treadmill deck and handle depth, and deck angle in degrees. Non-motorized treadmill dimensions, *right panel*; deck width and length and individual belt widths. *B.* Split-belt treadmill trasnfer paradigm. All subjects were recorded in baseline conditions for slow, medium, fast, and non-motorized treadmill (familiarization, not recorded and not shown) conditions in random order. The split-belt adaptation period was always immediately preceded by a second non-motorized treadmill baseline period used for non-motorized treadmill baseline measures. After all baseline periods were completed, all subjects adapted for 10 minutes with one limb on a belt traveling at the slow speed (0.67 m/s) and the other at the fast speed (1.34 m/s). Subjects then moved to the non-motorized treadmill without forward stepping to assess for the transfer to the non-motorized treadmill. Transfer was 5 minutes at preferred speed. Subjects then moved without forward stepping back to the motorized treadmill to assess for residual after-effects specific to the motorized treadmill for an additional 5 minutes. Finally, subjects then returned to the non-motorized once more to ensure all residual after-effects had been washout out for an additional 2 minutes.

Figure 2. Analytical calculation of step length. Visual representation of walking parameters for the analytical step length derivation. ***A.*** Walker demonstrating the position of the reflective markers on the fifth metatarsal, lateral malleolus, and lateral condyle. Gray crossed circle represents the pelvic centroid used to calculate the reference frame for ankle marker data on the motorized treadmill. Slow instantaneous step length shown (red arrow line) as distance between ankle markers at time of slow limb heel strike. Gray and black arrows indicate belt speeds. ***B.*** Slow step length visualized during early adaptation. Ankle marker trajectories in the pelvic centroid reference frame. Positive values indicate positions in front of the pelvic centroid and negative indicate positions behind the pelvic centroid. Dashed lines indicate individual ankle marker trajectories for the fast (black, x_f), and slow (gray, x_s) limbs. Ankle marker trajectories are plotted with time on the x-axis, therefore distance between dots indciates distance traveled every 10 milliseconds. Open circles indicate the position and time of the respective limbs heel strike. Filled circles indicate the position and time of the trailing limb ankle at the leading limbs heel strike. Solid vertical lines connecting open and closed circles indicate the instantaneous fast step length and the vertical dashed lines connecting open and closed circles indicate the instantaneous slow step length (i.e. SL_{slow}). The slow analytical step length (dashed red line) is the summation of the (negative) spatial component (α_s), and the product of the velocity of the fast limb (v_f) and the fast step time (t_f), the distance traveled by the fast limb from fast heel strike to fast position at slow heel strike. v_f is the difference between the fast limb ankle (x_f) at fast heel strike, and the fast limb ankle at slow limb heel strike, divided by the fast step time (t_f). ***C.*** Fast step length visualized dyring late adaptation. The fast analytical step length (solid red line) is the summation of the spatial component (α_f), and the product of the velocity of the slow limb (v_s) and the slow step time (t_s).

Figure 3. Adaptation; Progression of step length difference and contribution of spatial, temporal, and velocity asymmetries to motorized split-belt treadmill walking during adaptation. **A.** Group average step length difference during the first 100 and last 20 strides of the adaptation period. Shaded area represents plus/minus standard error. Negative values indicated the slow step length is larger than the fast step length. **B.** Individual subject step length difference data (small gray filled circles) comparing baseline (BA) slow walking to early adapt (AE) and late adapt (AL). Light gray lines connect individual subject data points between AE and AL. Group means are represented by large black circles, error bars are standard deviations. Thick black line connecting AE and AL indicates group change. Horizontal bars with star indicate significant differences between time points ($p < 0.05$). **C.** Stride-by-stride changes in individual components for the first 100 and last 20 strides; spatial (red), temporal (blue), and velocity (magenta) differences. Shaded areas are plus/minus standard error. **D.** Individual subject component data (gray filled colored circles) comparing AE and AL. Bars are group means, error bars are standard error. "*" indicates significant difference to baseline slow, "#" indicates late time point significantly different from early ($p < 0.05$).

Figure 4. Transfer; Progression of step length difference and contribution of spatial, temporal, and velocity asymmetries to non-motorized split-belt treadmill walking during transfer. *A.* Group average step length difference during the first 100 and last 20 strides of the transfer period. Shaded area represents plus/minus standard error. Positive values indicated the fast step length is larger than the slow step length. *B.* Individual subject step length difference data (small gray filled circles) comparing baseline (BA) walking to early transfer (TE) and late transfer (TL). Light gray lines connect individual subject data points between TE and TL. Group means are represented by large black circles, error bars are standard deviations. Thick black line connecting TE and TL indicates group change. Horizontal bars with star indicate significant differences between time points (p < 0.05). *C.* Stride-by-stride changes in individual components for the first 100 and last 20 strides; spatial (red), temporal (blue), and velocity (magenta) differences. Shaded areas are plus/minus standard error. *D.* Individual subject component data (gray filled colored circles) comparing TE and TL. Bars are group means, error bars are standard error. "*" indicates significant difference to baseline non-motorized, "#" indicates late time point significantly different from early (p < 0.05).

Figure 5. Washout; Progression of step length difference and contribution of spatial, temporal, and velocity asymmetries to motorized treadmill walking during washout. *A.* Group average step length difference during the first 100 and last 20 strides of the adaptation period. Shaded area represents plus/minus standard error. Positive values indicated the fast step length is larger than the slow step length. *B.* Individual subject step length difference data (small gray filled circles) comparing baseline (BA) slow walking to early washout (WE) and late washout (WL). Light gray lines connect individual subject data points between WE and WL. Group means are represented by large black circles, error bars are standard deviations. Thick black line connecting WE and WL indicates group change. Horizontal bars with star indicate significant differences between time points ($p < 0.05$). *C.* Stride-by-stride changes in individual components for the first 100 and last 20 strides; spatial (red), temporal (blue), and velocity (magenta) differences. Shaded areas are plus/minus standard error. *D.* Individual subject component data (gray filled colored circles) comparing WE and WL. Bars are group means, error bars are standard error. "*" indicates significant difference to baseline slow, "#" indicates late time point significantly different from early ($p < 0.05$).

Figure 6. Linear regression to assess interaction between transfer after-effect and washout after-effect magnitude. Data points are individual subject averages from first 5 strides during early transfer (TE) and washout (WE), respectively.

93
94
95

Figure 7. Group averaged muscle activity data during adaptation (first column), transfer (second column, gray), and washout (third column) for the VL (A, E, and I), BF (B, F, and J), MG (C, G, and K) and TA (D, H, and L). Left panels display group average muscle activity tracings during early (solid) and late (dashed line) time points across conditions compared to baseline speed and condition-matched values for each limb (gray shaded area) from -20 to 80% of the gait cycle. Vertical dashed line centered at zero indicates heel strike. Top panels in black are the fast limb with the lower panel in green for the slow limb. Dash-dot boxes within EMG traces indicate area of interest used in statistical analysis. Right panels within the columns are the group average difference curves from condition and speed-matched baseline values. Black dots represent fast limb during early (left) and late (right) time points, green dots are slow limb, and error bars are plus/minus standard deviation. "*" indicates significant difference to baseline, "#" indicates late time point significantly different from early (p < 0.05). Several subjects had corrupted EMG activity due to excessive movement artifact, therefore number of remaining subjects for each muscle is indicated with the "n =" in the upper right corner of the first column EMG trace.

Figure 8. Individual (small cirlces) and group average (large circles, ± sd) non-motorized treadmill belt speeds by limb. Fast limb instantaneous belt velocities are calculated as the instanteanous speed of the ankle marker in the global reference frame for the trailing limb at the instant of the leading limbs heel contact. Time points are during baseline non-motorized treadmill walking, early transfer, and early post-washout. Stars indicated significant difference (p < 0.05). Comparison between baseline and post are composite of both fast and slow limbs.

120