1 **Cryptogams signify key transition of bacteria and fungi in Arctic sand**

2 **dune succession**

3 Heli Juottonen[1*], Minna Männistö[2], Marja Tiirola[1] & Minna-Maarit Kytöviita[1]

4 [1]Department of Biological and Environmental Science, University of Jyväskylä, Finland

5 [2] Natural Resources Institute Finland (Luke), Rovaniemi, Finland

*Corresponding author. Heli Juottonen, Department of Biological and Environmental

Science, P.O. Box 35, 40014 University of Jyväskylä, Finland.

email: heli.juottonen@alumni.helsinki.fi, tel: +358 408053842

Running title: Cryptogams and microbes in Arctic dune succession

Summary

Microbes are the first colonizers of barren soils and an important factor in vegetation

establishment. Primary succession models focus on vascular plants, although most soil

successions include mosses and lichens. Here, we aim to clarify the role of these cryptogam-

dominated stages in plant-microbe interactions by covering the successional trajectory of

bacterial and fungal communities in eroded Arctic inland dunes. The earliest successional

stages with sparse or no plant cover (bare sand, grass) had low microbial biomass per organic

matter. In the cryptogam stages with the moss *Polytrichum piliferum* and lichens, microbial

biomass and community structure shifted towards the vascular vegetated stages (heath,

mountain birch forest). Soil chemistry explained microbial community variation better than

vegetation, but the transitions in chemistry and in plant and microbial community structure

did not align. We detected potential autotrophic and trace gas oxidizing bacteria, such as

WD272 (WPS-2), Chloroflexi and Cyanobacteria, which could form the basis of biomass

accumulation from bare sand to cryptogam stages. The results indicate that when considering

also belowground microbes, the ecosystem transition from vegetation loss to vegetation

recovery starts at the moss stage. *Polytrichum piliferum* and the associated microbes may

help stabilize the Arctic dune sand and facilitate vascular vegetation development.

Keywords: Arctic soil, bacterial community, cryptogam, fungal community, *Polytrichum*,

primary succession, sand dune

Introduction

Development of vegetation on bare land and the sequence of vegetative stages that
predictably follow each other have been described in several theories of succession (Egler,
1954, Wilson *et al.*, 1992). According to widely accepted views, primary succession is
influenced by site abiotic conditions (climate, soil parent material, topography) and biotic
factors (species pool present, order of arrival, species interactions). Despite the fact that
microbes are the first colonizers of any barren surface and shape soil chemistry, models
predicting successional trajectories of bacteria and fungi are few (Jackson, 2003, Fierer *et al.*,
2010, Dini-Andreote *et al.*, 2015, Tripathi *et al.*, 2018, Ortiz-Álvarez *et al.*, 2018). Most
models of terrestrial primary succession overlook bryophyte communities (e.g. Connell and
Slatyer, 1977, Baker and Walford, 1995), although most reports of primary succession
include moss and lichen dominated stages (Chapin *et al.*, 1994, Lichter, 1998, Hodkinson *et
al.*, 2003, Jones and Henry, 2003, Sparrius *et al.*, 2012).

Cryptogams, here mosses and lichens, may have diverse but unexplored effects on the
development of vascular plant and belowground microbial communities. Firstly, dispersal by
spores minimizes dispersal limitation – a feature classically included in models explaining
succession of vascular plant communities (Makoto and Wilson, 2019). The rapid arrival of
mosses warrants them a priority effect in establishing plant communities. Secondly, although
mosses and lichens lack roots and root-mediated effects on microbial communities, they
affect soil temperature, moisture, and nitrogen availability, which may promote or inhibit
vascular vegetation (Van der Wal and Brooker, 2004, Gornall *et al.*, 2007, Cornelissen *et al.*,
2007, Gornall *et al.*, 2011). Many lichens characteristic of primary succession fix nitrogen in
specific structures called cephalodia (Vitousek, 1994), whereas moss leaves contain
diazotrophs (DeLuca *et al.*, 2002, Arróniz-Crespo *et al.*, 2014). In addition to nitrogen,
rhizoids of mosses provide carbon to the belowground soil (Bowden, 1991). Thirdly, mosses
and lichens produce unique secondary metabolites that may have adverse effects on vascular
plants and beneficial or adverse effects on microbes (Cornelissen *et al.*, 2007, Xie and Lou,
2009).

Microbial community shifts in early succession are considered to be driven by soil chemistry,

external resources and dispersal of microbes, and in later stages by biotic factors, such as

establishment and changes in plant community (Jiang *et al.*, 2018, Brown and Jumpponen,

2014). During succession microbes modify soil chemistry, and biotic interactions including

plant-microbe interactions can aid plant establishment. Bacterial versus fungal communities

can be expected to show distinct trajectories in primary succession: Bacteria are smaller and

believed to disperse easier than fungi, which may lead to more deterministic community

assembly for bacteria (Schmidt *et al.*, 2014, Powell *et al.*, 2015) and higher influence of

priority effects and stochastic effects for fungi (Brown, S. P. and Jumpponen, 2014, Schmidt

et al., 2014, Jiang *et al.*, 2018). Further, the physiological diversity in bacteria enables some

of them to survive as autotrophs in bare oligotrophic soils of early successional stages in the

absence of plants fixing carbon (Nemergut *et al.*, 2007, Schmidt *et al.*, 2008, Duc *et al.*,

2009). Fungi in these habitats depend on already fixed carbon and nitrogen from wind-blown,

ancient, or microbially fixed sources (Schmidt *et al.*, 2014). The critical shift to ecosystem

based on production by resident plants is associated with changes in microbial community

(Bardgett and Walker, 2004, Edwards *et al.*, 2006, Blaalid *et al.*, 2012, Knelman *et al.*, 2012).

Along succession, fungi to bacteria ratio generally increases as soil organic matter increases

and pH decreases (Pennanen *et al.*, 2001, Tscherko *et al.*, 2004, Bardgett and Walker, 2004,

Bardgett *et al.*, 2007, Baldrian *et al.*, 2008). These results do not, however, identify the only

factors responsible for vascular plant establishment. As early colonizers, cryptogams with

their influence on soil chemical and physical properties and biotic interactions could be an

additional key factor in the ecosystem shift from early to late succession.

In the world where ecosystems are declining due to soil erosion and loss of plant cover

(Montgomery, 2007, Vanwalleghem *et al.*, 2017), it is critical to identify the key shifts

between non-vegetated and vegetated stages in ecosystem succession. Identifying the

microbial and possibly plant communities that signal for the shift from autotrophic to

allotrophic helps identify the key factors that drive the shift back to vascular plant cover, soil

stabilization and a productive ecosystem. Arctic inland sand dunes are an example of areas

where vegetation has been lost due to environmental changes and lack of recovery. The dunes

have formed from wind-blown sand after the last ice age (Koster, 1988). Plants colonized the

dunes later on, but today large areas lack vascular plant cover. Forest fires, insect outbreaks,

and overgrazing may have caused the ecosystem degradation (Seppälä, 1995). In our study

site in Northern Finland, wind erosion has carved deflation basins into the dunes. Inside the

basin, primary succession vegetation stages have developed, and dune slopes surrounding the

basins are covered by mountain birch forest, the climax stage in the present climate. Eroding

inland sand dunes provide an excellent opportunity to study primary succession due to the

mosaic of closely situated successional stages within deflation basins. Previous work

comparing the start and end points of succession proposed that Arctic sand dune bacteria and

fungi may show distinct successional patterns (Poosakkannu *et al.*, 2017).

In this study, we compared two cryptogam-dominated stages with non-vegetated and vascular

plant covered stages in an eroded Arctic dune ecosystem. We hypothesized that the

cryptogam-covered stages contain soil microbial community more similar to stages covered

by vascular plants and thereby drive succession towards vascular plant dominance through

microbially mediated facilitation. Secondly, we compared the belowground responses of two

microbial groups - fungi and bacteria - to successional stages defined by aboveground

characteristics. We hypothesized that bacteria would be more prominent and diverse in the

early successional stages and fungi in the later stages with vascular vegetation. By linking the

aboveground vegetation, the belowground microbial communities, and the associated soil

chemical environment, we aimed to define the role of cryptogam stages in the Arctic dune

ecosystem succession.

Materials and methods

Study site and sampling

The study site is located in an inland aeolian (i.e. wind-deposited) dune area in Northern

Fennoscandia (68° 29′ 16″ N, 24° 42′ 13″ E). The 1981–2000 average annual temperature in

the region was -1.3 °C, with extreme air temperatures between +30 °C and -52 °C and the

average annual precipitation 550 mm (Pirinen *et al.*, 2012). We defined six successional

stages in the dunes based on vegetation (Fig. S1). Although the soil in all the stages was

aeolian sand, in three the largest part of the surface was mostly exposed sand (92-100 %),

while in the other three vegetation covered the sand (0-52 % sand). The sand-exposed 'sandy

stages' were: 1) sand: bare loose sand without plants or visible soil crust, 2) grass: bare loose

sand with the grass *Deschampsia flexuosa* and without visible soil crust, and 3) moss: bare

loose sand with the moss *Polytrichum piliferum*. The stages with extensive vegetation cover

('vegetated stages') were 4) lichen: the main species lichens *Stereocaulon* spp. and *Cladonia*

spp., and mosses *P. piliferum* and *Racomitrium ericoides*, 5) heath: the dominant species

dwarf shrub crowberry *Empetrum nigrum* with *Cladonia* spp. and/or *Stereocaulon* spp., and

6) forest: mountain birch (*Betula pubescens* subsp. czerepanovii) forest, the main species in

study plots the shrubs *E. nigrum* and *Vaccinium vitis-idaea*,the moss *Pleurozium schreberi*,

D. flexuosa, and the herb *Linnaea borealis*. The sand, grass, moss, and lichen stages are

located at the bottom or on the inner slopes of the deflation basins and heath at the edge of

the basins. The remnant mountain birch forests are located around the basins at the crest of

the dune or on the outer slope. Permanent study plots of 1×1 m were established in 2008 and

included each of the six vegetation stages in seven deflation basins. The vegetation in the

plots was recorded in detail in 2009 by point frequency method (100 points within 50×50

cm, presence/absence and frequency of species). The plots have been photographed every

year from 2010 to 2018 and the vegetation has not changed notably.

We sampled soil in each successional stage in four of the basins in July 2011. The soil

samples were taken in triplicate from the depth of 0-10 cm with a soil corer (72 samples in

total: six successional stages $\times$ four basins $\times$ three replicates). Soil was sieved with a 4-mm

sieve and frozen at -20°C.

Soil chemistry and temperature

Total C and N of soil organic matter were analysed on a LECO CHN 1000 elemental analyser

(LECO Corporation, USA). For analyzing soil P, Ca, K, Na, Fe, Mn, Al, and Mg contents,

soils were extracted with acid (pH 4.65) 1 M ammonium acetate (15 ml soil sample /150 ml

extraction solution) and the elements were determined with an ICP emission spectrometer

iCAP 6500 DUO 2011 (Thermo Scientific, United Kingdom). The dry matter content of the

soil was determined by drying the samples (105 ºC) and organic matter (OM) content

analyzed by loss on ignition (550 ºC) using LECO TGA-701 analyzer (LECO Corporation,

USA). pH was determined in 1:2 (w:v) soil:water suspensions (10 g soil and 20 ml H2O).

Soil temperature at the 10 cm depth was recorded every two hours with Hobo Temp External

H08-002-02 loggers coupled with TMC6-HD soil temperature sensor during 2009-2015. We

defined maximum temperature as the average temperature of the warmest month (July) and

the minimum temperature as the average of the two coldest months (January and February).

Phospholipic fatty acid (PLFA) analysis

Phospholipid fatty acids were extracted as described previously (Francini *et al.*, 2014) from

freeze-dried soil (15 g for sand and grass stages, 10 g for moss, 8 g for lichen, 5 g for heath

and 3 g for forest stage corresponding to OM content). Briefly, lipids were extracted with

chloroform:methanol:citrate buffer (Bligh and Dyer, 1:2:0.8 v/v/v), fractionated using silica

columns (Bond Elut LRC, Varian), and phospholipids were methylated with mild alkaline

methanolysis. The resulting fatty acid methyl esters were analyzed by gas chromatography

and PLFAs identified based on a bacterial fatty acid standard mixture (Supelco, Bellefonte,

PA, USA) and fatty acids from reference strains. For community analysis, data was expressed

as peak area percentage of a single PLFA of the total area of the profile, and for biomass

analysis as nmol of PLFAs per gram dry weight (gdw) of soil or per gram OM. The sum of

PLFAs i15:0, a15:0, 15:0, i16:0, 16:1ω9, i17:0, a17:0/17:1ω8c (eluted in the same peak),

17:0, cyclo-17:0, 18:1ω7c/ω9t (eluted in the same peak) and cyclo-19:0 were used to

represent bacterial biomass and PLFA 18:2ω6 fungal biomass. Sum of all these PLFAs was

used to represent total microbial biomass. Biomass of selected bacterial groups was

calculated by multiplying the total microbial biomass of a sample with the relative abundance

of the bacterial group in the sample based on 16S rRNA gene sequencing.

DNA extraction and PCR

DNA was extracted from 0.85 g (0.4 g for forest stage) of soil with Powersoil DNA

extraction kit (MoBio). Bead beating was carried out with a FastPrep instrument at 5.0 m s^{-1}

for 40 s. DNA concentration was determined with Qubit fluorometer and Qubit dsDNA HS

kit (Invitrogen).

Bacterial and fungal PCR was carried out as two-step PCR where the second step introduced

barcodes and adapters for Ion Torrent sequencing. The first step of bacterial 16S rRNA gene

PCR (V1-V2 region) was carried out with primers 27F (5'-AGAGTTTGATCCTGGCTCAG-

3') and 338R (5'-TGCTGCCTCCCGTAGGAGT-3'). The 25-µl reactions were carried out in

duplicate and contained $1 \times$ buffer, 200 µM dNTPs, 0.4 µM each primer, 0.4µg µl-1 BSA,

0.75 U DreamTaq DNA polymerase (Thermo Fisher Scientific), and 1-2 µl of template DNA

(depending on concentration). The cycling conditions were initial denaturation at 95 °C for 3

min, followed by 30 cycles of 95 °C for 30 s, 53 °C for 30 s, and 72 °C for 45 s, and a final

extension of 72 °C for 7 min. The first step of intergenic transcribed region (ITS) PCR for

fungi was carried out with primers fITS7 (5'-GTGARTCATCGAATCTTTG-3') and ITS4

(5'-TCCTCCGCTTATTGATATGC-3') (Ihrmark *et al.*, 2012), where the fITS7 primer

contained an M13 adapter for the second step (Mäki *et al.*, 2016). Reaction composition was

the same as in bacterial PCR but including additional $MgCl_2$ to total concentration of 3 mM.

The cycling conditions were initial denaturation at 94 °C for 3 min, followed by 30 cycles of

94 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s, and a final extension of 72 °C for 7 min.

The second step primers introduced barcodes and Ion Torrent adapters A and P1. For

bacterial 2nd step PCR, the forward primer was adapter A-barcode-27f, and the reverse

primer adapter P1-338R. For the fungal 2nd step PCR, the forward primer was adapter A-

barcode-M13 adapter and the reverse primer adapter P1-ITS4. Second step PCR reactions (25

µl) for both bacteria and fungi contained $1 \times$ buffer, 200 µM dNTPs, 0.3 µM each primer, 0.5

U DreamTaq DNA polymerase, and 1 µl (bacteria) or 2 µl (fungi) of 1:10-diluted first step

PCR product as template. The cycling conditions were as in the first step PCR but with 7

cycles. After the second PCR, the duplicate products were pooled, purified with Agencourt

AMPure XP purification system (Beckman Coulter), and quantified with Qubit fluorometer

and Qubit dsDNA HS kit. Equimolar amounts (10 ng) of products were pooled for

sequencing. The fungal product pool was size-separated in precast 1% Pippin Prep gel (Sage

Science) to remove oversized products > 400 bp, which occurred in some of the samples.

These products were cloned and sequenced to confirm they were not of fungal origin (data

not shown).

Ion Torrent sequencing and sequence analysis

Pooled bacterial and fungal PCR products were sequenced after emulsion PCR with the Ion

OneTouch system and Ion OT2 400 kit (Life Technologies) on Ion 314 chips with Ion PGM

Sequencing 400 Kit (Life Technologies) according to the manufacturer's instructions. The

sequences were processed in mothur software package v. 1.38 (Schloss *et al.*, 2009)

following the relevant parts of MiSeq SOP protocol (https://mothur.org/wiki/MiSeq_SOP;

accessed in January 2017; (Kozich *et al.*, 2013). Sequences were quality-filtered using

average quality of 25 and a window size of 50 bases, a minimum sequence length of 200 bp,

a maximum length of 400 bp for bacteria and 410 bp for fungi, and further settings

maxhomop=6, maxambig=0, pdiffs=1, and bdiffs=0. Bacterial sequences were aligned

against Silva database v. 1.23 (Quast *et al.*, 2013). Chimeras were detected with Uchime

(Edgar *et al.*, 2011) in mothur. Fungal ITS sequences were identified, ITS2 region was

extracted and non-fungal sequences were removed with ITSx (Bengtsson-Palme *et al.*, 2013).

After quality filtering, alignment for bacteria, removal of chimeras and non-target sequences,

we had 121711 bacterial sequence reads and 279885 fungal reads. Reads were preclustered

with setting diffs=2 (bacteria) or diffs=1 (fungi). The sequences were clustered into 97%

operational taxonomic units (OTUs) using the average neighbor method for bacteria, and the

vsearch algorithm for fungi, yielding 5242 bacterial and 1742 fungal OTUs after removal of

singletons. Bacterial OTUs were classified against the Silva v. 1.23 database and fungi

against the Unite database (Kõljalg *et al.*, 2013). Data was subsampled to the read number of

the sample with the lowest number of reads, which was 1055 reads for bacteria and 1605

reads for fungi. Shannon diversity index and coverage were calculated in mothur. Coverage

was $76 \pm 6\%$ for bacteria and $97 \pm 1\%$ for fungi. The sequence data were submitted to NCBI

under BioProject accession PRJNA471306.

Statistical analyses

Chemistry and microbial biomass were compared with the successional stage using linear

mixed models with R package nlme with successional stage as a fixed factor and deflation

basin as a random blocking factor. Chemistry and biomass values except pH and Al were log

transformed.

Bacterial and fungal sequence data were converted to relative abundances of OTUs and

square root transformed. Only OTUs with 5 ≥ reads were included. Variation of community

structure was visualized as non-metric multidimensional scaling (NMDS) plots with function

metaMDS using Bray-Curtis distance measure. Multivariate dispersion between the

successional stages was tested with betadisper (Bray-Curtis distances). Permutational analysis

of variance (PERMANOVA, (Anderson, 2001, McArdle and Anderson, 2001) with adonis

(Bray-Curtis distances) was used to compare community structure between the successional

stages and with environmental variables (OM, pH, and in addition Al, Na and minimum and

maximum temperature which did not correlate with OM and pH but differed with

successional stage). We used Mantel test to estimate the correlation between microbial

community structure and soil chemistry (Euclidean distances) and microbial community

structure (Bray-Curtis distances) and vegetation (Bray-Curtis distances of square root

transformed data). All multivariate analyses were carried out with the vegan package (v.2.4.2,

(Oksanen *et al.*, 2017) in R. Significance tests were based on 999 permutations.

We identified bacterial and fungal OTUs characteristic to specific successional stages or

combinations of stages by indicator species analysis with package indicspecies using function

multipatt and correction for unequal site numbers (54). Only OTUs that occurred in the

subsampled data at least 12 times, i.e., once in each sample of a successional stage, were

included in the analyses. The combinations of stages included in the analyses were each

successional stage on its own, all six stages together, all stages except sand, all stages except

forest, all stages except sand and grass, all stages except heath and forest, sand+grass,

moss+lichen, moss+grass, lichen+heath, heath+forest, sand+grass+moss, moss+lichen+heath,

and lichen+heath+forest.(De Cáceres and Legendre, 2009). Instead of using all possible

combinations, we selected the combinations to limit the number of tests and to focus on the

combinations most relevant for the aims of the study. OTUs were considered indicator OTUs

if their indicator value was >0.3 and p <0.05. Significance was evaluated with a permutation

test (999 permutations). P values were adjusted for multiple comparisons with function

p.adjust (method "fdr"). Generalist OTUs present in all successional stages were defined at

indicator values >0.5, because no p values are generated for taxa present in all sites. The

identified indicator OTUs were grouped based on their taxonomic classification at order

(bacteria) or class (fungi) level. Unclassified OTUs were included in these comparisons by

re-clustering all reads assigned to unclassified bacteria or fungi in mothur at 75% sequence

similarity to represent phylum level similarity for bacteria and class level similarity for fungi
(Tedersoo *et al.*, 2014).

To evaluate responses of bacterial and fungal indicator taxa to environmental variables, we
used relative abundances of indicator OTUs grouped at order level (bacteria) and class level
(fungi), including those unclassified 75% similarity groups that represented >0.4% of all
unclassified OTUs. The tested environmental variables were those that best explained
microbial community variation (OM, pH, Al for bacteria; OM, pH, minimum temperature for
fungi). The analysis was carried out with the function traitglm in the package mvabund (v.
3.13.1, (Wang *et al.*, 2012) and function traitglm. The function fits fourth corner models to
relate sites and their traits to species and species traits. When used without species trait data,
it becomes a predictive model of responses of different OTUs to variation of environmental
factors (57). The method used was glm1path, fitting a generalized linear model with LASSO
penalty, which means that fourth corner efficients of OTUs not decreasing Bayesian
information criterion (BIC) are forced to 0, i.e. no response to environmental variation, to
simplify the model (Brown, A. M. *et al.*, 2014) (see supplementary materials for details).

All statistical analyses were carried out in R v.3.2.5. or v.3.4.0 (RCoreTeam, 2018).

Results

Soil chemistry and microbial biomass along Arctic sand dune succession

The increasing aboveground vegetation cover from bare sand to mountain birch forest was
mirrored belowground in increasing soil OM content and decreasing pH (Pearson correlation
r=-0.78, p< 0.001, Fig. 1). Organic matter content was low in the sand, grass, and moss stage
soils (0.4±0.2%, mean±SD) but increased to 0.9±0.3% in the lichen stage and up to 4.5% ±
1.7% in the forest stage (Fig. 1A, Table S1). pH was highest in the grass stage (5.8 ± 0.2) and
lowest in the forest stage (4.7 ± 0.1) (Fig. 1B). Soil contents of N, P, K, Ca, Mg, Fe and S
correlated with OM (Table S1).

Microbial biomass measured as PLFAs per soil dry weight followed closely the OM gradient

(Fig. 1D). The moss stage had similar OM content as bare sand, but its microbial biomass

was higher than in the other sandy stages. When related to soil OM, microbial biomass in the

moss stage reached a level similar to the lichen and vascular plant stages (Fig. 1C). Similarly,

a shift to a higher proportion of fungal to bacterial biomass occurred in the moss stage (Fig.

1E). Overall, the proportion of fungi was highest in the heath stage.

Microbial community structure

Bacterial and fungal community structure based on sequencing (Fig. 2) and PLFA (Fig. S2)

followed the order of the vegetation stages on the OM-pH gradient and differed with

successional stage (PERMANOVA bacteria $r^2=0.40$ P<0.001, fungi $r^2=0.38$ p<0.001). The

moss stage grouped with the other sandy stages (sand, grass) with partly overlapping

communities, whereas the lichen stage grouped with the vascular plant stages (heath, forest)

(Fig. 2). It should be noted that multivariate dispersion also differed between the successional

stages (betadisper p=0.007 bacteria, p=0.001 fungi). The highest bacterial OTU richness (Fig.

1F) and Shannon diversity (Table S1) were detected in the heath stage, whereas fungal

richness and diversity were highest though very variable in bare sand (Fig. 1F, Table S1).

The largest shift in bacterial community structure was detected from moss to lichen stage

(Fig. 3). In contrast, the extent of fungal community change from stage to stage remained

rather constant. Plant community composition showed increasingly larger shifts with

succession, with the largest shift from lichen to heath. Soil chemistry changed most markedly

from heath to forest. Overall, microbial community structure correlated better with soil

chemistry than with plant species composition (Mantel test; bacteria chemistry r=0.71 vs.

vegetation r=0.53; fungi r=0.40 vs. r=0.25, p=0.001 for all). Community variation of bacteria

and fungi was best explained by OM content (Table 1), especially in the vegetated stages,

which comprised the largest shifts in OM content. In the sandy stages, pH, OM, and Al

content explained the highest share of bacterial and fungal variation.

Bacterial taxonomic distribution with succession

Overall, the most abundant bacterial phyla or classes were Alphaproteobacteria,

Actinobacteria, and Acidobacteria (Fig. 4A, Fig. S3). The most abundant bacterial OTU

belonged to the candidate phylum WD272 (WPS-2). This phylum was the fourth or fifth most

abundant phylum in the sandy and lichen stages (on average 4.6-8.1%), decreasing in the

heath (3.5%) and forest (0.9%) (Fig. 4A, Fig. S3). When related to microbial biomass, the

abundance of WD272 increased in the moss stage (Fig. 5). WD272 showed abundant

indicator OTUs for the sandy stages but also less abundant forest-specific indicators, which

possibly explains why WD272 indicators could not be related to environmental variables

(Fig. 6). On the other hand, most indicator groups specific for the sandy stages, such as

Cytophagales, Deinococcales, Acidobacteria subgroup 4, Pseudonocardiales (Actinobacteria),

Oligoflexales (Deltaproteobacteria) and unclassified group 9 were strongly associated with

low OM content. Cyanobacteria, three Chloroflexi groups and unclassified group 2 were

indicators for the moss and lichen stages and associated with both low OM and low pH,

which reflected their occurrence at the pH shift in the succession. Another sign that the moss

soil was a bacterial transition stage was that it showed unclassified clusters typical to both

sandy and vegetated stages (Fig. S5A). Most indicator groups for the forest stage were

indicators also for the other vegetated stages. For example, Acidobacteria subgroup 2

associated with low pH and subgroup 6 associated with high OM appeared in the moss and

lichen stages and had indicators also in the heath and forest stages. Other bacterial indicators

in the vegetated stages associated with low pH, high OM or both included Planctomycetes

(WD2101 soil group), Frankiales and Streptomycetales (Actinobacteria), and

Zanthomonadales (Gammaproteobacteria). Acidobacteriales (Acidobacteria) and

Corynebacteriales (Actinobacteria) occurred throughout the gradient but increased in relative

abundance towards the forest and were associated with low pH. The overall abundance of

Acidobacteria increased with decreasing pH (Pearson correlation r = -0.81, p<0.001).

Fungal taxonomic distribution with succession

The fungi at class level were dominated by Ascomycetes, followed by Basidiomycetes,

unclassified fungi, and smaller shares of Zygomycota, Rozellomycota, Chytridiomycota, and

Glomeromycota (Fig. 4B, Fig. S4). The most abundant Ascomycota class was Leotiomycetes,

followed by unclassified ascomycetes and Dothideomycetes. Bare sand had the highest

relative abundance of unclassified fungi up to 35% (Fig. 4B, Fig. S4), and several

unclassified clusters were specific to the sandy stages (Fig. S5B) and associated with high pH

or low OM (Fig. 7). Other strong indicators of sandy stages included Helotiales and

Thelebolales (Leotiomycetes), and Polysporales and Boletales (Agaricomycetes).

Lecanoromycetes, containing lichenized fungi, was most abundant in the moss stage (Fig.

4B). Similarly to bacteria, the fungal groups that were the best indicators of the moss and

lichen stages were associated with low OM and low pH or both, including Lecanoromycetes

(Lecanorales, Agyriales, Ostropomycetidae), Agaricomycetes (Cantharellales, Atheliales,

Agaricales), and Dothideomycetes (Pleosporales, Capnodiales). The most abundant fungal

OTU overall and one of the fungal generalists (Table S2) was classified as *Pezoloma ericae*

(Leotiomycetes; also known as *Rhizoscyphus ericae*) known to form mycorrhiza with ericoid

plants such as the dominant heath species *Empetrum nigrum* (Read, 1991, Smith and Read,

2008). Accordingly, this OTU was particularly dominant in the heath (34±5% of reads). Its

relative abundance was lowest in bare sand (2±1%) and the grass stage (12±14%), but in the

moss and lichen stages it increased to 26±16% despite the lack of known host plants. In the

forest, its relative abundance was lower than in the cryptogam-dominated stages (17±14%).

Venturiales (Dothideomycetes) was an indicator of the heath stage. More general indicator

groups for the vegetated stages associated with low pH, high OM or both were

Archaeorhizomycetes, Mortierellales (Zygomycotina), Rozellomycota, Russulales,

Sebacinales and Thelophorales (Agaricomycetes), and unclassified groups 1 and 11.

Hysteriales (Dothideomycetes) and Geoglossales (Geoclossomycetes) showed high OM

associated indicator OTUs exclusive for the forest stage.

Discussion

We hypothesized that the cryptogam-dominated stages of an Arctic sand dune succession

represent key ecosystem transitions towards vascular vegetation. Supporting this hypothesis,

the bacterial and fungal communities in the lichen stage resembled the communities in the

later vascular plant dominated stages. Although the moss stage microbial communities

overlapped with the earlier sandy stages, the shift towards vascular plant covered soils started

with the appearance of moss: Microbial biomass per OM and the proportion of fungi

increased in the moss stage, and later stage microbial groups appeared despite absence of

their presumed host plants.

The dune sand is a harsh habitat with low carbon and nitrogen availability. The initial

microbial colonizers must be able to tolerate the oligotrophic conditions either by using the

low levels of allocthonous organic matter (Hodkinson *et al.*, 2002) or as autotrophs

(Nemergut *et al.*, 2007, Fierer *et al.*, 2010). Carbon fixed by these initial colonizers then

allows the establishment of the first heterotrophs. However, comparison of biomass per OM

with biomass per soil dry weight (Fig. 1) shows that the microbial biomass increased in the

moss stage when OM potentially available for heterotrophs still remained low. A

characteristic group for the early stages from bare sand to lichen was candidate phylum

WD272, currently known as WPS-2 and recently named *Candidatus* Eremiobacteraeota,

'desert bacterial phylum' (Ji *et al.*, 2017). Studies on mosses, volcanic ice caves and

Antarctic desert soils have proposed members of WD272 to be autotrophs fixing carbon with

ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCo; (Tebo *et al.*, 2015, Ji *et al.*,

2017, Holland-Moritz *et al.*, 2018). Based on metagenome-assembled genomes, WD272

members have been proposed to use atmospheric H_2 and carbon monoxide (CO) as energy

sources (Ji *et al.*, 2017) or to be anoxygenic photoautotrophs (Holland-Moritz *et al.*, 2018).

As another potential trace gas oxidizing group, Chloroflexi in the moss and lichen stages

were classified as Ktedonobacteria, a class with several CO-oxidizers (King and King, 2014).

Many Chloroflexi can use H_2 and CO to persist under low nutrient conditions (Islam *et al.*,

2019). The potential of WD272 and Chloroflexi to survive in the harsh early stages as

autotrophs or trace gas oxidizers and their increased biomass in the moss and lichen stages

(Fig. 5) leads us to speculate that they form a basis of biomass accumulation in the Arctic

sand dune succession. These microbes may therefore facilitate cryptogam cover development

and further vascular vegetation development (Connell and Slatyer, 1977, Brooker *et al.*,

2008).

Many moss stage microbial groups that were rare or absent in the earlier sandy stages are

linked to biological soil crusts. Biocrusts are consortia of microbes, mosses, and lichens that

are recognized as the major stabilizing factor in early successional soils worldwide (Weber *et*

al., 2016). The rhizoids of *Polytrichum* mosses themselves stabilize soil, facilitate N_2

fixation, and provide OM to the soil (Bowden, 1991). The moss stage showed the highest

proportion of Cyanobacteria, N_2-fixing early colonizers and components of biocrusts (Weber

et al., 2016), and of Lecanoromycetes (lichenized fungi, Ascomycetes). Lichens were

therefore detected in the belowground soil before they were detected on the soil surface.

Chloroflexi were distinctive indicators for the cryptogam stages, and in dryland and grassland

ecosystems they occur in moss and lichen biocrusts and in sub-crust soil (Kuske *et al.*, 2012,

Navarro-Noya *et al.*, 2014, Maier *et al.*, 2018). Although *Polytrichum* biomass was small in

the moss stage, *Polytrichum piliferum* appeared to have a disproportionate effect at the

ecosystem level, suggesting it is a keystone plant species in Arctic sand dunes. The

mechanism could be soil stabilization by rhizoids and moss-associated, soil crust-forming

microbes. Traditionally, grasses are credited with the initial physical stabilization of sand in

dune succession (e.g. (Olson, 1958, Ellenberg, 2009). Inspecting soil microbial community

structure, it seems that in these Arctic dunes the grass *Deschampsia flexuosa* may not drive

the belowground ecosystem succession. Based on the results, we speculate that *Polytrichum*

mosses could participate in the physical stabilization of Arctic inland dunes and direct

succession towards vascular plant cover. Verifying this proposition requires further studies on

dune *Polytrichum* and its rhizoids.

The relative abundance of the dominant fungal OTU *Pezoloma ericae* increased in the moss

stage to levels similar to those in the later stages despite no host plants detected. In addition

to ericoid mycorrhiza, *P. ericae* forms mycorrhiza-like associations in liverwort rhizoids

(Pressel *et al.*, 2010, Kowal *et al.*, 2018), and it has been found as an Antarctic moss

endophyte (Zhang, T. *et al.*, 2013). However, no liverworts were visible among the moss at

our site, and no association of *P. ericae* with *Polytrichum* rhizoids is known. Nevertheless,

we interpret the increase of *P. ericae* in the moss stage as a possible sign of fungal

community shift towards a stage where vascular plant establishment is possible. An

alternative explanation would be that propagules of *P. ericae* accumulate as the early

succession proceeds.

We hypothesized that bacteria would be more prominent and diverse in early succession and

fungi in late succession but found no clear support for this. The proportion of fungi in PLFA

data, assumed to represent living biomass (Zhang, Y. *et al.*, 2019), increased already in the

moss stage. Bacterial richness peaked in the heath and fungal richness in the bare sand, and

neither correlated with ecosystem productivity defined as plant richness or soil OM. In

general, richness and diversity link to ecosystem functions through the general effects of

diversity on ecosystem stability and productivity (Loreau and de Mazancourt, 2013). Plant

richness peaks in the middle or late succession (Bonet and Pausas, 2004, Peyrat and Fichtner,

2011, Prach *et al.*, 2014). For microbes, few general patterns of richness or diversity with

primary succession have emerged. In very young soils, richness is often low and then

increases and stabilizes or even decreases in later stages (Jackson, 2003, Nemergut *et al.*,

2007, Fierer *et al.*, 2010, Williams *et al.*, 2013, Jiang *et al.*, 2018). However, primary

succession gradients where both bacteria and fungi have been studied show conflicting

patterns. In glacier forefield successions, bacterial richness has increased or remained stable

and fungal richness remained stable through succession (Brown, S. P. and Jumpponen, 2014,

Jiang *et al.*, 2018). In a lava flow succession, fungal richness increased but bacterial richness

was not related to terrain age (Cutler *et al.*, 2014). In a salt marsh chronosequence, fungal

richness was not consistently stable with soil age due to a dominant OTU at one stage,

whereas bacterial richness first remained stable and then decreased (Dini-Andreote *et al.*,

2014, Dini-Andreote *et al.*, 2016). Our results are best in line with studies showing stable

microbial richness through primary succession, as shown for fungi in a dune succession

(Roy-Bolduc *et al.*, 2015).

The low bacterial richness in the bare sand stage can be related to the oligotrophic conditions;

richness already increased with the scattered presence of the grass *D. flexuosa*. Pioneering

plants can influence bulk soil when plants are of sufficient size (Tscherko *et al.*, 2004,

Edwards *et al.*, 2006, Miniaci *et al.*, 2007). The high fungal richness in bare sand together

with the low fungal biomass may represent dormant fungi, spores or hostless biotrophic fungi

(Jumpponen, 2003, Rime *et al.*, 2015). On the other hand, our observation that the

unclassified fungi can be grouped into class level clusters that occurred consistently in

replicate samples and across the sandy successional stages also hints that the fungal

community in bare sand may be in one way or other selected.

The high bacterial richness in the heath coincided with the highest fungal biomass and

highest occurrence of *P. ericae*, presumably forming mycorrhizae with the dominant ericoid

E. nigrum. We speculate that the high bacterial richness could be promoted by resources and

niches provided by fungal mycelium and *Empetrum* litter, despite its high phenolic content

and proposed adverse properties on microbial activity (Gallet *et al.*, 1999, Wardle *et al.*,

1998). In the forest, where the highest microbial richness would be expected if richness

correlated with OM, plant species richness and productivity, bacterial diversity was lower

than in the heath, possibly due to the environmental filtering effect of the low pH in the forest

soil (Tripathi *et al.*, 2018). The varied responses of bacterial and fungal richness and diversity

to primary succession highlight the need to cover successional gradients with different parent

material, vegetation and soil chemistry to understand microbial community patterns with

succession.

Both bacterial and fungal community variation were better explained by soil chemistry than

by plant community composition. Effects of soil chemistry on microbial communities can

override biotic factors such as plant richness (Chen *et al.*, 2016, Jiang *et al.*, 2018), and the

role of vegetation is emphasized in successional gradients where soil chemistry does not

change markedly (Knelman *et al.*, 2012, Cutler *et al.*, 2014). Soil chemistry and vegetation

are strongly intertwined in succession, and the present gradient from bare sand to mountain

birch forest represents the common pattern where organic matter increases and the soil

acidifies with increasing plant cover (Odum, 1969, Merilä *et al.*, 2010). However, the largest

microbial community shifts did not align with the largest shifts in soil chemistry or vegetation

(Fig. 3). No measured soil chemistry variable changed markedly with the large bacterial

community shift between moss and lichen stages (Fig. 3). Both cryptogam stages showed

indicator microbial groups associated with low OM and low pH. These results suggest that

the microbial shift from moss to lichen is due to unmeasured effects of the lichen crust

formation. Such factors could be better long-term moisture retention by the lichen crust than

the sparse mosses in the moss stage, microenvironments provided by lichen crumbs in the

belowground soil (de los Ríos *et al.*, 2011) or the different chemical composition and

secondary metabolites of mosses versus lichens.

Conclusions

This study on inland sand dunes illuminates bacterial and fungal succession in a less studied

Arctic soil ecosystem. Our results highlight the importance of considering not only vascular

plants but also cryptogams, i.e. mosses and lichens and particularly the potential dune key

stone species *Polytrichum piliferum*, when defining the plant-microbe interactions in

succession. Taken together, the results indicate that the microbiological transition towards the

later stages with vascular vegetation and soil stabilization starts at the moss stage. The

biomass representing early stage microbes increased and soil crust-associated microbes and

later stage fungi appeared, although the plant cover and OM were still low. Whether the sand

stage microbes initiate the transition by facilitating the appearance of moss or whether the

sand microbes linger and proliferate in the soil after the moss has appeared should be

determined in future experiments. This work adds to evidence that bacterial and fungal

richness are not necessarily related to soil ecosystem productivity or successional change in

plant species richness (Waldrop *et al.*, 2006, Porazinska *et al.*, 2018, Delgado-Baquerizo and

Eldridge, 2019). Furthermore, soil chemistry and the effects of plants on microbial resources

and niches are important factors driving succession. However, the key transitions in soil

chemistry, plant community and microbial community did not align, suggesting additional

unrecognized drivers. Bacterial and fungal indicators identified in this work, including novel

unclassified diversity, can signpost ecosystem transitions towards vascular plant cover.

Understanding the drivers of the shift from bare to vegetated is crucial for restoration of

vegetation on cold-climate soils suffering from erosion and vegetation loss under changing

environment.

Acknowledgements

550 We thank Helena Jauhiainen, Riitta Nissinen and Manoj Kumar for help with sampling and

551 DNA extraction, Anita Mäki for assistance with sequencing, and Outi Manninen for the

552 vegetation inventory. This work was funded by Maj and Tor Nessling Foundation and the

553 Academy of Finland (project 287545 to M.-M.K.).

References

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecol. **26**: 32-46.

Arróniz-Crespo M, Pérez-Ortega S, De los Ríos A, Green TA, Ochoa-Hueso R,

Casermeiro MÁ, de la Cruz, María Teresa, Pintado A, Palacios D, and Rozzi R. 2014.

Bryophyte-cyanobacteria associations during primary succession in recently deglaciated areas

of Tierra del Fuego (Chile). *Plos One* **9**: e96081.

Baker WL, and Walford GM. 1995. Multiple stable states and models of riparian

vegetation succession on the Animas River, Colorado. *Ann. Assoc. Am. Geogr.* **85**: 320-338.

Baldrian P, Trögl J, Frouz J, Šnajdr J, Valášková V, Merhautová V, Cajthaml T, and

Herinková J. 2008. Enzyme activities and microbial biomass in topsoil layer during

spontaneous succession in spoil heaps after brown coal mining. *Soil Biol. Biochem.* **40**: 2107-

2115.

Bardgett RD, and Walker LR. 2004. Impact of coloniser plant species on the development

of decomposer microbial communities following deglaciation. *Soil Biol. Biochem.* **36**: 555-

559.

Bardgett RD, Richter A, Bol R, Garnett MH, Baumler R, Xu X, Lopez-Capel E,

Manning DA, Hobbs PJ, Hartley IR, and Wanek W. 2007. Heterotrophic microbial

communities use ancient carbon following glacial retreat. *Biol. Lett.* **3**: 487-490.

Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, Wit P,

Sánchez-García M, Ebersberger I, and Sousa F. 2013. Improved software detection and

extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for

analysis of environmental sequencing data. *Methods Ecol. Evol.* **4**: 914-919.

Blaalid R, Carlsen T, Kumar S, Halvorsen R, Ugland KI, Fontana G, and Kauserud H.

2012. Changes in the root-associated fungal communities along a primary succession gradient

analysed by 454 pyrosequencing. *Mol. Ecol.* **21**: 1897-1908.

Bonet A, and Pausas JG. 2004. Species richness and cover along a 60-year chronosequence

in old-fields of southeastern Spain. *Plant Ecol.* **174**: 257-270.

Bowden RD. 1991. Inputs, outputs, and accumulation of nitrogen in an early successional

moss (*Polytrichum*) ecosystem. *Ecol. Monogr.* **61**: 207-223.

Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G,

Liancourt P, Tielbörger K, Travis JM, and Anthelme F. 2008. Facilitation in plant

communities: the past, the present, and the future. *J. Ecol.* **96**: 18-34.

Brown SP, and Jumpponen A. 2014. Contrasting primary successional trajectories of fungi

and bacteria in retreating glacier soils. *Mol. Ecol.* **23**: 481-497.

Brown AM, Warton DI, Andrew NR, Binns M, Cassis G, and Gibb H. 2014. The fourth-corner solution – using predictive models to understand how species traits interact with the environment. *Methods Ecol. Evol.* **5**: 344-352.

Chapin FS, Walker LR, Fastie CL, and Sharman LC. 1994. Mechanisms of primary succession following deglaciation at Glacier Bay, Alaska. *Ecol. Monogr.* **64**: 149-175.

Chen Y, Ding J, Peng Y, Li F, Yang G, Liu L, Qin S, Fang K, and Yang Y. 2016. Patterns and drivers of soil microbial communities in Tibetan alpine and global terrestrial ecosystems. *J. Biogeogr.* **43**: 2027-2039.

Connell JH, and Slatyer RO. 1977. Mechanisms of succession in natural communities and their role in community stability and organization. *Am. Nat.* **111**: 1119-1144.

Cornelissen JH, Lang SI, Soudzilovskaia NA, and During HJ. 2007. Comparative cryptogam ecology: a review of bryophyte and lichen traits that drive biogeochemistry. *Ann. Bot.* **99**: 987-1001.

Cutler NA, Chaput DL, and van der Gast CJ. 2014. Long-term changes in soil microbial communities during primary succession. *Soil Biol. Biochem.* **69**: 359-370.

De Cáceres M, and Legendre P. 2009. Associations between species and groups of sites: indices and statistical inference. *Ecology* **90**: 3566-3574.

de los Ríos A, Raggio J, Pérez-Ortega S, Vivas M, Pintado A, Green TA, Ascaso C, and Sancho LG. 2011. Anatomical, morphological and ecophysiological strategies in *Placopsis pycnotheca* (lichenized fungi, Ascomycota) allowing rapid colonization of recently deglaciated soils. *Flora* **206**: 857-864.

Delgado-Baquerizo M, and Eldridge DJ. 2019. Cross-biome drivers of soil bacterial alpha diversity on a worldwide scale. *Ecosystems* https://doi.org/10.1007/s10021-018-0333-2.

DeLuca TH, Zackrisson O, Nilsson M, and Sellstedt A. 2002. Quantifying nitrogen-fixation in feather moss carpets of boreal forests. *Nature* **419**: 917-920.

Dini-Andreote F, de Cássia Pereira e Silva, Michele, Triadó-Margarit X, Casamayor EO, van Elsas JD, and Salles JF. 2014. Dynamics of bacterial community succession in a salt marsh chronosequence: evidences for temporal niche partitioning. *ISME J.* **8**: 1989-2001.

Dini-Andreote F, Pylro VS, Baldrian P, van Elsas JD, and Salles JF. 2016. Ecological

succession reveals potential signatures of marine-terrestrial transition in salt marsh fungal

communities. *ISME J.* **10**: 1984-1997.

Dini-Andreote F, Stegen JC, van Elsas JD, and Salles JF. 2015. Disentangling

mechanisms that mediate the balance between stochastic and deterministic processes in

microbial succession. *Proc. Natl. Acad. Sci. USA* **112**: E1326-E1332.

Duc L, Noll M, Meier BE, Bürgmann H, and Zeyer J. 2009. High diversity of diazotrophs

in the forefield of a receding alpine glacier. *Microb. Ecol.* **57**: 179-190.

Edgar RC, Haas BJ, Clemente JC, Quince C, and Knight R. 2011. UCHIME improves

sensitivity and speed of chimera detection. *Bioinformatics* **27**: 2194-2200.

Edwards I, Bürgmann H, Miniaci C, and Zeyer J. 2006. Variation in microbial

community composition and culturability in the rhizosphere of *Leucanthemopsis alpina* (L.)

Heywood and adjacent bare soil along an alpine chronosequence. *Microb. Ecol.* **52**: 679-692.

Egler FE. 1954. Vegetation science concepts I. Initial floristic composition, a factor in old-

field vegetation development. *Plant Ecol.* **4**: 412-417.

Ellenberg H. 2009. *Vegetation ecology of Central Europe.* Cambridge: Cambridge

University Press.

Fierer N, Nemergut D, Knight R, and Craine JM. 2010. Changes through time: integrating

microorganisms into the study of succession. *Res Microbiol* **161 8**: 635-642.

Francini G, Liiri M, Männistö M, Stark S, and Kytöviita M. 2014. Response to reindeer

grazing removal depends on soil characteristics in low Arctic meadows. *Appl. Soil Ecol.* **76**:

14-25.

Gallet C, Nilsson M, and Zackrisson O. 1999. Phenolic metabolites of ecological

significance in *Empetrum hermaphroditum* leaves and associated humus. *Plant Soil* **210**: 1-9.

Gornall JL, Jónsdóttir IS, Woodin SJ, and van der Wal R. 2007. Arctic mosses govern

below-ground environment and ecosystem processes. *Oecologia* **153**: 931-941.

Gornall JL, Woodin SJ, Jónsdóttir IS, and van der Wal R. 2011. Balancing positive and
negative plant interactions: how mosses structure vascular plant communities. *Oecologia*
166: 769-782.

Hodkinson ID, Coulson SJ, and Webb NR. 2003. Community assembly along proglacial
chronosequences in the high Arctic: vegetation and soil development in north-west Svalbard.
J. Ecol. **91**: 651-663.

Hodkinson ID, Webb NR, and Coulson SJ. 2002. Primary community assembly on land –
the missing stages: why are the heterotrophic organisms always there first? *J. Ecol.* **90**: 569-
577.

Holland-Moritz H, Stuart J, Lewis LR, Miller S, Mack MC, McDaniel SF, and Fierer N.
2018. Novel bacterial lineages associated with boreal moss species. *Environ. Microbiol.* **20**:
2625-2638.

Ihrmark K, Bödeker I, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y,
Stenlid J, Brandström-Durling M, and Clemmensen KE. 2012. New primers to amplify
the fungal ITS2 region–evaluation by 454-sequencing of artificial and natural communities.
FEMS Microbiol. Ecol. **82**: 666-677.

Islam ZF, Cordero PRF, Feng J, Chen Y, Bay SK, Jirapanjawat T, Gleadow RM,
Carere CR, Stott MB, Chiri E, and Greening C. 2019. Two Chloroflexi classes
independently evolved the ability to persist on atmospheric hydrogen and carbon monoxide.
ISME J 10.1038/s41396-019-0393-0.

Jackson CR. 2003. Changes in community properties during microbial succession. *Oikos*
101: 444-448.

Ji M, Greening C, Vanwonterghem I, Carere CR, Bay SK, Steen JA, Montgomery K,
Lines T, Beardall J, and van Dorst J. 2017. Atmospheric trace gases support primary
production in Antarctic desert surface soil. *Nature* **552**: 400-403.

Jiang Y, Lei Y, Yang Y, Korpelainen H, Niinemets Ü, and Li C. 2018. Divergent
assemblage patterns and driving forces for bacterial and fungal communities along a glacier
forefield chronosequence. *Soil Biol Biochem* **118**: 207-216.

Jones GA, and Henry GHR. 2003. Primary plant succession on recently deglaciated terrain
in the Canadian High Arctic. *J. Biogeogr.* **30**: 277-296.

Jumpponen A. 2003. Soil fungal community assembly in a primary successional glacier
forefront ecosystem as inferred from rDNA sequence analyses. *New Phytol.* **158**: 569-578.

King C, and King G. 2014. Description of *Thermogemmatispora carboxidivorans* sp. nov., a
carbon-monoxide-oxidizing member of the class *Ktedonobacteria* isolated from a
geothermally heated biofilm, and analysis of carbon monoxide oxidation by members of the
class *Ktedonobacteria*. *Int. J. Syst. Evol. Microbiol.* **64**: 1244-1251.

Knelman JE, Legg TM, O'Neill SP, Washenberger CL, González A, Cleveland CC, and
Nemergut DR. 2012. Bacterial community structure and function change in association with
colonizer plants during early primary succession in a glacier forefield. *Soil Biol. Biochem.* **46**:
172-180.

Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST,
Bruns TD, Bengtsson-Palme J, and Callaghan TM. 2013. Towards a unified paradigm for
sequence-based identification of fungi. *Mol. Ecol.* **22**: 5271-5277.

Koster EA. 1988. Ancient and modern cold-climate aeolian sand deposition: a review. *J*
Quaternary Sci **3**: 69-83.

Kowal J, Pressel S, Duckett JG, Bidartondo MI, and Field KJ. 2018. From rhizoids to
roots? Experimental evidence of mutualism between liverworts and ascomycete fungi. *Ann.*
Bot. **121**: 221-227.

Kozich JJ, Westcott SL, Baxter NT, Highlander SK, and Schloss PD. 2013. Development
of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence
data on the MiSeq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**: 5112-5120.

Kuske CR, Yeager CM, Johnson S, Ticknor LO, and Belnap J. 2012. Response and
resilience of soil biocrust bacterial communities to chronic physical disturbance in arid
shrublands. *ISME J* **6**: 886-897.

Lichter J. 1998. Primary succession and forest development oncoastal Lake Michigan sand
dunes. *Ecol. Monogr.* **68**: 487-510.

Loreau M, and de Mazancourt C. 2013. Biodiversity and ecosystem stability: a synthesis of

underlying mechanisms. *Ecol. Lett.* **16**: 106-115.

Maier S, Tamm A, Wu D, Caesar J, Grube M, and Weber B. 2018. Photoautotrophic

organisms control microbial abundance, diversity, and physiology in different types of

biological soil crusts. *ISME J.* **12**: 1032-1046.

Mäki A, Rissanen AJ, and Tiirola M. 2016. A practical method for barcoding and size-

trimming PCR templates for amplicon sequencing. *BioTechniques* **60**: 88-90.

Makoto K, and Wilson SD. 2019. When and where does dispersal limitation matter in

primary succession? *J. Ecol.* **107**: 559-565. 10.1111/1365-2745.12988.

McArdle BH, and Anderson MJ. 2001. Fitting multivariate models to community data: a

comment on distance-based redundancy analysis. *Ecology* **82**: 290-297.

Merilä P, Malmivaara-Lämsä M, Spetz P, Stark S, Vierikko K, Derome J, and Fritze H.

2010. Soil organic matter quality as a link between microbial community structure and

vegetation composition along a successional gradient in a boreal forest. *App.l Soil Ecol.* **46**:

259-267.

Miniaci C, Bunge M, Duc L, Edwards I, Bürgmann H, and Zeyer J. 2007. Effects of

pioneering plants on microbial structures and functions in a glacier forefield. *Biol. Fertility*

Soils **44**: 289-297.

Montgomery DR. 2007. Soil erosion and agricultural sustainability. *Proc. Natl. Acad. Sci.*

USA **104**: 13268-13272.

Navarro-Noya YE, Jiménez-Aguilar A, Valenzuela-Encinas C, Alcántara-Hernández

RJ, Ruíz-Valdiviezo VM, Ponce-Mendoza A, Luna-Guido M, Marsch R, and

Dendooven L. 2014. Bacterial communities in soil under moss and lichen-moss crusts.

Geomicrobiol. J. **31**: 152-160.

Nemergut DR, Anderson SP, Cleveland CC, Martin AP, Miller AE, Seimon A, and

Schmidt SK. 2007. Microbial community succession in an unvegetated, recently deglaciated

soil. *Microb. Ecol.* **53**: 110-122.

Odum EP. 1969. The strategy of ecosystem development. *Science* **164**: 262-270.

Oksanen J, Blanchet FG, Friendly M, Kindt R, McGlinn D, Legendre P, Minchin PR,

O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, and Wagner H. 2017.

vegan: Community Ecology Package. R package version 2.4-2. https://CRAN.R-

project.org/package=vegan.

Olson JS. 1958. Rates of succession and soil changes on southern Lake Michigan sand

dunes. *Bot. Gaz.* **119**: 125-170.

Ortiz-Álvarez R, Fierer N, de los Ríos A, Casamayor EO, and Barberán A. 2018.

Consistent changes in the taxonomic structure and functional attributes of bacterial

communities during primary succession. *ISME J.* **12**: 1658-1667.

Pennanen T, Strommer R, Markkola A, and Fritze H. 2001. Microbial and plant

community structure across a primary succession gradient. *Scand. J. For. Res.* **16**: 37-43.

Peyrat J, and Fichtner A. 2011. Plant species diversity in dry coastal dunes of the southern

Baltic coast. *Community Ecol.* **12**: 220-226.

Pirinen P, Simola H, Aalto J, Kaukoranta J, Karlsson P, and Ruuhela R. 2012.

Climatological statistics of Finland 1981–2010. *Finnish Meteorological Institute Reports* **1**:

1-96.

Poosakkannu A, Nissinen R, Männistö M, and Kytöviita M. 2017. Microbial community

composition but not diversity changes along succession in arctic sand dunes. *Environ.*

Microbiol. **19**: 698-709.

Porazinska DL, Farrer EC, Spasojevic MJ, Bueno de Mesquita, Clifton P, Sartwell SA,

Smith JG, White CT, King AJ, Suding KN, and Schmidt SK. 2018. Plant diversity and

density predict belowground diversity and function in an early successional alpine ecosystem.

Ecology **99**: 1942-1952.

Powell JR, Karunaratne S, Campbell CD, Yao H, Robinson L, and Singh BK. 2015.

Deterministic processes vary during community assembly for ecologically dissimilar taxa.

Nat. Commun. **6**: 8444.

Prach K, Řehounková K, Lencová K, Jírová A, Konvalinková P, Mudrák O, Študent V,

Vaněček Z, Tichý L, and Petřík P. 2014. Vegetation succession in restoration of disturbed

sites in Central Europe: the direction of succession and species richness across 19 seres. *Appl.*

Veg. Sci. **17**: 193-200.

Pressel S, Bidartondo MI, Ligrone R, and Duckett JG. 2010. Fungal symbioses in

bryophytes: new insights in the twenty first century. *Phytotaxa* **9**: 238-253.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, and Glockner

FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. *Nucleic Acids Res.* **41**: D590-6.

RCoreTeam. 2018. *R: A language and environment for statistical computing.*,Vienna,

Austria: Foundation for Statistical Computing. https://www.R-project.org/

Read DJ. 1991. Mycorrhizas in ecosystems. *Experientia* **47**: 376-391.

Rime T, Hartmann M, Brunner I, Widmer F, Zeyer J, and Frey B. 2015. Vertical

distribution of the soil microbiota along a successional gradient in a glacier forefield. *Mol.*

Ecol. **24**: 1091-1108.

Roy-Bolduc A, Bell TH, Boudreau S, and Hijri M. 2015. Comprehensive sampling of an

isolated dune system demonstrates clear patterns in soil fungal communities across a

successional gradient. *En.v Microbiol. Rep.* **7**: 839-848.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski

RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn

DJ, and Weber CF. 2009. Introducing mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial communities. *Appl.*

Environ. Microbiol. **75**: 7537-7541.

Schmidt SK, Nemergut DR, Darcy JL, and Lynch R. 2014. Do bacterial and fungal

communities assemble differently during primary succession? *Mol. Ecol.* **23**: 254-258.

Schmidt SK, Reed SC, Nemergut DR, Grandy AS, Cleveland CC, Weintraub MN, Hill

AW, Costello EK, Meyer AF, Neff JC, and Martin AM. 2008. The earliest stages of

ecosystem succession in high-elevation (5000 metres above sea level), recently deglaciated

soils. *P Roy Soc B-Biol Sci* **275**: 2793-2802.

Seppälä M. 1995. Deflation and redeposition of sand dunes in Finnish Lapland. *Quaternary*

Sci. Rev. **14**: 799-809.

785 **Smith SE, and Read DJ. 2008.** *Mycorrhizal symbiosis.* London, UK: Academic Press.

786 **Sparrius L, Sevink J, and Kooijman A. 2012.** Effects of nitrogen deposition on soil and
787 vegetation in primary succession stages in inland drift sands. *Plant Soil* **353**: 261-272.

788 **Tebo BM, Davis RE, Anitori RP, Connell LB, Schiffman P, and Staudigel H. 2015.**
789 Microbial communities in dark oligotrophic volcanic ice cave ecosystems of Mt. Erebus,
790 Antarctica. *Front. Microbiol.* **6**: 179.

791 **Tedersoo L, Bahram M, Polme S, Koljalg U, Yorou NS, Wijesundera R, Villarreal Ruiz**
792 **L, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A,**
793 **Rosas M, Riit T, Ratkowsky D, Pritsch K, Poldmaa K, Piepenbring M, Phosri C,**
794 **Peterson M, Parts K, Partel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH,**
795 **Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS, Larsson KH, Kohout**
796 **P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo LD, Greslebin A, Grelet G, Geml**
797 **J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen**
798 **X, Buegger F, Brearley FQ, Bonito G, Anslan S, Abell S, and Abarenkov K. 2014.**
799 Global diversity and geography of soil fungi. *Science* **346**: 1256688.

800 **Tripathi BM, Stegen JC, Kim M, Dong K, Adams JM, and Lee YK. 2018.** Soil pH
801 mediates the balance between stochastic and deterministic assembly of bacteria. *ISME J.* **12**:
802 1072-1083.

803 **Tscherko D, Hammesfahr U, Marx M, and Kandeler E. 2004.** Shifts in rhizosphere
804 microbial communities and enzyme activity of *Poa alpina* across an alpine chronosequence.
805 *Soil Biol. Biochem.* **36**: 1685-1698.

806 **Van der Wal R, and Brooker R. 2004.** Mosses mediate grazer impacts on grass abundance
807 in arctic ecosystems. *Funct. Ecol.* **18**: 77-86.

808 **Vanwalleghem T, Gómez JA, Infante Amate J, González de Molina M, Vanderlinden K,**
809 **Guzmán G, Laguna A, and Giráldez JV. 2017.** Impact of historical land use and soil
810 management change on soil erosion and agricultural sustainability during the Anthropocene.
811 *Anthropocene* **17**: 13-29.

812 **Vitousek PM. 1994.** Potential nitrogen fixation during primary succession in Hawaii
813 Volcanoes National Park. *Biotropica* **26**: 234-240.

Waldrop MP, Zak DR, Blackwood CB, Curtis CD, and Tilman D. 2006. Resource
availability controls fungal diversity across a plant diversity gradient. *Ecol. Lett.* **9**: 1127-
1135.

Wang Y, Naumann U, Wright ST, and Warton DI. 2012. mvabund–an R package for
model-based analysis of multivariate abundance data. *Methods Ecol. Evol.* **3**: 471-474.

Wardle DA, Nilsson M, Gallet C, and Zackrisson O. 1998. An ecosystem-level
perspective of allelopathy. *Biol. Rev.* **73**: 305-319.

Weber B, Büdel B, and Belnap J. 2016. Biological soil crusts: An organizing principle in
drylands. Springer.

Williams MA, Jangid K, Shanmugam SG, and Whitman WB. 2013. Bacterial
communities in soil mimic patterns of vegetative succession and ecosystem climax but are
resilient to change between seasons. *Soil Biol. Biochem.* **57**: 749-757.

Wilson JB, Gitay H, Roxburgh SH, King WM, and Tangney RS. 1992. Egler's concept of
'initial floristic composition' in succession: Ecologists citing it don't agree what it means.
Oikos **64**: 591-593.

Xie C, and Lou H. 2009. Secondary metabolites in bryophytes: an ecological aspect. *Chem.*
Biodivers. **6**: 303-312.

Zhang T, Zhang Y, Liu H, Wei Y, Li H, Su J, Zhao L, and Yu L. 2013. Diversity and cold
adaptation of culturable endophytic fungi from bryophytes in the Fildes Region, King George
Island, maritime Antarctica. *FEMS Microbiol. Lett.* **341**: 52-61.

Zhang Y, Zheng N, Wang J, Yao H, Qiu Q, and Chapman SJ. 2019. High turnover rate of
free phospholipids in soil confirms the classic hypothesis of PLFA methodology. *Soil Biol.*
Biochem. **135**: 323-330.

Figure legends

Figure 1 (A) Organic matter, (B) pH, (C) microbial biomass based on phospholipid fatty acids (PLFAs) per soil organic matter (OM) content, (D) microbial biomass per gram soil dry weight (dw), (E) ratio of fungal to bacterial biomass based on PLFA, and (F) bacterial 16S rRNA gene (upper row, darker colours) and fungal ITS amplicon (lower row, lighter colours) OTU numbers in sand dune successional stages. Data are averages of four deflation basins and three replicates per basin. Ends of whiskers represent minimum and maximum values excluding outliers. Different letters indicate significant differences at $p<0.05$.

Figure 2 Non-metric multidimensional scaling (NMDS) plots of (A) bacterial communities based on 16S rRNA gene amplicon sequencing (stress=0.10) and (B) fungal communities based on ITS2 region amplicon sequencing (stress=0.13) in sand dune successional stages. Data points represent three replicate samples from four deflation basins per successional stage.

Figure 3 Average distances of successional stage pairs for bacterial, fungal and plant communities and for soil chemistry. Different letters indicate significant differences at $p<0.05$. Ends of whiskers represent minimum and maximum values excluding outliers. Shading indicates the largest transitions.

Figure 4 (A) Phylum level (class level for Proteobacteria) community structure of bacteria based on 16S rRNA gene sequencing and (B) class-level community structure of fungi based on ITS2 region sequencing with sand dune successional stage. Data are averages of four deflation basins and three replicates per basin. For values of individual samples and groups combined in 'other', see Fig. S3 and S4.

Figure 5 Biomass of selected early and cryptrogam stage bacteria with sand dune successional stage. Relative abundances from sequencing data were converted into biomass by multiplying with the total microbial PLFA concentration. Horizontal lines indicate mean values. Labelled points are outliers that would be outside the plot area.

Figure 6 Bacterial indicator OTUs for successional stages grouped at order level. Thicker horizontal lines show where the successional stage changes, and each group is placed in the stage where it was the most abundant. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients describing the association of the indicator OTU groups with environmental variables (env. variables). Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter.

Figure 7 Fungal indicator OTUs for successional stages grouped at class level. Thicker horizontal lines show where successional stage changes, and each group is placed in the stage where it was the most abundant. A, Ascomycota; B, Basidiomycota; R, Rozellomycota; Z, Zygomycota; cls, class; ord, order; Inc. sed., Incertae sedis. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients describing the association of the indicator OTU groups with environmental variables (env. variables). Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter, minT, minimum soil temperature.

Figures

Figure 1 (A) Organic matter, (B) pH, (C) microbial biomass based on phospholipid fatty acids (PLFAs) per soil organic matter (OM) content, (D) microbial biomass per gram soil dry weight (dw), (E) ratio of fungal to bacterial biomass based on PLFA, and (F) bacterial 16S rRNA gene (upper row, darker colours) and fungal ITS amplicon (lower row, lighter colours) OTU numbers in sand dune successional stages. Data are averages of four deflation basins and three replicates per basin. Ends of whiskers represent minimum and maximum values excluding outliers. Different letters indicate significant differences at p<0.05.

Figure 2 Non-metric multidimensional scaling (NMDS) plots of (A) bacterial communities based on 16S rRNA gene amplicon sequencing (stress=0.10) and (B) fungal communities based on ITS2 region amplicon sequencing (stress=0.13) in sand dune successional stages. Data points represent three replicate samples from four deflation basins per successional stage.

Figure 3 Average distances of successional stage pairs for bacterial, fungal and plant communities and for soil chemistry. Different letters indicate significant differences at p<0.05. Ends of whiskers represent minimum and maximum values excluding outliers. Shading indicates the largest transitions.

Figure 4 (A) Phylum level (class level for Proteobacteria) community structure of bacteria based on 16S rRNA gene sequencing and (B) class-level community structure of fungi based on ITS2 region sequencing with sand dune successional stage. Data are averages of four deflation basins and three replicates per basin. For values of individual samples and groups combined in 'other', see Fig. S3 and S4.

Figure 5 Biomass of selected early and cryptrogam stage bacteria with sand dune successional stage. Relative abundances from sequencing data were converted into biomass by multiplying with the total microbial PLFA concentration. Horizontal lines indicate mean values. Labelled points are outliers that would be outside the plot area.

Figure 6 Bacterial indicator OTUs for successional stages grouped at order level. Thicker horizontal lines show where the successional stage changes, and each group is placed in the stage where it was the most abundant. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients describing the association of the indicator OTU groups with environmental variables (env. variables). Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter.

Figure 7 Fungal indicator OTUs for successional stages grouped at class level. Thicker horizontal lines show where successional stage changes, and each group is placed in the stage where it was the most abundant. A, Ascomycota; B, Basidiomycota; R, Rozellomycota; Z, Zygomycota; cls, class; ord, order; Inc. sed., Incertae sedis. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients describing the association of the indicator OTU groups with environmental variables (env. variables). Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter, minT, minimum soil temperature.