Cryptogams signify key transition of bacteria and fungi in Arctic sand

dune succession

Heli Juottonen[1*], Minna Männistö[2], Marja Tiirola[1] & Minna-Maarit Kytöviita[1]

[1]Department of Biological and Environmental Science, University of Jyväskylä, Finland

[2] Natural Resources Institute Finland (Luke), Rovaniemi, Finland

*Corresponding author. Heli Juottonen, Department of Biological and Environmental

Science, P.O. Box 35, 40014 University of Jyväskylä, Finland.

email: heli.juottonen@alumni.helsinki.fi

Summary

• Primary succession models focus on aboveground vascular plants. However, the

prevalence of mosses and lichens, i.e. cryptogams, suggests they play a role in soil

successions. Here, we explore whether effects of cryptogams on belowground

microbes can facilitate progressive shifts in sand dune succession.

• We linked aboveground vegetation, belowground bacterial and fungal community,

and soil chemistry in six successional stages in Arctic inland sand dunes: bare sand,

grass, moss, lichen, ericoid heath and mountain birch forest.

• Compared to the bare sand and grass stages, microbial biomass and the proportion of

fungi increased in the moss stage, and later stage microbial groups appeared despite

the absence of their host plants. The microbial communities of the lichen stage

resembled the communities in the vascular plant stages. Bacterial community

correlated better with soil chemistry than with vegetation, whereas the correlation of

fungi with vegetation increased with vascular vegetation.

• Distinct bacterial and fungal patterns of biomass, richness, and plant-microbe

interaction showed that the aboveground vegetation change structured the bacterial

and fungal community differently. The nonalignment of aboveground vs.

belowground changes suggests that cryptogams can drive succession towards vascular

plant dominance through microbially mediated facilitation in eroded Arctic soil.

Keywords: Arctic soil, bacterial community, cryptogam, erosion, fungal community,

Polytrichum, primary succession, sand dune

Introduction

Development of vegetation on bare land and the sequence of vegetative stages that

predictably follow each other have been described in several theories of succession (Egler,

1954; Wilson *et al.*, 1992). According to widely accepted views, primary succession is

influenced by site abiotic conditions (climate, soil parent material, topography) and biotic

factors (plant species pool present, order of arrival, species interactions). Although focusing

on plants, most models of terrestrial primary succession overlook bryophyte communities

(e.g. Connell and Slatyer, 1977; Baker and Walford, 1995), whereas most reports of primary

succession include moss and lichen dominated stages (Chapin *et al.*, 1994; Lichter, 1998;

Hodkinson *et al.*, 2003; Jones and Henry, 2003).

Ecosystem succession includes stochastic elements for instance through priority effects and,

on the other hand, deterministic influences such as competition and facilitation (Måren *et al.*,

2018). Cryptogams, here mosses and lichens, may have diverse but unexplored competitive

and facilitative effects on the development of vascular plant and belowground microbial

communities. Firstly, dispersal of cryptogams by spores minimizes dispersal limitation – a

feature classically included in models explaining succession of vascular plant communities

(Makoto and Wilson, 2019). The rapid arrival of mosses warrants them a priority effect in

establishing plant communities. Secondly, although mosses and lichens lack roots and root-

mediated effects on microbial communities, they affect soil temperature, moisture, and

carbon and nitrogen availability, which may promote or inhibit vascular vegetation (Van der

Wal and Brooker, 2004; Gornall *et al.*, 2007; Cornelissen *et al.*, 2007; Gornall *et al.*, 2011).

Moss rhizoids and litter provide carbon to the belowground soil (Bowden, 1991), and mosses

can release carbon also in drying and wetting cycles (Wilson and Coxon, 1999). Many

lichens characteristic of primary succession fix atmospheric nitrogen in specific structures

called cephalodia (Vitousek, 1994), whereas moss leaves are a habitat of N_2-fixing bacteria

(DeLuca *et al.*, 2002; Arróniz-Crespo *et al.*, 2014). Thirdly, mosses and lichens produce

unique secondary metabolites that may have adverse effects on vascular plants and beneficial

or adverse effects on microbes (Cornelissen *et al.*, 2007; Xie and Lou, 2009).

Despite the fact that microbes are the first colonizers of any barren surface and modify soil

chemistry during succession, there are only few models predicting successional trajectories of

bacteria and fungi (Jackson, 2003; Fierer *et al.*, 2010 Dini-Andreote *et al.*, 2015; Tripathi *et*

al., 2018; Ortiz-Álvarez *et al.*, 2018). Belowground microbial community shifts in early

succession are considered to be driven by soil chemistry, external resources and dispersal of

microbes, and in later stages by biotic factors, such as establishment and changes in the

vegetation (Brown and Jumpponen, 2014; Jiang *et al.*, 2018). Bacterial versus fungal

communities can be expected to show distinct trajectories in primary succession: Bacteria are

smaller and believed to disperse easier than fungi, which may lead to more deterministic

community assembly for bacteria (Schmidt *et al.*, 2014; Powell *et al.*, 2015) and higher

influence of priority effects and stochastic effects for fungi (Brown and Jumpponen, 2014;

Schmidt *et al.*, 2014; Jiang *et al.*, 2018). Further, the physiological diversity in bacteria

enables some of them to survive as autotrophs in bare oligotrophic soils of early successional

stages in the absence of plants fixing carbon (Nemergut *et al.*, 2007; Schmidt *et al.*, 2008;

Duc *et al.*, 2009). Fungi in these habitats depend on already fixed carbon and nitrogen from

wind-blown, ancient, or microbially fixed sources (Schmidt *et al.*, 2014). The critical shift to

an ecosystem based on the biomass production by the resident plants is associated with

changes in the microbial community (Bardgett and Walker, 2004; Edwards *et al.*, 2006;

Blaalid *et al.*, 2012; Knelman *et al.*, 2012). Along succession, the fungi to bacteria ratio

generally increases as soil organic matter increases and pH decreases (Pennanen *et al.*, 2001;

Tscherko *et al.*, 2004; Bardgett and Walker, 2004). These results do not, however, fully

predict vegetation development in succession. We suggest that belowground changes

associated with cryptogams are a key factor in the ecosystem shift from early to late

succession. We link belowground community development with that aboveground and expect

the interaction strength to vary and that the community changes do not take place in complete

synchrony as is inferred in facilitation models (Brooker *et al.*, 2008).

Ecosystems are declining due to soil erosion and loss of vegetation cover (Montgomery,

2007; Vanwalleghem *et al.*, 2017), and it is critical to identify the key factors promoting

shifts between non-vegetated and vegetated stages in succession. Identifying microbial and

plant communities that signal for the shift from autotrophic to mainly heterotrophic microbial

biomass production can help identify factors that drive the shift back to vascular plant cover,

soil stabilization, and a productive ecosystem. One example of areas where vegetation has

been lost due to environmental changes and lack of recovery are Arctic inland sand dunes.

The dunes have formed from wind-blown sand after the last ice age (Koster, 1988). Plants

colonized the dunes later on, but today large areas lack vascular plant cover. Forest fires,

insect outbreaks, and overgrazing may have caused the ecosystem degradation (Seppälä,

1995). In our study site in Northern Finland, wind erosion has carved deflation basins into the

dunes. Inside the basin, primary succession vegetation stages have developed, and dune

slopes surrounding the basins are covered by mountain birch forest, the climax stage in the

present climate. Eroding inland sand dunes provide an excellent opportunity to study primary

succession due to the mosaic of closely situated successional stages within deflation basin,

including moss- and lichen-dominated areas. In temperate inland dunes, the cryptogam stages

have been identified as pioneer vegetation stages of the succession (Sparrius *et al.*, 2012).

Previous work comparing the start and end points of the Arctic inland dune succession

proposed that bacteria and fungi may show distinct successional patterns (Poosakkannu *et al.*,

2017). The role of the cryptogam stages as part of successional trajectories above- and

belowground has not been elucidated, although the prevalence of lichens and mosses suggests

they are important.

In this study, we explored whether the belowground and aboveground community changes

take place in synchrony in Arctic inland sand dune succession. We hypothesized that,

compared to bare soil, the cryptogam-covered stages contain belowground soil microbial

community more similar to the stages covered by vascular plants. Such nonalignment of

belowground vs. aboveground communities could indicate that the cryptogam stages drive

succession towards vascular plant dominance through microbially mediated facilitation.

Secondly, we compared the belowground responses of two microbial groups – bacteria and

fungi - to successional stages defined by aboveground vegetation. We hypothesized that i) in

comparison to fungi, bacteria would be more influenced by cryptogam-associated changes in

the soil chemical environment because of their limited saprophytic capacity and potential

autotrophy, and ii) as powerful and obligate decomposers, fungal community would respond

particularly to the appearance of vascular vegetation and consequently organic matter

accumulation. Such a pattern would indicate differences in the way vegetation change and

plant-microbe interaction structure bacterial vs. fungal communities along the succession. By

linking the aboveground vegetation, the belowground microbial communities, and the

associated soil chemical environment, we aimed to define the multi-faceted role of

cryptogams in Arctic ecosystem succession.

Materials and methods

Study site and sampling

The study site is located in an inland aeolian (i.e. wind-deposited) dune area in Northern

Fennoscandia (68° 29′ 16″ N, 24° 42′ 13″ E). The 1981–2000 average annual temperature in

the region was -1.3 °C, with extreme air temperatures between +30 °C and -52 °C and the

average annual precipitation 550 mm (Pirinen *et al.*, 2012). We defined six successional

stages in the dunes based on vegetation (Fig. 1a). Although the soil in all the stages was

aeolian sand, in three stages the largest part of the surface was mostly exposed sand (92-100

% sand), while in the other three stages vegetation covered the sand (0-52 % sand). The sand-

exposed stages ('sandy stages') were: 1) sand: bare loose sand without plants or visible soil

crust, 2) grass: bare loose sand with the grass *Deschampsia flexuosa* and without visible soil

crust, and 3) moss: bare loose sand with the moss *Polytrichum piliferum*. The stages with

extensive vegetation cover ('vegetated stages') were 4) lichen: the main species lichens

Stereocaulon spp. and *Cladonia* spp., and the mosses *P. piliferum* and *Racomitrium*

ericoides, 5) heath: the dominant species dwarf shrub crowberry *Empetrum nigrum* with

Cladonia spp. and/or *Stereocaulon* spp., and 6) forest: mountain birch (*Betula pubescens*

subsp. czerepanovii) forest, the main species in study plots the shrubs *E. nigrum* and

Vaccinium vitis-idaea, the moss *Pleurozium schreberi*, *D. flexuosa*, and the herb *Linnaea*

borealis. The sand, grass, moss, and lichen stages are located at the bottom or on the inner

slopes of the deflation basins and heath at the edge of the basins. The remnant mountain birch

forests are located around the basins at the crest of the dune or on the outer slope. Permanent

study plots of 1 × 1 m were established in 2008 and included each of the six vegetation stages

in seven deflation basins. The vegetation in the plots was recorded in detail in 2009 by point

frequency method (100 points within 50 × 50 cm, presence/absence and frequency of

species).

We sampled soil in each successional stage in four of the basins in July 2011. To take into

account that plants and microbes vary at different spatial scales, the soil samples were

collected in triplicate. The samples were taken from the depth of 0-10 cm with a 12.5 cm ×

12.5 cm soil corer (72 samples in total: six successional stages × four basins × three soil cores

per 1×1 m study plot). Soil was sieved with a 4-mm sieve and frozen at -20°C.

Soil chemistry and temperature

Total C and N of soil organic matter, soil P, Ca, K, Na, Fe, Mn, Al, and Mg contents, and pH

were measured as detailed in Methods S1. Soil temperature at the 10 cm depth was recorded

every two hours with Hobo Temp External H08-002-02 loggers coupled with TMC6-HD soil

temperature sensor during 2009-2015. We defined maximum temperature as the average

temperature of the warmest month (July) and the minimum temperature as the average of the

two coldest months (January and February).

Phospholipid fatty acid (PLFA) analysis

Phospholipid fatty acids were extracted as described previously (Francini *et al.*, 2014) from

freeze-dried soil (15 g for sand and grass stages, 10 g for moss, 8 g for lichen, 5 g for heath

and 3 g for forest stage corresponding to OM content). Briefly, lipids were extracted with

chloroform:methanol:citrate buffer (Bligh and Dyer, 1:2:0.8 v/v/v), fractionated using silica

columns (Bond Elut LRC, Varian), and phospholipids were methylated with mild alkaline

methanolysis. The resulting fatty acid methyl esters were analyzed by gas chromatography

and PLFAs identified based on a bacterial fatty acid standard mixture (Supelco, Bellefonte,

PA, USA) and fatty acids from reference strains. For community analysis, data was expressed

as peak area percentage of a single PLFA of the total area of the profile, and for biomass

analysis as nmol of PLFAs per gram dry weight (gdw) of soil or per gram OM. The sum of

PLFAs i15:0, a15:0, 15:0, i16:0, 16:1ω9, i17:0, a17:0/17:1ω8c (eluted in the same peak),

17:0, cyclo-17:0, 18:1ω7c/ω9t (eluted in the same peak) and cyclo-19:0 were used to

represent bacterial biomass and PLFA 18:2ω6 fungal biomass. Sum of all these PLFAs was

used to represent total microbial biomass. Biomass of selected bacterial groups was

calculated by multiplying the total microbial biomass of a sample with the relative abundance

of the bacterial group in the sample based on 16S rRNA gene sequencing.

DNA extraction and PCR

DNA was extracted from 0.85 g (0.4 g for forest stage) soil with Powersoil DNA extraction

kit (MoBio). Bead beating was carried out with a FastPrep instrument at 5.0 m s^{-1} for 40 s.

DNA concentration was determined with Qubit fluorometer and Qubit dsDNA HS kit

(Invitrogen).

Bacterial and fungal PCRs were carried out as two-step PCR where the second step

introduced barcodes and adapters for Ion Torrent sequencing. The first step of bacterial 16S

rRNA gene PCR (V1-V2 region) was carried out with primers 27F (5'-

AGAGTTTGATCCTGGCTCAG-3') and 338R (5'-TGCTGCCTCCCGTAGGAGT-3').The

first step of intergenic transcribed region (ITS) PCR for fungi was carried out with primers

fITS7 (5'-GTGARTCATCGAATCTTTG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-

3') (Ihrmark *et al.*, 2012), where the fITS7 primer contained an M13 adapter for the second

step (Mäki *et al.*, 2016).

The second step primers introduced barcodes and Ion Torrent adapters A and P1. For

bacterial 2nd step PCR, the forward primer was adapter A-barcode-27F, and the reverse

primer adapter P1-338R. For the fungal 2nd step PCR, the forward primer was adapter A-

barcode-M13 adapter and the reverse primer adapter P1-ITS4. Details for PCR are given in

Methods S1. After the second PCR, the duplicate products were pooled, purified with

Agencourt AMPure XP purification system (Beckman Coulter), and quantified with Qubit

fluorometer and Qubit dsDNA HS kit. Equimolar amounts (10 ng) of products were pooled

for sequencing.

Ion Torrent sequencing and sequence analysis

Pooled bacterial and fungal PCR products were sequenced after emulsion PCR with the Ion

OneTouch system and Ion OT2 400 kit (Life Technologies) on Ion 314 chips with Ion PGM

Sequencing 400 Kit (Life Technologies) according to the manufacturer's instructions. The

sequences were processed in mothur software package v. 1.38 (Schloss *et al.*, 2009)

following the relevant parts of MiSeq SOP protocol (https://mothur.org/wiki/MiSeq_SOP;

accessed in January 2017; Kozich *et al.*, 2013). Sequences were quality-filtered using average

quality of 25 and a window size of 50 bases, a minimum sequence length of 200 bp, a

maximum length of 400 bp for bacteria and 410 bp for fungi, and further settings maximum

homopolymer length maxhomop=6, maximum number of ambigious bases maxambig=0,

maximum number of differences to primer sequence pdiffs=1, and maximum number of

differences to barcode sequence bdiffs=0. Bacterial sequences were aligned against Silva

database v. 1.23 (Quast *et al.*, 2013). Chimeras were detected with Uchime in de novo mode

(Edgar *et al.*, 2011) in mothur. Fungal ITS2 region was extracted and non-fungal sequences

were removed with ITSx (Bengtsson-Palme *et al.*, 2013). After quality filtering, alignment

for bacteria, and removal of chimeras and non-target sequences, we had 121711 bacterial

sequence reads and 279885 fungal reads. Reads were preclustered with setting diffs=2

(bacteria) or diffs=1 (fungi). The sequences were clustered into 97% operational taxonomic

units (OTUs) using the average neighbor method for bacteria, and the vsearch algorithm for

fungi, yielding 5242 bacterial and 1742 fungal OTUs after removal of singletons. Bacterial

OTUs were classified against the Silva v. 1.23 database and fungi against the Unite database

(Kõljalg *et al.*, 2013). Data was subsampled to the read number of the sample with the lowest

number of reads, which was 1055 reads for bacteria and 1605 reads for fungi. Shannon

diversity index and coverage were calculated in mothur. Coverage was $76 \pm 6\%$ for bacteria

and $97 \pm 1\%$ for fungi. The sequence data were submitted to NCBI under BioProject

accession PRJNA471306.

Statistical analyses

Chemistry and microbial biomass were compared among the successional stages using linear

mixed models with R package nlme and function lme with successional stage as a fixed

factor and deflation basin as a random blocking factor. Chemistry and biomass values except

pH and Al were log transformed to normalize their distribution.

Bacterial and fungal sequence data were converted to relative abundances of OTUs and

square root transformed to achieve Hellinger transformation that reduces the weight of OTUs

with low counts and zeros. Only OTUs with $5 \geq$ reads were included. Variation of

community structure was visualized as non-metric multidimensional scaling (NMDS) plots

with function metaMDS using the Bray-Curtis distance measure. Dispersion was visualized

in the NMDS plots as ellipses based on standard deviation of sample scores using the

ordiellipse function. Multivariate dispersion between the successional stages was tested with

betadisper (Bray-Curtis distances). Permutational analysis of variance (PERMANOVA,

Anderson, 2001; McArdle and Anderson, 2001) with the adonis2 function (Bray-Curtis

distances) was used to compare community structure between the successional stages and in

response to environmental variables. In the univariate model for successional stage, the stage

was nested within deflation basin. The multivariate PERMANOVA model for environmental

variables included OM-pH gradient and in addition Al and minimum soil temperature, which

did not correlate with OM and pH (Pearson $|r|<0.7$) but differed with successional stage. The

OM-pH gradient was defined with detrended correspondence analysis, and the first axis

scores were used as a variable in PERMANOVA. Impact of the individual environmental

variables was determined with marginal tests. Transitions between successional stages were

quantified by calculating average distances of the successional stages from bare sand for

bacterial and fungal community structure (Bray-Curtis distances), vegetation (Bray-Curtis

distances of square root transformed data) and soil chemistry (Euclidean distances). Using the

same distance measures, we carried out Procrustes analysis with the function procrustes to

correlate microbial community structure with vegetation, and microbial community structure

with soil chemistry (Lisboa *et al*., 2014). The analysis was based on the first four axes of

NMDS analysis of bacterial community, fungal community, vegetation and soil chemistry

(stress for all ordinations ≤ 0.07). The strength and the significance of the Procrustes

correlation was tested with the function protest (Peres-Neto and Jackson, 2001). Residuals

from the Procrustes analysis were used to compare the strength of the plant-microbe and soil

chemistry-microbe correlation along the succession by comparing the successional stages

with linear mixed models as above for soil chemistry. To separate the effects of vegetation

and soil chemistry on the microbial community, we carried out partial Procrustes analysis

where the variation due to soil chemistry was first partitioned out by multiple regression of

vegetation and microbial data against soil chemistry, and the residuals were then used in

Procrustes analysis (Peres-Neto and Jackson, 2001; Lisboa *et al*., 2014). All multivariate

analyses were carried out with the vegan package (v.2.4.2, Oksanen *et al.*, 2017) in R.

Significance tests were based on 999 permutations.

We identified bacterial and fungal OTUs characteristic to specific successional stages or

combinations of stages by indicator species analysis with the package indicspecies using the

function multipatt and the association index IndVal.g which corrects for unequal site numbers

(De Cáceres and Legendre, 2009). Only OTUs that occurred in the subsampled data at least

12 times, i.e., once in each sample of a successional stage, were included in the analyses. The

combinations of stages included in the analyses were each successional stage on its own, all

six stages together, all stages except sand, all stages except forest, all stages except sand and

grass, all stages except heath and forest, sand+grass, moss+lichen, moss+grass, lichen+heath,

heath+forest, sand+grass+moss, moss+lichen+heath, and lichen+heath+forest. Instead of

using all possible combinations, we selected the combinations to limit the number of tests and

to focus on the combinations most relevant for the aims of the study. OTUs were considered

indicator OTUs if their indicator value was >0.3 and p <0.05. Significance was evaluated

with a permutation test (999 permutations). P values were adjusted for multiple comparisons

with function p.adjust (method "fdr"). Generalist OTUs present in all successional stages

were defined at indicator values >0.5, because no p values are generated for taxa present in

all sites. The identified indicator OTUs were grouped based on their taxonomic classification

at order (bacteria) or class (fungi) level. Unclassified OTUs were included in these

comparisons by re-clustering all reads assigned to unclassified bacteria or fungi in mothur at

75% sequence similarity to represent phylum level similarity for bacteria and class level

similarity for fungi (Tedersoo *et al.*, 2014).

To evaluate responses of bacterial and fungal indicator taxa to environmental variables, we

used relative abundances of indicator OTUs grouped at order level (bacteria) and class level

(fungi), including those unclassified 75% similarity groups that represented >0.4% of all

unclassified OTUs. The tested environmental variables were those that best explained

microbial community variation (OM, pH, Al for bacteria; OM, pH, minimum temperature for

fungi). The analysis was carried out with the function traitglm in the package mvabund (v.

3.13.1, Wang *et al.*, 2012). The function fits fourth corner models to relate sites and their

traits to species and species traits. When used without species trait data, it becomes a

predictive model of responses of different OTUs to variation of environmental factors

(Brown *et al.*, 2014). The method used was glm1path, fitting a generalized linear model with

LASSO penalty, which means that fourth corner coefficients of OTUs not decreasing

Bayesian information criterion (BIC) are forced to 0, i.e. no response to environmental

variation, to simplify the model (Brown *et al.*, 2014).

All statistical analyses were carried out in R v.3.2.5. or v.3.4.0 (RCoreTeam, 2018).

Results

Soil chemistry and microbial biomass along Arctic sand dune succession

The increasing aboveground vegetation cover from bare sand to mountain birch forest was

mirrored belowground in increasing soil OM content and decreasing pH (Fig. 1; Pearson

correlation r=-0.78, p< 0.001). Organic matter content was low in the sand, grass, and moss

stage soils (0.4±0.2%, mean±SD) but increased to 0.9±0.3% in the lichen stage and up to

4.5% ± 1.7% in the forest stage (Fig. 1b, Table S1). pH was highest in the grass stage (5.8 ±

0.2) and lowest in the forest stage (4.7 ± 0.1) (Fig. 1c). Soil contents of N, P, K, Ca, Mg, Fe

and S increased with increasing OM (Table S1).

Microbial biomass measured as PLFAs per soil dry weight followed closely the OM-pH

gradient (Fig. 1e, Fig. S1). The moss stage had similar OM content as bare sand, but its

microbial biomass was higher than in the other sandy stages. When related to soil OM,

microbial biomass in the moss stage reached a level similar to the lichen and vascular plant

stages (Fig. 1d). Similarly, a shift to a higher proportion of fungal to bacterial biomass

occurred in the moss stage and was maintained during the later successional stages (Fig. 1f).

Overall, the proportion of fungi was highest in the heath stage.

Microbial community structure

Bacterial and fungal community structure based on sequencing (Fig. 2) and PLFA (Fig. S2)

followed the order of the vegetation stages on the OM-pH gradient and differed with

successional stage (PERMANOVA bacteria r^2=0.58 P<0.001, fungi r^2=0.59 p<0.001). The

moss stage grouped with the other sandy stages (sand, grass) with partly overlapping

communities, whereas the lichen stage grouped with the vascular plant stages (heath, forest)

(Fig. 2a, 2c). It should be noted that multivariate dispersion also differed between the

successional stages (betadisper p=0.007 bacteria, p=0.001 fungi). The highest bacterial OTU

richness (Fig. 1g) and Shannon diversity (Table S1) were detected in the heath stage, whereas

fungal richness and diversity were highest though very variable in bare sand (Fig. 1g, Table

S1). The soil parameter that best explained the community variation of bacteria and fungi was

the OM-pH gradient, especially in the vegetated stages (Table S2).

To compare successional shifts in bacteria, fungi, vegetation, and soil chemistry, we

calculated distance measures from bare sand to the other successional stages (Fig. 3). The

largest shift in bacterial and fungal community structure was detected from moss to lichen

stage (Fig. 3). Soil chemistry changed most markedly from heath to forest. Vegetation

showed increasingly larger shifts in later succession, with the largest shift from heath to

forest (Fig. 3). To determine if the microbial communities correlated with vegetation and soil

chemistry, we carried out Procrustes analysis (Fig. S3). The bacterial community correlated

better with soil chemistry than vegetation (bacteria-vegetation r=0.71, bacteria-soil chemistry

r=0.80, p=0.001 for both). The fungal community correlated tighter with vegetation than with

soil chemistry (fungi-vegetation r=0.77, fungi-soil chemistry r=0.71, p=0.001 for both).

When the variation due to soil chemistry was partitioned out, correlation with vegetation

remained with lower correlation coefficients (bacteria-vegetation r=0.51, fungi-vegetation

r=0.52, both p=0.001). We then used the Procrustes residuals to compare the strength of the

relationship between the plant and microbial community along the succession. The strength

of the vegetation-bacteria correlation did not vary with successional stage (Fig. 4). In

contrast, vegetation determined fungal community composition more strongly in the heath

stage than in the other stages (Fig. 4)

Bacterial taxonomic distribution with succession

Overall, the most abundant bacterial phyla or classes were Alphaproteobacteria,

Actinobacteria, and Acidobacteria (Fig. 2b, Fig. S4). The most abundant bacterial OTU

belonged to the candidate phylum WD272 (WPS-2). This phylum was the fourth or fifth most

abundant phylum in the sandy and lichen stages (on average 4.6-8.1%), decreasing in the

heath (3.5%) and forest (0.9%) (Fig. 2b, Fig. S4). When related to microbial biomass, the

abundance of WD272 increased in the moss stage (Fig. S5). WD272 showed abundant

indicator OTUs for the sandy stages but also less abundant forest-specific indicators (Fig. 5).

Most indicator groups specific for the sandy stages were strongly associated with low OM

content (Cytophagales, Deinococcales, Acidobacteria subgroup 4, Pseudonocardiales

(Actinobacteria), Oligoflexales (Deltaproteobacteria), unclassified group 9) (Fig. 5).

Cyanobacteria, three Chloroflexi groups and unclassified group 2 were indicators for the

moss and lichen stages and associated with both low OM and low pH. Another sign that the

moss stage was a key transition stage for bacterial community was that it showed unclassified

clusters typical to both sandy and vegetated stages (Fig. S6a). Most indicator groups for the

forest stage were indicators also for the other vegetated stages. For example, Acidobacteria

subgroup 2 associated with low pH and subgroup 6 associated with high OM appeared in the

moss and lichen stages and had indicators also in the heath and forest stages (Fig. 6). The

overall abundance of Acidobacteria increased with decreasing pH (Pearson correlation r = -

0.81, p<0.001).

Fungal taxonomic distribution with succession

The fungi at class level were dominated by Ascomycetes, followed by Basidiomycetes,

unclassified fungi, and smaller shares of Zygomycota, Rozellomycota, Chytridiomycota, and

Glomeromycota (Fig. 2d, Fig. S7). The most abundant Ascomycota class was Leotiomycetes,

followed by unclassified ascomycetes and Dothideomycetes. Bare sand had the highest

relative abundance of unclassified fungi up to 35% (Fig. 2d, Fig. S7), and several unclassified

clusters were specific to the sandy stages (Fig. S6b) and associated with high pH or low OM

(Fig. 6). Other strong indicators of sandy stages included Helotiales and Thelebolales

(Leotiomycetes), and Polysporales and Boletales (Agaricomycetes). Lecanoromycetes,

containing lichenized fungi, was most abundant in the moss stage (Fig. 2d). Similarly to

bacteria, the fungal groups that were the best indicators of the moss and lichen stages were

associated with low OM and low pH or both, including Lecanoromycetes (Lecanorales,

Agyriales, Ostropomycetidae), Agaricomycetes (Cantharellales, Atheliales, Agaricales), and

Dothideomycetes (Pleosporales, Capnodiales) (Fig. 6). The most abundant fungal OTU

overall and one of the fungal generalists (Table S3) was classified as *Pezoloma ericae*

(Leotiomycetes; also known as *Rhizoscyphus ericae*). This OTU was particularly dominant in

the heath (34±5% of reads). Its relative abundance was lowest in bare sand (2±1%) and the

grass stage (12±14%), but in the moss and lichen stages it increased to 26±16%. In the forest,

its relative abundance was lower than in the cryptogam-dominated stages (17±14%).

Discussion

We hypothesized that the belowground and aboveground succession in Arctic sand dunes are

not completely synchronous. Specifically, we hypothesized that the cryptogam-dominated

stages of an Arctic sand dune succession involve key belowground components that facilitate

transitions towards vascular vegetation. Supporting this hypothesis, the bacterial and fungal

communities in the lichen stage soil resembled the communities in the later, vascular plant

dominated stages. In the other cryptogam stage studied – the moss stage – the overall

microbial communities overlapped with the earlier, sandy stages. Nevertheless, indicative of

belowground transitions preceding aboveground succession, microbial biomass per OM and

the proportion of fungi increased in the moss stage, and later stage microbial groups appeared

despite the absence of their presumed host plants. The higher correlation of bacterial

community with soil chemistry than with vegetation and the increase in the strength of plant-

fungal interaction with vascular vegetation supported the hypothesis on the different

influences of vegetation succession on bacteria and fungi.

The dune sand is a harsh habitat with low carbon and nitrogen availability. The initial

microbial colonizers must be able to tolerate the oligotrophic conditions either by using the

low levels of allocthonous organic matter (Hodkinson *et al.*, 2002) or as autotrophs

(Nemergut *et al.*, 2007; Fierer *et al.*, 2010). Comparison of biomass per OM with biomass per

soil dry weight shows that the microbial biomass increased in the moss stage when OM

potentially available for heterotrophs still remained low. A characteristic group for the early

stages from bare sand to lichen was candidate phylum WD272, currently known as WPS-2

and recently named *Candidatus* Eremiobacterota, 'desert bacterial phylum', which are

proposed to be autotrophs able to use atmospheric H_2 and carbon monoxide (CO) as energy

sources (Ji *et al.*, 2017; Holland-Moritz *et al.*, 2018) . Another potential H_2- and CO-

oxidizing group in the moss and lichen stages were members of Chloroflexi classified as

Ktedonobacteria (King and King, 2014; Islam *et al.*, 2019). The increased occurrence of

WD272 and Chloroflexi in the moss and lichen stages suggests that they may form a basis of

biomass accumulation in the Arctic sand dune succession. These microbes may therefore

facilitate development of cryptogam cover and heterotrophic microbial community (Connell

and Slatyer, 1977; Brooker *et al.*, 2008).

Many moss stage microbial groups that were rare or absent in the earlier sandy stages are

linked to biological soil crusts. Biocrusts are consortia of microbes, mosses, and lichens that

are recognized as the major stabilizing factor in early successional soils worldwide (Weber *et*

al., 2016). The rhizoids of *Polytrichum* mosses themselves stabilize soil, facilitate N_2

fixation, and provide OM to the soil (Bowden, 1991). The moss stage showed the highest

proportion of Cyanobacteria, known as N_2-fixing early colonizers and components of

biocrusts. Another distinctive indicator for the cryptogam stages were Chloroflexi, which in

dryland and grassland ecosystems occur in moss and lichen biocrusts and in sub-crust soil

(Kuske *et al.*, 2012; Navarro-Noya *et al.*, 2014; Maier *et al.*, 2018). In a temperate inland

dune succession, a thick *P. piliferum* cover increased microbial biomass compared to bare

sand (Sparrius and Kooijman, 2013). In our ecosystem, *P. piliferum* biomass was small in the

moss stage but nonetheless affected the microbial biomass and communities. Therefore *P.*

piliferum appeared to have a disproportionate effect on the ecosystem level, suggesting it is a

keystone plant species in Arctic sand dunes. The mechanism could be soil stabilization by

rhizoids and moss-associated, soil crust-forming microbes. Traditionally, grasses are credited

with the initial physical stabilization of sand in dune succession (e.g. Olson, 1958; Ellenberg,

2009). Inspecting the soil microbial community structure, it seems that in these Arctic dunes

the grass *Deschampsia flexuosa* may not drive ecosystem succession. Based on our results,

we propose that *P. piliferum* could participate in the physical stabilization of Arctic inland

dunes and direct succession towards vascular vegetation. Verifying this proposition requires

further studies on *P. piliferum* with emphasis on the role of rhizoids.

Indicative of nonaligned belowground vs. aboveground succession, lichen-forming fungi

(Lecanoromycetes, Ascomycetes) were detected in the moss stage soil before the succession

stage where they formed lichens. Furthermore, the relative abundance of the dominant fungal

OTU *Pezoloma ericae* increased in the moss stage to levels similar to those in the later

stages. We interpret this increase as a possible sign of fungal community shift at the moss

stage towards a status where vascular plant establishment is possible. *Pezoloma ericae* forms

mycorrhiza with ericoid plants such as the dominant heath species *Empetrum nigrum* (Read,

1991; Smith and Read, 2008). However, no known host plants of *P. ericae* were detected in

the moss stage. In addition to ericoid mycorrhiza, *P. ericae* forms mycorrhiza-like

associations in liverwort rhizoids (Pressel *et al.*, 2010; Kowal *et al.*, 2018), and it has been

found as an Antarctic moss endophyte (Zhang *et al.*, 2013). However, no liverworts were

visible among the moss at our site, and no association of *P. ericae* with *Polytrichum* rhizoids

is known.

Supporting the hypothesis of different controls, bacteria and fungi showed distinct richness

and biomass patterns along the succession. The proportion of fungi in the PLFA data,

assumed to represent the living biomass (Zhang *et al.*, 2019), increased already in the moss

stage. Bacterial richness peaked in the heath and fungal richness in the bare sand, and neither

correlated with ecosystem productivity or community maturity defined as soil OM or plant

richness. In general, richness and diversity link to ecosystem functions through the complex

interaction between diversity and ecosystem stability and productivity (Loreau and de

Mazancourt, 2013). Plant richness peaks in the middle or late succession (Bonet and Pausas,

2004; Peyrat and Fichtner, 2011; Prach *et al.*, 2014). For microbes, few general patterns of

richness or diversity with primary succession have emerged. In very young soils, richness is

often low and then increases and stabilizes or even decreases in later stages (Jackson, 2003;

Nemergut *et al.*, 2007; Fierer *et al.*, 2010; Williams *et al.*, 2013; Jiang *et al.*, 2018). Primary

succession gradients where both bacteria and fungi have been studied show conflicting

patterns with each other but generally agree on different trajectories for bacterial and fungal

richness (Brown and Jumpponen, 2014; Cutler, *et al.*, 2014; Dini-Andreote, *et al.* 2014; Dini-

Andreote, *et al.* 2016; Jiang, *et al.* 2018). In our case, fungal richness remained rather stable

under vegetation covered stages, while bacterial richness had a unimodal response. What can

be concluded from above and the present study is that fungi and bacteria do not follow the

same trajectories during succession and are not equally shaped by the same factors.

The low bacterial richness in the bare sand stage can be related to the oligotrophic conditions

(i.e. low environmental resource availability); richness already increased with the scattered

presence of the grass *D. flexuosa*. Pioneering plants can influence bulk soil when plants are of

sufficient size (Tscherko *et al.*, 2004; Edwards *et al.*, 2006; Miniaci *et al.*, 2007). The high

fungal richness in the bare sand together with the low fungal biomass, however, is

unpredicted by theories that species richness should increase with the increase in resources

and environmental heterogeneity (Stein *et al.*, 2014; Cline *et al.*, 2018). Some of the fungal

richness in bare soil may represent dormant fungi, spores or hostless biotrophic fungi

(Jumpponen, 2003; Rime *et al.*, 2015). On the other hand, our observation that the

unclassified fungi can be grouped into class level clusters that occurred consistently in

replicate samples and across the sandy successional stages also hints that the fungal

community in bare sand may be in one way or other selected or specialized.

The high bacterial richness in the heath coincided with the highest fungal biomass and

highest occurrence of *P. ericae*, presumably forming mycorrhizae with the dominant ericoid

plant *E. nigrum*. We suppose that the high bacterial richness is promoted by resources and

niches provided by fungal mycelium and *Empetrum* litter, despite its high phenolic content

and proposed adverse properties on microbial activity (Wardle *et al.*, 1998; Gallet *et al.*,

1999). In the forest, where the highest microbial richness would be expected if richness

correlated positively with OM, plant species richness, and productivity, bacterial diversity

was lower than in the heath, possibly due to the environmental filtering effect of the low pH

in the forest soil (Tripathi *et al.*, 2018). The varied responses of bacterial and fungal richness

and diversity to primary succession highlight the need to cover successional gradients with

different parent material, vegetation and soil chemistry to understand microbial community

patterns along succession.

Effects of soil chemistry on microbial communities can override biotic factors such as plant

richness (Chen *et al.*, 2016; Jiang *et al.*, 2018), and the role of vegetation is emphasized in

successional gradients where soil chemistry does not change markedly (Knelman *et al.*, 2012;

Cutler *et al.*, 2014). Soil chemistry, vegetation and microbes are strongly intertwined in

succession, and the present gradient from bare sand to mountain birch forest represents the

common pattern where organic matter increases and the soil acidifies with increasing plant

cover (Odum, 1969; Merilä *et al.*, 2010). However, the largest microbial community shifts

did not align with the largest shifts in soil chemistry or vegetation. No measured soil

chemistry variable changed in parallel with the large bacterial community shift between moss

and lichen stages. Both cryptogam stages showed indicator microbial groups associated with

low OM and low pH. These results suggest that the microbial shift from moss to lichen stage

is due to unmeasured effects of the lichen crust formation. Such factors could be better long-

term moisture retention by the lichen crust than the sparse mosses in the moss stage,

microenvironments provided by lichen crumbs in the belowground soil (de los Ríos *et al.*,

2011) or the different chemical composition and secondary metabolites of mosses versus

lichens.

Conclusions

This study on inland sand dunes illuminates bacterial and fungal succession in relation to

vegetation and soil chemistry in a less studied Arctic soil ecosystem. Successional dynamics

may not be linear nor deterministic, which makes the space-for-time substitution in mosaics

of repeated plant communities that are thought to represent chronosequences challenging

(Walker *et al.*, 2010). In our case, the soil organic matter helps link vegetation development

in the lichen, heath and forest with time. However, the moss and grass stages are equally

devoid of soil organic matter. The microbial community composition in the grass stage

suggests that this stage is not well connected to the progressive successional pathway from

bare sand to mountain birch forest. Our results highlight the importance of considering not

only vascular plants but also cryptogams*,* i.e. mosses and lichens and particularly the

potential dune keystone species *Polytrichum piliferum*, when defining the plant-microbe

interactions in succession. Taken together, the results indicate that the belowground transition

towards the later stages starts before the establishment of vascular vegetation. This means

that the belowground succession may have a conditioning effect on the vascular plant

succession in the early primary succession. This work complements evidence that bacterial

and fungal richness are not necessarily related to soil ecosystem productivity or change in

plant species richness (Waldrop *et al.*, 2006; Porazinska *et al.*, 2018; Delgado-Baquerizo and

Eldridge, 2019). Soil chemistry and the effects of plants on microbial resources and niches

seem to be important factors driving succession. However, the key transitions in soil

chemistry, vegetation and microbial community did not align, suggesting additional drivers

such as competition between microbial taxa. Bacterial and fungal indicators identified in this

work, including novel unclassified diversity, can signpost ecosystem transitions towards

vascular plant cover. Understanding the drivers of the shift from bare to vegetated stage is

crucial for restoration of vegetation on cold-climate soils suffering from erosion and

vegetation loss under changing environment.

Acknowledgements

568   We thank Helena Jauhiainen, Riitta Nissinen and Manoj Kumar for help with sampling and

569   DNA extraction, Anita Mäki for assistance with sequencing, and Outi Manninen for the

570   vegetation inventory. This work was funded by Maj and Tor Nessling Foundation and the

571   Academy of Finland (project 287545 to MMK).

Author contributions

574   MMK conceived the research and led the field work, HJ and MM conducted PLFA analyses,

575   HJ conducted DNA work, analyzed all the data and performed the statistical analyses, MT

576   provided expertise and resources for Ion Torrent sequencing, HJ and MMK wrote the

577   manuscript, and all authors commented on the manuscript.

References

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecology **26**: 32-46.

Arróniz-Crespo M, Pérez-Ortega S, De los Ríos A, Green TA, Ochoa-Hueso R,

Casermeiro MÁ, de la Cruz, María Teresa, Pintado A, Palacios D, Rozzi R. 2014.

Bryophyte-cyanobacteria associations during primary succession in recently deglaciated areas

of Tierra del Fuego (Chile). *PloS One* **9**: e96081.

Baker WL, Walford GM. 1995. Multiple stable states and models of riparian vegetation

succession on the Animas River, Colorado. *Annals of the Association of American*

Geographers **85**: 320-338.

Bardgett RD, Walker LR. 2004. Impact of coloniser plant species on the development of
decomposer microbial communities following deglaciation. *Soil Biology and Biochemistry*
36: 555-559.

Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, Wit P,
Sánchez-García M, Ebersberger I, Sousa F. 2013. Improved software detection and
extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for
analysis of environmental sequencing data. *Methods in Ecology and Evolution* **4**: 914-919.

Blaalid R, Carlsen T, Kumar S, Halvorsen R, Ugland KI, Fontana G, Kauserud H.
2012. Changes in the root-associated fungal communities along a primary succession gradient
analysed by 454 pyrosequencing. *Molecular Ecology* **21**: 1897-1908.

Bonet A, Pausas JG. 2004. Species richness and cover along a 60-year chronosequence in
old-fields of southeastern Spain. *Plant Ecology* **174**: 257-270.

Bowden RD. 1991. Inputs, outputs, and accumulation of nitrogen in an early successional
moss (*Polytrichum*) ecosystem. *Ecological Monographs* **61**: 207-223.

Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G,
Liancourt P, Tielbörger K, Travis JM, Anthelme F. 2008. Facilitation in plant
communities: the past, the present, and the future. *Journal of Ecology* **96**: 18-34.

Brown SP, Jumpponen A. 2014. Contrasting primary successional trajectories of fungi and
bacteria in retreating glacier soils. *Molecular Ecology* **23**: 481-497.

Brown AM, Warton DI, Andrew NR, Binns M, Cassis G, Gibb H. 2014. The fourth-
corner solution – using predictive models to understand how species traits interact with the
environment. *Methods in Ecology and Evolution* **5**: 344-352.

Chapin FS, Walker LR, Fastie CL, Sharman LC. 1994. Mechanisms of primary
succession following deglaciation at Glacier Bay, Alaska. *Ecological Monographs* **64**: 149-
175.

Chen Y, Ding J, Peng Y, Li F, Yang G, Liu L, Qin S, Fang K, Yang Y. 2016. Patterns and

drivers of soil microbial communities in Tibetan alpine and global terrestrial ecosystems.

Journal of Biogeography **43**: 2027-2039.

Cline LC, Hobbie SE, Madritch MD, Buyarski CR, Tilman D, Cavender-Bares JM.

2018. Resource availability underlies the plant-fungal diversity relationship in a grassland

ecosystem. *Ecology* **99**: 204-216.

Connell JH, Slatyer RO. 1977. Mechanisms of succession in natural communities and their

role in community stability and organization. *The American Naturalist* **111**: 1119-1144.

Cornelissen JH, Lang SI, Soudzilovskaia NA, During HJ. 2007. Comparative cryptogam

ecology: a review of bryophyte and lichen traits that drive biogeochemistry. *Annals of Botany*

99: 987-1001.

Cutler NA, Chaput DL, van der Gast CJ. 2014. Long-term changes in soil microbial

communities during primary succession. *Soil Biology and Biochemistry* **69**: 359-370.

De Cáceres M, Legendre P. 2009. Associations between species and groups of sites: indices

and statistical inference. *Ecology* **90**: 3566-3574.

de los Ríos A, Raggio J, Pérez-Ortega S, Vivas M, Pintado A, Green TA, Ascaso C,

Sancho LG. 2011. Anatomical, morphological and ecophysiological strategies in *Placopsis*

pycnotheca (lichenized fungi, Ascomycota) allowing rapid colonization of recently

deglaciated soils. *Flora-Morphology, Distribution, Functional Ecology of Plants* **206**: 857-

864.

Delgado-Baquerizo M, Eldridge DJ. 2019. Cross-biome drivers of soil bacterial alpha

diversity on a worldwide scale. *Ecosystems* **22**: 1220–1231.

DeLuca TH, Zackrisson O, Nilsson M, Sellstedt A. 2002. Quantifying nitrogen-fixation in

feather moss carpets of boreal forests. *Nature* **419**: 917-920.

Dini-Andreote F, de Cássia Pereira e Silva, Michele, Triadó-Margarit X, Casamayor

EO, van Elsas JD, Salles JF. 2014. Dynamics of bacterial community succession in a salt

marsh chronosequence: evidences for temporal niche partitioning. *The ISME Journal* **8**:
1989-2001.

Dini-Andreote F, Pylro VS, Baldrian P, van Elsas JD, Salles JF. 2016. Ecological
succession reveals potential signatures of marine-terrestrial transition in salt marsh fungal
communities. *The ISME Journal* **10**: 1984-1997.

Dini-Andreote F, Stegen JC, van Elsas JD, Salles JF. 2015. Disentangling mechanisms
that mediate the balance between stochastic and deterministic processes in microbial
succession. *Proceedings of the National Academy of Sciences* **112**: E1326-E1332.

Duc L, Noll M, Meier BE, Bürgmann H, Zeyer J. 2009. High diversity of diazotrophs in
the forefield of a receding alpine glacier. *Microbial Ecology* **57**: 179-190.

Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves
sensitivity and speed of chimera detection. *Bioinformatics* **27**: 2194-2200.

Edwards I, Bürgmann H, Miniaci C, Zeyer J. 2006. Variation in microbial community
composition and culturability in the rhizosphere of *Leucanthemopsis alpina* (L.) Heywood
and adjacent bare soil along an alpine chronosequence. *Microbial Ecology* **52**: 679-692.

Egler FE. 1954. Vegetation science concepts I. Initial floristic composition, a factor in old-
field vegetation development. *Plant Ecology* **4**: 412-417.

Ellenberg H. 2009. *Vegetation ecology of Central Europe.* Cambridge: Cambridge
University Press.

Fierer N, Nemergut D, Knight R, Craine JM. 2010. Changes through time: integrating
microorganisms into the study of succession. *Research in Microbiology* **161 8**: 635-642.

Francini G, Liiri M, Männistö M, Stark S, Kytöviita M. 2014. Response to reindeer
grazing removal depends on soil characteristics in low Arctic meadows. *Applied Soil Ecology*
76: 14-25.

Gallet C, Nilsson M, Zackrisson O. 1999. Phenolic metabolites of ecological significance in
Empetrum hermaphroditum leaves and associated humus. *Plant and Soil* **210**: 1-9.

Gornall JL, Jónsdóttir IS, Woodin SJ, van der Wal R. 2007. Arctic mosses govern below-
ground environment and ecosystem processes. *Oecologia* **153**: 931-941.

Gornall JL, Woodin SJ, Jónsdóttir IS, van der Wal R. 2011. Balancing positive and
negative plant interactions: how mosses structure vascular plant communities. *Oecologia*
166: 769-782.

Hodkinson ID, Coulson SJ, Webb NR. 2003. Community assembly along proglacial
chronosequences in the high Arctic: vegetation and soil development in north-west Svalbard.
Journal of Ecology **91**: 651-663.

Hodkinson ID, Webb NR, Coulson SJ. 2002. Primary community assembly on land – the
missing stages: why are the heterotrophic organisms always there first? *Journal of Ecology*
90: 569-577.

Holland-Moritz H, Stuart J, Lewis LR, Miller S, Mack MC, McDaniel SF, Fierer N.
2018. Novel bacterial lineages associated with boreal moss species. *Environmental*
Microbiology **20**: 2625-2638.

Ihrmark K, Bödeker I, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y,
Stenlid J, Brandström-Durling M, Clemmensen KE. 2012. New primers to amplify the
fungal ITS2 region–evaluation by 454-sequencing of artificial and natural communities.
FEMS Microbiology Ecology **82**: 666-677.

Islam ZF, Cordero PRF, Feng J, Chen Y, Bay SK, Jirapanjawat T, Gleadow RM,
Carere CR, Stott MB, Chiri E et al. 2019. Two Chloroflexi classes independently evolved
the ability to persist on atmospheric hydrogen and carbon monoxide. *The ISME Journal* **13**:
1801–1813.

Jackson CR. 2003. Changes in community properties during microbial succession. *Oikos*
101: 444-448.

Ji M, Greening C, Vanwonterghem I, Carere CR, Bay SK, Steen JA, Montgomery K,
Lines T, Beardall J, van Dorst J. 2017. Atmospheric trace gases support primary
production in Antarctic desert surface soil. *Nature* **552**: 400-403.

Jiang Y, Lei Y, Yang Y, Korpelainen H, Niinemets Ü, Li C. 2018. Divergent assemblage
patterns and driving forces for bacterial and fungal communities along a glacier forefield
chronosequence. *Soil Biology and Biochemistry* **118**: 207-216.

Jones GA, Henry GHR. 2003. Primary plant succession on recently deglaciated terrain in
the Canadian High Arctic. *Journal of Biogeography* **30**: 277-296.

Jumpponen A. 2003. Soil fungal community assembly in a primary successional glacier
forefront ecosystem as inferred from rDNA sequence analyses. *New Phytologist* **158**: 569-
578.

King C, King G. 2014. Description of *Thermogemmatispora carboxidivorans* sp. nov., a
carbon-monoxide-oxidizing member of the class *Ktedonobacteria* isolated from a
geothermally heated biofilm, and analysis of carbon monoxide oxidation by members of the
class *Ktedonobacteria*. *International Journal of Systematic and Evolutionary Microbiology*
64: 1244-1251.

Knelman JE, Legg TM, O'Neill SP, Washenberger CL, González A, Cleveland CC,
Nemergut DR. 2012. Bacterial community structure and function change in association with
colonizer plants during early primary succession in a glacier forefield. *Soil Biology and*
Biochemistry **46**: 172-180.

Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST,
Bruns TD, Bengtsson-Palme J, Callaghan TM. 2013. Towards a unified paradigm for
sequence-based identification of fungi. *Molecular Ecology* **22**: 5271-5277.

Koster EA. 1988. Ancient and modern cold-climate aeolian sand deposition: a review.
Journal of Quaternary Science **3**: 69-83.

Kowal J, Pressel S, Duckett JG, Bidartondo MI, Field KJ. 2018. From rhizoids to roots?
Experimental evidence of mutualism between liverworts and ascomycete fungi. *Annals of*
Botany **121**: 221-227.

Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data

on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology* **79**:
5112-5120.

Kuske CR, Yeager CM, Johnson S, Ticknor LO, Belnap J. 2012. Response and resilience
of soil biocrust bacterial communities to chronic physical disturbance in arid shrublands. *The*
ISME Journal **6**: 886-897.

Lichter J. 1998. Primary succession and forest development oncoastal Lake Michigan sand
dunes. *Ecological Monographs* **68**: 487-510.

Lisboa FJG, Peres-Neto PR, Chaer GM, Jesus EdC, Mitchell RJ, Chapman SJ, Berbera
RLL. 2014. Much beyond Mantel: bringing Procrustes association metric to the plant and
soil ecologist's toolbox. *PLoS ONE* **9**: e101238.

Loreau M, de Mazancourt C. 2013. Biodiversity and ecosystem stability: a synthesis of
underlying mechanisms. *Ecology Letters* **16**: 106-115.

Maier S, Tamm A, Wu D, Caesar J, Grube M, Weber B. 2018. Photoautotrophic
organisms control microbial abundance, diversity, and physiology in different types of
biological soil crusts. *The ISME Journal* **12**: 1032-1046.

Mäki A, Rissanen AJ, Tiirola M. 2016. A practical method for barcoding and size-trimming
PCR templates for amplicon sequencing. *BioTechniques* **60**: 88-90.

Makoto K, Wilson SD. 2019. When and where does dispersal limitation matter in primary
succession? *Journal of Ecology* **107**: 559-565.

Måren IE, Kapfer J, Aarrestad PA, Grytnes JA, Vandvik V. 2018. Changing
contributions of stochastic and deterministic processes in community assembly over a
successional gradient. *Ecology* **99**, 148-157.

McArdle BH, Anderson MJ. 2001. Fitting multivariate models to community data: a
comment on distance-based redundancy analysis. *Ecology* **82**: 290-297.

Merilä P, Malmivaara-Lämsä M, Spetz P, Stark S, Vierikko K, Derome J, Fritze H.
2010. Soil organic matter quality as a link between microbial community structure and

vegetation composition along a successional gradient in a boreal forest. *Applied Soil Ecology* **46**: 259-267.

Miniaci C, Bunge M, Duc L, Edwards I, Bürgmann H, Zeyer J. 2007. Effects of pioneering plants on microbial structures and functions in a glacier forefield. *Biology and Fertility of Soils* **44**: 289-297.

Montgomery DR. 2007. Soil erosion and agricultural sustainability. *Proceedings of the National Academy of Sciences* **104**: 13268-13272.

Navarro-Noya YE, Jiménez-Aguilar A, Valenzuela-Encinas C, Alcántara-Hernández RJ, Ruíz-Valdiviezo VM, Ponce-Mendoza A, Luna-Guido M, Marsch R, Dendooven L. 2014. Bacterial communities in soil under moss and lichen-moss crusts. *Geomicrobiology Journal* **31**: 152-160.

Nemergut DR, Anderson SP, Cleveland CC, Martin AP, Miller AE, Seimon A, Schmidt SK. 2007. Microbial community succession in an unvegetated, recently deglaciated soil. *Microbial Ecology* **53**: 110-122.

Odum EP. 1969. The strategy of ecosystem development. *Science* **164**: 262-270.

Oksanen J, Blanchet FG, Friendly M, Kindt R, McGlinn D, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P et al. 2017. vegan: Community Ecology Package. R package version 2.4-2. https://CRAN.R-project.org/package=vegan.

Olson JS. 1958. Rates of succession and soil changes on southern Lake Michigan sand dunes. *Botanical Gazette* **119**: 125-170.

Ortiz-Álvarez R, Fierer N, de los Ríos A, Casamayor EO, Barberán A. 2018. Consistent changes in the taxonomic structure and functional attributes of bacterial communities during primary succession. *The ISME Journal* **12**: 1658-1667.

Pennanen T, Strommer R, Markkola A, Fritze H. 2001. Microbial and plant community structure across a primary succession gradient. *Scandinavian Journal of Forest Research* **16**: 37-43.

Peres-Neto PR, Jackson DA. 2001. How well do multivariate data sets match? The
advantages of a Procrustean superimposition approach over the Mantel test. *Oecologia*
129:169-178.

Peyrat J, Fichtner A. 2011. Plant species diversity in dry coastal dunes of the southern
Baltic coast. *Community Ecology* **12**: 220-226.

Pirinen P, Simola H, Aalto J, Kaukoranta J, Karlsson P, Ruuhela R. 2012.
Climatological statistics of Finland 1981–2010. *Finnish Meteorological Institute Reports* **1**:
1-96.

Poosakkannu A, Nissinen R, Männistö M, Kytöviita M. 2017. Microbial community
composition but not diversity changes along succession in arctic sand dunes. *Environmental*
Microbiology **19**: 698-709.

Porazinska DL, Farrer EC, Spasojevic MJ, Bueno de Mesquita, Clifton P, Sartwell SA,
Smith JG, White CT, King AJ, Suding KN, Schmidt SK. 2018. Plant diversity and density
predict belowground diversity and function in an early successional alpine ecosystem.
Ecology **99**: 1942-1952.

Powell JR, Karunaratne S, Campbell CD, Yao H, Robinson L, Singh BK. 2015.
Deterministic processes vary during community assembly for ecologically dissimilar taxa.
Nature Communications **6**: 8444.

Prach K, Řehounková K, Lencová K, Jírová A, Konvalinková P, Mudrák O, Študent V,
Vaněček Z, Tichý L, Petřík P. 2014. Vegetation succession in restoration of disturbed sites
in Central Europe: the direction of succession and species richness across 19 seres. *Applied*
Vegetation Science **17**: 193-200.

Pressel S, Bidartondo MI, Ligrone R, Duckett JG. 2010. Fungal symbioses in bryophytes:
new insights in the twenty first century. *Phytotaxa* **9**: 238-253.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO.
2013. The SILVA ribosomal RNA gene database project: improved data processing and web-
based tools. *Nucleic Acids Research* **41**: D590-6.

RCoreTeam. 2018. R: A language and environment for statistical computing. Vienna,
Austria: Foundation for Statistical Computing. https://www.R-project.org/

Read DJ. 1991. Mycorrhizas in ecosystems. *Experientia* **47**: 376-391.

Rime T, Hartmann M, Brunner I, Widmer F, Zeyer J, Frey B. 2015. Vertical distribution
of the soil microbiota along a successional gradient in a glacier forefield. *Molecular Ecology*
24: 1091-1108.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski
RA, Oakley BB, Parks DH, Robinson CJ et al. 2009. Introducing mothur: open-source,
platform-independent, community-supported software for describing and comparing
microbial communities. *Applied and Environmental Microbiology* **75**: 7537-7541.

Schmidt SK, Nemergut DR, Darcy JL, Lynch R. 2014. Do bacterial and fungal
communities assemble differently during primary succession? *Molecular Ecology* **23**: 254-
258.

Schmidt SK, Reed SC, Nemergut DR, Grandy AS, Cleveland CC, Weintraub MN, Hill
AW, Costello EK, Meyer AF, Neff JC et al. 2008. The earliest stages of ecosystem
succession in high-elevation (5000 metres above sea level), recently deglaciated soils.
Proceedings of the Royal Society B-Biological Sciences **275**: 2793-2802.

Seppälä M. 1995. Deflation and redeposition of sand dunes in Finnish Lapland. *Quaternary*
Science Reviews **14**: 799-809.

Smith SE, and Read DJ. 2008. *Mycorrhizal symbiosis.* London, UK: Academic Press.

Sparrius L, Sevink J, Kooijman A. 2012. Effects of nitrogen deposition on soil and
vegetation in primary succession stages in inland drift sands. *Plant and Soil* **353**: 261-272.

Sparrius LB, Kooijman AM. 2013. Nitrogen deposition and soil carbon content affect
nitrogen mineralization during primary succession in acid inland drift sand vegetation. *Plant*
and Soil **364:** 219-228.

Stein A, Gerstner K, Kreft H. 2014. Environmental heterogeneity as a universal driver of
species richness across taxa, biomes and spatial scales. *Ecology Letters* **17**: 866–880.

Tedersoo L, Bahram M, Polme S, Koljalg U, Yorou NS, Wijesundera R, Villarreal Ruiz
L, Vasco-Palacios AM, Thu PQ, Suija A et al. 2014. Global diversity and geography of soil
fungi. *Science* **346**: 1256688.

Tripathi BM, Stegen JC, Kim M, Dong K, Adams JM, Lee YK. 2018. Soil pH mediates
the balance between stochastic and deterministic assembly of bacteria. *The ISME Journal* **12**:
1072-1083.

Tscherko D, Hammesfahr U, Marx M, Kandeler E. 2004. Shifts in rhizosphere microbial
communities and enzyme activity of *Poa alpina* across an alpine chronosequence. *Soil*
Biology and Biochemistry **36**: 1685-1698.

Van der Wal R, Brooker R. 2004. Mosses mediate grazer impacts on grass abundance in
arctic ecosystems. *Functional Ecology* **18**: 77-86.

Vanwalleghem T, Gómez JA, Infante Amate J, González de Molina M, Vanderlinden K,
Guzmán G, Laguna A, Giráldez JV. 2017. Impact of historical land use and soil
management change on soil erosion and agricultural sustainability during the Anthropocene.
Anthropocene **17**: 13-29.

Vitousek PM. 1994. Potential nitrogen fixation during primary succession in Hawaii
Volcanoes National Park. *Biotropica* **26**: 234-240.

Waldrop MP, Zak DR, Blackwood CB, Curtis CD, Tilman D. 2006. Resource availability
controls fungal diversity across a plant diversity gradient. *Ecology Letters* **9**: 1127-1135.

Walker LR, Wardle DA, Bardgett RD, Clarkson BD. 2010. The use of chronosequences
in studies of ecological succession and soil development. *Journal of Ecology* **98**: 725-736.

Wang Y, Naumann U, Wright ST, Warton DI. 2012. mvabund–an R package for model-
based analysis of multivariate abundance data. *Methods in Ecology and Evolution* **3**: 471-474.

Wardle DA, Nilsson M, Gallet C, Zackrisson O. 1998. An ecosystem-level perspective of
allelopathy. *Biological Reviews* **73**: 305-319.

Weber B, Büdel B, and Belnap J, eds. 2016. Biological soil crusts: An organizing principle
in drylands. *Ecological Studies* 226, Springer.

Williams MA, Jangid K, Shanmugam SG, Whitman WB. 2013. Bacterial communities in
soil mimic patterns of vegetative succession and ecosystem climax but are resilient to change
between seasons. *Soil Biology and Biochemistry* **57**: 749-757.

Wilson JB, Gitay H, Roxburgh SH, King WM, Tangney RS. 1992. Egler's concept of
'initial floristic composition' in succession: Ecologists citing it don't agree what it means.
Oikos **64**: 591-593.

Xie C, Lou H. 2009. Secondary metabolites in bryophytes: an ecological aspect. *Chemistry*
& Biodiversity **6**: 303-312.

Zhang T, Zhang Y, Liu H, Wei Y, Li H, Su J, Zhao L, Yu L. 2013. Diversity and cold
adaptation of culturable endophytic fungi from bryophytes in the Fildes Region, King George
Island, maritime Antarctica. *FEMS Microbiology Letters* **341**: 52-61.

Zhang Y, Zheng N, Wang J, Yao H, Qiu Q, Chapman SJ. 2019. High turnover rate of free
phospholipids in soil confirms the classic hypothesis of PLFA methodology. *Soil Biology and*
Biochemistry **135**: 323-330.

Figure legends

Figure 1 (a) Six successional stages defined in deflation basins of an Arctic inland sand dune area, (b) organic matter, (c) pH, (d) microbial biomass based on phospholipid fatty acids (PLFAs) per soil organic matter (OM) content, (e) microbial biomass per gram soil dry weight (dw), (f) ratio of fungal to bacterial biomass based on PLFA, and (g) bacterial 16S rRNA gene (upper row, darker colours) and fungal ITS amplicon (lower row, lighter colours) OTU numbers in sand dune successional stages. Data are averages of four deflation basins and three soil cores per basin. Ends of whiskers represent minimum and maximum values excluding outliers, and outliers are shown as separate points. Different letters indicate significant differences at $p < 0.05$.

Figure 2 Community structure of bacteria (a) as a non-metric multidimensional scaling (NMDS) plot (stress=0.10) and (b) at phylum level (class level for Proteobacteria) taxonomical classification based on 16S rRNA gene amplicon sequencing, and community structure of fungi (c) as an NMDS plot (stress=0.13) and (d) at class-level taxonomical classification based on ITS2 region amplicon sequencing. Data points represent three soil cores from four deflation basins per sand dune successional stage. Shapes of the symbols represent separate deflation basins. In b and d, data are averages of four deflation basins and three soil cores per basin. For values of individual samples and the groups combined in 'other', see Fig. S4 and S7.

Figure 3 Average distances of successional stages from bare sand for bacterial, fungal, and plant communities and for soil chemistry. Different letters indicate significant differences at $p < 0.05$. Ends of whiskers represent minimum and maximum values excluding outliers, and outliers are shown as separate points.

Figure 4 Strength of correlation of vegetation with (a) bacterial community and (b) fungal community determined as residuals of partial Procrustes analysis when variation due to soil chemistry was partitioned out. Low values of the residuals indicate stronger correlation.

Different letters indicate significant differences between the sand dune successional stages at
p<0.05.

Figure 5 Bacterial indicator OTUs for successional stages grouped at order level. Grey
shading shows where the successional stage changes, and each group is placed in the stage
where it was the most abundant (vertical successional stage labels; for., forest). Only groups
with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows
fourth corner coefficients (coeff.) describing the association of the indicator OTU groups
with environmental variables (env. variables). Positive coefficient indicates that the microbial
group occurred at high values of the environmental factor and negative coefficients indicate
occurrence at low values of the environmental factor. Ind. OTUs, number of indicator OTUs;
rel. abund., relative abundance; OM, organic matter.

Figure 6 Fungal indicator OTUs for successional stages grouped at class level. Grey shading
shows where successional stage changes, and each group is placed in the stage where it was
the most abundant (vertical successional stage labels: lich., lichen). A, Ascomycota; B,
Basidiomycota; R, Rozellomycota; Z, Zygomycota; cls, class; ord, order; Inc. sed., Incertae
sedis. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap
on the right shows fourth corner coefficients (coeff.) describing the association of the
indicator OTU groups with environmental variables (env. variables). Positive coefficient
indicates that the microbial group occurred at high values of the environmental factor and
negative coefficients indicate occurrence at low values of the environmental factor. Ind.
OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter,
minT, minimum soil temperature.

Figure 1 (a) Six successional stages defined in deflation basins of an Arctic inland sand dune area, (b) organic matter, (c) pH, (d) microbial biomass based on phospholipid fatty acids (PLFAs) per soil organic matter (OM) content, (e) microbial biomass per gram soil dry weight (dw), (f) ratio of fungal to bacterial biomass based on PLFA, and (g) bacterial 16S rRNA gene (upper row, darker colours) and fungal ITS amplicon (lower row, lighter colours) OTU numbers in sand dune successional stages. Data are averages of four deflation basins and three soil cores per basin. Ends of whiskers represent minimum and maximum values excluding outliers, and outliers are shown as separate points. Different letters indicate significant differences at p<0.05.

Figure 2 Community structure of bacteria (a) as a non-metric multidimensional scaling (NMDS) plot (stress=0.10) and (b) at phylum level (class level for Proteobacteria) taxonomical classification based on 16S rRNA gene amplicon sequencing, and community structure of fungi (c) as an NMDS plot (stress=0.13) and (d) at class-level taxonomical classification based on ITS2 region amplicon sequencing. Data points represent three soil cores from four deflation basins per sand dune successional stage. Shapes of the symbols represent separate deflation basins. In b and d, data are averages of four deflation basins and three soil cores per basin. For values of individual samples and the groups combined in 'other', see Fig. S4 and S7.

Figure 3 Average distances of successional stages from bare sand for bacterial, fungal, and plant communities and for soil chemistry. Different letters indicate significant differences at p<0.05. Ends of whiskers represent minimum and maximum values excluding outliers, and outliers are shown as separate points.

Figure 4 Strength of correlation of vegetation with (a) bacterial community and (b) fungal community determined as residuals of partial Procrustes analysis when variation due to soil chemistry was partitioned out. Low values of the residuals indicate stronger correlation. Different letters indicate significant differences between the sand dune successional stages at p<0.05.

Figure 5 Bacterial indicator OTUs for successional stages grouped at order level. Grey shading shows where the successional stage changes, and each group is placed in the stage where it was the most abundant (vertical successional stage labels; for., forest). Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients (coeff.) describing the association of the indicator OTU groups with environmental variables (env. variables). Positive coefficient indicates that the microbial group occurred at high values of the environmental factor and negative coefficients indicate occurrence at low values of the environmental factor. Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter.

Figure 6 Fungal indicator OTUs for successional stages grouped at class level. Grey shading shows where successional stage changes, and each group is placed in the stage where it was the most abundant (vertical successional stage labels: lich., lichen). A, Ascomycota; B, Basidiomycota; R, Rozellomycota; Z, Zygomycota; cls, class; ord, order; Inc. sed., Incertae sedis. Only groups with total abundance of indicator OTUs >0.05% are shown. The heatmap on the right shows fourth corner coefficients (coeff.) describing the association of the indicator OTU groups with environmental variables (env. variables). Positive coefficient indicates that the microbial group occurred at high values of the environmental factor and negative coefficients indicate occurrence at low values of the environmental factor. Ind. OTUs, number of indicator OTUs; rel. abund., relative abundance; OM, organic matter, minT, minimum soil temperature.