Viral expression of a SERCA2a-activating PLB mutant improves calcium cycling and synchronicity in dilated cardiomyopathic hiPSC-CMs

Daniel R. Stroik[a], Delaine K. Ceholski[b], Justyna Mleczko[b], Paul F. Thanel[a], Philip A. Bidwell[a], Joseph M. Autry[a], Razvan L. Cornea[a], David D. Thomas[a,*]

[a]Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, Minneapolis, Minnesota 55455, United States
[b]Cardiovascular Research Center, Icahn School of Medicine at Mount Sinai, New York City, New York 10029, United States

[*]Corresponding author at: David D. Thomas, University of Minnesota, Dept. of Biochemistry, Molecular Biology, and Biophysics, 6-155 Jackson Hall, 321 Church St. S.E., Minneapolis, MN 55455, USA. E-mail address: ddt@umn.edu (D.D. Thomas). Tel: +1 (612) 625-0957.

Highlights

- FRET-based assays demonstrate that the combination of two PLB mutations: L31A-PLB_M (loss of inhibitory function) and I40A-PLB_M (associated with increased SERCA binding), result in a PLB_M (L31A/I40A) capable of displacing PLB_{WT}.
- Ectopic expression of the L31A/I40A-PLB_M construct in HEK293-6E cells expressing PLB_{WT} increased Ca-ATPase activity.
- In hiPSC-CMs, direct measurement of the effect(s) of virally-delivered L31A/I40A-PLB_M on SR Ca^{2+} transport showed significantly enhancement in Ca-release amplitude and removal compared to controls.
- rAAV2.L31A/I40A-PLB_M corrected irregular Ca^{2+} transients and improved Ca^{2+} transport in a hereditary dilated cardiomyopathy model in hiPSC-CMs (R14del-PLB).
- These results suggest that gene transfer of PLB_M is viable therapy for cardiac diseases associated with impaired Ca^{2+} transport and muscle contractility.

Abstract

There is increasing momentum toward the development of gene therapy for heart failure (HF), cardiomyopathy, and other progressive cardiac diseases that correlate with impaired calcium (Ca^{2+}) transport and reduced contractility. We have used FRET between fluorescently-tagged SERCA2a (the cardiac Ca^{2+} pump) and PLB (its ventricular peptide regulator) to directly test the effectiveness of loss-of-function/gain-of-binding PLB mutants (PLB_M) that were engineered to compete with the binding of inhibitory wild type PLB (PLB_{WT}). Our therapeutic strategy is to relieve PLB_{WT} inhibition of SERCA2a by utilizing the reserve adrenergic capacity of PLB to enhance non-stimulated cardiac contractility. We determined that the combination of a loss-of-inhibitory function PLB mutation (L31A) and a gain-of-binding PLB mutation (I40A) results in a novel engineered PLB_M (L31A/I40A) that effectively displaces PLB_{WT} binding to cardiac SERCA2a in HEK293-6E cells, as assessed by a structural assay based on FRET spectroscopy. Functional assays demonstrated that co-expression of L31A/I40A-PLB_M enhances SERCA Ca-ATPase activity by increasing enzyme Ca^{2+} affinity (1/K_{Ca}) in PLB_{WT}-inhibited HEK cell homogenates. For an initial assessment of physiological relevance, human induced pluripotent stem cell derived cardiomyocytes (hiPSC-CMs) from a healthy individual were used to determine the effect of PLB_M in a native SERCA2a/WT-PLB cardiac system, finding that recombinant adeno-associated virus 2 (rAAV2)-driven expression of L31A/I40A-PLB_M enhances the amplitude of SR Ca^{2+} release and the rate of SR Ca^{2+} re-uptake in cardiomyocytes. To assess therapeutic potential, a hiPSC-CM dilated cardiomyopathy model (DCM) containing PLB mutation R14del was used to characterize Ca^{2+} transients and the effect of viral-driven expression of L31A/I40A-PLB_M, demonstrating that PLB_M rescues arrhythmic Ca^{2+} transients and alleviates decreased Ca^{2+} transport in DCM cultured cells. We propose that PLB_M transgene expression is an effective gene therapy strategy for cardiomyopathies associated with impaired Ca^{2+} transport and decreased cardiac contractility.

Keywords: calcium transport, cardiomyocyte, dilated cardiomyopathy, gene therapy, phospholamban, SERCA.

Abbreviations:

Ca^{2+} (calcium)
DCM (dilated cardiomyopathy)
ER (endoplasmic reticulum)
FLT (fluorescence lifetime)
FRET (fluorescence resonance energy transfer)
GOI (gain of inhibitory functionbinding)
GFP (green fluorescent protein)
hiPSC-CM (human induced pluripotent stem cell-derived cardiomyocyte)
HEK (human embryonic kidney)
HF (heart failure)
LOI (loss of inhibitory function)
mAb (monoclonal antibody)
miRNA (microRNA)
pAb (polyclonal antibody)
PLB (phospholamban)
PLB_M (phospholamban mutant)
PLB_{WT} (wild type phospholamban)
rAAV (recombinant adeno-associated virus)
RFP (red fluorescent protein)
SERCA2a (sarcoendoplasmic reticulum calcium ATPase 2a)
SR (sarcoplasmic reticulum)

1. Introduction

Heart failure (HF) is a pathology characterized by the impaired capacity of ventricles to fill with or eject blood, constituting a leading cause of morbidity and mortality worldwide.[1] Over the past few decades, researchers have made significant progress in identifying intracellular and molecular mechanisms that are altered during HF progression. Defective sarcoplasmic reticulum (SR) Ca^{2+} transport has been identified as a potential determinant of decreased contractility in the failing hearts.[2-4] Ca^{2+} transport from the cytosol to the SR lumen in human cardiomyocytes (CM) is largely performed (~70% of Ca^{2+} ions transported) via the SR Ca-ATPase (SERCA2a).[5] Despite recent advances in device and pharmacological therapies, rates of heart disease continue to rise due to an aging population; thus, new treatment options are urgently needed. Increasing knowledge of the molecular mechanisms fundamental to cardiac disease – including HF – has expanded the therapeutic potential to target key players involved in Ca^{2+} handling, including SERCA2a.

The activity of SERCA2a is regulated by phospholamban (PLB), a 52-residue single-pass transmembrane protein expressed in the SR of cardiac muscle, which binds to SERCA2a and reduces its apparent Ca^{2+} affinity.[6] PLB is in equilibrium between homopentamers and monomers, where the oligomeric state is proposed to function as a reservoir.[7] Previous studies have identified two functional regions on the transmembrane (TM) helix of PLB.[8-10] Single-residue mutations on one side of the TM helix diminish inhibitory function without significantly affecting PLB oligomerization state(s), whereas mutations on the other side of the TM helix modulate PLB oligomerization and can lead to enhanced SERCA inhibition.[8,11] The regulation of gene expression has been investigated in healthy individuals and HF patients, and decreased SERCA-to-PLB ratio is associated with deteriorated cardiac function, indicating that SERCA/PLB stoichiometry and SERCA2a activity are potential therapeutic targets.[12-14] To this end, SERCA2a transgene expression via recombinant adeno-associated virus (rAAV) was achieved in HF animal models to increase the SERCA-to-PLB ratio in the heart. Exogenous SERCA2a expression was tolerated and significantly enhanced cardiac function in disease models by multiple metrics.[15,16] Recently, gene therapy designed to increase SERCA2a expression in the heart underwent clinical trials in patients diagnosed with end-stage HF.[17] Despite positive preliminary results in Phase 1/2a,[18] the trial did not meet its primary end goals in phase 2b and this has been ascribed to dosage constraints.[19] One factor may be the large size of SERCA2a, which limits steady-state expression levels of the gene therapy construct. Other promising strategies to increase the SERCA-to-PLB ratio include expression of miRNAs that directly target PLB expression[20-22] and oligonucleotide-based drugs[23,24], although these strategies have not yet produced an effective therapeutic agent.

In this study, we have pursued an alternative approach to activate SERCA2a by expressing a loss-of-inhibition (LOI) PLB mutant (PLB_M) that is engineered to compete with the inhibitory wild type PLB (PLB_{WT}) (Fig. 1A) via viral delivery. In this approach, the ratio between endogenous SERCA and PLB is not targeted, rather the endogenous inhibitory PLB_{WT} is competed by PLB_M to relieve SERCA2a inhibition and enhance its Ca^{2+} transport function. Thus, this strategy is designed to overcome limitations associated with SERCA2a overexpression-based gene therapy.

2. Materials and methods

2.1. Molecular biology

eGFP and tagRFP were fused to the N-terminus of human SERCA2a and human PLB, respectively. We have demonstrated that attachment of the fluorescent proteins at these sites does not interfere with SERCA activity or PLB inhibition.[25,26] PLB cDNA mutations were introduced using the QuikChange mutagenesis kit (Agilent Technologies, Santa Clara, CA), and all expression plasmids were sequenced for confirmation.

2.2. Cell culture

Human embryonic kidney cells 293 (HEK293-6E, NRC, Canada) were cultured in FreeStyle F17 expression medium (Thermo Fisher Scientific, Waltham, MA) supplemented with 2 mM L-glutamine (Invitrogen, Waltham, MA). Cell cultures were maintained in a circular shaker (125 rpm, 37°C, 5% CO_2 (Forma Series II Water Jacket CO_2 Incubator, Thermo Fisher Scientific, Waltham, MA). For displacement assays, HEK293-6E cells were transiently transfected using 293fectamine with GFP-SERCA2a, RFP-PLB_{WT}, and PLB_M or empty vector in a 1:7:7 molar ratio. Cells were then assayed 48 hours post-transfection.

The control hiPSC line (SKiPS-31.3) was cultured as previously described.[27] The R14del hiPSC line was derived from the SKiPS-31.3 line using homologous recombination via CRISPR/Cas9. Monolayer cardiac differentiation was performed as described,[27] yielding beating cardiomyocytes within 7-10 days.

2.3. Gene expression

RNA was extracted from uninfected and rAAV-infected hiPSC-CMs at day 25, and cDNA was synthesized as previously described.[27] Gene expression compared to the housekeeping gene β2-microglobulin was determined using qRT-PCR, as assessed by $\Delta\Delta C_t$ analysis. See Supplemental Table 1 for the list of primers.

2.4. Immunoblot analysis

Samples were separated on a 4-20% polyacrylamide gradient gel (Bio-Rad, Hercules, CA) and transferred to polyvinylidene difluoride (PVDF) membrane. The PVDF membrane was blocked in Odyssey Blocking Buffer (LI-COR Biosciences, Lincoln, NE) followed by overnight incubation at 4 °C with the primary target antibody: rabbit anti-GFP polyclonal antibody (pAb) (1:1000; ab290, Abcam, Cambridge, United Kingdom), mouse anti-SERCA2 monoclonal antibody (mAb) (1:1000; 2A7-A1, Abcam), rabbit anti-tagRFP pAb (1:1000; ab233, Evrogen), mouse anti-PLB mAb (1:1000, 2D12, Abcam), or rabbit anti-β-actin pAb (1:5000, ab8227, Abcam). Blots were incubated with anti-mouse or anti-rabbit secondary antibodies conjugated to IRDye 680RD or IRDye 800CW, respectively, for 1 h at 23 °C (1:20,000; LI-COR Biosciences). Blots were quantified on the Odyssey scanner (LI-COR Biosciences).

2.5. Fluorescence data acquisition and analysis

Fluorescence lifetime (FLT) measurements were conducted in a top-read FLT plate reader designed and built by Fluorescence Innovations, Inc. (St. Paul, MN) in 384-well plate formats. GFP donor fluorescence was excited with a 473 nm microchip laser from Concepts Research Corporation (Belgium, WI), and emission was acquired with 490 nm long-pass and 520/17 nm band-pass filters (Semrock, Rochester, NY). We previously validated the performance of this FLT plate reader with known fluorescence standards, as well as with a FRET-based high-throughput screening strategy that that targeted 2-color SERCA.[28,29] FLT waveforms from donor- and donor/acceptor-labeled samples were analyzed as described in our previous publications and the supplemental methods.[28,29]

2.6. Ca-ATPase activity assay

An enzyme-coupled, NADH-linked ATPase assay was used to measure Ca^{2+}-activated ATPase activity of SERCA in 96-well microplates. Each well contained HEK293-6E homogenates, 50 mM MOPS (pH 7.0), 100 mM KCl, 1 mM EGTA, 0.2 mM NADH, 1 mM phosphoenol pyruvate, 10 IU/mL of pyruvate kinase, 10 IU/mL of lactate dehydrogenase, 3.5 μg/mL of the Ca^{2+} ionophore A23187, and $CaCl_2$ added to set $[Ca^{2+}]_i$ to the desired values.[11] The assay was started by addition of Mg-ATP at a final concentration of 5 mM, and well absorbances were read in a SpectraMax 384 Plus microplate spectrophotometer (Molecular Devices, Sunnyvale, CA). The Ca-ATPase assays were conducted over a range of $[Ca^{2+}]_i$, and the ATPase activities were fitted using the Hill function

$$V = \frac{V_{max}}{1 + 10 - \left(n(pK_{Ca} - pCa)\right)}$$

Eq. (1)

where V is the initial rate of Ca^{2+}-dependent ATPase activity at a specified Ca^{2+} concentration (pCa), V_{max} is the rate of Ca-ATPase activity at saturating $[Ca^{2+}]_i$, n is the Hill coefficient, pK_{Ca} is the fitted Ca^{2+} dissociation constant, and pCa is the concentration of ionized Ca^{2+} per specific well and V.

2.7. Live-cell Ca^{2+} transient measurements

hiPSC-CMs (day 25 of differentiation) were enzymatically dissociated using the Detach 2 kit (Promocell, Heidelberg, Germany) and plated on Matrigel-coated German glass coverslips. After 2 days, the plated hiPSC-CMs were loaded with a Ca^{2+}-sensitive fluorescent dye (Fura-2 AM, cell permeant, ThermoFisher, Rockville,

MD, USA), and the ratio of fluorescence intensities (excitation ratio of 340/380 nm) were recorded using the IonOptix system (Ionoptix, Milton, MA). The electrically-induced Ca^{2+} transients were triggered by pulses from a MyoPacer (IonOptix, Milton, MA) at 40 V and 0.5 Hz, with cells at 23 °C. Ca^{2+} traces were analyzed using IonWizard software (IonOptix) to calculate the release amplitude (peak height relative to baseline) and tau (time of Ca^{2+} removal). The number of irregular Ca^{2+} transients was quantified using IonOptix software.

Using rAAV2.L31A, rAAV2.I40A, and rAAV2.L31A/I40A PLB viruses, hiPSC-CMs were infected at day 20 of differentiation (1e4 MOI), enzymatically dissociated on day 23, and plated on Matri-gel coated German glass coverslips, as described above. Fura-2 fluorescence measurements were recorded in AAV-infected hiPSC-CMs on day 25 and compared to non-infected control hiPSC-CMs. AAV viruses were purified according to the two-plasmid method with iodixanol gradient as described.[30]

2.8. Statistical analyses

All statistics were performed using Prism (GraphPad, La Jolla, CA), and analysis was done by one-way ANOVA followed by the Bonferroni post hoc test; analysis of two group comparisons was done by Student's t-test (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, #$P < 0.0001$, and 'ns' is not significant). Data is presented as mean ± standard error of the mean (SEM), and all statistical values were calculated from a minimum of three separate experiments.

3. Results

3.1. FRET assay demonstrates SERCA-binding competition between PLB_{WT} and loss-of-function PLB_M

Previously, two classes of point mutations within the PLB TM domain had been identified via alanine-scanning mutagenesis with disparate functional outcomes: loss-of-inhibitory function (LOI) and gain-of-inhibitory function (GOI).[8] We hypothesized that combining LOI and GOI mutations in a single PLB would result in a LOI-PLB that can effectively compete with the inhibitory PLB_{WT}. To test this, we used a SERCA2a-PLB biosensor system that consists of co-expressed GFP-tagged SERCA2a and RFP-tagged PLB_{WT} in HEK293-6E cells and a fluorescence lifetime (FLT) readout of FRET. FLT-FRET is used to resolve changes in SERCA2a-PLB complex structure and binding.[28] We varied the ratio between the donor molecule (GFP-SERCA2a) and acceptor molecule (RFP-PLB_{WT}), and found that the maximal energy transfer efficiency E (fractional decrease of the fluorescence lifetime) in this live-cell based system saturates at 0.10, as previously reported.[28] We transfected cells expressing GFP-SERCA2a and RFP-PLB_{WT} with either untagged PLB_{WT} or PLB_M containing TM mutations (L31A, I40A, or L31A/I40A). Displacement of the RFP-PLB_{WT} from its interaction with GFP-SERCA2a was observed as a decrease in the FRET efficiency relative to that measured in control cells expressing only the GFP-SERCA/RFP-PLB_{WT} donor-acceptor pair (Fig. 1B). FRET between GFP-SERCA2a and RFP-PLB_{WT} decreased significantly upon co-expression of unlabeled PLB_{WT}, indicating that the RFP-PLB_{WT} and unlabeled PLB_{WT} are in equilibrium for hetero-dimeric binding to GFP-SERCA2a. An alanine substitution at L31A (LOI mutation) did not significantly alter the FRET value relative to the wild-type control. In contrast, an alanine substitution at I40A (associated with increased SERCA inhibition via decreased Ca^{2+} sensitivity) further decreased FRET, indicating that I40A-PLB_M competes effectively with RFP-PLB_{WT} with a potency (affinity) comparable to or greater than that of unlabeled PLB_{WT}. The combination of L31A and I40A mutations resulted in a PLB_M with potency similar to that of I40A-PLB_M, supporting our hypothesis that PLB_M with both mutation types retains high affinity toward SERCA2a.

Expression of the constructs was confirmed by SDS-PAGE and immunoblot in "boiled" samples, which disrupts PLB pentamers, vs. non-"boiled" heated samples, which maintains the pentameric PLB state (Fig. 1C, D). GFP-tagged SERCA2a was resolved from endogenous SERCA2b, as verified by binding of a GFP-specific antibody. There were no bands of lower mobility and no apparent aggregation in intact cells. Expression of RFP-PLB produced a single band recognized by RFP- and PLB-specific antibodies. The I40A mutation disrupts pentamer formation, and we observed the disappearance of the band corresponding to PLB_M pentamer when the I40A mutation is present (Fig. 1D; lanes 11 and 12). PLB pentamer was detected for PLB_{WT} and L31A-PLB (Fig. 1D; lanes 13 and 14). As the expression of GFP-SERCA2a, RFP-PLB_{WT}, and PLB_M is not significantly different between respective samples, the observed changes in FRET are probably due to displacement of the RFP-tagged PLB_{WT} via competition (to GFP-SERCA2a) by untagged PLB_M. The Ca-

ATPase function of GFP-SERCA is similar to untagged SERCA, so the tag does not disrupt endogenous function.[26]

3.2. Effects of loss-of-function and gain-of-binding PLB mutants on SERCA2a regulation

The effects of PLB_M overexpression on the apparent Ca^{2+} affinity of SERCA2a were determined in HEK293-6E homogenate samples, enabling co-expression of PLB_{WT} and PLB_M in a cell-based system similar to the FRET assays. Ca-ATPase activity in HEK293-6E cells expressing SERCA2a alone (Fig. 2A, black) increases with Ca^{2+} at physiological Ca^{2+} concentrations (*e.g.,* between pCa 7 and pCa 6) and saturates at higher Ca^{2+} concentrations (pCa 5), consistent with previous measurements.[28] Concomitant expression of PLB_{WT} (Fig. 2A,B, dark gray) decreases the apparent Ca^{2+} affinity of SERCA2a, and the additional expression of PLB_{WT} (light blue) or L31A-PLB_M (light gray) does not significantly decrease the apparent Ca^{2+} affinity; indicating that SERCA2a is fully inhibited and that L31A-PLB_M is not sufficient to relieve PLB_{WT} inhibition under these experimental conditions. We observed a further decrease in the apparent Ca^{2+} affinity upon co-expression of SERCA2a, PLB_{WT}, and I40A-PLB_M (red), suggesting that I40A-PLB_M binds to SERCA2a in the presence of PLB_{WT} (consistent with FRET measurements) and acts as a super-inhibitor. Co-expression with L31A/I40A-PLB_M (green) increased Ca^{2+} affinity to values similar to the SERCA-only sample, indicating that the PLB inhibition was relieved. Taken together, the structural (FRET) and functional (Ca-ATPase) assay results demonstrate that L31A/I40A-PLB_M is non-inhibitory and binds to SERCA2a in HEK293-6E cells.

3.3. rAAV-driven expression of L31A/I40A-PLB_M enhances Ca^{2+} transients in hiPSC-CMs

We differentiated hiPSCs derived from a healthy individual (SKiPS-31.3 line) into cardiomyocytes using established protocols that yields a predominant ventricular-like population.[27] To assess physiological relevance in this native SERCA-PLB cardiac system, we tested the effects of rAAV-delivered PLB_M by infecting hiPSC-CMs with empty virus, rAAV2.L31A-, rAAV2.I40A-, or rAAV2.L31A/I40A-PLB_M at day 20 of differentiation and recording Ca^{2+} transients via fura-2AM upon electrical stimulation at day 25 (Fig. 3A). Ca^{2+} transients were regular in appearance (Fig. 3B). hiPSC-CMs expressing rAAV2.L31A-PLB_M or rAAV2.I40A-PLB_M did not show significant changes in peak amplitude or the rate of Ca^{2+} removal relative to control cells expressing empty vector (Fig. 3C). However, there were striking improvements in Ca^{2+} transient parameters (increased peak amplitude, increased Ca^{2+} removal rate) in hiPSC-CMs infected with rAAV2.L31A/I40A-PLB_M (Fig. 3B and C). Expression of the exogenous PLB_M was confirmed by qRT-PCR where virally-infected hiPSC-CMs had approximately two PLB_M per one PLB_{WT} (Fig. 3D). We observed no differences in SERCA2a or cardiac Troponin T expression levels (Fig. 3D).

3.4. Relief of SERCA2a inhibition by PLB_M rescues irregular Ca^{2+} transients in cardiomyopathic R14del-PLB hiPSC-CMs

Autosomal dominant mutations in PLB have been linked to DCM and these mutations include R9C[31], R14del[32-34], R25C[35], and L39stop[36]. Recent work using hiPSC-CMs derived from patients heterozygous for the R14del-PLB mutation show mislocalization of R14del-PLB into aggregates leading to higher levels of autophagy and accompanying irregular Ca^{2+} transients.[37,38] Mice heterozygous for the R14del-PLB mutation have severe defects in SERCA2a activity and Ca^{2+} transport. We sought to characterize the effects of viral expression of PLB_M in this dilated cardiomyopathic model to evaluate therapeutic potential. To accomplish this, we generated a R14del-PLB knock-in hiPSC line using homologous recombination via CRISPR/Cas9 to insert the R14del-PLB mutation into the control SKiPS-31.3 line (Fig. 3), producing an isogenic control to test our therapies.[27] The R14del-PLB hiPSC-CMs exhibited an arrhythmic (irregular) Ca^{2+} transient profile [37] after day 35 of differentiation, characterized by a failure to respond to electric stimulation. This irregularity occurred at a frequency of 58 ± 19% of Ca^{2+} transient recorded under basal conditions. (n = 85 cells) (Fig. 4A/D). Upon infection with rAAV2.L31A/I40A-PLB, we observed an improvement in Ca^{2+} handling properties of the R14del-PLB hiPSC-CMs. Representative Ca^{2+} tracings show a significant reduction in irregular Ca^{2+} transients and improvements in Ca^{2+} amplitude and tau upon rAAV2.L31A/I40A-PLB infection (Fig. 4B/C). Finally, no irregular Ca^{2+} transients (n = 20 cells) were observed after infection with rAAV2.L31A/I40A-PLB (Fig. 4D). Altogether, these data indicate an impairment in Ca^{2+} cycling in R14del-PLB hiPSC-CMs that is corrected by exogenous expression of the SERCA activating L31A/I40A-PLB_M.

4. Discussion

We demonstrated previously that the interaction between the cardiac Ca^{2+} pump, SERCA2a, and its inhibitor, PLB, can be measured via FRET in HEK293 cell system.[28] In the present study, we applied this FRET assay to measure the competition of non-inhibitory PLB_M and RFP-labeled PLB_{WT} for binding to the GFP-SERCA2a. We used this assay to measure the relative affinity of non-inhibitory PLB_M and identified a double mutant (L31A/I40A) that binds with affinity greater than PLB_{WT}. Expression of L31A/I40A-PLB_M increased Ca-ATPase activity in PLB_{WT}-expressing HEK293 cells. Mechanistically, this suggests that L31A/I40A-PLB_M enhances SERCA activity by competitively displacing the inhibitory form of PLB. There were previous attempts to increase SR Ca-influx in HF-animal models by expressing non-inhibitory PLB_M, including a pseudo-phosphorylated mutant (S16E)[39] and a dominant-negative mutant (R3E/R14E)[40]. Although there were significant improvements in Ca transients and cardiac output, these mutations preclude PLB phosphorylation by PKA and β-adrenergic stimulation.[39,40] In hiPSC-CMs, we found that rAAV2-dependent expression of L31A/I40A-PLB_M significantly improved Ca^{2+} release amplitude and removal in both healthy and pathogenic cell lines. This work paves the way for a potential therapeutic for heart diseases associated with impaired Ca^{2+} transport and decreased contractility.

Acknowledgments

This study was supported by National Institutes of Health grants to D.D.T. (GM27906, HL129814, and AG26160). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank Bengt Svensson for help with figures. Spectrophotometric assays were performed in the Biophysical Technology Center at the University of Minnesota Department of Biochemistry, Molecular Biology, and Biophysics.

Fig. 1. Co-expression of SERCA, PLB, and PLB$_M$. (A): Schematic of gene therapy strategy: a tight-binding, non-inhibitory PLB mutant (PLB$_M$, dark green) displaces inhibitory RFP-PLB (PLB$_{WT}$, dark red), resulting in a FRET decrease. (B) FRET assay of PLB$_M$ competition in HEK293-6E. GFP-SERCA-RFP-PLB$_{WT}$ FRET (binding, *white bar*) is decreased (arrow) by PLB variants. LOI = Loss-of-inhibitory function mutant. GOI = Gain-of-inhibition mutant. (C/D) Immunoblots of HEK homogenate samples incubated at 95°C ("boiled", lanes 1-7) or at 70°C (lanes 8–14) from untransfected HEK293-6E cells (lane 1&8), cells expressing GFP-SERCA2a (lanes 2&9), or cells expressing GFP-SERCA2a+RFP-PLB (lanes 3–7&11–14) +L31A/I40A-PLB (lanes 4&11), +I40A-PLB (lanes 5&12), +L31A-PLB (lanes 6&13), and +WT-PLB (lanes 7&14). Primary antibodies used (from top to bottom of 1D panels) are anti-GFP, anti-SERCA2, anti-RFP, anti-PLB, and anti-β-actin. Quantification of steady-state expression levels by densitometry for boiled samples (lanes 1–7). Error bars indicate SEM (n=3). *P < 0.05. *JMCC note: this is a 2-column wide figure and it should use color in print*

Fig. 2. SERCA activation by PLB$_M$ competition. (A) Normalized Ca-ATPase activity of HEK293-6E cells was measured 48 hours after transfection using +empty vector (negative control, not shown), +WT-SERCA2a (black), +WT-SERCA2a/WT-PLB (dark gray, dotted line), +WT-SERCA2a/WT-PLB +WT-PLB (blue), +WT-SERCA2a/WT-PLB +L31A-PLB (light grey), WT-SERCA2a/WT-PLB +I40A-PLB (red), or WT-SERCA2a/WT-PLB +L31A/I40A-PLB (green). (B) Quantification of apparent Ca^{2+} affinity (1/pK$_{Ca}$) and PLB$_M$-induced change (Δ) from (A). Error bars indicate SEM (n = 3). *P < 0.05. *JMCC note: this is a 2-column wide figure, and it should use color in print.*

Fig. 3. In normal, healthy hiPSC-CMs, rAAV2-driven expression of L31A/I40A-PLB enhances Ca^{2+} release amplitude and Ca^{2+} removal rate. (A) Schematic of experimental design for hiPSC differentiation, transduction, and cytosolic Ca^{2+} assay. (B) Representative Ca^{2+} transients in control hiPSC-CMs *(left)* compared to rAAV2-L31A/I40A PLB transduced hiPSC-CMs *(right)*. (C) Quantification of Ca^{2+} release amplitude (peak height) normalized to uninfected control (CTRL) and Ca^{2+} removal time constant (Tau) of hiPSC-CMs normalized to uninfected control (CTRL). Error bars indicate SEM (n = 3). *$P < 0.05$. (D) qRT-PCR analysis of gene expression of Ca^{2+} transport and contractility proteins: SERCA2a *(left)*, cardiac troponin T *(center)*, and PLB *(right;* black = endogenous PLB$_{WT}$, gray = exogenous PLB$_{M}$) of uninfected (CTRL) or infected (rAAV2-L31A/I40A PLB) hiPSC-CMs. Gene expression values are normalized to uninfected control. ***JMCC note: this is a 2-column wide figure, and it should use color in print.***

Fig. 4. In cardiomyopathic R14del hiPSC-CMs, viral delivery of L31A/I40A-PLB rescues arrhythmogenic Ca^{2+} transients and enhances cellular Ca^{2+} cycling. (A) Representative Ca^{2+} transients in R14del PLB hiPSC-CMs (left) andR14del hiPSC-CMs infected with rAAV2.L31A/I40A-PLB (right). Arrhythmogenic Ca^{2+} transients are identified by red arrow. (B) Ca^{2+} release peak amplitude (fluorescence intensity) normalized to uninfected R14del hiPSC-CMs. (C) Ca^{2+} re-uptake time constant (Tau) normalized to uninfected R14del hiPSC-CMs. (D) Irregular Ca^{2+} transients determined by quantifying the number of individual cells exhibiting irregular Ca^{2+} transients. Error bars indicate SEM (n = 3). *P < 0.05. *JMCC note: this is a 2-column wide figure, and it should use color in print.*

Supplementary Methods:

Fluorescence Data Acquisition and Analysis[28]

The measured FLT waveform

$$I(t) = \int_{-\infty}^{\infty} IRF(t - t') \cdot F(t')dt' \qquad \text{Eq. (2)}$$

is a function of the nanosecond decay time t, and is modeled as the convolution integral of the measured instrument response function, IRF(t), and the fluorescence decay model, *F*(t). The fluorescence decay model

$$F_{D+A}(t) = x_D F_D(t) + (1 - x_D) F_{DA}(t) \qquad \text{Eq. (3)}$$

is a linear combination of a donor-only fluorescence decay function F_D(t) and an energy transfer-decreased donor fluorescence decay F_{DA}(t). The donor decay F_D(t) is a sum of *n* exponentials

$$F_D(t) = \sum_{i=1}^{n} A_i \, exp(-t/\tau_i) \qquad \text{Eq. (4)}$$

with discrete FLT species τ_i and pre-exponential mole fractions A_i. For the GFP donor, two exponentials (*n* = 2) are required to fit the observed fluorescence decay waveform. The energy transfer-decreased donor decay function, F_{DA}(t),

$$F_{DA}(t) = \sum_{j=1}^{N} X_j \cdot T_j(t) \qquad \text{Eq. (5)}$$

is the sum of the distribution of multiple structural states (*j*) with mole fractions X_j, represented by the FRET-quenched donor fluorescence decays T_j(t). The increase in the donor decay rate (inverse donor FLT) due to FRET is given by the Förster equation

$$k_{Ti} = k_{Di}(R/R_{0i})^{-6}, \text{ where} \qquad \text{Eq. (6)}$$

$$k_{DAi} = k_{Di} + k_{Ti}, \text{ and} \qquad \text{Eq. (7)}$$

$$k_{Di} = 1/\tau_i \qquad \text{Eq. (8)}$$

Supplementary Table 1. Primers used in qPCR.

Gene	Forward Primer (5`→3`)	Reverse Primer (5`→3`)
SERCA2a	CTGTCCATGTCACTCCACTTCC	AGCGGTTACTCCAGTATTGCAG
Cardiac TnT	AAGAGGCAGACTGAGCGGGAAA	AGATGCTCTGCCACAGCTCCTT
Endogenous PLB	ACAGCTGCCAAGGCTACCTA	GCTTTTGACGTGCTTGTTGA
Exogenous PLB (AAV-expressed)	TCAATTACGGGGTCATTAGTTC	ACTAATACGTAGATGTACTGCC
B2M	GGGATCGAGACATGTAAGCAG	CAAGCAAGCAGAATTTGGAA

References

Writing Group, M. *et al.* Heart Disease and Stroke Statistics-2016 Update: A Report From the American Heart Association. *Circulation* **133**, e38-360, doi:10.1161/CIR.0000000000000350 (2016).

Marks, A. R. Calcium cycling proteins and heart failure: mechanisms and therapeutics. *J Clin Invest* **123**, 46-52, doi:10.1172/JCI62834 (2013).

Ablorh, N. A. & Thomas, D. D. Phospholamban phosphorylation, mutation, and structural dynamics: a biophysical approach to understanding and treating cardiomyopathy. *Biophysical reviews* **7**, 63-76, doi:10.1007/s12551-014-0157-z (2015).

Gorski, P. A., Ceholski, D. K. & Hajjar, R. J. Altered myocardial calcium cycling and energetics in heart failure--a rational approach for disease treatment. *Cell Metab* **21**, 183-194, doi:10.1016/j.cmet.2015.01.005 (2015).

Bers, D. M. Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol* **70**, 23-49, doi:10.1146/annurev.physiol.70.113006.100455 (2008).

MacLennan, D. H. & Kranias, E. G. Phospholamban: a crucial regulator of cardiac contractility. *Nat Rev Mol Cell Biol* **4**, 566-577, doi:10.1038/nrm1151 (2003).

Robia, S. L. *et al.* Forster transfer recovery reveals that phospholamban exchanges slowly from pentamers but rapidly from the SERCA regulatory complex. *Circ Res* **101**, 1123-1129 (2007).

Kimura, Y., Kurzydlowski, K., Tada, M. & MacLennan, D. H. Phospholamban inhibitory function is activated by depolymerization. *J Biol Chem* **272**, 15061-15064, doi:10.1074/jbc.272.24.15061 (1997).

Cornea, R. L., Jones, L. R., Autry, J. M. & Thomas, D. D. Mutation and phosphorylation change the oligomeric structure of phospholamban in lipid bilayers. *Biochemistry* **36**, 2960-2967, doi:10.1021/bi961955q (1997).

Simmerman, H. K., Kobayashi, Y. M., Autry, J. M. & Jones, L. R. A leucine zipper stabilizes the pentameric membrane domain of phospholamban and forms a coiled-coil pore structure. *J Biol Chem* **271**, 5941-5946, doi:10.1074/jbc.271.10.5941 (1996).

Cornea, R. L., Autry, J. M., Chen, Z. & Jones, L. R. Reexamination of the role of the leucine/isoleucine zipper residues of phospholamban in inhibition of the Ca2+ pump of cardiac sarcoplasmic reticulum. *J Biol Chem* **275**, 41487-41494, doi:10.1074/jbc.M008195200 (2000).

Hasenfuss, G. *et al.* Relation between myocardial function and expression of sarcoplasmic reticulum Ca(2+)-ATPase in failing and nonfailing human myocardium. *Circ Res* **75**, 434-442 (1994).

Mercadier, J. J. *et al.* Altered sarcoplasmic reticulum Ca2(+)-ATPase gene expression in the human ventricle during end-stage heart failure. *J Clin Invest* **85**, 305-309, doi:10.1172/JCI114429 (1990).

Nagai, R. *et al.* Regulation of myocardial Ca2+-ATPase and phospholamban mRNA expression in response to pressure overload and thyroid hormone. *Proc Natl Acad Sci U S A* **86**, 2966-2970, doi:10.1073/pnas.86.8.2966 (1989).

Schmidt, U. *et al.* Restoration of diastolic function in senescent rat hearts through adenoviral gene transfer of sarcoplasmic reticulum Ca(2+)-ATPase. *Circulation* **101**, 790-796 (2000).

Byrne, M. J. *et al.* Recirculating cardiac delivery of AAV2/1SERCA2a improves myocardial function in an experimental model of heart failure in large animals. *Gene Ther* **15**, 1550-1557, doi:10.1038/gt.2008.120 (2008).

Jessup, M. *et al.* Calcium Upregulation by Percutaneous Administration of Gene Therapy in Cardiac Disease (CUPID): a phase 2 trial of intracoronary gene therapy of sarcoplasmic reticulum Ca2+-ATPase in patients with advanced heart failure. *Circulation* **124**, 304-313, doi:10.1161/CIRCULATIONAHA.111.022889 (2011).

Zsebo, K. *et al.* Long-term effects of AAV1/SERCA2a gene transfer in patients with severe heart failure: analysis of recurrent cardiovascular events and mortality. *Circulation research* **114**, 101-108, doi:10.1161/CIRCRESAHA.113.302421 (2014).

Greenberg, B. *et al.* Prevalence of AAV1 neutralizing antibodies and consequences for a clinical trial of gene transfer for advanced heart failure. *Gene Ther* **23**, 313-319, doi:10.1038/gt.2015.109 (2016).

Andino, L. M. *et al.* AAV-mediated knockdown of phospholamban leads to improved contractility and calcium handling in cardiomyocytes. *J Gene Med* **10**, 132-142, doi:10.1002/jgm.1131 (2008).

Grobetal, T. *et al.* A novel artificial microRNA expressing AAV vector for phospholamban silencing in cardiomyocytes improves Ca2+ uptake into the sarcoplasmic reticulum. *PLoS One* **9**, e92188, doi:10.1371/journal.pone.0092188 (2014).

Bish, L. T. *et al.* Cardiac gene transfer of short hairpin RNA directed against phospholamban effectively knocks down gene expression but causes cellular toxicity in canines. *Hum Gene Ther* **22**, 969-977, doi:10.1089/hum.2011.035 (2011).

Soller, K. J. *et al.* Rheostatic Regulation of the SERCA/Phospholamban Membrane Protein Complex Using Non-Coding RNA and Single-Stranded DNA oligonucleotides. *Sci Rep* **5**, 13000, doi:10.1038/srep13000 (2015).

Soller, K. J., Yang, J., Veglia, G. & Bowser, M. T. Reversal of Phospholamban Inhibition of the Sarco(endo)plasmic Reticulum Ca2+-ATPase (SERCA) Using Short, Protein-interacting RNAs and Oligonucleotide Analogs. *J Biol Chem* **291**, 21510-21518, doi:10.1074/jbc.M116.738807 (2016).

Gruber, S. J. *et al.* Discovery of enzyme modulators via high-throughput time-resolved FRET in living cells. *Journal of biomolecular screening* **19**, 215-222, doi:10.1177/1087057113510740 (2014).

Bidwell, P., Blackwell, D. J., Hou, Z., Zima, A. V. & Robia, S. L. Phospholamban binds with differential affinity to calcium pump conformers. *J Biol Chem* **286**, 35044-35050, doi:10.1074/jbc.M111.266759 (2011).

Ceholski, D. K. *et al.* Functional and transcriptomic insights into pathogenesis of R9C phospholamban mutation using human induced pluripotent stem cell-derived cardiomyocytes. *J Mol Cell Cardiol* **119**, 147-154, doi:10.1016/j.yjmcc.2018.05.007 (2018).

Stroik, D. R. *et al.* Targeting protein-protein interactions for therapeutic discovery via FRET-based high-throughput screening in living cells. *Sci Rep* **8**, 12560, doi:10.1038/s41598-018-29685-z (2018).

Schaaf, T. M. *et al.* High-Throughput Spectral and Lifetime-Based FRET Screening in Living Cells to Identify Small-Molecule Effectors of SERCA. *SLAS Discov* **22**, 262-273, doi:10.1177/1087057116680151 (2017).

Lopez-Gordo, E., Kohlbrenner, E., Katz, M. G. & Weber, T. AAV Vectors for Efficient Gene Delivery to Rodent Hearts. *Methods Mol Biol* **1950**, 311-332, doi:10.1007/978-1-4939-9139-6_19 (2019).

Schmitt, J. P. *et al.* Dilated cardiomyopathy and heart failure caused by a mutation in phospholamban. *Science* **299**, 1410-1413, doi:10.1126/science.1081578 299/5611/1410 [pii] (2003).

Haghighi, K. *et al.* A mutation in the human phospholamban gene, deleting arginine 14, results in lethal, hereditary cardiomyopathy. *Proc Natl Acad Sci U S A* **103**, 1388-1393, doi:0510519103 [pii] 10.1073/pnas.0510519103 (2006).

Karakikes, I. *et al.* Correction of human phospholamban R14del mutation associated with cardiomyopathy using targeted nucleases and combination therapy. *Nat Commun* **6**, 6955, doi:10.1038/ncomms7955 (2015).

van der Heijden, J. F. & Hassink, R. J. The phospholamban p.Arg14del founder mutation in Dutch patients with arrhythmogenic cardiomyopathy. *Neth Heart J* **21**, 284-285, doi:10.1007/s12471-013-0413-z (2013).

Liu, G. S. *et al.* A novel human R25C-phospholamban mutation is associated with super-inhibition of calcium cycling and ventricular arrhythmia. *Cardiovasc Res* **107**, 164-174, doi:10.1093/cvr/cvv127 (2015).

Haghighi, K. *et al.* Human phospholamban null results in lethal dilated cardiomyopathy revealing a critical difference between mouse and human. *J Clin Invest* **111**, 869-876, doi:10.1172/JCI17892 (2003).

Karakikes, I. *et al.* Correction of human phospholamban R14del mutation associated with cardiomyopathy using targeted nucleases and combination therapy. *Nat Commun* **6**, 6955, doi:10.1038/ncomms7955 (2015).

Te Rijdt, W. P. *et al.* Phospholamban p.Arg14del cardiomyopathy is characterized by phospholamban aggregates, aggresomes, and autophagic degradation. *Histopathology* **69**, 542-550, doi:10.1111/his.12963 (2016).

Hoshijima, M. *et al.* Chronic suppression of heart-failure progression by a pseudophosphorylated mutant of phospholamban via in vivo cardiac rAAV gene delivery. *Nat Med* **8**, 864-871, doi:10.1038/nm739 (2002).

Ziolo, M. T., Martin, J. L., Bossuyt, J., Bers, D. M. & Pogwizd, S. M. Adenoviral gene transfer of mutant phospholamban rescues contractile dysfunction in failing rabbit myocytes with relatively preserved SERCA function. *Circ Res* **96**, 815-817, doi:10.1161/01.RES.0000163981.97262.3b (2005).