Two C-terminal sequence variations determine differential neurotoxicity between human and mouse α-synuclein

Natalie Landeck,[1,¶] Katherine E. Strathearn,[2, ¶, #a] Daniel Ysselstein,[2, ¶] Kerstin Buck,[1, #b], Sayan Dutta,[2,3] Siddhartha Banerjee,[4] Zhengjian Lv,[4, #c] John D. Hulleman,[2, #d] Jagadish Hindupur,[2, #e] Li-Kai Lin,[5] Sonal Padalkar,[5, #f] George P. McCabe,[6] Lia A. Stanciu,[5] Yuri L. Lyubchenko,[4] Deniz Kirik,[1,*] and Jean-Christophe Rochet[2,3,*]

[1]Brain Repair and Imaging in Neural Systems, Department of Experimental Medical Science, Lund University, Lund, Sweden

[2]Department of Medicinal Chemistry and Molecular Pharmacology, Purdue University, West Lafayette, IN, USA

[3]Purdue Institute for Integrative Neuroscience, Purdue University, West Lafayette, IN, USA

[4]Department of Pharmaceutical Sciences, University of Nebraska Medical Center, Omaha, NE, USA

[5]Department of Materials Engineering, Purdue University, West Lafayette, IN, USA

[6]Department of Statistics, Purdue University, West Lafayette, IN, USA

[¶]These authors contributed equally to this work

[#a]Current address: Oxford Genetics Ltd., USA
[#b]Current address: AbbVie Deutschland GmbH & Co KG, Ludwigshafen, Germany
[#c]Current address: Bruker Nanosurfaces Division, Goleta, Santa Barbara, CA, USA
[#d]Current address: Departments of Ophthalmology and Pharmacology, University of Texas Southwestern Medical Center, Dallas, TX, USA
[#e]Current address: Liveon Biolabs Pvt. Ltd., Tumakuru, Karnataka, India
[#f]Current address: Department of Mechanical Engineering, Iowa State University, Ames, IA, USA

*Correspondence: deniz.kirik@med.lu.se (D.K.), jrochet@purdue.edu (J.-C.R.)

Abstract

α-Synuclein (aSyn) aggregation is thought to play a central role in neurodegenerative disorders termed synucleinopathies, including Parkinson's disease (PD). Mouse aSyn contains a threonine residue at position 53 that mimics the human familial PD substitution A53T, yet in contrast to A53T patients, mice show no evidence of aSyn neuropathology even after aging. Here we studied the neurotoxicity of human A53T, mouse aSyn, and various human-mouse chimeras in cellular and *in vivo* models as well as their biochemical properties relevant to aSyn pathobiology. We report that mouse aSyn is less neurotoxic than the human A53T variant as a result of inhibitory effects of two C-terminal amino acid substitutions on membrane-induced aSyn aggregation and aSyn-mediated vesicle permeabilization. Our findings highlight the importance of membrane-induced self-assembly in aSyn neurotoxicity and suggest that inhibiting this process by targeting the C-terminal domain could slow neurodegeneration in PD and other synucleinopathy disorders.

Introduction

Parkinson's disease (PD) is a common progressive neurodegenerative disorder characterized

clinically by motor symptoms attributed to a loss of dopaminergic neurons in the *substantia nigra*

(SN). At post-mortem examination neurons in various brain regions of PD patients present with

cytosolic inclusions named Lewy bodies that contain amyloid-like fibrils of the presynaptic

protein α-synuclein (aSyn) (1). A number of patients with familial forms of PD have been found

to harbor mutations in the SNCA gene, including point mutations encoding the substitutions

A30P, E46K, H50Q, G51D, A53E, and A53T and gene multiplications (2). Genetic and

neuropathological findings in humans and data from animal model studies suggest that aSyn

self-assembly plays a central role in the pathogenesis of PD and other neurodegenerative

disorders involving an accumulation of aSyn aggregates in the brain, collectively referred to as

synucleinopathies (3). A detailed understanding of molecular mechanisms by which aSyn forms

neurotoxic aggregates is critical for developing therapies aimed at slowing neurodegeneration in

the brains of patients with PD and other synucleinopathy disorders.

aSyn has been reported to adopt a natively unfolded, monomeric structure in solution (4)

and to exist as a compact disordered monomer in mammalian cells (5), although additional

evidence suggests that it can also exist as an oligomer in the cytosol (6). The protein is typically

expressed as a 14.4 kDa polypeptide and consists of 3 domains. The N-terminal domain

spanning residues 1-67 contains five conserved, lysine-rich repeats. The central region

spanning residues 61-95 contains a sixth lysine-rich repeat and is highly hydrophobic. A key

feature of this region is the presence of a segment spanning residues 71-82 that is required for

aSyn aggregation (7). The C-terminal region spanning residues 96-140 is enriched with proline

and acidic residues and is thought to regulate aSyn aggregation through auto-inhibitory long-

range interactions (8, 9), with electrostatic interactions mediated by the acidic residues playing a

major role in increasing the fibrillization lag time (10). aSyn binds to anionic phospholipid

vesicles by forming an amphipathic α-helix with varying lengths, including a short N-terminal

helix spanning residues ~1-25 and a longer helix spanning residues ~1-97 encompassing both

the N-terminal and central hydrophobic domains (11, 12). Although membrane binding

apparently plays an important role in the normal function of aSyn related to regulation of

synaptic vesicle trafficking (12, 13), the protein has also been shown to undergo accelerated

aggregation when incubated in the presence of phospholipid vesicles at high protein:lipid ratios

(14-16). aSyn aggregation at the membrane surface is likely stimulated by the exposure of

hydrophobic residues as the membrane-bound protein shifts from the long-helix form to the

short-helix form (11, 16), and by the fact that molecular interactions needed for aSyn self-

assembly likely occur with a higher probability on the two dimensional surface of the lipid bilayer

than in solution (17, 18). Evidence from our laboratory suggests that membrane-induced aSyn

aggregation plays a key role in neurotoxicity (16), potentially via a mechanism involving

membrane permeabilization (19-22).

An interesting observation regarding aSyn biology is the fact that mouse aSyn (m-aSyn)

naturally contains a threonine residue at position 53 and is thus identical to the human familial

PD mutant A53T (h-aSyn A53T) at this site (23). This observation raises the question of why h-

aSyn A53T triggers accelerated neurodegeneration in the brains of familial PD patients,

whereas even aged mice show no evidence of neurodegeneration or the presence of Lewy

bodies (24). This paradox is particularly striking given that recombinant m-aSyn forms amyloid-

like fibrils more rapidly than h-aSyn WT or h-aSyn A53T (25, 26), apparently as a result of the

disruption of long-range interactions involving the C-terminal region and the N-terminal and

central hydrophobic domains (27). In fact, however, m-aSyn differs from h-aSyn-A53T by six

mismatches, one in the central hydrophobic domain (S87N) and five in the C-terminal region

(L100M, N103G, A107Y, D121G and D122S) (Figure 1A), and each of these could contribute to

the apparent difference in neurotoxicity between the two variants.

The goal of this study was to investigate the impact of the mismatches between m-aSyn

and h-aSyn A53T on aSyn neurotoxicity. Previously it was reported that rat aSyn (which differs

from the mouse protein by a single G-to-S mismatch at position 121) failed to trigger neuronal

cell death when expressed from a lentiviral vector infused in rat SN, in contrast to h-aSyn WT

and h-aSyn A53T (28). Similarly, we found that rat aSyn had a reduced ability to trigger nigral

dopaminergic cell death compared to h-aSyn A53T in the rAAV-mediated overexpression model

in the rat (29). Based on the close sequence similarity between mouse and rat aSyn, we

hypothesized that the mouse protein is less neurotoxic than h-aSyn A53T, potentially because

the toxic consequences of having a threonine residue at position 53 are alleviated by mismatch

residues elsewhere in the murine sequence. To address this hypothesis, we compared h-aSyn

A53T, m-aSyn, and various human-mouse chimeras in terms of their ability to elicit

neurodegeneration in cellular and animal models relevant to PD and other synucleinopathy

disorders. The human and mouse proteins and a subset of chimeras were further examined for

their ability to form amyloid-like fibrils in the absence of membranes, undergo membrane-

induced aSyn aggregation, and trigger membrane permeabilization.

Results

Neurotoxicity of aSyn variants in primary midbrain culture

In the first part of our study, we used a primary mesencephalic cell culture model to

examine the effects of amino acid variations between m-aSyn and h-aSyn A53T on aSyn

neurotoxicity. In addition to h-aSyn A53T and m-aSyn, two chimeric variants were analyzed: (i)

h-aSyn Chimera, consisting of h-aSyn with the human-to-mouse substitutions A53T, D121G,

and N122S; and (ii) m-aSyn Chimera, consisting of m-aSyn with the mouse-to-human

substitutions G121D and S122N (Figure 1B). We chose to examine the effects of substitutions

at positions 121 and 122 based on data showing that post-translational modifications near these

residues, including C-terminal truncation and Y125 phosphorylation, modulate the protein's

aggregation propensity and neurotoxicity (30-34).

Primary midbrain cultures were processed as untreated control or transduced with

adenovirus encoding each of the aSyn variants at multiplicities of infection (MOIs) that were

adjusted to ensure equal expression of the different variants. We have previously shown that

the adenoviral transduction efficiency is >90% for dopaminergic and non-dopaminergic neurons

(16). The cells were fixed and stained for microtubule associated protein 2 (MAP2), a general

neuronal marker, and tyrosine hydroxylase (TH), a marker of dopaminergic neurons. Relative

dopaminergic cell viability was evaluated by determining the percentage of MAP2$^+$ neurons that

were also TH$^+$. We found a significant reduction in the percentage of TH$^+$ neurons in cultures

transduced with h-aSyn A53T virus compared to untreated cultures or cultures expressing m-

aSyn (Figure 2A, B). Cultures expressing m-aSyn Chimera exhibited a decrease in relative TH$^+$

cell viability, similar to A53T-expressing cultures (Figure 2B). In contrast, h-aSyn Chimera had

no impact on dopaminergic cell survival, similar to m-aSyn (Figure 2A). In additional

experiments we found that cultures expressing A53T/D121G (but not A53T/N122S) exhibited a

significant decrease in TH$^+$ neuron loss compared to cultures expressing A53T (Figure 2C).

Together these results suggested that (i) h-aSyn A53T is more neurotoxic than m-aSyn, and (ii)

the human-to-mouse substitutions D121G and N122S (and in particular D121G) are at least

partially responsible for the decrease in neurotoxicity of m-aSyn compared to h-aSyn A53T.

In an independent set of experiments, we examined the impact of the mismatch between

m-aSyn and h-aSyn A53T at position 87 (Figure 1) on aSyn-mediated dopaminergic cell death.

Here our rationale was that residue 87 is located in the central hydrophobic domain, and

substitutions and deletions in this region have been shown to modulate aSyn aggregation and

neurotoxicity (7, 31). Cultures expressing h-aSyn A53T/S87N, h-aSyn Chimera S87N, or m-

aSyn Chimera N87S exhibited no difference in relative dopaminergic cell viability compared to

cultures expressing h-aSyn A53T (Figure 2D), h-aSyn Chimera (Figure 2E), or m-aSyn Chimera

(Figure 2F), respectively. From these results, we inferred that the mismatch between m-aSyn

and h-aSyn A53T at position 87 does not affect aSyn neurotoxicity.

Taken together, these findings suggest that residues D121 and N122 are important for

the ability of h-aSyn A53T to elicit dopaminergic cell death. Moreover, we infer that residues

G121 and S122 (but not N87) of m-aSyn interfere with the expected neurotoxic effect of the

threonine residue at position 53 of the mouse sequence.

Neurotoxicity of aSyn variants in rat nigrostriatal projection neurons

To further examine the effects of mismatches at positions 121 and 122 of m-aSyn and h-

aSyn A53T on aSyn neurotoxicity, we characterized h-aSyn A53T, m-aSyn, h-aSyn Chimera,

and m-aSyn Chimera in terms of their toxicity to nigral dopamine neurons *in vivo*. rAAV vectors

encoding these variants were injected unilaterally into the rat SN at two different titers,

corresponding to a low-toxicity titer at 1.3E13 gc/mL (n=4-6 per group) and a high-toxicity titer at

3.5E14 gc/mL (n=4-10 per group). Animals were killed 8 weeks after transduction, and serial

coronal midbrain sections were stained for histological analysis to assess transgene expression

and the extent of neurodegeneration in the nigral dopamine neurons. Representative specimens

of animals injected with a high-titer vector are shown in Figure 3A-Y. The aSyn antibody applied

here was used to evaluate transgene expression and is specific for aSyn but does not

distinguish between human and rodent variants as seen in Figure 3A, where endogenous aSyn

can be visualized especially in the axon terminal located in the SN pars reticulata contralateral

to the AAV injection. On the injected side, transduced cells over-expressing h-aSyn A53T

(Figure 3B), h-aSyn Chimera (Figure 3C), m-aSyn (Figure 3D), or m-aSyn Chimera (Figure 3E)

were visible in the SN pars compacta. The human aSyn-specific syn211 antibody resulted in

specific staining in groups expressing h-aSyn A53T or m-aSyn Chimera, whereas the

contralateral side of rat midbrain sections as well as in sections expressing m-aSyn or h-aSyn

Chimera were devoid of any immunoreactivity (Figure S1). Notably, staining with a pan aSyn

antibody, Syn-1, showed that the overexpression was present in all groups, suggesting that the

affinity of syn211 to the aSyn variants was dependent on the identity of the residues at positions

121 and 122 (Figure S1F-J).

Immunohistochemical analyses were carried out to evaluate the loss of dopaminergic

neurons in rat midbrain sections stained for TH (Figure 3F-J) and vesicular monoamine

transporter 2 (VMAT2) (Figure 3K-O). A clear loss of TH$^+$ as well as VMAT2$^+$ neurons was seen

in all groups when compared to the uninjected contralateral side (Figures 3F and 3K,

respectively). The reduction in cell numbers was not due to a loss of expression of

dopaminergic markers, as it was clearly visible also in specimens stained with antibodies

against the neuronal marker proteins Hu (Figure 3P-T) or with cresyl violet (CV) (Figure 3U-Y).

In order to quantify dopaminergic cell loss in the SN, we counted TH$^+$ cells in serial sections

from both the low-titer and high-titer vector groups. Results obtained for the low-titer group

showed no cell death for any of the aSyn variants and no significant differences among groups

(Figure 3Z). In contrast, in the 3.5E14 gc/mL titer groups, the expression of h-aSyn A53T

resulted in a significantly higher percentage of cell death when compared to the h-aSyn

Chimera group (75 ± 4 versus 50 ± 5% cell loss, respectively; Figure 3AA). Moreover, the m-

aSyn group displayed significantly reduced neurotoxicity in TH$^+$ neurons than the m-aSyn

Chimera group (56 ± 8 versus 17 ± 3% cell survival, respectively). These results suggested a

considerable effect of the human/mouse mismatches at positions 121 and 122 on aSyn-

mediated dopaminergic cell death in rat SN.

To determine whether the significant differences in dopaminergic cell death observed in

the SN were also evident in the nigrostriatal projections, we evaluated the TH$^+$ and VMAT2$^+$

fiber intensity in the striatum of the high titer groups. In serial sections throughout the

rostocaudal axis of the striatum, we found a reduction in staining intensity for both TH (Figure

4A-E, A'-E') and VMAT2 (Figure 4G-K, G'-K') for all groups when compared to contralateral

striatal areas. A significant reduction of innervation density was observed for the m-aSyn

Chimera group when compared to the m-aSyn group for both TH$^+$ (Figure 4F) and VMAT2$^+$

projections (Figure 4L). These observations are in accordance with dopaminergic cell loss seen

in the corresponding SN sections of the same animals. There was a trend for a difference in the

h-aSyn Chimera group when compared to the h-aSyn A53T group, although this did not reach

significance [TH (p=0.11); VMAT2 (p=0.27)].

Aggregate formation by aSyn variants in rat midbrain

To determine whether the aSyn variants differed in terms of aggregation propensity in rat

midbrain, we treated striatal sections with Proteinase K (PK) to digest non-aggregated species

and reveal digestion-resistant aSyn assemblies. As a reference for the PK incubation we chose

the endogenous aSyn staining in the CA3 region of the hippocampus (Figure S2A, F, K), which

would not be affected by nigral transduction. Whereas untreated striatal sections stained for

aSyn showed both punctate and fiber-like staining (Figure S2B-E), a clearance of digested and

therefore non-aggregated aSyn could already be appreciated after 5 min of PK incubation

(Figure S2F-J). At a later incubation time (45 min) when endogenous aSyn staining could no

longer be detected in the CA3 region (Figure S2K), and therefore all non-aggregated synuclein

would be digested, resistant aSyn aggregates were detected in striatal sections in all groups

(Figure S2L-O). These results suggested that the human/mouse mismatches at positions 121

and 122 do not have a major impact on the ability of aSyn to form PK-resistant aggregates in

the striatal fiber terminals of dopaminergic neurons.

Effects of human/mouse aSyn mismatches on aSyn fibrillization rates

The next set of experiments was aimed at determining the effects of human/mouse aSyn

mismatches on rates of aSyn fibrillization. Based on evidence that m-aSyn forms amyloid-like

fibrils more rapidly than h-aSyn WT or A53T (25, 26), we hypothesized that a more rapid

conversion of natively unfolded aSyn to mature, amyloid-like fibrils could lead to a depletion of

toxic oligomers (26, 35, 36). As a corollary, we reasoned that the human-to-mouse substitutions

could stimulate aSyn fibrillization, which in turn could account for the reduced neurotoxicity of m-

aSyn and h-aSyn Chimera compared to h-aSyn A53T. To address this hypothesis, we

monitored the fibrillization of human, mouse, and chimeric aSyn variants using a thioflavin T

fluorescence assay. Consistent with previous results, we found that m-aSyn and h-aSyn A53T

formed fibrils with a markedly reduced lag time compared to h-aSyn WT, although the decrease

in lag time was more pronounced for the mouse protein (Figure 5A). Analysis of the chimeric

aSyn variants revealed that h-aSyn Chimera formed fibrils with a longer lag time compared to h-

aSyn A53T, whereas the rate of fibrillization was only slightly decreased for m-aSyn Chimera

compared to m-aSyn (Figure 5B).

Secondly, we investigated the effects of the human/mouse mismatch at position 87 on

fibrillization rates, based on the premise that residue 87 is in the central hydrophobic region, and

substitutions or deletions in this region have been shown to modulate aSyn self-assembly (7,

31). In contrast to our observation that substitutions at position 87 had no impact on aSyn

neurotoxicity in primary midbrain cultures (Figure 2), A53T/S87N and h-aSyn Chimera/S87N

exhibited a reduced fibrillization lag time compared to h-aSyn A53T and h-aSyn Chimera,

respectively (Figure S3A, B). Moreover, m-aSyn N87S and m-aSyn Chimera/N87S formed

amyloid-like fibrils with a longer lag time compared to m-aSyn and m-aSyn Chimera,

respectively (Figure S3C, D).

Collectively, these results suggest that differences in neurotoxicity among the human,

mouse, and chimeric aSyn variants (Figures 2-4) correlate weakly with differences in

fibrillization rates.

Effects of human/mouse aSyn mismatches on fibril morphology and dimensions

Next, we examined the impact of human/mouse aSyn mismatches on fibril morphology

and dimensions based on evidence that different types of aSyn fibrils (e.g. cylindrical fibrils

versus flatter ribbons) can trigger different levels of neurotoxicity in cellular and animal models

(37). Examination of incubated samples of human, mouse, and chimeric aSyn by atomic force

microscopy (AFM) revealed that all of the variants formed straight, rigid, unbranched fibrils,

organized either as individual fibrils or in fibril bundles (Figure 6A-D). Two types of fibrils

('twisted' and 'non-twisted') were detected in samples of m-aSyn and h-aSyn-Chimera, whereas

only non-twisted fibrils were observed in samples of h-aSyn-A53T and m-aSyn-Chimera under

our experimental conditions. The heights of the non-twisted fibrils ranged from ~8 to ~11 nm,

whereas the heights of the 'valley' and 'peak' regions of the twisted fibrils ranged from ~7 to ~9

nm and ~11 to ~12 nm, respectively. The average pitch of twisted fibrils formed by m-aSyn and

h-aSyn Chimera was 123 ± 5 nm and 91 ± 8 nm, respectively (Figure 6E,F).

Further analysis of the fibrils by TEM also revealed the presence of straight, rigid,

unbranched fibrils for all of the variants (Figure 6G-L). The fibrils had similar properties as those

described for previously reported aSyn amyloid-like fibrils (e.g. diameter of ~10-15 nm; apparent

winding of two fibrils around each other via a helical twist) (26). Similar features were also

observed for fibrils formed by aSyn variants with the S87N or N87S substitution. There were no

striking differences in fibril morphology among the different aSyn variants, except that individual

fibrils formed by h-aSyn-Chimera had a twisted appearance similar to that observed for this

variant by AFM (in contrast, twisted fibrils of m-aSyn were not observed by TEM). It is unclear

why different results were obtained for m-aSyn and h-aSyn Chimera via AFM versus TEM,

although this discrepancy can potentially be explained by the fact that (i) the variants formed

two different types of twisted fibrils, as implied by their different pitch values (Figure 6E,F), and

(ii) different types of fibrils adsorb with different efficiencies to substrates with different

properties such as the hydrophilic mica and hydrophobic carbon-coated grids used here for

AFM and TEM, respectively (38).

Overall, these data suggest that differences in the morphologies of aSyn fibrils formed

under the self-assembly conditions used in this study could potentially contribute to differences

in neurotoxicity among the human, mouse, and chimeric aSyn variants (Figures 2-4).

Effects of human/mouse aSyn mismatches on membrane-induced aSyn aggregation.

Based on our earlier finding that lipid-induced self-assembly plays a key role in aSyn

neurotoxicity (16), we hypothesized that human-to-mouse substitutions should lead to a

decrease in aSyn aggregation propensity at membrane surfaces. Our rationale was that

although aSyn aggregation is primarily driven by the central hydrophobic region (7, 11, 16),

modifications to the C-terminal domain have been shown to modulate aSyn self-assembly (39)

and aSyn-membrane interactions (40) through effects on long-range interactions.

As a first step, we characterized the human, mouse, and chimeric variants in terms of

membrane affinities, based on the premise that differences in affinity could lead to differences in

the variants' propensity to undergo membrane-induced aggregation (16). Recombinant h-aSyn

WT, h-aSyn A53T, h-aSyn Chimera, m-aSyn, and m-aSyn Chimera in a monomeric state were

titrated with small unilamellar vesicles (SUVs) composed of egg phosphatidylglycerol (PG) and

egg phosphatidylcholine (PC) (1:1 mol/mol). Analysis of the samples via far-UV CD, a method

used to monitor the increase in aSyn α-helical structure that results from binding of the protein

to phospholipids, yielded similar lipid titration curves, K_d values, and maximum α-helical

contents for all of the aSyn variants (Figure 7A; Table S1). These results suggested that the C-

terminal human/mouse mismatches have little effect on the affinity of aSyn for phospholipid

membranes, or on the degree of α-helical structure adopted by the protein in the presence of

saturating lipid.

Next, we compared the human, mouse, and chimeric aSyn variants in terms of their

ability to undergo membrane-induced aggregation using a lipid-flotation assay. The proteins

were incubated with PG:PC SUVs under conditions that promote aSyn self-assembly, and the

membrane fraction was isolated by gradient centrifugation and analyzed via Western blotting.

The lane loaded with h-aSyn WT contained immunoreactive bands at ~45 and 60 kDa, whereas

the lane containing h-aSyn A53T or m-aSyn Chimera displayed more extensive laddering,

including bands at 75 and 100 kDa, and an intense smear of aSyn immunoreactivity in the

region of the gel above 245 kDa (Figure 7B). In contrast, high molecular-weight species were

markedly less abundant in the lanes loaded with h-aSyn Chimera or m-aSyn. Densitometry

analysis revealed a ~2-fold greater ratio of total oligomer band intensity to monomer band

intensity in the h-aSyn A53T and m-aSyn Chimera samples compared to the h-aSyn Chimera

and m-aSyn samples, respectively (Figure 7B). Although the results revealed a trend towards

an increase in the oligomer/monomer ratios for h-aSyn A53T and m-aSyn Chimera compared to

the human WT protein, these effects did not reach statistical significance. Collectively, these

data suggested that the human-to-mouse substitutions at positions 121 and 122 interfere with

membrane-induced aSyn aggregation.

Effects of human/mouse aSyn mismatches on a-Syn-mediated membrane disruption.

271   A number of groups have reported that aSyn aggregation at the membrane can affect

bilayer integrity via a mechanism involving phospholipid extraction coupled with aggregate

growth (19, 21). Recently, we found that two familial aSyn mutants with a high propensity to

form oligomers at membrane surfaces (A30P and G51D) (16) efficiently triggered vesicle

disruption under incubation conditions suitable for membrane-induced aggregation, whereas a

variant with a low propensity to undergo self-assembly at the membrane (A29E) exhibited no

vesicle permeabilization activity (22). Thus, we hypothesized that h-aSyn A53T and m-aSyn

Chimera should elicit greater membrane disruption than h-aSyn Chimera and m-aSyn, based on

the relative propensities of these variants to undergo membrane-induced aggregation (Figure

7B). To examine this hypothesis, we compared the human, mouse, and chimeric aSyn variants

in terms of their ability to disrupt vesicles using a dye release assay. PG-PC SUVs pre-loaded

with the fluorescent dye calcein at a concentration leading to self-quenching were incubated

with the aSyn variants in monomer form at 37 °C. Membrane permeabilization was assessed by

monitoring the increase in fluorescence intensity resulting from dequenching of the dye after its

release from the vesicles and dilution in the surrounding buffer. Upon extended incubation, all of

the aSyn-SUV mixtures displayed a time-dependent increase in fluorescence, in contrast to the

control sample consisting of vesicles without protein (Figure 7C). After 120 h, the fluorescence

emission observed for h-aSyn A53T and m-aSyn Chimera (corresponding to a yield of ~55% of

the maximal signal obtained by lysing the vesicles with detergent) was 2.5- and 1.5-fold greater

than that observed for h-aSyn Chimera and m-aSyn, respectively (Figure 7C). H-aSyn WT

produced a similar increase in fluorescence compared to h-aSyn A53T and m-aSyn Chimera.

Together, these data suggested that the human-to-mouse substitutions at positions 121 and

122 interfere with aSyn-mediated vesicle disruption under conditions compatible with aSyn

aggregation at the membrane surface.

Discussion

Understanding how human and mouse aSyn differ in terms of pathobiology is a high priority in

the synucleinopathy field not only because of the A/T mismatch at position 53 relevant to familial

PD, but also because of the widespread use of genetically engineered mice to investigate

molecular underpinnings of aSyn neuropathology *in vivo*. Previously, it was reported that

transgenic mice expressing m-aSyn downstream of the Thy-1 promoter show evidence at post-

mortem examination of extranigral aSyn accumulations similar to those observed in mice

expressing WT or mutant forms of h-aSyn (41). Moreover, Lashuel and colleagues (42) showed

that endogenous m-aSyn interferes with the formation of Lewy-like inclusions by h-aSyn WT in

cultured neurons and in mouse brain. Here we report that m-aSyn has a reduced propensity to

elicit dopaminergic cell death compared to h-aSyn A53T in primary midbrain cultures and

exhibits a similar effect when expressed from an AAV vector in rat SN. Furthermore, we show

for the first time that this differential toxicity is associated with differences in aSyn-mediated

vesicle disruption and aggregation at the membrane surface, rather than differences in aSyn

fibrillization previously suggested to be important for modulating its toxicity (43). Most

importantly, we provide evidence that these characteristics of the overtly toxic h-aSyn A53T

variant can be blunted, down to the same level of m-aSyn, by modification of only 2 amino acid

residues in positions 121 and 122 of the protein.

Our findings are consistent with the idea that m-aSyn has a reduced propensity to elicit

neurotoxicity compared to h-aSyn A53T as a result of attenuation of the toxic effects of

threonine 53 in the mouse protein by variations in other sites of the protein. Moreover, these

results support the hypothesis that mice show no signs of neurodegeneration despite

expressing an aSyn variant with a threonine residue at position 53 at least in part because m-

aSyn is intrinsically less toxic than h-aSyn A53T. The lack of nigral degeneration in mouse

models with altered expression of proteins that interact functionally with aSyn, including

transgenic mice over-expressing LRRK2 (44) and DJ-1 knockout mice (45), could also

potentially be explained by the reduced neurotoxicity of endogenous m-aSyn. Nevertheless,

despite its lower intrinsic toxicity, m-aSyn can also contribute to dopaminergic cell death in other

models relevant to nigral degeneration, including mice treated with MPTP (46). In addition to the

intrinsic biochemical differences between h-aSyn A53T and m-aSyn reported here, other factors

including differences in neurochemistry between the nigrostriatal systems of mice and humans

(47, 48) could also account at least in part for the apparent absence of early-onset aSyn

neuropathology in mice in contrast to human A53T patients.

Our data are in line with previous results showing that rat aSyn, identical to m-aSyn

except for a single G-to-S mismatch at position 121, triggers less extensive nigral degeneration

compared to h-aSyn WT, h-aSyn A30P, or h-aSyn A53T when expressed from a lentiviral vector

in rat midbrain (28). Moreover, we observed that rat aSyn has a similar blunted response of

nigral dopaminergic cell death compared to mouse aSyn in the rAAV-mediated overexpression

model in the rat (29). Another group reported evidence of TH[+] neuron loss and an accumulation

of Lewy-like inclusions in the SN of rats injected intrastriatally with preformed fibrils (PFFs)

prepared from recombinant m-aSyn (49). A similar but less pronounced neuropathological

phenotype was observed upon injecting a sample of non-fibrillized m-aSyn in rat midbrain,

although the interpretation of the results was complicated by uncertainty about the degree of

aSyn oligomerization in the injected, non-fibrillar protein sample.

A key finding of this study was the observation in both primary mesencephalic cultures

and rat midbrain that (i) h-aSyn Chimera, a variant of h-aSyn A53T with m-aSyn residues

introduced at positions 121 and 122, was less neurotoxic than h-aSyn A53T; and (ii) m-aSyn

Chimera, a variant of m-aSyn with human aSyn residues introduced at positions 121 and 122,

was more neurotoxic than the mouse protein. From these results, we infer that the presence of

glycine and serine residues at positions 121 and 122 of m-aSyn (respectively) is largely

responsible for offsetting the expected neurotoxic effects of threonine 53 in the murine

sequence. In turn, these findings reveal for the first time that residues 121 and 122 play a key

role in modulating aSyn-mediated neurodegeneration. Notably, in contrast to the striking effects

of human/mouse substitutions at positions 121 and 122, the S87N or N87S substitution had no

impact on the ability of human, mouse, or chimeric aSyn variants to trigger dopaminergic cell

death in primary midbrain cultures, suggesting that the variations in the two C-terminal residues

are unique in their impact on the toxicity. In support of these findings, a number of groups have

shown that modifications in the C-terminal domain, including truncation (31, 34) and Y125

phosphorylation (32), affect aSyn neurotoxicity in cellular and animal models.

Previously, m-aSyn was shown to form amyloid-like fibrils with a reduced lag time and a

greater elongation rate compared to h-aSyn WT or h-aSyn A53T (25, 26). Here we confirmed

these earlier findings and found that two chimeric variants, h-aSyn Chimera and m-aSyn

Chimera, undergo less rapid fibrillization than the mouse protein. These results indicate that the

human-to-mouse substitutions at positions 121 and 122 are necessary but not sufficient for the

enhanced fibrillization rate of m-aSyn relative to h-aSyn A53T. Because the human-to-mouse

and mouse-to-human substitutions at positions 121 and 122 do not have opposite effects on the

rate of aSyn fibrillization, we infer that the impact of each of these substitutions on aSyn self-

assembly is dependent on the sequence context, potentially because residues 121 and 122 are

involved in long-range intramolecular interactions that influence the rate of aSyn self-assembly

(8, 9). Additional data revealed that variants of h-aSyn A53T and h-aSyn Chimera with the S87N

substitution form fibrils with a decreased lag time, consistent with previously reported effects of

this mutation (25), whereas variants of m-aSyn and m-aSyn Chimera with the N87S substitution

form fibrils with an increased lag time.

Importantly, however, differences in fibrillization rates among the human, mouse, and

chimeric aSyn variants correlate only weakly with differences in neurotoxicity. For example, h-

aSyn A53T, h-aSyn Chimera, m-aSyn, and m-aSyn Chimera exhibit the following order of

fibrillization rates:

m-aSyn ≥ m-aSyn Chimera > h-aSyn A53T > h-aSyn Chimera

In contrast, the same variants exhibit the following order of neurotoxicity:

h-aSyn A53T ≈ m-aSyn Chimera > m-aSyn ≈ h-aSyn Chimera

Moreover, despite causing changes in aSyn fibrillization lag times, the S87N and N87S

substitutions have no impact on aSyn neurotoxicity in primary midbrain cultures, further

highlighting the weak correlation between fibrillization propensity and aSyn-mediated

neurodegeneration revealed by our study. The lack of a tight correlation may reflect the fact that

in mammalian cells non-membrane-bound aSyn adopts a compact disordered conformation that

is apparently incompatible with efficient fibrillization (5). Thus, a perturbation that favors or

hinders fibril formation by recombinant, natively unfolded aSyn in solution may have a very

different effect on the protein's conversion to neurotoxic species in its native intracellular

environment.

AFM and TEM analyses revealed that human, mouse, and chimeric aSyn formed

amyloid-like fibrils with different morphologies under the experimental conditions used here.

Namely, m-aSyn and h-aSyn Chimera, the two variants with reduced neurotoxicity, formed both

twisted and non-twisted fibrils, whereas only non-twisted fibrils were formed by h-aSyn A53T

and m-aSyn Chimera. These results are consistent with solid-state NMR data showing that

fibrils formed by h-aSyn WT and m-aSyn differ in terms of protofilament substructure (50, 51),

although previous EM analyses revealed no major morphological differences between fibrils

prepared from human A53T and mouse aSyn (25, 52). Multiple groups have shown that

injecting PFFs into the brains of WT or transgenic rodents initiates the aggregation of

endogenous aSyn via a seeding mechanism, leading to the formation of Lewy-like inclusions

(49, 53, 54). Moreover, evidence suggests that different types of fibrils (e.g. cylindrical fibrils

versus ribbons distinguishable by EM) can act as prion-like 'strains' with different abilities to

undergo cell-to-cell transmission throughout the brain (37). Our AFM and TEM data imply that

m-aSyn and h-aSyn Chimera could potentially form different fibrillar strains (characterized by

increased amounts of twisted fibrils) compared to the other aSyn variants, and, therefore, the

differences in *in vivo* neurotoxicity reported here could be due to differences in the variants'

abilities to seed aSyn neuropathology in rat brain. However, this interpretation is challenged by

evidence that (i) h-aSyn A53T can form twisted fibrils under certain incubation conditions (55);

and (ii) PFFs formed from recombinant m-aSyn induce inclusion formation and dopaminergic

cell death with greater or similar efficiency (respectively) compared to h-aSyn A53T PFFs

following their injection in mouse SN (56), in contrast to the reduced neurotoxicity of m-aSyn

compared to h-aSyn A53T reported here. Additional studies are needed to assess whether

human, mouse, or chimeric aSyn variants produce different fibrillar strains with different levels of

neurotoxicity *in vivo*.

Previously we reported that two familial substitutions, A30P and G51D, increase aSyn

neurotoxicity by enhancing the protein's ability to form aggregates at membrane surfaces (16).

Here we show that two aSyn variants with a higher propensity to elicit dopaminergic cell death

in primary culture and *in vivo*, h-aSyn A53T and m-aSyn Chimera, undergo membrane-induced

self-assembly with greater efficiency compared to two variants with lower neurotoxicity, h-aSyn

Chimera and m-aSyn. These results suggest that mouse-to-human substitutions at positions

121 and 122 cause an increase in aSyn neurotoxicity by promoting Syn aggregation at

membrane surfaces, whereas human-to-mouse substitutions have the opposite effect. Because

all of the aSyn variants examined in this study exhibited similar affinities for PG:PC SUVs, we

infer that human-to-mouse substitutions interfere with lipid-induced aSyn aggregation not by

favoring binding of the central hydrophobic domain to the membrane, but rather by inducing a

conformational change in the C-terminal region that disrupts aSyn self-assembly, potentially via

perturbation of long-range interactions (8, 9, 40). Consistent with this scenario, data from

several studies suggest that modifications in the C-terminal region (namely, Y125

phosphorylation or nitration, or the nitration of all three C-terminal tyrosine residues) can

modulate the conformation of the N-terminal and central hydrophobic domains of the

membrane-bound form of the protein (40, 57, 58). These findings support the idea that long-

range interactions can occur in the membrane-associated form of aSyn, in turn suggesting that

the human-to-mouse substitutions at positions 121 and 122 could potentially influence

membrane-induced aSyn aggregation by altering these intramolecular contacts. Moreover,

Burré et al. (59) have proposed a model wherein oligomers formed by aSyn at the membrane

surface consist of neighboring aSyn molecules with their C-terminal alpha-helices arranged in

an anti-parallel configuration. In this arrangement, the aspartate residue at position 121 of

human aSyn could potentially participate in electrostatic interactions with positively charged

residues on a neighboring subunit, and in turn these interactions could favor the formation of

pathological aSyn aggregates at the membrane surface. This idea is also supported by

computational data from our group suggesting that intermolecular interactions between the C-

terminal domain and central hydrophobic region play a key role in aSyn dimerization at the

surface of anionic phospholipid membranes (60).

In the current study and in our earlier work (16, 22), SDS-resistant aSyn oligomers were

detected in the lipid fraction of protein/SUV mixtures isolated by lipid flotation. These data

indicate that membrane-bound aSyn oligomers were generated upon incubating the protein with

SUVs. The formation of these membrane-bound oligomers could occur as a result of (i) aSyn

aggregation at the membrane surface, or (ii) aSyn aggregation in solution, followed by the

binding of the aggregates to the membrane. Three observations argue against the second

scenario. First, we find that aSyn does not form aggregates when incubated in the absence of

SUVs under the same conditions as those used in the membrane-induced aggregation assay.

Second, the relative rates at which aSyn variants form oligomers in the presence of SUVs

correlate poorly with the relative rates at which the variants undergo self-assembly in the

absence of phospholipids (16). Third, we recently showed that ENSA, a protein that interacts

selectively with membrane-bound aSyn (61), inhibits aSyn aggregation in the presence of

SUVs, whereas it does not interfere with aSyn fibrillization in the absence of lipids (22). The fact

that ENSA does not inhibit aSyn fibril formation in the absence of SUVs suggests that the

inhibitory effect of ENSA on aSyn self-assembly in the presence of lipids must arise from its

interaction with aSyn at the membrane surface. In turn, this evidence supports the idea that

aSyn forms oligomers on the membrane when incubated with SUVs under the conditions

described here.

Importantly, our data reveal a strong correlation between aggregation propensity at

membrane surfaces and neurotoxicity among the human, mouse, and chimeric aSyn variants,

and thus they further support the hypothesis that lipid-induced self-assembly plays a key role in

aSyn-mediated neurodegeneration (11, 16, 18, 22). The importance of membrane-induced aSyn

aggregation to the pathogenesis of synucleinopathy disorders is further underscored by

evidence that non-membrane-associated aSyn adopts a compact, aggregation-resistant

conformation, with little exposure of the central hydrophobic region, in mammalian cells (5).

Although it is uncertain why all of the aSyn variants examined here formed PK-resistant

aggregates in rat striatum, one explanation could be that the inclusions consist of fibrils derived

from both membrane-bound and non-membrane-bound forms of aSyn (20, 21). Accordingly, a

decrease in the levels of fibrils formed in the presence of lipids could be offset by an increase in

the levels of fibrils formed away from the membrane in the case of m-aSyn, whereas the

opposite scenario could be true in the case of h-aSyn A53T. Consistent with this idea, data

reported by multiple groups imply that aSyn can undergo various forms of self-assembly,

leading to the formation of different types of aggregated species with different degrees of

neurotoxicity, in cell culture models, aSyn transgenic mice, and the brains of synucleinopathy

patients (62-66). Although our results highlight the importance of membrane-induced aSyn

aggregation to aSyn-mediated dopaminergic cell death, we suggest that aSyn aggregates

formed both at membrane surfaces and in solution likely contribute to the overall

neuropathology of synucleinopathy disorders. Assessing whether differences in the rates or

extents of aSyn oligomerization in the absence of membranes also account at least in part for

differences in neurotoxicity among the aSyn variants is an important goal for future studies.

A final key outcome of this study was our finding that the relatively neurotoxic variants h-

aSyn A53T and m-aSyn Chimera elicit the disruption of synthetic vesicles under conditions

favoring membrane-induced aSyn aggregation with greater efficiency than the less toxic

variants h-aSyn Chimera and m-aSyn. These data suggest that mouse-to-human substitutions

at positions 121 and 122 enhance aSyn neurotoxicity by promoting aSyn-mediated membrane

permeabilization, whereas human-to-mouse substitutions have the opposite effect.

Unexpectedly, we found that h-aSyn WT triggered vesicle disruption to a similar extent as h-

aSyn A53T and m-aSyn Chimera. This observation was inconsistent with the fact that h-aSyn

WT exhibited a non-significant trend towards less extensive membrane-induced self-assembly,

or with evidence from our earlier studies that h-aSyn WT is considerably less toxic than h-aSyn

A53T in primary midbrain cultures (16) and causes neurodegeneration less rapidly than h-aSyn

A53T in rat midbrain (67). As one possibility, the current findings can be interpreted to mean

that h-aSyn WT engages in different molecular interactions during aggregation at the membrane

surface compared to h-aSyn A53T or m-aSyn Chimera, resulting in an increase in the extent of

membrane permeabilization relative to the total amount of aggregate formed. Consistent with

this idea, evidence from multiple studies suggests that h-aSyn WT and familial aSyn mutants

can form different types of amyloid-like fibrils (68-70). Overall, our data reveal a strong

correlation between vesicle disruption and neurotoxicity among the human, mouse, and

chimeric aSyn variants, and thus they support the notion that membrane permeabilization

associated with lipid-induced self-assembly plays a key role in aSyn-mediated

neurodegeneration. These findings are consistent with evidence that h-aSyn E57K, a variant

with a high propensity to form aggregates on a supported lipid bilayer and elicit membrane

damage, causes extensive dopaminergic cell death when expressed in rat SN (19, 35).

Previous findings reported by Volles et al. (71, 72) suggest that aSyn oligomers formed

in the absence of membranes fail to permeabilize low-diameter vesicles composed of a mixture

of anionic and zwitterionic lipids. Accordingly, we infer that the vesicle disruption observed in

aSyn-SUV mixtures in this study and in our earlier work (22) results from aSyn self-assembly at

the membrane surface. These observations further corroborate the idea that aSyn forms

oligomers on the membrane when incubated with SUVs under the conditions described in this

study. Moreover, oligomeric m-aSyn prepared in the absence of membranes was reported by

Volles et al. (71) to permeabilize vesicles composed of anionic lipids with a similar efficiency

compared to oligomeric h-aSyn A53T, contrary to the difference in the variants' neurotoxicity

reported here. This observation suggests that vesicle disruption associated with lipid-induced

aSyn aggregation plays a more important role than membrane permeabilization by preformed

oligomers in aSyn-mediated neurodegeneration *in vivo*.

Conclusions We report for the first time that m-aSyn has a reduced ability to elicit

dopaminergic cell death compared to h-aSyn A53T in primary mesencephalic cultures and in rat

midbrain. Our results highlight the central importance of membrane-induced aSyn aggregation

and vesicle disruption in aSyn neurotoxicity, and they reveal a C-terminal sequence motif

encompassing residues 121 and 122 with an important role in modulating these activities.

Collectively, these findings provide a strong rationale for developing therapies aimed at

inhibiting aSyn aggregation at membrane surfaces, including interventions that target the C-

terminal domain, as a strategy to slow neurodegeneration in the brains of synucleinopathy

patients (16, 18, 22).

Experimental Procedures

Materials. Unless otherwise stated all chemicals were purchased from Sigma-Aldrich (St. Louis,

MO or Stockholm, Sweden). DMEM, FBS, penicillin-streptomycin, Lipofectamine 2000,

Optimem, Trypsin-EDTA, ProLong Gold Antifade Reagent with DAPI, Proteinase K (25530-

049), and the Virapower Adenoviral Expression System were obtained from Invitrogen

(Carlsbad, CA). RPMI medium, phenol red-free RPMI medium, Nuserum, Plus slides

(J3800AMNZ), Phosphatase Inhibitor Mini Tablets, bicinchoninic acid (BCA) assay reagents,

Coomassie Brilliant Blue, and the Immobilon-FL PVDF membrane was purchased from Thermo

Fisher Scientific (Rockford, IL). The QuickTiter Adenovirus Titer Elisa kit was purchased from

Cell Biolabs (San Diego, CA). Antisedan, Temgesic, and the Fentanyl/Dormitor mixture were

obtained from Apoteksbolaget, (Stockholm, Sweden). The Vectastain ABC kit was purchased

from Vector Laboratories, Inc. (Burlingame, CA), and 3,3'-diaminobenzidine (DAB Safe) was

obtained from Saveen Werner (Limhamnsvägen, Sweden). Isopropyl β-D-1-

thiogalactopyranoside, ampicillin, and PMSF were purchased from Gold Biotechnology (St.

Louis, MO). Recombinant Syk, Src, and Fyn kinases were obtained from SignalChem

(Richmond, BC, Canada). L-α-phosphatidylcholine (egg PC), L-α-phosphatidylglycerol (egg

PG), and phospholipid extrusion membranes were purchased from Avanti Polar Lipids

(Alabaster, AL). Iodixanol and the enhanced chemifluorescence (ECF) substrate were obtained

from GE Life Sciences (Pittsburgh, PA). The nitrocellulose membrane was purchased from Bio-

Rad Laboratories (Hercules, CA). 100 kDa spin filters and 10 kDa concentration filters were

obtained from Millipore (Billerica, MA or Solna, Sweden). SH-SY5Y cells were purchased from

ATCC (Manassas, VA) (catalog # CRL-2266, RRID:CVCL_0019), and the PC12 Tet-Off cell line

(RRID:CVCL_V361) was obtained from Clontech Laboratories (Mountain View, CA). Cell line

authentication and mycoplasma testing were carried out by IDEXX BioResearch (Columbia,

MO) (cell line identities were confirmed via species-specific PCR evaluation and STR profiling,

and both cell lines were shown to be mycoplasma negative). A plasmid encoding murine Syk

fused at its C-terminus with EGFP (Syk-GFP) (73) was provided by Dr. Robert Geahlen (Purdue

University), and vectors encoding human GST-Src and GST-Fyn (74) were provided by Dr.

Laurie Parker (University of Minnesota), with permission from Dr. Benjamin Turk (Yale

University).

Antibodies. The following antibodies were used in these studies:

Primary cell culture – chicken anti-MAP2 (CPCA-MAP2, RRID:AB_2138173, EnCor

Biotechnology, Gainesville, FL); rabbit anti-tyrosine hydroxylase (TH) (AB152,

RRID:AB_390204, Millipore, Bellerica, MA); anti-rabbit IgG-Alexa Fluor 488 (R37116,

RRID:AB_2556544) and anti-chicken IgG-Alexa Fluor 594 (A11042, RRID:AB_142803)

(Thermo Fisher Scientific, Rockford, IL).

In vivo experiments – human-specific syn211 (36-008, RRID:AB_310817, Millipore,

Germany); mouse anti-aSyn, clone 42 (Syn-1) (610787, RRID:AB_398108, BD Biosciences,

UK); rabbit anti-TH (P40101-0, RRID:AB_461064**,** Pel-freeze, USA); rabbit anti-VMAT2

(ab81855, RRID:AB_2188123, Abcam, UK or 20042, RRID:AB_10013884, Immunostar, USA);

mouse anti-HuC/HuD, clone 16A11 (A21271, RRID:AB_221448, Thermo Scientific, USA);

biotinylated secondary antibodies (Vector Laboratories Inc., USA).

Western blotting – mouse anti-aSyn, clone 42 (Syn-1) (610787, RRID:AB_398108, BD

Biosciences, San Jose, CA); mouse anti-phosphotyrosine, clone 27B10 (APY03-HRP,

Cytoskeleton, Denver, CO); rabbit anti-pY125-aSyn (ab131466, RRID:AB_11158765, Abcam,

Cambridge, MA, or A1219, BioVision, Milpitas, CA); mouse anti-β-actin, clone AC-74 (A5316,

RRID:AB_476743, Sigma-Aldrich); mouse anti-GFP, clone B-2 (sc-9996, RRID:AB_627695,

Santa Cruz Biotechnology, Dallas, TX); rabbit anti-Src (2108, RRID:AB_331137), rabbit anti-Fyn

(4023, RRID:AB_2108441), AP-linked anti-mouse IgG (7056, RRID:AB_330921), and AP-linked

anti-rabbit IgG (7054, RRID:AB_2099235) (Cell Signaling Technology, Danvers, MA); goat anti-

mouse IR-Dye 680RD (92568070, RRID:AB_2651128) and goat anti-rabbit IR-Dye 800CW

(92532211, RRID:AB_2651127) (Li-COR, Lincoln, NE).

Preparation of bacterial expression constructs. Bacterial expression constructs encoding h-

aSyn WT (pT7-7-h-aSyn), h-aSyn A53T (pT7-7-h-aSyn-A53T), and m-aSyn (pT7-7-m-aSyn)

were described previously (26). A construct encoding h-aSyn Chimera was generated by

replacing the BamHI-HindIII fragment of pT7-7-h-aSyn-A53T (encompassing base pairs 354-

423 of the h-aSyn A53T cDNA) with the equivalent fragment from pT7-7-m-aSyn. A construct

encoding m-aSyn Chimera was generated by replacing the BamHI-HindIII fragment of pT7-7-m-

aSyn (encompassing base pairs 354-423 of the m-aSyn cDNA) with the equivalent fragment

from pT7-7-h-aSyn-A53T. pT7-7 constructs encoding A53T/D121G, A53T/N122S, A53T/S87N,

h-aSyn Chimera S87N, m-aSyn N87S, and m-aSyn Chimera N87S were generated via site-

directed mutagenesis using the QuikChange method (Stratagene). The sequence of the DNA

insert in each bacterial expression construct was verified using an Applied Biosystems

(ABI3700) DNA sequencer (Purdue Genomics Core Facility).

Adenoviral vector design and virus production. Adenoviral constructs encoding h-aSyn WT

and h-aSyn A53T were described previously (16, 75). cDNAs encoding h-aSyn Chimera,

A53T/D121G, A53T/N122S, A53T/S87N, and h-aSyn Chimera S87N (obtained using the pT7-7

constructs outlined above as PCR templates) were subcloned as KpnI-XhoI fragments into the

entry vector pENTR1A. cDNAs encoding m-aSyn, m-aSyn Chimera, and m-aSyn Chimera

N87S were subcloned as SalI-XhoI fragments into pENTR1A. Inserts from the pENTR1A

constructs were then transferred into the pAd/CMV/V5 adenoviral expression vector via

recombination using Gateway LR Clonase. The sequence of the DNA insert in each adenoviral

construct was verified using an Applied Biosystems (ABI3700) DNA sequencer (Purdue

Genomics Core Facility). Adenoviral constructs were packaged into virus via lipid-mediated

transient transfection of the HEK 293A packaging cell line. Purified adenoviral particles were

titered using the QuickTiter Adenovirus Titer Elisa kit.

AAV vector design and virus production. Adeno-associated viral vectors of serotype 5

(AAV5) were comprised of a genome cassette flanked by inverted terminal repeats (ITR2). The

cassette contained a chicken beta-actin promoter in combination with a CMV enhancer element

(CBA) leading to expression of one of the four different transgenes. Genes were PCR-amplified

with the plasmids pENTR1A h-aSyn A53T, pENTR1A h-aSyn Chimera, pENTR1A m-aSyn, and

pENTR1A m-aSyn Chimera as templates and subcloned as HindIII-SpeI fragments in the pTR-

UF20 plasmid. The validity of the constructs was confirmed by restriction analysis and Sanger

sequencing. Downstream of each cDNA insert was an h-SV40 polyA transcription termination

sequence. HEK293 cells were co-transfected at a confluency of 70-80% using the calcium-

phosphate precipitation method. Plasmids used here encode essential adenoviral packaging

and AAV5 capsid genes as previously described (76). Three days after transfection, cells were

harvested in PBS and lysed by performing three freeze-thaw cycles in a dry ice/ethanol bath.

The lysate was then treated with benzonase and purified using a discontinuous iodixanol

gradient followed by Sepharose Q column chromatography (77). Vectors were concentrated

using a 100 kDa molecular weight cut-off column, and titers of the stock solution were

determined by qPCR using primers and probes targeting the ITR sequence. Before being used

in an experiment, vectors were diluted in PBS, pH 7.4 and re-titered, yielding the values

reported in the Results and in Figure 3.

Preparation and treatment of primary mesencephalic cultures. Primary midbrain cultures

were prepared via dissection of day 17 embryos obtained from pregnant Sprague-Dawley rats

(Harlan, Indianapolis, IN, USA) as described (16, 75). All of the procedures involving animal

handling were approved by the Purdue Animal Care and Use Committee. The cells were plated

on poly-L-lysine-treated 48-well plates at a density of 163,500 cells per well. Five days after

plating, the cells were treated with cytosine arabinofuranoside (AraC) (20 µM, 48 h) to inhibit the

growth of glial cells. At this stage (i.e., 7 days in vitro (DIV)), the neurons appeared differentiated

with extended processes. The cultures were transduced with adenoviruses encoding the aSyn

variants for 72 h at a multiplicity of infection (MOI) ranging from 7 to 15 to ensure equal

expression levels as determined via Western blotting. After incubation for an additional 24 h in

fresh medium without virus, the cells were fixed in 4% (w/v) PFA in PBS (pH 7.4). The cells

were then washed in PBS and incubated with chicken anti-MAP2 (1:2000) and rabbit anti-

tyrosine hydroxylase (TH) 1:500) for 24 h. After an additional wash in PBS, the cultures were

incubated with AlexaFluor 594-conjugated goat anti-chicken and AlexaFluor 488-conjugated

goat anti-rabbit secondary antibodies (1:1000) for 1 h at 22 °C. ProLong Gold Antifade Reagent

with DAPI was applied to each well, and a coverslip was added.

Measurement of neuronal viability in midbrain cultures. Relative dopaminergic cell viability

was evaluated as described (16, 75). MAP2$^+$ and TH$^+$ cells were counted in a blinded manner in

at least 10 randomly chosen observational fields (approximately 500-1000 neurons total) for

each experimental condition using a Nikon TE2000-U inverted fluorescence microscope (Nikon

Instruments, Melville, NY) equipped with a 20x objective. The data are expressed as the

percentage of MAP2$^+$ neurons that are also TH$^+$ to correct for variation in cell density. Each

experiment was repeated a minimum of 3 times with cultures isolated from different pregnant

rats.

Animals for in vivo experiments. Young adult Sprague-Dawley rats received from Charles

River (Kisslegg, Germany) were housed on a 12 h light-dark cycle with *ad libitum* access to

water and food. All experimental procedures were approved by the Ethical Committee for Use of

Laboratory Animals in the Lund-Malmö region.

Stereotaxic surgery. Rats were anesthetized via i.p. injection of a 20:1 mixture of Fentanyl and

Dormitor (6 mL/kg). After placing each animal into a stereotaxic frame (Stoelting, Wood Dale,

USA), the SN was targeted unilaterally using the following coordinates: anteroposterior (AP), -

5.0 mm; mediolateral (ML), -2.0 mm from bregma; and dorsoventral (DV), -7.2 mm from the

dura. The tooth bar was adjusted to -2.3 mm. A pulled glass capillary (about 60-80 μm in

diameter) was attached to a 5 μL Hamilton syringe equipped with a 22s gauge needle and used

to deliver 2 μL of rAAV5 vector solution at a pulsed injection of 0.1 μL every 15 sec. After

delivery of the vector, the capillary was left in place for 5 min, retracted 0.1 mm, and after 1 min

it was slowly removed from the brain. After closing the wound with clips, Antisedan and

Temgesic were administered s.c. as an analgesic treatment and to reverse the anesthesia.

Histology. An overdose of sodium pentobarbital was used to kill rats 8 weeks after vector

delivery. Animals were perfused via the ascending aorta first with 50 mL of 0.9% (w/v) NaCl

followed by 250 mL of ice-cold 4% (w/v) PFA in 0.1 M phosphate buffer, pH 7.4, for 5 min.

Brains were taken out and post-fixed in 4% (w/v) PFA for 24 h and then incubated in 25% (w/v)

sucrose for cryoprotection. Brains were cut on a microtome (Microm HM450, Thermo Scientific,

USA) in 35 µm-thick coronal sections and placed into antifreeze solution (0.5 M phosphate

buffer, 30% (v/v) glycerol, 30% (v/v) ethylene glycol) at -20 °C for long term storage until further

processed.

Immunohistochemical staining was performed on free-floating brain sections. Specimens

were rinsed with Tris-buffered saline (TBS) buffer (5 mM Tris-HCl, 15 mM NaCl, pH 7.6). Only

TH staining required an antigen retrieval step carried out for 30 min at 80 °C using Tris/EDTA

buffer (10 mM Tris-HCl, 1 mM EDTA, pH 9.1). This step was terminated by repeated washes in

TBS at 22 °C. Endogenous peroxidase activity was quenched using a solution of 3% (v/v) H_2O_2

and 10% (v/v) methanol in TBS buffer for 30 min, and the reaction was stopped by washing

sections 3 times with TBS. To block unspecific binding sites, sections were incubated for 30 min

in 0.05% (v/v) Triton X-100 in TBS buffer (TBS-T) containing 5% (v/v) normal serum from the

same species as that in which the secondary antibody was raised. The primary antibodies were

then applied in 1% (w/v) BSA in TBS-T overnight at 22 °C: human-specific syn211 at 1:100,000;

Syn-1 at 1:10,000; TH at 1:5,000; VMAT2 ab81855 at 1:4,000, or 20042 at 1:10,000; HuC/HuD

1:200. The next day sections were washed in TBS-T, and the appropriate biotinylated

secondary antibody (1:200) was applied in 1% (w/v) BSA in TBS-T for 1 h. Specimens were

rinsed once more with TBS-T and further incubated in a solution containing avidin-biotin-

peroxidase complexes (Vectastain ABC kit, Vector Laboratories Inc., USA) for 1 h. Staining was

visualized using 3,3'-diaminobenzidine (DAB Safe) and 0.01% (v/v) H_2O_2. Sections were then

mounted on chromatin-gelatin coated glass slides, dehydrated in increasing alcohol solutions,

cleared with xylene, and applied to coverslips using DPX (06522, Sigma-Aldrich, Sweden).

For cresyl violet staining, sections were mounted on chromatin-gelatin coated glass

slides and dried overnight. Sections were then hydrated in decreasing alcohol solutions and

stained for 30 sec in 0.5% (w/v) cresyl violet + 0.1% (v/v) acetic acid. After rinsing specimens

with H_2O, sections were dehydrated in increasing alcohol solutions, cleared in xylene, and

coverslipped using DPX.

To visualize insoluble aggregates, sections were treated with Proteinase K. For this

purpose, specimens were first rinsed with potassium-containing phosphate-buffered saline

(KPBS) and then heat-treated at 80°C for 30 min in KPBS. Sections were then quenched using

3% (v/v) H_2O_2 and 10% (v/v) methanol in KPBS for 30 min. After mounting sections on plus

slides, the sections were incubated in a KPBS solution containing 10 µg/mL Proteinase K for

either 5 or 45 min. The same staining protocol was applied as described above using the aSyn-

specific primary antibody Syn-1 at 1:1,000.

Stereological analysis. In order to estimate surviving TH[+] nigral neurons, an unbiased

stereological quantification using the optical fractionator method (78, 79) was applied. The

implementation of a random starting position and systematic sampling was done using the

Visiopharm VIS software (version 4.5.5.433 Visiopharm A/S, Denmark) and a Nikon Eclipse 80i

light microscope fitted with an xy motorized stage, a motorized z-axis (Prior, UK), and a

microcator (Heidenheim, Germany). Through the rostro-caudal axis of the SN, every 6[th] section

was stained for TH. Regions of interest were outlined with a 4x objective, and counting was

executed using a 60x oil-immersion objective with a numerical aperture of 1.4. Counting

parameters were set to reach at least 100 cell counts per hemisphere. Estimation of the total

number of TH[+] cells was performed by a single blinded investigator. The coefficient of error due

to the sampling was calculated based on Gundersen and Jensen (80), and values lower than

0.1 were considered acceptable. The contralateral uninjected side of each animal served as an

internal control, and values were expressed as a percentage of cell counts on the injected side

relative to the intact side (mean ± SEM).

Densitometry. The optical density of TH^+ and $VMAT2^+$ fibers was measured on digital images

of coronal striatal sections using the Zeiss microscope (Axio Zoom.V16, Zeiss, Germany). The

striatum of every 24^{th} section in the rostro-caudal axis – in total 6 sections per brain – was

outlined using ImageJ, and optical density readings were corrected for non-specific background

using density measurements from the *corpus callosum* of each animal. Data consisting of

means ± SEM from all brains are expressed as a percentage of contralateral side values.

Protein purification. Recombinant aSyn variants were purified from BL21 (DE3) cells

transformed with pT7-7 constructs as described (16). The final purity of each protein was found

to be > 98% as determined by SDS-PAGE with Coomassie Blue staining.

aSyn fibrillization. Lyophilized aSyn was dissolved in PBS (pH 7.4) with 0.02% [v/v] NaN_3, and

the solution was filtered by successive centrifugation steps through a 0.22 µm spin filter and a

100 kDa centrifugal filter to isolate monomeric protein. The stock solution was dialyzed against

the same buffer (24 h at 4 °C) to remove excess salt. Fibrillization solutions of aSyn were

prepared by diluting the appropriate stock solutions of dialyzed protein with buffer to a final

concentration of 35 µM in the wells of a 96-well plate. To determine the extent of aSyn

fibrillization, thioflavin T (ThioT, final concentration, 20 µM) was added to the fibrillization

solutions, which were incubated at 37 °C with shaking at the 'normal' setting in a Spectra Fluor

Plus or Genios plate reader (Tecan, Upsala, Sweden). ThioT fluorescence was measured with

excitation at 440 nm and emission at 490 nm. Mean ThioT fluorescence data determined from 3

or 4 technical replicates (defined here as samples treated identically in a single experiment)

were normalized using the following equation:

$$normalized\ fluorescence = \frac{F_t - F_{min}}{F_{max} - F_{min}}$$ (Eq. 1)

where F_t is the ThioT fluorescence emission intensity at time t, and F_{min} and F_{max} are the

minimum and maximum fluorescence intensities during the incubation, respectively.

Lipid vesicle preparation. Small unilamellar vesicles (SUVs) (diameter = 50 nm) were

prepared as described (16). We opted to examine SUVs composed of egg PG and egg PC (1:1

mol/mol) because they contain anionic lipids necessary for aSyn membrane interactions (12)

and this lipid composition is compatible with producing stable 50 nm vesicles (16). Egg PG and

egg PC suspended in chloroform were mixed in a round bottom flask. The chloroform was

evaporated under a nitrogen stream and further dried under vacuum for 1 h. Dried lipids were

suspended in PBS, and 50 nm SUVs were prepared by extruding the suspension through a

Whatman membrane. The size of the vesicles was confirmed by dynamic light scattering. Lipid

vesicles were stored at 4 °C prior to use.

Circular dichroism. Far-UV CD measurements were performed as described (16) using a

Chirascan SC spectrometer (Applied Photophysics, Leatherhead, UK). Solutions of recombinant

aSyn (5 µM in 20 mM KPi, pH 7.4) in the absence or presence of SUVs (protein-to-lipid ratio,

1:50 to 1:1600, mol/mol) were analyzed in a 1 mm quartz cuvette at 25°C. The ellipticity at 222

nm was recorded and background corrected to eliminate scattering signals arising from the

SUVs. The mean residue molar ellipticity at 222 nm ($[\theta]_{MR,222}$) was calculated using the

following equation:

$$[\theta]_{MR,222} = \theta_{222}/10Cnl \qquad \text{(Eq. 2)}$$

772 where θ_{222} is the observed (background corrected) ellipticity at 222 nm in millidegrees, C is the

773 protein concentration in M, n is the number of amino acid residues in the protein, and l is the

774 path length of the cuvette in cm. Binding curves were generated by plotting $[\theta]_{MR,222}$ versus the

775 lipid concentration. Curves were analyzed by fitting to the following equation:

$$R = R_0 - (R_0 - R_f)\frac{K_d + C + L/N - \sqrt{(K_d + C + L/N)^2 - 4CL/N}}{2C} \qquad \text{(Eq. 3)}$$

779 where R is the measured $[\theta]_{MR,222}$ at a given lipid concentration, R_0 is $[\theta]_{MR,222}$ in the absence

780 of lipid, R_f is $[\theta]_{MR,222}$ in the presence of saturating lipid, L is the total lipid concentration, C is

781 the total protein concentration, Kd is the apparent macroscopic dissociation equilibrium

782 constant, and N is the binding stoichiometry (lipids/protein) (16, 81).

783   To determine the maximal helical content of each aSyn variant, we used the following

784 equations:

$$H = 100\%\frac{(\theta - \theta_{coil})}{(\theta_\alpha - \theta_{coil})} \qquad \text{(Eq. 4)}$$

$$\theta_\alpha = -40000\left(1 - \frac{2.5}{n}\right) + 100t \qquad \text{(Eq. 5)}$$

$$\theta_{coil} = 640 - 45t \qquad \text{(Eq. 6)}$$

where H is the maximal % helicity, θ is $[\theta]_{MR,222}$ in the presence of saturating lipid (R_f in Eq. 3),

θ_α is the $[\theta]_{MR,222}$ value of an idealized α-helical peptide, θ_{coil} is the $[\theta]_{MR,222}$ value of an

idealized random coil peptide, n is the number of amino acid residues in the protein (= 140), and

t is the temperature (25 °C) (16, 82, 83).

Lipid flotation assay. Lipid flotation analyses were carried out as described (16, 22).

Lyophilized aSyn was resuspended in PBS with 0.02% (w/v) NaN_3, and the solution was

dialyzed against the same buffer to remove salts and filtered by successive centrifugation steps

through a 0.22 μm spin filter and a 100 kDa centrifugal filter to isolate monomeric protein. aSyn

(40 μM) was incubated with PG:PC SUVs (protein-to-lipid ratio, 1:20 mol/mol) at 37°C for 72 h

in a total volume of 60 μL. After incubation, the sample was mixed with 4 mL of 30% (v/v)

iodixanol solution and overlaid with 7.0 mL of 25% (v/v) iodixanol and 350 μL of 5% (v/v)

iodixanol in a polyalomar tube (Beckman, Miami, FL). All of the above iodixanol solutions were

prepared in lipid flotation buffer (10 mM HEPES, pH 7.4, 150 mM NaCl). The samples were

spun at 200,000 x g in a Beckman SW 41 Ti rotor for 4 h. The membrane fraction was carefully

collected from the 5% iodixanol fraction at the top of the gradient, concentrated using a 10 kDa

spin filter, and analyzed via Western blotting using a primary antibody specific for aSyn (Syn-1)

(1:1,500) (16, 22). In general, we found that the rate of aSyn aggregation increased in the

presence of SUVs that had been stored for ~2-4 weeks at 4 °C (16).

Synthetic vesicle disruption assay. Calcein-loaded egg PG:PC SUVs were prepared as

described (84) with some modifications. A solution of calcein (170 mM) was prepared in H_2O

using NaOH to adjust the pH to 7.4. The final osmolality was 280 mOsm/kg. The calcein

solution was used to resuspend dried egg PG:PC lipids, and the suspension was extruded

through a Whatman membrane to generate 50 nm SUVs. Calcein-containing vesicles were

isolated from free calcein via gel filtration through a Sephadex G-50 column pre-equilibrated

with PBS, pH 7.4, 0.02% (w/v) NaN_3 (280 mOsm/kg). Fractions containing isolated vesicles

were pooled and stored at 4 °C until use. Calcein-loaded vesicles were found to be very stable,

with no spontaneous dye leakage observed over several weeks.

For membrane disruption experiments, monomeric aSyn variants were isolated as

described above (see 'Lipid Flotation Assay'). Each aSyn variant (40 µM) was incubated with

calcein-loaded PG:PC SUVs (protein-to-lipid ratio, 1:20 mol/mol) at 37°C in a total volume of 40

µL. At each time point, 3 µL of the reaction mixture was diluted into 180 µL of PBS, and the

diluted samples were analyzed with a Fluoromax-3 spectrofluorometer (Horiba Scientific,

Edison, New Jersey) (excitation wavelength, 485 nm; emission wavelengths, 505-530 nm; slit

width, 1 nm). To determine the maximum dye release, vesicles were lysed by adding 5 µL of 1%

(v/v) Triton X-100. The % leakage at time t was determined using the following equation:

$$\% \ leakage = \frac{I_t - I_0}{I_{max} - I_0} \times 100\% \hspace{3cm} \text{(Eq. 7)}$$

where I_t is the fluorescence emission intensity at time t, I_0 is the intensity at time 0, and I_{max} is

the maximal intensity determined after detergent lysis of the vesicles (intensity values were

determined at 515 nm).

AFM analysis of aSyn amyloid-like fibrils. A mica surface functionalized with 1-(3-

aminopropyl) silatrane (APS) was used to image aSyn amyloid-like fibrils by AFM (18, 85-87).

APS-mica was prepared by incubating freshly cleaved mica in a solution of APS (167 µM) for 30

min and then rinsed with deionized water and dried with an argon stream. A suspension of aSyn

fibrils (10 µL, prepared as described above under 'aSyn fibrillization' with an incubation time of

~100 h and diluted 1/20 in deionized water) was deposited onto the APS-mica surface, and the

sample was incubated for 2 min, rinsed with deionized water, and dried under an argon stream.

The sample was imaged with an AFM Nanoscope VIII system (Bruker, Santa Barbara, CA)

using MSNL probes (Cantilever F with spring constant 0.6 N/m), operating in air in peak force

mode. Images were acquired over a few randomly selected locations. Images were analyzed

using Gwyddion and FemtoScan online software (Advanced Technologies Center, Moscow,

Russia) (88, 89).

TEM analysis of aSyn amyloid-like fibrils. The morphology of aSyn amyloid-like fibrils

(prepared as described above under 'aSyn fibrillization' with an incubation time of ~100 h) was

analyzed by negative stain biological TEM (90). For biological sample preparation by negative

staining, which is a sample preparation technique that imparts the necessary contrast for

viewing during biological TEM imaging, 3 µL of aSyn sample solution (35 µM) was pipetted on a

discharged carbon-coated copper TEM grid substrate. Subsequently, the sample was washed

with deionized water carefully without letting it dry and stained with a 1% (w/v) phosphotungstic

acid solution (3 µL), which was left in contact with the protein on the grid for 1 min. The excess

solution was then removed by blotting with filter paper, and the sample was imaged using an

FEI Tecnai G2 20 Transmission Electron Microscope operating at 200 KV.

Sample size estimation. Samples sizes for studies of primary neuron viability, membrane-

induced aSyn aggregation, and aSyn-mediated membrane permeabilization were estimated via

power analysis based on our previous studies (16, 22). Power analyses were designed to

determine the number of biological replicates required to detect a ~20% or ~30% difference

from control, with α = 0.05 (type I error) and power = 0.80 (statistical power). Biological

replicates are defined here as samples treated identically in independent experiments carried

out on different days. For *in vivo* experiments, sample sizes were estimated based on previous

data collected by our group and others for the A53T variant.

Statistical analysis. Statistical analyses were carried out using GraphPad Prism 6.0 (La Jolla,

CA). Primary neuron viability data, in vivo stereology and densitometry data, densitometry data

from Western blots, and calcein dye release data were analyzed via ANOVA followed by

Tukey's multiple comparisons *post hoc* test for normally distributed measurements. In analyzing

percentage dye release data and percentage cell viability data by ANOVA, square root

transformations were carried out to conform to ANOVA assumptions. Normalized FFN-102

fluorescence data were subjected to a log transformation to account for skewness in the data.

The log-transformed data were analyzed using an approach that accounts for comparison

across experiments conducted on different days. Log-transformed fluorescence emission values

for multiple treatment groups were compared using a general linear model implemented in the

GLM procedure of SAS Version 9.3 followed by Tukey's multiple comparisons *post hoc* test

(Cary, NC). Kruskal-Wallis followed by a Dunn's multiple comparisons test or a two-tailed Mann-

Whitney test were used as non-parametric tests. Differences were considered significant for

$p < 0.05$. Unless otherwise stated, values of n indicated in the Figure Legends refer to the

number of independent biological replicates.

Acknowledgments

885 We thank Björn Anzelius, Anneli Josefsson, Ulla Samuelsson, and Ulrika Sparrhult-Björk (Lund

886 University) for excellent technical assistance and Aswathy Chandran (Purdue University) for

887 valuable discussions. We are also grateful to Dr. Christopher Gilpin (Purdue University) for

888 advice on TEM analyses.

References

1. Spillantini MG, Schmidt ML, Lee VM-Y, Trojanowski JQ, Jakes R, Goedert M. a-Synuclein
in Lewy bodies. Nature. 1997;388:839-40.

2.   Petrucci S, Ginevrino M, Valente EM. Phenotypic spectrum of alpha-synuclein mutations:
New insights from patients and cellular models. Parkinsonism & related disorders. 2016;22
Suppl 1:S16-20.
3.   Deleersnijder A, Gerard M, Debyser Z, Baekelandt V. The remarkable conformational
plasticity of alpha-synuclein: blessing or curse? Trends in molecular medicine.
2013;19(6):368-77.
4.   Weinreb PH, Zhen W, Poon AW, Conway KA, Lansbury PT, Jr. NACP, a protein implicated
in Alzheimer's disease and learning, is natively unfolded. Biochemistry. 1996;35(43):13709-
15.
5.   Theillet FX, Binolfi A, Bekei B, Martorana A, Rose HM, Stuiver M, et al. Structural disorder
of monomeric alpha-synuclein persists in mammalian cells. Nature. 2016;530(7588):45-50.
6.   Bartels T, Choi JG, Selkoe DJ. alpha-Synuclein occurs physiologically as a helically folded
tetramer that resists aggregation. Nature. 2011;477(7362):107-10.
7.   Giasson BI, Murray IV, Trojanowski JQ, Lee VM. A hydrophobic stretch of 12 amino acid
residues in the middle of a-synuclein is essential for filament assembly. J Biol Chem.
2001;276(4):2380-6.
8.   Dedmon MM, Lindorff-Larsen K, Christodoulou J, Vendruscolo M, Dobson CM. Mapping
long-range interactions in alpha-synuclein using spin-label NMR and ensemble molecular
dynamics simulations. Journal of the American Chemical Society. 2005;127(2):476-7.
9.   Bertoncini CW, Jung YS, Fernandez CO, Hoyer W, Griesinger C, Jovin TM, et al. Release
of long-range tertiary interactions potentiates aggregation of natively unstructured alpha-
synuclein. P Natl Acad Sci USA. 2005;102(5):1430-5.
10. Levitan K, Chereau D, Cohen SI, Knowles TP, Dobson CM, Fink AL, et al. Conserved C-
terminal charge exerts a profound influence on the aggregation rate of alpha-synuclein. J
Mol Biol. 2011;411(2):329-33.
11. Bodner CR, Dobson CM, Bax A. Multiple tight phospholipid-binding modes of alpha-
synuclein revealed by solution NMR spectroscopy. J Mol Biol. 2009;390(4):775-90.
12. Davidson WS, Jonas A, Clayton DF, George JM. Stabilization of alpha-synuclein secondary
structure upon binding to synthetic membranes. J Biol Chem. 1998;273(16):9443-9.
13. Venda LL, Cragg SJ, Buchman VL, Wade-Martins R. alpha-Synuclein and dopamine at the
crossroads of Parkinson's disease. Trends Neurosci. 2010;33(12):559-68.
14. Lee H-J, Choi C, Lee S-J. Membrane-bound a-synuclein has a high aggregation propensity
and the ability to seed the aggregation of the cytosolic form. J Biol Chem. 2002;277:671-8.
15. Galvagnion C, Buell AK, Meisl G, Michaels TC, Vendruscolo M, Knowles TP, et al. Lipid
vesicles trigger alpha-synuclein aggregation by stimulating primary nucleation. Nat Chem
Biol. 2015;11(3):229-34.
16. Ysselstein D, Joshi M, Mishra V, Griggs AM, Asiago JM, McCabe GP, et al. Effects of
impaired membrane interactions on alpha-synuclein aggregation and neurotoxicity.
Neurobiol Dis. 2015;79:150-63.
17. Abedini A, Raleigh DP. A role for helical intermediates in amyloid formation by natively
unfolded polypeptides? Phys Biol. 2009;6(1):15005.
18. Banerjee S, Hashemi M, Lv Z, Maity S, Rochet JC, Lyubchenko YL. A novel pathway for
amyloids self-assembly in aggregates at nanomolar concentration mediated by the
interaction with surfaces. Scientific reports. 2017;7:45592.
19. Reynolds NP, Soragni A, Rabe M, Verdes D, Liverani E, Handschin S, et al. Mechanism of
membrane interaction and disruption by alpha-synuclein. J Am Chem Soc.
2011;133(48):19366-75.
20. Comellas G, Lemkau LR, Zhou DH, George JM, Rienstra CM. Structural intermediates
during alpha-synuclein fibrillogenesis on phospholipid vesicles. J Am Chem Soc.
2012;134(11):5090-9.

21. Lee JH, Hong CS, Lee S, Yang JE, Park YI, Lee D, et al. Radiating amyloid fibril formation
on the surface of lipid membranes through unit-assembly of oligomeric species of alpha-
synuclein. PLoS ONE. 2012;7(10):e47580.
22. Ysselstein D, Dehay B, Costantino IM, McCabe GP, Frosch MP, George JM, et al.
Endosulfine-alpha inhibits membrane-induced alpha-synuclein aggregation and protects
against alpha-synuclein neurotoxicity. Acta neuropathologica communications. 2017;5(1):3.
23. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, et al. Mutation in the
alpha-synuclein gene identified in families with Parkinson's disease. Science.
1997;276(5321):2045-7.
24. Ahlemeyer B, Halupczok S, Rodenberg-Frank E, Valerius KP, Baumgart-Vogt E.
Endogenous Murine Amyloid-beta Peptide Assembles into Aggregates in the Aged
C57BL/6J Mouse Suggesting These Animals as a Model to Study Pathogenesis of
Amyloid-beta Plaque Formation. J Alzheimers Dis. 2018;61(4):1425-50.
25. Kang L, Wu KP, Vendruscolo M, Baum J. The A53T mutation is key in defining the
differences in the aggregation kinetics of human and mouse alpha-synuclein. J Am Chem
Soc. 2011;133(34):13465-70.
26. Rochet JC, Conway KA, Lansbury PT, Jr. Inhibition of fibrillization and accumulation of
prefibrillar oligomers in mixtures of human and mouse a-synuclein. Biochemistry.
2000;39(35):10619-26.
27. Wu KP, Kim S, Fela DA, Baum J. Characterization of conformational and dynamic
properties of natively unfolded human and mouse alpha-synuclein ensembles by NMR:
implication for aggregation. J Mol Biol. 2008;378(5):1104-15.
28. Lo Bianco C, Ridet JL, Schneider BL, Deglon N, Aebischer P. alpha -Synucleinopathy and
selective dopaminergic neuron loss in a rat lentiviral-based model of Parkinson's disease.
Proc Natl Acad Sci U S A. 2002;99(16):10813-8.
29. Landeck N, Buck K, Kirik D. Toxic effects of human and rodent variants of alpha-synuclein
in vivo. The European journal of neuroscience. 2017;45(4):536-47.
30. Murray IV, Giasson BI, Quinn SM, Koppaka V, Axelsen PH, Ischiropoulos H, et al. Role of
alpha-synuclein carboxy-terminus on fibril formation in vitro. Biochemistry.
2003;42(28):8530-40.
31. Periquet M, Fulga T, Myllykangas L, Schlossmacher MG, Feany MB. Aggregated alpha-
synuclein mediates dopaminergic neurotoxicity in vivo. J Neurosci. 2007;27(12):3338-46.
32. Chen L, Periquet M, Wang X, Negro A, McLean PJ, Hyman BT, et al. Tyrosine and serine
phosphorylation of alpha-synuclein have opposing effects on neurotoxicity and soluble
oligomer formation. J Clin Invest. 2009;119(11):3257-65.
33. Negro A, Brunati AM, Donella-Deana A, Massimino ML, Pinna LA. Multiple phosphorylation
of a-synuclein by protein tyrosine kinase Syk prevents eosin-induced aggregation. FASEB
J. 2002;16(2):210-2.
34. Bassil F, Fernagut PO, Bezard E, Pruvost A, Leste-Lasserre T, Hoang QQ, et al. Reducing
C-terminal truncation mitigates synucleinopathy and neurodegeneration in a transgenic
model of multiple system atrophy. Proc Natl Acad Sci U S A. 2016;113(34):9593-8.
35. Winner B, Jappelli R, Maji SK, Desplats PA, Boyer L, Aigner S, et al. In vivo demonstration
that alpha-synuclein oligomers are toxic. Proc Natl Acad Sci U S A. 2011;108(10):4194-9.
36. Karpinar DP, Balija MB, Kugler S, Opazo F, Rezaei-Ghaleh N, Wender N, et al. Pre-fibrillar
alpha-synuclein variants with impaired beta-structure increase neurotoxicity in Parkinson's
disease models. Embo J. 2009;28(20):3256-68.
37. Bousset L, Pieri L, Ruiz-Arlandis G, Gath J, Jensen PH, Habenstein B, et al. Structural and
functional characterization of two alpha-synuclein strains. Nature communications.
2013;4:2575.

38.   Gorman PM, Yip CM, Fraser PE, Chakrabartty A. Alternate aggregation pathways of the
Alzheimer beta-amyloid peptide: Abeta association kinetics at endosomal pH. J Mol Biol.
2003;325(4):743-57.
39.   Paleologou KE, Schmid AW, Rospigliosi CC, Kim HY, Lamberto GR, Fredenburg RA, et al.
Phosphorylation at Ser-129 but not the phosphomimics S129E/D inhibits the fibrillation of
alpha-synuclein. J Biol Chem. 2008;283(24):16895-905.
40.   Sevcsik E, Trexler AJ, Dunn JM, Rhoades E. Allostery in a disordered protein: oxidative
modifications to alpha-synuclein act distally to regulate membrane binding. J Am Chem
Soc. 2011;133(18):7152-8.
41.   Rieker C, Dev KK, Lehnhoff K, Barbieri S, Ksiazek I, Kauffmann S, et al. Neuropathology in
mice expressing mouse alpha-synuclein. PloS one. 2011;6(9):e24834.
42.   Fares MB, Maco B, Oueslati A, Rockenstein E, Ninkina N, Buchman VL, et al. Induction of
de novo alpha-synuclein fibrillization in a neuronal model for Parkinson's disease. Proc Natl
Acad Sci U S A. 2016;113(7):E912-21.
43.   Burre J, Sharma M, Sudhof TC. Definition of a Molecular Pathway Mediating alpha-
Synuclein Neurotoxicity. J Neurosci. 2015;35(13):5221-32.
44.   Li Y, Liu W, Oo TF, Wang L, Tang Y, Jackson-Lewis V, et al. Mutant LRRK2(R1441G) BAC
transgenic mice recapitulate cardinal features of Parkinson's disease. Nat Neurosci.
2009;12(7):826-8.
45.   Goldberg MS, Pisani A, Haburcak M, Vortherms TA, Kitada T, Costa C, et al. Nigrostriatal
dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-
linked gene DJ-1. Neuron. 2005;45(4):489-96.
46.   Fornai F, Schluter OM, Lenzi P, Gesi M, Ruffoli R, Ferrucci M, et al. Parkinson-like
syndrome induced by continuous MPTP infusion: Convergent roles of the ubiquitin-
proteasome system and {alpha}-synuclein. Proc Natl Acad Sci U S A. 2005;102(9):3413-8.
47.   Burbulla LF, Song P, Mazzulli JR, Zampese E, Wong YC, Jeon S, et al. Dopamine
oxidation mediates mitochondrial and lysosomal dysfunction in Parkinson's disease.
Science. 2017;357(6357):1255-61.
48.   Carballo-Carbajal I, Laguna A, Romero-Gimenez J, Cuadros T, Bove J, Martinez-Vicente
M, et al. Brain tyrosinase overexpression implicates age-dependent neuromelanin
production in Parkinson's disease pathogenesis. Nat Commun. 2019;10(1):973.
49.   Paumier KL, Luk KC, Manfredsson FP, Kanaan NM, Lipton JW, Collier TJ, et al.
Intrastriatal injection of pre-formed mouse alpha-synuclein fibrils into rats triggers alpha-
synuclein pathology and bilateral nigrostriatal degeneration. Neurobiol Dis. 2015;82:185-99.
50.   Lv G, Kumar A, Huang Y, Eliezer D. A Protofilament-Protofilament Interface in the Structure
of Mouse alpha-Synuclein Fibrils. Biophys J. 2018;114(12):2811-9.
51.   Hwang S, Fricke P, Zinke M, Giller K, Wall JS, Riedel D, et al. Comparison of the 3D
structures of mouse and human alpha-synuclein fibrils by solid-state NMR and STEM. J
Struct Biol. 2019;206(1):43-8.
52.   Tanaka G, Yamanaka T, Furukawa Y, Kajimura N, Mitsuoka K, Nukina N. Sequence- and
seed-structure-dependent polymorphic fibrils of alpha-synuclein. Biochimica et biophysica
acta Molecular basis of disease. 2019;1865(6):1410-20.
53.   Volpicelli-Daley LA, Luk KC, Patel TP, Tanik SA, Riddle DM, Stieber A, et al. Exogenous
alpha-synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and
neuron death. Neuron. 2011;72(1):57-71.
54.   Luk KC, Kehm V, Carroll J, Zhang B, O'Brien P, Trojanowski JQ, et al. Pathological alpha-
synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice.
Science. 2012;338(6109):949-53.
55.   Conway KA, Harper JD, Lansbury PT, Jr. Fibrils formed *in vitro* from a-synuclein and two
mutant forms linked to Parkinson's disease are typical amyloid. Biochemistry.
2000;39:2552-63.

56. Luk KC, Covell DJ, Kehm VM, Zhang B, Song IY, Byrne MD, et al. Molecular and Biological
Compatibility with Host Alpha-Synuclein Influences Fibril Pathogenicity. Cell reports.
2016;16(12):3373-87.
57. Hejjaoui M, Butterfield S, Fauvet B, Vercruysse F, Cui J, Dikiy I, et al. Elucidating the role of
C-terminal post-translational modifications using protein semisynthesis strategies: alpha-
synuclein phosphorylation at tyrosine 125. J Am Chem Soc. 2012;134(11):5196-210.
58. Burai R, Ait-Bouziad N, Chiki A, Lashuel HA. Elucidating the Role of Site-Specific Nitration
of alpha-Synuclein in the Pathogenesis of Parkinson's Disease via Protein Semisynthesis
and Mutagenesis. J Am Chem Soc. 2015;137(15):5041-52.
59. Burre J, Sharma M, Sudhof TC. alpha-Synuclein assembles into higher-order multimers
upon membrane binding to promote SNARE complex formation. Proc Natl Acad Sci U S A.
2014;111(40):E4274-83.
60. Lv Z, Hashemi M, Banerjee S, Zagorski K, Rochet JC, Lyubchenko YL. Assembly of alpha-
synuclein aggregates on phospholipid bilayers. Biochimica et biophysica acta Proteins and
proteomics. 2019;1867(9):802-12.
61. Woods WS, Boettcher JM, Zhou DH, Kloepper KD, Hartman KL, Ladror DT, et al.
Conformation-specific binding of alpha-synuclein to novel protein partners detected by
phage display and NMR spectroscopy. J Biol Chem. 2007;282(47):34555-67.
62. Parkkinen L, O'Sullivan SS, Collins C, Petrie A, Holton JL, Revesz T, et al. Disentangling
the relationship between lewy bodies and nigral neuronal loss in Parkinson's disease.
Journal of Parkinson's disease. 2011;1(3):277-86.
63. Raiss CC, Braun TS, Konings IB, Grabmayr H, Hassink GC, Sidhu A, et al. Functionally
different alpha-synuclein inclusions yield insight into Parkinson's disease pathology.
Scientific reports. 2016;6:23116.
64. Tompkins MM, Hill WD. Contribution of somal Lewy bodies to neuronal death. Brain
research. 1997;775(1-2):24-9.
65. Tsika E, Moysidou M, Guo J, Cushman M, Gannon P, Sandaltzopoulos R, et al. Distinct
region-specific alpha-synuclein oligomers in A53T transgenic mice: implications for
neurodegeneration. J Neurosci. 2010;30(9):3409-18.
66. Surmeier DJ, Obeso JA, Halliday GM. Selective neuronal vulnerability in Parkinson
disease. Nat Rev Neurosci. 2017;18(2):101-13.
67. Maingay M, Romero-Ramos M, Kirik D. Viral vector mediated overexpression of human
alpha-synuclein in the nigrostriatal dopaminergic neurons: a new model for Parkinson's
disease. CNS spectrums. 2005;10(3):235-44.
68. Heise H, Celej MS, Becker S, Riedel D, Pelah A, Kumar A, et al. Solid-state NMR reveals
structural differences between fibrils of wild-type and disease-related A53T mutant alpha-
synuclein. J Mol Biol. 2008;380(3):444-50.
69. Lemkau LR, Comellas G, Lee SW, Rikardsen LK, Woods WS, George JM, et al. Site-
specific perturbations of alpha-synuclein fibril structure by the Parkinson's disease
associated mutations A53T and E46K. PloS one. 2013;8(3):e49750.
70. Yonetani M, Nonaka T, Masuda M, Inukai Y, Oikawa T, Hisanaga S, et al. Conversion of
wild-type alpha-synuclein into mutant-type fibrils and its propagation in the presence of
A30P mutant. J Biol Chem. 2009;284(12):7940-50.
71. Volles MJ, Lansbury PT, Jr. Vesicle permeabilization by protofibrillar a-synuclein is
sensitive to Parkinson's disease-linked mutations and occurs by a pore-like mechanism.
Biochemistry. 2002;41:4595-602.
72. Volles MJ, Lee S-J, Rochet J-C, Shtilerman MD, Ding TT, Kessler JC, et al. Vesicle
permeabilization by protofibrillar a-synuclein: implications for the pathogenesis and
treatment of Parkinson's disease. Biochemistry. 2001;40(26):7812-9.
73. Ma H, Yankee TM, Hu J, Asai DJ, Harrison ML, Geahlen RL. Visualization of Syk-antigen
receptor interactions using green fluorescent protein: differential roles for Syk and Lyn in

the regulation of receptor capping and internalization. Journal of immunology (Baltimore,
Md : 1950). 2001;166(3):1507-16.
74. Deng Y, Alicea-Velazquez NL, Bannwarth L, Lehtonen SI, Boggon TJ, Cheng HC, et al.
Global analysis of human nonreceptor tyrosine kinase specificity using high-density peptide
microarrays. Journal of proteome research. 2014;13(10):4339-46.
75. Liu F, Hindupur J, Nguyen JL, Ruf KJ, Zhu J, Schieler JL, et al. Methionine sulfoxide
reductase A protects dopaminergic cells from Parkinson's disease-related insults. Free
Radic Biol Med. 2008;45(3):242-55.
76. Grimm D, Kern A, Rittner K, Kleinschmidt JA. Novel tools for production and purification of
recombinant adenoassociated virus vectors. Human gene therapy. 1998;9(18):2745-60.
77. Zolotukhin S, Byrne BJ, Mason E, Zolotukhin I, Potter M, Chesnut K, et al. Recombinant
adeno-associated virus purification using novel methods improves infectious titer and yield.
Gene therapy. 1999;6(6):973-85.
78. Schmitz C, Hof PR. Design-based stereology in neuroscience. Neuroscience.
2005;130(4):813-31.
79. West MJ. Stereological methods for estimating the total number of neurons and synapses:
issues of precision and bias. Trends Neurosci. 1999;22(2):51-61.
80. Gundersen HJ, Jensen EB. The efficiency of systematic sampling in stereology and its
prediction. Journal of microscopy. 1987;147(Pt 3):229-63.
81. Shvadchak VV, Falomir-Lockhart LJ, Yushchenko DA, Jovin TM. Specificity and kinetics of
alpha-synuclein binding to model membranes determined with fluorescent excited state
intramolecular proton transfer (ESIPT) probe. J Biol Chem. 2011;286(15):13023-32.
82. Scholtz JM, Qian H, York EJ, Stewart JM, Baldwin RL. Parameters of helix-coil transition
theory for alanine-based peptides of varying chain lengths in water. Biopolymers.
1991;31(13):1463-70.
83. Shvadchak VV, Yushchenko DA, Pievo R, Jovin TM. The mode of alpha-synuclein binding
to membranes depends on lipid composition and lipid to protein ratio. FEBS Lett.
2011;585(22):3513-9.
84. Allen TM, Cleland LG. Serum-induced leakage of liposome contents. Biochimica et
biophysica acta. 1980;597(2):418-26.
85. Lovas S, Zhang Y, Yu J, Lyubchenko YL. Molecular mechanism of misfolding and
aggregation of Abeta(13-23). The journal of physical chemistry B. 2013;117(20):6175-86.
86. Lv Z, Condron MM, Teplow DB, Lyubchenko YL. Nanoprobing of the effect of Cu(2+)
cations on misfolding, interaction and aggregation of amyloid beta peptide. Journal of
neuroimmune pharmacology : the official journal of the Society on NeuroImmune
Pharmacology. 2013;8(1):262-73.
87. Lyubchenko YL, Shlyakhtenko LS. AFM for analysis of structure and dynamics of DNA and
protein-DNA complexes. Methods (San Diego, Calif). 2009;47(3):206-13.
88. Miyagi A, Ando T, Lyubchenko YL. Dynamics of nucleosomes assessed with time-lapse
high-speed atomic force microscopy. Biochemistry. 2011;50(37):7901-8.
89. Sun Z, Tan HY, Bianco PR, Lyubchenko YL. Remodeling of RecG Helicase at the DNA
Replication Fork by SSB Protein. Scientific reports. 2015;5:9625.
90. Brenner S, Horne RW. A negative staining method for high resolution electron microscopy
of viruses. Biochimica et biophysica acta. 1959;34:103-10.

1138

Figure Legends

Figure 1. Human and mouse aSyn have seven sequence mismatches. (A) Sequence alignment of human, mouse, and chimeric aSyn variants showing mismatches between h-aSyn WT and m-aSyn at positions 53, 87, 100, 103, 107, 121, and 122. Also highlighted are the positions of familial PD substitutions and known serine and tyrosine phosphorylation sites. (B) Schematic representation of human, mouse, and chimeric aSyn variants examined in this study.

Figure 2. Human/mouse mismatches at positions 121 and 122 affect aSyn neurotoxicity in cell culture. Primary midbrain cultures were transduced with adenovirus encoding human, mouse, and chimeric aSyn variants at MOIs adjusted to ensure equal expression levels. Additional cultures were untransduced ('control'). The cells were fixed, stained with antibodies specific for MAP2 and TH, and scored for dopaminergic cell viability. The data are presented as the mean ± SEM, n = 4 (A, B), n = 5 (C), or n = 3 (D-F). $*p<0.05$, $**p<0.01$, $***p<0.001$; square root transformation, one-way ANOVA followed by Tukey's multiple comparisons *post hoc* test.

Figure 3. Human/mouse mismatches at positions 121 and 122 affect aSyn-mediated dopaminergic cell death in rat midbrain. Rats (4-10 per group) were injected unilaterally into the SN with vectors encoding h-aSyn A53T, h-aSyn Chimera, m-aSyn, or m-aSyn Chimera at two different titers (1.3E13 gc/ml or 3.5E14 gc/ml). Eight weeks after surgery, coronal midbrain sections of the high-dose group were stained for aSyn in order to visualize transgene expression (A-E). Adjacent specimens were stained for TH (F-J), VMAT2 (K-O), Hu (P-T) or cresyl violet (U-Y). The first panel in each row shows a representative image of the contralateral uninjected side (A, F, K, P, U). Counts of TH+ nigral dopaminergic neurons are shown in Z for the lower vector dose and in AA for the higher vector dose. Data are presented as the

1163   percentage of intact side cell numbers ± SEM. **p<0.01; two-tailed Mann-Whitney U test. Scale

1164   bar, 50 µm (panels A-Y). *See also Figure S1.*

1165

Figure 4. Mouse-to-human substitutions at positions 121 and 122 enhance the ability of

m-aSyn to elicit dopaminergic fiber loss in rat striatum. Representative striatal rat brain

sections stained for TH or VMAT2 (4-6 per group) are shown in panels A-E or G-K, respectively,

as low-magnification images, and in panels A'-E' or G'-K', respectively, as corresponding high-

magnification images. The imaged sections were from animals injected with the highest vector

dose. The first panel of each row shows a representative image of the contralateral uninjected

side (A, A', G, G'). Quantified staining intensities of the striatum are displayed in F for TH and L

for VMAT2. Data are presented as the percentage of corresponding contralateral staining

intensity ± SEM. *p<0.05; two-tailed Mann-Whitney U test. Scale bar, 1 mm (panels A-K) or 50

µm (panels A'-K'). *See also Figure S2.*

1176

Figure 5. M-aSyn forms amyloid-like fibrils with a shorter lag time compared to human

and chimeric aSyn variants. The formation of amyloid-like fibrils was monitored in solutions of

monomeric h-aSyn WT, h-aSyn A53T, and m-aSyn (A), or h-aSyn A53T, h-aSyn Chimera, m-

aSyn Chimera, and m-aSyn (B) (35 µM of each). The protein solutions were incubated at 37 °C

with constant agitation and analyzed at various times for thioflavin T fluorescence. The graphs

show the mean normalized fluorescence (determined from 3 independent experiments; 3

technical replicates per experiment) plotted against the incubation time. The dotted lines above

and below each solid line correspond to the mean signal ± SEM. *See also Figure S3.*

1185

Figure 6. Human, mouse, and chimeric aSyn variants form amyloid-like fibrils with similar

dimensions but different morphologies. (A-D) Results of AFM analysis of fibrils prepared

from recombinant h-aSyn A53T (A), m-aSyn (B), h-aSyn Chimera (C), and m-aSyn Chimera (D).

Panels A(i) – D(i) on the left show typical AFM topographic images of fibrils formed by each

aSyn variant. Dashed boxes indicate areas shown at a higher magnification in a spectral color

scheme in panels A(ii) – D(ii). Panels A(iii)-D(iii) on the right show cross-section profiles.

Regions of the fibrils from which the profiles are taken are shown with white solid lines on the

AFM images in panels A(ii)-D(ii). Twisted fibrils were observed for m-aSyn (B) and h-aSyn

Chimera (C). Twisted and non-twisted fibrils account for 65% and 35% (respectively) of total

fibrils formed by m-aSyn, and 76% and 24% (respectively) of total fibrils formed by h-aSyn

Chimera. Cross-section profiles in panels A(iii) and D(iii) reveal heights of non-twisted fibrils (~8-

11 nm). Cross-section profiles 1 and 2 reveal heights of twisted fibrils at valley regions (~7-9

nm) and peak regions (~11-12 nm), respectively, in panels B(iii) and C(iii). (E, F) Histograms

showing distances between peaks of twisted fibrils formed by m-aSyn (E) and h-aSyn Chimera

(F). The distances were determined from cross-section profiles obtained along the long axis of

the fibrils. (G-L) Representative TEM images of fibrils formed by h-aSyn WT (G), h-aSyn A53T

(H), m-aSyn (I), h-aSyn Chimera (J), and m-aSyn Chimera (K). Scale bar, 100 nm. (L) Higher-

magnification images generated from boxed regions of panels G, I, and J show a pair of fibrils

wound around each other via a helical twist (I), two fibrils aligned in parallel (II), or individual

fibrils with a twisted morphology (III and IV).

1206

Figure 7. Human/mouse mismatches at positions 121 and 122 affect membrane-induced

aSyn aggregation and aSyn-mediated vesicle disruption. (A) Solutions of recombinant aSyn

variants were incubated with increasing concentrations of SUVs composed of egg PG:PC (1:1

mol:mol) and analyzed by far-UV CD to determine $[\theta]_{MR,222}$. The data were fit to equation 3 (see

'Experimental Procedures'), and values for KD, minimum $[\theta]_{MR,222}$, and maximum α-helical

content were determined from the values of the fit parameters (Table S1). (B) Monomeric aSyn

variants were incubated with PG:PC SUVs (protein-to-lipid ratio, 1:20 mol/mol) at 37 °C for 72 h.

Left: Representative Western blot image (the vertical line to the right of the image shows the

region of the blot with bands corresponding to oligomeric forms of aSyn). Right: Bar graph

showing the ratio of total oligomer band intensity to monomer band intensity determined for

each sample. (C) Calcein-loaded SUVs were incubated in the absence ('ctrl') or presence of

aSyn variants. Calcein release was monitored via fluorescence measurements using excitation

and emission wavelengths of 485 nm and 515 nm, respectively. Left: graph showing % dye

release versus time, with 100% release determined as the signal obtained from vesicles treated

with Triton X-100. Right: table listing P values determined for vesicle permeabilization data

obtained at the 120 h time point (a more complete listing of P values at different time points is

presented in Table S2). The data in (B) and (C) are presented as the mean ± SEM, n = 4 (B) or

n = 3 (C). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, one-way ANOVA followed by Tukey's

multiple comparisons *post hoc* test (a square root transformation was carried out on the data in

panel C).

Figure 1

A.

	1	10	20	30	40	50	60	70
h-asyn	MDVFMKGLSK	AKEGVVAAAE	KTKQGVAEA**A**	GKTKEGVLYV	GSKTK**E**GVVH	**GVA**TVAEKTK	EQVTNVGGAV
m-asyn	....	....	....	....	....	..T.......	....

	71	80	90	100	110	120	130	140
h-asyn	VTGVTAVAQK	TVEGAG**S**IAA	ATGFVKKDQL	GKNEEGAPQE	GILEDMPVDP	DNEA**Y**EMP**S**E	EG**Y**QD**Y**EPEA
m-asyn	....	......N...	.........M	..G...Y...	....	GS........	....

X = disease causing mutation A30P, E46K, G51D, A53E and A53T

S = known serine phosphorylation residues

Y = known tyrosine phosphorylation residues

h-asyn aa sequence

m-asyn aa sequence

B.

Figure 2

Figure 3

Z. Stereology - 1.3E13 gc/ml

AA. Stereology - 3.5E14 gc/ml

Figure 4

Figure 5.

A

B

Figure 6

Figure 7