Title: *Synechocystis* KaiC3 displays temperature and KaiB dependent ATPase activity and is

important for growth in darkness

Running title: Biochemical characterization of *Synechocystis* KaiC3

Authors: Anika Wiegard[1,*,#], Christin Köbler[2], Katsuaki Oyama[3,‡], Anja K. Dörrich[4], Chihiro Azai[3,5],

Kazuki Terauchi[3,5,#], Annegret Wilde[2], Ilka M. Axmann[1]

Author affiliations:

[1] Institute for Synthetic Microbiology, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich

Heine University Duesseldorf, Duesseldorf, Germany

[2] Institute of Biology III, Faculty of Biology, University of Freiburg, Freiburg, Germany

[3] Graduate School of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

[4] Institute for Microbiology and Molecular Biology, Justus-Liebig University, Giessen, Germany

[5] College of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

* present address: Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm,

Sweden

‡ present address: Graduate School of Medicine, Kobe University, Kobe, Hyougo, Japan

Correspondence and requests for materials should be addressed to A. Wiegard (email:

anika.wiegard@ki.se) or K. Terauchi (email: terauchi@fc.ritsumei.ac.jp)

Abstract:

Cyanobacteria form a heterogeneous bacterial group with diverse lifestyles, acclimation strategies and differences in the presence of circadian clock proteins. In *Synechococcus elongatus* PCC 7942, a unique posttranslational KaiABC oscillator drives circadian rhythms. ATPase activity of KaiC correlates with the period of the clock and mediates temperature compensation. *Synechocystis* sp. PCC 6803 expresses additional Kai proteins, of which KaiB3 and KaiC3 proteins were suggested to fine-tune the standard KaiAB1C1 oscillator. In the present study, we therefore characterized the enzymatic activity of KaiC3 as a representative of non-standard KaiC homologs *in vitro*. KaiC3 displayed ATPase activity, which were lower compared to the *Synechococcus elongatus* PCC 7942 KaiC protein. ATP hydrolysis was temperature-dependent. Hence, KaiC3 is missing a defining feature of the model cyanobacterial circadian oscillator. Yeast two-hybrid analysis showed that KaiC3 interacts with KaiB3, KaiC1 and KaiB1. Further, KaiB3 and KaiB1 reduced *in vitro* ATP hydrolysis by KaiC3. Spot assays showed that chemoheterotrophic growth in constant darkness is completely abolished after deletion of Δ*kaiAB1C1* and reduced in the absence of *kaiC3*. We therefore suggest a role for adaptation to darkness for KaiC3 as well as a crosstalk between the KaiC1 and KaiC3 based systems.

Importance: The circadian clock influences the cyanobacterial metabolism and deeper understanding of its regulation will be important for metabolic optimizations in context of industrial applications. Due to the heterogeneity of cyanobacteria, characterization of clock systems in organisms apart from the circadian model *Synechococcus elongatus* PCC 7942 is required. *Synechocystis* PCC 6803 represents a major cyanobacterial model organism and harbors phylogenetically diverged homologs of the clock proteins, which are present in various other non-

43   cyanobacterial prokaryotes. By our *in vitro* studies we unravel the interplay of the multiple

44   *Synechocystis* Kai proteins and characterize enzymatic activities of the non-standard clock

45   homolog KaiC3. We show that the deletion of *kaiC3* affects growth in constant darkness

46   suggesting its involvement in the regulation of non-photosynthetic metabolic pathways.

Introduction:

Cyanobacteria have evolved the circadian clock system to sense, anticipate and respond to

predictable environmental changes based on the rotation of Earth and the resulting day-night

cycle. Circadian rhythms are defined by three criteria: (i) oscillations with a period of about 24

hours without external stimuli, (ii) synchronization of the oscillator with the environment and (iii)

compensation of the usual temperature dependence of biochemical reactions, so that the period

of the endogenous oscillation does not depend on temperature in a physiological range (2). The

cyanobacterial circadian clock system has been studied in much detail in the unicellular model

cyanobacterium *Synechococcus elongatus* PCC 7942 (hereafter *S. elongatus*). Its core oscillator is

composed of three proteins, which are unique to prokaryotes: KaiA, KaiB, and KaiC (from now on

KaiA, KaiB and KaiC; please note that some of the hereafter cited information were gained

studying proteins from *Thermosynechococcus elongatus* BP-1 though) (3). The level of KaiC

phosphorylation and KaiC's ATPase activity represent the key features of the biochemical

oscillator. KaiA stimulates autophosphorylation and ATPase activity of KaiC, whereas KaiB binding

to the complex inhibits KaiA action, stimulates autodephosphorylation activity and reduces

ATPase activity of KaiC (4-6). As a consequence of dynamic interactions with KaiA and KaiB, KaiC

rhythmically phosphorylates and dephosphorylates with a 24-hour period (3).

KaiC consists of two replicated domains (CI and CII) which assemble into a hexamer forming an N-

terminal CI ring and a C-terminal CII ring (7-9). Phosphorylation takes place in the CII ring (10),

whereas ATP hydrolysis occurs in both rings (11). In the CII ring, ATP hydrolysis is part of the

dephosphorylation mechanism (12, 13). ATP hydrolysis in the CI ring correlates with the period of

the clock and temperature compensation and is further required for a conformational change of

KaiC, which allows binding of KaiB (5, 14, 15). The levels of phosphorylation and ATP hydrolysis of

KaiC serve as the read-out for regulatory proteins, which orchestrate the circadian output (16,

17). For a recent review on the functioning of the KaiABC system, see Swan *et al.* (18).

In a natural day and night cycle, *S. elongatus* orchestrates its metabolism in a precise temporal

schedule, with the metabolism being adjusted by the clock but also feeding input information to

the clock (19-21). Knockout of the *S. elongatus kai* genes leads to a growth disadvantage. When

grown in competition with the wild type under light-dark conditions, the clock deficient mutant

cells are eliminated from the culture (22). Growth in light-dark cycles is also impaired by deletion

of the regulator of phycobilisome association A (RpaA) (19, 23), the master transcription factor of

circadian gene expression. In addition, studies on single nucleotide polymorphisms identified the

rpaA gene as one of three genes responsible for the faster growth of *Synechococcus elongatus*

UTEX 2973 compared to *S. elongatus* (24). Using a transposon library, Welkie *et al.* (25) showed

that KaiA, despite being non-essential for growth in light-dark cycles, strongly contributes to the

fitness of *S. elongatus* under these conditions. Decreased fitness most likely occurs due to

reduced phosphorylation of RpaA in the *kaiA* knockout strain (25). Overall, these data

demonstrate the value of the cyanobacterial clock system for metabolic orchestration under

natural conditions, with the clock system considered especially important for the transition from

light to darkness (26).

Cyanobacteria represent one of the most diverse prokaryotic phyla (27) and little is known about

timekeeping mechanisms in other cyanobacteria than *S. elongatus*. A core diurnal genome has

been described (28), but temporal coordination varies and, based on genomic analyses, large

variations in the cyanobacterial clock systems can be expected (29-34).

The cyanobacterial model strain *Synechocystis* sp. PCC 6803 (from now on *Synechocystis*) contains

a standard *kai* operon, encoding homologs of the *kaiA*, *kaiB* and *kaiC* genes (in the following

designated *kaiA$_{6803}$, kaiB1* and *kaiC1*) as well as two additional copies each of *kaiB* and *kaiC*

(designated *kaiB2*, *kaiB3* and *kaiC2, kaiC3* (35). Please note that naming of KaiC3/KaiC2 and

KaiB3/KaiB2 is not consistent in the literature. In this paper, we name the *Synechocystis kai* genes

according to Aoki and Onai, Wiegard *et al.* and Schmelling *et al.* (28, 30, 32). The *kaiC2* and *kaiB2*

genes form an operon, whereas *kaiC3* and *kaiB3* are orphan genes. Based on phylogenetic

reconstruction analysis, the KaiB and KaiC proteins of *Synechocystis* were allocated in three

phylogenetically different subclasses (32, 36). KaiC1 and KaiB1 display 82 % and 88 % amino acid

identity to *S. elongatus* KaiC and KaiB, respectively. Amino acid identity to *S. elongatus* proteins

is lower for the other *Synechocystis* Kai homologs, namely, 37 % for KaiC2, 52 % for KaiB2, 55 %

for KaiC3 and 48 % for KaiB3. The KaiC homologs show differences in their C-terminus: Both, KaiC2

and KaiC3 show low conservation of the A-loop. KaiC2 further differs from KaiC3 by displaying

modified phosphorylation sites (two serine residues instead of serine and threonine in KaiC1 and

KaiC3) (32).

Nevertheless, all three KaiC proteins from *Synechocystis* show high conservation of the kinase

motif in the CII domain and all three proteins were shown to exhibit autophosphorylation activity

(28, 32). KaiA$_{6803}$ stimulates the autophosphorylation of KaiC1, but does not affect the

phosphorylation of the two other KaiC homologs. Based on sequence analysis and experimental

validation of kinase activity, Schmelling *et al.* (28) concluded that autokinase and

autophosphatase activities are highly conserved features of all cyanobacterial KaiC homologs.

Likewise, ATPase motifs in the N-terminal CI domain of KaiC are highly conserved in all

cyanobacterial KaiC homologs (28). However, ATP hydrolysis was only characterized for several

KaiC1 homologs and a KaiC2 homolog from *Legionella pneumophila* (11, 37-39).

In *Synechocystis* deletion of the *kaiAB1C1* gene cluster causes reduced growth in light-dark

rhythms, even when grown as single culture, demonstrating a more pronounced phenotype

compared to *S. elongatus*, where only a competitive growth disadvantage was observed (40).

Deletion of *rpaA* also reduced growth in light-dark cycles, which further implies metabolic

orchestration by the *Synechocystis* timer (23). The reported number of oscillating genes in

Synechocystis varies largely between different studies, which is likely due to differences in growth

conditions and strain variations (41-44). Aoki and Onai (30) suggested that KaiC3 and KaiB3

modulate the amplitude and period of the KaiAB1C1 oscillator, whereas disruption of *kaiC2B2*

implied a non-clock related function for KaiC2 and KaiB2. There are different *Synechocystis* lab

strain variants in use, which show phenotypic variation, for example in their glucose tolerance,

motility and tolerance to abiotic stress (45). In the *Synechocystis* strain used in this study (PCC-M,

re-sequenced, (46)) the *kaiC2B2* cluster cannot be deleted (40), which further implies a non-clock-

related essential function. The function of KaiC3 has not been addressed so far. BLAST analysis

identified KaiC3 homologs to be present in addition to KaiC1 in about one third of cyanobacterial

genera. KaiC3 homologs were also found in non-cyanobacterial eubacteria and archaea, where

they show a higher distribution than the phylogenetically different KaiC1 and KaiC2 homologs

(28).

In this study, we therefore aim at a detailed biochemical characterization of the putative clock

component KaiC3 and the role of *Synechocystis* KaiB homologs in modulation of KaiC3 function.

Characterization of the ATPase activity of KaiC3 was of special interest, because ATP hydrolysis

defines the period length and temperature compensation of the Kai oscillator (5). We further

demonstrate that the Δ*kaiC3* mutant strain has a growth defect under chemoheterotrophic

growth conditions, which is similar, but less pronounced compared to Δ*kaiAB1C1* and Δ*rpaA*

strains (23, 40). Our data support the idea of a function of KaiC3 and KaiB3 in fine-tuning the

central oscillator composed of KaiA$_{6803}$, KaiB1 and KaiC1 in *Synechocystis*.

(This article was submitted to an online preprint archive (1))

Results:

Recombinant KaiC3 displays only low ATP synthase activity

Previous bioinformatic analysis predicted that kinase, dephosphorylation and ATPase activities

are conserved in KaiC3 (28). So far, only the kinase activity of KaiC3 was experimentally confirmed

which differed from the activity of *S. elongatus* KaiC by lacking stimulation by KaiA and KaiA$_{6803}$

(32). Therefore, we aimed at the characterization of further predicted activities of KaiC3.

The first step in KaiC dephosphorylation is regarded as a reversal of phosphorylation: KaiC

transfers its bound phosphoryl group to ADP and thereby synthesizes ATP (12, 13). To investigate

reversibility of intrinsic phosphorylation, we tested whether KaiC3 can synthesize [α-^{32}P]ATP from

[α-^{32}P]ADP (Fig. 1). As control, we used phosphorylated and dephosphorylated KaiC (Fig. S1.).

Immediately after adding [α-^{32}P]ADP, all KaiC proteins started to synthesize [α-^{32}P]ATP.

Phosphorylated as well as non-phosphorylated KaiC both showed higher initial ATP synthesis than

KaiC3 (Fig. 1). After incubating phosphorylated and dephosphorylated KaiC for 2 hours at 30°C

with [α-^{32}P]ADP, a relative [α-^{32}P]ATP level of 26-28 % was detected. KaiC3 showed low intrinsic

ATP synthesis, but formation of relative [α-^{32}P]ATP was not significantly higher than in the control

with cold ATP. This can be explained by two hypotheses: (i) compared to *S. elongatus* KaiC, KaiC3

showed lower phosphotransferase activity or (ii) higher consumption of the produced ATP. The

latter would require that KaiC3 exhibits higher ATPase activity than KaiC. Therefore, we next

examined ATP hydrolysis activity of KaiC3.

KaiC3 displays ATPase activity

ATP hydrolysis can be measured by quantifying ADP production by KaiC over time (5).

Conservation of WalkerA and WalkerB motifs in the CI domain of KaiC3 proteins suggested their

capability to hydrolyze ATP (28). We purified recombinant hexameric KaiC3 wild type protein

fused to an N-terminal Strep-tag (Strep-KaiC3, Fig. S4) and confirmed the predicted ATPase

activity *in vitro*. Based on the measured bulk ATPase activity we calculated that 8.5 ± 1.0 ADP

molecules per monomer and day (mean ± s.d., Fig. 2) were produced by Strep-KaiC3. We were

not able to purify stable recombinant KaiC1 with sufficient purity and therefore used the highly

similar KaiC ortholog from *S. elongatus* for comparison. The ATPase activity of Strep-KaiC3 was

about 45 % of the value for KaiC from *S. elongatus* (19.1 ± 3.3 ADP per day (5)). Hence, the above-

observed lower level of net ATP production by KaiC3 was not based on a higher ATP consumption,

but due to lower dephosphorylation activity *per se*. In KaiC, the ATPase activity changes after

substitution of the phosphorylation sites (5). We therefore generated variants of Strep-KaiC3, in

which residues S423 and T424 are either replaced with aspartate and glutamate (Strep-KaiC3-DE)

or replaced with two alanine residues (Strep-KaiC3-AA). Strep-KaiC3-AA, showed more than two

fold increased ATPase activity, whereas ATP hydrolysis by Strep-KaiC3-DE was only slightly

different from the wild-type protein (Fig. 2) and not reduced as reported for *S. elongatus* KaiC-DE

(5).

KaiC3 interacts with KaiB3 and components of the standard KaiAB1C1 oscillator

In vitro and *in silico* studies suggested that KaiA$_{6803}$, KaiB1 and KaiC1 form the standard clock

system of *Synechocystis* (30, 32). Since Aoki and Onai (30) suggested that KaiC3 might modulate

the main oscillator function, we performed protein-protein interaction studies in order to reveal

a possible crosstalk between the multiple Kai proteins. First, interaction was determined by yeast-

two hybrid analysis using KaiC1, KaiC3, KaiB1 and KaiB3 fused to AD and BD domains, respectively.

The color change of the colonies indicates β-galactosidase activity and is an estimate for the

interaction of the respective two proteins. As expected, these experiments showed self-

interaction of KaiC3 (Fig. 3A), since KaiC homologs are known to form hexamers (7, 9). In addition,

an interaction of KaiC3 with KaiB3 was detected (Fig. 3A) which is in line with bioinformatic

analysis showing frequent co-occurrence of *kaiC3* and *kaiB3* genes in genomes (28). We could not

detect an interaction between KaiC3 and KaiA$_{6803}$ by yeast-two hybrid analysis (Fig. 3B), but

detected a heteromeric interaction between KaiC1 and KaiC3 using different protein fusion

variants (Fig. 3C). These results confirm previously published data, which showed (i) co-

purification of KaiA with KaiC1 but not with KaiC3 and (ii) a weak interaction between the two

KaiC homologs KaiC1 and KaiC3 in ex-vivo pulldown analysis followed by Western Blot analysis

(32). Besides KaiC1, also the cognate KaiB protein, KaiB1, showed an interaction with KaiC3 in our

analysis (Fig. 3C), corroborating the hypothesis that there is a cross talk between the putative

KaiC3-B3 system and the core oscillator KaiAB1C1. Such a possible cross talk via the KaiB proteins

was further supported by our *in vitro* pull-down assays, in which KaiC3 interacted with KaiB3 (Fig.

S3B,D) and further with KaiB1 (Fig. S3B,D). Also KaiC1 interacted with both, KaiB1 and KaiB3

homologs (Fig. S3,A,C). One must take into account that based on the *in vitro* pull-down assays

using *Synechocystis* whole cell extracts we cannot exclude indirect interactions. As we have shown

that KaiC1 and KaiC3 interact with each other, both proteins could bind as a hetero-hexamer to

the GST-tagged KaiB proteins and therefore co-elute from the affinity matrix.

ATPase activity of KaiC3 is reduced in the presence of KaiB1 and KaiB3

The interaction of KaiC3 with KaiB1 and KaiB3 (Fig. 3 and Fig. S3) suggested a regulation of KaiC3

activity by these two KaiB proteins. We therefore measured ATP hydrolysis by Strep-KaiC3 in the

presence of 0.04 mg ml^{-1} KaiB1 and KaiB3 proteins, respectively (Fig. 4A). After size exclusion

chromatography KaiB1 was only eluted as a tetramer (44 and 72 kDa, respectively, depending on

the column), whereas KaiB3 was eluted as monomer (13/23 kDa) and tetramer (41/70 kDa) (Fig.

S4). Therefore, the monomeric and tetrameric KaiB3 fractions were tested separately. In all

measurements, the ATPase activity was linear and showed no oscillations. ATP hydrolysis was

reduced by 55 % after the addition of the KaiB3 monomer, but not affected by the KaiB3 tetramer.

The KaiB1 tetramer also inhibited the ATPase activity of Strep-KaiC3 (35 % reduction compared

to Strep-KaiC3 alone, Fig. 4). These data imply a trend, that KaiB1 has less effect on KaiC3's ATPase

activity than the KaiB3 monomer, but the differences were not statistically significant (Fig. 4).

Because ATPase activity of KaiC is influenced by KaiA (5), we also investigated the effect of $KaiA_{6803}$

on KaiC3, but in line with the lack of interaction in our yeast two-hybrid analysis data (Fig. 3B),

ATPase activity of Strep-KaiC3 was not significantly affected by $KaiA_{6803}$ (Fig. 4A). Fig. 4C

summarizes the Kai protein interactions and regulation of KaiC3's ATPase activity by KaiB proteins.

ATPase activity of KaiC3 is temperature dependent

True circadian clocks are characterized by temperature compensated oscillations, which ensure

robust time measurements under temperature fluctuations. In the *S. elongatus* KaiABC clock,

overall temperature compensation is derived from KaiC's ATPase activity, which is stable between

25 and 35 °C (Q_{10}=1.2, (5)). We therefore asked whether N-Strep-KaiC3, as a representative of

non-standard KaiC homologs, shows temperature compensation as well. Measurements at 25, 30

and 35 °C, however, revealed a temperature-dependent ADP production by KaiC3 (Q_{10}=2.4, Fig.

4B; activation energy = 66.5 kJ mol^{-1}, Fig. S5A). Hence, KaiC3 is lacking a characteristic feature of

circadian oscillators. In accordance, dephosphorylation of Strep-KaiC3 was higher at 25 °C than at

30 and 35 °C (Fig. S5B)

The non-standard KaiC3 protein supports growth of *Synechocystis* cells in the dark

As *in vitro* analyses suggested an interaction of KaiC3 with KaiB1 and KaiC1, we aimed at

elucidating a putative role of KaiC3 and its crosstalk with the KaiC1-based system in the cell. Clock

factors are reported to be essential for cell viability in light-dark cycles (19, 23, 25). For assessing

growth of mutant strains lacking *kai* genes, we used a spot plating assay (48). In this assay, the

amount of cells plated on agar in serial dilutions is compared to the amount of cells which are

able to form a colony. The plating efficiency then allows us to compare the sensitivity of cells to

changing light conditions. Notably, this method allows only limited assertions on the growth rate

of different strains. Deletion of the *kaiAB1C1* cluster of *Synechocystis* resulted in lower cell

viability in light-dark cycles (40). This was even more pronounced under photomixotrophic (0.2 %

glucose) compared to photoautotrophic conditions, whereas deletion of *kaiC3* had no effect on

cell viability in light-dark cycles (40). As KaiC3 homologs are also present in non-photosynthetic

bacteria (28), we were interested in the growth of the Δ*kaiC3* strain in the dark. The *Synechocystis*

wild-type strain used here, in contrast to previous studies (47), is able to grow in complete

darkness when supplemented with glucose (23). We therefore analyzed the viability of Δ*kaiC3*

cells via spot assays in constant light and in complete darkness on agar plates containing 0.2 %

glucose (Fig. 5A). Under photomixotrophic growth conditions with continuous illumination, wild

type and Δ*kaiC3* showed similar viability, whereas the viability of the Δ*kaiAB1C1* strain was

reduced, reflected by a lower plating efficiency. Further, in complete darkness, wild type

displayed detectable growth, while growth of Δ*kaiAB1C1* was abolished completely. Sensitivity of

the Δ*kaiAB1C1* mutant to mixotrophic conditions in the light and lack of growth in the dark

highlights the importance of the KaiAB1C1 oscillator for the switch between these two different

metabolic modes of *Synechocystis*. The Δ*kaiC3* strain could grow in the dark but showed some

impairments. The plating efficiency in serial dilutions indicates similar viability of the wild-type

and Δ*kaiC3* deletion strain (Fig. 5A). However, dark growth seemed to be more impaired in the

Δ*kaiC3* strain when compared to the wild type. For a more quantitative assessment, intensity

measurements of single spots of the *Synechocystis* wild type and *kaiC3* deletion strain were

performed (Fig. 5B), validating the growth impairment of the *kaiC3* deletion strain in complete

darkness. Thus, KaiC3 seems to be linked to dark adaption of *Synechocystis* cells, yet not as

essential as the core oscillator KaiAB1C1.

Discussion:

Characterization of enzymatic activities of KaiC3

Previous sequence analysis suggested that the three enzymatic activities of KaiC, which are autophosphorylation, autodephosphorylation and ATPase activity, are conserved in all cyanobacterial KaiC proteins (28). This hypothesis is supported by experiments demonstrating autophosphorylation of all *Synechocystis* KaiC proteins (32), different cyanobacterial KaiC1 and KaiC3 homologs (28, 37, 39), and even non-cyanobacterial KaiC homologs from *Rhodopseudomonas palustris* (38), *Legionella pneumophila* (48) as well as two thermophilic Archaea (28). In the current paper, we provide experimental evidence that ATPase activity is conserved in KaiC3 as well. However, the level of the ATPase activity seems to vary between KaiC proteins. While ATPase activity of the KaiC from *Prochlorococcus marinus* MED4 was reported to have the same activity as KaiC (37), we show here that the ATPase activity of KaiC3 is lower. In contrast, ATP hydrolysis activity of the KaiC2 homolog from *Legionella pneumophila* and KaiC1 homolog from *Gloeocapsa* sp. PCC 7428 were reported to be elevated when compared to the standard KaiC protein (38, 39).

ATPase activity of KaiC3 further differed from that of KaiC, by lacking temperature compensation (Fig. 4). Therefore, KaiC3 cannot be the core component of a true circadian oscillator. This is further supported by differences in the ATP hydrolysis rate between the phosphomimetic variants of KaiC and KaiC3. ATPase activity of the true clock protein KaiC from *S. elongatus* depends on its phosphorylation status. In the KaiC-AA variant, which mimics the dephosphorylated state of KaiC, the ATPase activity is higher than average, whereas, in KaiC-DE, mimicking the phosphorylation state, it is diminished (5). In our analysis, ATP hydrolysis by Strep-KaiC3-AA was similarly increased

in comparison to Strep-KaiC3. However, Strep-KaiC3-DE did not show reduced activity as was

shown for KaiC-DE, implying that the phosphorylation state of KaiC3 does not influence KaiC3's

ATPase activity.

The ATPase activity of Strep-KaiC3 was affected decisively by the addition of KaiB3 and KaiB1. This

points at a putative mechanism how KaiC3 might fine-tune the standard oscillator: Binding of

KaiB1 to KaiC3 will reduce the free KaiB1 concentration in the cell and thereby influence the

KaiAB1C1 oscillator. Both KaiB homologs led to decreased ATPase activity of Strep-KaiC3 (Fig. 4A).

One needs to take into consideration that we used 0.04 mg ml^{-1} KaiB protein for all assays. As

KaiB1 was present only as tetramer in the used fractions, this resulted in a molar concentration

of 0.82 µM KaiB1 tetramers versus 3.36 µM KaiB3 monomers, respectively. *S. elongatus* KaiB and

KaiB3 both exist as monomer or tetramer in solution and KaiB is known to bind as a monomer to

KaiC, (49-51). In our analysis where we separated monomeric and tetrameric KaiB3 forms, only

monomeric KaiB3 showed an effect on ATP hydrolysis by KaiC3. Therefore, addition of a similar

monomeric concentration (3.28 µM KaiB1 and 3.36 µM KaiB3) was important to compare the

effect of KaiB1 and KaiB3. As a tetramer, KaiB subunits adopt a different unique fold, which was

also observed in crystals of KaiB1 (52). Residues K57, G88 and D90, which are important for fold

switching and hence transition between tetrameric and monomeric KaiB (51), are also conserved

in KaiB3. We, therefore, cannot exclude that addition of a higher molar concentration of KaiB3

tetramers might affect ATPase activity of KaiC3. However, we show only *in vitro* data here. In the

cell, protein concentrations as well as spatial and temporal separation of the different proteins

might have an influence on these interactions. In addition, we do not take possible heteromeric

interactions among the KaiB proteins into account.

Function of KaiC3 *in an extended network*

The interaction studies shown here imply a crosstalk between KaiC1 and KaiC3 (Fig. 4C). KaiC1 is

believed to form a standard oscillator together with $KaiA_{6803}$ and KaiB1 (30, 32), which is

supported by the KaiC1-KaiB1 interaction observed here (Fig. 3C, Fig. S3). We further hypothesize

that KaiC3 acts in a separate non-circadian regulatory system together with KaiB3. In accordance

with bioinformatic analysis, which showed a significant co-occurrence of KaiB3 and KaiC3 in

Cyanobacteria (28), KaiB3 had a stronger effect on the ATPase activity of KaiC3 than KaiB1 (Fig.

4A). In the cell, the KaiAB1C1 and KaiB3C3 systems could work independently from each other.

However, KaiC3 is able to form hetero-oligomers with KaiC1 and was shown to interact with KaiB1

as well. Therefore, interference between the KaiC3-KaiB3 system and the proteins of the standard

oscillator is very likely. Further, we exclude that $KaiA_{6803}$ is involved in the putative crosstalk by

following reasons: (i) $KaiA_{6803}$ did neither stimulate ATP hydrolysis nor kinase activity of KaiC3 (this

paper and Wiegard *et al.* (32)); (ii) We were not able to show an interaction of KaiC3 and $KaiA_{6803}$

using different approaches (this paper and Wiegard *et al.* (32)); (iii) KaiA interacting residues are

not conserved in cyanobacterial KaiC3 homologs, and (iv) KaiC3 homologs are present in

organisms which do not harbor KaiA (28).

The growth defect of the mutant strains, Δ*kaiAB1C1* and Δ*kaiC3,* in complete darkness suggests

that the here proposed putative KaiC3-KaiB3 system and the KaiAB1C1 oscillator may target

similar cellular functions. The metabolic network of cyanobacteria is described as temporally

partitioned with extensive effects of day-night transitions, involving shifts in ATP and reductant

levels and alterations of the carbon flux (25). In *S. elongatus*, environmental signals can be fed

into the main clock output system via the transcriptional regulator RpaB (53). In contrast to *S.*

elongatus, *Synechocystis* is able to grow chemoheterotrophically, which adds another layer of

complexity to day-night transitions and demand for further regulatory elements. The observed

impaired growth of the *kaiC3* mutant in darkness supports the idea of KaiC3 and KaiB3 functioning

as such additional elements to adjust the state of the main *Synechocystis* oscillator. Conversely,

it is also possible that KaiC3 function is controlled by the KaiAB1C1 clock system. Köbler *et al.* (23)

demonstrated that solely KaiC1, but not KaiC3, interacts with the main output histidine kinase

SasA in the *Synechocystis* timing system. Thus, in *Synechocystis*, only the main oscillator feeds

timing information into the SasA-RpaA output system to control the expression of many genes

involved in dark growth (23). The output pathway for KaiC3 is unknown so far and it might be

possible that the only function of KaiC3 is to modulate the function of the main oscillator in

response to a yet unknown input factor.

Material and Methods:

Cloning, expression and purification of recombinant Kai proteins

Genes encoding KaiB1 (ORF *slr0757*) and KaiB3 (ORF *sll0486*) were amplified from genomic

Synechocystis wild type DNA using specific primers (Table S1) and Phusion Polymerase (New

England Biolabs). After restriction digest with BamHI and NotI, amplified fragments were inserted

into pGEX-6P1 (GE Healthcare) and the resulting plasmids were used for heterologous expression.

For production of recombinant KaiC as well as $KaiA_{6803}$, KaiB1, KaiB3 and KaiC3, we used pGEX-

based plasmids described in Wiegard *et al.* (32) (see also Table S2 for a list of all plasmids used in

this study). A detailed protocol of expression and purification can be found on protocols.io

(https://dx.doi.org/10.17504/protocols.io.48ggztw). Briefly, proteins were expressed as GST-

fusion proteins in *E. coli* BL21 [DE3] or NEB Express (New England Biolabs) and lysed in 50 mM

Tris/HCl (pH8), 150 mM NaCl, 0.5 mM EDTA, 1 mM DTT (+ 5 mM $MgCl_2$, 1 mM ATP for KaiC

proteins). Purification was performed via *batch* affinity chromatography using glutathione

agarose 4B (Macherey and Nagel) or glutathione sepharose 4B (GE Healthcare) in the same buffer.

Finally, the GST-tag was cleaved off using PreScission protease (GE Healthcare) in 50 mM Tris/HCl

(pH8), 150 mM NaCl, 1 mM EDTA, 1 mM DTT (+ 5 mM $MgCl_2$, 1 mM ATP for KaiC proteins). If

homogeneity of the proteins was not sufficient, they were further purified via anion exchange

chromatography using a MonoQ or ResourceQ column (GE Healthcare Life Sciences) using 50 mM

Tris/HCl (pH8), 1 mM EDTA, 1 mM DTT (+ 5 mM $MgCl_2$, 1 mM ATP for KaiC proteins) and a 0-1M

NaCl gradient.

To produce Strep-KaiC3 variants with amino acid substitutions, the *kaiC3* gene in the pGEX-kaiC3

vector was modified by site directed mutagenesis using the Quick-Change Site-Directed

Mutagenesis Kit (Stratagene) or Q5 Site-Directed Mutagenesis Kit (New England Biolabs). Base

triplets encoding S423 and T424 were changed to code for alanine or for aspartate and glutamate

resulting in *kaiC3-AA* and *kaiC3-DE* genes, respectively. To generate *kaiC3-catE1⁻catE2*, two

subsequent site-directed mutagenesis reactions were performed to exchange nucleotides

encoding E67 and E68 as well as nucleotides encoding E310 and E311 all with bases encoding

glutamine (all primers used for mutagenesis are listed in Table S1). Afterwards, *kaiC3* WT and

modified *kaiC3* genes were amplified with KOD-Plus-Neo polymerase (Toyobo) using pASK-kaiC3

primers (Table S1). Amplicons were digested with SacII and HindIII and ligated into the respective

restriction sites of pASK-IBA5plus (IBA Life sciences). For purification of recombinant Strep-KaiC,

the pASK-IBA-5plus based vector described in Oyama *et al.* (15) was used. Strep-KaiC3 proteins

were expressed in *E. coli* Rosetta gamiB (DE3) or Rosetta gami2 (DE3) cells (Novagen). Expression

of Strep-KaiC was carried out in *E. coli* DH5α. Cells were cultured in LB medium containing 100 µg

ml^{-1} ampicillin with vigorous agitation at 37 °C. At OD$_{600nm}$ 0.33-0.68 protein expression was

induced with 200 ng ml^{-1} anhydrotetracycline and the strains were further incubated as following:

Strep-KaiC: 7h at 37°C, Strep-KaiC3-WT: 5h at 35°C or 3.5h at 37°C, Strep-KaiC3-AA and Strep-

KaiC3-catE1$^-$catE2$^-$: 18°C overnight, Strep-KaiC3-DE: 25°C overnight. Cells were harvested and

lysed by sonication in ice-cold buffer W [20mM Tris/HCl (pH8), 150 mM NaCl, 5 mM MgCl$_2$, 1 mM

ATP (+ 2 mM DTT for Strep-KaiC3 proteins)] including protease inhibitors (protease inhibitor

cocktail, Roche or Nacalai). Soluble proteins were loaded on self-prepared columns packed with

Strep-Tactin XT superflow or Strep-Tactin Sepharose (IBA lifesciences) and purified under gravity

flow. After washing with buffer W, Strep-KaiC proteins were eluted with ice cold buffer W + 50

mM D(+)biotin (for Strep-Tactin XT Superflow) or ice cold buffer W + 2.5 mM desthiobiotin (for

Strep tactin Superflow). See https://dx.doi.org/10.17504/protocols.io.meac3ae for a detailed

protocol.

All proteins used for ATPase activity measurements were further purified via size exclusion

chromatography. Strep-KaiC3 and Strep-KaiC proteins were applied on a Sephacryl S300 HR

HiPrep 16/60 Sephacryl column (GE Healthcare) and separated in 20 or 50 mM Tris/HCl (pH 8.0),

150 mM NaCl, 2 mM DTT, 1 mM ATP and 5 mM MgCl2. For separation of KaiB1 and KaiB3, a

Sephacryl S200 HR HiPrep 16/60 column (GE Healthcare) or Superdex 200 Increase 10/30 GL

column (GE Healthcare) and 20 or 50 mM Tris/HCl (pH 8.0), 150 mM NaCl, 2 mM DTT as running

buffer were used. KaiA$_{6803}$ was purified on a Superdex 200 Increase 10/30 GL column (GE

Healthcare) in 20 mM Tris/HCl (pH 8.0), 150 mM NaCl, 2 mM DTT. See

https://dx.doi.org/10.17504/protocols.io.mdtc26n for further details.

ATPase activity

For ATPase measurements, KaiC proteins fused to an N-terminal Strep-tag were used and all Kai

proteins were purified via size exclusion chromatography (see above). 3.45 μM of Strep-KaiC3

variants were incubated in ATPase buffer (20 mM Tris/HCl (pH8), 150 mM NaCl, 1 mM ATP, 5 mM

$MgCl_2$) at 25 °C, 30 °C or 35 °C for 24 hours. To analyze the influence of $KaiA_{6803}$ and KaiB proteins

on KaiC3 ATPase activity, we mixed 0.2 mg ml^{-1} KaiC3 with 0.04 mg ml^{-1} $KaiA_{6803}$ or 0.04 mg ml^{-1}

KaiB and incubated the mixtures for 24 hours at 30 °C. Monomeric and oligomeric KaiB3 were

analyzed separately. To monitor ADP production, every 3, 4 or 6 hours 2 μl of the reaction mixture

were applied on a Shim-Pack-VP-ODS column (Shimadzu) and separated using 100 mM

phosphoric acid, 150 mM triethylamine, 1 % acetonitrile as running buffer. ADP production per

monomer KaiC and 24 hours was calculated using a calibration curve. A detailed protocol can be

found on protocols.io (https://dx.doi.org/10.17504/protocols.io.mebc3an). The Q_{10} value was

calculated from ATPase measurements at 25 °C and 35 °C, using the formula $Q_{10} =$

$\left(\frac{R_2}{R_1}\right)^{10°C/(T_2-T_1)}$

ATP synthase activity

To investigate dephosphorylation via ATP synthesis, we used KaiC proteins which were expressed

as GST-fusion proteins and subsequently cleaved off their GST-tag. 3 μM KaiC in ATP synthesis

buffer (20 mM Tris/HCl(pH8), 150 mM NaCl, 0.5 mM EDTA, 5 mM $MgCl_2$, 0.5 mM ATP) was mixed

with 0.8 μCi ml^{-1} [$α^{32}P$]ADP and stored at -20 °C or incubated for 2 hours at 30 °C. As a control,

the same experiment was performed in the presence of 0.5 mM ADP. After 20-fold dilution,

nucleotides in a 0.5 μl reaction mixture were separated via thin layer chromatography using TLC

PEI Cellulose F plates (Merck Millipore) and 1 M LiCl as solvent. [α-^{32}P]ADP and [γ-^{32}P]ATP were

separated in parallel to identify the signals corresponding to ADP and ATP, respectively. Dried

plates were subjected to autoradiography and signals were analyzed using a Personal Molecular

Imager FX system (Bio-Rad) and ImageLab software (Bio-Rad). For each reaction mixture, the

relative intensity of [α-^{32}P]ATP was calculated as percentage of all signals in the corresponding

lane. Because [α-^{32}P]ATP was already synthesized during mixing of the samples, the relative ATP

intensity measured in the -20 °C sample containing 0.5 mM ADP was subtracted for normalization.

The principle of this method is based on Egli *et al.* (13). A detailed protocol is available on

protocols.io (https://dx.doi.org/10.17504/protocols.io.48qgzvw).

Yeast two-hybrid assays

For yeast two-hybrid assays, vectors containing the GAL4 activation domain (AD) and the GAL4

DNA-binding domain (BD) were used. Genes of interest were amplified from *Synechocystis* wild

type genomic DNA with the Phusion Polymerase (NEB) according to manufacturer's guidelines.

Indicated restriction sites were introduced via the oligonucleotides listed in Table S1. Vectors and

PCR fragments were cut with the respective restriction enzymes (Thermo Fisher Scientific) and

the gene of interest was ligated into the vector, leading to a fusion protein with an AD- or BD-tag

either at the N- or C-terminus. All constructed plasmids are listed in Table S2. Transformation of

yeast cells was performed according to manufacturer's guidelines using the Frozen-EZ Yeast

Transformation II Kit (Zymo Research) and cells were selected on complete supplement mixture

(CSM) lacking leucine and tryptophan (-Leu -Trp) dropout medium (MP Biochemicals) at 30 °C for

3–4 days. Y190 (Clontech) cells were used for measuring β-galactosidase activity. Formed colonies

were spotted on a second plate (CSM -Leu -Trp) and incubated for 2 days. Afterwards a colony-

lift filter assay was performed as described by Breeden *et al*. (54). A detailed protocol can be

found on protocols.io (https://dx.doi.org/10.17504/protocols.io.v7ve9n6).

Bacterial strains and growth conditions

Wild type *Synechocystis* sp. PCC 6803 (PCC-M, re-sequenced, (46)) and the *kaiC3* deletion mutant

(40) were cultured photoautotrophically in BG11 medium (55) supplemented with 10 mM TES

buffer (pH 8) under constant illumination with 50 µmol photons $m^{-2}s^{-1}$ of white light (Philips TLD

Super 80/840) at 30 °C. Cells were grown either in Erlenmeyer flasks with constant shaking (140

rpm) or on plates (0.75 % Bacto-Agar, Difco) supplemented with 0.3 % thiosulfate. Detailed

recipes can be found on protocols.io (https://dx.doi.org/10.17504/protocols.io.wj5fcq6).

Spot assays

Experiments were performed as previously described (56). Strains were propagated

mixotrophically on BG11 agar plates with the addition of 0.2 % (w/v) glucose. Dilution series of

cell cultures started with OD_{750nm} 0.2 and OD_{750nm} 0.4, followed by incubation of the plates for 6

or 28 days under constant light conditions and in complete darkness, respectively. Plates were

scanned and single spot intensities were quantified using Quantity One (Bio-Rad). Measured

intensities were normalized to the intensity of the wild-type spot of the respective control grown

under continuous light.

Acknowledgement:

We thank Junko Moriwaki, Nancy Sauer, Jennifer Andres, Lukas Pohlig, Thomas Volkmer and

Megumi Fujimoto for technical assistance. This work was funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence

Strategy – EXC 2048/1 – Project ID: 390686111 to A. Wiegard and I.M. Axmann. The work was

financially supported by Heine Research Academies - Travel Grants, and EMBO ASTF 145 to A.

Wiegard, by grants AX 84/1-3 and EXC 1028 from the German Research Foundation to I.M.

Axmann, by JSPS Grants-in-Aid 16H00784, 17K19247 and 19K05833 to K. Terauchi, and by grant

WI2014/5-3 and WI2014/10-1 from the German Research Foundation to A. Wilde.

Author contributions:

I.M. Axmann, A. Wilde, A. Wiegard, C. Köbler and K. Terauchi conceived the project. A. Wiegard,

C. Köbler A.K. Dörrich and K. Oyama designed, performed and analyzed experiments. I.M Axmann,

A. Wilde and K. Terauchi supervised the study. All authors interpreted and discussed the data. A.

Wiegard, C. Köbler and A. Wilde wrote the manuscript. I.M. Axmann, C. Azai, K. Terauchi, A.K.

Dörrich, and K. Oyama commented essentially on the manuscript. All authors approved the

manuscript.

This manuscript contains supplementary information:

Supplementary methods

Table S1. Oligonucleotides used in this study.

Table S2. Plasmids used in this study.

Figure S1. Phosphorylation level of KaiC proteins used for ATP synthase activity assay shown in

Fig. 1.

Figure S2. Original scans of the KaiC3 interaction with KaiB3 and the proteins of the main oscillator

KaiC1, KaiB1.

Figure S3. Interaction of KaiB and KaiC proteins in pull down analysis.

Figure S4. Purification of KaiB and KaiC3 proteins used in this study.

Figure S5. ATPase activity and dephosphorylation of KaiC3 are temperature dependent

References:

References:

1.   Wiegard A, Köbler C, Oyama, K, Dörrich, AK, Azai, C, Terauchi, K, Wilde, A, Axmann, IM. 2019.

Synechocystis KaiC3 displays temperature and KaiB dependent ATPase activity and is important

for viability in darkness. bioRxiv doi:10.1101/700500:700500.

2.   Ditty JL, Williams SB, Golden SS. 2003. A cyanobacterial circadian timing mechanism. Annu Rev

Genet 37:513-43.

3.   Nakajima M, Imai K, Ito H, Nishiwaki T, Murayama Y, Iwasaki H, Oyama T, Kondo T. 2005.

Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation *in vitro*. Science

308:414-5.

4.   Iwasaki H, Nishiwaki T, Kitayama Y, Nakajima M, Kondo T. 2002. KaiA-stimulated KaiC

phosphorylation in circadian timing loops in cyanobacteria. Proc Natl Acad Sci U S A 99:15788-93.

5.   Terauchi K, Kitayama Y, Nishiwaki T, Miwa K, Murayama Y, Oyama T, Kondo T. 2007. ATPase activity

of KaiC determines the basic timing for circadian clock of cyanobacteria. Proc Natl Acad Sci U S A

104:16377-81.

6.   Kitayama Y, Iwasaki H, Nishiwaki T, Kondo T. 2003. KaiB functions as an attenuator of KaiC

phosphorylation in the cyanobacterial circadian clock system. EMBO J 22:2127-34.

7.   Hayashi F, Suzuki H, Iwase R, Uzumaki T, Miyake A, Shen JR, Imada K, Furukawa Y, Yonekura K,

Namba K, Ishiura M. 2003. ATP-induced hexameric ring structure of the cyanobacterial circadian

clock protein KaiC. Genes Cells 8:287-96.

8.   Mori T, Saveliev SV, Xu Y, Stafford WF, Cox MM, Inman RB, Johnson CH. 2002. Circadian clock

protein KaiC forms ATP-dependent hexameric rings and binds DNA. Proc Natl Acad Sci U S A

99:17203-8.

9.   Pattanayek R, Wang J, Mori T, Xu Y, Johnson CH, Egli M. 2004. Visualizing a circadian clock protein:

crystal structure of KaiC and functional insights. Mol Cell 15:375-88.

10.   Nishiwaki T, Iwasaki H, Ishiura M, Kondo T. 2000. Nucleotide binding and autophosphorylation of

508   the clock protein KaiC as a circadian timing process of cyanobacteria. Proc Natl Acad Sci U S A

509   97:495-9.

510   11.   Murakami R, Miyake A, Iwase R, Hayashi F, Uzumaki T, Ishiura M. 2008. ATPase activity and its

511   temperature compensation of the cyanobacterial clock protein KaiC. Genes Cells 13:387-95.

512   12.   Nishiwaki T, Kondo T. 2012. Circadian autodephosphorylation of cyanobacterial clock protein KaiC

513   occurs via formation of ATP as intermediate. J Biol Chem 287:18030-5.

514   13.   Egli M, Mori T, Pattanayek R, Xu Y, Qin X, Johnson CH. 2012. Dephosphorylation of the core clock

515   protein KaiC in the cyanobacterial KaiABC circadian oscillator proceeds via an ATP synthase

516   mechanism. Biochemistry 51:1547-58.

517   14.   Phong C, Markson JS, Wilhoite CM, Rust MJ. 2013. Robust and tunable circadian rhythms from

518   differentially sensitive catalytic domains. Proc Natl Acad Sci U S A 110:1124-9.

519   15.   Oyama K, Azai C, Nakamura K, Tanaka S, Terauchi K. 2016. Conversion between two

520   conformational states of KaiC is induced by ATP hydrolysis as a trigger for cyanobacterial circadian

521   oscillation. Sci Rep 6:32443.

522   16.   Shultzaberger RK, Boyd JS, Diamond S, Greenspan RJ, Golden SS. 2015. Giving Time Purpose: The

523   Synechococcus elongatus Clock in a Broader Network Context. Annu Rev Genet 49:485-505.

524   17.   Dong G, Yang Q, Wang Q, Kim YI, Wood TL, Osteryoung KW, van Oudenaarden A, Golden SS. 2010.

525   Elevated ATPase activity of KaiC applies a circadian checkpoint on cell division in *Synechococcus*

526   *elongatus*. Cell 140:529-39.

527   18.   Swan JA, Golden SS, LiWang A, Partch CL. 2018. Structure, function, and mechanism of the core

528   circadian clock in cyanobacteria. J Biol Chem 293:5026-5034.

529   19.   Diamond S, Jun D, Rubin BE, Golden SS. 2015. The circadian oscillator in Synechococcus elongatus

530   controls metabolite partitioning during diurnal growth. Proc Natl Acad Sci U S A 112:E1916-25.

531   20.   Welkie DG, Rubin BE, Diamond S, Hood RD, Savage DF, Golden SS. 2019. A Hard Day's

532   Night: Cyanobacteria in Diel Cycles. Trends Microbiol 27:231-242.

533   21.   Pattanayak G, Rust MJ. 2014. The cyanobacterial clock and metabolism. Curr Opin Microbiol 18:90-

534   5.

535   22.   Woelfle MA, Ouyang Y, Phanvijhitsiri K, Johnson CH. 2004. The adaptive value of circadian clocks:

536   an experimental assessment in cyanobacteria. Curr Biol 14:1481-6.

537   23.   Köbler C, Schultz SJ, Kopp D, Voigt K, Wilde A. 2018. The role of the Synechocystis sp. PCC 6803

538   homolog of the circadian clock output regulator RpaA in day-night transitions. Mol Microbiol

539   110:847-861.

540   24.   Ungerer J, Wendt KE, Hendry JI, Maranas CD, Pakrasi HB. 2018. Comparative genomics reveals the

541   molecular determinants of rapid growth of the cyanobacterium Synechococcus elongatus UTEX

542   2973. Proc Natl Acad Sci U S A doi:10.1073/pnas.1814912115.

543   25.   Welkie DG, Rubin BE, Chang YG, Diamond S, Rifkin SA, LiWang A, Golden SS. 2018. Genome-wide

544   fitness assessment during diurnal growth reveals an expanded role of the cyanobacterial circadian

545   clock protein KaiA. Proc Natl Acad Sci U S A 115:E7174-e7183.

546   26.   Diamond S, Rubin BE, Shultzaberger RK, Chen Y, Barber CD, Golden SS. 2017. Redox crisis underlies

547   conditional light-dark lethality in cyanobacterial mutants that lack the circadian regulator, RpaA.

548   Proc Natl Acad Sci U S A 114:E580-e589.

549   27.   Shih PM, Wu D, Latifi A, Axen SD, Fewer DP, Talla E, Calteau A, Cai F, Tandeau de Marsac N, Rippka

550   R, Herdman M, Sivonen K, Coursin T, Laurent T, Goodwin L, Nolan M, Davenport KW, Han CS, Rubin

551   EM, Eisen JA, Woyke T, Gugger M, Kerfeld CA. 2013. Improving the coverage of the cyanobacterial

552   phylum using diversity-driven genome sequencing. Proc Natl Acad Sci U S A 110:1053-8.

553   28.   Schmelling NM, Lehmann R, Chaudhury P, Beck C, Albers S-VV, Axmann IMM, Wiegard A. 2017.

554   Minimal Tool Set for a Prokaryotic Circadian Clock. bioRxiv doi:10.1101/075291.

555   29.   Axmann IM, Hertel S, Wiegard A, Dorrich AK, Wilde A. 2014. Diversity of KaiC-based timing systems

556   in marine Cyanobacteria. Mar Genomics 14C:3-16.

557   30.   Aoki S, Onai K. 2009. Circadian clocks of *Synechocystis* sp. strain PCC 6803, *Thermosynechococcus*

558   *elongatus*, *Prochlorococcus* spp., *Trichodesmium* spp. and other species, p 259-282. *In* Ditty JL,

559   Mackey SR, Johnson CH (ed), Bacterial circadian programs. Springer Berlin Heidelberg.

560   31.   Dvornyk V, Vinogradova O, Nevo E. 2003. Origin and evolution of circadian clock genes in

561   prokaryotes. Proc Natl Acad Sci U S A 100:2495-500.

562   32.   Wiegard A, Dörrich AK, Deinzer HT, Beck C, Wilde A, Holtzendorff J, Axmann IM. 2013. Biochemical

563   analysis of three putative KaiC clock proteins from *Synechocystis* sp. PCC 6803 suggests their

564   functional divergence. Microbiology 159:948-958.

565   33.   Johnson CH, Egli M. 2014. Metabolic compensation and circadian resilience in prokaryotic

566   cyanobacteria. Annu Rev Biochem 83:221-47.

567   34.   Holtzendorff J, Partensky F, Mella D, Lennon JF, Hess WR, Garczarek L. 2008. Genome streamlining

568   results in loss of robustness of the circadian clock in the marine cyanobacterium *Prochlorococcus*

569   *marinus* PCC 9511. J Biol Rhythms 23:187-99.

570   35.   Kanesaki Y, Shiwa Y, Tajima N, Suzuki M, Watanabe S, Sato N, Ikeuchi M, Yoshikawa H. 2012.

571   Identification of substrain-specific mutations by massively parallel whole-genome resequencing of

572   Synechocystis sp. PCC 6803. DNA Res 19:67-79.

573   36.   Dvornyk V, Knudsen B. 2005. Functional divergence of the circadian clock proteins in prokaryotes.

574   Genetica 124:247-54.

575   37.   Axmann IM, Dühring U, Seeliger L, Arnold A, Vanselow JT, Kramer A, Wilde A. 2009. Biochemical

576   evidence for a timing mechanism in *Prochlorococcus*. J Bacteriol 191:5342-7.

577   38.   Ma P, Mori T, Zhao C, Thiel T, Johnson CH. 2016. Evolution of KaiC-Dependent Timekeepers: A

578   Proto-circadian Timing Mechanism Confers Adaptive Fitness in the Purple Bacterium

579   Rhodopseudomonas palustris. PLoS Genet 12:e1005922.

39. Mukaiyama A, Ouyang D, Furuike Y, Akiyama S. 2019. KaiC from a cyanobacterium Gloeocapsa sp.

PCC 7428 retains functional and structural properties required as the core of circadian clock

system. International Journal of Biological Macromolecules 131:67-73.

40. Dörrich AK, Mitschke J, Siadat O, Wilde A. 2014. Deletion of the Synechocystis sp. PCC 6803

kaiAB1C1 gene cluster causes impaired cell growth under light-dark conditions. Microbiology

doi:10.1099/mic.0.081695-0.

41. Kucho K-i, Okamoto K, Tsuchiya Y, Nomura S, Nango M, Kanehisa M, Ishiura M. 2005. Global

analysis of circadian expression in the cyanobacterium *Synechocystis* sp. strain PCC 6803. J

Bacteriol 187:2190-9.

42. Beck C, Hertel S, Rediger A, Lehmann R, Wiegard A, Kolsch A, Heilmann B, Georg J, Hess WR,

Axmann IM. 2014. A daily expression pattern of protein-coding genes and small non-coding RNAs

in Synechocystis sp. PCC 6803. Appl Environ Microbiol doi:10.1128/AEM.01086-14.

43. van Alphen P, Hellingwerf KJ. 2015. Sustained Circadian Rhythms in Continuous Light in

Synechocystis sp. PCC6803 Growing in a Well-Controlled Photobioreactor. PLoS One 10:e0127715.

44. Saha R, Liu D, Hoynes-O'Connor A, Liberton M, Yu J, Bhattacharyya-Pakrasi M, Balassy A, Zhang F,

Moon TS, Maranas CD, Pakrasi HB. 2016. Diurnal Regulation of Cellular Processes in the

Cyanobacterium Synechocystis sp. Strain PCC 6803: Insights from Transcriptomic, Fluxomic, and

Physiological Analyses. MBio 7.

45. Zavřel T, Očenášová P, Červený J. 2017. Phenotypic characterization of Synechocystis sp. PCC 6803

substrains reveals differences in sensitivity to abiotic stress. PLoS One 12:e0189130-e0189130.

46. Trautmann D, Voss B, Wilde A, Al-Babili S, Hess WR. 2012. Microevolution in cyanobacteria: re-

sequencing a motile substrain of Synechocystis sp. PCC 6803. DNA Res 19:435-48.

47. Anderson SL, McIntosh L. 1991. Light-activated heterotrophic growth of the cyanobacterium

Synechocystis sp. strain PCC 6803: a blue-light-requiring process. J Bacteriol 173:2761-7.

48.   Loza-Correa M, Sahr T, Rolando M, Daniels C, Petit P, Skarina T, Gomez Valero L, Dervins-Ravault

605   D, Honore N, Savchenko A, Buchrieser C. 2014. The Legionella pneumophila kai operon is

606   implicated in stress response and confers fitness in competitive environments. Environ Microbiol

607   16:359-81.

608   49.   Snijder J, Schuller JM, Wiegard A, Lossl P, Schmelling N, Axmann IM, Plitzko JM, Forster F, Heck AJ.

609   2017. Structures of the cyanobacterial circadian oscillator frozen in a fully assembled state. Science

610   355:1181-1184.

611   50.   Tseng R, Goularte NF, Chavan A, Luu J, Cohen SE, Chang YG, Heisler J, Li S, Michael AK, Tripathi S,

612   Golden SS, LiWang A, Partch CL. 2017. Structural basis of the day-night transition in a bacterial

613   circadian clock. Science 355:1174-1180.

614   51.   Chang YG, Cohen SE, Phong C, Myers WK, Kim YI, Tseng R, Lin J, Zhang L, Boyd JS, Lee Y, Kang S,

615   Lee D, Li S, Britt RD, Rust MJ, Golden SS, LiWang A. 2015. Circadian rhythms. A protein fold switch

616   joins the circadian oscillator to clock output in cyanobacteria. Science 349:324-8.

617   52.   Hitomi K, Oyama T, Han S, Arvai AS, Getzoff ED. 2005. Tetrameric architecture of the circadian

618   clock protein KaiB. A novel interface for intermolecular interactions and its impact on the circadian

619   rhythm. J Biol Chem 280:19127-35.

620   53.   Espinosa J, Boyd JS, Cantos R, Salinas P, Golden SS, Contreras A. 2015. Cross-talk and regulatory

621   interactions between the essential response regulator RpaB and cyanobacterial circadian clock

622   output. Proc Natl Acad Sci U S A 112:2198-203.

623   54.   Breeden L, Nasmyth K. 1985. Regulation of the yeast HO gene. Cold Spring Harb Symp Quant Biol

624   50:643-50.

625   55.   Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. 1979. Generic assignments, strain

626   histories and properties of pure cultures of cyanobacteria. J Gen Microbiol 111:1-61.

627   56.   Dörrich AK, Wilde A. 2015. Spot Assays for Viability Analysis of Cyanobacteria. Bio-protocol

628   5:e1574.

Figures:

Figure 1: KaiC3 showed no or highly decreased ATP synthase activity compared to KaiC. Prior to

the experiment, KaiC was incubated for 2 weeks at 4 °C or overnight at 30 °C to generate fully

phosphorylated (P-KaiC) and dephosphorylated (NP-KaiC) protein, respectively (Fig. S1). **A:**

Representative autoradiograph of separation of [γ-^{32}P]ATP (product) and [α-^{32}P]ADP (substrate)

via thin layer chromatography after incubation with indicated KaiC proteins for 2h. Controls show

the ATP signal in the presence of an excess of cold ADP. For size control [γ-^{32}P]ATP and [α-^{32}P]ADP

were separated on the same cellulose F plate. **B:** Relative ATP signals after 2 hours incubation

displayed as percentage of all radioactive signals in the corresponding lane (mean with standard

deviation from two experiments, each analyzed in duplicates). All values are normalized to the

relative ATP signal at 0h incubation time in the presence of an excess of cold ADP. Statistical

significance was calculated by Brown-Forsythe and Welch ANOVA tests followed by a Dunnett T3

multiple comparisons test using GraphPad Prism 8. Significance of the mean difference is

indicated as follows: non-significant ($p \geq 0.05$, ns), significant with $p = 0.01$ to 0.05 (*), very

significant with $p = 0.001$ to 0.01 (**), extremely significant with $p = 0.0001$ to 0.001 (***) and

extremely significant with $p < 0.0001$ (****).

Figure 2: ATPase activity of Strep-KaiC3 and variants Strep-KaiC3-DE and Strep-KaiC3-AA. Strep-

KaiC3, Strep-KaiC3-DE and Strep-KaiC3-AA were incubated for 24 hours at 30 °C and ADP

production per day and per Strep-KaiC3 monomer was calculated. As a control we used Strep-

KaiC3-catE1⁻catE2⁻, which lacks both ATPase motifs due to replacement of catalytic glutamates

67,68, 310 and 311 with glutamine residues. For comparison ADP production by KaiC was

monitored for 24 or 48 hours. Strep-KaiC was dephosphorylated prior to the experiment by

incubation at 30°C overnight. Shown are mean values with standard deviation of at least five

replicates. Statistical significance was calculated by Brown-Forsythe and Welch ANOVA tests

followed by a Dunnett T3 multiple comparisons test using GraphPad Prism 8. Significance of the

mean difference is indicated as follows: non-significant (p ≥ 0.05, ns), significant with p = 0.01 to

0.05 (*), very significant with p = 0.001 to 0.01 (**), extremely significant with p = 0.0001 to 0.001

(***) and extremely significant with p < 0.0001 (****). Please note that we averaged all

measurements for Strep-KaiC3 WT at 30 °C and therefore repeatedly display them in Fig. 2, Fig.

4A and Fig. 4B.

Figure 3: KaiC3 interacts with KaiB3 and the proteins of the main oscillator KaiC1, KaiB1. A, B,

C: Yeast two-hybrid reporter strains carrying the respective bait and prey plasmids were selected

by plating on complete supplement medium (CSM) lacking leucine and tryptophan (-Leu -Trp).

Physical interaction between bait and prey fusion proteins is indicated by a color change in the

assays using 5-brom-4-chlor-3-indoxyl-β-D-galactopyranoside. **AD**: GAL4 activation domain; BD,

GAL4 DNA-binding domain. Shown are representative results of two replicates. For clear

presentation, spots were assembled from several assays (original scans are shown in Fig. S2).

Figure 4: ATPase activity of Strep-KaiC3 is regulated by KaiB proteins and is temperature dependent. A: ATPase activity of Strep-KaiC3 was decreased in the presence of KaiB1 and monomeric KaiB3 but was not influenced by oligomeric KaiB3 and by KaiA$_{6803}$. Shown are mean values with standard deviation of at least three replicates. Controls show that KaiA$_{6803}$ and KaiB proteins alone did not display ATPase activity (referring to Strep-KaiC3 monomer activity). Statistical significance was calculated by Brown-Forsythe and Welch ANOVA tests followed by a Dunnett T3 multiple comparisons test using GraphPad Prism 8. Significance of the mean difference is indicated as follows: non-significant ($p \geq 0.05$, ns), significant with $p = 0.01$ to 0.05 (*), very significant with $p = 0.001$ to 0.01 (**), extremely significant with $p = 0.0001$ to 0.001 (***)

and extremely significant with p < 0.0001 (****). **B:** ATPase activity of Strep-KaiC3 is temperature-

dependent. Strep-KaiC3 was incubated for 24 hours at the indicated temperatures and ADP

production per day and monomer Strep-KaiC3 was calculated. Shown are mean values with

standard deviation of at least three replicates. A temperature-compensated activity would result

in flat horizontal line. Please note that we averaged all measurements for Strep-KaiC3 WT at 30

°C and therefore repeatedly display them in Fig. 2, Fig. 4A and Fig. 4B. **C:** Summary of the crosstalk

between the KaiAB1C1 standard oscillator and KaiC3 in *Synechocystis*. We hypothesize that KaiC1

forms a standard oscillator together with KaiA$_{6803}$ and KaiB1. KaiC3 might form an additional

regulatory mechanism together with KaiB3. The two systems intertwine by interactions of the

KaiC proteins with each other and the non-corresponding KaiB homologs. Solid lines indicate

experimentally verified physical interactions (this paper and Wiegard *et al*.(32)), dotted arrows

indicate stimulation of phosphorylation (32) and dotted lines with bars show inhibition of ATPase

activity.

Figure 5: The Δ*kaiC3* strain shows growth defects in complete darkness. A: Proliferation of the

Synechocystis wild type, the *kaiAB1C1* deletion mutant (Δ*kaiAB1C1*) and the *kaiC3* deletion

mutant (Δ*kaiC3*) was tested under mixotrophic conditions in continuous light and under

heterotrophic dark conditions. Strains were grown in liquid culture under constant light and

different dilutions were spotted on agar plates containing 0.2 % glucose. Plates were analyzed

after further incubation for 6 or 28 days of continuous light and darkness, respectively. A

representative result of three independent experiments is shown. **B:** Measurement of single spot

densities of the *Synechocystis* wild-type and *kaiC3* deletion strains. Single spot intensities were

determined with Quantity One (Bio-Rad) and normalized to the intensity of the wild-type spot of

the respective control (continuous light condition). Shown are mean values with standard

deviation of at least three replicates. Statistical significance was calculated by Brown-Forsythe

and Welch ANOVA tests followed by a Dunnett T3 multiple comparisons test using GraphPad

Prism 8. Significance of the mean difference is indicated as follows: non-significant ($p \geq 0.05$, ns),

significant with $p = 0.01$ to 0.05 (*), very significant with $p = 0.001$ to 0.01 (**), extremely

significant with $p = 0.0001$ to 0.001 (***) and extremely significant with $p < 0.0001$ (****). CL,

continuous light; D, darkness.

Supplemental material

Title: *Synechocystis* KaiC3 displays temperature and KaiB dependent ATPase activity and is important for growth in darkness

Running title: Biochemical characterization of *Synechocystis* KaiC3

Authors: Anika Wiegard[1,*,#], Christin Köbler[2], Katsuaki Oyama[3,‡], Anja K. Dörrich[4], Chihiro Azai[3,5], Kazuki Terauchi[3,5,#], Annegret Wilde[2], Ilka M. Axmann[1]

Author affiliations:

[1] Institute for Synthetic Microbiology, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich Heine University Duesseldorf, Duesseldorf, Germany

[2] Institute of Biology III, Faculty of Biology, University of Freiburg, Freiburg, Germany

[3] Graduate School of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

[4] Institute for Microbiology and Molecular Biology, Justus-Liebig University, Giessen, Germany

[5] College of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

* present address: Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm, Sweden

‡ present address: Graduate School of Medicine, Kobe University, Kobe, Hyougo, Japan

Correspondence and requests for materials should be addressed to A. Wiegard (email: anika.wiegard@ki.se) or K. Terauchi (email: terauchi@fc.ritsumei.ac.jp)

 Supplementary methods

 Pull down analysis of KaiB and KaiC proteins

GST-KaiB1 and GST-KaiB3 proteins were expressed in *E. coli* BL21 using the pGEX-6P1 vector and

extracted as described in the main text. Protein concentration of the generated whole cell lysate

was determined using the Bradford method (Bradford MM. 1976. Anal Biochem 72:248-54). For

immobilization of each KaiB protein, a volume corresponding to 40 mg whole protein content was

incubated with 100 µl glutathione sepharose 4B (GE Healthcare) for 20 min at room temperature.

The resin was thoroughly washed four times with extraction buffer [50 mM Tris/HCl (pH8), 150

mM NaCl, 0.5 mM EDTA, 1 mM DTT (+ 5 mM $MgCl_2$, 1 mM ATP for KaiC proteins)] and

subsequently incubated with *Synechocystis* wild type whole cell lysate, which had been generated

from 200 ml cell culture grown to an OD_{750nm} of ~0.8. Briefly, cells were lysed in cold thylakoid

buffer [50 mM Hepes/NaOH (pH7), 5 mM $MgCl_2$, 25 mM $CaCl_2$, 10% Glycerol] using glass beads

(mixture of 0.1–0.11 mm and 0.25-0.5 mm size) in a bead beater (Reetsch) at 4 °C. 1 ml of the

lysate was used for incubation with glutathione sepharose-bound GST-KaiB1 and GST-KaiB3

proteins to co-precipitate Kai interaction partners. Afterwards, the resin material was thoroughly

washed with extraction buffer, mixed with SDS loading dye and incubated at 50 °C for 30 min

before being subjected to SDS-PAGE and Western Blot analysis.

Expression and purification of 3xFLAG-tagged KaiC1 and KaiC3 proteins was performed as

described earlier (Wiegard A, Dörrich AK, Deinzer HT, Beck C, Wilde A, Holtzendorff J, Axmann IM.

2013. Microbiology 159:948-958) from *Synechocystis*. After binding of 3xFLAG-KaiC1/3 to ANTI-

FLAG M2 affinity gel (Sigma Aldrich) over night at 4 °C, the resin was thoroughly washed with cold

FLAG buffer containing 0.03 % β-DM, followed by one washing step without the addition of β-

DM. For pull down of KaiB proteins, 150 µl resin-bound 3x-FLAG-KaiC1 and 3xFLAG-KaiC3 were

incubated with 100 µl of *E. coli* BL21 whole cell lysate (adjusted to a protein content of 10 µg/µl)

containing GST-KaiB1 and GST-KaiB3, respectively. After removing unbound proteins by washing

with FLAG buffer, the samples were subjected to SDS-PAGE and Western Blot analysis.

Antibodies

Antibodies against KaiB1 and KaiB3 were produced as whole-rabbit-IgG fraction by Pineda

Antikörper Services using synthetic peptides coupled to keyhole limpet haemocyanin as antigens.

For the synthetic peptides, epitopes were completed N- and C-terminally to 15 AA. The peptide

sequences were CIDVLK**NPQLAEEDKIL**AT (bold epitope corresponds to AA 50-59 in KaiB1) and

CLDIVPEG**LQVRL**PED (bold epitope corresponds to AA 95-98 in KaiB3), respectively. For detection

of KaiC1 and KaiC3 the specific peptide derived antibodies described in Wiegard *et al.* were used

(Wiegard A, Dörrich AK, Deinzer HT, Beck C, Wilde A, Holtzendorff J, Axmann IM. 2013.

Microbiology 159:948-958). HemA antibody was produced by Pineda Antikörper Services using

the recombinant protein and has been described in Sobotka *et al.* (Sobotka R, Tichy M, Wilde A,

Hunter CN. 2011. Plant Physiol 155:1735-47). The antibody against AtpB was kindly provided by

K.-D. Irrgang (Technical University Berlin, Germany).

Oligonucleotide Name	Sequence (5' – 3')	Purpose[#]
Construction of yeast two-hybrid expression vectors		
KaiC3-AD-fw	ta<u>GGATCC</u>ATGATCGACCAAGAGACAG	Y2H
KaiC3-AD-rev	gc<u>TCTAGA</u>TATTTTCTCATCGAATAAACCG	Y2H
AD-KaiC3-fw	ta<u>GGATCC</u>ATATCGACCAAGAGACAGATG	Y2H
AD-KaiC3-rev	at<u>CTCGAG</u>TATTTTCTCATCGAATAAACCG	Y2H
BD-KaiC3-fw	ta<u>GGATCC</u>AATCGACCAAGAGACAGATG	Y2H
BD-KaiC3-rev	gc<u>ACTAGT</u>TATTTTCTCATCGAATAAACCG	Y2H
AD-KaiA-fw	ta<u>GGATCC</u>GTCAGTCTCCCCTCTCCC	Y2H
AD-KaiA-rev	ta<u>CTCGAG</u>ATCCGTCTGATAATATACATCAAAG	Y2H
BD-KaiA-fw	ta<u>GGATCC</u>GCAGTCTCCCCTCTCCC	Y2H
BD-KaiA-rev	gc<u>ACTAGT</u>ATCCGTCTGATAATATACATCAAAG	Y2H
KaiB3-AD-fw	ta<u>GGATCC</u>ATGGATATGAATAGGATTGTGTTAAG	Y2H
KaiB3-AD-rev	gc<u>TCTAGA</u>ATCCTCCGGCAAACG	Y2H
AD-KaiB3-fw	ta<u>GGATCC</u>CTGATATGAATAGGATTGTGTTAAGAC	Y2H
AD-KaiB3-rev	ta<u>CTCGAG</u>ATCCTCCGGCAAACG	Y2H
BD-KaiC1-fw	aa<u>GGATCC</u>GAACTTACCGATTGTTAACGAAC	Y2H
BD-KaiC1-rev	gc<u>ACTAGT</u>CTCAGCGGTCTTGTCC	Y2H
KaiB1-AD-fw	ta<u>GGATCC</u>ATGAGCCCCTTTAAAAAAAC	Y2H
KaiB1-AD-rev	gc<u>ACTAGT</u>TTGGTCTTCTGCTTCCC	Y2H
AD-KaiB1-fw	ta<u>GGATCC</u>GGAGCCCCTTTAAAAAAACTTAC	Y2H
AD-KaiB1-rev	tt<u>CTCGAG</u>TTGGTCTTCTGCTTCCC	Y2H
KaiB1-BD-rev	tt<u>GTCGAC</u>TTTTGGTCTTCTGCTTCCC	Y2H
BD-KaiB1-fw	ta<u>GGATCC</u>GAGCCCCTTTAAAAAAACTTAC	Y2H
Construction of vectors for heterologous expression in *E. coli*		
kaiB1-fw	CCGT<u>GGATCC</u>AGCCCCTTTAAAAAAACT	E
kaiB1-rev	CCTG<u>GCGGCCGC</u>TTTTCTATTGGTC	E
kaiB3-fw	GTTATCA<u>GGATCC</u>GATATGAATAGGATTGTG	E
kaiB3-rev	CTAGGG<u>GCGGCCGC</u>TTAATCCTCC	E
MU-kaiC3-AA-fw	GGGGAGCTTCCATTACCGATGCCCATATTGCAGCAATTACCGATTCG	MU
MU-kaiC3-AA-rev	CGAATCGGTAATTGCTGCAATATGGGCATCGGTAATGGAAGCTCCCC	MU
MU-kaiC3-DE-fw	GGGGAGCTTCCATTACCGATGCCCATATTGACGAAATTACCGATTCG	MU
MU-kaiC3-DE-rev	CGAATCGGTAATTTCGTCAATATGGGCATCGGTAATGGAAGCTCCCC	MU
MU-kaiC3-E67QE68Q-fw	TGTCACTTTTCAGCAACCCCCCAAGG	MU
MU-kaiC3-E67QE68Q-fw	AAAACACCGTTTTCCCCC	MU
MU-kaiC3-E310QE311Q-fw	TTTTGCCTTTCAGCAAAGTCGAGAACAATTAATTC	MU
MU-kaiC3-E310QE311Q-fw	ACTAAACAGCGCTCTCCATTG	MU
pASK-kaiC3-fw	AT<u>CCGCGG</u>TCATCGACCAAGAGACAGATG	E
pASK-kaiC3-rev	TCGC<u>AAGCTT</u>TATATTTTCTCATCGAATAAACC	E

775 **Table S1.** Oligonucleotides used in this study. Restriction sites are underlined. [#] Y2H, expression
776 in yeast cells; E, expression in *E. coli* cells; MU, mutagenesis

Plasmid Name	Description	Reference
pCGADT7ah	Expression of fusion proteins with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	(1)
pGADT7ah	Expression of fusion proteins with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	(2)
pD153	Expression of fusion proteins with a C-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	(3)
pGBKT7	Expression of fusion proteins with an N-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	Clontech, Germany
pCGAD-*kaiC3*-AD	Expression of KaiC3$_{6803}$ with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pGAD-AD-*kaiC3*	Expression of KaiC3$_{6803}$ with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pD153-*kaiC3*-BD	Expression of KaiC3$_{6803}$ with a C-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	(4)
pGBK-BD-*kaiC3*	Expression of KaiC3$_{6803}$ with an N-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	This study
pCGAD-*kaiA*-AD	Expression of KaiA$_{6803}$ with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	(4)
pGAD-AD-*kaiA*	Expression of KaiA$_{6803}$ with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pD153-*kaiA*-BD	Expression of KaiA$_{6803}$ with a C-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	(4)
pGBK-BD-*kaiA*	Expression of KaiA$_{6803}$ with an N-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	This study
pCGAD-*kaiB3*-AD	Expression of KaiB3$_{6803}$ with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pGAD-AD-*kaiB3*	Expression of KaiB3$_{6803}$ with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pCGAD-*kaiC1*-AD	Expression of KaiC1$_{6803}$ with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	(4)
pGAD-AD-*kaiC1*	Expression of KaiC1$_{6803}$ with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	(4)
pD153-*kaiC1*-BD	Expression of KaiC1$_{6803}$ with a C-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	(4)
pGBK-BD-*kaiC1*	Expression of KaiC1$_{6803}$ with an N-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	This study
pCGAD-*kaiB1*-AD	Expression of KaiB1$_{6803}$ with a C-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pGAD-AD-*kaiB1*	Expression of KaiB1$_{6803}$ with an N-terminal *GAL4*$_{(768–881)}$ AD-tag in yeast cells, *LEU2*, HA epitope tag	This study
pD153-*kaiB1*-BD	Expression of KaiB1$_{6803}$ with a C-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	This study

Plasmid Name	Description	Reference
pGBK-BD-*kaiB1*	Expression of KaiB1 with an N-terminal GAL4$_{(1–147)}$ DNA-BD-tag in yeast cells, *TRP1*, c-Myc epitope tag	This study
pGEX-*kaiA$_{6803}$*	Expression of KaiA$_{6803}$ with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells	(5)
pGEX-*kaiB1*	Expression of KaiB1 with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells	This study
pGEX-*kaiB3*	Expression of KaiB3 with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells	This study
pGEX-*kaiC*	Expression of KaiC with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells	(6)
pGEX-*kaiC3*	Expression of KaiC3 with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells	(5)
pGEX-*kaiC3-AA*	Can be used for expression of KaiC3-AA with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells. Was only used for cloning in this study	This study
pGEX-*kaiC3-DE*	Can be used for expression of KaiC3-DE with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells. Was only used for cloning in this study	This study
pGEX-*kaiC3-catE1$^-$catE2$^-$*	Can be used for expression of KaiC3- catE1$^-$catE2$^-$ with an N-terminal GST-tag$_{(1-231)}$) in *E.coli* cells. Was only used for cloning in this study	This study
pASK-kaiC	Expression of KaiC with an N-terminal Strep-tag$_{(1-11)}$) in *E.coli* cells	(7)
pASK-kaiC3	Expression of KaiC with an N-terminal Strep-tag$_{(1-11)}$) in *E.coli* cells	This study
pASK-kaiC3-*AA*	Expression of KaiC-AA with an N-terminal Strep-tag$_{(1-11)}$) in *E.coli* cells	This study
pASK-kaiC3-*DE*	Expression of KaiC-DE with an N-terminal Strep-tag$_{(1-11)}$) in *E.coli* cells	This study
pASK-kaiC3-*catE1$^-$catE2$^-$*	Expression of KaiC3 -catE1$^-$catE2$^-$ with an N-terminal Strep-tag$_{(1-11)}$) in *E.coli* cells	This study

Table S2. Plasmids used in this study.

Table S2 references

1.   Rausenberger J, Tscheuschler A, Nordmeier W, Wust F, Timmer J, Schafer E, Fleck C,
Hiltbrunner A. 2011. Photoconversion and nuclear trafficking cycles determine
phytochrome A's response profile to far-red light. Cell 146:813-25.

2. Hiltbrunner A, Viczian A, Bury E, Tscheuschler A, Kircher S, Toth R, Honsberger A, Nagy F,
Fankhauser C, Schafer E. 2005. Nuclear accumulation of the phytochrome A
photoreceptor requires FHY1. Curr Biol 15:2125-30.

3. Shimizu-Sato S, Huq E, Tepperman JM, Quail PH. 2002. A light-switchable gene promoter
system. Nat Biotechnol 20:1041-4.

4. Axmann IM, Dühring U, Seeliger L, Arnold A, Vanselow JT, Kramer A, Wilde A. 2009.
Biochemical evidence for a timing mechanism in *Prochlorococcus*. J Bacteriol 191:5342-7.

5. Wiegard A, Dörrich AK, Deinzer HT, Beck C, Wilde A, Holtzendorff J, Axmann IM. 2013.
Biochemical analysis of three putative KaiC clock proteins from *Synechocystis* sp. PCC
6803 suggests their functional divergence. Microbiology 159:948-958.

6. Nishiwaki T, Satomi Y, Nakajima M, Lee C, Kiyohara R, Kageyama H, Kitayama Y,
Temamoto M, Yamaguchi A, Hijikata A, Go M, Iwasaki H, Takao T, Kondo T. 2004. Role of
KaiC phosphorylation in the circadian clock system of *Synechococcus elongatus* PCC
7942. Proc Natl Acad Sci U S A 101:13927-32.

7. Oyama K, Azai C, Nakamura K, Tanaka S, Terauchi K. 2016. Conversion between two
conformational states of KaiC is induced by ATP hydrolysis as a trigger for cyanobacterial
circadian oscillation. Sci Rep 6:32443.

Figure S1: Phosphorylation level of KaiC proteins used for ATP synthase activity assay shown in Fig. 1. Proteins were separated via SDS-PAGE using a polyacrylamide gel with 11 %T, 0.67 %C (see dx.doi.org/10.17504/protocols.io.gysbxwe for method description). Fully phosphorylated (P-KaiC) and dephosphorylated (NP-KaiC) *S. elongatus* KaiC proteins were generated by incubating the protein for 2 weeks at 4 °C or overnight at 30 °C, respectively.

Figure S2. Original scans of the KaiC3 interaction with KaiB3 and the proteins of the main oscillator KaiC1, KaiB1. Yeast two-hybrid reporter strains carrying the respective bait and prey plasmids, were selected by plating on complete supplement medium (CSM) lacking leucine and tryptophan (-Leu -Trp). As a positive control, KaiA$_{6803}$ dimer interaction was used. AD, GAL4 activation domain; BD, GAL4 DNA-binding domain; n.r., interactions not relevant for this study. **A, C-F:** Physical interaction between bait and prey fusion proteins is indicated by a color change in the assays using 5-brom-4-chlor-3-indoxyl-β-D-galactopyranoside. **B:** Physical interaction between bait and prey fusion proteins is determined by growth on complete medium lacking leucine, tryptophan and histidine (-Leu -Trp -His) and addition of 6.25 mM 3-amino-1,2,4-triazole (3-AT). A detailed protocol can be found on protocols.io (https://dx.doi.org/10.17504/protocols.io.wcnfave).

Figure S3. Interaction of *Synechocystis* KaiB and KaiC proteins in pull down analysis. A,B GST-tagged KaiB1 and KaiB3 proteins were expressed in *E. coli* BL21 cells, bound to glutathione sepharose and incubated with *Synechocystis* WT whole cell lysate (WCL). In the eluate, KaiC1 (A) and KaiC3 (B) were detected by Western Blot analysis using specific antibodies. As negative control, blots were incubated with an antiserum against the ferrochelatase HemH, because we did not expect an interaction between the KaiB proteins and HemH. **C,D** FLAG- KaiC1 (C) and FLAG-KaiC3 (D) were expressed in *Synechocystis*, bound to Anti-FLAG-agarose and incubated with whole cell lysate (WCL) from *E. coli* BL21 cells expressing GST-KaiB1 and GST-KaiB3 proteins, respectively. To detect whether GST-KaiB proteins were co-eluted we used an antibody raised against GST. Incubation with an AtpB antibody served as negative control. Both experiments were performed once.

Figure S4: Purification of KaiB and KaiC3 proteins used in this study. AB: KaiB proteins were subsequently purified via affinity chromatography, anion exchange chromatography and size exclusion chromatography. Shown are the chromatograms after size exclusion chromatography using a superdex 200 Increase 10/30 GL column (A) or Sephacryl S200 HR HiPrep 16/60 column (B). On both columns KaiB3 was separated into a monomer (red asterisk) and tetramer (blue asterisk), whereas KaiB1 was mainly eluted as tetramer (blue asterisk). **CD:** KaiC3 proteins were purified via affinity chromatography followed by size exclusion chromatography on a Sephacryl S300 HR HiPrep 16/60 Sephacryl column. All KaiC3 proteins eluted as oligomer (green asterisk). Arrows indicate the size of standard proteins in kDa.

Figure S5. ATPase activity and dephosphorylation of KaiC3 are temperature dependent. A. Temperature-dependent ATPase activity of Strep-KaiC3 (also displayed in Fig. 4B) shown in an Arrhenius plot. Strep-KaiC3 was incubated for 24 hours at 25°C, 30°C and 35°C and ADP production per day and monomer Strep-KaiC3 was calculated. To determine the activation energy, the natural logarithm of the mean ADP production rate was plotted against the reciprocal temperature. The activation energy of KaiC3 ATPase was calculated from the slope of a linear regression equation as E_a=66.5 x kJ mol^{-1}. **B,C.** Relative dephosphorylation of Strep-KaiC3. Strep-KaiC3 was incubated for 24 hours in 20 mM Tris-HCl/pH8, 150 mM NaCl, 5 mM MgCl$_2$, 1 mM ATP at the indicated temperatures. Proteins were separated via SDS-PAGE using a polyacrylamide gel with 11 %T, 0.67 %C. The phosphorylation level at each temperature was determined as the ratio of P-Strep-KaiC3 to total Strep-KaiC3 (P-Strep-KaiC3 + NP-Strep-KaiC3) using ImageJ. A representative gel image is shown in **B**. In **C**, the relative phosphorylation levels compared to the stock protein (0h) of 5 experiments are plotted as mean values with standard deviation. Incubation at 25°C resulted in lower phosphorylation levels, demonstrating higher net-dephosphorylation.