Title: *Synechocystis* **KaiC3 displays temperature and KaiB dependent ATPase activity and is**

important for viability in darkness

Running title: Biochemical characterization of *Synechocystis* **KaiC3**

Authors: Anika Wiegard[1,*,#], Christin Köbler[2], Katsuaki Oyama[3,‡], Anja K. Dörrich[4], Chihiro Azai[3,5],

Kazuki Terauchi[3,5,#], Annegret Wilde[2], Ilka M. Axmann[1]

Author affiliations:

[1] Institute for Synthetic Microbiology, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich

Heine University Duesseldorf, Duesseldorf, Germany

[2] Institute of Biology III, Faculty of Biology, University of Freiburg, Freiburg, Germany

[3] Graduate School of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

[4] Institute for Microbiology and Molecular Biology, Justus-Liebig University, Giessen, Germany

[5] College of Life Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

* present address: Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm,

Sweden

‡ present address: Graduate School of Medicine, Kobe University, Kobe, Hyougo, Japan

Correspondence and requests for materials should be addressed to A. Wiegard (email:

anika.wiegard@ki.se) or K. Terauchi (email: terauchi@fc.ritsumei.ac.jp)

Abstract:

Cyanobacteria form a heterogeneous bacterial group with diverse lifestyles, acclimation strategies and differences in the presence of circadian clock proteins. In *Synechococcus elongatus* PCC 7942, a unique posttranslational KaiABC oscillator drives circadian rhythms. ATPase activity of KaiC correlates with the period of the clock and mediates temperature compensation. *Synechocystis* sp. PCC 6803 expresses additional Kai proteins, of which KaiB3 and KaiC3 proteins were suggested to fine-tune the standard KaiAB1C1 oscillator. In the present study, we therefore characterised the enzymatic activity of KaiC3 as a representative of non-standard KaiC homologs *in vitro*. KaiC3 displayed ATP synthase and ATPase activities, which were lower compared to the *Synechococcus* 7942 KaiC protein. ATP hydrolysis showed no temperature compensation. Hence, KaiC3 is missing a defining feature of the model cyanobacterial circadian oscillator. Yeast two-hybrid analysis showed that KaiC3 interacts with KaiB3, KaiC1 and KaiB1. Further, KaiB3 and KaiB1 reduced *in vitro* ATP hydrolysis by KaiC3. Spot assays showed that chemoheterotrophic growth in constant darkness is completely abolished after deletion of Δ*kaiAB1C1* and reduced in the absence of *kaiC3*. We therefore suggest a role for adaptation to darkness for KaiC3 as well as a crosstalk between the KaiC1 and KaiC3 based systems.

Importance: The circadian clock is known to influence the cyanobacterial metabolism globally and a deeper understanding of its regulation and fine-tuning will be important for metabolic optimizations in context of industrial applications. Due to the heterogeneity of cyanobacteria, the characterization of clock systems in organisms apart from the standard circadian model *Synechococcus elongatus* sp. PCC 7942 is required. *Synechocystis* sp. PCC 6803 is of special interest, because it represents a major cyanobacterial model organism and harbors diverged

43 homologs of the clock proteins, which are present in various other non-cyanobacterial

44 prokaryotes. By our *in vitro* studies we unravel the interplay of the multiple *Synechocystis* Kai

45 proteins and characterize the enzymatic activities of the non-standard clock homolog KaiC3. We

46 show that the deletion of *kaiC3* affects growth in constant darkness suggesting its involvement in

47 the regulation of non-photosynthetic metabolic pathways.

Introduction:

49 Cyanobacteria have evolved the circadian clock system to sense, anticipate and respond to

50 predictable environmental changes based on the rotation of Earth and the resulting day-night

51 cycle. Circadian rhythms are defined by three criteria: (i) oscillations with a period of about 24

52 hours without external stimuli, (ii) synchronization of the oscillator with the environment and (iii)

53 compensation of the usual temperature dependence of biochemical reactions, so that the period

54 of the endogenous oscillation does not depend on temperature in a physiological range (1). The

55 cyanobacterial circadian clock system has been studied in much detail in the unicellular model

56 cyanobacterium *Synechococcus elongatus* PCC 7942 (hereafter *Synechococcus* 7942). Its core

57 oscillator is composed of three proteins, which are unique to prokaryotes: KaiA, KaiB, and KaiC

58 (from now on $KaiA_{7942}$, $KaiB_{7942}$ and $KaiC_{7942}$; please note that some of the hereafter cited

59 information were gained studying proteins from *Thermosynechococcus elongatus* BP-1 though)

60 (2). The level of $KaiC_{7942}$ phosphorylation and $KaiC_{7942}$'s ATPase activity represent the key features

61 of the biochemical oscillator. $KaiA_{7942}$ stimulates autophosphorylation and ATPase activity of

62 $KaiC_{7942}$, whereas $KaiB_{7942}$ binding to the complex inhibits $KaiA_{7942}$ action, stimulates

63 autodephosphorylation activity and reduces ATPase activity of $KaiC_{7942}$ (3-5). As a consequence of

dynamic interactions with KaiA$_{7942}$ and KaiB$_{7942}$, KaiC$_{7942}$ rhythmically phosphorylates and

dephosphorylates with a 24-hour period (2).

KaiC$_{7942}$ consists of two replicated domains (CI and CII) which assemble into a hexamer forming

an N-terminal CI ring and a C-terminal CII ring (6-8). Phosphorylation takes place in the CII ring (9),

whereas ATP hydrolysis occurs in both rings (10). In the CII ring, ATP hydrolysis is part of the

dephosphorylation mechanism (11, 12). ATP hydrolysis in the CI ring correlates with the period of

the clock and temperature compensation and is further required for a conformational change of

KaiC$_{7942}$, which allows binding of KaiB$_{7942}$ (4, 13, 14). The levels of phosphorylation and ATP

hydrolysis of the KaiC$_{7942}$ serve as the read-out for regulatory proteins, which orchestrate the

circadian output (15, 16). For a recent review on the functioning of the KaiABC$_{7942}$ system, see

Swan *et al.* (17).

Cyanobacteria represent one of the most diverse prokaryotic phyla(18) and little is known about

timekeeping mechanisms in other cyanobacteria than *Synechococcus* 7942. A core diurnal

genome has been described (19), but temporal coordination varies and, based on genomic

analyses, large variations in the cyanobacterial clock systems can be expected (20-25).

The cyanobacterial model strain *Synechocystis* sp. PCC 6803 (from now on *Synechocystis* 6803)

contains a standard *kai* operon, encoding homologs of the *kaiA*, *kaiB* and *kaiC* genes (in the

following designated *kaiA$_{6803}$, kaiB1$_{6803}$* and *kaiC1$_{6803}$*) as well as two additional *kaiB* and *kaiC*

genes (26). The *kaiC2$_{6803}$* and *kaiB2$_{6803}$* genes form an operon, whereas *kaiC3$_{6803}$* and *kaiB3$_{6803}$* are

orphan genes. Bioinformatic studies revealed that all cyanobacteria, which do contain at least one

kaiB gene, always encode orthologs of KaiB1 and KaiC1. Additional KaiB3 and KaiC3 homologs are

present in about one third of cyanobacterial genera, with a slightly higher occurrence of KaiC3.

KaiC3 is also present in bacteria and archaea, showing a higher distribution than KaiC1 and KaiC2

homologs (19). The reported number of oscillating genes in *Synechocystis* 6803 varies largely

between different studies, which is likely due to differences in growth conditions (27-30). These

results imply that *Synechocystis* 6803 might be able to fine-tune its circadian rhythm in response

to environmental conditions. In line with this, Aoki and Onai (21) suggested that $KaiC3_{6803}$ and

$KaiB3_{6803}$ modulate the amplitude and period of the $KaiAB1C1_{6803}$ oscillator, whereas, to our

knowledge, no function for $kaiC2B2_{6803}$ has been described so far.

The three KaiC proteins from *Synechocystis* 6803 show high conservation of the kinase motif in

the CII domain and all three proteins were shown to exhibit autophosphorylation activity (19, 23).

$KaiA_{6803}$ stimulates the autophosphorylation of $KaiC1_{6803}$, but does not affect the phosphorylation

of the two other KaiC homologs. Based on sequence analysis and experimental validation of

kinase activity, Schmelling *et al.* (19) concluded that autokinase and autophosphatase activities

are highly conserved features of all cyanobacterial KaiC homologs. Likewise, ATPase motifs in the

N-terminal CI domain of KaiC are highly conserved in all cyanobacterial KaiC homologs (19).

However, ATP hydrolysis was only characterised for several KaiC1 homologs and one non-

cyanobacterial KaiC2 protein (10, 31-33).

In a natural day and night cycle, *Synechococcus* 7942 orchestrates its metabolism in a precise

temporal schedule, with the metabolism being adjusted by the clock but also feeding input

information to the clock (34-36). Knockout of the *Synechococcus* 7942 *kai* genes leads to a growth

disadvantage. When grown in competition with the wild type under light-dark conditions, the

clock deficient mutant cells are eliminated from the culture (37). In *Synechocystis* 6803 deletion

of the *kaiAB1C1* gene cluster causes reduced growth in light-dark rhythms, even when grown as

single culture (38). A comparably strong impact on growth in a light-dark cycle was observed in

both strains after deletion of the regulator of phycobilisome association A (RpaA) (34, 39), the

master transcription factor of circadian gene expression. In addition, studies on single nucleotide

polymorphisms identified the *rpaA* gene as one of three genes responsible for the faster growth

of *Synechococcus elongatus* UTEX 2973 compared to *Synechococcus* 7942(40). Using a transposon

library, David *et al.* (41) showed that KaiA, despite being non-essential for growth in light-dark

cycles, strongly contributes to the fitness of *Synechococcus* 7942 under these conditions.

Decreased fitness most likely occurs due to reduced phosphorylation of RpaA in the *kaiA* knockout

strain (41). Overall, these data demonstrate the value of the cyanobacterial clock system for

metabolic orchestration under natural conditions, with the clock system considered especially

important for the transition from light to darkness (42).

Here, we aim at a detailed biochemical characterization of $KaiC3_{6803}$ and the role of *Synechocystis*

6803 $KaiB_{6803}$ homologs in modulation of $KaiC3_{6803}$ function. Further, we demonstrate that the

$\Delta kaiC3$ mutant strain has a growth defect under chemoheterotrophic growth conditions, which is

similar, but less pronounced compared to $\Delta kaiAB1C1$ and $\Delta rpaA$ strains (38, 39). Our data support

the idea of a function of $KaiC3_{6803}$ and $KaiB3_{6803}$ in fine-tuning the central oscillator composed of

$KaiA_{6803}$, $KaiB1_{6803}$ and $KaiC1_{6803}$ in *Synechocystis* 6803.

Results:

Recombinant $KaiC3_{6803}$ displays ATP synthase activity

Previous bioinformatic analysis predicted that kinase, dephosphorylation and ATPase activities

are conserved in $KaiC3_{6803}$ (19). So far, only the kinase activity of $KaiC3_{6803}$ was experimentally

confirmed which differed from the activity of $KaiC_{7942}$ by lacking stimulation by $KaiA_{7942}$ and

KaiA$_{6803}$ (23). Therefore, we aimed at the characterization of further predicted activities of

KaiC3$_{6803}$.

The first step in KaiC$_{7942}$ dephosphorylation is regarded as a reversal of phosphorylation: KaiC$_{7942}$

transfers its bound phosphoryl group to ADP and thereby synthesizes ATP (11, 12). To investigate

reversibility of intrinsic phosphorylation, we tested whether KaiC3$_{6803}$ can synthesize [α-^{32}P]ATP

from [α-^{32}P]ADP (Fig. 1). As control, we used phosphorylated and dephosphorylated KaiC$_{7942}$ (Fig.

S1.). Immediately after adding [α-^{32}P]ADP, all KaiC proteins started to synthesize [α-^{32}P]ATP.

Phosphorylated as well as non-phosphorylated KaiC$_{7942}$ both showed higher initial ATP synthesis

than KaiC3$_{6803}$ (Fig. 1). After incubating phosphorylated and dephosphorylated KaiC$_{7942}$ for 2 hours

at 30°C with [α-^{32}P]ADP, a relative [α-^{32}P]ATP level of 26-28 % was detected. KaiC3$_{6803}$ was also

capable of intrinsic ATP synthesis, but compared to KaiC$_{7942}$ less than one third of relative [α-

^{32}P]ATP was formed. This can be explained by two hypotheses: (i) KaiC3$_{6803}$ showed lower

phosphotransferase activity or (ii) higher consumption of the produced ATP. The latter would

require that KaiC3$_{6803}$ exhibits higher ATPase activity than KaiC$_{7942}$. Therefore, we next examined

ATP hydrolysis activity of KaiC3$_{6803}$.

KaiC3$_{6803}$ displays ATPase activity

ATP hydrolysis can be measured by quantifying ADP production in the presence of KaiC over time

(4). Conservation of WalkerA and WalkerB motifs in the CI domain of KaiC3$_{6803}$ proteins suggested

their capability to hydrolyze ATP (19). We confirmed the predicted ATPase activity *in vitro* using

Strep-KaiC3$_{6803}$, which produced 8.5 ± 1.0 ADP molecules per monomer and day (mean ± s.d., Fig.

2). This activity was about 45 % lower than the one measured for KaiC$_{7942}$ (19.1 ± 3.3 ADP per

day (4)). Hence, the above-observed lower level of net ATP production by KaiC3$_{6803}$ was not based

on a higher ATP consumption, but due to lower dephosphorylation activity *per se*. In KaiC$_{7942}$, the

ATPase activity changes after substitution of the phosphorylation sites (4). We therefore

generated variants of Strep-KaiC3$_{6803}$, in which residues S423 and T424 are either replaced with

aspartate and glutamate (Strep-KaiC3$_{6803}$-DE) or replaced with two alanine residues (Strep-

KaiC3$_{6803}$-AA). Strep-KaiC3$_{6803}$-AA, showed more than two fold increased ATPase activity, whereas

ATP hydrolysis by Strep-KaiC3$_{6803}$-DE was only slightly different from the wild-type protein (Fig. 2)

and not reduced as reported for KaiC$_{7942}$ (4).

KaiC3$_{6803}$ interacts with KaiB3$_{6803}$ and components of the standard KaiAB1C1$_{6803}$ oscillator

In vitro and *in silico* studies suggested that KaiA$_{6803}$, KaiB1$_{6803}$ and KaiC1$_{6803}$ form the standard

clock system of *Synechocystis* 6803 (21, 23). Since Aoki and Onai (21) suggested that KaiC3$_{6803}$

might modulate the main oscillator function, we performed protein-protein interaction studies in

order to reveal a possible crosstalk between the multiple Kai proteins. First, interaction was

determined by yeast-two hybrid analysis using KaiC1$_{6803}$, KaiC3$_{6803}$, KaiB1$_{6803}$ and KaiB3$_{6803}$ fused

to AD and BD domains, respectively. The color change of the colonies indicates β-galactosidase

activity and is an estimate for the interaction of the respective two proteins. As expected, these

experiments showed self-interaction of KaiC3$_{6803}$ (Fig. 3A), since KaiC homologs are known to form

hexamers (6, 8). In addition, an interaction of KaiC3$_{6803}$ with KaiB3$_{6803}$ was detected (Fig. 3A) which

is in line with bioinformatic analysis showing frequent co-occurrence of *kaiC3* and *kaiB3* genes in

genomes (19). We could not detect an interaction between KaiC3$_{6803}$ and KaiA$_{6803}$ by yeast-two

hybrid analysis (Fig. 3B), but detected a heteromeric interaction between KaiC1$_{6803}$ and KaiC3$_{6803}$

using different protein fusion variants (Fig. 3C). These results confirm previous *in vivo* data, which

showed co-purification of KaiA$_{6803}$ with KaiC1$_{6803}$ but not with KaiC3$_{6803}$, and a weak interaction

between the two KaiC homologs $KaiC1_{6803}$ and $KaiC3_{6803}$ (23). Besides $KaiC1_{6803}$, also its respective

KaiB protein, $KaiB1_{6803}$, showed an interaction with $KaiC3_{6803}$ in our analysis (Fig. 3C),

corroborating the hypothesis that there is a cross talk between the putative $KaiC3_{6803}$-$B3_{6803}$

system and the core oscillator $KaiAB1C1_{6803}$. Such a possible cross talk via the KaiB proteins was

further supported by our *in vitro* pull-down assays, in which $KaiC3_{6803}$ interacted with $KaiB3_{6803}$

(Fig. S3B,D) and further with $KaiB1_{6803}$ (Fig. S3B,D). Also $KaiC1_{6803}$ interacted with both, $KaiB1_{6803}$

and $KaiB3_{6803}$ homologs (Fig. S3,A,C). One must take into account that based on the *in vitro* pull-

down assays using *Synechocystis* 6803 whole cell extracts we cannot exclude indirect interactions.

As we have shown that $KaiC1_{6803}$ and $KaiC3_{6803}$ interact with each other, both proteins could bind

as a hetero-hexamer to the GST-tagged $KaiB_{6803}$ proteins and therefore co-elute from the affinity

matrix.

ATPase activity of $KaiC3_{6803}$ is reduced in the presence of $KaiB1_{6803}$ and $KaiB3_{6803}$

The interaction of $KaiC3_{6803}$ with $KaiB1_{6803}$ and $KaiB3_{6803}$ (Fig. 3 and Fig. S3) suggested a regulation

of $KaiC3_{6803}$ activity by these two KaiB proteins. We therefore measured ATP hydrolysis by Strep-

$KaiC3_{6803}$ in the presence of $KaiB1_{6803}$ and $KaiB3_{6803}$ proteins, respectively (Fig. 4A). After size

exclusion chromatography $KaiB1_{6803}$ was mainly eluted as tetramer (44 and 72 kDa, respectively,

depending on the column), whereas $KaiB3_{6803}$ was eluted as monomer (13/23 kDa) and tetramer

(41/70 kDa) (Fig. S4). Therefore, the monomeric and tetrameric $KaiB3_{6803}$ fractions were tested

separately. In all measurements, the ATPase activity was linear and showed no oscillations. ATP

hydrolysis was reduced by 55 % after the addition of the $KaiB3_{6803}$ monomer, but not affected by

the $KaiB3_{6803}$ tetramer. The $KaiB1_{6803}$ tetramer also inhibited the ATPase activity of Strep-

$KaiC3_{6803}$, though to a lesser extend (35 % reduction compared to Strep-$KaiC3_{6803}$ alone, Fig. 4).

Because ATPase activity of $KaiC_{7942}$ is influenced by $KaiA_{7942}$ (4), we also investigated the effect of

$KaiA_{6803}$ on $KaiC3_{6803}$, but in line with the lack of interaction in our yeast two-hybrid analysis data

(Fig. 3B), ATPase activity of Strep-$KaiC3_{6803}$ was not clearly affected by $KaiA_{6803}$ (Fig. 4A).

ATPase activity of $KaiC3_{6803}$ is temperature dependent

True circadian clocks are characterized by temperature compensated oscillations, which ensure

robust time measurements under temperature fluctuations. In the *Synechococcus* 7942 KaiABC

clock, overall temperature compensation is derived from $KaiC_{7942}$'s ATPase activity, which is stable

between 25 and 35 °C (Q_{10}=1.2, (4)). We therefore asked whether N-Strep-$KaiC3_{6803}$, as a

representative of non-standard KaiC homologs, shows temperature compensation as well.

Measurements at 25, 30 and 35 °C, however, revealed a temperature-dependent ADP production

by $KaiC3_{6803}$ (Q_{10}=2.4, Fig. 4B; activation energy = 66.5 kJ mol^{-1}, Fig. S5A). Hence, $KaiC3_{6803}$ is

lacking a characteristic feature of circadian oscillators. In accordance, dephosphorylation of Strep-

$KaiC3_{6803}$ was higher at 25 °C than at 30 and 35 °C (Fig. S5B)

The non-standard $KaiC3_{6803}$ protein supports growth of *Synechocystis* 6803 cells in the dark

As *in vitro* analyses suggested an interaction of $KaiC3_{6803}$ with $KaiB1_{6803}$ and $KaiC1_{6803}$, we aimed

at elucidating the role of $KaiC3_{6803}$ and its crosstalk with the putative main oscillator in the cell.

Clock factors are reported to be essential for cell viability in light-dark cycles (34, 39, 41).

Accordingly, deletion of the *kaiAB1C1* cluster of *Synechocystis* 6803 resulted in growth defects in

light-dark cycles (38). Lower viability of the Δ*kaiAB1C1* strain was more pronounced under

photomixotrophic (0.2 % glucose) compared to photoautotrophic conditions, whereas deletion

of *kaiC3* had no effect on cell viability in light-dark cycles (38). As KaiC3 homologs are also present

in non-photosynthetic bacteria (19), we were interested in the growth of the Δ*kaiC3* strain in the

dark. The *Synechocystis* 6803 WT strain used here, in contrast to previous studies (43), is able to

grow in complete darkness when supplemented with glucose (39). We therefore analyzed the

viability of Δ*kaiC3* cells via spot assays (44) in constant light and in complete darkness on agar

plates containing 0.2 % glucose (Fig. 5). Under photomixotrophic growth conditions with

continuous illumination, wild type and Δ*kaiC3* showed similar viability, whereas the viability of

the Δ*kaiAB1C1* strain was reduced. Further, in complete darkness, wild type and Δ*kaiC3* displayed

detectable growth, while growth of Δ*kaiAB1C1* was abolished completely. Growth inhibition of

the Δ*kaiAB1C1* mutant under mixotrophic conditions in the light and in the dark highlights the

importance of the KaiAB1C1$_{6803}$ oscillator for the switch between these two different metabolic

modes of *Synechocystis* 6803. Comparing wild-type and Δ*kaiC3* strains, the latter showed reduced

growth in complete darkness. Thus, KaiC3$_{6803}$ seems to be linked to dark adaption of *Synechocystis*

6803 cells, yet not as essential as the core oscillator KaiAB1C1$_{6803}$.

Discussion:

Characterization of enzymatic activities of KaiC3$_{6803}$

Previous sequence analysis suggested that the three enzymatic activities of KaiC$_{7942}$, which are

autophosphorylation, autodephosphorylation and ATPase activity, are conserved in all

cyanobacterial KaiC proteins (19). This hypothesis is supported by experiments demonstrating

autophosphorylation of all KaiC$_{6803}$ proteins (23), different cyanobacterial KaiC1 and KaiC3

homologs (19, 31, 33), and even non-cyanobacterial KaiC homologs from *Rhodopseudomonas*

palustris (32), *Legionella pneumophila* (45) as well as two thermophilic Archaea (19). In the

current paper, we provide experimental evidence that dephosphorylation and ATPase activities

are conserved in KaiC3$_{6803}$ as well. However, the level of the ATPase activity seems to vary

between KaiC proteins. While ATPase activity of the KaiC from *Prochlorococcus marinus* MED4

was reported to have the same activity as KaiC$_{7942}$ (31), we show here that the ATPase activity of

KaiC3$_{6803}$ is lower. In contrast, ATP hydrolysis activity of the KaiC2 homolog from *Legionella*

pneumophila and KaiC1 homolog from *Gloeocapsa* sp. PCC 7428 were reported to be elevated

when compared to the standard KaiC$_{7942}$ protein (32, 33).

ATPase activity of KaiC3$_{6803}$ further differed from that of KaiC$_{7942}$, by lacking temperature

compensation (Fig. 4). Therefore, KaiC3$_{6803}$ cannot be the core component of a true circadian

oscillator. This is further supported by differences in the ATP hydrolysis rate between the

phosphomimetic variants of KaiC$_{7942}$ and KaiC3$_{6803}$. ATPase activity of the true clock protein

KaiC$_{7942}$ depends on its phosphorylation status. In the KaiC$_{7942}$-AA variant, which mimics the

dephosphorylated state of KaiC$_{7942}$, the ATPase activity is higher than average, whereas, in

KaiC$_{7942}$-DE, mimicking the phosphorylation state, it is diminished (4). In our analysis, ATP

hydrolysis by Strep-KaiC3$_{6803}$-AA was similarly increased in comparison to Strep-KaiC3$_{6803}$.

However, Strep-KaiC3$_{6803}$-DE did not show reduced activity as was shown for KaiC$_{7942}$-DE, implying

that the phosphorylation state of KaiC3$_{6803}$ does not influence KaiC3$_{6803}$'s ATPase activity.

However, the ATPase activity of Strep-KaiC3$_{6803}$ was affected decisively by the addition of

KaiB3$_{6803}$ and KaiB1$_{6803}$. Both KaiB homologs led to decreased ATPase activity of Strep-KaiC3$_{6803}$,

with KaiB3$_{6803}$ having a stronger impact (Fig. 4A). KaiB$_{7942}$ exists as a monomer or a tetramer in

solution (46). As a tetramer, KaiB$_{7942}$ subunits adopt a unique fold, which was also observed in

crystals of KaiB1$_{6803}$ (47). Notably, KaiB$_{7942}$ binds as a monomer to KaiC$_{7942}$, in which it adopts a

different, thioredoxin-like fold (48-50). The formation of KaiB$_{7942}$ tetramers is concentration-

dependent and thereby ensures that the effective monomer concentration is stable over a broad

range of total $KaiB_{7942}$ concentrations (46). This points at a putative mechanism how $KaiC3_{6803}$

might fine-tune the standard oscillator: Binding of $KaiB1_{6803}$ to $KaiC3_{6803}$ will reduce the free

$KaiB1_{6803}$ concentration in the cell and thereby influence the $KaiAB1C1_{6803}$ oscillator. In contrast,

only the monomeric form of $KaiB3_{6803}$ had a regulatory effect on ATP hydrolysis of $KaiC3_{6803}$

suggesting that $KaiB3_{6803}$ bound as a monomer to $KaiC3_{6803}$, but no transition from tetramer to

monomer occurred. This finding is surprising, because residues K57, G88 and D90, which are

important for fold switching of $KaiB_{7942}$ (50), are conserved in $KaiB3_{6803}$. However, we show only

in vitro data here. In the cell, protein concentrations as well as spatial and temporal separation of

the different proteins might have an influence on these interactions. In addition, we do not take

possible heteromeric interactions among the KaiB proteins into account.

Function of $KaiC3_{6803}$ *in an extended network*

Our interaction studies imply a crosstalk between $KaiC1_{6803}$ and $KaiC3_{6803}$ (Fig. 6). KaiC1 is believed

to form a standard oscillator together with $KaiA_{6803}$ and $KaiB1_{6803}$ (21, 23), which is supported by

the interaction with $KaiB1_{6803}$ observed in this paper (Fig. 3C, Fig. S3). We further hypothesize that

$KaiC3_{6803}$ acts in a separate non-circadian regulatory system together with $KaiB3_{6803}$. In

accordance with bioinformatic analysis, which showed a significant co-occurrence of KaiB3 and

KaiC3 in Cyanobacteria (19), $KaiB3_{6803}$ had a stronger effect on the ATPase activity of $KaiC3_{6803}$

than $KaiB1_{6803}$ (Fig. 4A). Since $KaiC3_{6803}$ is able to form hetero-oligomers with $KaiC1_{6803}$ and was

shown to interact with $KaiB1_{6803}$ as well, interference between the $KaiC3_{6803}$-$KaiB3_{6803}$ system and

the standard oscillator is very likely. However, we exclude that $KaiA_{6803}$ is involved in the putative

crosstalk by four reasons: (i) $KaiA_{6803}$ did neither stimulate ATP hydrolysis nor kinase activity of

$KaiC3_{6803}$ (this paper and Wiegard *et al.* (23)); (ii) We were not able to show an interaction of

KaiC3$_{6803}$ and KaiA$_{6803}$ using different approaches (this paper and Wiegard *et al.* (23)); (iii) KaiA

interacting residues are not conserved in cyanobacterial KaiC3 homologs, and (iv) KaiC3 homologs

are present in organisms which do not harbor KaiA (19).

The growth defect of the mutant strains, Δ*kaiAB1C1* and Δ*kaiC3,* in complete darkness suggests

that the here proposed putative KaiC3-KaiB3 system and the KaiAB1C1$_{6803}$ oscillator may target

similar cellular functions. The metabolic network of cyanobacteria is described as temporally

partitioned with extensive effects of day-night transitions, involving shifts in ATP and reductant

levels and alterations of the carbon flux (41). In *Synechococcus* 7942, environmental signals can

be fed into the main clock output system via the transcriptional regulator RpaB (51). In contrast

to *Synechococcus* 7942, *Synechocystis* 6803 is able to grow in complete darkness, which adds

another layer of complexity to day-night transitions and demand for further regulatory elements.

The here observed impaired mixotrophic growth of the *kaiC3$_{6803}$* mutant in darkness supports the

idea of KaiC3$_{6803}$ and KaiB3$_{6803}$ functioning as such additional elements to adjust the state of the

main *Synechocystis* 6803 oscillator. Conversely, it is also possible that KaiC3$_{6803}$ function is

controlled by the KaiAB1C1$_{6803}$ clock system. Köbler *et al.* (39) demonstrated that solely KaiC1$_{6803}$,

but not KaiC3$_{6803}$, interacts with the main output histidine kinase SasA in the *Synechocystis* 6803

timing system. Thus, in *Synechocystis* 6803, only the main oscillator feeds timing information into

the SasA-RpaA output system to control the expression of many genes involved in dark growth

(39). The output pathway for KaiC3$_{6803}$ is unknown so far and it might be possible that the only

function of KaiC3$_{6803}$ is to modulate the function of the main oscillator in response to a yet

unknown input factor.

Material and Methods:

Cloning, expression and purification of recombinant Kai proteins

In this paper, we name the *Synechocystis kai* genes according to the definition introduced by Schmelling *et al.* (19) (Please note that naming of KaiB3/KaiB2 and KaiC3/KaiC2 is not consistent in the literature). Genes encoding KaiB1$_{6803}$ (ORF *slr0757*) and KaiB3$_{6803}$ (ORF *sll0486*) were amplified from genomic *Synechocystis* 6803 wild type DNA using specific primers (Table S1) and Phusion Polymerase (New England Biolabs). After restriction digest with BamHI and NotI, amplified fragments were inserted into pGEX-6P1 (GE Healthcare) and the resulting plasmids were used for heterologous expression. For production of recombinant KaiC$_{7942}$ as well as KaiA$_{6803}$, KaiB1$_{6803}$, KaiB3$_{6803}$ and KaiC3$_{6803}$, we used pGEX-based plasmids described in Wiegard *et al.* (23) (see also Table S2 for a list of all plasmids used in this study). A detailed protocol of expression and purification can be found on protocols.io (https://dx.doi.org/10.17504/protocols.io.48ggztw). Briefly, proteins were expressed as GST-fusion proteins in *E. coli* BL21 [DE3] or NEB Express (New England Biolabs) and lysed in 50 mM Tris/HCl (pH8), 150 mM NaCl, 0.5 mM EDTA, 1 mM DTT (+ 5 mM MgCl$_2$, 1 mM ATP for KaiC proteins). Purification was performed via *batch* affinity chromatography using glutathione agarose 4B (Macherey and Nagel) or glutathione sepharose 4B (GE Healthcare) in the same buffer. Finally, the GST-tag was cleaved off using PreScission protease (GE Healthcare) in 50 mM Tris/HCl (pH8), 150 mM NaCl, 1 mM EDTA, 1 mM DTT (+ 5 mM MgCl$_2$, 1 mM ATP for KaiC proteins). If homogeneity of the proteins was not sufficient, they were further purified via anion exchange chromatography using a MonoQ or ResourceQ column (GE Healthcare Life Sciences) using 50 mM

Tris/HCl (pH8), 1 mM EDTA, 1 mM DTT (+ 5 mM MgCl$_2$, 1 mM ATP for KaiC proteins) and a 0-1M

NaCl gradient.

To produce Strep-KaiC3$_{6803}$ variants with amino acid substitutions, the *kaiC3$_{6803}$* gene in the pGEX-

kaiC3$_{6803}$ vector was modified by site directed mutagenesis using the Quick-Change Site-Directed

Mutagenesis Kit (Stratagene) or Q5 Site-Directed Mutagenesis Kit (New England Biolabs). Base

triplets encoding S423 and T424 were changed to code for alanine or for aspartate and glutamate

resulting in *kaiC3$_{6803}$-AA* and *kaiC3$_{6803}$-DE* genes, respectively. To generate *kaiC3$_{6803}$-catE1$^-$catE2*,

two subsequent site-directed mutagenesis reactions were performed to exchange nucleotides

encoding E67 and E68 as well as nucleotides encoding E310 and E311 all with bases encoding

glutamine (all primers used for mutagenesis are listed in Table S1). Afterwards, *kaiC3$_{6803}$* WT and

modified *kaiC3$_{6803}$* genes were amplified with KOD-Plus-Neo polymerase (Toyobo) using pASK-

kaiC3 primers (Table S1). Amplicons were digested with SacII and HindIII and ligated into the

respective restriction sites of pASK-IBA5plus (IBA Life sciences). For purification of recombinant

Strep-KaiC$_{7942}$, the pASK-IBA-5plus based vector described in Oyama *et al.* (14) was used. Strep-

KaiC3$_{6803}$ proteins were expressed in *E. coli* Rosetta gamiB (DE3) or Rosetta gami2 (DE3) cells

(Novagen). Expression of Strep-KaiC$_{7942}$ was carried out in *E. coli* DH5α. Cells were cultured in LB

medium containing 100 μg ml^{-1} ampicillin with vigorous agitation at 37 °C. At OD$_{600nm}$ 0.33-0.68

protein expression was induced with 200 ng ml^{-1} anhydrotetracycline and the strains were further

incubated as following: Strep-KaiC$_{7942}$: 7h at 37°C, Strep-KaiC3$_{6803}$-WT: 5h at 35°C or 3.5h at 37°C,

Strep-KaiC3$_{6803}$-AA and Strep-KaiC3$_{6803}$-catE1$^-$catE2$^-$: 18°C overnight, Strep-KaiC3$_{6803}$-DE: 25°C

overnight. Cells were harvested and lysed by sonication in ice-cold buffer W [20mM Tris/HCl

(pH8), 150 mM NaCl, 5 mM MgCl$_2$, 1 mM ATP (+ 2 mM DTT for Strep-KaiC3$_{6803}$ proteins)] including

protease inhibitors (protease inhibitor cocktail, Roche or Nacalai). Soluble proteins were loaded

on self-prepared columns packed with Strep-Tactin XT superflow or Strep-Tactin Sepharose (IBA

lifesciences) and purified under gravity flow. After washing with buffer W, Strep-KaiC proteins

were eluted with ice cold buffer W + 50 mM D(+)biotin (for Strep-Tactin XT Superflow) or ice cold

buffer W + 2.5 mM desthiobiotin (for Strep tactin Superflow). See

https://dx.doi.org/10.17504/protocols.io.meac3ae for a detailed protocol.

All proteins used for ATPase activity measurements were further purified via size exclusion

chromatography. Strep-KaiC3$_{6803}$ and Strep-KaiC$_{7942}$ proteins were applied on a Sephacryl S300

HR HiPrep 16/60 Sephacryl column (GE Healthcare) and separated in 20 or 50 mM Tris/HCl (pH

8.0), 150 mM NaCl, 2 mM DTT, 1 mM ATP and 5 mM MgCl2. For separation of KaiB1$_{6803}$ and

KaiB3$_{6803}$, a Sephacryl S200 HR HiPrep 16/60 column (GE Healthcare) or Superdex 200 Increase

10/30 GL column (GE Healthcare) and 20 or 50 mM Tris/HCl (pH 8.0), 150 mM NaCl, 2 mM DTT as

running buffer were used. KaiA$_{6803}$ was purified on a Superdex 200 Increase 10/30 GL column (GE

Healthcare) in 20 mM Tris/HCl (pH 8.0), 150 mM NaCl, 2 mM DTT. See

https://dx.doi.org/10.17504/protocols.io.mdtc26n for further details.

ATPase activity

For ATPase measurements, KaiC proteins fused to an N-terminal Strep-tag were used and all Kai

proteins were purified via size exclusion chromatography (see above). 3.45 µM of Strep-KaiC3$_{6803}$

variants were incubated in ATPase buffer (20 mM Tris/HCl (pH8), 150 mM NaCl, 1 mM ATP, 5 mM

MgCl$_2$) at 25 °C, 30 °C or 35 °C for 24 hours. To analyze the influence of KaiA$_{6803}$ and KaiB$_{6803}$

proteins on KaiC3$_{6803}$ ATPase activity, we mixed 0.2 mg ml^{-1} KaiC3$_{6803}$ with 0.04 mg ml^{-1} KaiA$_{6803}$

or 0.04 mg ml^{-1} KaiB$_{6803}$ and incubated the mixtures for 24 hours at 30 °C. Monomeric and

oligomeric KaiB3$_{6803}$ were analyzed separately. To monitor ADP production, every 3, 4 or 6 hours

2 µl of the reaction mixture were applied on a Shim-Pack-VP-ODS column (Shimadzu) and

separated using 100 mM phosphoric acid, 150 mM triethylamine, 1 % acetonitrile as running

buffer. ADP production per monomer KaiC and 24 hours was calculated using a calibration curve.

A detailed protocol can be found on protocols.io

(https://dx.doi.org/10.17504/protocols.io.mebc3an). The Q$_{10}$ value was calculated from ATPase

measurements at 25 °C and 35 °C, using the formula $Q_{10} = \left(\frac{R_2}{R_1}\right)^{10°C/(T_2 - T_1)}$

ATP synthase activity

To investigate dephosphorylation via ATP synthesis, we used KaiC proteins which were expressed

as GST-fusion proteins and subsequently cleaved off their GST-tag. 3 µM KaiC in ATP synthesis

buffer (20 mM Tris/HCl(pH8), 150 mM NaCl, 0.5 mM EDTA, 5 mM MgCl$_2$, 0.5 mM ATP) was mixed

with 0.8 µCi ml^{-1} [α^{32}P]ADP and stored at -20 °C or incubated for 2 hours at 30 °C. As a control,

the same experiment was performed in the presence of 0.5 mM ADP. After 20-fold dilution,

nucleotides in a 0.5 µl reaction mixture were separated via thin layer chromatography using TLC

PEI Cellulose F plates (Merck Millipore) and 1 M LiCl as solvent. [α-^{32}P]ADP and [γ-^{32}P]ATP were

separated in parallel to identify the signals corresponding to ADP and ATP, respectively. Dried

plates were subjected to autoradiography and signals were analysed using a Personal Molecular

Imager FX system (Bio-Rad) and ImageLab software (Bio-Rad). For each reaction mixture, the

relative intensity of [α-^{32}P]ATP was calculated as percentage of all signals in the corresponding

lane. Because [α-^{32}P]ATP was already synthesized during mixing of the samples, the relative ATP

intensity measured in the -20 °C sample containing 0.5 mM ADP was subtracted for normalization.

The principle of this method is based on Egli *et al.* (12). A detailed protocol is available on

protocols.io (https://dx.doi.org/10.17504/protocols.io.48qgzvw).

Yeast two-hybrid assays

For yeast two-hybrid assays, vectors containing the GAL4 activation domain (AD) and the GAL4

DNA-binding domain (BD) were used. Genes of interest were amplified from *Synechocystis* 6803

wild type genomic DNA with the Phusion Polymerase (NEB) according to manufacturer's

guidelines. Indicated restriction sites were introduced via the oligonucleotides listed in Table S1.

Vectors and PCR fragments were cut with the respective restriction enzymes (Thermo Fisher

Scientific) and the gene of interest was ligated into the vector, leading to a fusion protein with an

AD- or BD-tag either at the N- or C-terminus. All constructed plasmids are listed in Table S2.

Transformation of yeast cells was performed according to manufacturer's guidelines using the

Frozen-EZ Yeast Transformation II Kit (Zymo Research) and cells were selected on complete

supplement mixture (CSM) lacking leucine and tryptophan (-Leu -Trp) dropout medium (MP

Biochemicals) at 30 °C for 3–4 days. Y190 (Clontech) cells were used for measuring

β-galactosidase activity. Formed colonies were spotted on a second plate (CSM -Leu -Trp) and

incubated for 2 days. Afterwards a colony-lift filter assay was performed as described by Breeden

et al. (52). A detailed protocol can be found on protocols.io

(https://dx.doi.org/10.17504/protocols.io.v7ve9n6).

Bacterial strains and growth conditions

Wild type *Synechocystis* 6803 (PCC-M, re-sequenced, (53)) and the *kaiC3* deletion mutant (38)

were cultured photoautotrophically in BG11 medium (54) supplemented with 10 mM TES buffer

(pH 8) under constant illumination with 50 µmol photons $m^{-2}s^{-1}$ of white light (Philips TLD Super

80/840) at 30 °C. Cells were grown either in Erlenmeyer flasks with constant shaking (140 rpm) or

on plates (0.75 % Bacto-Agar, Difco) supplemented with 0.3 % thiosulfate. Detailed recipes can

be found on protocols.io (https://dx.doi.org/10.17504/protocols.io.wj5fcq6).

Spot assays

Experiments were performed as previously described (44). Strains were propagated

mixotrophically on BG11 agar plates with the addition of 0.2 % (w/v) glucose. Dilution series of

cell cultures started with OD_{750nm} 0.2 and OD_{750nm} 0.4, followed by incubation of the plates for 6

or 28 days under constant light conditions and in complete darkness, respectively.

Acknowledgement:

We thank Junko Moriwaki, Nancy Sauer, Jennifer Andres, Lukas Pohlig, Thomas Volkmer and

Megumi Fujimoto for technical assistance. This work was funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence

Strategy – EXC 2048/1 – Project ID: 390686111 to A. Wiegard and I.M. Axmann. The work was

financially supported by Heine Research Academies - Travel Grants, and EMBO ASTF 145 to A.

Wiegard, by grants AX 84/1-3 and EXC 1028 from the German Research Foundation to I.M.

Axmann, by JSPS Grants-in-Aid 16H00784, 17K19247 and 19K05833 to K. Terauchi, and by grant

WI2014/5-3 from the German Research Foundation to A. Wilde.

Author contributions:

I.M. Axmann, A. Wilde, A. Wiegard, C. Köbler and K. Terauchi conceived the project. A. Wiegard,

C. Köbler A.K. Dörrich and K. Oyama designed, performed and analyzed experiments. I.M Axmann,

A. Wilde and K. Terauchi supervised the study. All authors interpreted and discussed the data. A.

Wiegard, C. Köbler and A. Wilde wrote the manuscript. I.M. Axmann, C. Azai, K. Terauchi, A.K.

Dörrich, and K. Oyama commented essentially on the manuscript. All authors approved the

manuscript.

This manuscript contains supplementary information:

Supplementary methods

Table S1. Oligonucleotides used in this study.

Table S2. Plasmids used in this study.

Figure S1. Initial phosphorylation level of KaiC proteins.

Figure S2. Original scans of the KaiC3 yeast two-hybrid interactions.

Figure S3. Interaction of $KaiB_{6803}$ and $KaiC_{6803}$ proteins in pull down analysis.

Figure S4. Purification of KaiB1 and KaiB3 proteins used for ATPase analysis

Figure S5. ATPase activity and dephosphorylation of $KaiC3_{6803}$ are temperature dependent

References:

1. Ditty JL, Williams SB, Golden SS. 2003. A cyanobacterial circadian timing mechanism. Annu Rev Genet 37:513-43.

2. Nakajima M, Imai K, Ito H, Nishiwaki T, Murayama Y, Iwasaki H, Oyama T, Kondo T. 2005. Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation *in vitro*. Science 308:414-5.

3. Iwasaki H, Nishiwaki T, Kitayama Y, Nakajima M, Kondo T. 2002. KaiA-stimulated KaiC phosphorylation in circadian timing loops in cyanobacteria. Proc Natl Acad Sci U S A 99:15788-93.

4. Terauchi K, Kitayama Y, Nishiwaki T, Miwa K, Murayama Y, Oyama T, Kondo T. 2007. ATPase activity of KaiC determines the basic timing for circadian clock of cyanobacteria. Proc Natl Acad Sci U S A 104:16377-81.

5. Kitayama Y, Iwasaki H, Nishiwaki T, Kondo T. 2003. KaiB functions as an attenuator of KaiC phosphorylation in the cyanobacterial circadian clock system. EMBO J 22:2127-34.

6. Hayashi F, Suzuki H, Iwase R, Uzumaki T, Miyake A, Shen JR, Imada K, Furukawa Y, Yonekura K, Namba K, Ishiura M. 2003. ATP-induced hexameric ring structure of the cyanobacterial circadian clock protein KaiC. Genes Cells 8:287-96.

7. Mori T, Saveliev SV, Xu Y, Stafford WF, Cox MM, Inman RB, Johnson CH. 2002. Circadian clock protein KaiC forms ATP-dependent hexameric rings and binds DNA. Proc Natl Acad Sci U S A 99:17203-8.

8. Pattanayek R, Wang J, Mori T, Xu Y, Johnson CH, Egli M. 2004. Visualizing a circadian clock protein: crystal structure of KaiC and functional insights. Mol Cell 15:375-88.

9. Nishiwaki T, Iwasaki H, Ishiura M, Kondo T. 2000. Nucleotide binding and autophosphorylation of the clock protein KaiC as a circadian timing process of cyanobacteria. Proc Natl Acad Sci U S A 97:495-9.

10.   Murakami R, Miyake A, Iwase R, Hayashi F, Uzumaki T, Ishiura M. 2008. ATPase activity and its

temperature compensation of the cyanobacterial clock protein KaiC. Genes Cells 13:387-95.

11.   Nishiwaki T, Kondo T. 2012. Circadian autodephosphorylation of cyanobacterial clock protein KaiC

occurs via formation of ATP as intermediate. J Biol Chem 287:18030-5.

12.   Egli M, Mori T, Pattanayek R, Xu Y, Qin X, Johnson CH. 2012. Dephosphorylation of the core clock

protein KaiC in the cyanobacterial KaiABC circadian oscillator proceeds via an ATP synthase

mechanism. Biochemistry 51:1547-58.

13.   Phong C, Markson JS, Wilhoite CM, Rust MJ. 2013. Robust and tunable circadian rhythms from

differentially sensitive catalytic domains. Proc Natl Acad Sci U S A 110:1124-9.

14.   Oyama K, Azai C, Nakamura K, Tanaka S, Terauchi K. 2016. Conversion between two

conformational states of KaiC is induced by ATP hydrolysis as a trigger for cyanobacterial circadian

oscillation. Sci Rep 6:32443.

15.   Shultzaberger RK, Boyd JS, Diamond S, Greenspan RJ, Golden SS. 2015. Giving Time Purpose: The

Synechococcus elongatus Clock in a Broader Network Context. Annu Rev Genet 49:485-505.

16.   Dong G, Yang Q, Wang Q, Kim YI, Wood TL, Osteryoung KW, van Oudenaarden A, Golden SS. 2010.

Elevated ATPase activity of KaiC applies a circadian checkpoint on cell division in *Synechococcus*

elongatus. Cell 140:529-39.

17.   Swan JA, Golden SS, LiWang A, Partch CL. 2018. Structure, function, and mechanism of the core

circadian clock in cyanobacteria. J Biol Chem 293:5026-5034.

18.   Shih PM, Wu D, Latifi A, Axen SD, Fewer DP, Talla E, Calteau A, Cai F, Tandeau de Marsac N, Rippka

R, Herdman M, Sivonen K, Coursin T, Laurent T, Goodwin L, Nolan M, Davenport KW, Han CS, Rubin

EM, Eisen JA, Woyke T, Gugger M, Kerfeld CA. 2013. Improving the coverage of the cyanobacterial

phylum using diversity-driven genome sequencing. Proc Natl Acad Sci U S A 110:1053-8.

19.   Schmelling NM, Lehmann R, Chaudhury P, Beck C, Albers S-VV, Axmann IMM, Wiegard A. 2017.

Minimal Tool Set for a Prokaryotic Circadian Clock. bioRxiv doi:10.1101/075291.

20. Axmann IM, Hertel S, Wiegard A, Dorrich AK, Wilde A. 2014. Diversity of KaiC-based timing systems

in marine Cyanobacteria. Mar Genomics 14C:3-16.

21. Aoki S, Onai K. 2009. Circadian clocks of *Synechocystis* sp. strain PCC 6803, *Thermosynechococcus*

elongatus, *Prochlorococcus* spp., *Trichodesmium* spp. and other species, p 259-282. *In* Ditty JL,

Mackey SR, Johnson CH (ed), Bacterial circadian programs. Springer Berlin Heidelberg.

22. Dvornyk V, Vinogradova O, Nevo E. 2003. Origin and evolution of circadian clock genes in

prokaryotes. Proc Natl Acad Sci U S A 100:2495-500.

23. Wiegard A, Dörrich AK, Deinzer HT, Beck C, Wilde A, Holtzendorff J, Axmann IM. 2013. Biochemical

analysis of three putative KaiC clock proteins from *Synechocystis* sp. PCC 6803 suggests their

functional divergence. Microbiology 159:948-958.

24. Johnson CH, Egli M. 2014. Metabolic compensation and circadian resilience in prokaryotic

cyanobacteria. Annu Rev Biochem 83:221-47.

25. Holtzendorff J, Partensky F, Mella D, Lennon JF, Hess WR, Garczarek L. 2008. Genome streamlining

results in loss of robustness of the circadian clock in the marine cyanobacterium *Prochlorococcus*

marinus PCC 9511. J Biol Rhythms 23:187-99.

26. Kanesaki Y, Shiwa Y, Tajima N, Suzuki M, Watanabe S, Sato N, Ikeuchi M, Yoshikawa H. 2012.

Identification of substrain-specific mutations by massively parallel whole-genome resequencing of

Synechocystis sp. PCC 6803. DNA Res 19:67-79.

27. Kucho K-i, Okamoto K, Tsuchiya Y, Nomura S, Nango M, Kanehisa M, Ishiura M. 2005. Global

analysis of circadian expression in the cyanobacterium *Synechocystis* sp. strain PCC 6803. J

Bacteriol 187:2190-9.

28. Beck C, Hertel S, Rediger A, Lehmann R, Wiegard A, Kolsch A, Heilmann B, Georg J, Hess WR,

Axmann IM. 2014. A daily expression pattern of protein-coding genes and small non-coding RNAs

in Synechocystis sp. PCC 6803. Appl Environ Microbiol doi:10.1128/AEM.01086-14.

29. van Alphen P, Hellingwerf KJ. 2015. Sustained Circadian Rhythms in Continuous Light in

Synechocystis sp. PCC6803 Growing in a Well-Controlled Photobioreactor. PLoS One 10:e0127715.

30. Saha R, Liu D, Hoynes-O'Connor A, Liberton M, Yu J, Bhattacharyya-Pakrasi M, Balassy A, Zhang F,

Moon TS, Maranas CD, Pakrasi HB. 2016. Diurnal Regulation of Cellular Processes in the

Cyanobacterium Synechocystis sp. Strain PCC 6803: Insights from Transcriptomic, Fluxomic, and

Physiological Analyses. MBio 7.

31. Axmann IM, Dühring U, Seeliger L, Arnold A, Vanselow JT, Kramer A, Wilde A. 2009. Biochemical

evidence for a timing mechanism in *Prochlorococcus*. J Bacteriol 191:5342-7.

32. Ma P, Mori T, Zhao C, Thiel T, Johnson CH. 2016. Evolution of KaiC-Dependent Timekeepers: A

Proto-circadian Timing Mechanism Confers Adaptive Fitness in the Purple Bacterium

Rhodopseudomonas palustris. PLoS Genet 12:e1005922.

33. Mukaiyama A, Ouyang D, Furuike Y, Akiyama S. 2019. KaiC from a cyanobacterium Gloeocapsa sp.

PCC 7428 retains functional and structural properties required as the core of circadian clock

system. International Journal of Biological Macromolecules 131:67-73.

34. Diamond S, Jun D, Rubin BE, Golden SS. 2015. The circadian oscillator in Synechococcus elongatus

controls metabolite partitioning during diurnal growth. Proc Natl Acad Sci U S A 112:E1916-25.

35. Welkie DG, Rubin BE, Diamond S, Hood RD, Savage DF, Golden SS. 2019. A Hard Day's

Night: Cyanobacteria in Diel Cycles. Trends Microbiol 27:231-242.

36. Pattanayak G, Rust MJ. 2014. The cyanobacterial clock and metabolism. Curr Opin Microbiol 18:90-

5.

37. Woelfle MA, Ouyang Y, Phanvijhitsiri K, Johnson CH. 2004. The adaptive value of circadian clocks:

an experimental assessment in cyanobacteria. Curr Biol 14:1481-6.

38. Dorrich AK, Mitschke J, Siadat O, Wilde A. 2014. Deletion of the Synechocystis sp. PCC 6803

kaiAB1C1 gene cluster causes impaired cell growth under light-dark conditions. Microbiology

doi:10.1099/mic.0.081695-0.

39.   Kobler C, Schultz SJ, Kopp D, Voigt K, Wilde A. 2018. The role of the Synechocystis sp. PCC 6803

homolog of the circadian clock output regulator RpaA in day-night transitions. Mol Microbiol

110:847-861.

40.   Ungerer J, Wendt KE, Hendry JI, Maranas CD, Pakrasi HB. 2018. Comparative genomics reveals the

molecular determinants of rapid growth of the cyanobacterium Synechococcus elongatus UTEX

2973. Proc Natl Acad Sci U S A doi:10.1073/pnas.1814912115.

41.   Welkie DG, Rubin BE, Chang YG, Diamond S, Rifkin SA, LiWang A, Golden SS. 2018. Genome-wide

fitness assessment during diurnal growth reveals an expanded role of the cyanobacterial circadian

clock protein KaiA. Proc Natl Acad Sci U S A 115:E7174-e7183.

42.   Diamond S, Rubin BE, Shultzaberger RK, Chen Y, Barber CD, Golden SS. 2017. Redox crisis underlies

conditional light-dark lethality in cyanobacterial mutants that lack the circadian regulator, RpaA.

Proc Natl Acad Sci U S A 114:E580-e589.

43.   Anderson SL, McIntosh L. 1991. Light-activated heterotrophic growth of the cyanobacterium

Synechocystis sp. strain PCC 6803: a blue-light-requiring process. J Bacteriol 173:2761-7.

44.   Dörrich AK, Wilde A. 2015. Spot Assays for Viability Analysis of Cyanobacteria. Bio-protocol

5:e1574.

45.   Loza-Correa M, Sahr T, Rolando M, Daniels C, Petit P, Skarina T, Gomez Valero L, Dervins-Ravault

D, Honore N, Savchenko A, Buchrieser C. 2014. The Legionella pneumophila kai operon is

implicated in stress response and confers fitness in competitive environments. Environ Microbiol

16:359-81.

46.   Snijder J, Burnley RJ, Wiegard A, Melquiond AS, Bonvin AM, Axmann IM, Heck AJ. 2014. Insight

into cyanobacterial circadian timing from structural details of the KaiB-KaiC interaction. Proc Natl

Acad Sci U S A 111:1379-84.

47. Hitomi K, Oyama T, Han S, Arvai AS, Getzoff ED. 2005. Tetrameric architecture of the circadian

clock protein KaiB. A novel interface for intermolecular interactions and its impact on the circadian

rhythm. J Biol Chem 280:19127-35.

48. Snijder J, Schuller JM, Wiegard A, Lossl P, Schmelling N, Axmann IM, Plitzko JM, Forster F, Heck AJ.

2017. Structures of the cyanobacterial circadian oscillator frozen in a fully assembled state. Science

355:1181-1184.

49. Tseng R, Goularte NF, Chavan A, Luu J, Cohen SE, Chang YG, Heisler J, Li S, Michael AK, Tripathi S,

Golden SS, LiWang A, Partch CL. 2017. Structural basis of the day-night transition in a bacterial

circadian clock. Science 355:1174-1180.

50. Chang YG, Cohen SE, Phong C, Myers WK, Kim YI, Tseng R, Lin J, Zhang L, Boyd JS, Lee Y, Kang S,

Lee D, Li S, Britt RD, Rust MJ, Golden SS, LiWang A. 2015. Circadian rhythms. A protein fold switch

joins the circadian oscillator to clock output in cyanobacteria. Science 349:324-8.

51. Espinosa J, Boyd JS, Cantos R, Salinas P, Golden SS, Contreras A. 2015. Cross-talk and regulatory

interactions between the essential response regulator RpaB and cyanobacterial circadian clock

output. Proc Natl Acad Sci U S A 112:2198-203.

52. Breeden L, Nasmyth K. 1985. Regulation of the yeast HO gene. Cold Spring Harb Symp Quant Biol

50:643-50.

53. Trautmann D, Voss B, Wilde A, Al-Babili S, Hess WR. 2012. Microevolution in cyanobacteria: re-

sequencing a motile substrain of Synechocystis sp. PCC 6803. DNA Res 19:435-48.

54. Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. 1979. Generic assignments, strain

histories and properties of pure cultures of cyanobacteria. J Gen Microbiol 111:1-61.

 Figure legends:

 Figure 1: KaiC3$_{6803}$ showed decreased ATP synthase activity compared to KaiC$_{7942}$. Prior to the

 experiment, KaiC$_{7942}$ was incubated for 2 weeks at 4 °C or overnight at 30 °C to generate fully

 phosphorylated (P-KaiC$_{7942}$) and dephosphorylated (NP-KaiC$_{7942}$) protein, respectively (Fig. S1). **A:**

 Representative autoradiograph of separation of [γ-^{32}P]ATP (product) and [α-^{32}P]ADP (substrate)

 via thin layer chromatography after incubation with indicated KaiC proteins for 2h. Controls show

 the ATP signal in the presence of an excess of cold ADP. For size control [γ-^{32}P]ATP and [α-^{32}P]ADP

 were separated on the same cellulose F plate. **B:** Relative ATP signals after 2 hours incubation

 displayed as percentage of all radioactive signals in the corresponding lane (mean with standard

 deviation from two experiments, each analysed in duplicates). All values are normalised to the

 relative ATP signal at 0h incubation time in the presence of an excess of cold ADP.

 Figure 2: ATPase activity of Strep-KaiC3$_{6803}$ and variants Strep-KaiC3$_{6803}$-DE and Strep-KaiC3$_{6803}$-

 AA. Strep-KaiC3$_{6803}$, Strep-KaiC3$_{6803}$-DE and Strep-KaiC3$_{6803}$-AA were incubated for 24 hours at 30

 °C and ADP production per day and per Strep-KaiC3$_{6803}$ monomer was calculated. As a control we

 used Strep-KaiC3$_{6803}$-catE1$^-$catE2$^-$, which lacks both ATPase motifs due to replacement of catalytic

 glutamates 67,68, 310 and 311 with glutamine residues. For comparison ADP production by

 KaiC$_{7942}$ was monitored for 24 or 48 hours. Strep-KaiC$_{7942}$ was dephosphorylated prior to the

 experiment by incubation at 30°C overnight. Shown are mean values with standard deviation of

 at least five replicates. Please note that we averaged all measurements for Strep-KaiC3$_{6803}$ WT at

 30 °C and therefore repeatedly display them in Fig. 2, Fig. 4A and Fig. 4B.

 Figure 3. KaiC3$_{6803}$ interacts with KaiB3$_{6803}$ and the proteins of the main oscillator KaiC1$_{6803}$,

 KaiB1$_{6803}$. A, B, C: Yeast two-hybrid reporter strains carrying the respective bait and prey plasmids

were selected by plating on complete supplement medium (CSM) lacking leucine and tryptophan

(-Leu -Trp). Physical interaction between bait and prey fusion proteins is indicated by a colour

change in the assays using 5-brom-4-chlor-3-indoxyl-β-D-galactopyranosid. **AD:** GAL4 activation

domain; BD, GAL4 DNA-binding domain. Shown are representative results of two replicates. For

clear presentation, spots were assembled from several assays (original scans are shown in Fig.

S2).

Figure 4: ATPase activity of Strep-KaiC3$_{6803}$ is regulated by KaiB proteins and is temperature

dependent. A: ATPase activity of Strep-KaiC3$_{6803}$ was decreased in the presence of KaiB1$_{6803}$ and

monomeric KaiB3$_{6803}$, but was not influenced by oligomeric KaiB3$_{6803}$ and by KaiA$_{6803}$. Shown are

mean values with standard deviation of at least three replicates. Controls show that KaiA$_{6803}$ and

KaiB$_{6803}$ proteins alone did not display ATPase activity (referring to Strep-KaiC3$_{6803}$ monomer

activity) **B:** ATPase activity of Strep-KaiC3$_{6803}$ is not temperature-compensated. Strep-KaiC3$_{6803}$

was incubated for 24 hours at the indicated temperatures and ADP production per day and

monomer Strep-KaiC3$_{6803}$ was calculated. Shown are mean values with standard deviation of at

least three replicates. Please note that we averaged all measurements for Strep-KaiC3$_{6803}$ WT at

30 °C and therefore repeatedly display them in Fig. 2, Fig. 4A and Fig. 4B.

Figure 5. Reduced viability of the Δ*kaiC3* strain in complete darkness. Viability of the

Synechocystis 6803 wild type, the *kaiAB1C1* deletion mutant (Δ*kaiAB1C1*) and the *kaiC3* deletion

mutant (Δ*kaiC3*) was tested under mixotrophic conditions in continuous light and dark conditions.

Strains were grown in liquid culture under constant light and different dilutions were spotted on

agar plates containing 0.2 % glucose. Plates were analysed after further incubation for 6 or 28

days of continuous light and darkness, respectively. A representative result of three independent

experiments is shown.

Figure 6: Model of the crosstalk between the KaiAB1C1$_{6803}$ standard oscillator and KaiC3$_{6803}$ in

***Synechocystis* 6803.** KaiC1$_{6803}$ forms a standard oscillator together with KaiA$_{6803}$ and KaiB1$_{6803}$.

The KaiAB1C1$_{6803}$ oscillator allows temporal metabolic orchestration. KaiC3$_{6803}$ forms an

additional regulatory mechanism together with KaiB3$_{6803}$. The two systems intertwine by

interactions of the KaiC$_{6803}$ proteins with each other and the non-corresponding KaiB$_{6803}$

homologs. Solid lines indicate experimentally verified physical interactions (this paper and

Wiegard *et al*.(23)), dotted arrows indicate stimulation of phosphorylation (23) and dotted lines

with bars show inhibition of ATPase activity.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6