1 **Title:**
2 Extent and context dependence of pleiotropy revealed by high-throughput single-cell
3 phenotyping
4
Author list:

Kerry A Geiler-Samerotte[a,b]

Shuang Li[a]

Charalampos Lazaris[a]

Austin Taylor[a]

Naomi Ziv[a]

Chelsea Ramjeawan[a]

Annalise B Paaby[c]

Mark L Siegal[a]

14
Author affiliations:

[a] Center for Genomics and Systems Biology, Department of Biology, New York
University, New York, NY 10003

[b] Center for Mechanisms of Evolution, Biodesign Institutes, School of Life Sciences,
Arizona State University, Tempe, AZ 85287

[c] School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332
21
22
23 **Corresponding author:**
24 Kerry Geiler-Samerotte: kerry.samerotte@asu.edu

Abstract:

Pleiotropy – when a single mutation affects multiple traits – is a controversial topic with far-reaching implications. Pleiotropy plays a central role in debates about how complex traits evolve and whether biological systems are modular or are organized such that every gene has the potential to affect many traits. Pleiotropy is also critical to initiatives in evolutionary medicine that seek to trap infectious microbes or tumors by selecting for mutations that encourage growth in some conditions at the expense of others. Research in these fields, and others, would benefit from understanding the extent to which pleiotropy reflects inherent relationships among phenotypes that correlate no matter the perturbation (vertical pleiotropy). Alternatively, pleiotropy may result from genetic changes that impose correlations between otherwise independent traits (horizontal pleiotropy). We distinguish these possibilities by using clonal populations of yeast cells to quantify the inherent relationships between single-cell morphological features. Then, we demonstrate how often these relationships underlie vertical pleiotropy and how often these relationships are modified by genetic variants (QTL) acting via horizontal pleiotropy. Our comprehensive screen measures thousands of pairwise trait correlations across hundreds of thousands of yeast cells and reveals ample evidence of both vertical and horizontal pleiotropy. Additionally, we observe that the correlations between traits can change with the environment, genetic background and cell-cycle position. These changing dependencies suggest a nuanced view of pleiotropy: biological systems demonstrate limited pleiotropy in any given context, but across contexts (e.g., across diverse environments and genetic backgrounds) each genetic change has the potential to influence a larger number of traits. Our method suggests that exploiting pleiotropy for applications in evolutionary medicine would benefit from focusing on traits with correlations that are less dependent on context.

Introduction

Pleiotropy exists when a single mutation affects multiple traits (1,2). Often, pleiotropy is defined instead as a single gene contributing to multiple traits, although what is implied is the original definition — that a single change at the genetic level can have multiple consequences at the phenotypic level (2). As our ability to survey the influence of genotype on phenotype improves, examples of pleiotropy are growing (3-8). For example, individual genetic variants have been associated with seemingly disparate immune, neurological, and digestive symptoms in humans and mice (9,10). Genes affecting rates of cell division across diverse environments and drug treatments have been identified in microbes and cancers (11,12). A view emerging from genome-wide association studies is that variation in complex traits is "omnigenic" in the sense that many loci indirectly contribute to variation in many traits (13,14).

However, the extent of pleiotropy remains a major topic of debate because, despite its apparent prevalence, pleiotropy is thought to be evolutionarily disadvantageous. The more traits a mutation affects, the more likely it is that the mutation will have a negative impact on at least one. Pervasive pleiotropy should therefore constrain evolution (15), exacting what is known as a cost of complexity or cost of pleiotropy (11,16-19). This cost may bias which mutations underlie adaptation, for example, toward less-pleiotropic *cis*-regulatory changes over more-pleiotropic changes in *trans*-acting factors (20,21), or toward changes to proteins that participate in relatively few biological processes (22,23). Over long periods, the cost of pleiotropy may influence the organization of biological systems, favoring a modular structure in which genetic changes influencing one group of traits have minimal impact system-wide (24-29).

At stake in the ongoing debate about the extent of pleiotropy (30-33) are some of modern biology's prime objectives, including the prediction of complex phenotypes from genotype data (18,34,35) and the prediction of how organisms will adapt to environmental change (36,37). These predictions are more challenging if genetic changes influence a large number of traits with complex interdependencies. Nonetheless, understanding how a given mutation influences multiple traits could be powerful, allowing prediction of some phenotypic responses given others (38,39). Indeed, recent strategies in medicine called evolutionary traps aim to exploit pleiotropy, for example by finding genetic changes that provide resistance to one treatment while promoting susceptibility to another (40-42).

The lack of consensus about the extent of pleiotropy in natural systems is in part due to poorly defined expectations for how to test for it experimentally. One key issue is that defining a phenotype is not trivial (43,44). Consider a variant in the *apolipoprotein B* gene that increases low-density lipoprotein (LDL) cholesterol levels as well as the risk of heart disease. Elevated LDL promotes heart disease (45), so are these two phenotypes or one? Alternatively, consider a mutation in the *phenylalanine hydroxylase* gene that affects nervous system function and skin pigmentation. These dissimilar effects, both symptoms of untreated phenylketonuria (PKU), originate from the same problem: a deficiency in converting phenylalanine to tyrosine (46). Is it appropriate to call mutations that have this single metabolic effect pleiotropic? Likewise, shall one call pleiotropic a mutation that makes tomatoes both ripen uniformly and taste bad, when the effect of the

mutation is to reduce the function of a transcription factor that promotes chloroplast
development, which in turn necessarily affects both coloration and sugar accumulation
(47)?

The LDL, PKU and tomato cases are examples of vertical pleiotropy, *i.e.*
pleiotropy that results when one phenotype influences another or both are influenced by a
shared factor (5,43). The alternative to vertical pleiotropy is horizontal pleiotropy, in
which genetic differences induce correlations between otherwise independent
phenotypes. It might be tempting to discard vertical pleiotropy as less "genuine" (48) or
less important than horizontal pleiotropy, but that would be a mistake because vertical
pleiotropy reveals important information about the underlying biological systems that
produce the phenotypes in question. Consider the value in identifying yet-unknown
factors in heart disease by finding traits that correlate with it, or in understanding where
in a system an intervention is prone to produce undesirable side effects. Consider also
that the extent and nature of vertical pleiotropy speak directly to the question of
modularity: modularity is implied if vertical pleiotropy either is rare or manifests as small
groups of correlated traits that are isolated from other such groups. If there is modularity
then there can be horizontal pleiotropy, when particular genetic variants make links
between previously unconnected modules.

The above considerations suggest that a unified analysis that distinguishes and
compares horizontal and vertical pleiotropy is needed to make sense of the organization
and evolution of biological systems. However, existing methods of distinguishing
horizontal and vertical pleiotropy are problematic because judgments must be made about
which traits are independent from one another. Such judgments differ between
researchers and over time. Indeed, the tomato example can be viewed as a case of
horizontal pleiotropy transitioning recently to vertical pleiotropy as knowledge of the
underlying system advanced.

In this study, we propose and apply an empirical and analytical approach to
measuring pleiotropy that relies far less on subjective notions of what constitutes an
independent phenotype. The key principle is that the distinction between vertical and
horizontal pleiotropy lies in whether traits are correlated in the absence of genetic
variation (43). For vertical pleiotropy, the answer is yes: because one trait influences the
other or the two share an influence, non-genetic perturbations that alter one phenotype are
expected to alter the other. For horizontal pleiotropy, the answer is no: genetic variation
causes the trait correlation. In this study, we determined how traits correlate in the
absence of genetic variation by measuring single-cell traits in clonal populations of cells.

We used high-throughput morphometric analysis (49-53) of hundreds of
thousands of single cells of the budding yeast *Saccharomyces cerevisiae* to measure how
dozens of cell-morphology traits (thousands of pairs of traits) co-vary within clonal
populations and between such populations representing different genotypes. Within-
genotype correlations report on vertical pleiotropy, whereas between-genotype
correlations report on horizontal pleiotropy to the extent that they exceed the
corresponding within-genotype correlations. For one set of genotypes, we used 374
progeny of a cross of two natural isolates (54), which enabled not only the estimation of

vertical and horizontal pleiotropy but also the identification of quantitative trait loci
(QTL) with pleiotropic effects. For another set of genotypes, we used a collection of
mutation-accumulation lines, each of which contains a small number of unique
spontaneous mutations (55,56), which enabled a more direct test of the ability of
mutations to alter trait correlations.

The traits we study – morphological features of single cells – represent important
fitness-related traits (51,57,58) that contribute to processes such as cell division and
tissue invasion (*e.g.* cancer metastasis (59)). Cell-morphological features may correlate
across cells for a variety of vertical or horizontal reasons. Vertical reasons include: (1)
inherent geometric constraints (*e.g.* on cell circumference and area); (2) constraints
imposed by gene-regulatory networks (*e.g.* if the genes influencing a group of traits are
all under control of the same transcription factor); and (3) constraints induced by
developmental processes (*e.g.* as a yeast cell divides or "buds", many morphological
features are affected). Horizontal pleiotropy might be evident because genetic variants
each affecting two or more traits (that are otherwise weakly correlated) are segregating in
the progeny of the cross between two natural isolates. Alternatively, horizontal pleiotropy
might be evident because a particular allele strengthens the trait correlation so that
genetic variation affecting one trait is more likely to affect another when that allele is
present. These alternatives can be distinguished by examining trait correlations in two
subsets of progeny strains defined by which natural isolate's allele they possess at a QTL
of interest.

In addition to genetic variation, non-genetic variation may also alter the
correlations between traits. We rely on non-genetic heterogeneity within clonal
populations to serve as perturbations that reveal inherent trait correlations. However, the
correlations themselves might be heterogeneous within these populations. For example,
the dependencies between morphological features may change as cells divide. To control
for this possibility, we performed our trait mapping and subsequent analysis after binning
cells into three stages (unbudded, small-budded and large-budded cells). We further
examined whether trait correlations change across the cell cycle by using a machine-
learning approach to more finely bin the imaged cells into 48 stages of division.
207
Collectively, the results we present here demonstrate that both types of pleiotropy,
vertical and horizontal, are prevalent for single-cell morphological traits, suggesting that
biological systems occupy a middle ground between extreme modularity and extreme
interconnectedness. Perhaps more surprisingly, we find that trait correlations are often
context dependent, and can be altered by mutations as well as cell-cycle state and drug
treatments. The dynamic nature of trait correlations encourages caution when attempting
to quantify and interpret the extent of pleiotropy in nature or when making predictions
about correlated phenotypic responses to the same selection pressure, as is done when
crafting evolutionary traps. However, applying our approach may suggest which trait
correlations are less context dependent and therefore more useful in setting such traps.
218
219
220
221

Results:

QTLs with pleiotropic effects influence yeast single-cell morphology

To detect genes with pleiotropic effects on cell morphology, we measured 167 single-cell morphological features (*e.g.* cell size, bud size, bud angle, distance from nucleus to bud neck; **Table S1**) in 374 yeast strains that were generated in a previous study from a mating between two wild yeast isolates (54,60). These wild isolates, one obtained from soil near an oak tree, the other from a wine barrel, differ by 0.006 SNPs per site (61) and have many heritable differences in single cell morphology (62). For example, we find that yeast cells from the wine strain, on average, are smaller, are rounder, and have larger nuclei during budding than yeast cells from the oak strain (**Fig S1**).

To measure their morphologies, we harvested exponentially growing cells from three replicate cultures of each of these 374 recombinant strains, and imaged on average 800 fixed, stained cells per strain using high-throughput microscopy in a 96-well plate format (**Fig S2**). We used control strains present on each plate to correct for plate-to-plate variation (see *Methods*), and quantified morphological features using CalMorph software (53), which divides cells into three categories based on their progression through the cell cycle (*i.e.* unbudded, small-budded, and large-budded cells) and measures phenotypes specific to each category.

A simple way to measure pleiotropy would be to identify QTL that contribute to variation in these phenotypes, and then to count the number of phenotypes to which each QTL contributes. However, such a measure is sensitive to the statistical thresholds that are used, and therefore risks yielding false inferences about trait modularity. Using a liberal threshold would cause false-positive cases of pleiotropy (less apparent modularity), whereas a conservative threshold would cause failures to detect pleiotropic QTL when they exist (more apparent modularity). Such a measure also assumes the counted traits are somehow independent except for correlations induced by genetic variants. Statistically independent traits could be constructed (and then counted) as principal components of the original traits, but the concern about too-liberal or too-conservative QTL-detection thresholds would remain. Moreover, as we explore extensively below, trait correlations are hierarchical (differing within and between genotypes and conditions), making application of principal components analysis problematic. For these reasons, we do not focus on counting the number of phenotypes influenced by a given locus. Still, to begin to dissect vertical and horizontal pleiotropy we must start with candidate examples of pleiotropic loci.

To detect QTL, we used 225 markers spread throughout the genome (54) and Haley-Knott regression implemented in the R package R/qtl (63,64). We used a standard permutation-based method to estimate statistical significance (63-65), with permutations performed separately for each trait such that the per-trait probability of detecting a false positive is 0.05. With this cutoff, we identified 41 QTL that contribute to variation in 155 of the surveyed morphological features (**Fig 1A**). This approach does not correct for the testing of multiple traits. When we do so using a false-discovery rate set to 5%, results do not change qualitatively. Indeed, the majority of QTL-trait associations that are eliminated using this more-stringent cutoff involve QTL that are detected regardless of this correction (**Fig 1A**; 80% of red points are present at QTL that also possess black

points). This observation suggests many of the associations detected with the less-
stringent cutoff are not spurious. We therefore present subsequent analyses using the
QTL-trait associations based on the less-stringent cutoff, but we also report analyses
using the reduced set of associations based on the more-stringent cutoff to establish that
qualitative conclusions did not change.
Most of the QTL we detect are pleiotropic, meaning each contributes to variation
in more than one morphological feature (**Fig 1A**; 36/41 QTL influence multiple traits,
20/26 after correcting for testing multiple traits). The median number of traits to which
each QTL contributes is 5 (5.5 after correction). This finding provides some support for
the idea that biological systems demonstrate limited pleiotropy, in that the median
number of traits affected per QTL is low (5/167). This median number of traits is similar
to that found in previous high-dimensional QTL screens (17,31), and in analyses of gene
knockouts in yeast and mouse (11). Further evidence that biological systems demonstrate
limited pleiotropy, and thus a modular organization, comes from previous studies of this
mapping family that show these same QTL do not contribute to variation in sporulation
efficiency (60).
However, as noted above, conclusions drawn about modularity from studies that
count traits are subject to criticism: some QTL influences may be too small to detect,
even with less-stringent significance thresholds, creating the appearance of modularity
even if every QTL influences every trait to at least some small degree. Further, we detect
some QTL that influence large numbers of traits, up to 73 (68 after correction). Although
such QTL mitigate to some extent concerns about detection power, they highlight other
potential problems: some QTL might contain multiple genetic differences that impact
different traits, and some morphological features might be inherently correlated and
therefore should not be counted as independent traits. Next, we focus on the first of these
problems by asking whether these QTL represent single genes that contribute to
phenotypic variation in many morphological features. We return to the issue of trait
independence after that.
296
Single genes with pleiotropic effects influence yeast single-cell morphology
When a QTL affects multiple traits, it might not mean that variation in a single
gene is contributing to variation in these traits but instead that linked genes are
contributing to variation in distinct, individual traits. We partitioned genotype-phenotype
associations on the same chromosome into separate QTL when they were greater than 5
cM apart, except in genome regions where many genotype-phenotype associations are
present and there is no clear break point. A distance of 5 cM corresponds on average to 8
protein-coding genes in the yeast genome. The largest QTL we detected spans 17 cM,
which is roughly half the window size utilized in a previous study of this same QTL
mapping family (63). This approach reduces but does not eliminate the possibility that
QTL represent the action of linked loci.
For a small number of QTL with high pleiotropy (highlighted in **Fig 1A**), we
sought to test whether the effects on different morphological features were due to the
action of a single gene. We performed these tests by swapping the parental versions of
candidate genes (*i.e.* we genetically modified the wine strain to carry the oak version of a
given gene, and vice versa). We used the *delitto perfetto* technique to perform these
swaps (66), such that the only difference between a parental genome and the swapped

genome is the coding sequence of the single candidate gene plus up to 1 kb of flanking
sequence (see *Methods*). Candidate genes were selected based on descriptions of the
single-cell morphologies of their knockout mutants (67) and the presence of at least one
non-synonymous amino acid difference between the wine and oak alleles (62).
When a candidate gene contributes to the morphological differences between the
wine and oak parents, we expect yeast strains that differ at only that locus to recapitulate
some of the morphological differences between the wine and oak parents. Indeed, this is
what we observe for *PXL1*, a candidate for the QTL on chromosome 11, and *HOF1*, a
candidate for the QTL on chromosome 13 (**Fig 1B**; compare each plot on the right to the
leftmost plot; see also **Table S2**). This influence is most pervasive for *HOF1*; both the
oak and the wine alleles have a strong effect on the morphology of the opposite parent,
and their effects recapitulate the parental difference to a large extent. The pervasive
influence of *HOF1* on various morphological features is consistent with the fact that this
gene's product affects actin-cable organization and is involved in both polar cell growth
and cytokinesis (68). The effect of *PXL1* on cell morphology is also apparent across
many single-cell features, although only the oak allele has a strong effect that
recapitulates the parental difference. We evaluated *RAS1*, a candidate for the QTL on
chromosome 15, but initial tests indicated that it did not have a significant impact on
most morphological features (**Table S2**). We also attempted to swap alleles for a
candidate gene corresponding to the QTL on chromosome 8, but were unsuccessful (see
Methods).
A previous screen for QTL influencing single-cell morphology in the progeny of a
genetically distinct pair of yeast strains (a different vineyard strain and a laboratory
strain) found some of the same pleiotropic QTL that we detect in the wine and oak cross
(69) (compare their Table 2 to our **Table S1**). In particular, we both find a QTL in the
same position on chromosome 15 that influences many morphological features related to
nucleus size, shape, and position in the cell (**Fig 1A**; orange). We also both detect a QTL
near base pair 100,000 on chromosome 8 that influences cell size and shape (**Fig 1A**;
pink). In the previous screen, the genetic basis of this QTL was shown to be a single
nucleotide change within the *GPA1* gene (69).
The main conclusion from our gene-swapping experiments, which is consistent
with the previous cell-morphology QTL study (69) as well as with comprehensive
surveys of how gene deletions affect the morphology of a laboratory yeast strain (11,49),
is that single genes with pleiotropic effects on cell morphology are readily detected in
budding yeast. Moreover, the morphological traits involved were previously shown to
influence fitness (51,57,58), which raises the question: why do so many genetic analyses
(including ours) detect pleiotropy (5,9-12,14) when other work suggests that pleiotropy
exacts a cost (17,18,20,21)?
352
Dissecting pleiotropy using clonal populations of cells
One hypothesis to explain pervasive pleiotropy may be that the phenotypes we
chose to measure are not independent. Instead, many of these single-cell morphological
features may be inherently related such that perturbing one will have unavoidable
consequences on another and thus any associated limitation of adaptation will be
unavoidable as well. In other words, the hypothesis is that much of the pleiotropy we
observe is vertical pleiotropy. A test of this hypothesis is to ask whether traits that are

jointly affected by the same QTL are correlated in the absence of genetic differences. Our
dataset provides a unique opportunity to perform such a test because we quantified
single-cell traits for, on average, 800 clonal cells per yeast strain (**Fig S2**).
We leverage the hierarchical structure and large sample size of our dataset to
obtain precise estimates of the correlations that exist within and between strains. Thereby
we learn about the underlying relationships between morphological traits, which we use
to distinguish vertical from horizontal pleiotropy. Because we are studying clonal
families without a complicated pedigree structure, these within- and between-strain
correlations are equivalent to the so-called environmental and genetic correlations of
quantitative genetics (70). Here, we use a simple (and fast) method that is appropriate for
two-level hierarchical data to partition the total correlation into a pooled within-strain
component (r_W) and a between-strain component (r_B) (71). One caveat of this correlation-
partitioning approach is that r_B is effectively the correlation between strain means, which
can bias estimates of genetic covariance (70). This bias is most pronounced at small
sample sizes (70), so our large sample sizes allay concern. Nonetheless, for a subset of
traits, we tested whether estimates obtained from correlation partitioning are similar to
those obtained from mixed-effect linear models that specify the variance-covariance
structure of the experimental design. Environmental correlations estimated using both
methods are nearly identical (**Fig S3**). Genetic correlations estimated by correlation
partitioning are sometimes slightly smaller in magnitude than those obtained by linear
modeling (**Fig S3**). This bias is conservative; it may prevent us from identifying cases
where the environmental and genetic correlations significantly differ but will not tend to
create such cases. Despite this reduced power, we rely on the correlation-partitioning
approach, which is substantially faster, because our goal is to estimate environmental and
genetic correlations for thousands of trait pairs.
Unlike the mapping analysis, which considered phenotypes across all three
classes of cell type (unbudded, small-budded and large-budded), this correlation-
partitioning analysis can only be applied to pairs of phenotypes that can be measured in
the same cell. Three of the 36 pleiotropic QTL exclusively affect traits from different cell
types. For example, a QTL on chromosome 4 affects the shape of the nucleus in
unbudded cells as well as in large-budded cells. The correlation between these traits
cannot be partitioned into a within-strain component because these traits are never
measured in the same single cell. Excluding these three QTL leaves 33 pleiotropic QTL.
The 167 single-cell morphological features we measured represent 5645 pairs of
traits (378, 1081, and 4186 pairs of morphological features pertaining to unbudded,
small-budded, and large-budded cells respectively). Because some of these traits are
related, these thousands of trait pairs are not independent. This dependence prevents us
from reliably counting the absolute number of traits that are influenced by vertical vs.
horizontal pleiotropy. Still, partitioning correlations into a non-genetic (r_W) and a genetic
(r_B) component for thousands of trait pairs enables us to: 1) analyze a network describing
the degree to which morphological traits are interconnected or modular, and 2) detect
examples of horizontal pleiotropy if they exist. This approach differs from that of
previous studies of pleiotropy that used principal component analysis (PCA) to
understand which traits are correlated (17). Performing PCA on the individual-cell data is
not the same as controlling for r_W because PCA would ignore the strain groupings, which
can then dominate the analysis (the classic "heterogeneous subgroup" problem in

correlation analysis (72)). As a consequence, PCA can miss cases of horizontal pleiotropy
and obscure, rather than reveal, inherent trait relationships.
408
Inherent relationships between traits contribute to pleiotropy
We focus first on vertical pleiotropy by analyzing correlations that exist in the
absence of any genetic differences (r_W). In the analyses that follow, when we refer to r_W
(or r_B), we mean the magnitude of the correlation, as the sign has no relevance for
arbitrary pairs of traits. The distribution of r_W across traits that are influenced by the same
QTL reflects the degree to which that QTL acts via vertical pleiotropy. The overall
pattern of r_W values (*i.e.*, whether there are isolated clusters of highly correlated traits
versus a densely interconnected network of traits) reflects the modularity of the
underlying biological system. These within-strain correlations are estimated with
extremely high precision because of our large sample size of hundreds of thousands of
clonal cells (800 per each of 374 strains).
Most pairs of single-cell morphological traits are not strongly correlated across
clonal cells (**Fig 2A**). Median r_W is < 0.1, and 74% of pairs have r_W < 0.2. Even if we
allow for nonlinear correlations by transforming data using a nonparametric model that
finds the fixed point of maximal correlation (73), r_W is less than 0.2 for roughly 65% of
pairs. These observations suggest that most of the morphological traits we surveyed are
not inherently related; i.e. for any individual cell, the value of one trait does not predict
well the values of most other traits.
Nonetheless, the distribution of r_W has a prominent right tail (**Fig 2A**) indicating
that some morphological features are strongly correlated across clonal cells. These
correlated features are more likely to be influenced by pleiotropic QTL. Among pairs
represented by this right tail (specifically, those with r_W > 0.2), 75% consist of traits that
share at least one QTL influence; the same is true for only 36% of pairs with r_W < 0.2.
These percentages are similar after changing our QTL detection threshold to correct for
having tested multiple phenotypes (66% and 21%, respectively). Further, the number of
pleiotropic QTL influencing both traits in a pair correlates with that pair's r_W (Pearson's *r*
is 0.52 before correction and 0.54 after). These results suggest that inherent correlations
among morphological features often cause genetic perturbations to one feature to have
consequences on another. In other words, we observe evidence of vertical pleiotropy.
Next, we studied each of the 33 pleiotropic QTL one at a time, asking whether
they influence pairs of traits with higher r_W than expected by chance. Most QTL have a
higher median r_W for the pairs of traits they influence than the median r_W given by all
possible pairs of traits (**Fig 2B**). This difference suggests that vertical pleiotropy drives a
large portion of the pleiotropy we detect.
We also used network analysis to move beyond the pairwise comparisons in **Fig
2A** and ask if morphological traits tend to be clustered into modules. Traits with higher
r_W do indeed tend to group into clusters in networks in which the single-cell
morphological traits are nodes and the r_W magnitudes are edge weights (**Fig 2C**). This
need not have been the case; pairs of traits with high r_W could have been distributed
throughout the network without necessarily being clustered near other high r_W pairs.
Instead, networks representing single-cell morphological features demonstrate more
clustering than do random networks drawn from the same values of r_W (**Fig 2D**; for
corresponding figures from unbudded and small-budded trait networks, see **Fig S4**). This

observation might indicate that morphological phenotypes have a modular organization,
whereby phenotypes within a module exert influence on one another, but exert less
influence on phenotypes from other modules. However, this observation could also result
from human bias when enumerating phenotypes that can be measured, in the sense that
phenotypes that bridge modules might somehow be absent from the data set. The
comprehensive nature of CalMorph diminishes this concern. A related concern is that
apparent modules are formed by trivially related phenotypes, such as the radius and
diameter of a circular object, but we do not find such trivial relationships among the
CalMorph phenotypes. Even a high correlation between the length and area of the
nucleus implies a constraint on nuclear aspect ratio.
462   Our analysis of within-strain trait correlations so far suggests that natural
variation contributing to variation in multiple single-cell morphological features often
acts via vertical pleiotropy. Still, there are hints of another mechanism at play. Some
QTL tend to influence traits that are not clustered in the correlation network (*e.g.* **Fig
2E**). And many pleiotropic QTL influence some pairs of traits with negligible r_W (**Fig
2B**). To investigate how often pleiotropy is not predicted by the degree to which
morphological features correlate in the absence of genetic variation, in the next section
we compare trait correlations present across clones (r_W) to those present between
genetically diverse strains (r_B).
471
472   ***Many traits are more strongly correlated across strains than they are across clones***
473   When genetic changes that perturb one trait have collateral effects on another, we
expect the way traits correlate across genetically diverse strains to reflect trait
correlations across clones (*i.e.* $r_B = r_W$). When this condition is met, pleiotropy can be
viewed as an expected consequence of inherent relationships between traits, *i.e.* vertical
pleiotropy. On the other hand, if a QTL influences two traits that do not correlate across
clones, it may cause these traits to correlate across strains in which this QTL is
segregating. In this case, we expect r_B will be greater than r_W, suggesting horizontal
pleiotropy.
481   After correcting for testing thousands of trait pairs, r_B significantly exceeds r_W in
25% of all trait pairs, and 43% of pairs in which at least one pleiotropic QTL influences
both traits (**Fig 3**; left panel; 43% of points are above the envelope, which represents a
Bonferroni-corrected significance threshold of p < 0.01). This percentage grows to 53%
in the smaller set of QTL that are detected after correcting for testing many traits. In the
majority of cases in which r_B significantly differs from r_W, r_B is greater than r_W (**Fig 3**;
left panel; 78% of points outside the envelope are above it). The magnitude of the
increase in r_B vs. r_W tends to scale with the number of pleiotropic QTL that jointly
influence both traits in a pair (**Fig 3**; left panel; colors get warmer farther above the
envelope). These observations are consistent with the hypothesis that QTL acting via
horizontal pleiotropy increase r_B relative to r_W.
492   However, horizontal pleiotropy is not the only reason traits may correlate
differently across strains versus across clones. We find significant deviations in r_B
relative to r_W in 14% of pairs for which no pleiotropic QTL influence both traits (**Fig 3**;
right panel), or 16% such pairs when using the smaller set of QTL that are detected after
correcting for testing many traits. This observation may suggest the presence of
pleiotropic genetic variants that we did not have statistical power to detect in our QTL

screen. But an alternate explanation for the observed increases in r_B over r_W is that
perhaps we sometimes underestimate r_W.
One reason r_W could be underestimated is that single-cell measurements are
noisier than group-level averages. To test this possibility, we randomly assigned
individual cells to groups (pseudo-strains) having the same number of cells as the actual
strains, and found that in these permuted data, r_B and r_W never significantly differ (**Fig 3**;
insets). Because detection of r_W was not underpowered relative to r_B, we conclude that
measurement noise does not meaningfully obscure r_W. Another reason r_W could be
underestimated is if trait correlations across strains are more linear than those across
clones. To test this possibility, for every pair of traits we transformed the single-cell trait
measurements using a nonparametric model that finds their maximal correlation (73).
This transformation abrogated significant differences in r_B relative to r_W for fewer than
5% of affected trait pairs. Another reason r_W might be less than r_B is if there tends to be
less phenotypic variation within strains than between strains. Contrary to this prediction,
every morphological trait we surveyed varies more within strains than between strains.
Indeed, broad-sense heritability of the traits did not exceed 15% (Fig S1B), reflecting that
within-strain phenotypic variation (*e.g.* variation in a cell's progress through the division
cycle) accounted for at least 85% of the total variation. A final reason r_W could be poorly
estimated is if non-genetic heterogeneity across different subpopulations within clonal
populations causes variation in r_W. Therefore, next we investigated whether the
relationship between single-cell features varies for clonal cells in different stages of the
cell-division cycle.
520
Inferring a cell's progress through division from fixed cell images
Pairs of traits for which r_B is strong whereas r_W is not should reflect horizontal
pleiotropy, but a closer examination of some of these pairs revealed traits that should
correlate due to simple geometric constraints. For example, cell size and the width of the
bud neck should correlate due to the constraint that, even at its maximum, bud neck width
cannot be larger than the diameter of the cell. When measured in small-budded cells,
these two traits are correlated across yeast strains ($r_B = 0.40$) but are significantly less
correlated across clones ($r_W = 0.15$). Given the simple geometric constraint coupling the
width of the bud neck to the cell's size, why is there a discrepancy between r_B and r_W?
We reasoned that this discrepancy exists because the correlation between cell size and
neck width is disrupted during particular moments of cell division; *e.g.* the width of the
bud neck starts small even for large cells (**Fig 4A**; cell micrographs outlined in blue show
two cells in the progress of budding). If the relationship between morphological features
varies during cell division, r_W may represent a poor summary statistic.
How often does the relationship between morphological traits change during cell
division? Our single cell measurements are primed to address this question: we fixed
cells during exponential growth and imaged hundreds of thousands of single cells,
thereby capturing the full spectrum of morphologies as cells divide. A remaining
challenge is sorting these images according to progress through cell division, and then re-
measuring the correlation between morphological features within narrow windows along
that progression.
We performed this sorting using the Wishbone algorithm (74). This algorithm
extracts developmental trajectories from high-dimensional phenotype data (typically

single-cell transcriptome data). We applied Wishbone separately to cells belonging to
each of the three cell types defined by morphometric analysis (unbudded, small-budded,
and large-budded cells). The trends describing how morphological features vary across
Wishbone-defined cell-division trajectories are consistent with previous observations of
how morphology changes as yeast cells divide (75,76) (**Fig 4A**; line plots). For example,
Wishbone sorts fixed-cell images in such a way that cell area increases throughout the
course of cell division (**Fig4A**; upper left panel), and nuclear elongation occurs just
before nuclear division (**Fig4A**; lower left panel). These trajectories also match our own
observations of how morphological features change as live cells divide, which we tracked
by imaging at 1-minute intervals one of the 374 progeny strains that we had engineered to
express a fluorescently tagged nuclear protein (HTB2-GFP) (**Fig 4A**; micrographs). We
chose this particular strain because it does not deviate from the average morphology of all
374 recombinants by more than one standard deviation for any of the phenotypes we
measure.
To further validate Wishbone's performance, we asked whether it could
reconstruct the time series of live-cell images from the HTB2-GFP strain. We obtained
time series for 78 single dividing cells, each imaged over at least 20 timepoints.
Quantifying morphological phenotypes from live-cell images in a high-throughput
fashion proved difficult because the morphometric software was optimized for fixed-cell
images and as cells grow and bud, the cells and their nuclei can move out of the focal
plane. Also, although we used short exposure times when imaging GFP fluorescence,
there are concerns about photo-toxicity and associated growth and morphology defects
(77). For these reasons, we expect Wishbone to perform better on fixed-cell images than
on time series of live cell images. Still, Wishbone's cell-division trajectories recapitulate
the time course. When we align time series data across live cells by centering on each
cell's average predicted progress through division, Spearman's r is 0.65, 0.91, and 0.77
for time series corresponding to each of the three cell types (**Fig 4B**; see **Fig S5** for
recapitulation of 78 individual time series). These correlations are substantially higher
than those obtained by repeating the merging procedure after randomly permuting each
time series (corresponding Spearman's r of 0.42, 0.43. and 0.56). These observations
suggest that Wishbone is effective at properly assigning single-cell images to their
position in the cell cycle.
576
Cell cycle state can influence the relationship between morphological features
To identify cases where significant differences in r_B vs. r_W might result because
r_W is sensitive to cell-cycle state, we first assigned each imaged yeast cell from the QTL-
mapping population to one of 16 equal-sized bins based on Wishbone's estimation of
how far that cell had progressed through division. Because we did this separately for each
of the previously defined cell stages (unbudded, small-budded, and large-budded), this
additional binning finely partitions cell division into 48 (16 x 3) stages. To hold genotype
representation constant across each of the 48 bins, we performed binning separately for
each of the 374 mapping-family strains, then merged like bins across strains. We then
performed correlation partitioning on each bin separately.
Binning cells by cell-cycle state typically decreased the amount of phenotypic
variation per bin, which we expect in turn to obscure the correlation between traits.
Consider an extreme example: if there is no phenotypic variation remaining for a given
trait, it cannot covary with any other traits. Indeed, for most pairs of traits, the binning

procedure either decreases r_W or does not have a dramatic effect on it; decreases in r_W are
especially evident for trait pairs where variation of at least one of the traits shows a
relatively large decrease upon binning (**Fig 4C**). However, for some pairs of traits,
despite the decrease in phenotypic variation for at least one trait, the correlation between
traits improves upon binning. For example, binning by cell division increases the
correlation between cell size and the width of the bud neck (**Fig 4D**; leftmost plot) such
that it approaches r_B. This increased correlation is consistent with our hypothesis that the
process of cell division was obscuring the dependency of bud neck width on cell size.
Examining more pairs of traits for which binning tends to increase r_W (**Fig 4C**; red,
orange, and yellow points) reveals additional cases where the process of cell division
decouples traits that are otherwise correlated, and where binning reveals the underlying
correlation (**Fig 4D**; leftmost three plots).
603   Despite the evidence that cell asynchrony alters some trait correlations, many
cases remain where heterogeneity in cell-cycle state does not explain the observed
discrepancy between r_W and r_B (**Fig 4D**; rightmost three plots). We previously
demonstrated that r_B significantly exceeds r_W in 24% of all trait pairs (1389/5645) (**Fig
3**)**.** For almost half of these pairs (689 pairs), binning by cell division does not resolve the
discrepancy between r_B and r_W to any extent; in other words, r_W does not increase in any
of the 16 bins. For an additional 193 pairs, binning by cell division resolves the
discrepancy by at most 5% in any bin. These results imply that cell-cycle heterogeneity
does not cause the discrepancy between r_W and r_B in the majority of cases, suggesting
that the elevation of r_B over r_W could be explained by QTL demonstrating horizontal
pleiotropy.
614
615   ***Many QTL demonstrate horizontal pleiotropy***
616   To test horizontal pleiotropy further, we asked whether pleiotropic QTL cause
increases in r_B relative to r_W. Not all pleiotropic QTL affect pairs of traits for which r_B is
significantly greater than r_W, so we focus this analysis on the 27 out of 33 QTL that do.
We divided our yeast strains into sets in which a given QTL is not segregating, then re-
measured the difference between r_B and r_W. More specifically, for each QTL, we split the
374 phenotyped yeast strains into two groups based on whether they inherited the wine or
the oak parent's allele at the genotyped marker closest to the estimated QTL location.
Then we repeated correlation partitioning on each subset of strains and compared the
results to those obtained from the complete set. For each QTL, we focused on trait pairs
in which: (1) both traits are affected by this QTL, and (2) r_B is significantly greater than
r_W. Across all such pairs, median r_B tends to decrease upon eliminating allelic variation at
the marker nearest the QTL (**Fig 5A**). No similar reduction in r_B is observed when we
focus on pairs of traits that are not affected by each QTL (**Fig 5A**) and no similar
reduction is observed in r_W (median reduction in r_W is 0.0001).
630   There appear to be two ways in which a QTL may affect r_B. In some cases,
eliminating genetic variation at the marker nearest a QTL decreases r_B in both resulting
subpopulations. Such cases are consistent with a straightforward scenario in which
horizontal pleiotropy results when a QTL that influences two or more traits (that are
otherwise weakly correlated) is segregating in a population (**Fig 5B**; top row shows that
the correlation is strongest in the mixed population where both oak and wine alleles are
segregating). In other cases, eliminating allelic variation at a QTL site decreases r_B in

only one of the two resulting subpopulations (*i.e.* the subpopulation possessing either the
oak or the wine allele). This observation demonstrates that horizontal pleiotropy can
emerge by virtue of a QTL allele strengthening a correlation between two traits so that
genetic variation affecting one trait is more likely to affect the other when that allele is
present (78,79) (**Fig 5B;** bottom row).
642   How many cases where r_B significantly exceeds r_W can be explained, to some
extent, by horizontal pleiotropy (*i.e.* a QTL increasing the between-genotype
correlation)? For every trait pair where r_B significantly exceeds r_W and at least one QTL
influences both traits in the pair (1108 pairs total), eliminating allelic variation at the
marker nearest at least one of the shared QTL causes r_B to decrease in one or both of the
resulting subpopulations (**Fig 5C**: solid black line in rightmost plot). About 60% of these
decreases affect both subpopulations (*e.g.* **Fig 5B**; top row) and 40% affect only one
subpopulation (*e.g.* **Fig 5B**; bottom row). These decreases in r_B appear to resolve the
discrepancies in r_B vs. r_W more often and to a greater extent than does accounting for
cell-cycle heterogeneity (**Fig 5C**; leftmost plot). Some QTL have larger impacts on r_B
than do others (**Fig 5C**). Eliminating allelic variation near a QTL on chromosome 13
decreases r_B in the largest number of traits pairs (658). Subtracting the influence of a
QTL on chromosome 15 decreases r_B to the greatest extent; the average decrease across
342 affected trait pairs is 0.07.
656   One caveat that remains is whether these examples of pleiotropy represent single
genetic changes that influence multiple traits, or the presence of multiple nearby genetic
variants within each QTL. Our finding that there are two types of horizontal pleiotropy
(**Fig 5B**), provides some insight. The first type of horizontal pleiotropy (**Fig 5B;** upper
row) may result from the presence of multiple genetic variants segregating together
because recombination has not broken them apart. However, the second type of
horizontal pleiotropy (**Fig 5B**; lower row) cannot be explained in the same way because a
strong trait correlation exists in the absence of allelic variation at that QTL (blue points).
Therefore, although multiple closely linked variants might underlie the difference in the
trait correlation between the oak and wine alleles of the QTL, they would act at the level
of the correlation rather than of individual traits.
667
668   ***Spontaneous mutations alter the relationships between morphological features***
669   Our finding that some QTL alleles appear to strengthen correlations between
otherwise weakly correlated traits (**Fig 5B**; lower panel) lends credence to the idea that
the relationships between phenotypes, and thus the extent of phenotypic modularity (or
integration), are mutable traits (80). This finding has implications for evolutionary
medicine, in particular evolutionary traps, *e.g.* strategies to contain microbial populations
by encouraging them to evolve resistance to one treatment so that they become
susceptible to another (40-42). These traps will fail if targeted correlations can be broken
by mutations. To test whether spontaneous mutations can alter trait correlations, we
analyzed the cell-morphology phenotypes of a collection of yeast mutation-accumulation
(MA) lines (55). These MA lines were derived from repeated passaging through
bottlenecks, which dramatically reduced the efficiency of selection and thereby allowed
retention of the natural spectrum of mutations irrespective of effect on fitness (56). We
previously imaged these lines in high throughput (>1000 clonal cells imaged per each of
94 lines) (51).

Because MA lines contain private mutations unique to each strain, they are not
amenable to QTL mapping and between-strain trait correlations have less meaning.
Instead, we focused on within-strain correlations, which we expected to be consistent
across strains because of the limited number of mutations distinguishing the strains (an
average of 4 single-nucleotide mutations per line (56)), except if a rare mutation does
indeed alter the correlation. To determine if such correlation-altering mutations exist, we
calculated within-strain correlations for each strain separately and asked, for each trait
pair, whether any strains had extreme correlations relative to the other strains. For most
trait pairs, the MA lines trait correlations did not vary much from each other or from that
of the ancestor strain (**Fig 6**). However, in several instances, we observed a trait-pair
correlation dramatically outside the range of the other trait pairs and more than four
standard deviations from the mean (**Fig 6A**). Some mutations appear to influence many
trait-trait relationships (mutations found in blue- and purple-colored strains in **Fig 6B &**
C), whereas others influence fewer (mutations found in magenta-colored strain in **Fig**
6C). Mutations that alter trait correlations are not necessarily the result of rare events
such as aneuploidies or copy number variants; all five strains highlighted in **Fig 6** do not
possess these types of mutations and instead possess at least one single nucleotide
mutation (56).
Given that in the small sampling of spontaneous mutations captured by the MA
strain collection, we found several that appear to alter the relationship between
morphological features, we think such mutations are common enough to merit further
consideration in evolutionary models. The mutations in the outlier lines provide candidate
correlation-altering mutations for future mechanistic studies as well.
706
Different environments alter the relationships between morphological features
We have used non-genetic heterogeneity within clonal populations to uncover
inherent trait correlations. One might consider achieving the same aim by using instead
the non-genetic perturbations represented by different environmental treatments.
However, our results suggest that trait correlations can be highly context dependent,
changing across cell cycle state (**Fig 4**) and genetic background (**Fig 5B lower panel &**
Fig 6). If trait correlations change across environments, then the intricacies of the
environment-specific effects would need to be incorporated into any inferences about
vertical and horizontal pleiotropy, adding a complicating dimension to the analysis.
To investigate the potential utility of across-environment trait correlations for
distinguishing horizontal from vertical pleiotropy, and to further explore the context
dependence of trait correlations, we analyzed trait correlations across a range of
concentrations of the Hsp90-inhibiting drug geldanamycin (GdA). We showed previously
that GdA affects cell morphology (51), so it presents an opportunity to analyze how
correlations among these traits vary across environments. We performed this analysis
using a subset of the yeast strains from our QTL mapping family (**Fig S6**), partitioning
trait correlations into a pooled within-strain component (r_W) and a between-strain
component (r_B).
GdA alters the correlations between morphological traits. The impact of GdA on
r_W increases with the concentration of GdA (**Fig 7**), suggesting that more extreme
environmental differences are more likely to result in changes in r_W. We conclude that
looking across diverse environments is not a good way to understand the inherent

relationships between traits that exist in a single environment. Indeed, previous studies of
pleiotropy have treated growth parameters in different environments as different traits
(81) rather than as a way to estimate inherent trait correlations.
732
Discussion:
Although evolutionary biologists and medical geneticists alike appreciate that
organismal traits can rarely be understood in isolation, the extent and implications of
pleiotropy have remained difficult to assess. A common approach to measuring
pleiotropy has been to count phenotypes influenced by individual genetic loci (18,34,35).
For example, the median number of skeletal traits affected per QTL in a mouse cross was
six (out of 70 traits measured); this small median fraction of traits suggests that variation
in skeletal morphology is modular (17,31). Of course, for a count of traits to be
meaningful the full trait list must be comprehensive, and correlations between traits must
be properly accounted for (18,34,35). We aimed for comprehensiveness in a very similar
way to the studies of mouse skeletal traits, by systematic phenotyping of a large number
of morphological traits. However, we addressed the need for a principled approach to
separating inherent trait correlations from those induced by genetic differences in a new
way: by extending the analysis to include within-genotype correlations and thereby
enabling an operational definition of the distinction between vertical and horizontal
pleiotropy.
Our comprehensive analysis of how thousands of trait pairs co-vary within and
between mapping strains yields an unprecedently quantitative and nuanced view of
pleiotropy. We found support for modularity, not only in the low median number of traits
affected per QTL (five out of 167), but also in the way that within-genotype correlations
grouped traits into relatively isolated clusters (**Fig 2**). We also found ample evidence of
horizontal pleiotropy layered on top of that modularity, with many cases of between-
genotype trait correlations that exceeded within-genotype correlations (**Fig 3**).
Our results do not speak directly to whether modularity results from selection
against pleiotropy in nature because we survey only two natural genetic backgrounds
(wine and oak). In other words, the presence of modularity is not necessarily evidence
that it is adaptive or that it is maintained by natural selection. Future work comparing MA
lines to a larger collection of natural isolates might help answer questions about the
extent to which selection purges pleiotropic mutations.
Our partitioning of between-strain (genetic) and within-strain (environmental)
correlations relates to another approach to understanding trait interdependencies, the
estimation of the so-called **G** matrix. This genetic variance-covariance matrix
summarizes the joint pattern of heritable variation in a population of the traits that
compose its rows and columns, and is central to understanding how trait correlations
constrain evolution. The **G** matrix arises in the multivariate breeder's equation, which
describes the responses to selection of correlated traits (82). If breeding is the goal, the
distinction between vertical and horizontal pleiotropy is not so important, because both
can impede selection. Indeed, any philosophical concern about what constitutes a
biologically meaningful trait is irrelevant to the breeder, who actually cares about
particular traits (*e.g.*, milk yield and fat content).
G matrices are not only relevant to breeders, but to evolutionary biologists as
well, and it is worthwhile to place our results into this context. A major evolutionary

question in the **G**-matrix literature is whether the **G** matrix itself can evolve. In other
words, do short-term responses to selection (as captured in the breeder's equation) predict
long-term responses or do constraints shift through time, perhaps in a way that facilitates
(or is part of) adaptation (83)? Our results with MA lines add to evidence that the **G**
matrix readily changes (84), in that individual mutations have major effects on particular
trait correlations (*e.g.* **Fig 6A**). Our QTL-mapping results also support this view, in that
some cases of horizontal pleiotropy appear to be caused by alleles that alter trait
correlations (*e.g.* **Fig 5B**; bottom panel).
Another prominent question in the **G**-matrix literature is the extent to which the **P**
matrix, which includes all sources of phenotypic variation and covariation, predicts the **G**
matrix, which only includes additive genetic effects (*i.e.*, those that respond to selection).
If **P** predicts **G** well, as proposed by Cheverud (85), then inference of selection responses
from patterns of trait covariation in a population would suffice when genetic analysis
would be difficult or costly. Our results do not speak directly to this question, because we
did not estimate **G** itself and instead estimated genetic correlations that include non-
additive effects. However, our results are informative from another angle, which is the
comparison of genetic and environmental correlations. As we showed (**Fig 3**), although
there are cases in which the environmental and genetic correlations have different signs,
the environmental correlations do tend to match the signs of the genetic correlations and
predict their magnitudes to some extent as well, consistent with similarity between **P** and
G. Future experiments using clones embedded in a more complicated crossing scheme
could properly partition **P** into **G**, **E**, and the non-additive genetic components, to address
Chevrud's conjecture (85) more directly. There are only a few reports of comparisons of
E matrices (86), but we encourage increased attention to the **E** matrix to understand
inherent trait correlations and to contextualize **G** in a way that diminishes concerns about
which traits are biologically meaningful and therefore merit status as the matrix's rows
and columns.
A major and unforeseen conclusion of our work is the extent to which context is
crucial. We have shown that trait correlations change through the cell-division cycle, in
different genetic backgrounds, and across a drug gradient. It is likely that
macroenvironmental differences alter trait correlations as well (87). These findings
provide insight as to how biological systems appear to be modular, as evolutionary theory
predicts (24-29), yet in other studies appear to be highly interconnected (13,14). Our
results suggest that biological systems are modular, but that these modules change across
contexts such that the potential phenotypic impacts of a genetic change can be extensive.
These results support the idea that predicting the phenotypic impact of a genetic
change requires a paradigm shift (88,89), away from merely mapping the relationships
between traits and toward unfurling the range of contexts across which those
relationships persist. Future work in this direction will not only advance understanding of
the evolution of complex traits but will have practical benefits. For example, our
approach demonstrates a potentially fruitful way to design evolutionary traps: studying
within-genotype correlations across contexts to identify particularly immutable
correlations between traits.
818
Acknowledgments:

We are grateful to Dmitri Petrov, Grant Kinsler and Michael Lynch for helpful
discussions. We also thank Barak Cohen and David Hall for providing strains used in this
study. This work was supported by National Institutes of Health grant R35GM118170 (to
MLS), National Institutes of Health fellowship F32GM103166 (to KGS), National
Institutes of Health grant R35GM133674 (to KGS), a New York University Graduate
School of Arts and Science Dean's Dissertation Fellowship (to SL), and National
Institutes of Health grant R35GM119744 (to ABP). The funders had no role in study
design, data collection and analysis, decision to publish, or preparation of the manuscript.
828

829   **Data Availability:**
830   All data presented in this study are available at Open Science Framework
831   (DOI 10.17605/OSF.IO/B7NY5).
832

Materials and Methods:
Measuring the morphology of single yeast cells
Recombinant yeast strains were generated from a cross between the oak parent (BC233:
SPS2:EGFP:kanMX4/SPS2:EGFP:kanMX4) and the wine parent (BC240:
SPS2:EGFP:natMX4/SPS2:EGFP:natMX4) then genotyped at 225 markers in a previous
study (54,60); each resulting recombinant strain is a homozygous diploid. We prepared
yeast cells from these strains for microscopy using published methods (50-52,90).
Briefly, yeast strains were grown in minimal media with 0.08% glucose in 96-well plates
(91), harvested during exponential phase, fixed in 4% paraformaldehyde, stained for cell-
surface manno-protein (with FITC-concanavalin A) and nuclear DNA (with DAPI),
sonicated, mounted on 96-well glass-bottom microscopy plates, and imaged with a Nikon
Eclipse TE-2000E epifluorescence automated microscope using a 40× objective and
appropriate fluorescence filters. Three biological replicate experiments were performed,
typically yielding a total of between 500 to 1,000 imaged cells per strain (**Fig S2**).
847

Statistical analysis and processing of cell image data
Cell image processing was performed similarly to previous studies (50-52,90). Imaged
cells were analyzed for quantitative morphological traits using the CalMorph software
package (53), which reports on hundreds of morphological features that are each specific
to one of three cell types: unbudded, small-budded, and large-budded cells. We excluded
phenotypes for which >10% of cells had missing values, leaving 167 morphological
features. Any cell that was not scored for all features pertaining to its type was
eliminated. Each morphological trait was transformed via a Box-Cox transformation of
the raw data with the value of lambda that makes the residuals of a linear regression of
phenotype on strain most normal using the EnvStats package in R (92). Internal controls
(several wells representing the wine and oak parents) were present on every 96-well plate
and were used to correct for effects on phenotypic variation that resulted from differences
among replicate experiments, such as differences in the brightness of the cell stain. We
calculated the mid-parent value for each phenotype on every plate, then calculated the
average mid-parent value across all plates. For each phenotype, we found the difference
between the plate-specific mid-parent value and the average mid-parent value across all
plates. Then we subtracted this difference from each plate for the corresponding
phenotype. After correction, any cell with a morphological feature that deviated from the

average by more than 5 standard deviations was then eliminated, as investigation of such
cells typically revealed these were CalMorph miscalls or cellular debris.
868
QTL mapping
QTL interval mapping was performed similarly to previous studies (63) using the R/qtl
package (64). We performed a QTL scan using the function "scanone", which finds at
most one QTL per chromosome, followed by the "scantwo" function which allowed us to
identify potential second additive QTL per chromosome. The yeast strains, which are
homozygous diploids, were modeled as doubled haploids and QTL models were fit using
Haley-Knott regression. When comparing QTL across traits, QTL greater than 5 cM apart
on the same chromosome were counted as separate QTL. We estimate that a region of 5
cM contains on average 8 genes, since there are 6,746 genes in the yeast genome (67),
and the map length we calculated using R/qtl is 4076 cM. In some cases, we detected a
QTL in between two others on the same chromosome and within 5 cM of both. In these
cases, there were typically many QTL found within a narrow region without any gaps of
greater than 2 cM. We counted these as single QTL that affect many traits. Using this
method, the largest QTL we detect spans 17 cM. A summary of all significant QTL
effects, including their chromosomal locations in cM and which QTL on the same
chromosome we considered unique, is provided in **Table S1** (also see **Fig 1A**).
885
To determine significance thresholds, we employed the method of (65) as implemented in
R/qtl (64). QTL were assigned a p-value based on a trait-specific empirical distribution of
genome-wide LOD score maximums from 10,000 (1,000 for the two dimensional scan)
randomly permuted datasets. We used a p-value cutoff of 0.05 to determine significant
QTL. We also obtained a more stringent set of significant QTLs by correcting for the
testing of multiple traits by controlling the false discovery rate across phenotypes. For
each trait we took the position and p-value of the maximum LOD score on every
chromosome. We calculated qvalues for this set of loci using the R 'qvalue' package (93)
and used a 0.05 q-value threshold to call significant QTL.
895
Candidate gene swaps
All yeast transformations were performed using the lithium acetate (94) and *delitto*
perfetto (66) methods. For each candidate gene, the gene was first deleted from haploid
variants of both the wine and oak parental strains and replaced with a selectable marker,
the yeast gene encoding orotidine-5'-phosphate decarboxylase (*URA3*). Gene knockouts
were confirmed by growth on plates lacking uracil and DNA sequencing of the affected
region. Next, the *URA3* selectable marker was replaced with the other parent's version of
the candidate gene. These candidate gene 'swaps' were selected by growth on 5-
Fluoroorotic acid and confirmed by sequencing of the affected region. For each candidate
gene, we swapped a region containing the coding sequence plus 5 – 750 bp up and
downstream. We used the following regions of homology to define the boundaries of
each swapped segment:
908
~300bp upstream of *PXL1*: TTATAATTGTGGTTTAGCGTTTCATAGTCGC
~300bp downstream of *PXL1*: CCTTATTCTCTATTCTTAGGCTCCTGTTCC
~5bp upstream of *HOF1*: GAAAGAATGAGCTACAGTTATGAAGCTTG

~ 300bp downstream of *HOF1*: GTATTCGTAACAAGTGACTCTAATGATAT
~ 750bp upstream of *RAS1*: CGACTAAAGGAATTATACCATCATGCATC
~ 300bp downstream of *RAS1*: GCATTTCTAAAAACAGAGCTTTGCCG
915
These regions of homology were chosen by searching for regions of higher GC content
nearby the start and end of each gene's coding sequence. In addition, we attempted to
swap the wine and oak parents' versions of the *GPA1* gene on chromosome 8. Despite
trying various regions of homology, we could not successfully replace *GPA1* with the
URA3 selectable marker in the oak parent. *GPA1* is known to be essential in some genetic
backgrounds (95).
922
Though the recombinant strains we studied are homothallic diploids, the strains in **Fig 1B**
(both the parental strains and the strains possessing the gene swaps) are haploid. Because
the analyses in **Fig 1B** compare pairs of strains (*e.g.* the oak haploid parent to the wine
haploid parent, or the wine haploid parent to the wine haploid parent possessing the oak
allele of *PXL1*), we only considered experiments where both strains in the pair were
imaged in the same replicate experiment. To account for differences among replicate
experiments, for each phenotype, we subtracted the value in one strain from the value in
the other to calculate the phenotypic difference between strains in that replicate
experiment; the reported value is the average of these differences across replicate
experiments (**Table S2, Fig 1B**).
933
934   *Calculation of correlation coefficients*
935   We used WABA II as implemented in the multilevel package in R (71) to
calculate cell-level (r_W) and strain-level (r_B) Pearson correlation coefficients for each pair
of traits. We used an r-to-z transformation to determine whether differences in r_B vs. r_W
are significant at a Bonferroni corrected p-value of 0.01 (this is a z-score cutoff of 4.63,
given 5645 pairs of traits were tested). To assess whether correlations across single cells
generally result in different values than correlations across group-level averages, we
assigned yeast cells to groups (pseudo-strains) randomly, maintaining the same number
of cells per strain as in the actual data. To assess whether results would differ if we
allowed for non-linear correlations, we transformed the single-cell data using a
nonparametric model that finds the fixed point of maximal correlation, implemented in
the R package acepack (73). To assess whether results from WABA differed from those
obtained using a standard quantitative genetics model (**Fig S3**), we implemented the latter
using the nlme package in R (96) to specify a mixed-effects model with cells nested
within strains. We specified a covariance structure that allows covariance between two
traits but no covariance between cells or between strains. We used this model to calculate
the environmental and genetic correlations for 350 pairs of randomly chosen traits.
951
Live imaging single cells as they divide
For live imaging the morphology of dividing yeast cells, we chose one of the
recombinant yeast strains, F2_292. This strain was chosen because it does not deviate
from the average morphology of all 374 recombinants by more than one standard
deviation for any of the phenotypes we measured. F2_292 was transformed to express a
fusion protein of GFP and a nuclear protein (histone H2B encoded by *HTB2*). Two

independent transformants were imaged in the GFP channel (for nuclei) and in brightfield
(for cell outlines). We prepared live cells for imaging following published methods
(91,97,98), in a similar way to that described above, except cells were neither fixed nor
stained. Cells were taken during mid-log phase growth, seeded in 96-well glass bottom
microscopy plates containing minimal media with 0.08% glucose, and imaged over a
period of 3 hours. In each of four replicate experiments, cells were imaged either every
minute, every 90 seconds, or every 2 minutes. We used short exposure times (afforded by
the highly abundant HTB2-GFP) and took only a single image per well per timepoint to
reduce photo-toxicity. We processed images with CalMorph then matched cells across
timepoints by their centroid locations in the imaging fields. Overall we obtained time
series for 78 cells that each: (1) were longer than 20 timepoints, (2) contained no gaps
where the cell was not phenotyped for many consecutive timepoints, and (3) contained no
images that appeared to be very out of focus potentially resulting in misestimation of
phenotype values. Because CalMorph divides cells into unbudded, small-budded and
large-budded stages, these 78 time series are also divided this way (11, 23, and 44 cells,
respectively).
974
We used the Wishbone algorithm implemented in python (74) to estimate progression
through the cell-division cycle. Wishbone recapitulates each of these 78 time series (**Fig
S5**) with Spearman correlations between the actual and inferred image orders that average
0.42, 0.85, 0.40 across all unbudded, small-budded or large-budded series, respectively.
The lower correlations between Wishbone's predicted progress through division and time
for the unbudded and large-budded cells may result because each time series captured
only a part of the cell-division cycle and, during some stretches in the cycle, there are
fewer morphological changes taking place. To estimate Wishbone's accuracy across a
longer stretch of time, we merged the Wishbone predictions within the classes of
unbudded, small-budded or large-budded cell time series. To do so, we had to contend
with the fact that the first timepoint for each imaged cell often represents a different
moment in division. For example, some time series for unbudded cells start from an
image that is already far along the division process (**Fig S5**; values close to 1 on the
vertical axis) while others start from a cell image that has just begun its division cycle
(**Fig S5**; values close to zero on the vertical axis). Therefore, we aligned the time series
by subtracting from each the difference between Wishbone's estimate of the average
percent progress through division and the average time elapsed.
992
Note that, because this merging procedure utilized information from Wishbone, it
imposes a correlation between time and Wishbone's estimated progress through division.
To reduce the impact of this induced correlation, we eliminated the cell images in the
middle of each time series, which represent the images that are most affected by this
induced correlation. Eliminating 25% or 50% of cell images in this way reduced the
correlations by at most 0.05, suggesting these correlations are not driven by our merging
procedure.
1000
Assigning cells to a bin based on progression through cell division
We used Wishbone to estimate how far each fixed-cell image had progressed through cell
division. Wishbone software requires input about which "start" cell has features

resembling those present at the start of the cell cycle. To identify such features, we used
the data from the live-imaged cell time series. We plotted how single-cell features change
over the course of live imaging, and chose several features that correlate best with
progress through cell division (*e.g.* cell size, bud size, location of the nucleus). Complete
datasets provided at Open Science Framework (DOI 10.17605/OSF.IO/B7NY5) include
information on which fixed-cell image was chosen as the start cell.
1010
Using Wishbone's estimation of how far each fixed cell had progressed through division,
we assigned each cell to one of 16 equal-sized bins. We did this separately for each of the
374 yeast strains, then merged like bins across strains, such that genetic diversity was
constant across each of the final 16 bins. We obtained very similar results to those
reported in **Figs 4C**, **4D**, and **5C** when we used 8 instead of 16 bins. The names of the
traits plotted in **Fig 4** represent succinct summaries of single-cell morphologies
quantified using CalMorph (53). For fuller descriptions of these traits, see the following
trait designations in the CalMorph software manual: **Fig 4A** upper left: C11.1 in
unbudded cells, C101 in budded cells; **Fig 4A** lower left: D184 in small-budded cells,
D182 in unbudded and large-budded cells; **Fig 4A** upper right: C12.2; **Fig 4A** lower
right: D116; **Fig 4C** upper left: C101 and C109 in small-budded cells; **Fig 4C** upper
middle: C11.2 and D132 in small-budded cells; **Fig 4C** upper right: C105 and C113 in
small-budded cells; **Fig 4C** lower left: C114 and D145 in large-budded cells; **Fig 4C**
lower middle: C109 and C126 in large-budded cells; **Fig 4C** lower right: D14.2 and D169
in large-budded cells.
1026
Eliminating genetic variation at the marker nearest a QTL
For each of the 27 QTL suspected of horizontal pleiotropy (*i.e.* pleiotropic QTL that
influence at least one pair of traits for which r_B significantly exceeds r_W), we divided the
374 phenotyped yeast strains into two groups based on whether they inherited the wine or
the oak parent's allele at the genotyped marker closest to the QTL. In some cases, a QTL
spans multiple markers; for example, a QTL on chromosome 15 that influences 64
morphological features spans 14 cM and 4 markers (**Table S1**). These 64 genotype-
phenotype associations are mainly clustered around the ninth marker on chromosome 15,
though a few are closer to the eighth, tenth, or eleventh. To avoid redundancy, for QTL
spanning multiple markers we study the one that is most represented. After dividing
strains into two groups based on which allele they inherited at that marker, we performed
correlation partitioning separately for each group of strains.
1039
1040 The names of the traits plotted in **Fig 5B** represent succinct summaries of single-cell
1041 morphologies quantified using CalMorph. For fuller descriptions of these traits, see the
1042 following trait designations in the CalMorph software manual: upper: D128 and C114 in
1043 large-budded cells; lower: D197 and D17.1 in large-budded cells.
1044
Quantifying trait correlations within each MA line
Mutation accumulation occurred in a diploid laboratory yeast strain with genotype *ade2,*
lys2-801, his3-ΔD200, leu2–3.112, ura3–52 (55). Resulting diploid MA lines were
sporulated to create haploids, which were sequenced in a previous study (56). We
previously imaged these haploid lines in high throughput (>1000 clonal cells imaged per

each of 94 lines) (51). Fewer morphological traits were analyzed in that study than in the
current study, such that there were only 3731 pairs of traits to survey, as opposed to 5645
in the QTL-mapping family. We calculated Pearson correlations between every pair of
traits, separately within each MA line. The names of the traits plotted in **Fig 6A** represent
succinct summaries of single-cell morphologies quantified using CalMorph. For fuller
descriptions of these traits, see the following trait designations in the CalMorph software
manual: upper left: D185 and D186 in large-budded cells; upper right: C102 and D132 in
small-budded cells; lower left: C108 and D167 in large-budded cells; lower right: D135
and D169 in large-budded cells.
1059
Quantifying trait correlations across drug concentrations
For this analysis, we imaged the single-cell morphologies of 78 of the 374 strains that
comprised our QTL mapping family. We chose these strains because they were stored
together on a single 96-well plate (the rest of the 96 wells represent blanks or internal
controls), removing concerns about batch effects. We imaged these strains after
exponential growth in three concentrations of geldanamycin (GdA) (8.5 µM, 25 µM and
100 µM). We chose these concentrations because of their wide-ranging impacts on cell
growth rate (51). We obtained single-cell morphology measurements for cells grown in
the lowest concentrations of GdA from our previous study (51) and collected data for
cells grown in higher concentrations following the procedures outlined in that study,
which was very similar to those outlined above, but with a control for the solvent in
which GdA is dissolved. Specifically, cells exposed to GdA were compared to cells
imaged in identical conditions (containing the same concentration of the solvent DMSO)
but lacking GdA. GdA+/– paired experiments are performed side-by-side, with cells
grown in each condition being imaged in adjacent wells on a 384-well microscopy plate.
1075
Resulting morphological data were analyzed following similar procedures as described
above. Briefly, each trait was transformed via a Box-Cox transformation of the raw data
based on the residuals of a linear model with strain, environment, and replicate as effects.
Two replicates were performed for both the 8.5 and 100 µM environments, and a single
replicate for the 25 µM environment. Internal controls (wells representing the wine and
oak parents) were used to correct for effects on phenotypic variation that resulted from
differences among replicate experiments. We used WABA II to calculate cell-level (r_W)
correlation coefficients in each of the three GdA concentrations, as well as the
corresponding three control conditions. To calculate the impact of GdA on r_W, we
compared r_W in each drug vs. control condition.
1086
1087

Author Contributions:

KGS: Conceptualization, Funding acquisition, Formal analysis, Investigation, Writing – original
draft, Writing – review & editing
SL: Investigation, Writing – review & editing
CL: Conceptualization, Formal analysis, Writing – review & editing
AT: Investigation, Writing – review & editing
NZ: Formal analysis, Writing – review & editing
CR: Investigation, Writing – review & editing
AP: Conceptualization, Writing – review & editing

MLS: Conceptualization, Funding acquisition, Formal analysis, Writing – original draft, Writing –
review & editing
1099
1100
1101

1102 **References:**
1103

1.   Plate L. Genetics and evolution. In: Festschrift zum sechzigsten Geburtstag
Richard Hertwigs. Jena: Verlag von Gustav Fischer; 1910. pp. 536–610.

2.   Stearns FW. One Hundred Years of Pleiotropy: A Retrospective. Genetics;
2010 Nov 1;186(3):767–73.

3.   Richardson TG, Harrison S, Hemani G, Smith GD. An atlas of polygenic risk
score associations to highlight putative causal relationships across the human
phenome. Elife. 2019;8:e43657.

4.   Simons YB, Bullaughey K, Hudson RR, Sella G. A population genetic
interpretation of GWAS findings for human quantitative traits. PLoS Biol.
2018 Mar;16(3):e2002985.

5.   Tyler AL, Asselbergs FW, Williams SM, Moore JH. Shadows of complexity:
what biological networks reveal about epistasis and pleiotropy. Bioessays.
2009 Feb 1;31(2):220–7.

6.   Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, et al.
10 Years of GWAS Discovery: Biology, Function, and Translation. Am J
Hum Genet. 2017 Jul 6;101(1):5–22.

7.   Visscher PM, Yang J. A plethora of pleiotropy across complex traits. Nat
Genet. 2016 Jun 28;48(7):707–8.

8.   Chesmore K, Bartlett J, Williams SM. The ubiquity of pleiotropy in human
disease. Hum Genet. 137(1):39–44.

9.   Sivakumaran S, Agakov F, Theodoratou E, Prendergast JG, Zgaga L,
Manolio T, et al. Abundant pleiotropy in human complex diseases and traits.
Am J Hum Genet. 2011 Nov 11;89(5):607–18.

10.   White JK, Gerdin A-K, Karp NA, Ryder E, Buljan M, Bussell JN, et al.
Genome-wide generation and systematic phenotyping of knockout mice
reveals new roles for many genes. Cell. 2013 Jul 18;154(2):452–64.

11.   Wang Z, Liao B-Y, Zhang J. Genomic patterns of pleiotropy and the
evolution of complexity. Proc Natl Acad Sci USA. 2010 Oct
19;107(42):18034–9.

12.   Housman G, Byler S, Heerboth S, Lapinska K, Longacre M, Snyder N, et al.
Drug resistance in cancer: an overview. Cancers (Basel). 2014 Sep
5;6(3):1769–92.

13.   Rockman MV. The QTN program and the alleles that matter for evolution: all
that's gold does not glitter. Evolution. 2012 Jan;66(1):1–17.

14.   Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. Cell. 2017 Jun;169(7):1177–86.

15.   Fisher RA. The Genetical Theory of Selection. Oxford: Claredon; 1930.

16.   Orr HA. Adaptation and the cost of complexity. Evolution. 2000;54(1):13.

17.   Wagner GP, Kenney-Hunt JP, Pavlicev M, Peck JR, Waxman D, Cheverud
JM. Pleiotropic scaling of gene effects and the 'cost of complexity'. Nature.
2008 Mar 27;452(7186):470–2.

18.   Wagner GP, Zhang J. The pleiotropic structure of the genotype–phenotype
map: the evolvability of complex organisms. Nat Rev Genet. 2011 Mar
1;12(3):204–13.

19.   McGee LW, Sackman AM, Morrison AJ, Pierce J, Anisman J, Rokyta DR.
Synergistic Pleiotropy Overrides the Costs of Complexity in Viral
Adaptation. Genetics. 2016 Jan;202(1):285–95.

20.   Wittkopp PJ, Haerum BK, Clark AG. Regulatory changes underlying
expression differences within and between Drosophila species. Nat Genet.
2008 Mar 1;40(3):346–50.

21.   Stern DL. Perspective: evolutionary developmental biology and the problem
of variation. Evolution. 2009 Jan 20;54(4):1079.

22.   He X, Zhang J. Toward a Molecular Understanding of Pleiotropy. Genetics.
2006 Aug 1;173(4):1885–91.

23.   Papakostas S, Vøllestad LA, Bruneaux M, Aykanat T, Vanoverbeke J, Ning
M, et al. Gene pleiotropy constrains gene expression changes in fish adapted
to different thermal conditions. Nat Commun. 2014 Jun 3;5:4071.

24.   Altenberg L. Modularity in Evolution: Some Low-Level Questions. In:
Callebaut W, Rasskin-Gutman D, editors. Modularity: Understanding the
Development and Evolution of Natural Complex Systems. Cambridge: MIT
Press; 2005. pp. 99–128.

25.   Wagner GP, Altenberg L. Perspective: Complex Adaptations and the
Evolution of Evolvability. Evolution. 1996 Jun;50(3):967.

26.   Welch JJ, Waxman D. Modularity and the cost of complexity. Evolution.
1168   2003 Aug;57(8):1723–34.

1169   27.   Wagner GP, Pavlicev M, Cheverud JM. The road to modularity. Nat Rev
1170   Genet. 2007 Dec;8(12):921–31.

1171   28.   Melo D, Porto A, Cheverud JM, Marroig G. Modularity: genes, development
1172   and evolution. Annu Rev Ecol Evol Syst. 2016;47(1):463–86.

1173   29.   Collet JM, McGuigan K, Allen SL, Chenoweth SF, Blows MW. Mutational
1174   Pleiotropy and the Strength of Stabilizing Selection Within and Between
1175   Functional Modules of Gene Expression. Genetics. 2018 Apr;208(4):1601–
1176   16.

1177   30.   Paaby AB, Rockman MV. Pleiotropy: what do you mean? Reply to Zhang
1178   and Wagner. Trends Genet. 2013 Jul;29(7):384.

1179   31.   Wagner GP, Zhang J. Universal pleiotropy is not a valid null hypothesis:
1180   reply to Hill and Zhang. Nat Rev Genet. 2012 Apr 1;13(4):296–6.

1181   32.   Hill WG, Zhang X-S. Assessing pleiotropy and its evolutionary
1182   consequences: pleiotropy is not necessarily limited, nor need it hinder the
1183   evolution of complexity. Nat Rev Genet. Nature Publishing Group; 2012 Apr
1184   1;13(4):296.

1185   33.   Zhang J, Wagner GP. On the definition and measurement of pleiotropy.
1186   Trends Genet. 2013 Jul;29(7):383–4.

1187   34.   Gibson G. Decanalization and the origin of complex disease. Nat Rev Genet.
1188   2009 Feb;10(2):134–40.

1189   35.   Nuzhdin SV, Friesen ML, McIntyre LM. Genotype-phenotype mapping in a
1190   post-GWAS world. Trends Genet. 2012 Sep;28(9):421–6.

1191   36.   Li Y, Venkataram S, Agarwala A, Dunn B, Petrov DA, Sherlock G, et al.
1192   Hidden Complexity of Yeast Adaptation under Simple Evolutionary
1193   Conditions. Curr Biol. 2018 Feb 19;28(4):515–6.

1194   37.   Gorter FA, Aarts MGM, Zwaan BJ, de Visser JAGM. Local Fitness
1195   Landscapes Predict Yeast Evolutionary Dynamics in Directionally Changing
1196   Environments. Genetics. 2018 Jan;208(1):307–22.

1197   38.   Baym M, Stone LK, Kishony R. Multidrug evolutionary strategies to reverse
1198   antibiotic resistance. Science. 2016 Jan 1;351(6268):aad3292–2.

1199   39.   Hartley SW, Monti S, Liu C-T, Steinberg MH, Sebastiani P. Bayesian
1200   methods for multivariate modeling of pleiotropic SNP associations and
1201   genetic risk prediction. Front Genet. 2012;3:176.

1202   40.   Chen G, Mulla WA, Kucharavy A, Tsai H-J, Rubinstein B, Conkright J, et al.
1203   Targeting the adaptability of heterogeneous aneuploids. Cell. 2015 Feb
1204   12;160(4):771–84.

1205   41.   Nichol D, Jeavons P, Fletcher AG, Bonomo RA, Maini PK, Paul JL, et al.
1206   Steering Evolution with Sequential Therapy to Prevent the Emergence of
1207   Bacterial Antibiotic Resistance. PLoS Comput Biol. 2015
1208   Sep;11(9):e1004493.

1209   42.   Kaznatcheev A, Peacock J, Basanta D, Marusyk A, Scott JG. Fibroblasts and
1210   alectinib switch the evolutionary games played by non-small cell lung cancer.
1211   Nat Ecol Evol. 2019 Mar;3(3):450–6.

1212   43.   Paaby AB, Rockman MV. The many faces of pleiotropy. Trends Genet. 2013
1213   Feb;29(2):66–73.

1214   44.   Walsh B, Blows MW. Abundant Genetic Variation + Strong Selection =
1215   Multivariate Genetic Constraints: A Geometric View of Adaptation. Annu
1216   Rev Ecol Evol Syst. 2009 Nov 9;40(1):41–59.

1217   45.   Ference BA, Ginsberg HN, Graham I, Ray KK, Packard CJ, Bruckert E, et al.
1218   Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1.
1219   Evidence from genetic, epidemiologic, and clinical studies. A consensus
1220   statement from the European Atherosclerosis Society Consensus Panel. Eur
1221   Heart J. 2017 Aug 21;38(32):2459–72.

1222   46.   Al Hafid N, Christodoulou J. Phenylketonuria: a review of current and future
1223   treatments. Transl Pediatr. 2015 Oct;4(4):304–17.

1224   47.   Powell ALT, Nguyen CV, Hill T, Cheng KL, Figueroa-Balderas R, Aktas H,
1225   et al. Uniform ripening encodes a Golden 2-like transcription factor
1226   regulating tomato fruit chloroplast development. Science. 2012 Jun
1227   29;336(6089):1711–5.

1228   48.   Grüneberg H. An analysis of the "pleiotropic" effects of a new lethal
1229   mutation in the rat (Mus norvegicus). Proc R Soc Lond B Biol Sci. 1938 Mar
1230   3;125(838):123–144.

1231   49.   Ohya Y, Sese J, Yukawa M, Sano F, Nakatani Y, Saito TL, et al. High-
1232   dimensional and large-scale phenotyping of yeast mutants. Proceedings of the
1233   National Academy of Sciences. 2005 Dec 27;102(52):19015–20.

1234   50.   Bauer CR, Li S, Siegal ML. Essential gene disruptions reveal complex
1235   relationships between phenotypic robustness, pleiotropy, and fitness. Mol
1236   Syst Biol. 2015 Jan;11(1):773–3.

51. Geiler-Samerotte KA, Zhu YO, Goulet BE, Hall DW, Siegal ML. Selection
Transforms the Landscape of Genetic Variation Interacting with Hsp90. PLoS
Biol. 2016 Oct;14(10):e2000465.

52. Richardson JB, Uppendahl LD, Traficante MK, Levy SF, Siegal ML. Histone
variant HTZ1 shows extensive epistasis with, but does not increase robustness
to, new mutations. PLoS Genet. 2013;9(8):e1003733.

53. Negishi T, Nogami S, Ohya Y. Multidimensional quantification of subcellular
morphology of Saccharomyces cerevisiae using CalMorph, the high-
throughput image-processing program. J Biotechnol. 2009 May 20;141(3-
4):109–17.

54. Gerke J, Lorenz K, Cohen B. Genetic interactions between transcription
factors cause natural variation in yeast. Science. 2009 Jan 23;323(5913):498–
501.

55. Hall DW, Mahmoudizad R, Hurd AW, Joseph SB. Spontaneous mutations in
diploid Saccharomyces cerevisiae: another thousand cell generations. Genet
Res (Camb). 2008 Jun;90(3):229–41.

56. Zhu YO, Siegal ML, Hall DW, Petrov DA. Precise estimates of mutation rate
and spectrum in yeast. Proc Natl Acad Sci USA. 2014 Jun 3;111(22):E2310–
8.

57. Cooper TF, Ostrowski EA, Travisano M. A negative relationship between
mutation pleiotropy and fitness effect in yeast. Evolution. 2007
Jun;61(6):1495–9.

58. Yang M, Ohnuki S, Ohya Y. Unveiling nonessential gene deletions that
confer significant morphological phenotypes beyond natural yeast strains.
BMC Genomics. 2014;15(1):932.

59. Sahai E. Mechanisms of cancer cell invasion. Curr Opin Genet Dev. 2005
Feb;15(1):87–96.

60. Gerke JP, Chen CTL, Cohen BA. Natural isolates of Saccharomyces
cerevisiae display complex genetic variation in sporulation efficiency.
Genetics. 2006 Oct;174(2):985–97.

61. Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, et al.
Population genomics of domestic and wild yeasts. Nature. 2009 Mar
19;458(7236):337–41.

62. Skelly DA, Merrihew GE, Riffle M, Connelly CF, Kerr EO, Johansson M, et
al. Integrative phenomics reveals insight into the structure of phenotypic
diversity in budding yeast. Genome Res. 2013 Sep;23(9):1496–504.

63.   Ziv N, Shuster BM, Siegal ML, Gresham D. Resolving the Complex Genetic
1274   Basis of Phenotypic Variation and Variability of Cellular Growth. Genetics.
1275   2017 May 11;:genetics.116.195180.

1276   64.   Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in
1277   experimental crosses. Bioinformatics. 2003 May 1;19(7):889–90.

1278   65.   Churchill GA, Doerge RW. Empirical threshold values for quantitative trait
1279   mapping. Genetics. 1994 Nov;138(3):963–71.

1280   66.   Stuckey S, Storici F. Gene knockouts, in vivo site-directed mutagenesis and
1281   other modifications using the delitto perfetto system in Saccharomyces
1282   cerevisiae. Meth Enzymol. 2013;533:103–31.

1283   67.   Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, et
1284   al. Saccharomyces Genome Database: the genomics resource of budding
1285   yeast. Nucleic Acids Res. 2012 Jan;40:D700–5.

1286   68.   Graziano BR, Yu H-YE, Alioto SL, Eskin JA, Ydenberg CA, Waterman DP,
1287   et al. The F-BAR protein Hof1 tunes formin activity to sculpt actin cables
1288   during polarized growth. Mol Biol Cell. 2014 Jun;25(11):1730–43.

1289   69.   Nogami S, Ohya Y, Yvert G. Genetic Complexity and Quantitative Trait Loci
1290   Mapping of Yeast Morphological Traits. PLoS Genet. 2007 Feb 23;3(2):e31.

1291   70.   Lynch M, Walsh B. Correlations between characters. In: Genetics and
1292   Analysis of Quantitative Traits. Sinauer Associates Incorporated; 1998. pp.
1293   629–56.

1294   71.   Bliese P. Multilevel modeling in R (2.6). 2013. Available from: https://cran.r-
1295   project.org/doc/contrib/Bliese_Multilevel.pdf

1296   72.   Sockloff AL. Behavior of the Product-Moment Correlation Coefficient When
1297   Two Heterogeneous Subgroups Are Pooled. Educational and Psychological
1298   Measurement. 1975;35(2):267–276.

1299   73.   Breiman L, Friedman JH. Estimating Optimal Transformations for Multiple
1300   Regression and Correlation. Journal of the American Statistical Association.
1301   2012 Mar 12;80(391):580–98.

1302   74.   Setty M, Tadmor MD, Reich-Zeliger S, Angel O, Salame TM, Kathail P, et
1303   al. Wishbone identifies bifurcating developmental trajectories from single-cell
1304   data. Nat Biotechnol. 2016 Jun;34(6):637–45.

1305   75.   Turner JJ, Ewald JC, Skotheim JM. Cell size control in yeast. Curr Biol. 2012
1306   May 8;22(9):R350–9.

1307   76.   Wang R, Kamgoue A, Normand C, Léger-Silvestre I, Mangeat T, Gadal O.
1308   High resolution microscopy reveals the nuclear shape of budding yeast during
1309   cell cycle and in various biological states. J Cell Sci. 2016 Dec
1310   15;129(24):4480–95.

1311   77.   Logg K, Bodvard K, Blomberg A, Käll M. Investigations on light-induced
1312   stress in fluorescence microscopy using nuclear localization of the
1313   transcription factor Msn2p as a reporter. FEMS Yeast Res. 2009
1314   Sep;9(6):875–84.

1315   78.   Pavlicev M, Cheverud JM. Constraints Evolve: Context Dependency of Gene
1316   Effects Allows Evolution of Pleiotropy. Annu Rev Ecol Evol Syst.
1317   2015;46(1):413:434.

1318   79.   Pavlicev M, Wagner GP, Noonan JP, Hallgrímsson B, Cheverud JM.
1319   Genomic correlates of relationship QTL involved in fore- versus hind limb
1320   divergence in mice. Genome Biol Evol. 2013;5(10):1926–36.

1321   80.   Porto A, Schmelter R, VandeBerg JL, Marroig G, Cheverud JM. Evolution of
1322   the genotype-to-phenotype map and the cost of pleiotropy in mammals.
1323   Genetics. 2016;204(4):1601–12.

1324   81.   Dudley AM, Janse DM, Tanay A, Shamir R, Church GM. A global view of
1325   pleiotropy and phenotypically derived gene function in yeast. Mol Syst Biol.
1326   2005 Jan 1;1(1):E1–E11.

1327   82.   Lande R. Quantitative genetic analysis of multivariate evolution, applied to
1328   brain:body size allometry. Evolution. 2017 May 31;33(1 Part 2):402–16.

1329   83.   Arnold SJ, Bürger R, Hohenlohe PA, Ajie BC, Jones AG. Understanding the
1330   evolution and stability of the g-matrix. Evolution. 2008 Oct;62(10):2451–61.

1331   84.   Björklund M, Husby A, Gustafsson L. Rapid and unpredictable changes of
1332   the G-matrix in a natural bird population over 25 years. J Evol Biol. 2012 Dec
1333   13;26(1):1–13.

1334   85.   Cheverud JM. A comparison of genetic and phenotypic correlations.
1335   Evolution. 2017 Jun 5;42(5):958–68.

1336   86.   Arnold SJ, Phillips PC. Hierarchical comparison of genetic variance-
1337   covariance matrices. II coastal-inland divergence in the garter snake,
1338   thamnophis elegans. Evolution. 2017 May 31;53(5):1516–27.

1339   87.   Sgrò CM, Hoffmann AA. Genetic correlations, tradeoffs and environmental
1340   variation. Heredity. 2004;93:241–8.

88. Eguchi Y, Bilolikar G, Geiler-Samerotte K. Why and how to study genetic
changes with context-dependent effects. Curr Opin Genet Dev. 2019 Oct;58–
59:95–102.

89. Pavlicev M, Wagner GP. Evolutionary Systems Biology: Shifting Focus to
the Context-Dependency of Genetic Effects. In: Martin LB, Ghalambor CK,
Woods HA, editors. Integrative Organismal Biology. 2014. pp. 91–108.

90. Levy SF, Siegal ML. Network hubs buffer environmental variation in
Saccharomyces cerevisiae. PLoS Biol. 2008 Nov 4;6(11):e264.

91. Ziv N, Siegal ML, Gresham D. Genetic and nongenetic determinants of cell
growth variation assessed by high-throughput microscopy. Mol Biol Evol.
2013 Dec;30(12):2568–78.

92. Millard SP. EnvStats. New York, NY: Springer Science & Business Media;
2013.

93. Storey JD, Bass A, Dabney A, Robinson D. qvalue: Q-value estimation for
false discovery rate control. R package version 2.16.0. 2019.

94. Gietz RD, Woods RA. Transformation of yeast by lithium acetate/single-
stranded carrier DNA/polyethylene glycol method. Meth Enzymol.
2002;350:87–96.

95. Miyajima I, Nakafuku M, Nakayama N, Brenner C, Miyajima A, Kaibuchi K,
et al. GPA1, a haploid-specific essential gene, encodes a yeast homolog of
mammalian G protein which may be involved in mating factor signal
transduction. Cell. 1987 Sep 25;50(7):1011–9.

96. Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. nlme: linear and
nonlinear mixed effects models. R package version 3.1-137. 2018.

97. Levy SF, Ziv N, Siegal ML. Bet hedging in yeast by heterogeneous, age-
correlated expression of a stress protectant. PLoS Biol. 2012;10(5):e1001325.

98. Li S, Giardina DM, Siegal ML. Control of nongenetic heterogeneity in
growth rate and stress tolerance of Saccharomyces cerevisiae by cyclic AMP-
regulated transcription factors. PLoS Genet. 2018 Nov;14(11):e1007744.

99. Wickham H. ggplot2. New York, NY: Springer New York; 2009.

Figure 1: Pleiotropic QTL influence yeast single-cell morphology. The vertical axes in all plots represent the 155 CalMorph morphological traits for which we detect QTL. These traits are sorted, from top to bottom, based on the difference between the oak and wine parental strains. **(A)** Of 41 QTL that contribute to variation in single-cell morphology, 36 contribute to variation in multiple features. The horizontal axis indicates the chromosomal location of each QTL (in cM). Differently shaped points indicate separate QTL that are more than 5 cM apart on the same chromosome. The darkness of a point represents the effect size of a QTL; effect sizes range from 1.3% (lightest points) to 17.5% (darkest points) of the difference between parents. All points represent genotype-phenotype associations detected using a per-trait genome-wide type one error rate of 5%. The points highlighted in black are significant after correcting for testing multiple phenotypes using a false discovery rate of 5%. The QTL highlighted in pink, green, purple, and orange are very pleiotropic, contributing to 57, 30, 73, or 64 morphological features, respectively. **(B)** Gene-swapping experiments demonstrate that single genes contribute to multiple morphological features. The horizontal axis represents the relative phenotypic differences between the wine and oak parents (leftmost column) or one of these strains versus a derivative strain that differs in a single gene. The relative phenotypic differences between a pair of strains are calculated by scaling each trait to have a mean of 0 and standard deviation of 1 across all cells in both strains, and then subtracting the average value in one strain from that in the other. To control for variation among replicate experiments, this scaling was done independently for each replicate experiment in which both strains were imaged. Error bars represent 95% confidence intervals inferred from the replicate experiments. The two gene replacements shown, *PXL1* and *HOF1*, are respectively located within the QTL highlighted in green and purple in panel **A**. When calculating the difference between strains, we always subtracted the trait values of the strain possessing more wine genes from those of the strain possessing more oak genes, such that the effects of the wine or oak gene replacements appear in the same direction on all plots.

Figure 2: Pairs of traits with high correlation across clones are overrepresented among those influenced by pleiotropic QTL. Within-genotype correlations (r_W) are calculated for 5645 pairs of morphological traits. **(A)** Histogram showing distribution of r_W. **(B)** Each point represents r_W for a pair of traits. The null distribution displays r_W for all 5645 pairs of traits. Every other distribution displays r_W for pairs of traits influenced by a single QTL. Horizontal axis labels represent the chromosome on which a QTL resides, followed by the order in which that QTL appears on the chromosome. Colored points correspond to those QTL highlighted in the same color in **Fig 1A**. Each boxplot shows the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5 × IQR (whiskers). **(C)** Force-directed network visualizing how pairs of morphological features correlate across clones. Each node represents a single-cell morphological trait measured in large-budded cells. For networks representing traits from unbudded and small budded cells, see **Fig S4**. The thickness of the line connecting each pair of nodes is proportional to r_W. Node position in the network is determined using the Fruchterman-Reingold algorithm. Purple nodes correspond to traits influenced by a QTL on chromosome 13 containing the *HOF1* gene. **(D)** Cumulative distributions of weighted clustering coefficients (*wcc*) in a network created using measured values of r_W (red line) or in 100 permuted networks (grey lines) for traits corresponding to large-budded cells. Permutations were performed by sampling r_W, without replacement, and reassigning each value to a random pair of traits. For distributions summarizing *wcc* in networks representing traits from unbudded and small budded cells, see **Fig S4**. **(E)** The same network as in panel C with colored nodes corresponding to traits influenced by the leftmost QTL on chromosome 15.

Figure 3: Natural genetic variation affects the correlation between morphological features. The absolute value of the between-strain correlation (r_B), made negative when r_B and r_W have opposite signs, is plotted against the absolute value of the within-strain correlation (r_W), for each pair of traits. The plot at left shows pairs of traits that share at least one QTL influence. The color of each point represents the number of pleiotropic QTL that influence both traits in that pair. The plot at right shows pairs of traits that share no QTL influence. The dashed line represents a Bonferroni-corrected significance threshold of $p < 0.01$. Insets represent the results of correlation partitioning performed after randomly assigning individual cells to groups (pseudo-strains) having the same numbers of cells as the actual strains.

Figure 4: Morphological features vary as cells divide. The morphological features of unbudded (red), small-budded (blue), and large-budded (purple) cells change as these cells progress through the cell cycle. **(A)** Variation of four traits through the cell cycle. Line plots represent fixed-cell images from all 374 mapping family strains, positioned on the horizontal axis based on progression through the cell cycle as calculated by Wishbone (74). Regression lines are smoothed with cubic splines, calculated with the "gam" method in the R package ggplot2 (99), to depict trends describing how each displayed trait varies across the estimated growth trajectory. The displayed trends match those observed in micrographs of live cells progressing through division. Each series of micrographs displays a different live cell imaged over several minutes, which are displayed in the lower right corner of each micrograph. **(B)** Centered data for 11, 23, and 44 unbudded, small-budded and large-budded cells, respectively, show how Wishbone sorts live cells in a way that recapitulates the actual time series. Each point in these plots represents a cell image from a single timepoint. The horizontal axis represents Wishbone's estimation of how far that cell has progressed through division. The vertical axis displays time, as a percentage of the total time elapsed and adjusted in a way that controls for every cell having started at a different place in the cell division cycle at time zero (see *Methods*). Trend lines are smooth fits using the "loess" method in the R package ggplot (99). **(C)** The correlation between some morphological features changes throughout the course of cell division. The scatterplot shows how binning influences both the phenotypic correlation (vertical axis) and phenotypic variation (horizontal axis) across clones. Each point represents these values for a pair of traits as measured in 1 of 16 bins. The value on the horizontal axis represents whichever trait in each pair had the larger decrease in standard deviation, as such decreases are likely to reduce the correlation on the vertical axis. The blue line shows a smooth fit by loess regression. Colored points on the scatterplot correspond to bin 5 for each pair of traits represented by the line plots in panel **D**. **(D)** These line plots show three pairs of traits for which binning increases r_W such that it approaches r_B (leftmost three plots), and three pairs of traits for which r_W does not approach r_B even after binning (rightmost three plots). In each plot, r_B is shown as the horizontal green line, r_W (without binning) is shown as the horizontal purple line, and r_W for each bin is shown in black.

Figure 5: Many QTL demonstrate horizontal pleiotropy. (A) Eliminating allelic variation at the site of each QTL tends to reduce r_B. The vertical axis represents how r_B changes upon eliminating allelic variation at each QTL site. Each point represents the median change in r_B for all pairs of traits that are affected or unaffected by one of the 27 QTL suspected of horizontal pleiotropy. Boxplots summarize these changes in r_B when re-measured across strains possessing the wine (red) or the oak (blue) allele at the marker closest to the QTL. **(B)** The upper and lower series of three plots demonstrate two different ways that a QTL can increase the correlation between traits. Each point represents a yeast strain possessing either the wine (red) or the oak (blue) allele at a marker closest to a QTL on chromosome 15 (upper) or 8 (lower). In the upper plots, the QTL increases the correlation between nucleus shape and size ratio when it is segregating across strains. In the lower plots, the oak allele strengthens a correlation between bud shape and the position of the nucleus in the mother cell that is weak in the wine subpopulation. Numbers in the lower corner of each plot represent r_B for the strains displayed. **(C)** Cumulative distributions display the extent to which binning cells or splitting strains resolves the difference between r_B and r_W. When calculating percent resolved (horizontal axes) we always plot the value in whichever subset (*e.g.* wine or oak) this percent is greatest. If subsetting always worsens the discrepancy between r_B vs. r_W, we score this as 0% resolution. Only pairs of traits for which r_B is significantly greater than r_W are considered. The pink, green, purple and orange lines show the effect of splitting strains by whether they inherited the wine or oak allele at the marker closest to each of four QTL (colors correspond to QTL in **Fig 1A**). In these plots, comparing the solid vs dotted lines shows that splitting strains resolves the discrepancy between r_B and r_W more often for pairs in which both traits are affected by the QTL than pairs in which both traits are unaffected. The black lines in the leftmost plot summarize these effects across 27 QTL, displaying for each trait pair, the largest resolution in the r_B vs. r_W discrepancy observed across all QTL that affect the pair of traits (solid line) or all QTL that do not (dotted line). The red line shows the effect of binning cells by their progress through division, displaying the largest resolution in the r_B vs. r_W difference across all 16 bins.

Figure 6: Some MA lines display unique relationships between certain pairs of traits. In all plots, black represents the ancestor of the MA lines and colors represent MA lines with trait correlations that differ from other lines (strains: black = HAncestor, green = DHC81H1, red = DHC41H1, magenta = DHC40H1, blue = DHC66H1, purple = DHC84H1; see Table S2 in Geiler-Samerotte et al 2016 (51)). **(A)** Histograms display the number of MA lines with Pearson correlations corresponding to the values on the horizontal axis for four example pairs of traits; the number of bins is set to 30. **(B)** This plot displays, for each of the 94 MA lines, the cumulative distribution of the number of standard deviations away from the mean correlation across all trait pairs. **(C)** Plots display, for each MA line, the maximum deviation from the mean observed for any pair of traits (left) and the average standard deviation observed across all pairs of traits (right).

Figure 7: The correlations between traits changes depending on drug concentration. Plots display the density of trait pairs for which the within-strain correlation (r_W) changes by the amount shown on the horizonal axis. To calculate how each drug treatment changes r_W, we subtracted r_W observed for a pair of traits in the drug condition from that observed in a paired experiment that lacked the drug. The absolute value of this change is displayed. These changes are smallest in the null condition, which represents the change in r_W observed across replicate experiments lacking the drug. For clarity, we exclude regions of the plot for which density is less than 0.01.

Figure S1: Morphological differences exist between the parents of the QTL mapping family. (A–C) Each density plot displays the distribution of phenotype values from yeast cells corresponding to the wine parent (red), the oak parent (blue), or all of the 374 progeny (grey) for the trait listed on the horizontal axis. Trait names in parentheses correspond to those listed in the CalMorph manual (53). Each distribution represents at minimum 5,000 cells from three replicate experiments; distributions corresponding to progeny strains represent many more cells (70,000 – 200,000 depending on whether the trait was measured in unbudded, small-budded, or large-budded cells). **(D)** The broad sense heritability for each of the 155 morphological features for which QTL were detected. Heritability is low because cell morphology varies across the cell cycle, and so the amount of non-genetic phenotypic variation is high.

Figure S2: Total numbers of cells imaged per each of 374 progeny strains. Each point represents, for one of the 374 progeny strains, the number of unbudded, small-budded, or large-budded cells for which images passed filtering (see *Methods*). Each boxplot shows the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5 × IQR (whiskers).

Figure S3: Comparison of correlation estimates obtained from correlation partitioning with those obtained from a mixed-effect linear model. Each point represents one of 350 randomly sampled trait pairs of the 5645 total. Vertical axes display trait correlations estimated using the correlation-partitioning approach; horizontal axes display trait correlations estimated using a mixed-effect linear model that specifies the variance-covariance structure of the experimental design.

Figure S4: Single-cell morphological traits have higher weighted clustering coefficients (*wcc*) than expected given the distribution of r_W. (A – B) Force-directed networks visualizing how pairs of morphological features correlate across clones in unbudded (panel A) and small-budded (panel B) cells. Each node represents a single-cell morphological trait. The thickness of the line connecting each pair of nodes is proportional to r_W. Node position in the network is determined using the Fruchterman-Reingold algorithm. Purple nodes correspond to traits influenced by a QTL on chromosome 13 containing the *HOF1* gene. **(C – D)** Cumulative distributions of weighted clustering coefficients (*wcc*) in a network created using measured values of r_W (red line) or in 100 permuted networks (grey lines) for traits corresponding to unbudded (panel C) or small-budded (panel D) cells. Permutations were performed by sampling r_W, without replacement, and reassigning each value to a random pair of traits.

Figure S5: Wishbone recapitulates time series data obtained in live images of 78 cells undergoing exponential growth. Each point represents a cell image. Horizontal axes display the minute that image was captured during a three-hour window of exponential growth. Vertical axes display Wishbone's prediction of how far that cell image has passed through the cell cycle. Linear regression lines are calculated with the "lm" method in the R package ggplot2 (99), and are colored red for images corresponding to unbudded cells, blue for small-budded cells and purple for large budded cells. Plots are organized by cell type and then from earliest to latest average predicted progress through cell division.

Figure S6: Total numbers of cells imaged per strain in varying concentrations of GdA. Each point represents, for one of the strains, the number of unbudded, small-budded, or large-budded cells for which images passed filtering (see *Methods*). Each boxplot shows the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5 × IQR (whiskers).

S1 Table. Chromosomal locations, effects sizes and phenotypes affected by quantitative trait loci described in this study.

S2 Table. Impact of gene swaps on single-cell morphological traits including the corrected phenotypic difference between strains for each phenotype, and its standard deviation and standard error across replicate experiments.