1 **Title:**
2 Extent and context dependence of pleiotropy revealed by high-throughput single-cell
3 phenotyping
4
Author list:

Kerry A Geiler-Samerotte[a,b]

Shuang Li[a]

Charalampos Lazaris[a]

Austin Taylor[a]

Naomi Ziv[a]

Chelsea Ramjeawan[a]

Annalise B Paaby[c]

Mark L Siegal[a]

14
Author affiliations:

[a] Center for Genomics and Systems Biology, Department of Biology, New York
University, New York, NY 10003

[b] Center for Mechanisms of Evolution, Biodesign Institutes, School of Life Sciences,
Arizona State University, Tempe, AZ 85287

[c] School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332
21
22 **Corresponding author:**
23 Kerry Geiler-Samerotte: kerry.samerotte@asu.edu

Abstract:

Pleiotropy – when a single mutation affects multiple traits – is a controversial topic with far-reaching implications. Pleiotropy plays a central role in ongoing debates about how complex traits evolve and whether biological systems tend to be modular or organized such that every gene has the potential to affect many traits. Pleiotropy is also critical to initiatives in evolutionary medicine that seek to trap infectious microbes or tumors by selecting for mutations that encourage growth in some conditions at the expense of others. Research in these fields, and others, would benefit from understanding the extent to which pleiotropy reflects inherent relationships among phenotypes that correlate no matter the perturbation (vertical pleiotropy), versus the action of genetic changes that impose correlations between otherwise independent traits (horizontal pleiotropy). We tackle this question by using high-throughput single-cell phenotyping to measure thousands of pairwise trait correlations across hundreds of thousands of cells representing hundreds of genotypes of the budding yeast, *Saccharomyces cerevisiae*. We map pleiotropic quantitative trait loci using genotypes derived from a cross between natural strains, and we separate vertical and horizontal pleiotropy by partitioning trait correlations into within- and between-genotype correlations. We investigate how pleiotropy can change by using genotypes from mutation-accumulation lines that experienced minimal selection, and by tracking trait correlations through the cell-division cycle. We find ample evidence of both vertical and horizontal pleiotropy, and observe that trait correlations depend on both genetic background and cell-cycle position. Our results suggest a nuanced view of pleiotropy in which trait correlations are highly context dependent and biological systems occupy a middle ground between modularity and interconnectedness. These results also suggest an approach to select pairs of traits that are more likely to remain correlated across contexts for applications in evolutionary medicine.

Introduction

Pleiotropy exists when a single mutation affects multiple traits [1,2]. Often, pleiotropy is defined instead as a single gene contributing to multiple traits, although what is implied is the original definition — that a single change at the genetic level can have multiple consequences at the phenotypic level [2]. As our ability to survey the influence of genotype on phenotype improves, examples of pleiotropy are growing [3-7]. For example, individual genetic variants have been associated with seemingly disparate immune, neurological, and digestive symptoms in humans and mice [8,9]. Genes affecting rates of cell division across diverse environments and drug treatments have been identified in microbes and cancers [10,11]. A view emerging from genome-wide association studies is that variation in complex traits is "omnigenic" in the sense that many loci indirectly contribute to variation in many traits [12,13].

However, the extent of pleiotropy remains a major topic of debate because, despite its apparent prevalence, pleiotropy is thought to be evolutionarily disadvantageous. The more traits a mutation affects, the more likely it is that the mutation will have a negative impact on at least one. Pervasive pleiotropy should therefore constrain evolution [14], exacting what is known as a cost of complexity or cost of pleiotropy [10,15-18]. This cost may bias which mutations underlie adaptation, for example, toward less-pleiotropic *cis*-regulatory changes over more-pleiotropic changes in *trans*-acting factors [19,20], or toward changes to proteins that participate in relatively few biological processes [21,22]. Over long periods, the cost of pleiotropy may influence the organization of biological systems, favoring a modular structure in which genetic changes influencing one group of traits have minimal impact system-wide [23-28].

At stake in the ongoing debate about the extent of pleiotropy [29-32] are some of modern biology's prime objectives, including the prediction of complex phenotypes from genotype data [17,33,34] and the prediction of how organisms will adapt to environmental change [35,36]. These predictions are more challenging if genetic changes influence a large number of traits with complex interdependencies. Nonetheless, understanding how a given mutation influences multiple traits could be powerful, allowing prediction of some phenotypic responses given others [37,38]. Indeed, recent strategies in medicine called evolutionary traps aim to exploit pleiotropy, for example by finding genetic changes that provide resistance to one treatment while promoting susceptibility to another [39-41].

The lack of consensus about the extent of pleiotropy in natural systems is in part due to poorly defined expectations for how to test for it experimentally. One key issue is that defining a phenotype is not trivial [42,43]. Consider a variant in the *apolipoprotein B* gene that increases low-density lipoprotein (LDL) cholesterol levels as well as the risk of heart disease. Elevated LDL promotes heart disease [44], so are these two phenotypes or one? Alternatively, consider a mutation in the *phenylalanine hydroxylase* gene that affects nervous system function and skin pigmentation. These dissimilar effects, both symptoms of untreated phenylketonuria (PKU), originate from the same problem: a deficiency in converting phenylalanine to tyrosine [45]. Is it fair to call mutations that have this single metabolic effect pleiotropic? Likewise, shall one call pleiotropic a mutation that makes tomatoes both ripen uniformly and taste bad, when the effect of the

mutation is to reduce the function of a transcription factor that promotes chloroplast
development, which in turn necessarily affects both coloration and sugar accumulation
[46]?

The LDL, PKU and tomato cases are examples of vertical pleiotropy, *i.e.*
pleiotropy that results when one phenotype influences another or both are influenced by a
shared factor [4,42]. The alternative to vertical pleiotropy is horizontal pleiotropy, in
which genetic differences induce correlations between otherwise independent
phenotypes. It might be tempting to discard vertical pleiotropy as less "genuine" [47] or
less important than horizontal pleiotropy, but that would be a mistake because vertical
pleiotropy reveals important information about the underlying biological systems that
produce the phenotypes in question. Consider the value in identifying yet-unknown
factors in heart disease by finding traits that correlate with it, or in understanding where
in a system an intervention is prone to produce undesirable side effects. Consider also
that the extent and nature of vertical pleiotropy speak directly to the question of
modularity: modularity is implied if vertical pleiotropy either is rare or manifests as small
groups of correlated traits that are isolated from other such groups. If there is modularity
then there can be horizontal pleiotropy, when particular genetic variants make links
between previously unconnected modules.

The above considerations suggest that a unified analysis that distinguishes and
compares horizontal and vertical pleiotropy is needed to make sense of the organization
and evolution of biological systems. However, existing methods of distinguishing
horizontal and vertical pleiotropy are problematic because judgments must be made about
which traits are independent from one another. Such judgments differ between
researchers and over time. Indeed, the tomato example can be viewed as a case of
horizontal pleiotropy transitioning recently to vertical pleiotropy as knowledge of the
underlying system advanced.

In this study, we propose and apply an empirical and analytical approach to
measuring pleiotropy that relies far less on subjective notions of what constitutes an
independent phenotype. The key principle is that the distinction between vertical and
horizontal pleiotropy lies in whether traits are correlated in the absence of genetic
variation [42]. For vertical pleiotropy, the answer is yes: because one trait influences the
other or the two share an influence, non-genetic perturbations that alter one phenotype are
expected to alter the other. For horizontal pleiotropy, the answer is no: genetic variation
causes the trait correlation. In this study, we determined how traits correlate in the
absence of genetic variation by measuring single-cell traits in clonal populations of cells.

We used high-throughput morphometric analysis [48-52] of hundreds of
thousands of single cells of the budding yeast *Saccharomyces cerevisiae* to measure how
dozens of cell-morphology traits (thousands of pairs of traits) co-vary within clonal
populations and between such populations representing different genotypes. Within-
genotype correlations report on vertical pleiotropy, whereas between-genotype
correlations report on horizontal pleiotropy to the extent that they exceed the
corresponding within-genotype correlations. For one set of genotypes, we used 374
progeny of a cross of two natural isolates [53], which enabled not only the estimation of

vertical and horizontal pleiotropy but also the identification of quantitative trait loci
(QTL) with pleiotropic effects. For another set of genotypes, we used a collection of
mutation-accumulation lines, each of which contains a small number of unique
spontaneous mutations [54,55], which enabled a more direct test of the ability of
mutations to alter trait correlations.

The traits we study – morphological features of single cells – represent important
fitness-related traits [50,56,57] that contribute to processes such as cell division and
tissue invasion (*e.g.* cancer metastasis [58]). Cell-morphological features may correlate
across cells for a variety of vertical or horizontal reasons. Vertical reasons include: (1)
inherent geometric constraints (*e.g.* on cell circumference and area); (2) constraints
imposed by gene-regulatory networks (*e.g.* if the genes influencing a group of traits are
all under control of the same transcription factor); and (3) constraints induced by
developmental processes (*e.g.* as a yeast cell divides or "buds", many morphological
features are affected). Horizontal pleiotropy might be evident because genetic variants
affecting two or more traits (that are otherwise weakly correlated) are segregating in the
progeny of the cross between two natural isolates. Alternatively, horizontal pleiotropy
might be evident because a particular allele strengthens the trait correlation so that
genetic variation affecting one trait is more likely to affect another when that allele is
present. These alternatives can be distinguished by examining trait correlations in two
subsets of progeny strains defined by which natural isolate's allele they possess at a QTL
of interest.

In addition to genetic variation, non-genetic variation may also alter the
correlations between traits. We rely on non-genetic heterogeneity within clonal
populations to serve as perturbations that reveal inherent trait correlations. However, the
correlations themselves might be heterogeneous within these populations. For example,
the dependencies between morphological features may change as cells divide. To control
for this possibility, we performed our trait mapping and subsequent analysis after binning
cells into three stages (unbudded, small-budded and large-budded cells). We further
examined whether trait correlations change across the cell cycle by using a machine-
learning approach to more finely bin the imaged cells into 48 stages of division.
207
Collectively, the results we present here demonstrate that both types of pleiotropy,
vertical and horizontal, are prevalent for single-cell morphological traits, suggesting that
biological systems occupy a middle ground between extreme modularity and extreme
interconnectedness. Perhaps more surprisingly, we find that trait correlations are often
context dependent, and can be altered by mutations as well as cell-cycle state. The
dynamic nature of trait correlations at these different timescales encourages caution when
attempting to quantify and interpret the extent of pleiotropy in nature or when making
predictions about correlated phenotypic responses to the same selection pressure, as is
done when crafting evolutionary traps. However, applying our approach may suggest
which trait correlations are less context dependent and therefore more useful in setting
such traps.
219
220

Results:

QTLs with pleiotropic effects influence yeast single-cell morphology

To detect genes with pleiotropic effects on cell morphology, we measured 167 single-cell morphological features (*e.g.* cell size, bud size, bud angle, distance from nucleus to bud neck; **Table S1**) in 374 yeast strains that were generated in a previous study from a mating between two wild yeast isolates [53,59]. These wild isolates, one obtained from soil near an oak tree, the other from a wine barrel, differ by 0.006 SNPs per site [60] and have many heritable differences in single cell morphology [61]. For example, we find that yeast cells from the wine strain, on average, are smaller, are rounder, and have larger nuclei during budding than yeast cells from the oak strain (**Fig S1**).

To measure their morphologies, we harvested exponentially growing cells from three replicate cultures of each of these 374 recombinant strains, and imaged on average 800 fixed, stained cells per strain using high-throughput microscopy in a 96-well plate format (**Fig S2**). We used control strains present on each plate to correct for plate-to-plate variation (see *Methods*), and quantified morphological features using CalMorph software [52], which divides cells into three categories based on their progression through the cell cycle (*i.e.* unbudded, small-budded, and large-budded cells) and measures phenotypes specific to each category. Using 225 markers spread throughout the genome [53] and Haley-Knott regression implemented in the R package R/qtl [62,63], we identified 44 QTL that contribute to variation in 158 of the surveyed morphological features (FDR = 5%; **Fig 1A; Table S1**). Most (37) of the QTL we detected are pleiotropic, meaning each contributes to variation in more than one morphological feature (**Fig 1A**). The median number of traits to which each QTL contributes is six.

Single genes with pleiotropic effects influence yeast single-cell morphology

When a QTL affects multiple traits, it might not mean that variation in a single gene is contributing to variation in these traits but instead that linked genes are contributing to variation in distinct, individual traits. For several QTL with high pleiotropy (highlighted in **Fig 1A**), we sought to test whether the effects on different morphological features were due to the action of a single gene. We performed these tests by swapping the parental versions of candidate genes (*i.e.* we genetically modified the wine strain to carry the oak version of a given gene, and vice versa). We used the *delitto perfetto* technique to perform these swaps [64], such that the only difference between a parental genome and the swapped genome is the coding sequence of the single candidate gene plus up to 1 kb of flanking sequence (see *Methods*). Candidate genes were selected based on descriptions of the single-cell morphologies of their knockout mutants [65] and the presence of at least one non-synonymous amino acid difference between the wine and oak alleles [61].

When a candidate gene contributes to the morphological differences between the wine and oak parents, we expect yeast strains that differ at only that locus to recapitulate some of the morphological differences between the wine and oak parents. Indeed, this is what we observe for *PXL1*, a candidate for the QTL on chromosome 11, and *HOF1*, a candidate for the QTL on chromosome 13 (**Fig 1B**; compare each plot on the right to the leftmost plot; see also **Table S2**). This influence is most pervasive for *HOF1*; both the

oak and the wine alleles have a strong effect on the morphology of the opposite parent,
and their effects recapitulate the parental difference to a large extent. The pervasive
influence of *HOF1* on various morphological features is consistent with the fact that this
gene's product affects actin-cable organization and is involved in both polar cell growth
and cytokinesis [66]. The effect of *PXL1* on cell morphology is also apparent across
many single-cell features, although only the oak allele has a strong effect that
recapitulates the parental difference. We evaluated *RAS1*, a candidate for the QTL on
chromosome 15, but initial tests indicated that it did not have a significant impact on
most morphological features (**Table S2**). We also attempted to swap alleles for a
candidate gene corresponding to the QTL on chromosome 8, but were unsuccessful (see
Methods).
A previous screen for QTL influencing single-cell morphology in the progeny of a
genetically distinct pair of yeast strains (a different vineyard strain and a laboratory
strain) found some of the same pleiotropic QTL that we detect in the wine and oak cross
[67] (compare their Table 2 to our **Table S1**). In particular, we both find a QTL in the
same position on chromosome 15 that influences many morphological features related to
nucleus size, shape, and position in the cell (**Fig 1A**; orange). We also both detect a QTL
near base pair 100,000 on chromosome 8 that influences cell size and shape (**Fig 1A**;
pink). In the previous screen, the genetic basis of this QTL was shown to be a single
nucleotide change within the *GPA1* gene [67].
The main conclusion from our gene-swapping experiments, which is consistent
with the previous cell-morphology QTL study [67] as well as with comprehensive
surveys of how gene deletions affect the morphology of a laboratory yeast strain [10,48],
is that genes with pleiotropic effects on cell morphology are common in budding yeast.
Moreover, the morphological traits involved were previously shown to influence fitness
[50,56,57], which raises the question: why do so many genetic analyses (including ours)
detect pleiotropy [4,8-11,13] when other work suggests that pleiotropy exacts a cost
[16,17,19,20]?
295
Dissecting pleiotropy using clonal populations of cells
One hypothesis to explain pervasive pleiotropy may be that the phenotypes we
chose to measure are not independent. Instead, many of these single-cell morphological
features may be inherently related such that perturbing one will have unavoidable
consequences on another and thus any associated limitation of adaptation will be
unavoidable as well. In other words, the hypothesis is that much of the pleiotropy we
observe is vertical pleiotropy. A test of this hypothesis is to ask whether traits are
correlated in the absence of genetic differences. Our dataset provides a unique
opportunity to perform such a test because we quantified single-cell traits for, on average,
800 clonal cells per yeast strain (**Fig S2**).
We can leverage the hierarchical structure and large sample size of our dataset to
obtain precise estimates of the correlations that exist within and between strains, and
thereby to distinguish vertical from horizontal pleiotropy. Because we are studying clonal
families without a complicated pedigree structure, these within- and between-strain
correlations are equivalent to the so-called environmental and genetic correlations of
quantitative genetics [68]. Here, we use a simple (and fast) method that is appropriate for
two-level hierarchical data to partition the total correlation into a pooled within-strain

component (r_W) and a between-strain component (r_B) [69]. One caveat of this correlation-
partitioning approach is that r_B is effectively the correlation between strain means, which
can bias estimates of genetic covariance [68]. This bias is most pronounced at small
sample sizes [68], so our large sample sizes allay concern. Nonetheless, for a subset of
traits, we tested whether estimates obtained from correlation partitioning are similar to
those obtained from mixed-effect linear models that specify the variance-covariance
structure of the experimental design. Environmental correlations estimated using both
methods are nearly identical (**Fig S3**). Genetic correlations estimated by correlation
partitioning are sometimes slightly smaller in magnitude than those obtained by linear
modeling (**Fig S3**). This bias is conservative; it may prevent us from identifying cases
where the environmental and genetic correlations significantly differ but will not tend to
create such cases. Despite this reduced power, we rely on the correlation-partitioning
approach, which is substantially faster, because our goal is to estimate environmental and
genetic correlations for thousands of trait pairs.
Unlike the mapping analysis, which considered phenotypes across all three
classes of cell type (unbudded, small-budded and large-budded), this correlation-
partitioning analysis can only be applied to pairs of phenotypes measured in the same cell
type. Two of the 37 pleiotropic QTL exclusively affect traits from different cell types: a
QTL on chromosome 8 affects the short-axis length of unbudded cells and the short-axis
length of large-budded cells, and a QTL on chromosome 7 affects the cell axis ratio of
small-budded cells and the bud axis ratio of large-budded cells. The correlations for the
trait pairs affected by these two QTL cannot be partitioned because the traits are not
measured in the same cells within strains. Excluding these two QTL leaves 35 pleiotropic
QTL collectively contributing to 5645 pairs of traits (378, 1081, and 4186 pairs of
morphological features pertaining to unbudded, small-budded, and large-budded cells
respectively). For each of these trait pairs, we partitioned the correlation between traits.
In the analyses that follow, when we refer to r_W or r_B, we mean the magnitudes of these
correlations, as the sign has no relevance for arbitrary pairs of traits.
341
Inherent relationships between traits contribute to pleiotropy
We focus first on vertical pleiotropy by analyzing correlations that exist in the
absence of any genetic differences (r_W). The distribution of r_W values reflects the extent
of vertical pleiotropy, and the overall pattern of r_W values (*i.e.*, whether there are isolated
clusters of highly correlated traits versus a densely interconnected network of traits)
reflects the modularity of the underlying biological system. These within-strain
correlations are estimated with extremely high precision because of our large sample size
of hundreds of thousands of clonal cells (800 per each of 374 strains).
Most pairs of single-cell morphological traits are not strongly correlated across
clonal cells (**Fig 2A**). Median r_W is < 0.1, and 74% of pairs have r_W < 0.2. Even if we
allow for nonlinear correlations by transforming data using a nonparametric model that
finds the fixed point of maximal correlation [70], r_W is less than 0.2 for roughly 65% of
pairs. These observations suggest that most of the morphological traits we surveyed are
not inherently related; i.e. for any individual cell, the value of one trait does not predict
well the values of most other traits.
Nonetheless, the distribution of r_W has a prominent right tail (**Fig 2A**) indicating
that some morphological features are strongly correlated across clonal cells. These

correlated features are more likely to be influenced by pleiotropic QTL. Among pairs
represented by this right tail (specifically, those with $r_W > 0.2$), 78% consist of traits that
share at least one QTL influence; the same is true for only 40% of pairs with $r_W < 0.2$.
Further, the number of pleiotropic QTL influencing both traits in a pair correlates with
that pair's r_W (Pearson's $r = 0.52$). This result suggests that some of the pleiotropy we
observe is vertical and results from inherent correlations that cause genetic perturbations
to one morphological trait to have consequences on another.
Next, we asked how many pleiotropic QTL act exclusively via vertical
pleiotropy—that is, how many QTL, identified across genotypes, are only associated with
traits that correlate highly within genotypes. Of the 35 pleiotropic QTL that we
examined, 11 exclusively influence traits with $r_W > 0.2$ (**Fig 2B**). For example, a QTL on
chromosome 10 influences a single pair of traits – the area of the nucleus and the length
of the nucleus in large-budded cells – with an r_W of 0.9, suggesting that the aspect ratio
of the nucleus is constrained by vertical pleiotropy (**Fig 2B**). Another QTL on
chromosome 16 exclusively influences traits corresponding to unbudded cells, all with r_W
> 0.4 (**Fig 2B**). Excluding these 11 QTL, nearly all (21) of the remaining 24 QTL have a
median r_W for the pairs of traits they influence that is higher than the median r_W given by
all possible pairs of traits (**Fig 2C**, compare All QTL to All Pairs). In sum, pairs of traits
with stronger correlations across clones (higher r_W) are disproportionately represented
among those influenced by pleiotropic QTL, suggesting that vertical pleiotropy drives a
large portion of the pleiotropy we detect.
Next, we investigated the organization of the biological system underlying cell
morphology by using network analysis to move beyond pairwise comparisons and ask if
morphological traits tend to be clustered into modules. Traits with higher r_W do indeed
tend to group into clusters in networks in which the single-cell morphological traits are
nodes and the r_W magnitudes are edge weights (**Fig 2D** shows the network for traits of
large-budded cells). This need not have been the case; single pairs of traits with high r_W
could have been distributed throughout the network without necessarily being clustered
near other high r_W pairs. Instead, networks representing single-cell morphological
features demonstrate more clustering than do random networks drawn from the same
values of r_W (**Fig 2E**; for corresponding figures from unbudded and small-budded trait
networks, see **Fig S4**). This observation might indicate that morphological phenotypes
have a modular organization, whereby phenotypes within a module exert influence on
one another, but exert less influence on phenotypes from other modules. However, this
observation could also result from human bias when enumerating phenotypes that can be
measured, in the sense that phenotypes that bridge modules might somehow be absent
from the data set. The comprehensive nature of CalMorph diminishes this concern. A
related concern is that apparent modules are formed by trivially related phenotypes, such
as the radius and diameter of a circular object, but we do not find such trivial
relationships among the CalMorph phenotypes. Even a high correlation between the
length and area of the nucleus, as noted above, implies a constraint on nuclear aspect
ratio.
Some pleiotropic QTL tend to influence traits that are clustered in these networks.
Even when we focus on the 24 pleiotropic QTL that do not exclusively influence traits
with high r_W, we find that more influence traits with higher weighted clustering
coefficients (*wcc*) than expected given the distribution of *wcc* across all traits (**Fig 2F**,

compare All QTL to All Pairs). For example, the QTL containing the *HOF1* gene has a
slight tendency to influence traits with higher than average **wcc** (**Fig 2D**: purple nodes;
Fig 2F: purple points).
Together, these observations (**Fig 2**) suggest that natural genetic variation
contributing to the single-cell morphological features we measured often acts via vertical
pleiotropy. In other words, correlations among morphological features that are present in
the absence of genetic variation underlie a large portion of pleiotropic genetic influences
on single cell morphology. Still, there are hints of another mechanism at play. Some QTL
tend to influence traits that are among the most weakly clustered in the correlation
network (**Fig 2F**). Moreover, most of the pleiotropic QTL we surveyed (24/35) each
influence at least 2 traits with $r_W < 0.2$. To investigate how often pleiotropy is not
predicted by the degree to which morphological features correlate in the absence of
genetic variation, in the next section we compare trait correlations present across clones
(r_W) to those present between genetically diverse strains (r_B).
419
Many traits are more strongly correlated across strains than they are across clones
When genetic changes that perturb one trait have collateral effects on another, we
expect the way traits correlate across genetically diverse strains to reflect trait
correlations across clones (*i.e.* $r_B = r_W$). When this condition is met, pleiotropy can be
viewed as an expected consequence of inherent relationships between traits, *i.e.* vertical
pleiotropy. On the other hand, if a QTL influences two traits that do not correlate across
clones, it may cause these traits to correlate across strains in which this QTL is
segregating. In this case, we expect r_B will be greater than r_W, suggesting horizontal
pleiotropy.
After correcting for multiple hypothesis testing, r_B significantly exceeds r_W in
24% of all trait pairs, and 41% of pairs in which at least one pleiotropic QTL influences
both traits (**Fig 3**; left panel; 41% of points are above the envelope, which represents a
Bonferroni corrected significance threshold of p < 0.01). In the majority of cases in which
r_B significantly differs from r_W, r_B is greater than r_W (**Fig 3**; left panel; 83% of points
outside the envelope are above it). The magnitude of the increase in r_B vs. r_W tends to
scale with the number of pleiotropic QTL that jointly influence both traits in a pair (**Fig
3**; left panel; colors get warmer farther above the envelope). These observations are
consistent with the hypothesis that QTL acting via horizontal pleiotropy increase r_B
relative to r_W.
However, horizontal pleiotropy is not the only reason traits may correlate
differently across strains versus across clones. We find significant deviations in r_B
relative to r_W in 14% of pairs for which no pleiotropic QTL influence both traits, (**Fig 3**;
right panel). This observation may suggest the presence of pleiotropic genetic variants
that we did not have statistical power to detect with an FDR of 5% in our QTL screen.
But an alternate explanation for the observed increases in r_B over r_W is that perhaps we
sometimes underestimate r_W.
One reason r_W could be underestimated is that single-cell measurements are
noisier than group-level averages. To test this possibility, we randomly assigned
individual cells to groups (pseudo-strains) having the same number of cells as the actual
strains, and found that in these permuted data, r_B and r_W never significantly differ (**Fig 3**;
insets). Because detection of r_W was not underpowered relative to r_B, we conclude that

measurement noise does not meaningfully obscure r_W. Another reason r_W could be
underestimated is if trait correlations across strains are more linear than those across
clones. To test this possibility, for every pair of traits we transformed the single-cell trait
measurements using a nonparametric model that finds their maximal correlation [70].
This transformation abrogated significant differences in r_B relative to r_W for fewer than
5% of affected trait pairs. Another reason r_W might be less than r_B is if there tends to be
less phenotypic variation within strains than between strains. Contrary to this prediction,
every morphological trait we surveyed varies more within strains than between strains. A
final reason r_W could be poorly estimated is if non-genetic heterogeneity across different
subpopulations within clonal populations causes variation in r_W. Therefore, next we
investigated whether the relationship between single-cell features varies for clonal cells
in different stages of the cell-division cycle.
463
Inferring a cell's progress through division from fixed cell images
Pairs of traits for which r_B is strong whereas r_W is not should reflect horizontal
pleiotropy, but a closer examination of some of these pairs revealed traits that should
correlate due to simple geometric constraints. For example, cell size and the width of the
bud neck should correlate due to the constraint that, even at its maximum, bud neck width
cannot be larger than the diameter of the cell. When measured in small-budded cells,
these two traits are correlated across yeast strains ($r_B = 0.40$) but are significantly less
correlated across clones ($r_W = 0.15$). Given the simple geometric constraint coupling the
width of the bud neck to the cell's size, why is there a discrepancy between r_B and r_W?
We reasoned that this discrepancy exists because the correlation between cell size and
neck width is disrupted during particular moments of cell division; *e.g.* the width of the
bud neck starts small even for large cells (**Fig 4A**; cell micrographs outlined in blue show
two cells in the progress of budding). If the relationship between morphological features
varies during cell division, r_W may represent a poor summary statistic.
How often does the relationship between morphological traits change during cell
division? Our single cell measurements are primed to address this question: we fixed
cells during exponential growth and imaged hundreds of thousands of single cells,
thereby capturing the full spectrum of morphologies as cells divide. A remaining
challenge is sorting these images according to progress through cell division, and then re-
measuring the correlation between morphological features within narrow windows along
that progression.
We performed this sorting using the Wishbone algorithm [71]. This algorithm
extracts developmental trajectories from high-dimensional phenotype data (typically
single-cell transcriptome data). We applied Wishbone separately to cells belonging to
each of the three cell types defined by morphometric analysis (unbudded, small-budded,
and large-budded cells). The trends describing how morphological features vary across
Wishbone-defined cell-division trajectories are consistent with previous observations of
how morphology changes in as yeast cells divide [72,73] (**Fig 4A**; line plots). For
example, Wishbone sorts fixed-cell images in such a way that cell area increases
throughout the course of cell division (**Fig4A**; upper left panel), and nuclear elongation
occurs just before nuclear division (**Fig4A**; lower left panel). These trajectories also
match our own observations of how morphological features change as live cells divide,
which we tracked by imaging at 1-minute intervals one of the 374 progeny strains that we

had engineered to express a fluorescently tagged nuclear protein (HTB2-GFP) (**Fig 4A**;
micrographs). We chose this particular strain because it does not deviate from the average
morphology of all 374 recombinants by more than one standard deviation for any of the
phenotypes we measure.
To further validate Wishbone's performance, we asked whether it could
reconstruct the time series of live-cell images from the HTB2-GFP strain. We obtained
time series for 78 single dividing cells, each imaged over at least 20 timepoints.
Quantifying morphological phenotypes from live-cell images in a high-throughput
fashion proved difficult because the morphometric software was optimized for fixed-cell
images and as cells grow and bud, the cells and their nuclei can move out of the focal
plane. Also, although we used short exposure times when imaging GFP fluorescence,
there are concerns about photo-toxicity and associated growth and morphology defects
[74]. For these reasons, we expect Wishbone to perform better on fixed-cell images than
on time series of live cell images. Still, Wishbone's cell-division trajectories recapitulate
the time course. When we align time series data across live cells by centering on each
cell's average predicted progress through division, Spearman's r is 0.65, 0.91, and 0.77
for time series corresponding to each of the three cell types (**Fig 4B**; see **Fig S5** for
recapitulation of 78 individual time series). These correlations are substantially higher
than those obtained by repeating the merging procedure after randomly permuting each
time series (corresponding Spearman's r of 0.42, 0.43. and 0.56). These observations
suggest that Wishbone is effective at properly assigning single-cell images to their
position in the cell cycle.
519
Cell cycle state can influence the relationship between morphological features
To identify cases where significant differences in r_B vs. r_W might result because
r_W is sensitive to cell-cycle state, we first assigned each imaged yeast cell from the QTL-
mapping population to one of 16 equal-sized bins based on Wishbone's estimation of
how far that cell had progressed through division. Because we did this separately for each
of the previously defined cell stages (unbudded, small-budded, and large-budded), this
additional binning finely partitions cell division into 48 (16 x 3) stages. To hold genotype
representation constant across each of the 48 bins, we performed binning separately for
each of the 374 mapping-family strains, then merged like bins across strains. We then
performed correlation partitioning on each bin separately.
Binning cells by cell-cycle state typically decreased the amount of phenotypic
variation per bin, which we expect in turn to obscure the correlation between traits.
Consider an extreme example: if there is no phenotypic variation remaining for a given
trait, it cannot covary with any other traits. Indeed, for most pairs of traits, the binning
procedure either decreases r_W or does not have a dramatic effect on it; decreases in r_W are
especially evident for trait pairs where variation of at least one of the traits shows a
relatively large decrease upon binning (**Fig 4C**). However, for some pairs of traits,
despite the decrease in phenotypic variation for at least one trait, the correlation between
traits improves upon binning. For example, binning by cell division increases the
correlation between cell size and the width of the bud neck (**Fig 4D**; leftmost plot) such
that it approaches r_B. This increased correlation is consistent with our hypothesis that the
process of cell division was obscuring the dependency of bud neck width on cell size.
Examining more pairs of traits for which binning tends to increase r_W (**Fig 4C**; red,
orange, and yellow points) reveals additional cases where the process of cell division

decouples traits that are otherwise correlated, and where binning reveals the underlying
correlation (**Fig 4D**; leftmost three plots).
Despite the evidence that cell asynchrony alters some trait correlations, many
cases remain where heterogeneity in cell-cycle state does not explain the observed
discrepancy between r_W and r_B (**Fig 4D**; rightmost three plots). We previously
demonstrated that r_B significantly exceeds r_W in 24% of all trait pairs (1389/5645) (**Fig
3).** For almost half of these pairs (689 pairs), binning by cell division does not resolve the
discrepancy between r_B and r_W to any extent; in other words, r_W does not increase in any
of the 16 bins. For an additional 193 pairs, binning by cell division resolves the
discrepancy by at most 5% in any bin. These results imply that cell-cycle heterogeneity
does not cause the discrepancy between r_W and r_B in the majority of cases, and that
elevation of r_B over r_W is best explained by QTL demonstrating horizontal pleiotropy.
556
Many QTL demonstrate horizontal pleiotropy
To test horizontal pleiotropy further, we focused on the 24 QTL each found to
influence at least 2 traits with $r_W < 0.2$ (pleiotropic QTL not included in **Fig 2B**). To test
whether these pleiotropic QTL cause increases in r_B relative to r_W, we divided our yeast
strains into sets in which a given QTL is not segregating, then re-measured the difference
between r_B and r_W. More specifically, for each QTL, we split the 374 phenotyped yeast
strains into two groups based on whether they inherited the wine or the oak parent's allele
at the genotyped marker closest to the estimated QTL location. Then we repeated
correlation partitioning on each subset of strains and compared the results to those
obtained from the complete set. For each QTL, we focused on trait pairs in which: (1)
both traits are affected by this QTL, and (2) r_B is significantly greater than r_W. Across all
such pairs, median r_B tends to decrease upon eliminating allelic variation at the marker
nearest the QTL (**Fig 5A**). No similar reduction in r_B is observed when we focus on pairs
of traits that are not affected by each QTL (**Fig 5A**) and no similar reduction is observed
in r_W (median reduction in r_W is 0.0001).
There appear to be two ways in which a QTL may affect r_B. In some cases,
eliminating genetic variation at the marker nearest a QTL decreases r_B in both resulting
subpopulations. Such cases are consistent with a straightforward scenario in which
horizontal pleiotropy results when a QTL that influences two or more traits (that are
otherwise weakly correlated) is segregating in a population (**Fig 5B**; top row). In other
cases, eliminating allelic variation at a QTL site decreases r_B in only one of the two
resulting subpopulations (*i.e.* the subpopulation possessing either the oak or the wine
allele). This observation demonstrates that horizontal pleiotropy can emerge by virtue of
a QTL allele strengthening a correlation between two traits so that genetic variation
affecting one trait is more likely to affect the other when that allele is present [76,77] (**Fig
5B;** bottom row).
How many cases where r_B significantly exceeds r_W can be explained, to some
extent, by horizontal pleiotropy (*i.e.* a QTL increasing the between-genotype
correlation)? For every trait pair where r_B significantly exceeds r_W and at least one QTL
influences both traits in the pair (1153 pairs total), eliminating allelic variation at the
marker nearest at least one of the shared QTL causes r_B to decrease in one or both of the
resulting subpopulations (**Fig 5C**: solid black line in rightmost plot). About 60% of these
decreases affect both subpopulations (*e.g.* **Fig 5B**; top row) and 40% affect only one

subpopulation (*e.g.* **Fig 5B**; bottom row). These decreases in r_B appear to resolve the
discrepancies in r_B vs. r_W more often and to a greater extent than does accounting for
cell-cycle heterogeneity (**Fig 5C**; leftmost plot). Some QTL have larger impacts on r_B
than do others (**Fig 5C**). Eliminating allelic variation near a QTL on chromosome 13
decreases r_B in the largest number of traits pairs (681). Subtracting the influence of a
QTL on chromosome 15 decreases r_B to the greatest extent; the average decrease across
357 affected trait pairs is 0.07. Together these observations suggest: (1) many QTL
demonstrate horizontal pleiotropy (**Fig 5A**), (2) there are at least two ways for horizontal
pleiotropy to emerge (**Fig 5B**), and (3) horizontal pleiotropy is a major factor driving
increases in r_B over r_W in this study (**Fig 5C**).
600
Spontaneous mutations alter the relationships between morphological features
Our finding that some QTL alleles appear to strengthen correlations between
otherwise weakly correlated traits (**Fig 5B**; lower panel) lends credence to the idea that
the relationships between phenotypes, and thus the extent of phenotypic modularity (or
integration), are mutable traits [78]. This finding has implications for evolutionary
medicine, in particular evolutionary traps, *e.g.* strategies to contain microbial populations
by encouraging them to evolve resistance to one treatment so that they become
susceptible to another [39-41]. These traps will fail if targeted correlations can be broken
by mutations. To test whether spontaneous mutations can alter trait correlations, we
analyzed the cell-morphology phenotypes of a collection of yeast mutation-accumulation
(MA) lines [54]. These MA lines were derived from repeated passaging through
bottlenecks, which dramatically reduced the efficiency of selection and thereby allowed
retention of the natural spectrum of mutations irrespective of effect on fitness [55]. We
previously imaged these lines in high throughput (>1000 clonal cells imaged per each of
94 lines) [50].
Because MA lines contain private mutations unique to each strain, they are not
amenable to QTL mapping and between-strain trait correlations have less meaning.
Instead, we focused on within-strain correlations, which we expected to be consistent
across strains because of the limited number of mutations distinguishing the strains (an
average of 4 single-nucleotide mutations per line [55]), except if a rare mutation does
indeed alter the correlation. To determine if such correlation-altering mutations exist, we
calculated within-strain correlations for each strain separately and asked, for each trait
pair, whether any strains had extreme correlations relative to the other strains. For most
trait pairs, the MA lines trait correlations did not vary much from each other or from that
of the ancestor strain (**Fig 6**). However, in several instances, we observed a trait-pair
correlation dramatically outside the range of the other trait pairs and more than four
standard deviations from the mean (**Fig 6A**). Some mutations appear to influence many
trait-trait relationships (mutations found in blue- and purple-colored strains in **Fig 6B &**
C), whereas others influence fewer (mutations found in magenta-colored strain in **Fig**
6C).
Given that in this small sampling of spontaneous mutations, we found several that
appear to alter the relationship between morphological features, we think such mutations
are common enough to merit further consideration in evolutionary models. The mutations
in the outlier lines provide candidate correlation-altering mutations for future mechanistic
studies as well.
636

Discussion:

Although evolutionary biologists and medical geneticists alike appreciate that organismal traits can rarely be understood in isolation, the extent and implications of pleiotropy have remained difficult to assess. One approach to measuring pleiotropy has been to count phenotypes influenced by individual genetic loci [17,33,34]. For example, the median number of skeletal traits affected per QTL in a mouse cross was six (out of 70 traits measured); this small median fraction of traits suggests that variation in skeletal morphology is modular [16,30]. Of course, for a count of traits to be meaningful the full trait list must be comprehensive, and correlations between traits must be properly accounted for [17,33,34]. We aimed for comprehensiveness in a very similar way to the studies of mouse skeletal traits, by systematic phenotyping of a large number of morphological traits. However, we addressed the need for a principled approach to separating inherent trait correlations from those induced by genetic differences in a new way: by extending the analysis to include within-genotype correlations and thereby enabling an operational definition of the distinction between vertical and horizontal pleiotropy.

Our comprehensive analysis of how thousands of trait pairs co-vary within and between mapping strains yields an unprecedently quantitative and nuanced view of pleiotropy. We found support for modularity, not only in the striking correspondence between our median number of traits affected per QTL (six out of 167) and that found for mouse skeletal traits [16,30], but also in the way that within-genotype correlations grouped traits into relatively isolated clusters (**Fig 2**). We also found ample evidence of horizontal pleiotropy layered on top of that modularity, with many cases of between-genotype trait correlations that exceeded within-genotype correlations (**Fig 3**). Our results do not speak directly to whether modularity results from selection against pleiotropy in nature, because we sampled only two natural genetic backgrounds (wine and oak). However, future work comparing MA lines to a larger collection of natural isolates might help answer questions about the extent to which selection purges pleiotropic mutations.

Our partitioning of between-strain (genetic) and within-strain (environmental) correlations relates to another approach to understanding trait interdependencies, the estimation of the so-called **G** matrix. This genetic variance-covariance matrix summarizes the joint pattern of heritable variation in a population of the traits that compose its rows and columns, and is central to understanding how trait correlations constrain evolution. The **G** matrix arises in the multivariate breeder's equation, which describes the responses to selection of correlated traits [79]. If breeding is the goal, the distinction between vertical and horizontal pleiotropy is not so important, because both can impede selection. Indeed, any philosophical concern about what constitutes a biologically meaningful trait is irrelevant to the breeder, who actually cares about particular traits (*e.g.*, milk yield and fat content).

G matrices are not only relevant to breeders, but to evolutionary biologists as well, and it is worthwhile to place our results into this context. A major evolutionary question in the **G**-matrix literature is whether the **G** matrix itself can evolve. In other words, do short-term responses to selection (as captured in the breeder's equation) predict long-term responses or do constraints shift through time, perhaps in a way that facilitates (or is part of) adaptation [80]? Our results with MA lines add to evidence that the **G** matrix readily changes [81], in that individual mutations have major effects on particular

trait correlations (*e.g.* **Fig 6A**). Our QTL-mapping results also support this view, in that
some cases of horizontal pleiotropy appear to be caused by alleles that alter trait
correlations (*e.g.* **Fig 5B**; bottom panel).
Another prominent question in the **G**-matrix literature is the extent to which the **P**
matrix, which includes all sources of phenotypic variation and covariation, predicts the **G**
matrix, which only includes additive genetic effects (*i.e.*, those that respond to selection).
If **P** predicts **G** well, as proposed by Cheverud [82], then inference of selection responses
from patterns of trait covariation in a population would suffice when genetic analysis
would be difficult or costly. Our results do not speak directly to this question, because we
did not estimate **G** itself and instead estimated genetic correlations that include non-
additive effects. However, our results are informative from another angle, which is the
comparison of genetic and environmental correlations. As we showed (**Fig 3**), although
there are cases in which the environmental and genetic correlations have different signs,
the environmental correlations do tend to match the signs of the genetic correlations and
predict their magnitudes to some extent as well, consistent with similarity between **P** and
G. Future experiments using clones embedded in a more complicated crossing scheme
could properly partition **P** into **G**, **E**, and the non-additive genetic components, to address
Cheverud's conjecture [82] more directly. There are only a few reports of comparisons of
E matrices [83], but we encourage increased attention to the **E** matrix to understand
inherent trait correlations and to contextualize **G** in a way that diminishes concerns about
which traits have been granted status as its rows and columns.
A major conclusion of our work is that context is crucial. We have shown that
trait correlations change through the cell-division cycle and in different genetic
backgrounds. It is likely that macroenvironmental differences alter trait correlations as
well [84]. These results support the idea that predicting the mapping from genotype to
phenotype requires a paradigm shift [85], away from merely mapping the relationships
between traits and toward unfurling the range of contexts across which those
relationships persist. Future work in this direction will not only advance understanding of
the evolution of complex traits, but will have practical benefits. For example, our
approach demonstrates a potentially fruitful way to consider the design of evolutionary
traps: using within-genotype correlations to identify particularly immutable inherent
correlations between traits.
715

Acknowledgments:
We are grateful to Dmitri Petrov and Grant Kinsler for helpful discussions about
pleiotropy and comments on an earlier version of this manuscript. We also thank Barak
Cohen and David Hall for providing strains used in this study. This work was supported
by National Institutes of Health grant R35GM118170 (to MLS), National Institutes of
Health fellowship F32GM103166 (to KGS), a New York University Graduate School of
Arts and Science Dean's Dissertation Fellowship (to SL), and National Institutes of
Health grant R35GM119744 (to ABP). The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.
725

Materials and Methods:
Measuring the morphology of single yeast cells

Recombinant yeast strains were generated and genotyped at 225 markers in a previous
study [53,59]; each strain is a homothallic diploid. We prepared yeast cells from these
strains for microscopy using published methods [49-51,86]. Briefly, yeast strains were
grown in minimal media with 0.08% glucose in 96-well plates [87], harvested during
exponential phase, fixed in 4% paraformaldehyde, stained for cell-surface manno-protein
(with FITC-concanavalin A) and nuclear DNA (with DAPI), sonicated, mounted on 96-
well glass-bottom microscopy plates, and imaged with a Nikon Eclipse TE-2000E
epifluorescence automated microscope using a 40× objective and appropriate
fluorescence filters. Three biological replicate experiments were performed, typically
yielding a total of between 500 to 1,000 imaged cells per strain (**Fig S2**).
738
Statistical analysis and processing of cell image data
Cell image processing was performed similarly to previous studies [49-51,86]. Imaged
cells were analyzed for quantitative morphological traits using the CalMorph software
package [52], which reports on hundreds of morphological features that are each specific
to one of three cell types: unbudded, small-budded, and large-budded cells. We excluded
phenotypes for which >10% of cells had missing values, leaving 167 morphological
features. Any cell that was not scored for all features pertaining to its type was
eliminated. Each morphological trait was transformed via a Box-Cox transformation of
the raw data with the value of lambda that makes the residuals of a linear regression of
phenotype on strain most normal using the EnvStats package in R [88]. Internal controls
(several wells representing the wine and oak parents) were present on every 96-well plate
and were used to correct for effects on phenotypic variation that resulted from differences
among replicate experiments, such as differences in the brightness of the cell stain. We
calculated the mid-parent value for each phenotype on every plate, then calculated the
average mid-parent value across all plates. For each phenotype, we found the difference
between the plate-specific mid-parent value and the average mid-parent value across all
plates. Then we subtracted this difference from each plate for the corresponding
phenotype. After correction, any cell with a morphological feature that deviated from the
average by more than 5 standard deviations was then eliminated, as investigation of such
cells typically revealed these were CalMorph miscalls or cellular debris.
759
QTL mapping
QTL interval mapping was performed similarly to previous studies [62] using the R/qtl
package [63]. We performed a QTL scan using the function "scanone", which finds at
most one QTL per chromosome. Yeast strains, which are homozygous diploids, were
modeled as haploids and QTL models were fit using Haley-Knott regression. As in
previous work [62], when comparing QTL across traits, QTL within 30 cM on the same
chromosome were counted as the same QTL. In some cases, we detected a QTL in
between two others on the same chromosome and within 30 cM of both. In these cases,
we made decisions about the total number of unique QTL present by using our best
judgement and considering factors such as the proximity between QTL. A summary of all
significant QTL effects, including their chromosomal locations in cM and which QTL on
the same chromosome we considered unique, is provided in **Table S1** (also see **Fig 1A**).
QTL effects were counted as significant when they were stronger than any QTL effect

detected in 100 randomly permutated datasets, allowing for 5% false positives.
Permutations were performed separately for each trait.
775

Candidate gene swaps
All yeast transformations were performed using the lithium acetate [89] and *delitto*
perfetto [64] methods. For each candidate gene, the gene was first deleted from haploid
variants of both the wine and oak parental strains and replaced with a selectable marker,
the yeast gene encoding orotidine-5'-phosphate decarboxylase (*URA3*). Gene knockouts
were confirmed by growth on plates lacking uracil and DNA sequencing of the affected
region. Next, the *URA3* selectable marker was replaced with the other parent's version of
the candidate gene. These candidate gene 'swaps' were selected by growth on 5-
Fluoroorotic acid and confirmed by sequencing of the affected region. For each candidate
gene, we swapped a region containing the coding sequence plus 5 – 750 bp up and
downstream. We used the following regions of homology to define the boundaries of
each swapped segment:
788

~300bp upstream of *PXL1*: TTATAATTGTGGTTTAGCGTTTCATAGTCGC
~300bp downstream of *PXL1*: CCTTATTCTCTATTCTTAGGCTCCTGTTCC
~5bp upstream of *HOF1*: GAAAGAATGAGCTACAGTTATGAAGCTTG
~ 300bp downstream of *HOF1*: GTATTCGTAACAAGTGACTCTAATGATAT
~ 750bp upstream of *RAS1*: CGACTAAAGGAATTATACCATCATGCATC
~ 300bp downstream of *RAS1*: GCATTCTAAAAACAGAGCTTTTGCCG
795

These regions of homology were chosen by searching for regions of higher GC content
nearby the start and end of each gene's coding sequence. In addition, we attempted to
swap the wine and oak parents' versions of the *GPA1* gene on chromosome 8. Despite
trying various regions of homology, we could not successfully replace *GPA1* with the
URA3 selectable marker in the oak parent. *GPA1* is known to be essential in some genetic
backgrounds [90].
802

Though the recombinant strains we studied are homothallic diploids, the strains in **Fig 1B**
(both the parental strains and the strains possessing the gene swaps) are haploid. Because
the analyses in **Fig 1B** compare pairs of strains (*e.g.* the oak haploid parent to the wine
haploid parent, or the wine haploid parent to the wine haploid parent possessing the oak
allele of *PXL1*), we only considered experiments where both strains in the pair were
imaged in the same replicate experiment. To account for differences among replicate
experiments, for each phenotype, we subtracted the value in one strain from the value in
the other to calculate the phenotypic difference between strains in that replicate
experiment; the reported value is the average of these differences across replicate
experiments (**Table S2, Fig 1B**).
813

Calculation of correlation coefficients
We used WABA II as implemented in the multilevel package in R [69] to
calculate cell-level (r_W) and strain-level (r_B) Pearson correlation coefficients for each pair
of traits. We used an r-to-z transformation to determine whether differences in r_B vs. r_W
are significant at a Bonferroni corrected p-value of 0.01 (this is a z-score cutoff of 4.63,

given 5645 pairs of traits were tested). To assess whether correlations across single cells
generally result in different values than correlations across group-level averages, we
assigned yeast cells to groups (pseudo-strains) randomly, maintaining the same number
of cells per strain as in the actual data. To assess whether results would differ if we
allowed for non-linear correlations, we transformed the single-cell data using a
nonparametric model that finds the fixed point of maximal correlation, implemented in
the R package acepack [70]. To assess whether results from WABA differed from those
obtained using a standard quantitative genetics model (**Fig S3**), we implemented the latter
using the nlme package in R [91] to specify a mixed-effects model with cells nested
within strains. We specified a covariance structure that allows covariance between two
traits but no covariance between cells or between strains. We used this model to calculate
the environmental and genetic correlations for 350 pairs of randomly chosen traits.
831
Live imaging single cells as they divide
For live imaging the morphology of dividing yeast cells, we chose one of the
recombinant yeast strains, F2_292. This strain was chosen because it does not deviate
from the average morphology of all 374 recombinants by more than one standard
deviation for any of the phenotypes we measured. F2_292 was transformed to express a
fusion protein of GFP and a nuclear protein (histone H2B encoded by *HTB2*). Two
independent transformants were imaged in the GFP channel (for nuclei) and in brightfield
(for cell outlines). We prepared live cells for imaging following published methods
[87,92,93], in a similar way to that described above, except cells were neither fixed nor
stained. Cells were taken during mid-log phase growth, seeded in 96-well glass bottom
microscopy plates containing minimal media with 0.08% glucose, and imaged over a
period of 3 hours. In each of four replicate experiments, cells were imaged either every
minute, every 90 seconds, or every 2 minutes. We used short exposure times (afforded by
the highly abundant HTB2-GFP) and took only a single image per well per timepoint to
reduce photo-toxicity. We processed images with CalMorph then matched cells across
timepoints by their centroid locations in the imaging fields. Overall we obtained time
series for 78 cells that each: (1) were longer than 20 timepoints, (2) contained no gaps
where the cell was not phenotyped for many consecutive timepoints, and (3) contained no
images that appeared to be very out of focus potentially resulting in misestimation of
phenotype values. Because CalMorph divides cells into unbudded, small-budded and
large-budded stages, these 78 time series are also divided this way (11, 23, and 44 cells,
respectively).
854
We used the Wishbone algorithm implemented in python [71] to estimate progression
through the cell-division cycle. Wishbone recapitulates each of these 78 time series (**Fig**
S5) with Spearman correlations between the actual and inferred image orders that average
0.42, 0.85, 0.40 across all unbudded, small-budded or large-budded series, respectively.
The lower correlations between Wishbone's predicted progress through division and time
for the unbudded and large-budded cells may result because each time series captured
only a part of the cell-division cycle and, during some stretches in the cycle, there are
fewer morphological changes taking place. To estimate Wishbone's accuracy across a
longer stretch of time, we merged the Wishbone predictions within the classes of
unbudded, small-budded or large-budded cell time series. To do so, we had to contend

with the fact that the first timepoint for each imaged cell often represents a different
moment in division. For example, some time series for unbudded cells start from an
image that is already far along the division process (**Fig S5**; values close to 1 on the
vertical axis) while others start from a cell image that has just begun its division cycle
(**Fig S5**; values close to zero on the vertical axis). Therefore, we aligned the time series
by subtracting from each the difference between Wishbone's estimate of the average
percent progress through division and the average time elapsed.
872
Note that, because this merging procedure utilized information from Wishbone, it
imposes a correlation between time and Wishbone's estimated progress through division.
To reduce the impact of this induced correlation, we eliminated the cell images in the
middle of each time series, which represent the images that are most affected by this
induced correlation. Eliminating 25% or 50% of cell images in this way reduced the
correlations by at most 0.05, suggesting these correlations are not driven by our merging
procedure.
880
Assigning cells to a bin based on progression through cell division
We used Wishbone to estimate how far each fixed-cell image had progressed through cell
division. Wishbone software requires input about which "start" cell has features
resembling those present at the start of the cell cycle. To identify such features, we used
the data from the live-imaged cell time series. We plotted how single-cell features change
over the course of live imaging, and chose several features that correlate best with
progress through cell division (*e.g.* cell size, bud size, location of the nucleus).
888
Using Wishbone's estimation of how far each fixed cell had progressed through division,
we assigned each cell to one of 16 equal-sized bins. We did this separately for each of the
374 yeast strains, then merged like bins across strains, such that genetic diversity was
constant across each of the final 16 bins. We obtained very similar results to those
reported in **Figs 4C**, **4D**, and **5C** when we used 8 instead of 16 bins. The names of the
traits plotted in **Fig 4** represent succinct summaries of single-cell morphologies
quantified using CalMorph [52]. For fuller descriptions of these traits, see the following
trait designations in the CalMorph software manual: **Fig 4A** upper left: C11.1 in
unbudded cells, C101 in budded cells; **Fig 4A** lower left: D184 in small-budded cells,
D182 in unbudded and large-budded cells; **Fig 4A** upper right: C12.2; **Fig 4A** lower
right: D116; **Fig 4C** upper left: C101 and C109 in small-budded cells; **Fig 4C** upper
middle: C11.2 and D132 in small-budded cells; **Fig 4C** upper right: C105 and C113 in
small-budded cells; **Fig 4C** lower left: C114 and D145 in large-budded cells; **Fig 4C**
lower middle: C109 and C126 in large-budded cells; **Fig 4C** lower right: D14.2 and D169
in large-budded cells.
904
Eliminating genetic variation at the marker nearest a QTL
For each of the 24 QTL suspected of horizontal pleiotropy (*i.e.* pleiotropic QTL not in
Fig 2B), we divided the 374 phenotyped yeast strains into two groups based on whether
they inherited the wine or the oak parent's allele at the genotyped marker closest to the
QTL. We then performed correlation partitioning separately for each group of strains.
The names of the traits plotted in **Fig 5B** represent succinct summaries of single-cell

morphologies quantified using CalMorph. For fuller descriptions of these traits, see the following trait designations in the CalMorph software manual: upper: D128 and C114 in large-budded cells; lower: D197 and D17.1 in large-budded cells.

Quantifying trait correlations within each MA line
We used MA line data from our previous study [50]. Fewer traits were analyzed in that study than in the current study, such that there were only 3731 pairs of traits to survey, as opposed to 5645 in the QTL-mapping family. We calculated Pearson correlations between every pair of traits, separately within each MA line. The names of the traits plotted in **Fig 6A** represent succinct summaries of single-cell morphologies quantified using CalMorph. For fuller descriptions of these traits, see the following trait designations in the CalMorph software manual: upper left: D185 and D186 in large-budded cells; upper right: C102 and D132 in small-budded cells; lower left: C108 and D167 in large-budded cells; lower right: D135 and D169 in large-budded cells.

References:

1.	Plate L. Genetics and evolution. Fischer J, editor. Fest schrift zum sechzigsten Geburtstag Richard Hertwigs; 1910. pp. 536–610.

2.	Stearns FW. One Hundred Years of Pleiotropy: A Retrospective. Genetics. Genetics; 2010;186: 767–773. doi:10.1534/genetics.110.122549

3.	Simons YB, Bullaughey K, Hudson RR, Sella G. A population genetic interpretation of GWAS findings for human quantitative traits. Gibson G, editor. PLoS Biol. 2018;16: e2002985. doi:10.1371/journal.pbio.2002985

4.	Tyler AL, Asselbergs FW, Williams SM, Moore JH. Shadows of complexity: what biological networks reveal about epistasis and pleiotropy. Bioessays. WILEY-VCH Verlag; 2009;31: 220–227. doi:10.1002/bies.200800022

5.	Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, et al. 10 Years of GWAS Discovery: Biology, Function, and Translation. Am J Hum Genet. 2017;101: 5–22. doi:10.1016/j.ajhg.2017.06.005

6.	Visscher PM, Yang J. A plethora of pleiotropy across complex traits. Nat Genet. 2016;48: 707–708. doi:10.1038/ng.3604

7.	Chesmore K, Bartlett J, Williams SM. The ubiquity of pleiotropy in human disease. Hum Genet. 137: 39–44.

8.	Sivakumaran S, Agakov F, Theodoratou E, Prendergast JG, Zgaga L, Manolio T, et al. Abundant pleiotropy in human complex diseases and traits. Am J Hum Genet. Elsevier; 2011;89: 607–618. doi:10.1016/j.ajhg.2011.10.004

9.   White JK, Gerdin A-K, Karp NA, Ryder E, Buljan M, Bussell JN, et al. Genome-
wide generation and systematic phenotyping of knockout mice reveals new roles
for many genes. Cell. Elsevier; 2013;154: 452–464. doi:10.1016/j.cell.2013.06.022

10. Wang Z, Liao B-Y, Zhang J. Genomic patterns of pleiotropy and the evolution of
complexity. Proc Natl Acad Sci USA. National Acad Sciences; 2010;107: 18034–
18039. doi:10.1073/pnas.1004666107

11. Housman G, Byler S, Heerboth S, Lapinska K, Longacre M, Snyder N, et al. Drug
resistance in cancer: an overview. Cancers (Basel). Multidisciplinary Digital
Publishing Institute; 2014;6: 1769–1792. doi:10.3390/cancers6031769

12. Rockman MV. The QTN program and the alleles that matter for evolution: all
that's gold does not glitter. Evolution. 2012;66: 1–17. doi:10.1111/j.1558-
5646.2011.01486.x

13. Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. Cell. 2017;169: 1177–1186.

14. Fisher RA. The Genetical Theory of Selection. Claredon. 1930.

15. Allen Orr H. Adaptation and the cost of complexity. Evolution. 2000;54: 13.
doi:10.1554/0014-3820(2000)054[0013:AATCOC]2.0.CO;2

16. Wagner GP, Kenney-Hunt JP, Pavlicev M, Peck JR, Waxman D, Cheverud JM.
Pleiotropic scaling of gene effects and the 'cost of complexity'. Nature. Nature
Publishing Group; 2008;452: 470–472. doi:10.1038/nature06756

17. Wagner GP, Zhang J. The pleiotropic structure of the genotype–phenotype map:
the evolvability of complex organisms. Nat Rev Genet. Nature Publishing Group;
2011;12: 204–213. doi:10.1038/nrg2949

18. McGee LW, Sackman AM, Morrison AJ, Pierce J, Anisman J, Rokyta DR.
Synergistic Pleiotropy Overrides the Costs of Complexity in Viral Adaptation.
Genetics. 2016;202: 285–295. doi:10.1534/genetics.115.181628

19. Wittkopp PJ, Haerum BK, Clark AG. Regulatory changes underlying expression
differences within and between Drosophila species. Nat Genet. Nature Publishing
Group; 2008;40: 346–350. doi:10.1038/ng.77

20. Stern DL. Perspective: evolutionary developmental biology and the problem of
variation. Evolution. 2009;54: 1079. doi:10.1554/0014-
3820(2000)054[1079:PEDBAT]2.0.CO;2

21. He X, Zhang J. Toward a Molecular Understanding of Pleiotropy. Genetics.
Genetics; 2006;173: 1885–1891. doi:10.1534/genetics.106.060269

22.   Papakostas S, Vøllestad LA, Bruneaux M, Aykanat T, Vanoverbeke J, Ning M, et
al. Gene pleiotropy constrains gene expression changes in fish adapted to different
thermal conditions. Nat Commun. Nature Publishing Group; 2014;5: 4071.
doi:10.1038/ncomms5071

23.   Altenberg L. Modularity in Evolution: Some Low-Level Questions. In: Rasskin-
Gutman D, Callebaut W, editors. Modularity. 2005.

24.   Wagner GP, Altenberg L. Perspective: Complex Adaptations and the Evolution of
Evolvability. Evolution. 1996;50: 967. doi:10.2307/2410639

25.   Welch JJ, Waxman D. Modularity and the cost of complexity. Evolution. 2003;57:
1723–1734.

26.   Wagner GP, Pavlicev M, Cheverud JM. The road to modularity. Nat Rev Genet.
Nature Publishing Group; 2007;8: 921–931. doi:10.1038/nrg2267

27.   Melo D, Porto A, Cheverud JM, Marroig G. Modularity: genes, development and
evolution. Annu Rev Ecol Evol Syst. Annual Reviews; 2016;47: 463–486.
doi:10.1146/annurev-ecolsys-121415-032409

28.   Collet JM, McGuigan K, Allen SL, Chenoweth SF, Blows MW. Mutational
Pleiotropy and the Strength of Stabilizing Selection Within and Between
Functional Modules of Gene Expression. Genetics. 2018;208: 1601–1616.
doi:10.1534/genetics.118.300776

29.   Paaby AB, Rockman MV. Pleiotropy: what do you mean? Reply to Zhang and
Wagner. Trends Genet. Elsevier; 2013;29: 384. doi:10.1016/j.tig.2013.05.003

30.   Wagner GP, Zhang J. Universal pleiotropy is not a valid null hypothesis: reply to
Hill and Zhang. Nat Rev Genet. Nature Publishing Group; 2012;13: 296–296.
doi:10.1038/nrg2949-c2

31.   Hill WG, Zhang X-S. Assessing pleiotropy and its evolutionary consequences:
pleiotropy is not necessarily limited, nor need it hinder the evolution of
complexity. Nat Rev Genet. Nature Publishing Group; 2012;13: 296–296.
doi:10.1038/nrg2949-c1

32.   Zhang J, Wagner GP. On the definition and measurement of pleiotropy. Trends
Genet. 2013.

33.   Gibson G. Decanalization and the origin of complex disease. Nat Rev Genet.
2009;10: 134–140. doi:10.1038/nrg2502

34.   Nuzhdin SV, Friesen ML, McIntyre LM. Genotype-phenotype mapping in a post-
GWAS world. Trends Genet. 2012;28: 421–426. doi:10.1016/j.tig.2012.06.003

35.   Li Y, Venkataram S, agarwala A, dunn B, Petrov DA, Sherlock G, et al. Hidden
1017   Complexity of Yeast Adaptation under Simple Evolutionary Conditions. Curr Biol.
1018   2018;28: 515–525.e6. doi:10.1016/j.cub.2018.01.009

1019   36.   Gorter FA, Aarts MGM, Zwaan BJ, de Visser JAGM. Local Fitness Landscapes
1020   Predict Yeast Evolutionary Dynamics in Directionally Changing Environments.
1021   Genetics. 2018;208: 307–322. doi:10.1534/genetics.117.300519

1022   37.   Baym M, Stone LK, Kishony R. Multidrug evolutionary strategies to reverse
1023   antibiotic resistance. Science. 2016;351: aad3292–aad3292.
1024   doi:10.1126/science.aad3292

1025   38.   Hartley SW, Monti S, Liu C-T, Steinberg MH, Sebastiani P. Bayesian methods for
1026   multivariate modeling of pleiotropic SNP associations and genetic risk prediction.
1027   Front Genet. Frontiers; 2012;3: 176. doi:10.3389/fgene.2012.00176

1028   39.   Chen G, Mulla WA, Kucharavy A, Tsai H-J, Rubinstein B, Conkright J, et al.
1029   Targeting the adaptability of heterogeneous aneuploids. Cell. Elsevier; 2015;160:
1030   771–784. doi:10.1016/j.cell.2015.01.026

1031   40.   Nichol D, Jeavons P, Fletcher AG, Bonomo RA, Maini PK, Paul JL, et al. Steering
1032   Evolution with Sequential Therapy to Prevent the Emergence of Bacterial
1033   Antibiotic Resistance. PLoS Comput Biol. 2015;11: e1004493.
1034   doi:10.1371/journal.pcbi.1004493

1035   41.   Kaznatcheev A, Peacock J, Basanta D, Marusyk A, Scott JG. Fibroblasts and
1036   alectinib switch the evolutionary games played by non-small cell lung cancer. Nat
1037   Ecol Evol. 2019;3: 450–456. doi:10.1038/s41559-018-0768-z

1038   42.   Paaby AB, Rockman MV. The many faces of pleiotropy. Trends Genet. Elsevier;
1039   2013;29: 66–73. doi:10.1016/j.tig.2012.10.010

1040   43.   Walsh B, Blows MW. Abundant Genetic Variation + Strong Selection =
1041   Multivariate Genetic Constraints: A Geometric View of Adaptation.
1042   http://dxdoiorg/101146/annurevecolsys110308120232. Annual Reviews; 2009;40:
1043   41–59. doi:10.1146/annurev.ecolsys.110308.120232

1044   44.   Ference BA, Ginsberg HN, Graham I, Ray KK, Packard CJ, Bruckert E, et al.
1045   Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence
1046   from genetic, epidemiologic, and clinical studies. A consensus statement from the
1047   European Atherosclerosis Society Consensus Panel. Eur Heart J. 2017;38: 2459–
1048   2472. doi:10.1093/eurheartj/ehx144

1049   45.   Hafid Al N, Christodoulou J. Phenylketonuria: a review of current and future
1050   treatments. Transl Pediatr. 2015;4: 304–317. doi:10.3978/j.issn.2224-
1051   4336.2015.10.07

46.   Powell ALT, Nguyen CV, Hill T, Cheng KL, Figueroa-Balderas R, Aktas H, et al.
Uniform ripening encodes a Golden 2-like transcription factor regulating tomato
fruit chloroplast development. Science. 2012;336: 1711–1715.
doi:10.1126/science.1222218

47.   Grüneberg H. An analysis of the "pleiotropic" effects of a new lethal mutation in
the rat (Mus norvegicus). 1938. doi:10.2307/82266

48.   Ohya Y, Sese J, Yukawa M, Sano F, Nakatani Y, Saito TL, et al. High-
dimensional and large-scale phenotyping of yeast mutants. Proceedings of the
National Academy of Sciences. 2005;102: 19015–19020.
doi:10.1073/pnas.0509436102

49.   Bauer CR, Li S, Siegal ML. Essential gene disruptions reveal complex
relationships between phenotypic robustness, pleiotropy, and fitness. Mol Syst
Biol. 2015;11: 773–773. doi:10.15252/msb.20145264

50.   Geiler-Samerotte KA, Zhu YO, Goulet BE, Hall DW, Siegal ML. Selection
Transforms the Landscape of Genetic Variation Interacting with Hsp90. PLoS
Biol. 2016;14: e2000465. doi:10.1371/journal.pbio.2000465

51.   Richardson JB, Uppendahl LD, Traficante MK, Levy SF, Siegal ML. Histone
variant HTZ1 shows extensive epistasis with, but does not increase robustness to,
new mutations. PLoS Genet. 2013;9: e1003733. doi:10.1371/journal.pgen.1003733

52.   Negishi T, Nogami S, Ohya Y. Multidimensional quantification of subcellular
morphology of Saccharomyces cerevisiae using CalMorph, the high-throughput
image-processing program. J Biotechnol. 2009;141: 109–117.
doi:10.1016/j.jbiotec.2009.03.014

53.   Gerke J, Lorenz K, Cohen B. Genetic interactions between transcription factors
cause natural variation in yeast. Science. 2009;323: 498–501.
doi:10.1126/science.1166426

54.   Hall DW, Mahmoudizad R, Hurd AW, Joseph SB. Spontaneous mutations in
diploid Saccharomyces cerevisiae: another thousand cell generations. Genet Res
2008;90: 229–241. doi:10.1017/S0016672308009324

55.   Zhu YO, Siegal ML, Hall DW, Petrov DA. Precise estimates of mutation rate and
spectrum in yeast. Proc Natl Acad Sci USA. 2014;111: E2310–8.
doi:10.1073/pnas.1323011111

56.   Cooper TF, Ostrowski EA, Travisano M. A negative relationship between
mutation pleiotropy and fitness effect in yeast. Evolution. 2007;61: 1495–1499.
doi:10.1111/j.1558-5646.2007.00109.x

57.   Yang M, Ohnuki S, Ohya Y. Unveiling nonessential gene deletions that confer
1088   significant morphological phenotypes beyond natural yeast strains. BMC
1089   Genomics. 2014;15: 932. doi:10.1186/1471-2164-15-932

1090   58.   Sahai E. Mechanisms of cancer cell invasion. Curr Opin Genet Dev. 2005;15: 87–
1091   96. doi:10.1016/j.gde.2004.12.002

1092   59.   Gerke JP, Chen CTL, Cohen BA. Natural isolates of Saccharomyces cerevisiae
1093   display complex genetic variation in sporulation efficiency. Genetics. 2006;174:
1094   985–997. doi:10.1534/genetics.106.058453

1095   60.   Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, et al. Population
1096   genomics of domestic and wild yeasts. Nature. 2009;458: 337–341.
1097   doi:10.1038/nature07743

1098   61.   Skelly DA, Merrihew GE, Riffle M, Connelly CF, Kerr EO, Johansson M, et al.
1099   Integrative phenomics reveals insight into the structure of phenotypic diversity in
1100   budding yeast. Genome Res. 2013;23: 1496–1504. doi:10.1101/gr.155762.113

1101   62.   Ziv N, Shuster BM, Siegal ML, Gresham D. Resolving the Complex Genetic Basis
1102   of Phenotypic Variation and Variability of Cellular Growth. Genetics. 2017;
1103   genetics.116.195180. doi:10.1534/genetics.116.195180

1104   63.   Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental
1105   crosses. Bioinformatics. 2003;19: 889–890.

1106   64.   Stuckey S, Storici F. Gene knockouts, in vivo site-directed mutagenesis and other
1107   modifications using the delitto perfetto system in Saccharomyces cerevisiae. Meth
1108   Enzymol. 2013;533: 103–131. doi:10.1016/B978-0-12-420067-8.00008-8

1109   65.   Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, et al.
1110   Saccharomyces Genome Database: the genomics resource of budding yeast.
1111   Nucleic Acids Res. 2012;40: D700–5. doi:10.1093/nar/gkr1029

1112   66.   Graziano BR, Yu H-YE, Alioto SL, Eskin JA, Ydenberg CA, Waterman DP, et al.
1113   The F-BAR protein Hof1 tunes formin activity to sculpt actin cables during
1114   polarized growth. Mol Biol Cell. 2014;25: 1730–1743. doi:10.1091/mbc.E14-03-
1115   0850

1116   67.   Nogami S, Ohya Y, Yvert G. Genetic Complexity and Quantitative Trait Loci
1117   Mapping of Yeast Morphological Traits. PLoS Genet. 2007;3: e31.
1118   doi:10.1371/journal.pgen.0030031

1119   68.   Lynch M, Walsh B. Correlations between characters. Genetics and Analysis of
1120   Quantitative Traits. Sinauer Associates Incorporated; 1998. pp. 629–656.
1121   doi:10.1046/j.1420-9101.2003.00556.x/full

69.   Bliese P. Multilevel modeling in R (2.5). Retrieved September 2013.

70.   Breiman L, Friedman JH. Estimating Optimal Transformations for Multiple
Regression and Correlation. Journal of the American Statistical Association.
Taylor & Francis Group; 2012;80: 580–598.

71.   Setty M, Tadmor MD, Reich-Zeliger S, Angel O, Salame TM, Kathail P, et al.
Wishbone identifies bifurcating developmental trajectories from single-cell data.
Nat Biotechnol. 2016;34: 637–645. doi:10.1038/nbt.3569

72.   Turner JJ, Ewald JC, Skotheim JM. Cell size control in yeast. Curr Biol. 2012;22:
R350–9. doi:10.1016/j.cub.2012.02.041

73.   Wang R, Kamgoue A, Normand C, Léger-Silvestre I, Mangeat T, Gadal O. High
resolution microscopy reveals the nuclear shape of budding yeast during cell cycle
and in various biological states. J Cell Sci. 2016;129: 4480–4495.
doi:10.1242/jcs.188250

74.   Logg K, Bodvard K, Blomberg A, Käll M. Investigations on light-induced stress in
fluorescence microscopy using nuclear localization of the transcription factor
Msn2p as a reporter. FEMS Yeast Res. 2009;9: 875–884. doi:10.1111/j.1567-
1364.2009.00536.x

75.   Wickham H. ggplot2. New York, NY: Springer New York; 2009.
doi:10.1007/978-0-387-98141-3

76.   Pavlicev M, Cheverud JM. Constraints Evolve: Context Dependency of Gene
Effects Allows Evolution of Pleiotropy. Annu Rev Ecol Evol Syst. 2015;46:
413:434.

77.   Pavlicev M, Wagner GP, Noonan JP, Hallgrímsson B, Cheverud JM. Genomic
correlates of relationship QTL involved in fore- versus hind limb divergence in
mice. Genome Biol Evol. 2013;5: 1926–1936. doi:10.1093/gbe/evt144

78.   Porto A, Schmelter R, VandeBerg JL, Marroig G, Cheverud JM. Evolution of the
genotype-to-phenotype map and the cost of pleiotropy in mammals. Genetics.
2016;204: 1601–1612.

79.   Lande R. Quantitative genetic analysis of multivariate evolution, applied to
brain:body size allometry. Evolution. 2017;33: 402–416. doi:10.1111/j.1558-
5646.1979.tb04694.x

80.   Arnold SJ, Bürger R, Hohenlohe PA, Ajie BC, Jones AG. Understanding the
evolution and stability of the g-matrix. Evolution. 2008;62: 2451–2461.
doi:10.1111/j.1558-5646.2008.00472.x

81.   Björklund M, Husby A, Gustafsson L. Rapid and unpredictable changes of the G-
matrix in a natural bird population over 25 years. J Evol Biol. 2012;26: 1–13.
doi:10.1111/jeb.12044

82.　Cheverud JM. A comparison of genetic and phenotypic correlations. Evolution.
2017;42: 958–968. doi:10.1111/j.1558-5646.1988.tb02514.x

83.　Arnold SJ, Phillips PC. Hierarchical comparison of genetic variance-covariance
matrices. Ii coastal-inland divergence in the garter snake, thamnophis elegans.
Evolution. 2017;53: 1516–1527. doi:j.1558-5646.1999.tb05415.x

84.　Sgrò CM, Hoffmann AA. Genetic correlations, tradeoffs and environmental
variation. Heredity. 2004;93: 241–248.

85.　Pavlicev M, Wagner GP. Evolutionary Systems Biology: Shifting Focus to the
Context-Dependency of Genetic Effects. Integrative Organismal Biology. 2015. p.
91:108.

86.　Levy SF, Siegal ML. Network hubs buffer environmental variation in
Saccharomyces cerevisiae. PLoS Biol. 2008;6: e264.
doi:10.1371/journal.pbio.0060264

87.　Ziv N, Siegal ML, Gresham D. Genetic and nongenetic determinants of cell
growth variation assessed by high-throughput microscopy. Mol Biol Evol.
2013;30: 2568–2578. doi:10.1093/molbev/mst138

88.　Millard SP. EnvStats. New York, NY: Springer Science & Business Media; 2013.
doi:10.1007/978-1-4614-8456-1

89.　Gietz RD, Woods RA. Transformation of yeast by lithium acetate/single-stranded
carrier DNA/polyethylene glycol method. Meth Enzymol. 2002;350: 87–96.

90.　Miyajima I, Nakafuku M, Nakayama N, Brenner C, Miyajima A, Kaibuchi K, et
al. GPA1, a haploid-specific essential gene, encodes a yeast homolog of
mammalian G protein which may be involved in mating factor signal transduction.
Cell. 1987;50: 1011–1019.

91.　Pinheiro J, Bates D, DebRoy S, Sarkar D. R Core Team (2018). nlme: linear and
nonlinear mixed effects models. R package version 3.1-137. 2018.

92.　Levy SF, Ziv N, Siegal ML. Bet hedging in yeast by heterogeneous, age-correlated
expression of a stress protectant. PLoS Biol. 2012;10: e1001325.
doi:10.1371/journal.pbio.1001325

93.　Li S, Giardina DM, Siegal ML. Control of nongenetic heterogeneity in growth rate
and stress tolerance of Saccharomyces cerevisiae by cyclic AMP-regulated
transcription factors. PLoS Genet. 2018;14: e1007744.
doi:10.1371/journal.pgen.1007744

Figures:

Figure 1: Pleiotropic QTL influence yeast single-cell morphology. The vertical axes in all plots represent the 158 CalMorph morphological traits for which we detect QTL with a genome-wide FDR of 5%. These traits are sorted, from top to bottom, based on the difference between the oak and wine parental strains. **(A)** Of 44 QTL that contribute to variation in single-cell morphology, 37 contribute to variation in multiple features. The horizontal axis indicates the chromosomal location of each QTL (in cM). Differently shaped points indicate QTL that are more than 30 cM apart on the same chromosome. The darkness of a point represents the effect size of a QTL; effect sizes range from 0.3% (lightest points) to 18% (darkest points) of the difference between parents. The QTL highlighted in pink, green, purple, and orange contribute to 58, 33, 78, or 66 morphological features, respectively. **(B)** Single genes contribute to multiple morphological features. The horizontal axis represents the relative phenotypic differences between the wine and oak parents (leftmost column) or one of these strains versus a derivative strain that differs in a single gene. The relative phenotypic differences between a pair of strains are calculated by scaling each trait to have a mean of 0 and standard deviation of 1 across all individuals in both strains, and then subtracting the average value in one strain from that in the other. To control for variation among replicate experiments, this scaling was done independently for each replicate experiment in which both strains were imaged. Error bars represent 95% confidence intervals inferred from the replicate experiments. The two gene replacements shown, *PXL1* and *HOF1*, are respectively located within the QTL highlighted in green and purple in panel **A**. When calculating the difference between strains, we always subtracted the trait values of the strain possessing more wine genes from those of the strain possessing more oak genes, such that the effects of the wine or oak gene replacements appear in the same direction on all plots.

Figure 2: Pairs of traits with high correlation across clones are overrepresented among those influenced by pleiotropic QTL. Within-genotype correlations (r_W) are calculated for 5645 pairs of morphological traits. **(A)** Histogram showing distribution of r_W. **(B)** This plot displays the 11 of 35 pleiotropic QTL that exclusively influence pairs of traits with $r_W > 0.2$. Points represent r_W for pairs of traits influenced by a QTL; the shapes of these points match those in **Fig 1A** and distinguish a QTL from others on the same chromosome. **(C)** Points in grey represent r_W for all 5645 pairs of traits. Points in black each represent the median r_W across pairs of traits influenced by one of the 24 pleiotropic QTL not included in panel B. The next four sets of points each display r_W for pairs of traits influenced by a single QTL corresponding to those highlighted in the same color in **Fig 1A**. Each boxplot shows the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5 × IQR (whiskers). **(D)** A force-directed network visualizing how pairs of morphological features correlate across clones. Each node represents a single-cell morphological trait measured in large-budded cells. The thickness of the line connecting each pair of nodes is proportional to r_W. Node position in the network is determined using the Fruchterman-Reingold algorithm. Purple nodes correspond to traits influenced by a QTL on chromosome 13 containing the *HOF1* gene. **(E)** Cumulative distributions of weighted clustering coefficients (*wcc*) in a network created using measured values of r_W (red line) or in 100 permuted networks (grey lines) for traits corresponding to large-budded cells. Permutations were performed by sampling r_W, without replacement, and reassigning each value to a random pair of traits. **(F)** This panel is similar to panel C, except points represent *wcc* of traits rather than r_W of trait pairs, and black points each represent the median *wcc* across traits influenced by one of 24 pleiotropic QTL.

Figure 3: Natural genetic variation affects the correlation between morphological features. The absolute value of the between-strain correlation (r_B), made negative when r_B and r_W have opposite signs, is plotted against the absolute value of the within-strain correlation (r_W), for each pair of traits. The plot at left shows pairs of traits that share at least one QTL influence. The color of each point represents the number of pleiotropic QTL that influence both traits in that pair. The plot at right shows pairs of traits that share no QTL influence. The dashed line represents a Bonferroni-corrected significance threshold of $p < 0.01$. Insets represent the results of correlation partitioning performed after randomly assigning individual cells to groups (pseudo-strains) having the same numbers of cells as the actual strains.

Figure 4: Morphological features vary as cells divide. The morphological features of unbudded (red), small-budded (blue), and large-budded (purple) cells change as these cells progress through the cell cycle. **(A)** Variation of four traits through the cell cycle. Line plots represent fixed-cell images from all 374 mapping family strains, positioned on the horizontal axis based on progression through the cell cycle as calculated by Wishbone [71]. Regression lines are smoothed with cubic splines, calculated with the "gam" method in the R package ggplot2 [75], to depict trends describing how each displayed trait varies across the estimated growth trajectory. The displayed trends match those observed in micrographs of live cells progressing through division. Each series of micrographs displays a different live cell imaged over several minutes, which are displayed in the lower right corner of each micrograph. **(B)** Centered data for 11, 23, and 44 unbudded, small-budded and large-budded cells, respectively, show how Wishbone sorts live cells in a way that recapitulates the actual time series. Each point in these plots represents a cell image from a single timepoint. The horizontal axis represents Wishbone's estimation of how far that cell has progressed through division. The vertical axis displays time, as a percentage of the total time elapsed and adjusted in a way that controls for every cell having started at a different place in the cell division cycle at time zero (see *Methods*). Trend lines are smooth fits using the "loess" method in the R package ggplot [75]. **(C)** The correlation between some morphological features changes throughout the course of cell division. The scatterplot shows how binning influences both the phenotypic correlation (vertical axis) and phenotypic variation (horizontal axis) across clones. Each point represents these values for a pair of traits as

measured in 1 of 16 bins. The value on the horizontal axis represents whichever trait in each pair had the larger decrease in standard deviation, as such decreases are likely to reduce the correlation on the vertical axis. The blue line shows a smooth fit by loess regression. Colored points on the scatterplot correspond to bin 5 for each pair of traits represented by the line plots in panel **D**. **(D)** These line plots show three pairs of traits for which binning increases r_W such that it approaches r_B (leftmost three plots), and three pairs of traits for which r_W does not approach r_B even after binning (rightmost three plots). In each plot, r_B is shown as the horizontal green line, r_W (without binning) is shown as the horizontal purple line, and r_W for each bin is shown in black.

Figure 5: Many QTL demonstrate horizontal pleiotropy. (A) Eliminating allelic variation at the site of each QTL tends to reduce r_B. The vertical axis represents how r_B changes upon eliminating allelic variation at each QTL site. Each point represents the median change in r_B for all pairs of traits that are affected or unaffected by one of the 24 QTL suspected of horizontal pleiotropy. Boxplots summarize these changes in r_B when re-measured across strains possessing the wine (red) or the oak (blue) allele at the marker closest to the QTL. **(B)** The upper and lower series of three plots demonstrate two different ways that a QTL can increase the correlation between traits. Each point represents a yeast strain possessing either the wine (red) or the oak (blue) allele at a marker closest to a QTL on chromosome 15 (upper) or 8 (lower). In the upper plots, the QTL increases the correlation between nucleus shape and size ratio when it is segregating across strains. In the lower plots, the wine allele strengthens a correlation between bud shape and the position of the nucleus in the mother cell that is weak in the oak subpopulation. Numbers in the lower corner of each plot represent r_B for the strains displayed. **(C)** Cumulative distributions display the extent to which binning cells or splitting strains resolves the difference between r_B and r_W. When calculating percent resolved (horizontal axes) we always plot the value in whichever

subset (*e.g.* wine or oak) this percent is greatest. If subsetting always worsens the discrepancy between r_B vs. r_W, we score this as 0% resolution. Only pairs of traits for which r_B is significantly greater than r_W are considered. The pink, green, purple and orange lines show the effect of splitting strains by whether they inherited the wine or oak allele at the marker closest to each of four QTL (colors correspond to QTL in **Fig 1A**). In these plots, comparing the solid vs dotted lines shows that splitting strains resolves the discrepancy between r_B and r_W more often for pairs in which both traits are affected by the QTL than pairs in which both traits are unaffected. The black lines in the leftmost plot summarize these effects across 24 QTL, displaying for each trait pair, the largest resolution in the r_B vs. r_W discrepancy observed across all QTL that affect the pair of traits (solid line) or all QTL that do not (dotted line). The red line shows the effect of binning cells by their progress through division, displaying the largest resolution in the r_B vs. r_W difference across all 16 bins.

Figure 6: Some MA lines display unique relationships between certain pairs of traits. In all plots, black represents the ancestor of the MA lines and colors represent MA lines with trait correlations that differ from other lines (strains: black = HAncestor, green = DHC81H1, red = DHC41H1, magenta = DHC40H1, blue = DHC66H1, purple = DHC84H1; see Table S2 in Geiler-Samerotte et al 2016 [50]). **(A)** Histograms display the number of MA lines with Pearson correlations corresponding to the values on the horizontal axis for four example pairs of traits; the number of bins is set to 30. **(B)** This plot displays, for each of the 94 MA lines, the cumulative distribution of the number of standard deviations away from the mean correlation across all trait pairs. **(C)** Plots display, for each MA line, the maximum deviation from the mean observed for any pair of traits (left) and the average standard deviation observed across all pairs of traits (right).

Supplemental Figures:

Figure S1: Morphological differences exist between the parents of the QTL mapping family. Each density plot displays the distribution of phenotype values from yeast cells corresponding to the wine parent (red), the oak parent (blue), or all of the 374 progeny (grey) for the trait listed on the horizontal axis. Trait names in parentheses correspond to those listed in the CalMorph manual [52]. Before plotting, each morphological trait was transformed to have a mean of zero and a standard deviation of one across all strains. Each distribution represents at minimum 5,000 cells from three replicate experiments; distributions corresponding to progeny strains represent many more cells (70,000 – 200,000 depending on whether the trait was measured in unbudded, small-budded, or large-budded cells).

Figure S2: Total numbers of cells imaged per each of 374 progeny strains. Each point represents, for one of the 374 progeny strains, the number of unbudded, small-budded, or large-budded cells for which images passed filtering (see *Methods*). Each boxplot shows the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5 × IQR (whiskers).

Figure S3: Comparison of correlation estimates obtained from correlation partitioning with those obtained from a mixed-effect linear model. Each point represents one of 350 randomly sampled trait pairs of the 5645 total. Vertical axes display trait correlations estimated using the correlation-partitioning approach; horizontal axes display trait correlations estimated using a mixed-effect linear model that specifies the variance-covariance structure of the experimental design.

Figure S4: Single-cell morphological traits have higher weighted clustering coefficients (*wcc*) than expected given the distribution of r_W. (A – B) Force-directed networks visualizing how pairs of morphological features correlate across clones in unbudded (panel A) and small-budded (panel B) cells. Each node represents a single-cell morphological trait. The thickness of the line connecting each pair of nodes is proportional to r_W. Node position in the network is determined using the Fruchterman-Reingold algorithm. Purple nodes correspond to traits influenced by a QTL on chromosome 13 containing the *HOF1* gene. **(C – D)** Cumulative distributions of weighted clustering coefficients (*wcc*) in a network created using measured values of r_W (red line) or in 100 permuted networks (grey lines) for traits corresponding to unbudded (panel C) or small-budded (panel D) cells. Permutations were performed by sampling r_W, without replacement, and reassigning each value to a random pair of traits.

Figure S5: Wishbone recapitulates time series data obtained in live images of 78 cells undergoing exponential growth. Each point represents a cell image. Horizontal axes display the minute that image was captured during a three-hour window of exponential growth. Vertical axes display Wishbone's prediction of how far that cell image has passed through the cell cycle. Linear regression lines are calculated with the "lm" method in the R package ggplot2 [75], and are colored red for images corresponding to unbudded cells, blue for small-budded cells and purple for large budded cells. Plots are organized by cell type and then from earliest to latest average predicted progress through cell division.

S1 Table. Chromosomal locations, effects sizes and phenotypes affected by quantitative trait loci described in this study.

S2 Table. Impact of gene swaps on single-cell morphological traits including the corrected phenotypic difference between strains for each phenotype, and its standard deviation and standard error across replicate experiments.