1   **Characters of neoantigens in cancer immunotherapy**

2   Peng Bai[1], Yongzheng Li[1], Qiuping Zhou[1], Jiaqi Xia[1], Min Wu[2], Hexiang Deng[3], John W. Kappler[4,5,6,7],

3   Philippa Marrack[4,6,8], Yu Zhou[1], Lei Yin[1*]

1 State Key Laboratory of Virology, Hubei Key Laboratory of Cell Homeostasis, College of Life Sciences,

Wuhan University, Wuhan, 430072, Hubei, China

2 Hubei Key Laboratory of Cell Homeostasis, College of Life Sciences, Wuhan University, Wuhan,

430072, Hubei, China

3 Key Laboratory of Biomedical Polymers, Ministry of Education, the Institute for Advanced Studies,

College of Chemistry and Molecular Sciences, Wuhan University, Wuhan, 430072, China

4 Department of Biomedical Research, National Jewish Health, Denver, CO 80206, USA

5 Barbara Davis Center for Childhood Diabetes, University of Colorado, Aurora, CO 80045, USA

6 Department of Immunology and Microbiology, University of Colorado School of Medicine, Aurora,

CO 80045, USA

7 Structural Biology and Biochemistry program, University of Colorado Anschutz Medical Campus,

Aurora, CO 80045, USA

8 Department of Biochemistry and Molecular Genetics, University of Colorado Anschutz Medical

Campus, Aurora, CO 80045, USA

Abstract

Cancer mutations lead to the production of mutated proteins from which can be processed and

presented to CD8[+] T cells as neoantigens. Despite the promise of immunotherapy targeting

neoantigens, which mutant peptides are likely to induce immune responses remain elusive. Here, we

determined two sets of MHC-I bound neoantigen structures and performed bioinformatics investigation

based on neoantigen data. Structural and bioinformatic analyses reveal that some types of mutations in

immunogenic neoantigens are preferentially selected. We also developed a strategy to expand the usage

of a known neoantigen by scanning potential MHC alleles. Overall, our results will extend the ability to

discover neoantigens that may be useful for cancer immunotherapy.

Introduction

The immune system can sometimes recognize and destroy cancer cells[1]. CD8[+] T cells have been

demonstrated to be a major component in this process by both recognizing and destroying the target

cells[2–5]. For T cells to react with tumor cells two components are required, major histocompatibility

complex class I and possibly class II proteins (MHC-I, MHC-II) on the tumor cells and peptides which

processed by the tumors bound to these MHC proteins. CD8[+] T cells can then recognize the

combination of MHCI and tumor derived peptides on the cancer cells. Thus, it is crucial to identify

effective tumor antigenic peptides with their relevant MHCs.

During the last 25 years, great efforts have been made to identify tumor antigens[6]. These tumor

antigens can be classified into two broad categories, self-antigens which are seldom expressed in

normal tissue or expressed at much higher levels on the cancer cell versus normal tissues (like NY-

ESO-1, MART-1) and non-self-antigens called neoantigens which are derived from somatic mutations

in the tumor[7]. The neoantigens are not presented by normal tissues, hence the immune system is not

tolerant to them and views them, correctly, as foreign antigens and responds appropriately[8, 9].

Recently, next-generation sequencing (NGS) provide formidable identification of tumor-specific non-

synonymous mutations, frameshift mutations, and gene rearrangements, which have driven clinical

immunotherapeutic methods such as vaccination, adoptive cell transfer (ACT) and blockade of the

proteins that inhibit T cell activation, have demonstrated that immunotherapy targeting tumor

neoantigens can induce objective clinical responses in many patients with metastatic cancers[10–13]. It

appears that T cell targeting of tumor neoantigens might represent the "final common pathway" to treat

cancer*(14)*. However, neoantigen discovery and validation remains a daunting problem. Classical

cDNA expressing screening methods to identify tumor neoantigens without knowledge of the

properties needed for a mutated peptide to be a successful immunotherapeutic is not efficient*(15)*. To

solve this problem, recently, a study identified mouse neoantigens by combination of NGS and

prediction algorithms for antigen presentation by MHCI*(16)*. In brief, they compared mouse tumor and

normal DNA for identifying non-synonymous mutations and used peptide/MHCI binding prediction

algorithms to find mutated peptides which could bind MHC to be potential effective neoantigens.

These advances of discovering neoantigens bring us a step closer to personalize immunotherapy.

Indeed, neoantigen prediction lighten the burden of immunogenicity testing by reducing the number of

candidate peptides*(17–27)*. Nevertheless these algorithms are not robust enough to identify

immunogenic neoantigens accurately*(14, 28)*. There are several reasons for this. Firstly, many of these

algorithms do not consider the likelihood that the peptide in question will be properly processed and

transported to the endoplasmic reticulum, where they bind MHCI. This problem can be solved by

adding other methods, such as NetChop*(29)* (for antigen processing prediction) and NetCTL*(30)* (for

peptide transport prediction) to the prediction process. Secondly, the MHC binding prediction methods

based on artificial neural network (ANN) models need a lot of data ,such as those available in NetMHC

4.0, for training*(31)*. However, large amounts of these input data are lacking for many human MHC

alleles. Also, in some cases, most peptides in the training data were of non-tumor origin such as

cultured cell lines and microbial foreign peptides. Thus, it is unclear whether these non-self peptidome

data accurately represent the profile of mutated-self neoantigens and elicit T cell response*(32)*. Thirdly,

among neoantigens binding MHC actually, only a small portion of them are immunogenic and have

therapeutic effects*(33)*. Thus, it is crucial to investigate the details of neoantigen immunogenicity and

to improve the performance of *in silico* neoantigen prediction algorithm.

Recently, Tran *et al*. reported a remarkable case of a patient whose metastatic colorectal cancer

regressed after infusion of HLA-C*08:02-restricted tumor-infiltrating lymphocytes (TILs) targeting

KRAS G12D neoantigens. *Kras* encodes a small GTPase that mediates downstream signaling from

growth factor receptors, and it is one of the most infamous driver genes led to oncogenesis. The most

common KRAS mutation is a gain-of-function substitution of the glycine to aspartic acid at codon 12

(KRAS G12D), a change that is often present in colorectal or pancreatic cancer*(34–36)*. The KRAS

G12D mutant 9mer peptide GADGVGKSA and 10mer peptide GADGVGKSAL can both stimulate

autologous T cells in the context of HLA-C*08:02, while their wild type counterparts cannot*(23)*.

Adoptive cell transfer (ACT) of T cells specific for the single mutant neoantigen led to tumor

disappearance in patient. Thus, the KRAS G12D neoantigens have been acknowledged to be useful and

typical neoantigen that might be used as a model to illustrate the secret of therapeutic tumor

neoantigens.

Here we took structural analyses of several neoantigens with their relevant MHCs, combined with a

broad examination of immunotherapeutic and non-immunogenic tumor antigens. We concluded that a

universal type of tumor neoantigen peptide mutations from non-preferential residues to preferential

residues with their relevant MHCs at anchor positions is unexpectedly a good indicator to predict

neoantigen immunogenicity. Combined with other findings about effective neoantigens we thus

propose an optimized strategy for neoantigen identification.

Results

The structures of HLA-C*08:02 in complex with KRAS G12D neoantigens

As described above, Tran and colleagues reported a clinical case which showed substantial tumor

regression after adoptive T cell transfer of T cells specific for HLA-C*08:02 bound to the mutant

KRAS G12D 9mer and 10mer peptides*(23)*. HLA-C*08:02 is expressed in approximately 8% of

whites*(23)* and 7% of African Americans*(37)* in the U.S. The KRAS G12D mutation is most common

in pancreatic and colorectal cancers, being present in approximately 45% and 13% of these two tumor

types, respectively*(38)*. Thus, overall about 2.3% patients with colorectal cancer share the HLA-

C*08:02 allele and KRAS G12D mutation (Supplementary Table 1) and these results suggest that

thousands of patients per year in the United States alone could potentially be eligible for the therapy

targeting HLA-C*08:02-restricted KRAS G12D neoantigens.

With this in mind we decided it would be useful to know the structure of HLA-C*08:02 in complex

with four peptides: wild type KRAS G12D 9mer GAGGVGKSA (wt9m); wild type KRAS G12D

10mer peptide GAGGVGKSAL (wt10m); mutant KRAS G12D 9mer peptide GADGVGKSA (mut9m,

mutation site is indicated with the amino acid underlined) and mutant KRAS G12D 10mer peptide

GADGVGKSAL (mut10m). HLA-C*08:02 heavy chain and light chain β2-microglobulin (β2M) were

expressed in *E. coli* as inclusion bodies and refolded with peptides. C08-mut9m and C08-mut10m

complex were successfully refolded and detected on SDS-PAGE gels. Unfortunately, we failed to

refold C08-wt9m complex and C08-wt10mer complex even with a ten-fold increase in concentration of

the peptide (data not shown). This result suggested that the glycine to aspartic acid mutation of these

neoantigens is important to maintain the pMHC stability. However, the artificial neural network (ANN)

based peptide binding algorithm NetMHC4.0*(39)* predicted that binding affinity of KRAS G12D

mutated neoantigens (both 9mer and 10mer) and their wild type counterparts for HLA-C*08:02 would

all be very low (less than 15,000 nM, Supplementary Table 2). This could be due to relatively less

training data of HLA-C*08:02.

We solved the structures at 2.4 angstroms (Å) for the C08-mut9m complex and 1.9 Å for the C08-

mut10m complex (Supplementary Table 3). The high-quality resolution and unbiased electron density

at the region involving the peptides (Fig.1a, b) allowed reliable model building. The C08-mut9m

complex and the C08-mut10m complex shown highly conserved conformation except the peptide

region (Fig.1 c, d).

Peptide amino acid binding motif of HLA-C*08:02 shows that smaller residues such as alanine and

serine are preferably selected at P1 and P2 position (Fig.1e). This preference is due to the narrow cleft

formed by several C*08:02 aromatic residues (Tyr7, Phe33, Tyr67, Tyr99, Tyr59, Tyr171, Tyr159 and

Trp167), which limit the size of the P1 and P2 side chains that can bind at that site (Fig.1f). C08-

mut9m and C08-mut10m are relatively similar at the position 1 (P1) and position 2 (P2) of the

peptides. In C08-mut9m and C08-mut10m the N-terminal P1G (Glycine at position 1) residue of the

peptides form hydrogen bonds with Tyr7, Tyr159 and Tyr171 and the P2A residues interact with Glu63

and Lys66 (Fig. 2a).

Differences appeared at the P3-PΩ (PΩ: C-terminal position of peptide) regions of the peptides.

Although P3D in C08-mut9m and C08-mut10m both act as anchor residues, the side chain carboxyl

group of P3D forms an intra-chain hydrogen bond with P4G in mut9m peptide, while this hydrogen

bond is absent in the mut10m peptide structure. The hydrogen bond in the mut9m structure pulls down

the arch of the 9mer peptide, causing the region of the 9mer peptide that is accessible to TCRs to differ

from that of the 10mer peptide. The greatest difference between the two structures is apparent in the

center of the peptide (P4-P7), where each complex differs in the contacts it offers for TCR recognition.

The P8S has different orientations in the two structures and acts as an auxiliary anchor in C08-mut10m.

Differences were also observed at the C-termini of the peptides (Fig. 1c; Fig. 2b, c). The P9A of C08-

mut9m forms hydrogen bonds with Ser77, Asn80, Tyr84, Thr143 and Lys146, and acts as an anchor

residue (Fig. 2b). The P9A of C08-mut10m is totally different from the P9A of C08-mut9m with the

side chain up. However, the P10L of C08-mut10m has similar orientation towards HLA-C*08:02 and

plays a similar role as P9A in C08-mut9m (Fig. 1c; Fig. 2b, c).

HLA-C*08:02 presents KRAS G12D neoantigens with an unconventional MHC anchor motif

Typically, peptides always use anchor positions at P2 and PΩ to bind the B and F pockets respectively

of HLA class I alleles. The structures of the C08-mut9m and C08-mut10m reveal that G12D 9mer and

10mer peptides all use unconventional P3 and PΩ residues for anchors to form stable complexes with

HLA-C*08:02 (Fig.1c; Fig. 2b, c) thus differing from the canonical HLA Class I binding pattern (using

P2 and PΩ residues for anchors). In this case, the B pocket of HLA-C*08:02 binds the P3D anchor

residue via interactions with Arg97 and Arg156. Although the 9mer and 10mer peptides occupy the

same F pocket with their PΩ residues as anchors, the P10L side chain from the 10mer peptide is buried

more deeply than P9A in the 9mer structure (Fig.1c; Fig. 2b, c). Also, instead of the upward residue,

P8S, in 9mer peptide, the 10mer peptide uses P8S as an auxiliary anchor, binding the HLA protein via

hydrogen bonds with Glu152 and Arg156. These interactions squeeze the P4 to P7 residues together

with residue P3D (Fig. 2c). Also, together with P3D, the existence of auxiliary anchor residue P8S

arches the P4-P7 residues of the 10mer peptide to a higher position than that in the 9mer peptide (Fig.

2b, c). This alteration likely increases the binding affinity of the 10mer peptide with HLA and changes

the pHLA binding surface for the T cell receptor (TCR).

KRAS G12D 9mer and 10mer peptides were previously reported to stimulate different T-cell clones

with different TCR variable chains(10). Our structural analyses are in line with such a report since

C08-mut9m and C08-mut10m provide different binding surfaces for T cell recognition (Fig.1b; Fig. 2b,

c). Furthermore, both wild type counterparts of 9mer and 10mer peptides failed to refold with HLA-

C*08:02, which means that the G12D mutation is important for forming the pMHC complex. Structural

analysis shows that G12D mutations of the KRAS neoantigens act as anchors and stabilize contacts

with the MHC. Thus, in this case, the KRAS mutation shows the likelihood that anchor mutated

neoantigens will exist to stimulate T cells.

168   Structural basis of non-therapeutic mouse neoantigen DPAGT1 V213L in complex with H-2 Kb

Having verified that the P3D residue of KRAS G12D neoantigens plays an important role in formation

of the pMHC complex, we studied whether other cancer neopeptides with anchor mutations had similar

properties. One such neopeptide, which is not immunotherapeutic, is the mutation of the C terminal

amino acid anchor residue of the mouse mutant DPAGT1 V213L peptide*(7)*. To study this we solved

the structure of mouse MHCI H-2 Kb in complex with mutant DPAGT1 V213L 8mer peptide

(SIIVFNL<u>V</u>, mutation site is indicated with the amino acid underlined) and the wild type 8mer

counterpart (SIIVFNLL) (Supplementary Table 3). The last residue of the peptide, a valine in the wild

type 8mer peptide (wt8mV) was replaced by leucine in the mutant 8mer peptide (mut8mL). Soluble

complexes of Kb-wt8mV and Kb-mut8mL were separately expressed, refolded and purified for

crystallization trials. Crystal diffraction data of Kb-wt8mV and Kb-mut8mL are processed to 2.4 Å and

2.5 Å resolution respectively (Supplementary Table 3) and provides unambiguous electron density for

each peptide (Supplementary Fig.1a).

The overall structure of the Kb-wt8mV complex closely resembles that of Kb-mut8mL with the

exception of a slight difference at PΩ (P8) (Supplementary Fig.1b). The C-terminal PΩ residue acts as

an anchor in both the Kb-wt8mV and Kb-mut8mL complexes (Supplementary Fig.1b). Moreover, both

PΩ valine and leucine were preferably selected by H-2 Kb in a peptide library analysis (Supplementary

Fig.1c). Both of the PΩ residues formed hydrogen bonds with Asp77, Tyr84, Thr143 and Lys146 of Kb

but the side chain of leucine in the mut8mL peptide is insert more deeply into Kb than valine in wt8mV

peptide because of longer side chain of the Leucine residue. Thus, P8L is able to make Van der Waals

contact with Tyr116 on the floor of the F pocket (Supplementary Fig.1d, e). These alterations did not

significantly affect the formation of peptide/Kb complex and change the TCR binding surface,

suggesting that, as far as T cell recognition is concerned the mut8mL neoantigen would be treated as a

"self-peptide", thus minimizing mut8mL neoantigen immunogenicity *in vivo*. All these indicate the

wild type and mutant DPAGT1 V213L peptides interact with Kb in a similar way, thus the mutation

might provide neither a higher concentration of peptide/Kb on the cell surface nor a TCR interface that

differs. This is unlike the case of the KRAS G12D mutant peptides which appear to interact more

intensively with HLA, via the mutant acid residue P3D interaction with basic residues in the B pocket

of C*08:02, versus than the wild type counterpart.

Taken together, all these suggest that whether wild type or mutant amino acids at anchor positions

interacted differently with relevant HLA alleles might be key to neoantigen immunogenicity.

Curated datasets of human immunogenic neoantigens from publications

To extend these analyses, bioinformatic studies were performed on all available immunogenic and non-

immunogenic neoantigen peptide/MHC complexes. We manually gathered immunogenic neoantigens

associated with various tumor types from 34 recently published papers*(5, 11–13, 28, 40–66)*. This

immunogenic neoantigen dataset (IND) encompassed 128 neoepitopes (neoantigens and their relevant

HLAs) from 15 types of human cancer. Each of them was well tested by at least tested *in vitro* T cell

assays. Less well characterized neoantigens and the neoantigens without minimal length were not

included. Each entry included information such as the cancer type, HLA allele type, gene name, peptide

sequence and other characteristics which are important in cancer immunotherapy. Melanoma and non-

small cell lung cancer contain higher mutational burdens and have been more extensively studied than

others, thus nearly 80% entries in our dataset came from them.

We also gathered 11739 non-immunogenic neopeptides (peptides splice from mutant proteins) from 13

published papers termed raw non-immunogenic neopeptide dataset (RNND). All of these mutant

peptides were tested in *in vitro* T cell assays or clinical trials and did not elicit effective immune

responses. Notably, the vast majority of them were not the minimal peptides known to induce $CD8^+$ T

cells response (most of them are 25mer peptides). Thus, we use the prediction algorithm NetMHCpan

4.0*(67)* to predict the conventional minimal length (9mers and 10mers) non-immunogenic neopeptides

with relevant HLAs under a cut-off IC50=500 nM. This dataset was termed non-immunogenic

neopeptide dataset (NND). This dataset contains 2883 putative entries. It turns out that many

neoantigens predicted to bind patients' HLAs very well are experimentally tested to be non-

immunogenic and do not have the ability to evoke effective immune response. This fact compromised

the usage of predicting immunogenic neoantigen through prediction of neoantigen binding to patients'

HLAs.

Evident differences of mutation distribution between immunogenic and non-immunogenic

peptides

It is thought that immunogenic neoantigens would bind MHC well and the mutations in the

immunogenic neoantigens could increase the affinity for binding MHC. With our immunogenic

neoantigen data collected, we preformed binding prediction (NetMHCpan 4.0) of wild type and mutant

peptides in IND and showed the distribution of affinity ratio between them (Fig.3a). About 25.5% of

immunogenic neoantigens shows the predicted binding affinity above 500nM and about 24% do not

exhibit increased prediction binding affinity (WT/MUT affinity ratio <1). Thus, more characters of

immunogenic neoantigens which could help immunogenic neoantigen predictions are needed.

It is suspected that the position in the peptides of mutated amino acids could affects the

immunogenicity of neoantigens. To further investigate the mutation distribution at each peptide

position between immunogenic and non-immunogenic data, we calculated the mutation occurrence

frequency at each position of 9mer and 10mer peptides. Interestingly, both immunogenic 9mer and

10mer neoantigens show frequency fluctuation at each position with their mutations. The fluctuation of

non-immunogenic 9mer and 10mer neoantigens seems not as high as those from immunogenic dataset

(Fig. 3b; Supplementary Fig. 2). However the difference is not reaching significance probably due to

the limited size of so far available immunogenic data. Mutations in 9mer neoantigens were more

frequent at P2, P4, P5 and P8 position and relatively lower frequency at P3 and P7 positions, relative to

non-immunogenic 9mer data (Fig. 3b). Conceptually, when bound to HLA, peptides usually use P2 and

PΩ as anchor residues and use P4, P5 and P8 to contact TCR. Other residues in the peptide may contact

with MHC but usually contribute less to the affinity of the peptide-HLA interaction[68]. Also to be

taken into this account is the fact that our structural studies above also show that mutations at anchor

positions can evoke effective immune responses. Thus, these observations enlightened us to classify the

mutation based on whether it located on the anchor (contribute primary binding force to MHC), MHC

contact (contact MHC but contribute much less for binding MHC) or TCR contact positions for further

study.

To find out the mutation location (anchor, MHC contact position and TCR contact position) of each

peptide in IND an NND respectively, a strategy was developed to define the mutations location of

neoantigens by combining the SYFPEITHI database*(69)*, antigen binding motif from NetMHCpan*(67)*

and solved pMHC structures from Protein Data Bank (see **Methods**). We chose 9mer length of

neoantigens to do these studies since 9mer peptides are most frequent in both the IND and NND

collections. Peptides bound to MHC alleles for which structure models were not available in PDB were

excluded from these analyses.

The 9mer neoantigens (or neopeptides) with mutations from IND and NND were divided into three

functional classes based on the following: anchor mutation, MHC contact position mutation or TCR

contact position mutation (Fig. 3c). After calculating the frequency of each class, we found that the

immunogenic neoantigens were significantly more likely than NND to express mutations at TCR

contact region (Fig. 3d, $p<0.05$, Z-test) and less likely to include mutations at MHC contact regions

(excluding those at anchor positions) (Fig. 3d, $p<0.01$, Z-test). These results can partially support

proposals of others that neopeptides with TCR contact position mutation will preferentially be

immunogenic*(7)*. However, the odds of mutations at the anchor positions were similar between INDs

and NNDs, therefore the importance of anchor position mutations in immunogenicity is not clear (Fig.

3d). These results are consistent with our structural studies showed that the immunogenic neoantigen

KRAS G12D and the non-immunogenic neopeptide DPAGT1 V213L can both present by MHC.

Certain type of mutation change at anchor position could be a clear indicator to distinguish

immunogenic neoantigens and non-immunogenic neopeptides

It is thought that the nature of mutated amino acids affects the neoantigen's immunogenicity. To find

out if this is so, we calculated the frequency with which different amino acids appeared as the

mutations in either IND or NND neoantigens. So far Leu and Tyr are likely to show up in the mutation

position from immunogenic neoantigens (Supplementary Fig. 3). This could be due to the idea that

mutation to Leu and Tyr might supply good contacts with MHC to stabilize the binding of neoantigens

to MHC or good contacts with TCR to stimulate T cells. This is consistent with our previously studies

that Tyr is important in TCR-peptide/MHC recognition*(70–73)*.

Next, to check the similarity between wild type and the mutant amino acids of neoantigens and non-

immunogenic neopeptides, we defined similarity of those amino acids from IND and NND using the

normalized BLOSUM62 substitution matrix*(74)*. In this matrix, bigger normalized score represents

more similarity between the wild type and the mutant peptides. For the situation of immunogenic data

compare to non-immunogenic data, the peptides with immunogenicity did not shown significant

difference between wild type and the mutant amino acids (Fig. 4a, p=0.2077, student t-test). For the

situation of anchor mutated peptides, the similarity of immunogenic neoantigens to their wild type

counterparts turn out to be lower than the similarity of the non-immunogenic peptides (Fig. 4b,

p=0.001, student t-test). For the situation of 9mer peptides with TCR contact position mutation, similar

result occurs between immunogenic and non-immunogenic data (Fig. 4c, p=0.019, student t-test). We

did not test for mutations at MHC contact regions, because the available data was too limited so far.

These results demonstrated that through the mutation position classification quantifiable amino acid

differences between mutation and their wild type counterparts do indeed exist at anchor and TCR

contact positions between immunogenic and non-immunogenic data.

To test whether the mutant peptides in IND are more preferentially presented by HLAs than wild type

peptides, we generated amino acids half-bits of each HLA from P1 to P9 (we test 9mer peptides only

due to their relatively abundant quantity) which present in IND, based on the binders collected from

IEDB (see **Methods**). Then, merged bits of mutated neoantigens and their wild type peptides with their

relevant HLAs were calculated respectively (Fig. 4d-f). To our knowledge, peptide with higher merge

bits means preferentially presented by HLA. We first tested peptide merge bits in all 9mer neoantigens

and their wild type counterparts. The higher merged were shown in mutant neoantigens compare to

their wild type counterparts, which means immunogenic neoantigens more likely to be presented by

their relevant HLAs (Fig. 4d, p=0.0088, student t-test). We next tested this character from the data of

9mer anchor mutated neoantigen and 9mer TCR contact position mutated neoantigen. Surprisingly,

anchor mutated neoantigen showed significantly higher preference to be present by their relevant HLAs

than their wild type counterparts (Fig. 4e, ****p<0.0001, student t-test), while TCR contact position

mutated data did not shown any significance (Fig. 4f, p=0.9358, student t-test). From above results, it is

evident that mutations of immunogenic neoantigens in the anchor positions and in the TCR contact

positions shows different characters and should be studied separately. Moreover, the mutations of

immunogenic neoantigens in the anchor positions were needed for further studies.

Inspired by our structure studies of two sets of anchor position mutated neoantigens, we therefore

decided to find out how the mutations and wild type amino acids in immunogenic neoantigens were

related to the known residue preferences of different HLA alleles, and whether the associations were

similar for immunogenic and non-immunogenic peptides. The cut-off for preferential amino acids was

applied to be above 10% (include 10%) as described previouslly(75) based on the data from peptide

library (Supplementary Fig. 4). Neoantigen mutations changes from wild type to mutant at anchor

positions were divided into 4 types: non-preferential to non-preferential residues (N to N), non-

preferential to preferential residues (N to P), preferential to non-preferential residues (P to N) and

preferential to preferential residues (P to P) (Supplementary Table 5). Mutations at MHC contact

position mutation were not examined because, as above, the numbers of these types of mutations were

too small to achieve statistical significance. Mutations at TCR contact regions was also not examined

because no residue preferences were found at these sites (see above).

We analyzed the proportion of these 4 types of mutation in the immunogenic and non-immunogenic

datasets respectively. Interestingly, almost all (26 in 27) of the immunogenic neoantigens appear to be

in the class of N to P preference (Supplementary Table 5; Fig. 4e). Statistical analysis shown that the N

to P preference frequency of immunogenic neoantigens with mutations at anchor positions was

significant higher (Fig. 4e, $p<0.001$) than that for non-immunogenic neoantigens'. The other classes of

preference did not show significant differences between the immunogenic and non-immunogenic

neoantigens. Analyzing our results in more depth we found that one immunogenic neoantigen,

MYADM, in the immunogenic peptide collection did not follow the N to P preference at its C-terminal

binding site for MHC. However, the original paper describing this immunogenic neoantigen showed

that both the wild type and the mutant peptides of MYADM could elicit self T cell responses(40),

suggesting that the wild type peptide, with its C terminal anchor, was binding the patient's MHC

protein and immunogenic anyway. Thus, this neoantigen may not be a good target for further clinical

trial and can be regarded as an exception in our study.

With the classification of 4 types of neoantigen mutations changes at anchor positions, 96% (100% if

the exception MYADM is not counted) of the immunogenic neoantigen mutations belong to the class of

N to P preference and our detailed study of the KRAS G12D therapeutic neoantigen with the mutation

from the non-preferential residue G to the preferential residue D (N to P) is one of the clear examples.

That is, neoantigens with mutations at anchor positions not belonging to the class of N to P preference

would be very unlikely to be immunogenic and our detailed study of the DPAGT1 V213L non-

therapeutic neoantigen with the mutation from preferential residue V to the preferential residue L is one

of the clear examples. Thus it could be a clear indicator for the exclusion of non-immunogenic

neoantigen in the process of neoantigen prediction. From our NND dataset containing non-

immunogenic neopeptides predicted to bind relevant HLAs, about 42% of non-immunogenic

neoantigens with mutations at anchor positions were not N to P preference and could be eliminated

during immunogenic neoantigen prediction with this indicator (Fig. 4f). Thus, we suggest that the

processes we describe here could greatly help prediction of functional neoantigens for cancer

immunotherapy.

HLA-C*05:01 might be able to present KRAS G12D 10mer peptide

Because the frequency of KRAS mutations in tumors is so high, mutations in this protein are attractive

targets for immunotherapy. So far, it is thought that targeting the KRAS G12D neoantigen would be

useful only for patients expressing HLA-C*08:02. To expand use of the KRAS mutant we developed a

strategy to screen other HLA-C alleles for their ability to bind KRAS G12D and thus have the potential

to elicit antitumor responses.

Our strategy to identify new HLA candidates is divided into four steps. First, phylogenetic analysis was

performed for the 15 different HLA-C alleles with relatively high frequency in human*(76)*

(Supplementary Table 6; Supplementary Fig. 5). In the phylogenetic tree, three alleles (C*07:01,

C*07:02 and C* 17:01) were distantly related to the reference allele, C*08:02, and were excluded from

further analyses (Supplementary Fig. 5). The relative distance of the remaining alleles to C*08:02 was

calculated based on the phylogenetic tree. Secondly, their relative distance from C*08:02 was

normalized as shown on the heatmap (Fig. 5a). This showed that C*05:01 is closely related to C*08:02

and might therefore be able to present the KRAS neoantigen. Third, sequence alignment combined with

structure analyses was used to compare HLA-C*08:02 and other HLA-C alleles. Comparison of P3 and

PΩ binding motif of HLA-C alleles shown that C*05:01 is the only allele have the same sequence as

C*08:02 at its P3 anchor binding motif (Supplementary Table 7; Supplementary Fig. 6). Moreover

C*05:01 differed from C*08:02 by only two amino acids (at positions 77 and 80 on the α1 helix) at its

PΩ binding motif. Because of these observations we decided to test whether the changes would affect

the ability of C*05:01 to present the KRAS G12D peptide. We performed structural analyses based on

our solved HLA-C*08:02 structure (C08-mut9m).

Three residues from HLA -C*08:02 (position 97, position 99 and position 156) bind P3 of the peptide.

Two of these are arginine amino acids which promote engagement of the peptide acidic residue (Fig.

5b; Supplementary Table 7) and thus may play important roles in presenting acidic P3 mutated

neoantigen. These three amino acids are the same in C*05:01. The PΩ of peptide contacts amino acids

77, 80, 84, 143 and 146 in C*08:02 (Fig. 2c). Three of these (positions 84, 143, and 146) are invariable

in most HLA-C alleles (Supplementary Table 7). The two amino acids at positions 77 and 80 are also

the same in many of the HLA-C alleles but C*05:01 and several other HLA-C alleles differ from

C*08:02 by S77N and N80K. Amino acids at these positions pointed toward the TCR region and

peptide respectively (Fig. 5c). However, these five residues may not affect the presentation of peptides

which use leucine as a c-terminal residue because, regardless of the amino acids they express at these

five positions, most of HLA-C alleles select leucine as a preferential residue at the PΩ position of

peptide (Supplementary Fig. 4). Thus, after these structural comparisons of HLA-C*05:01 and HLA-

C*08:02, we inferred that HLA-C*05:01 might also present mutant KRAS G12D peptide and therefore

might be of potential use for T cell-based immunotherapy. We came to this conclusion in spite of the

fact that peptide binding predictions suggest that HLA-C*05:01 would bind with only low affinity to

both KRAS G12D 9mer and 10mer mutated peptides (Supplementary Table 2).

We next evaluated whether KRAS G12D can really be presented by C*05:01. We expressed C*05:01

heavy chain and β2M in E. coli and refolded them with 4 peptides: GADGVGKSA (mut9m, mutation

site is indicated with the amino acid underlined); KRAS G12D 10mer peptide GADGVGKSAL

(mut10m); wild type KRAS G12D 9mer peptide GAGGVGKSA (wt9m) and KRAS G12D 10mer

peptide GAGGVGKSAL (wt10m). Only the mutant KRAS G12D 10mer peptide successfully refolded

with HLA-C*05:01, suggesting that HLA-C*05:01 and HLA-C*08:02 are partially differ in their

antigen presentation abilities. We solved the crystal structure of C*05:01 in complex with the KRAS

G12D 10mer peptide to a resolution of 1.9 angstroms (Supplementary Table 3). High quality resolution

and electron density maps shown reliable placement of the 10mer peptide in C*05:01 complex (Fig.

5d). The TCR contact positions P5, P6, P7 and P9 of the peptides are flexible and can't form stable

conformations (Fig.5e). Nevertheless, the structure of the C05-mut10m closely resembles that of C08-

mut10m at the P1-P4 peptide region (Fig. 5e-g). Overlay of the C05-mut10m and C08-mut10m peptide

regions shows that an exception occurs at the P5-PΩ region of the peptides and position 77 and 80 of

the HLA proteins (Fig. 5e). The peptide P8S positions in both the C05-mut10m and C08-mut10m act

as axillary anchors. Differences also occur at the P10L position. Although C*05 and C*08 differ at

position 77, the Asn77 of C05 and the Ser77 of C08 form hydrogen bonds with P10L in a similar way.

However, Asn80 from C*05 binds the oxygen of P10L while Lys80 from C*08 does not with the

oxygen of P10L shifting towards the a2 helix and thus losing one of the hydrogen bonds from Tyr84

(Fig. 5h). These results explain to some extent why the GA<u>D</u>GVGKSA (mut9m) peptide did not

successfully refold with C*05:01. The most likely cause is the change at position 80, which will

weaken the interaction between the 9mer peptide and C*05:01, thus reducing the likelihood of forming

a stable complex. Excluding these differences, the peptide binding grooves of the HLA-C proteins had

relatively conserved interactions with the KRAS G12D 10mer peptide. It is, however, possible that

position 80 is located in the interface of the pMHC and T cell receptor, and there may change the

morphology of the epitope and impact T cell selection.

In line with our hypothesis, the structure of HLA-C*05:01 in complex with KRAS G12D indicates that

C*05:01 can also present the KRAS G12D neoantigen but in a slightly different way than C*08:02

does. Therefore HLA-C*05:01 might also be a potential target for T cell-based cancer immunotherapy.

An optimized procedure of neoepitope prediction

Nowadays, most in silico neoepitope (neoantigen and corresponding HLAs) prediction methods have

used neopeptide binding affinity to discover neoantigens for cancer immunotherapy. Although binding

to MHC-I is currently the most popular computational filter for removing nonantigenic neopeptides, it

still not robust enough to eliminate most of these candidates.

Our results emphasize the importance of anchor mutated neopeptides in cancer immunotherapy. We

demonstrated that almost all of anchor mutated neoantigens which have induced autologous T cell

response and are clinically effective have a mutation the demonstrates an N to P preference at anchor

positions. Thus, we suggest an optimized procedure that will improve the accuracy of neoepitope

prediction methods. The workflow of this procedure is shown in Fig. 6a. Tumor and normal samples

are first obtained from cancer patient for genomic DNA and RNA sequencing and tumor infiltrating

lymphocytes (TILs) isolation. Whole-exome sequencing analysis allows discovery of somatic

mutations present in the tumor cells, but not in normal cells. *In silico* binding prediction is performed

and can output a list of candidate peptides. The novelty of our procedure is in the following steps.

Neoantigen candidates will be classified into 3 types: MHC anchor mutation (contact MHC and is vital

for binding MHC), MHC contact position mutation (contact MHC but contribute much less) or TCR

contact position mutation. Each type may use a different filtration strategy during neoantigen candidate

prediction (Fig. 6b). This procedure may eliminate at least 42% of the anchor position mutated

candidates. Also, the similarity score could be used to help judge the possibility that the TCR contact

position mutated neopeptides will be immunogenic. These candidates can be further tested *in vitro* and

later used in clinical trials. Meanwhile, a validated neoantigen can be used to identify other HLA alleles

that might present it, as shown in our analyses of KRAS G12D and HLA-C*05:01.

Discussion

Recently T cells targeting tumor antigens, especially neoantigens, have proved to be one of the most

promising new methods of treating cancers. However, few investigations have reported the structures

of T cells reacting with cancer neoantigens. To fill this gap, here we solved the structure of one of the

most successful clinical cancer immunotherapy neoantigens, KRAS with the G12D mutation, presented

by HLA-C*0802. We found that the G12D mutation in these neoantigens (9mer and 10mer) drive

intensive interactions to MHC via acting as an unconventional anchor residue. This observation of the

mutation at the anchor position which cause the epitope to be immunogenic illustrates the complexity

of neoantigens beyond the previous evidences that mutations which were solvent-exposed and

therefore accessible to TCRs were more likely to be immunogenic. We next studied the structure of a

non-immunogenic neoantigen, DPAGT1 V213L discovered from previously publication*(19)*. Although

this peptide also had a mutation at the anchor position, it did not change the conformation of the TCR

contact region. Thus different types of mutations in the anchor position of neoantigens would lead to

different fates for immunogenicity.

To expand our findings more broadly and deeply we combined our structural work with the

bioinformatics research in which the latter examined all available clinical data comparing cancer

neopeptides that had proved to be immunogenic or non-immunogenic. Both types of mutated peptides

from IND and NND were assigned to one of three different categories, those that contained mutations

in anchor positions for MHC, or mutations in TCR contact regions or those that were mutated in

contact regions with MHC that were not in anchor positions. Based on our results, mutations prefer to

occur at TCR contact positions rather than occur at MHC contact position (contact MHC but not anchor

position) in IND. However, mutation frequency at anchor position did not show significant difference

between IND and NND. And in fact, whether mutations in peptide anchor positions are important for

the immunogenicity of the neoantigen was previously unclear. Different pieces of evidence led to

different conclusions(19, 20, 77, 78). These assignments took into account the fact that the peptide

anchor amino acids differ according to MHC allele with which they are engaged and the MHC anchor

mutations were further categorized by whether or not the changed amino acids would improve binding

of the peptide to MHC (N to P changes), or reduce binding to MHC (P to N changes) or would not

affect the ability of the peptide to bind MHC (N to N or P to P changes). Our analyses revealed a

remarkable finding, for the immunogenic anchor position mutated neoantigens nearly all the changes

fall into the non-preferential to preferential, N to P, category (100% if the exception one MYADM,

which is always reactive no matter it has the mutation or not, was not counted). Thus this indicator are

clear and practical for identifying anchor mutated neoantigens, called as "Non-preferential to

Preferential Conversion (NPC)". The amino acid turning from non-preferential to preferential may

create a new epitope which never seen by self-immune system. Given the fact that during the prediction

of viral antigen, most amino acids in the peptide are different from self-peptide, such indicator above is

not necessary for consideration. Hence, we believe NPC is an effective supplement indicator to

eliminate non-immunogenic peptides from cancer mutaome *in silico*.

The way of neoantigen study and the rules for immunogenic neoantigens could be very different with

normal antigens. Neoantigens are the altered-self between self-peptides and totally different foreign

peptides. In conventional antigen immunogenicity research, how amino acids of multiple positions get

together to interact with MHC and be recognized by TCR are main points. However, in neoantigen

research, how one residue change could reshape the peptide interactions with MHC and/or reshape

TCR contact region as to break the immune tolerance are main points. Thus, mutation positions,

mutation types, surrounding elements affecting mutations or affected by mutations and etc. should be

paid more attentions. More specific bioinformatic strategies and more specific rules for immunogenic

neoantigens are needed for research. In our study, certain rules were revealed only after large scale

studies on human tumor mutant peptides and careful classification according to the mutation position in

the peptides, which shows that the simple presence of a peptide mutated amino acid at an MHC binding

site is not predictive. Only if the mutation follows the NPC indicator does the change at an anchor

position become valuable.

On a related subject, here we tackled the issue of whether MHC alleles other than C*08:02 might be

useful in presenting the common KRAS G12D mutation as a neoantigen. We found that an MHC allele

that is closely related to C*08:02, C*05:01 might be such a possibility. Our structural work showed that

C*05:01 could indeed bind the KRAS G12D 10mer peptide and the configuration of the structure we

observed was quite similar to that of the same peptide bound to C*08:02. Thus, KRAS G12D mutation

could also be a good neoantigen in HLA-C*05:01 positive patients and the strategy we used to discover

this could be used to help enlarge knowledge of other suitable HLA alleles thus increasing the size of

the patient population in which such neoantigens could be used.

Our data suggest three possible immunogenic neoantigen models (Supplementary Fig. 7). Model A

represents the situation which peptide mutation occurs at TCR contact region and therefore appears

directly foreign to T cells. In this case, the neoantigens might be strongly immunogenic if the

corresponding wild type peptide is greatly dissimilar at the mutated site. Important differences might be

amino acid properties such as hydrophobicity, charge and polarity. Model B represents the situation in

which peptide mutation occurs at an MHC contact region (not including anchor sites). We found that

mutations at these sites are less likely to render the neoantigen immunogenic. However, our curated

data did show some bias for mutation at such regions, due to the relatively limited sample size

available now for this category, the relationship for such between mutation and immunogenicity is not

clear yet. Such mutants may still be immunogenic under certain conditions, if the mutant fits into an

MHC pocket of dramatically changes the TCR contact surface of the pMHC complex. Model C

represents the situation which the peptide mutation occurs at an MHC anchor position and increases

MHC binding.

Overall, more structural and biochemical experiments, with inclusion of more MHC alleles are needed

to improve bioanalytical predictive power in the important search for useful cancer neoantigens to treat

these devastating diseases. We hope the methods proposed here will help guide such future studies.

Materials and methods

Study design

This study was designed to use x-ray crystallography to examine the difference between immunogenic

and non-immunogenic mutant peptides bound to MHCI. On the basis of these results, we also took

bioinformatic analysis of our manually curated data form publications to see whether immunogenic

peptides have a certain character for neoantigen prediction. Moreover, we also performed phylogenetic

and structural analyses to extended neoantigen application.

Protein Expression, refolding and purification

The DNA encoding MHC heavy chain (HLA-C*08:02, HLA-C*05:01 and H-2 Kb) and light chain

(human β2M and mouse β2M) were synthesized (Idobio) and cloned into pET-22(b) vector (Novagen).

The vectors were transformed into the E. coli strain BL21 DE3 (Novagen). Transformants were

selected from Lurian broth (LB) agar plates containing ampicillin. A single colony was selected and

cultured in LB fluid medium with the antibiotics listed above at 37°C. Upon reaching an optical density

OD600 of 0.6, expression was induced with the addition of 1mM IPTG. Incubation continued at 37°C

for 5h. The cells were harvested by centrifugation and then resuspended in PBS buffer with 1 mM

PMSF at 4°C. The cells were lysed, and the lysate was clarified by centrifugation at 10,000 g to collect

inclusion bodies. Inclusion bodies were harvested and solubilized in 20 mM Tris (Vetec) pH 8.0, 8 M

urea (Vetec), 1 mM EDTA (BBI life sciences), 1 mM DTT (Sinopharm chemical reagent) and 0.2 mM

PMSF (Sinopharm chemical reagent).

Refolding was performed in the presence of MHC heavy chain, β2M and peptides as described

previously*(79)*. Briefly, the resolubilized heavy chain and light chain in the presence of the

corresponding peptide were added into 1 liter of refolding buffer. Refolding buffer contains 100 mM

Tris (pH 8.4), 0.5 mM oxidized glutathione (BBI life sciences), 5 mM reduced glutathione (BBI life

sciences), 400mM L-arginine (Vetec), 2mM EDTA (BBI life sciences). After 48h refolding, the 1 L

mixture dialyzed against 15 liters of 10 mM Tris buffer (pH 8.0) at 4 °C. for 24 h.

Soluble proteins were first purified by anion exchange chromatography with Q Sepharose HP (GE

Healthcare) column and Mono Q column (GE Healthcare). The purified protein sample was

concentrated by tangential flow filtration using Amicon Ultra centrifugal filters (Merck). For

desalination and purification, sample was loaded onto a Superdex 200 increase 10/300 GL column (GE

Healthcare) for size exclusion chromatography. Chromatography performed with BioLogic DuoFlow

system (Bio-rad) at a flow rate of 1 ml/min. Peak analysis was performed using the ASTRA software

package (BioLogic Chromatography Systems).

Crystallization, data collection and processing

Purified pMHC complex were concentrated to 10 mg/ml before the crystallization trials. For the HLA

complex, crystallizations were done by the sitting drop vapor diffusion technique at 4 °C.

Crystallization screens were performed using a series of kits from Hampton Research. Single crystal of

C08-mut9m, C08-mut10m and C05-mut10m complex were obtained in the condition of 0.2 M

ammonium acetate, 0.1 M HEPES (pH 7.5), 25% w/v polyethylene glycol 3,350. For the H-2 Kb

complex, single crystal of Kb-8mV and Kb-8mL complex were obtained when 4% v/v Tacsimate (pH

7.0), 12% w/v Polyethylene glycol 3,350 was used as the reservoir buffer.

Before data collection, crystals were transferred to crystallization buffer containing 20% (w/v) glycerol.

Next, the crystals were flash-cooled in liquid nitrogen immediately. The diffraction data were collected

at the Shanghai Synchrotron Radiation Facility (Shanghai, China) on beam line

BL17U1/BL18U1/BL19U1. The diffraction data were processed using the iMosflm program*(80)*, and

data reduction was performed with Aimless Pointless in the CCP4 software suite*(81)*. All structures

were determined by molecular replacement using the program Phaser*(82)*. The models from the

molecular replacement were built using the COOT (Crystallographic Object-Oriented Toolkit)

program*(83)* and subsequently subjected to refinement using Phenix software*(84)*. Data collection,

processing, and refinement statistics are summarized in (Supplementary Table 3). All the structural

figures were prepared using PyMOL program. The atomic coordinates and structure factors for the

reported crystal structures have been deposited on the Protein Data Bank (PDB;

http://www.rcsb.org/pdb/).

Neoantigens and non-immunogenic neopeptides data collection and correction.

Neoantigens and non-immunogenic neopeptides were manually curated with their corresponding HLA

alleles from 34 published literatures and recorded in immunogenic neoantigen dataset, IND and non-

immunogenic neopeptide dataset, NND.

To construct the immunogenic dataset IND, we collected 128 minimal peptides come from cancer

somatic mutations and their corresponding HLA class I alleles that reported to elicit T cell response in

humans. Those tumor neopeptides with uncertain response or not provided with minimal sequence

were not collected in our dataset. Each entry includes following information: tumor type, gene symbol,

wild type/ mutant peptide sequence, wild type/mutant peptide predicted binding affinity(31), single

amino acid change, HLA allele, and PubMed ID from NCBI.

For non-immunogenic dataset NND, we collected 11739 peptides (not the minimal peptide) from

cancer somatic mutations and their corresponding HLA class I alleles that reported not elicit T cell

response in humans. Each entry includes following information: tumor type, gene symbol, wild type/

mutant peptide sequence, wild type/mutant peptide predicted binding affinity(31), single amino acid

change, HLA allele, and PubMed ID from NCBI.

Construction of non-immunogenic neopeptide dataset (NND)

In order to generate minimal non-immunogenic neopeptide dataset, binding prediction was taken by

NetMHCpan 4.0 Server(67) based on the non-immunogenic neopeptide dataset (NND). Briefly,

different length of mutant peptides from NND were input in NetMHCpan 4.0 Server with custom

python scripts. The mutant peptides in 9- or 10-mer length with binding affinity under 500 nM (IC50)

were considered to be binders and then recorded in the IND. Related peptide information was added to

each entry like IND dataset.

Similarity comparison

Similarity comparison of wild type and mutant amino acids were taken based on normalized

BLOSUM62 matrix as described previously(74). This matrix provides a value from 0 to 1 for

representing the similarity of amino acid pairs (higher value indicates more similar). Tails of the violins

to the range of the data were not trimmed.

HLA present tendentiousness determination

Amino acid half-bit log-odds score in every position for different HLA alleles was calculated as

described by Morten *et al(86)*. Briefly, frequency of amino acid on distinct position was firstly

computed using Hobonm 1 algorithm to decrease the weight of similar sequence before calculation.

It is worth to note that, in some cases, if the sample size is too large, it may include similar sequences,

even duplicate sequences. Thus, clustering should be performed at first. For example, if one cluster

include 10 similar sequences, the weight of every sequence in this cluster will be 1/10. The similarity

score was the maximum percent of the same amino acids between one current sequence and every

sequence in one cluster, and the threshold of default similarity to decide whether current sequence

belongs to a cluster is 0.63.

In other cases, small sample related to amino acid depletion. Thus, we make use of pseudo counts to

solve this problem. The formula to calculate the pseudo counts is

610
$$g_i = \sum_j f_j \times q_{i|j}$$

611   where f_j is the real frequency of the amino acid j and $g_{i|j}$ is frequency of amino acid i replace amino

612   acid j based on BLOSUM62 matrix*(74)*.

613   After that, the estimated frequency is calculated according to equation

614
$$P_i = \frac{\alpha \cdot f_i + \beta \cdot g_i}{\alpha + \beta}$$

615   where α is the number of clusters subtracted by one. The default value β is 200.

616   Next, the half-bit log odds score is defined as

617
$$W_{i,a} = 2 \cdot \log_2 \frac{p_{i,a}}{q_a}$$

618   where q_a is the background frequency of amino acids. In our case, an equiprobable background amino

619   acid distribution is used.

620   The merged half-bits of one peptide is the summing half-bit score of amino acids in this peptide.

621   Statistical test of wild type peptides and mutant peptides merged half-bits from IND were tested using

622   t-test.

Nonameric peptide library obtained, peptide library sequence logo generation and HLA anchor

position preferential determination

The nonameric peptide library of 30 HLA alleles was downloaded from IEDB*(87)*. Sequence logos

were generated base on peptides library using the sequence logo generator*(88)*. Threshold for

preferential amino acids at anchor position of each HLA alleles was applied to be above 10%*(75)*

(include 10%) based on the data from nonameric peptide library.

HLA-C allele's frequency verification, phylogenetic analysis and heat map comparison

HLA-C allele frequency was verified from Allele Frequency Net Database (AFND;

http://www.allelefrequencies.net)*(76)*. HLA-C alleles sequences were obtained from The IPD and IPD-

IMGT/HLA Database*(89)*. Multiple sequence alignment was performed using MUSCLE (v3.8.31 for

windows)*(90)*. The input file and output file format were defaulted. We constructed phylogenetic trees

using MrBayes (v3.2.5 for windows) by following the instruction of 'Quick Start Version' section in

manual*(91)*.

For display the allele similarities on the heat map, we read the phylogenetic tree using R package ape,

and deleted some distantly related alleles (C*07:01, C*07:02, C*17:01). Similarity calculation was

performed according to the distance of each HLA allele to C*08:02. The similarity was normalized

between [0,1] and the heat map was plotted using R package heat map.

Statistical analysis

All data handling and nonlinear regression analyses were done using R, Microsoft Office Excel 2016

and GraphPad prism 6. P-values for difference were calculated using student's t test and Z-test.

References and Notes

1. M. D. Vesely, M. H. Kershaw, R. D. Schreiber, M. J. Smyth, Natural Innate and Adaptive Immunity

to Cancer, *Annu. Rev. Immunol.* **29**, 235–271 (2011).

2. Y. Sykulev, M. Joo, I. Vturina, T. J. Tsomides, H. N. Eisen, Evidence that a Single Peptide–MHC
Complex on a Target Cell Can Elicit a Cytolytic T Cell Response, *Immunity* **4**, 565–571 (1996).

3. P. C. Tumeh, C. L. Harview, J. H. Yearley, I. P. Shintaku, E. J. M. Taylor, L. Robert, B. Chmielowski,
M. Spasic, G. Henry, V. Ciobanu, A. N. West, M. Carmona, C. Kivork, E. Seja, G. Cherry, A. J. Gutierrez,
T. R. Grogan, C. Mateus, G. Tomasic, J. A. Glaspy, R. O. Emerson, H. Robins, R. H. Pierce, D. A. Elashoff,
C. Robert, A. Ribas, PD-1 blockade induces responses by inhibiting adaptive immune resistance,
Nature **515**, 568–571 (2014).

4. T. N. Schumacher, R. D. Schreiber, Neoantigens in cancer immunotherapy, *Science* **348**, 69–74
(2015).

5. N. A. Rizvi, M. D. Hellmann, A. Snyder, P. Kvistborg, V. Makarov, J. J. Havel, W. Lee, J. Yuan, P.
Wong, T. S. Ho, M. L. Miller, N. Rekhtman, A. L. Moreira, F. Ibrahim, C. Bruggeman, B. Gasmi, R.
Zappasodi, Y. Maeda, C. Sander, E. B. Garon, T. Merghoub, J. D. Wolchok, T. N. Schumacher, T. A.
Chan, Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer,
Science **348**, 124–128 (2015).

6. R.-F. Wang, H. Y. Wang, Immune targets and neoantigens for cancer immunotherapy and
precision medicine, *Cell Res.* **27**, 11–37 (2017).

7. M. Yadav, S. Jhunjhunwala, Q. T. Phung, P. Lupardus, J. Tanguay, S. Bumbaca, C. Franci, T. K.
Cheung, J. Fritsche, T. Weinschenk, Z. Modrusan, I. Mellman, J. R. Lill, L. Delamarre, Predicting
immunogenic tumour mutations by combining mass spectrometry and exome sequencing, *Nature*
515, 572–576 (2014).

8. P. G. Coulie, B. J. Van den Eynde, P. van der Bruggen, T. Boon, Tumour antigens recognized by T
lymphocytes: at the core of cancer immunotherapy, *Nat. Rev. Cancer* **14**, 135–146 (2014).

9. N. R. J. Gascoigne, V. Rybakin, O. Acuto, J. Brzostek, TCR Signal Strength and T Cell Development,
Annu. Rev. Cell Dev. Biol. **32**, 327–348 (2016).

10. E. Tran, P. F. Robbins, Y.-C. Lu, T. D. Prickett, J. J. Gartner, L. Jia, A. Pasetto, Z. Zheng, S. Ray, E. M.
Groh, I. R. Kriley, S. A. Rosenberg, T-Cell Transfer Therapy Targeting Mutant KRAS in Cancer, *N. Engl.*
J. Med. **375**, 2255–2262 (2016).

11. P. A. Ott, Z. Hu, D. B. Keskin, S. A. Shukla, J. Sun, D. J. Bozym, W. Zhang, A. Luoma, A. Giobbie-
Hurder, L. Peter, C. Chen, O. Olive, T. A. Carter, S. Li, D. J. Lieb, T. Eisenhaure, E. Gjini, J. Stevens, W.
J. Lane, I. Javeri, K. Nellaiappan, A. M. Salazar, H. Daley, M. Seaman, E. I. Buchbinder, C. H. Yoon, M.
Harden, N. Lennon, S. Gabriel, S. J. Rodig, D. H. Barouch, J. C. Aster, G. Getz, K. Wucherpfennig, D.
Neuberg, J. Ritz, E. S. Lander, E. F. Fritsch, N. Hacohen, C. J. Wu, An immunogenic personal
neoantigen vaccine for patients with melanoma, *Nature* **547**, 217–221 (2017).

12. U. Sahin, E. Derhovanessian, M. Miller, B.-P. Kloke, P. Simon, M. Löwer, V. Bukur, A. D. Tadmor,
U. Luxemburger, B. Schrörs, T. Omokoko, M. Vormehr, C. Albrecht, A. Paruzynski, A. N. Kuhn, J. Buck,

S. Heesch, K. H. Schreeb, F. Müller, I. Ortseifer, I. Vogler, E. Godehardt, S. Attig, R. Rae, A. Breitkreuz, C. Tolliver, M. Suchan, G. Martic, A. Hohberger, P. Sorn, J. Diekmann, J. Ciesla, O. Waksmann, A.-K. Brück, M. Witt, M. Zillgen, A. Rothermel, B. Kasemann, D. Langer, S. Bolte, M. Diken, S. Kreiter, R. Nemecek, C. Gebhardt, S. Grabbe, C. Höller, J. Utikal, C. Huber, C. Loquai, Ö. Türeci, Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer, *Nature* **547**, 222–226 (2017).

13. P. F. Robbins, Y.-C. Lu, M. El-Gamil, Y. F. Li, C. Gross, J. Gartner, J. C. Lin, J. K. Teer, P. Cliften, E. Tycksen, Y. Samuels, S. A. Rosenberg, Mining exomic sequencing data to identify mutated antigens recognized by adoptively transferred tumor-reactive T cells, *Nat. Med.* **19**, 747–752 (2013).

14. E. Tran, P. F. Robbins, S. A. Rosenberg, "Final common pathway" of human cancer immunotherapy: targeting random somatic mutations, *Nat. Immunol.* **18**, 255–262 (2017).

15. T. Wolfel, M. Hauer, J. Schneider, M. Serrano, C. Wolfel, E. Klehmann-Hieb, E. De Plaen, T. Hankeln, K. Meyer zum Buschenfelde, D. Beach, A p16INK4a-insensitive CDK4 mutant targeted by cytolytic T lymphocytes in a human melanoma, *Science* **269**, 1281 (1995).

16. H. Matsushita, M. D. Vesely, D. C. Koboldt, C. G. Rickert, R. Uppaluri, V. J. Magrini, C. D. Arthur, J. M. White, Y.-S. Chen, L. K. Shea, J. Hundal, M. C. Wendl, R. Demeter, T. Wylie, J. P. Allison, M. J. Smyth, L. J. Old, E. R. Mardis, R. D. Schreiber, Cancer exome analysis reveals a T-cell-dependent mechanism of cancer immunoediting, *Nature* **482**, 400–404 (2012).

17. N. van Rooij, M. M. van Buuren, D. Philips, A. Velds, M. Toebes, B. Heemskerk, L. J. A. van Dijk, S. Behjati, H. Hilkmann, D. el Atmioui, M. Nieuwland, M. R. Stratton, R. M. Kerkhoven, C. Keşmir, J. B. Haanen, P. Kvistborg, T. N. Schumacher, Tumor Exome Analysis Reveals Neoantigen-Specific T-Cell Reactivity in an Ipilimumab-Responsive Melanoma, *J. Clin. Oncol.* **31**, e439–e442 (2013).

18. P. F. Robbins, Y.-C. Lu, M. El-Gamil, Y. F. Li, C. Gross, J. Gartner, J. C. Lin, J. K. Teer, P. Cliften, E. Tycksen, Y. Samuels, S. A. Rosenberg, Mining exomic sequencing data to identify mutated antigens recognized by adoptively transferred tumor-reactive T cells, *Nat. Med.* **19**, 747–752 (2013).

19. M. Yadav, S. Jhunjhunwala, Q. T. Phung, P. Lupardus, J. Tanguay, S. Bumbaca, C. Franci, T. K. Cheung, J. Fritsche, T. Weinschenk, Z. Modrusan, I. Mellman, J. R. Lill, L. Delamarre, Predicting immunogenic tumour mutations by combining mass spectrometry and exome sequencing, *Nature* **515**, 572–576 (2014).

20. F. Duan, J. Duitama, S. Al Seesi, C. M. Ayres, S. A. Corcelli, A. P. Pawashe, T. Blanchard, D. McMahon, J. Sidney, A. Sette, B. M. Baker, I. I. Mandoiu, P. K. Srivastava, Genomic and bioinformatic profiling of mutational neoepitopes reveals new rules to predict anticancer immunogenicity, *J. Exp. Med.* **211**, 2231–2248 (2014).

21. C. J. Cohen, J. J. Gartner, M. Horovitz-Fried, K. Shamalov, K. Trebska-McGowan, V. V. Bliskovsky, M. R. Parkhurst, C. Ankri, T. D. Prickett, J. S. Crystal, Y. F. Li, M. El-Gamil, S. A. Rosenberg, P. F. Robbins, Isolation of neoantigen-specific T cells from tumor and peripheral lymphocytes, *J. Clin. Invest.* **125**, 3981–3991 (2015).

22. B. M. Carreno, V. Magrini, M. Becker-Hapak, S. Kaabinejadian, J. Hundal, A. A. Petti, A. Ly, W.-R.
Lie, W. H. Hildebrand, E. R. Mardis, G. P. Linette, A dendritic cell vaccine increases the breadth and
diversity of melanoma neoantigen-specific T cells, *Science* **348**, 803 (2015).

23. E. Tran, P. F. Robbins, Y.-C. Lu, T. D. Prickett, J. J. Gartner, L. Jia, A. Pasetto, Z. Zheng, S. Ray, E. M.
Groh, I. R. Kriley, S. A. Rosenberg, T-Cell Transfer Therapy Targeting Mutant KRAS in Cancer, *N. Engl.*
J. Med. **375**, 2255–2262 (2016).

24. E. Stronen, M. Toebes, S. Kelderman, M. M. van Buuren, W. Yang, N. van Rooij, M. Donia, M.-L.
Boschen, F. Lund-Johansen, J. Olweus, T. N. Schumacher, Targeting of cancer neoantigens with
donor-derived T cell receptor repertoires, *Science* **352**, 1337–1341 (2016).

25. V. Anagnostou, K. N. Smith, P. M. Forde, N. Niknafs, R. Bhattacharya, J. White, T. Zhang, V. Adleff,
J. Phallen, N. Wali, C. Hruban, V. B. Guthrie, K. Rodgers, J. Naidoo, H. Kang, W. Sharfman, C.
Georgiades, F. Verde, P. Illei, Q. K. Li, E. Gabrielson, M. V. Brock, C. A. Zahnow, S. B. Baylin, R. B.
Scharpf, J. R. Brahmer, R. Karchin, D. M. Pardoll, V. E. Velculescu, Evolution of Neoantigen Landscape
during Immune Checkpoint Blockade in Non–Small Cell Lung Cancer, *Cancer Discov.* **7**, 264–276
(2017).

26. U. Sahin, E. Derhovanessian, M. Miller, B.-P. Kloke, P. Simon, M. Löwer, V. Bukur, A. D. Tadmor,
U. Luxemburger, B. Schrörs, T. Omokoko, M. Vormehr, C. Albrecht, A. Paruzynski, A. N. Kuhn, J. Buck,
S. Heesch, K. H. Schreeb, F. Müller, I. Ortseifer, I. Vogler, E. Godehardt, S. Attig, R. Rae, A. Breitkreuz,
C. Tolliver, M. Suchan, G. Martic, A. Hohberger, P. Sorn, J. Diekmann, J. Ciesla, O. Waksmann, A.-K.
Brück, M. Witt, M. Zillgen, A. Rothermel, B. Kasemann, D. Langer, S. Bolte, M. Diken, S. Kreiter, R.
Nemecek, C. Gebhardt, S. Grabbe, C. Höller, J. Utikal, C. Huber, C. Loquai, Ö. Türeci, Personalized
RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer, *Nature* **547**,
222–226 (2017).

27. N. Zacharakis, H. Chinnasamy, M. Black, H. Xu, Y.-C. Lu, Z. Zheng, A. Pasetto, M. Langhan, T.
Shelton, T. Prickett, J. Gartner, L. Jia, K. Trebska-McGowan, R. P. Somerville, P. F. Robbins, S. A.
Rosenberg, S. L. Goff, S. A. Feldman, Immune recognition of somatic mutations leading to complete
durable regression in metastatic breast cancer, *Nat. Med.* **24**, 724–730 (2018).

28. E. Stronen, M. Toebes, S. Kelderman, M. M. van Buuren, W. Yang, N. van Rooij, M. Donia, M.-L.
Boschen, F. Lund-Johansen, J. Olweus, T. N. Schumacher, Targeting of cancer neoantigens with
donor-derived T cell receptor repertoires, *Science* **352**, 1337–1341 (2016).

29. M. Nielsen, C. Lundegaard, O. Lund, C. Keşmir, The role of the proteasome in generating
cytotoxic T-cell epitopes: insights obtained from improved predictions of proteasomal cleavage,
Immunogenetics **57**, 33–41 (2005).

30. M. V. Larsen, C. Lundegaard, K. Lamberth, S. Buus, O. Lund, M. Nielsen, Large-scale validation
of methods for cytotoxic T-lymphocyte epitope prediction, *BMC Bioinformatics* **8**, 424 (2007).

31. M. Andreatta, M. Nielsen, Gapped sequence alignment using artificial neural networks:
application to the MHC class I system, *Bioinformatics* **32**, 511–517 (2016).

32. M. M. Davis, P. Brodin, Rebooting Human Immunology, *Annu. Rev. Immunol.* **36**, 843–864 (2018).

33. A. Vitiello, M. Zanetti, Neoantigen prediction and the need for validation, *Nat. Biotechnol.* **35**,
815–817 (2017).

34. C. Kandoth, M. D. McLellan, F. Vandin, K. Ye, B. Niu, C. Lu, M. Xie, Q. Zhang, J. F. McMichael, M.
A. Wyczalkowski, M. D. M. Leiserson, C. A. Miller, J. S. Welch, M. J. Walter, M. C. Wendl, T. J. Ley, R.
K. Wilson, B. J. Raphael, L. Ding, Mutational landscape and significance across 12 major cancer
types, *Nature* **502**, 333–339 (2013).

35. C. H. June, Drugging the Undruggable Ras — Immunotherapy to the Rescue?, *N. Engl. J. Med.*
375, 2286–2289 (2016).

36. T. Kamisawa, L. D. Wood, T. Itoi, K. Takaori, Pancreatic cancer, *The Lancet* **388**, 73–85 (2016).

37. F. F. González-Galarza, L. Y. C. Takeshita, E. J. M. Santos, F. Kempson, M. H. T. Maia, A. L. S. da
Silva, A. L. T. e Silva, G. S. Ghattaoraya, A. Alfirevic, A. R. Jones, D. Middleton, Allele frequency net
2015 update: new features for HLA epitopes, KIR and disease and HLA adverse drug reaction
associations, *Nucleic Acids Res.* **43**, D784–D788 (2015).

38.   T-Cell Transfer Therapy Targeting Mutant KRAS, *N. Engl. J. Med.* **376**, e11 (2017).

39. M. Andreatta, M. Nielsen, Gapped sequence alignment using artificial neural networks:
application to the MHC class I system, *Bioinformatics* **32**, 511–517 (2016).

40. N. McGranahan, A. J. S. Furness, R. Rosenthal, S. Ramskov, R. Lyngaa, S. K. Saini, M. Jamal-
Hanjani, G. A. Wilson, N. J. Birkbak, C. T. Hiley, T. B. K. Watkins, S. Shafi, N. Murugaesu, R. Mitter, A.
U. Akarca, J. Linares, T. Marafioti, J. Y. Henry, E. M. Van Allen, D. Miao, B. Schilling, D. Schadendorf,
L. A. Garraway, V. Makarov, N. A. Rizvi, A. Snyder, M. D. Hellmann, T. Merghoub, J. D. Wolchok, S. A.
Shukla, C. J. Wu, K. S. Peggs, T. A. Chan, S. R. Hadrup, S. A. Quezada, C. Swanton, Clonal neoantigens
elicit T cell immunoreactivity and sensitivity to immune checkpoint blockade, *Science* **351**, 1463–
1469 (2016).

41. P. F. Robbins, A mutated beta-catenin gene encodes a melanoma-specific antigen recognized
by tumor infiltrating lymphocytes, *J. Exp. Med.* **183**, 1185–1192 (1996).

42. M. Guéguen, J.-J. Patard, B. Gaugler, F. Brasseur, J.-C. Renauld, P. J. Van Cangh, T. Boon, B. J. Van
den Eynde, An Antigen Recognized by Autologous CTLs on a Human Bladder Carcinoma, *J. Immunol.*
160, 6188 (1998).

43. K. T. Hogan, D. P. Eisinger, S. B. Cupp, K. J. Lekstrom, D. D. Deacon, J. Shabanowitz, D. F. Hunt, V.
H. Engelhard, C. L. Slingluff, M. M. Ross, The Peptide Recognized by HLA-A68.2-restricted,
Squamous Cell Carcinoma of the Lung-specific Cytotoxic T Lymphocytes Is Derived from a Mutated
Elongation Factor 2 Gene, *Cancer Res.* **58**, 5144 (1998).

44. E. Zorn, T. Hercend, A natural cytotoxic T cell response in a spontaneously regressing human
melanoma targets a neoantigen resulting from a somatic point mutation, *Eur. J. Immunol.* **29**, 592–

601 (1999).

45. R. Chiari, F. Foury, E. De Plaen, J.-F. Baurain, J. Thonnard, P. G. Coulie, Two Antigens Recognized
by Autologous Cytolytic T Lymphocytes on a Melanoma Result from a Single Point Mutation in an
Essential Housekeeping Gene, *Cancer Res.* **59**, 5785 (1999).

46. H. Echchakir, F. Mami-Chouaib, I. Vergnon, J.-F. Baurain, V. Karanikas, S. Chouaib, P. G. Coulie, A
Point Mutation in the α-Actinin-4 Gene Generates an Antigenic Peptide Recognized by Autologous
Cytolytic T Lymphocytes on a Human Lung Carcinoma, *Cancer Res.* **61**, 4078 (2001).

47. J. Zhou, M. E. Dudley, S. A. Rosenberg, P. F. Robbins, Persistence of Multiple Tumor-Specific T-
Cell Clones Is Associated with Complete Tumor Regression in a Melanoma Patient Receiving
Adoptive Cell Transfer Therapy:, *J. Immunother.* **28**, 53–62 (2005).

48. M. Sensi, G. Nicolini, M. Zanon, C. Colombo, A. Molla, I. Bersani, R. Lupetti, G. Parmiani, A.
Anichini, Immunogenicity without Immunoselection: A Mutant but Functional Antioxidant Enzyme
Retained in a Human Metastatic Melanoma and Targeted by CD8+ T Cells with a Memory
Phenotype, *Cancer Res* , 10 (2005).

49. V. Lennerz, M. Fatho, C. Gentilini, R. A. Frye, A. Lifke, D. Ferel, C. Wolfel, C. Huber, T. Wolfel, The
response of autologous T cells to a human melanoma is dominated by mutated neoantigens, *Proc.*
Natl. Acad. Sci. **102**, 16013–16018 (2005).

50. M. Takenoyama, J.-F. Baurain, M. Yasuda, T. So, M. Sugaya, T. Hanagiri, K. Sugio, K. Yasumoto, T.
Boon, P. G. Coulie, A point mutation in the NFYC gene generates an antigenic peptide recognized
by autologous cytolytic T lymphocytes on a human squamous cell lung carcinoma: NF-YC Mutated
Peptide-Reactive CTL in NSCLC, *Int. J. Cancer* **118**, 1992–1997 (2006).

51. D. Ito, C. Visus, T. K. Hoffmann, V. Balz, H. Bier, E. Appella, T. L. Whiteside, R. L. Ferris, A. B. DeLeo,
Immunological characterization of missense mutations occurring within cytotoxic T cell-defined
p53 epitopes in HLA-A*0201+ squamous cell carcinomas of the head and neck, *Int. J. Cancer* **120**,
2618–2624 (2007).

52. N. van Rooij, M. M. van Buuren, D. Philips, A. Velds, M. Toebes, B. Heemskerk, L. J. A. van Dijk,
S. Behjati, H. Hilkmann, D. el Atmioui, M. Nieuwland, M. R. Stratton, R. M. Kerkhoven, C. Keşmir, J.
B. Haanen, P. Kvistborg, T. N. Schumacher, Tumor Exome Analysis Reveals Neoantigen-Specific T-
Cell Reactivity in an Ipilimumab-Responsive Melanoma, *J. Clin. Oncol.* **31**, e439–e442 (2013).

53. D. A. Wick, J. R. Webb, J. S. Nielsen, S. D. Martin, D. R. Kroeger, K. Milne, M. Castellarin, K.
Twumasi-Boateng, P. H. Watson, R. A. Holt, B. H. Nelson, Surveillance of the Tumor Mutanome by
T Cells during Progression from Primary to Recurrent Ovarian Cancer, *Clin. Cancer Res.* **20**, 1125–
1134 (2014).

54. Y.-C. Lu, X. Yao, J. S. Crystal, Y. F. Li, M. El-Gamil, C. Gross, L. Davis, M. E. Dudley, J. C. Yang, Y.
Samuels, S. A. Rosenberg, P. F. Robbins, Efficient Identification of Mutated Cancer Antigens
Recognized by T Cells Associated with Durable Tumor Regressions, *Clin. Cancer Res.* **20**, 3401–3410
(2014).

55. C. J. Cohen, J. J. Gartner, M. Horovitz-Fried, K. Shamalov, K. Trebska-McGowan, V. V. Bliskovsky,
M. R. Parkhurst, C. Ankri, T. D. Prickett, J. S. Crystal, Y. F. Li, M. El-Gamil, S. A. Rosenberg, P. F. Robbins,
Isolation of neoantigen-specific T cells from tumor and peripheral lymphocytes, *J. Clin. Invest.* **125**,
3981–3991 (2015).

56. E. Tran, M. Ahmadzadeh, Y.-C. Lu, A. Gros, S. Turcotte, P. F. Robbins, J. J. Gartner, Z. Zheng, Y. F.
Li, S. Ray, J. R. Wunderlich, R. P. Somerville, S. A. Rosenberg, Immunogenicity of somatic mutations
in human gastrointestinal cancers, *Science* **350**, 1387–1390 (2015).

57. A. Snyder, V. Makarov, T. Merghoub, J. Yuan, J. M. Zaretsky, A. Desrichard, L. A. Walsh, M. A.
Postow, P. Wong, T. S. Ho, T. J. Hollmann, C. Bruggeman, K. Kannan, Y. Li, C. Elipenahli, C. Liu, C. T.
Harbison, L. Wang, A. Ribas, J. D. Wolchok, T. A. Chan, Genetic Basis for Clinical Response to CTLA-
4 Blockade in Melanoma, *N. Engl. J. Med.* **371**, 2189–2199 (2014).

58. S. Kalaora, E. Barnea, E. Merhavi-Shoham, N. Qutob, J. K. Teer, N. Shimony, J. Schachter, S. A.
Rosenberg, M. J. Besser, A. Admon, Y. Samuels, Use of HLA peptidomics and whole exome
sequencing to identify human immunogenic neo-antigens, *Oncotarget* **7** (2016),
doi:10.18632/oncotarget.6960.

59. A. Gros, M. R. Parkhurst, E. Tran, A. Pasetto, P. F. Robbins, S. Ilyas, T. D. Prickett, J. J. Gartner, J.
S. Crystal, I. M. Roberts, K. Trebska-McGowan, J. R. Wunderlich, J. C. Yang, S. A. Rosenberg,
Prospective identification of neoantigen-specific lymphocytes in the peripheral blood of melanoma
patients, *Nat. Med.* **22**, 433–438 (2016).

60. T. D. Prickett, J. S. Crystal, C. J. Cohen, A. Pasetto, M. R. Parkhurst, J. J. Gartner, X. Yao, R. Wang,
A. Gros, Y. F. Li, M. El-Gamil, K. Trebska-McGowan, S. A. Rosenberg, P. F. Robbins, Durable Complete
Response from Metastatic Melanoma after Transfer of Autologous T Cells Recognizing 10 Mutated
Tumor Antigens, *Cancer Immunol. Res.* **4**, 669–678 (2016).

61. M. Parkhurst, A. Gros, A. Pasetto, T. Prickett, J. S. Crystal, P. Robbins, S. A. Rosenberg, Isolation
of T-Cell Receptors Specifically Reactive with Mutated Tumor-Associated Antigens from Tumor-
Infiltrating Lymphocytes Based on CD137 Expression, *Clin. Cancer Res.* **23**, 2491–2505 (2017).

62. E. Tran, P. F. Robbins, Y.-C. Lu, T. D. Prickett, J. J. Gartner, L. Jia, A. Pasetto, Z. Zheng, S. Ray, E. M.
Groh, I. R. Kriley, S. A. Rosenberg, T-Cell Transfer Therapy Targeting Mutant KRAS in Cancer, *N. Engl.*
J. Med. **375**, 2255–2262 (2016).

63. V. Anagnostou, K. N. Smith, P. M. Forde, N. Niknafs, R. Bhattacharya, J. White, T. Zhang, V. Adleff,
J. Phallen, N. Wali, C. Hruban, V. B. Guthrie, K. Rodgers, J. Naidoo, H. Kang, W. Sharfman, C.
Georgiades, F. Verde, P. Illei, Q. K. Li, E. Gabrielson, M. V. Brock, C. A. Zahnow, S. B. Baylin, R. B.
Scharpf, J. R. Brahmer, R. Karchin, D. M. Pardoll, V. E. Velculescu, Evolution of Neoantigen Landscape
during Immune Checkpoint Blockade in Non–Small Cell Lung Cancer, *Cancer Discov.* **7**, 264–276
(2017).

64. S. Stevanović, A. Pasetto, S. R. Helman, J. J. Gartner, T. D. Prickett, B. Howie, H. S. Robins, P. F.
Robbins, C. A. Klebanoff, S. A. Rosenberg, C. S. Hinrichs, Landscape of immunogenic tumor antigens

in successful immunotherapy of virally induced epithelial cancer, *Science* **356**, 200–205 (2017).

65. D. T. Le, J. N. Durham, K. N. Smith, H. Wang, B. R. Bartlett, L. K. Aulakh, S. Lu, H. Kemberling, C.
Wilt, B. S. Luber, F. Wong, N. S. Azad, A. A. Rucki, D. Laheru, R. Donehower, A. Zaheer, G. A. Fisher,
T. S. Crocenzi, J. J. Lee, T. F. Greten, A. G. Duffy, K. K. Ciombor, A. D. Eyring, B. H. Lam, A. Joe, S. P.
Kang, M. Holdhoff, L. Danilova, L. Cope, C. Meyer, S. Zhou, R. M. Goldberg, D. K. Armstrong, K. M.
Bever, A. N. Fader, J. Taube, F. Housseau, D. Spetzler, N. Xiao, D. M. Pardoll, N. Papadopoulos, K. W.
Kinzler, J. R. Eshleman, B. Vogelstein, R. A. Anders, L. A. Diaz, Mismatch repair deficiency predicts
response of solid tumors to PD-1 blockade, *Science* **357**, 409–413 (2017).

66. D. C. Deniger, A. Pasetto, P. F. Robbins, J. J. Gartner, T. D. Prickett, B. C. Paria, P. Malekzadeh, L.
Jia, R. Yossef, M. M. Langhan, J. R. Wunderlich, D. N. Danforth, R. P. T. Somerville, S. A. Rosenberg,
T-cell Responses to *TP53* "Hotspot" Mutations and Unique Neoantigens Expressed by Human
Ovarian Cancers, *Clin. Cancer Res.* **24**, 5562–5573 (2018).

67. V. Jurtz, S. Paul, M. Andreatta, P. Marcatili, B. Peters, M. Nielsen, NetMHCpan-4.0: Improved
Peptide–MHC Class I Interaction Predictions Integrating Eluted Ligand and Peptide Binding Affinity
Data, *J. Immunol.* **199**, 3360–3368 (2017).

68. M. Matsumura, D. Fremont, P. Peterson, I. Wilson, Emerging principles for the recognition of
peptide antigens by MHC class I molecules, *Science* **257**, 927–934 (1992).

69. H.-G. Rammensee, J. Bachmann, N. P. N. Emmerich, O. A. Bachor, S. Stevanović, SYFPEITHI:
database for MHC ligands and peptide motifs, *Immunogenetics* **50**, 213–219 (1999).

70. K. Christopher Garcia, J. J. Adams, D. Feng, L. K. Ely, The molecular basis of TCR germline bias
for MHC is surprisingly simple, *Nat. Immunol.* **10**, 143 (2009).

71. L. Yin, J. Scott-Browne, J. W. Kappler, L. Gapin, P. Marrack, T cells and their eons-old obsession
with MHC, *Immunol. Rev.* **250**, 49–60 (2012).

72. L. Yin, E. Huseby, J. Scott-Browne, K. Rubtsova, C. Pinilla, F. Crawford, P. Marrack, S. Dai, J. W.
Kappler, A Single T Cell Receptor Bound to Major Histocompatibility Complex Class I and Class II
Glycoproteins Reveals Switchable TCR Conformers, *Immunity* **35**, 23–33 (2011).

73. J. J. Adams, S. Narayanan, M. E. Birnbaum, S. S. Sidhu, S. J. Blevins, M. H. Gee, L. V. Sibener, B.
M. Baker, D. M. Kranz, K. C. Garcia, Structural interplay between germline interactions and adaptive
recognition determines the bandwidth of TCR-peptide-MHC cross-reactivity, *Nat. Immunol.* **17**, 87
(2015).

74. S. Henikoff, J. G. Henikoff, Amino acid substitution matrices from protein blocks., *Proc. Natl.*
Acad. Sci. **89**, 10915–10919 (1992).

75. K. Falk, O. Rötzschke, S. Stevanovié, G. Jung, H.-G. Rammensee, Allele-specific motifs revealed
by sequencing of self-peptides eluted from MHC molecules, *Nature* **351**, 290–296 (1991).

76. F. F. González-Galarza, L. Y. C. Takeshita, E. J. M. Santos, F. Kempson, M. H. T. Maia, A. L. S. Da

Silva, A. L. Teles E Silva, G. S. Ghattaoraya, A. Alfirevic, A. R. Jones, D. Middleton, Allele frequency net 2015 update: New features for HLA epitopes, KIR and disease and HLA adverse drug reaction associations, *Nucleic Acids Res.* **43**, D784–D788 (2015).

77. E. F. Fritsch, M. Rajasagi, P. A. Ott, V. Brusic, N. Hacohen, C. J. Wu, HLA-Binding Properties of Tumor Neoepitopes in Humans, *Cancer Immunol. Res.* **2**, 522–529 (2014).

78. M. Yarchoan, B. A. Johnson, E. R. Lutz, D. A. Laheru, E. M. Jaffee, Targeting neoantigens to augment antitumour immunity, *Nat. Rev. Cancer* **17**, 209–222 (2017).

79. W. A. Macdonald, Z. Chen, S. Gras, J. K. Archbold, F. E. Tynan, C. S. Clements, M. Bharadwaj, L. Kjer-Nielsen, P. M. Saunders, M. C. J. Wilce, F. Crawford, B. Stadinsky, D. Jackson, A. G. Brooks, A. W. Purcell, J. W. Kappler, S. R. Burrows, J. Rossjohn, J. McCluskey, T Cell Allorecognition via Molecular Mimicry, *Immunity* **31**, 897–908 (2009).

80. T. G. G. Battye, L. Kontogiannis, O. Johnson, H. R. Powell, A. G. W. Leslie, *iMOSFLM* : a new graphical interface for diffraction-image processing with *MOSFLM*, *Acta Crystallogr. D Biol. Crystallogr.* **67**, 271–281 (2011).

81. M. D. Winn, C. C. Ballard, K. D. Cowtan, E. J. Dodson, P. Emsley, P. R. Evans, R. M. Keegan, E. B. Krissinel, A. G. W. Leslie, A. McCoy, S. J. McNicholas, G. N. Murshudov, N. S. Pannu, E. A. Potterton, H. R. Powell, R. J. Read, A. Vagin, K. S. Wilson, Overview of the *CCP* 4 suite and current developments, *Acta Crystallogr. D Biol. Crystallogr.* **67**, 235–242 (2011).

82. A. J. McCoy, R. W. Grosse-Kunstleve, P. D. Adams, M. D. Winn, L. C. Storoni, R. J. Read, *Phaser* crystallographic software, *J. Appl. Crystallogr.* **40**, 658–674 (2007).

83. P. Emsley, K. Cowtan, *Coot* : model-building tools for molecular graphics, *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132 (2004).

84. P. D. Adams, R. W. Grosse-Kunstleve, L.-W. Hung, T. R. Ioerger, A. J. McCoy, N. W. Moriarty, R. J. Read, J. C. Sacchettini, N. K. Sauter, T. C. Terwilliger, *PHENIX* : building new software for automated crystallographic structure determination, *Acta Crystallogr. D Biol. Crystallogr.* **58**, 1948–1954 (2002).

85. V. Jurtz, S. Paul, M. Andreatta, P. Marcatili, B. Peters, M. Nielsen, NetMHCpan-4.0: Improved Peptide–MHC Class I Interaction Predictions Integrating Eluted Ligand and Peptide Binding Affinity Data, *J. Immunol.* **199**, 3360–3368 (2017).

86. M. C. F. Thomsen, M. Nielsen, Seq2Logo: a method for construction and visualization of amino acid binding motifs and sequence profiles including sequence weighting, pseudo counts and two-sided representation of amino acid enrichment and depletion, *Nucleic Acids Res.* **40**, W281–W287 (2012).

87. R. Vita, J. A. Overton, J. A. Greenbaum, J. Ponomarenko, J. D. Clark, J. R. Cantrell, D. K. Wheeler, J. L. Gabbard, D. Hix, A. Sette, B. Peters, The immune epitope database (IEDB) 3.0, *Nucleic Acids Res.* **43**, D405–D412 (2015).

88. M. C. F. Thomsen, M. Nielsen, Seq2Logo: a method for construction and visualization of amino
acid binding motifs and sequence profiles including sequence weighting, pseudo counts and two-
sided representation of amino acid enrichment and depletion, *Nucleic Acids Res.* **40**, W281–W287
(2012).

89. J. Robinson, J. A. Halliwell, J. D. Hayhurst, P. Flicek, P. Parham, S. G. E. Marsh, The IPD and
IMGT/HLA database: allele variant databases, *Nucleic Acids Res.* **43**, D423–D431 (2015).

90. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput,
Nucleic Acids Res. **32**, 1792–1797 (2004).

91. F. Ronquist, M. Teslenko, P. van der Mark, D. L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M.
A. Suchard, J. P. Huelsenbeck, MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model
Choice Across a Large Model Space, *Syst. Biol.* **61**, 539–542 (2012).

92. E. Paradis, J. Claude, K. Strimmer, APE: Analyses of Phylogenetics and Evolution in R language,
Bioinformatics **20**, 289–290 (2004).

Figure Legends

Figure 1. Structural comparison of HLA-C*08:02 in complex with therapeutic KRAS G12D

9mer and 10mer neoantigens.

a-b, Unambiguous 2Fo-Fc electron density maps of (**a**) KRAS G12D 9mer (GA<u>D</u>GVGKSA, green)

and (**b**)10mer (GA<u>D</u>GVGKSAL, yellow) peptides from solved complex structures. The underlined

amino acids represent the mutation in the peptides.

c, Overlay of the KRAS G12D 9mer and 10mer peptides.

d, Overlay of Cα traces of HLA-C*08:02 in complex with 9mer and 10mer peptides. These two

structures show conformational differences at their peptide regions.

e, Sequence logo shows amino acid preference for 9mer peptide bounds to HLA-C*08:02. The peptide

library obtained from IEDB, n=2321. Sequence logo was generated from Seq2Logo server.

f, Aromatic residues (green) from HLA-C*08:02 accommodating P1 and P2 residues of KRAS G12D

9mer peptide.

Figure 2. Different presentation patterns between C08-mut9m and C08-mut10m

a, Polar interactions between the HLA-C*08:02 molecule (grey) and 9mer (green), or HLA-C*08:02

interacts with 10mer (yellow) peptide at P1G and P2A positions.

b, KRAS G12D 9mer peptide interact with HLA-C*08:02 with the side chains P3D and P9A pointing

down.

c, KRAS G12D 10mer peptide interact residues of HLA-C*08:02 with the side chains of P3D, P8S and

P9A pointing down.

Figure 3. Systematic analysis of IND and NND dataset.

a, Neoantigen predicted affinity ratio. Dots represents the ratio of predicted affinity of wild type

peptide and predicted affinity of mutant peptide. All tested neoantigens were from the IND (n=128).

Affinity prediction was performed using NetMHC 4.0pan. Orange dots (n=26) represent the

neoantigens with less than (or equal to) 500 nM predicted affinity. Green dots (n=102) represent the

neoantigens with more than 500 nM predicted affinity.

b, Comparation of the mutation occurrence frequency at each position of 9mer peptides from IND (red

line) and NND (dark blue line). The grey dotted line represents the mutation theoretical frequency (1/9)

of 9mer peptide. Positions of 9mer peptide show on horizontal axis.

c, Donut plots showing the percentage of 9mer peptide mutation distribution from IND and NND based

on the three different classes: anchor position mutation, MHC contact position mutation and TCR

contact position mutation. (immunogenic data, n =51; non-immunogenic data, n =763).

d, 9mer peptides mutation distribution of three different classes (anchor position mutation, MHC

contact position mutation and TCR contact position mutation) from IND and NND. The percentage of

mutation distribution at anchor position did not shown significant difference between immunogenic and

non-immunogenic data (ns, no significance). The frequency of mutation distribution at TCR contact

region and MHC contact region shown significant difference (*p<0.05, **P<0.01; immunogenic data,

n =51; non-immunogenic data, n =763, Z-test).

Figure 4. Similarity and preference of data from IND and NND.

a-c, a, Similarity between wild type and mutant amino acids from immunogenic and non-immunogenic

peptides. Normalized similarity score was calculated based on the BLOSUM62 matrix (ns, p=0.3077, n

(immunogenic data) =128; n (non-immunogenic data) =2883, student t-test). **b,** Similarity between

wild type and mutant amino acids from anchor position mutated immunogenic and non-immunogenic

peptides (**p=0.010, n (immunogenic data) =29; n (non-immunogenic data) =426, student t-test). **c,**

Similarity of amino acids from 9mer TCR contact position mutated immunogenic and non-

immunogenic peptides (**p<0.05, n (immunogenic data) =40; n (non-immunogenic data) =426, student

t-test).

d-f, d, Comparison of HLA present tendentiousness between wild type and mutant 9mer neoantigens

from IND (n=69, **p=0.0088, student t-test); **e,** Comparison of HLA present tendentiousness between

wild type and mutant 9mer anchor mutated neoantigens from IND (n=17, ****p<0.0001, student t-

test); **f,** Comparation of HLA present tendentiousness between wild type and mutant 9mer TCR contact

position mutated neoantigens from IND (n=39, p=0.9358, student t-test).

g, Frequency of 4 different anchor mutation classes (non-preferential to non-preferential residues (N to

N), non-preferential to preferential residues (N to P), preferential to non-preferential residues (P to N)

and preferential to preferential residues (P to P)) of immunogenic data (from IND) and non-

immunogenic data (from NND). Each class is representing the preference alteration from wild type

counterparts to mutant peptides. The frequency between immunogenic and non-immunogenic data on

N to P class shows significant difference (***p<0.001, n (immunogenic data) = 27; n (non-

immunogenic data) =425, Z-test).

h, Pie charts represent the percentage of different anchor mutation classes. N to P class (n=270 and not

N to P class (n=425).

Figure 5. HLA-C*05:01 is a potential allele to present KRAS G12D 10mer peptide.

a, Heatmap illustrating the relative sequence similarity of HLA-C alleles. HLA-C*08:02 acts as the

reference allele.

b, Top (left panel) and side (right panel) view of HLA-C*08:02 molecule. Three residues (Arg97,

Tyr99 and Arg156) side chains from HLA-C*08:02 molecule contact to peptide P3 were shown as

spheres.

c, Top (left panel) and side (right panel) view of HLA-C*08:02 molecule. Two residues (Ser77 and

Asn80) side chains contact to peptide PΩ, which different from HLA-C*05:01 molecule were shown as

spheres.

d, Unambiguous 2Fo-Fc electron density maps of KRAS G12D 10mer peptide (GA_D_GVGKSAL) from

solved C05-mut10m complex structures.

e, Overlay of the peptide region of C05-mut10m (cyan) and C08-mut10m (yellow) complexes.

f, Overlay of C05-mut10m (cyan) and C08-mut10m (yellow) complexes.

g, Similar peptide binding pattern at P1-P3 region of C05-mut10m and C08-mut10m complexes.

Similar bonds between these two complexes were shown in red dot lines.

h, Comparation of C05-mut10m and C08-mut10m complexes at peptides P10L. The upper panel shows

P10L from C08-mut10m. The peptide and bonds show in yellow. The lower panel shows P10L C05-

1036   mut10m. The peptide and bonds show in cyan.

1037

Figure 6. Optimized procedure of neoepitope (neoantigen and corresponding HLA) and new

HLA candidate discovery.

a, Workflow of neoepitope discovery. Conventional procedures were indicated in solid box.

Propositional procedures in our study were indicated in dotted box.

b, Interpretation of neoantigen candidate filtration procedures based on ours studies. Different classes

of neoantigen candidates filtrate with different strategies during neoantigen candidate filtration.

1044

Supplementary Figure 1. Structure of H-2 Kb in complex with DPAGT V213L wild type peptide

wt8mV and mutant mut8mL.

a, Unambiguous 2Fo-Fc electron density maps of DPAGT1 wild type 8mer peptide (wt8mV peptide

SIIVFNLV, magenta) and mutant 8mer peptide (mut8mL peptide SIIVFNL<u>L</u>, orange) from solved

structures.

b, Overlay of the DPAGT1 wt8mV (magenta) and mutant mut8mL (orange) peptides.

c, Sequence logo based on the data from IEDB showing amino acid preferences for 8mer peptides

bound to H-2 Kb. The peptide library was obtained from IEDB (n=4141).

d, Anchor residue P8V of wt8mV peptide interacts with H-2 Kb (grey).

e, Anchor residue P8L of mut8mLpeptide interacts with H-2 Kb (grey).

1055

1056 **Supplementary Figure 2. Comparison of the mutation distribution of 10mer peptides from IND**

1057 **and NND.**

1058 Comparation of the mutation occurrence frequency at each position of 10mer peptides from IND (red

1059 line) and NND (dark blue line). The grey dotted line represents the mutation theoretical frequency

1060 (1/10) of 10mer peptide. Positions of 10mer peptide show on horizontal axis.

1061

1062 **Supplementary Figure 3. Amino acids frequency of mutant peptides from IND and NND.**

1063 The horizontal axis depicts the amino acid codes of mutations. The vertical axis indicates the frequency

1064 of mutations (***$p<0.001$, n (immunogenic data) =129; n (non-immunogenic data) =11739, Z-test).

1065

1066 **Supplementary Figure 4. 9mer peptides binding profile of 29 HLA alleles.**

1067 Sequence logos showing amino acid binding profile for 9mer peptides bound to 29 HLA alleles. The

1068 peptide libraries were obtained from IEDB. The sample size of each allele is shown on the figure.

1069 Sequence logos were generated from peptide-binding matrices using the Seq2Logo server.

1070

1071 **Supplementary Figure 5. Phylogenetic analysis of 15 HLA-C alleles.**

1072 15 HLA-C sequences were aligned and then used to make phylogenetic relationship. Alleles in the red

1073 box represent the relatively far genetic relationship from HLA-C*08:02.

1074

1075 **Supplementary Figure 6. Multiple sequence alignment of HLA-C*08:02 and HLA-C*05:01**

1076 Sequence Alignment was done using Clustal Omega. Position 77 (Ser to Asn) and position 80 (Asn to

1077 Lys) are shown without underpainting.

1078

Supplementary Figure 7. Different peptide-MHC models with their potential immunogenicity.

Model A demonstrates the peptides which have mutations at TCR contact positions and presented by

MHC. The blue circle represents wild type residues. The red triangle represents the mutant residues

with far similarities with the wild type. Model B demonstrates the peptides which have mutations at

MHC contact position and presented by MHC. The red triangle represents the mutant residues with far

similarities with the wild type. Model C demonstrates the peptides which have mutations at anchor

position and are presented by MHC. The yellow rectangle represents wild type residues which are not

preferential in anchor position for related MHC alleles and thus hardly to bind and present by HLA.

The purple circle represents mutated residues which are preferential selected by related MHC alleles at

anchor position.

Figure 1

Figure 2

a

b

c

Figure 3

a

b

c

d

Figure 4

Figure 5

Figure 6

a

b

	Candidates classification	Candidates filtration methods	Candidates eliminated
Anchor mutated neopeptide	●●●	Preference selection	●● *Decrease ~42%
MHC contact position mutated neopeptide	●●	Importance selection Preference selection Morphologic selection	●
TCR contact position mutated neopeptide	●●●●●	Dissimilarity selection Morphologic selection	●●●●