Phenylacetyl-CoA, not phenylacetic acid, mediates quorum sensing-regulated virulence in

Burkholderia cenocepacia

Tasia Joy Lightly[a], Kara L. Frejuk[a], Marie-Christine Groleau[c], Laurent R. Chiarelli[d], Cor Ras[e],

Silvia Buroni[d], Eric Déziel[c], John L. Sorensen[f] and Silvia T. Cardona[a,b#]

[a]Department of Microbiology, University of Manitoba, Winnipeg, Canada

[b]Department of Medical Microbiology & Infectious Diseases, University of Manitoba,

Winnipeg, Canada

[c]Armand-Frappier Santé Biotechnologie Center, Institut National de la Recherche Scientifique

(INRS), Laval, Canada

[d]Department of Biology and Biotechnology "Lazzaro Spallanzani", University of Pavia, Pavia,

Italy

[e]Department of Biotechnology, Delft University of Technology, Delft, The Netherlands

[f]Department of Chemistry, University of Manitoba, Winnipeg, Canada

Running Head: Phenylacetyl-CoA inhibits LuxIR-type quorum sensing

#Address correspondence to Silvia T. Cardona, Silvia.Cardona@umanitoba.ca

Abstract word count: 244

Importance word count: 150

Text word count: 6,373

Abstract

In many bacteria, phenylalanine and other aromatic compounds are catabolized through phenylacetic acid (PAA) degradation. In this pathway, PAA is converted to phenylacetyl-CoA (PAA-CoA) by a ligase, PaaK, and then epoxidized by a multicomponent monooxygenase, PaaABCDE, before further degradation to the TCA cycle. As *Burkholderia cenocepacia paaABCDE* mutants release PAA and are attenuated for quorum sensing-regulated virulence in rat and nematode models of infection, overproduction of PAA in *B. cenocepacia* was linked to the regulation of the LuxIR-like quorum sensing system, CepIR. To further investigate this link, we created knockout mutants of the first step of the pathway (PAA-CoA synthesis by PaaK) and characterized them in comparison to wild type and PaaABCDE mutant strains using liquid chromatography/mass spectrometry (LC-MS/MS) and phenotypic assays. Both mutants released high levels of PAA. However, only the PaaABCDE mutant accumulated PAA-CoA intracellularly, had decreased virulence in *Caenorhabditis elegans,* and was defective for exoprotease activity. Conversely, the PaaK mutant did not accumulate PAA-CoA and was virulent, linking PAA-CoA to the attenuation of virulence in *B. cenocepacia*. While transcriptional analysis showed no evidence of *cepIR* downregulation by PAA-CoA, the attenuated pathogenicity of a *cepR* mutant was dependent on the presence of PAA-CoA in PAA pathway mutants, as it drastically changed to a virulent phenotype upon deletion of *paaK* but not when *paaABCDE* was deleted. Our results demonstrate the role of PAA-CoA in quorum sensing-regulated virulence and are consistent with a virulence regulatory mechanism that is evident when CepR is not active.

Importance

The opportunistic pathogen *Burkholderia cenocepacia* uses a chemical signal process called

quorum sensing to produce virulence factors. In *B. cenocepacia*, quorum sensing relies on the

presence of the transcriptional regulator CepR, which upon binding quorum sensing signal

molecules, activates virulence. In this work, we have found that even in the absence of CepR, *B.*

cenocepacia can elicit a pathogenic response if phenylacetyl-CoA, an intermediate of the

phenylacetic acid (PAA) degradation pathway, is not produced. Instead, accumulation of

phenylacetyl-CoA attenuates pathogenicity. Therefore, we have discovered that triggering of

virulence is possible in the absence of quorum sensing, challenging the classical view of

activation of virulence by quorum sensing. Our work provides new insight into the relationship

between metabolism and virulence in opportunistic bacteria. We propose that, in the event that

quorum sensing signaling molecules cannot accumulate to trigger a pathogenic response, a

metabolic signal can still activate virulence in *B. cenocepacia*.

Introduction

Intercellular communication in bacteria, or quorum sensing (QS), relies on the production

and detection of signalling molecules to regulate gene expression in a cell-density dependent

manner (1, 2). In many Gram-negative bacteria, LuxI homologs synthesize *N*-acyl homoserine

lactones (AHLs) as signalling molecules. These small diffusible molecules accumulate

extracellularly as the population increases until the concentration is sufficient for AHLs to return

into the cell where they bind LuxR-type transcriptional regulators. Complexing of the AHL to

the LuxR homolog results in a conformational change that alters the ability of the transcriptional

regulator to bind DNA and regulate gene expression (3). Many of the genes regulated by QS

control expression of virulence traits, which are important for bacteria to establish infection (4).

A bacterium with a LuxIR-type QS system is *Burkholderia cenocepacia*, a member of the

Burkholderia cepacia complex (5). The strain *B. cenocepacia* K56-2, isolated from the sputum

of a cystic fibrosis patient, has two complete LuxIR-type QS systems, CepIR and CciIR, and one

orphan transcriptional regulator, CepR2 (5, 6). The regulons of CepR and CciR have been

examined and a large number of the genes that are regulated are linked to virulence (7). While

CepR and CciR have separate regulons, certain genes are regulated by both in a reciprocal

manner (7). CepR positively regulates gene expression, whereas CciR is responsible for negative

gene regulation (7). While these two systems regulate certain genes in a reciprocal manner,

rather than hierarchically like those of *Pseudomonas aeruginosa,* the transcription of both *cepI*

and *cciIR* is dependent on CepR and in turn CciR negatively regulates the expression of *cepI* (7,

8). CepR and CciR both positively regulate their canonical autoinducer synthase and negatively

regulate their own transcription (8, 9). CepI mainly produces *N*-octanoyl homoserine lactone

(C8-HSL) with slight production of *N*-hexanoyl homoserine lactone (C6-HSL), whereas CciI

synthesizes C6-HSL with slight secondary production of C8-HSL (8, 10). CepR2 lacks a cognate

AHL synthase and has been shown to regulate known QS-regulated genes independent of a

signalling molecule (11).

Several lines of evidence point to metabolic adaptation as an important component of

bacterial virulence regulation (12–15) and the relationship between phenylacetic acid (PAA)

degradation and virulence in *B. cenocepacia* is the focus of this study. *B. cenocepacia* can use

the PAA degradation pathway to catabolize diverse aromatic compounds, such as phenylalanine,

to components of the three carboxylic acid (TCA) cycle (16). Based on the PAA degradation

pathway elucidated in *Escherichia coli* K12*,* PAA is converted to phenylacetyl-Coenzyme A

(PAA-CoA) by the action of a phenylacetyl-CoA ligase*,* PaaK (16). The conversion to PAA-

CoA by PaaK can be reverted to PAA by the action of PaaI, a thioesterase, to prevent

accumulation should downstream steps be disrupted (17). Next, PAA-CoA is epoxidized by the

multicomponent monooxygenase, PaaABCDE and then further degraded into succinyl-CoA and

acetyl-CoA (16).

We previously found that CepIR-regulated virulence traits and *cepI* and *cepR* promoter

activity were downregulated in a *B. cenocepacia* mutant of the *paaABCDE* operon that released

PAA (18). We also demonstrated that PAA could be produced to detectable levels in wild type *B.*

cenocepacia K56-2 under certain conditions, suggesting a potential physiological role of the

regulation of virulence by this molecule (19). However, the PaaABCDE mutant is still able to

degrade PAA to PAA-CoA. Therefore, the metabolite responsible and the mechanism of the

regulation of virulence by accumulated PAA-metabolites is not known. In this work, we further

characterized mutants of the PAA degradation pathway to show that the accumulation of PAA-

CoA, but not PAA, is responsible for the attenuation phenotype. While we found no evidence of

a direct effect of PAA-CoA on CepI activity or the formation of CepR:C8-HSL complexes, our

results are consistent with a model in which a parallel mechanism that uses PAA-CoA as a

regulatory ligand, competes with the CepIR complex to regulate virulence. We propose that

PAA-CoA acts as a co-repressor that binds to a yet unidentified transcriptional regulator,

competitively inhibiting the CepIR QS regulation.

Results

Creation and phenotypic characterization of a paaK *double mutant*

Previously, we demonstrated a link between the attenuation of *B. cenocepacia* K56-

2Δ*paaABCDE* and the accumulation of PAA (18). However, the interruption of the pathway at

this step could also result in an accumulation of PAA-CoA because PAA is converted to PAA-

CoA by a PaaK ligase (Fig. 1). To determine the role of each of these molecules in the

attenuation of virulence, we characterized a *paaK* mutant that should only produce PAA and

compared it to the *paaABCDE* mutant. In K56-2 there are two similar but not identical

phenylacetyl-CoA ligases, PaaK1 and PaaK2 (20). We created marker-less deletion mutants of

BCAL0404 (*paaK1*) and BCAM1711 (*paaK2*). Knockout mutants of the other *paaK* were then

made in each background resulting in two independently made double knockouts, which we have

named K56-2Δ*paaK1*Δ*paaK2* and K56-2Δ*paaK2*Δ*paaK1*.

Fig. 1: The proposed phenylacetic acid degradation (PAA) pathway in *B. cenocepacia.*

Based on the pathway described in *E. coli* and *Pseudomonas* by Teufel *et al.* (2010),

phenylalanine (Phe) is degraded to PAA and the first step of the pathway is the conversion of

PAA to phenylacetyl-CoA (PAA-CoA) by the phenylacetyl-CoA ligase, **PaaK** (shown in

purple). The phenylacetyl-CoA thioesterase, **PaaI,** can reversibly convert PAA-CoA back to

PAA to prevent toxic accumulation of the epoxide if the downstream pathway is blocked. PAA-

CoA is epoxidized by the ring 1,2 phenylacetyl-CoA epoxidase, **PaaABCDE** (shown in orange),

forming ring-1,2-epoxyphenylacetyl-CoA or epoxide (EP-CoA). The epoxide is further degraded

to components of the TCA cycle, acetyl-CoA and succinyl-CoA.

To confirm the interruption of the PAA pathway, wild type K56-2 and mutants were

grown on LB media and M9 minimal medium with glucose or PAA as a sole carbon source. No

growth defect was observed in LB or glucose (Fig. 2 A and B). Both *paaK* single knockout

mutants were also able to grow on PAA as a sole carbon source to wild type levels, indicating

that PAA can be used as a substrate for both ligases (Fig. 2B). While mutants were able to grow

normally on glucose as a sole carbon source, the *paaK* double knockout mutants (K56-

2Δ*paaK1*Δ*paaK2* and K56-2Δ*paaK2*Δ*paaK1*) were unable to grow on PAA (Fig. 2B). Growth

was complemented with a plasmid containing a rhamnose-inducible *paaK2* (BCAM1711) (Fig.

S2) confirming that the interruption of the pathway in these mutants was caused by the deletion

of *paaK*.

The PaaABCDE mutant has been shown to accumulate extracellular PAA due to the

interruption of the pathway (18) but the concentration of intracellular PAA and PAA-CoA was

unknown. Intracellular and extracellular concentrations of both PAA and PAA-CoA were

measured with ion pair reversed phase ultra-high performance liquid chromatography tandem

mass spectrometry (IP-RP-UHPLC-MS/MS) using the differential method (21–23). As expected,

mutants with complete interruption of the pathway (Δ*paaABCDE*, Δ*paaK1*Δ*paak2,* and

Δ*paaK2*Δ*paak1*) accumulated more PAA than wild type (Fig. 2C). PAA was released

extracellularly (Fig. 2C, Fig. S3A), whereas PAA-CoA accumulated intracellularly (Fig. 2D, Fig.

S3B). The only mutant strain that accumulated PAA-CoA was Δ*paaABCDE* (Fig. 2D).

Fig. 2: Characterization of the growth and metabolite production of *B. cenocepacia* PAA

pathway mutants. A) *B. cenocepacia* K56-2 PAA pathway mutants exhibited no growth defects

when grown on LB media. **B)** *B. cenocepacia* K56-2 PAA pathway mutants were grown on M9

with 25 mM Glucose as the sole carbon source or M9 with 5 mM PAA as the sole carbon source

for 48 hours and the OD_{600} was measured. Mutants with full interruption of the pathway

(*ΔpaaABCDE, ΔpaaK1ΔpaaK2,* and *ΔpaaK2ΔpaaK1*) were unable to grow on PAA as a sole

carbon source. Single deletion mutants of the *paaK* ligases (*ΔpaaK1* and *ΔpaaK2*) grew to wild

type levels on PAA indicating that the pathway is not interrupted in these strains. Error bars

represent SD for three biological replicates. b represents p values ≤ 0.05 as determined by

student's T-test (two tailed) when compared to growth on PAA of K56-2. **C)** IP-RP-UHPLC-

MS/MS revealed that PAA is released extracellularly in the *paaABCDE* and *paaK* knockout

mutants. PAA did not accumulate in the wild type and *paaK* single knockout strains where the

PAA pathway is still functional. **D)** PAA-CoA was present at detectable limits in all of the

strains, but PAA-CoA accumulates solely in the *paaABCDE* operon knockout mutant. For all IP-

RP-UHPLC-MS/MS two biological replicates with three technical replicates each were

performed. An Asterix denotes $p \leq 0.05$ as determined by student's T-test (two tailed) when

compared to K56-2.

Fig. 3: The Δ*paaK1*Δ*paaK2* mutant has wild type levels of virulence in *C. elegans* and

proteolytic activity whereas the Δ*paaABCDE* mutant is attenuated. A) Slow killing assays

with *C. elegans* over a period of seven days show that the Δ*paaK1*Δ*paaK2* mutant kills *C.*

elegans in a similar time frame as wild type, whereas the nematodes exposed to the Δ*paaABCDE*

mutant have increased survival. The increased killing ability of the Δ*paaK1*Δ*paaK2* mutant

compared to wild type was determined as significant according to the log-rank test. Three

biological replicates were performed. **B)** Proteolytic activity was measured as the area of the

zone of clearance (excluding colonies) on agar containing 2% skim milk with or without the

addition of 2 mM of phenylalanine. The error bars represent the SD of three independent

experiments.

PAA-CoA has no effect on CepR and CepI function in vitro

The above findings suggest that a mutant that releases PAA and accumulates PAA-CoA

is attenuated for virulence in *C. elegans* and has decreased proteolytic activity whereas a mutant

that only releases PAA is not attenuated for these QS-regulated virulence traits. Since the

accumulation of PAA-CoA is the major difference between these strains we hypothesized that

PAA-CoA or a derivative was responsible for attenuating these traits. Because the CepIR QS

system is required for full virulence of *B. cenocepacia* K56-2 in *C. elegans*, while the CciIR QS

system is not (25), we focused on the effect of PAA-CoA on the CepIR QS system. In addition,

Δ*paaABCDE* had decreased *cepI* and *cepR* promoter activity and AHL signaling (18).

Specifically, we asked whether PAA-CoA could: 1) inhibit the formation of CepR:AHL

complexes; 2) inhibit the enzymatic activity of CepI; and/or 3) indirectly affect an upstream

regulator.

To address the first question, we built a reporter system similar to those commonly used

to identify quorum sensing inhibitors (26). *E. coli* W14 was selected as our reporter strain as it

lacks the PAA pathway genes (27, 28) ensuring the exogenously added PAA-related compounds

would not be degraded. Using arabinose-inducible expression of *cepR* (BCAM1868) we created

a reporter strain that responds to CepR:AHL complexes by activating the *cepI* promoter which

controls the expression of luminescence genes *luxCDABE* (Fig. 4A). Expression of luminescence

is decreased in the presence of an inhibitor of the formation of CepR:AHL complexes. However,

PAA-CoA cannot be added exogenously to cultures because it is too large to cross the

membrane. Therefore, we created a plasmid that contains the *E. coli* W *paaK* ligase gene

(ECW_RS07845) under the control of a rhamnose-inducible promoter. In the presence of

rhamnose, PaaK converts exogenously added PAA to PAA-CoA, as confirmed by IP-RP-

UHPLC-MS/MS (Fig. 4B). To ensure that any effects we saw were not due to an inhibition of

the arabinose-inducible promoter or luminescence itself we created two control strains that

contain either the arabinose-inducible P_{BAD} promoter or a constitutive promoter, *dhfr*, controlling

the *luxCDABE* genes.

Using this reporter system (W14/pTL7/pTL5/pTL22 referred to from here on as CepR

reporter strain, Fig. 4A) we observed that although there was inhibition of luminescence in the

presence of PAA-CoA, the level of inhibition was similar to that of the control strains (Fig. 4C).

Similarly, PAA itself also had no effect on the system (Fig. 4D). In summary, these results show

that PAA-CoA and PAA do not directly affect CepI activity or CepR:C8-HSL formation.

Fig. 4: Neither PAA nor PAA-CoA affect the formation of CepR:C8-HSL complexes. A)

The CepR reporter strain and control strains used. Regulatory genes *araC* (for P_{BAD}) and *rhaSR*

(for P_{rhaBAD}) were excluded from the figure for simplicity. **B)** In the presence of 0.01% rhamnose

(rha), 0.5 mM of PAA was converted to PAA-CoA by the PaaK ligase in the CepR reporter

strain. **C)** PAA-CoA had no effect on the CepR reporter strain compared to that of the effect on

control strains. **D)** PAA itself was tested at various concentrations from 0.1-1mM (0.1 mM

shown here) and no effect was observed when compared to the control strains. The error bars

represent the SD of three independent experiments.

To analyze if PAA or PAA-CoA inhibits the enzymatic activity of CepI, we used

recombinant CepI in the presence of dichlorophenolindophenol (DCPIP) to

spectrophotometrically measure the activity of CepI (29). CepI catalyzes the formation of C8-

HSL from octanoyl-Acyl Carrier Protein (C8-ACP) and *S*-adenosyl methionine (SAM). During

this reaction DCPIP is reduced resulting in a colour change that can be used to measure activity

of CepI. Enzyme activity showed no change in the presence of 200 µM of PAA-CoA or PAA

(Table 1), suggesting that neither molecule directly inhibits the activity of CepI.

Table 1: CepI activity is not inhibited by the presence of PAA or PAA-CoA.

Conditions	Enzyme activity $\pm$ SD (min^{-1})
C8-ACP + SAM	3.46 ± 0.12
C8-ACP + SAM + 200 µM PAA	3.29 ± 0.21
C8-ACP + SAM + 200 µM PAA-CoA	3.04 ± 0.29

As our data did not show a direct effect of PAA-CoA on the CepIR QS system but *cepI*

and *cepR* promoter activity was decreased in the PaaABCDE mutant (18), we hypothesized that

PAA-CoA may indirectly affect an upstream regulator of the CepIR QS system. If this was the

case, we predicted that the differences in pathogenicity between the Δ*paaABCDE* and the

Δ*paaK1*Δ*paaK2* mutant strains would be explained by differential expression of *cepI* and *cepR*.

Using stationary phase cultures, we measured the transcript levels of *cepI* and *cepR* and the

levels of C8-HSL in the PAA pathway mutants and compared it to wild type. Surprisingly, the

Δ*paaABCDE* and the Δ*paaK1*Δ*paaK2* mutants had wild type levels of *cepI* and *cepR*

transcription (Fig. 5A and Fig. S5A) despite the striking differences in their virulence

phenotypes and contradicting our previous results where reporter systems measured decreased

cepI and *cepR* promoter activity and decreased AHL signaling (18). This suggested that the

attenuation of the Δ*paaABCDE* mutant is independent of the transcription of the CepIR QS

system. To confirm the wild type transcription levels of our mutants, we measured the levels of

AHL molecules in each strain. Similar to the transcription, the levels of C8-HSL, taken from

stationary phase samples did not correlate with the decreased virulence of our PAA pathway

mutants (Fig. 5B and Fig. S5B). However, both Δ*cepI* and Δ*cepR* had no measurable C8-HSL

which is consistent with their lack of virulence (Fig. 5B).

Fig. 5: Transcription of *cepI* and *cepR* and C8-HSL production is not decreased in any of

the PAA pathway mutants. A) RT-qPCR shows that *cepI* and *cepR* gene expression is not

decreased compared to wild type in the PAA pathway mutants. **B)** C8-HSL levels were not

decreased in any of the mutants, but C8-HSL levels were higher in the Δ*paaABCDE* mutant.

Virulence of strains in both *C. elegans* assays and exoprotease activity assays is denoted above

the graphs per the results from Fig. 3.

The CepR-independent virulence of PaaK knockout mutants is mediated by the presence/absence

of PAA-CoA

Transcription of *cepI* and *cepR* is not decreased in PAA pathway mutants and C8-HSL is

present in the mutants yet the virulence factors attenuated in the Δ*paaABCDE* mutant

(pathogenicity in *C. elegans* and exoprotease activity) are known to be under the control of the

CepIR system. To further understand how the PAA pathway may regulate QS related virulence

without affecting *cepIR* transcription, we created marker-less deletions of *cepI* and *cepR* in the

PAA pathway deletion mutant backgrounds. We reasoned that if PAA-related regulation of

virulence was completely independent of CepI and CepR activity, then deletion of *cepI* and *cepR*

in a Δ*paaABCDE* background should show an additive attenuation effect. Similarly, deletion of

cepI and *cepR* in a Δ*paaK1*Δ*paaK2* mutant background should show the same degree of

attenuation as the *cepI* and *cepR* deletions in a wild type background. Once we had created

knockout mutants of *cepI* and *cepR* for each PAA pathway mutant in parallel with *cepI* and *cepR*

deletions in wild type K56-2, we analyzed their growth and found no defect when grown in LB

media over 24 hours (Fig. S6). Next, we used virulence assays to determine if the CepIR QS

system was needed for the regulation of their QS. The virulence of the *cepI* and *cepR* knockout

mutants in wild type and PAA mutant backgrounds was compared in slow killing assays (Fig. 6A

and B, Fig. S7A and B). Both *cep* QS mutants in the Δ*paaABCDE* background were more

attenuated for virulence than Δ*paaABCDE* (Fig. S7A). However, they failed to show an additive

effect (Fig. S7A). Moreover, while the Δ*paaK1*Δ*paaK2*Δ*cepI* mutant was as attenuated as Δ*cepI*,

ΔpaaK1ΔpaaK2ΔcepR was not, showing that the pathogenic phenotype was independent of

CepR (Fig. 6A and B). This was mirrored in the *ΔpaaK2ΔpaaK1ΔcepR* mutant (Fig. S7B). This

is surprising considering the *cepR* deletion in wild type and the *paaABCDE* mutant has a severe

attenuation (Fig. S7A).

Corroborating the results of the *C. elegans* assay, *ΔpaaK1ΔpaaK2ΔcepR* had wild type

levels of proteolytic activity whereas the *ΔpaaK1ΔpaaK2ΔcepI* mutant had none (Fig. 6C and

Fig. S7C). The only difference between the *ΔpaaK1ΔpaaK2ΔcepR* mutant and the *cepR* mutant

is the presence of PAA. If PAA was responsible for the elimination of the *cepR* attenuation, the

ΔpaaABCDEΔcepR mutant should be pathogenic but our results show it is not. Therefore, we

conclude that it is the lack of PAA-CoA which causes the increase in pathogenicity in

ΔpaaK1ΔpaaK2ΔcepR, confirming the link between PAA-CoA and CepR mediated QS.

To further understand the virulence of the *ΔpaaK1ΔpaaK2ΔcepR* mutant, we examined

the expression levels of *cepI* and *cepR* and the levels of C8-HSL in the mutants. While

transcription of *cepR* is abolished in *ΔcepI*, it is restored to wild type levels when the deletion of

cepI occurs in a *ΔpaaK1ΔpaaK2* or *ΔpaaABCDE* background (Fig. 6D). Therefore PAA, which

is present in both PAA pathway mutants, restores *cepR* transcription but not pathogenicity,

because there is no CepI and no C8-HSL (Fig. 6F). When *cepI* transcript levels were analyzed in

ΔcepR in wild type and PAA degradation mutant backgrounds, the transcription of *cepI* was

downregulated in the CepR mutant, as expected. Similarly, *cepI* transcription was reduced in the

attenuated *ΔpaaABCDEΔcepR* and the pathogenic *ΔpaaK1ΔpaaK2ΔcepR* (Fig. 6E). The lack of

C8-HSL in both the *ΔpaaK1ΔpaaK2ΔcepI* and *ΔpaaK1ΔpaaK2ΔcepR* (Fig. 6F) mutants

suggests the pathogenicity of the *ΔpaaK1ΔpaaK2ΔcepR* mutant is independent of *cepI*. This was

mirrored in the *ΔpaaK2ΔpaaK1ΔcepR* mutant (Fig. S8).

Fig. 6: The virulence of Δ*paaK1*Δ*paaK2*Δ*cepR* is independent of the CepIR QS system. A)

The *cepI* and *cepR* mutations in a Δ*paaK1*Δ*paaK2* mutant have opposing levels of virulence.

Δ*paaK1*Δ*paaK2*Δ*cepR* is as virulent as wild type whereas Δ*paaK1*Δ*paaK2*Δ*cepI* is attenuated for

virulence in *C. elegans*. **B)** *cepI* and *cepR* mutants have similar levels of attenuation in *C.*

elegans. Figures **A)** and **B)** are representative of three biological replicates. **C)** The

Δ*paaK1*Δ*paaK2*Δ*cepR* mutant has wild type levels of proteolytic activity with or without

phenylalanine whereas the Δ*cepR* mutant has decreased proteolytic activity. Proteolytic activity

was measured as the area of the zone of clearance (excluding colonies) on agar containing 2%

skim milk with or without the addition of 2 mM of phenylalanine. **D)** *cepR* transcription is

restored to wild type levels in Δ*paaABCDE* and Δ*paaK1*Δ*paaK2*Δ*cepI* indicating that PAA

restores *cepR* transcription. **E)** *cepI* is downregulated in all of the Δ*cepR* mutants. **F)** C8-HSL

levels were measured and none of the QS mutants had detectable levels of C8-HSL. The error

bars represent the SD of three independent experiments.

Deletions of the *cepI* and *cepR* genes in the PAA pathway mutant backgrounds allowed

us to determine that the PAA pathway has an effect on CepIR-regulated virulence that is evident

when CepI or CepR is absent. In this context, our results suggest that the absence of PAA-CoA,

or a derivative of PAA-CoA, permits an alternative signalling pathway to activate virulence in

the Δ*paaK1*Δ*paaK2*Δ*cepR* mutant. The opposite effects of the PaaK and PaaABCDE mutants

with regards to pathogenicity and the striking differences that *paaK* and *paaABCDE* gene

deletions have in the *cepR* mutant background suggests that PAA-CoA, or a derivative, is the

central molecule implicated in this alternative signalling pathway.

Discussion

In *B. cenocepacia*, the PAA pathway has been linked to the regulation of virulence. The

attenuation of CepIR QS-regulated virulence traits in PAA pathway mutants was thought to be

due to the accumulation of PAA in these mutants (18). PAA has also been shown to regulate the

virulence of other Gram-negative bacteria, *Acinetobacter baumannii* (30) and *Pseudomonas*

aeruginosa (31, 32), two Gram-negative bacteria that also use LuxIR-type QS systems to

regulate their virulence (33, 34). Previously, we demonstrated that the release of PAA in a *B.*

cenocepacia mutant of the PAA pathway resulted in the inhibition of CepIR-regulated QS, a

LuxIR-type system (18). However, the Δ*paaABCDE* mutant has the genes necessary to convert

PAA to PAA-CoA (Fig. 1). Therefore, PAA or PAA-CoA could be the molecule responsible for

the inhibition of CepIR-regulated QS.

To determine the molecule responsible, we initially created a mutant unable to

accumulate PAA-CoA. Unlike the Δ*paaABCDE* mutant, the PaaK knockout mutant was as

virulent as wild type, suggesting that the attenuated virulence was due to the accumulation of

PAA-CoA, or a derivative. Another difference between the PaaK and the PaaABCDE mutant is

that the PaaK knockout mutant released less PAA than the PaaABCDE mutant. However, the

intracellular concentrations of PAA were similar and detected at levels very close to, or lower

than, the smallest standard used for the calibration curve (8 µM). We believe that differences in

intracellular PAA levels should have been found if PAA affected the regulation of virulence.

We hypothesized CepIR-regulated QS inhibition was achieved by one of three

mechanisms: 1) direct inhibition of the enzymatic activity of CepI or its products; 2) inhibition of

the formation of CepR:C8-HSL complex; or 3) an indirect effect on an upstream regulator. Using

an *E. coli* reporter system that can't metabolize PAA, we examined the ability of PAA-CoA to

act as a direct inhibitor of the CepIR QS system. We were able to observe strong but unspecific

inhibition of luminescence by PAA and PAA-CoA, highlighting the importance of control strains

when reporter systems are used to identify QS inhibitors: QS-regulated phenotypes often depend

on other factors; and reporter strains using bioluminescence can be affected by the overall

metabolic activity of the cell (26). In our system, the control strains had either the arabinose-

inducible P$_{BAD}$ promoter or the *dhfr* constitutive promoter controlling the expression of

luxCDABE luminescence genes. First, in the presence of higher concentrations of PAA (above 1

mM) the expression of luminescence by the constitutive promoter was inhibited suggesting a

general effect on luminescence. Second, the concentration of rhamnose used to express *paaK,*

0.01% w/v, inhibited the arabinose-inducible promoter. With these control strains we were able

to minimize the general effects on the reporter system and determine that neither PAA or PAA-

CoA have a direct effect on the formation of CepR:C8-HSL complexes or on CepI activity.

Interestingly, we found that the attenuation of virulence in Δ*paaABCDE* did not

correspond to decreased *cepI* and *cepR* transcription or C8-HSL production. In fact, the

concentration of C8-HSLs present in the strains did not correlate to the pathogenicity of the PAA

pathway mutants. While one of the pathogenic *paaK* knockout mutants had wild type levels of

C8-HSL, the independently made mutant had a similar concentration to that of the Δ*paaABCDE*

mutant despite their opposing pathogenicity phenotypes. Based on the current knowledge of the

CepIR QS system, transcription of *cepI* relies on CepR and CepR is negatively autoregulated (9).

This is in agreement with our findings that the transcription of *cepI* is dependent on CepR and

that *cepR* is downregulated in the absence of *cepI*. There was no detectable C8-HSL in either of

the *cepI* and *cepR* deletion strains supporting the interdependence of these two genes. Therefore

the lack of downregulation of either *cepI* or *cepR* in the Δ*paaABCDE* mutant was quite

surprising because this contradicted the results from our plasmid-based reporter systems that

showed decreased *cepI* and *cepR* promoter activity and decreased AHL signalling (18). To

investigate this discrepancy in results between the different methods, we introduced the plasmids

with P$_{cepI}$::*luxCDABE* or P$_{cepR}$::*luxCDABE* to the Δ*paaK1*Δ*paaK2* and Δ*paaK2*Δ*paaK1* mutants

and found decreased *cepI* and *cepR* promoter activity almost identical to that of the Δ*paaABCDE*

mutant with the same plasmid based reporter system (data not shown). This suggests that the

PAA present in both these mutants might have a general effect on the reporter system itself. We

thought that PAA may affect the copy number of the plasmid or affect the metabolism of the

cells in such a way that affects plasmid-based reporter strains. Previously PAA has been shown

to affect the copy number and increase the frequency of plasmid loss from an *E. coli* strain (35).

The concentration of PAA used in that assay was higher than what we measured with IP-RP-

UHPLC-MS/MS for these strains, but the effect of PAA on plasmid copy number in *B.*

cenocepacia has not been studied. These reporter strains also use bioluminescence, which

requires large amounts of energy, so any changes in metabolic activity could be reflected in the

reporter strain and misinterpreted without the use of proper controls (26).

To determine if the attenuated virulence of Δ*paaABCDE* could be independent of CepI

and CepR activity we created *cepI* and *cepR* mutants in the PAA pathway mutant backgrounds.

If PAA-related regulation of virulence was independent of CepI and CepR activity, then we

would see an additive effect on attenuation. Whereas, in a *paaK* knockout mutant background we

expected to see a virulence phenotype similar to that of the same deletion in a wild type strain. In

the Δ*paaABCDE* QS mutants we saw no evidence of an additive effect as the QS mutations were

dominant, leading to the same pathogenicity as a wild type Δ*cepI* or Δ*cepR* mutant. Because the

cepI/R mutations in Δ*paaABCDE* did not have an additive effect, this suggests the mechanism of

attenuation involves the CepIR QS system. In the *paaK* knockout background the QS mutations

had opposing effects. While a *cepI* deletion resulted in the expected attenuated virulence, a *cepR*

deletion had wild type levels of virulence. Transcription of *cepI* and *cepR* in these mutants

showed that PAA pathway mutants with a *cepI* deletion had increased transcription of *cepR*.

Therefore, the presence of PAA, present in both PAA pathway mutants, can restore the

transcription of *cepR* to wild type levels. Based on our findings, we propose that PAA-CoA

exerts its effect by forming a complex with a non-canonical transcriptional regulator XR that

competes with CepR:C8-HSL for binding in virulence gene regulatory elements (Fig. 7). In this

context, XR would increase virulence while XR:PAA-CoA would not. In wild type (Fig. 7, left

panel) *cepI* and *cepR* are transcribed and CepR complexes with C8-HSL and regulates virulence.

In conditions where the PAA pathway is functional PAA-CoA does not accumulate to levels that

would inhibit virulence. In the PaaABCDE mutant background, where PAA-CoA is abundant,

the XR:PAA-CoA complexes displace CepR:C8-HSL complexes. This competition results in

attenuated virulence in *C. elegans* (Fig. 7, right panel) and would be diminished by the addition

of C8-HSL as we determined previously (18). In the PaaABCDE QS mutants XR:PAA-CoA has

no CepR:AHL complexes to displace so the virulence is the same as a QS mutant in a wild type

background. CepR mutants have decreased *cepI* transcription so no CepR:C8-HSL complexes

are formed resulting in attenuated virulence (Fig. 7, bottom panel). This model would also

explain the CepR-independent virulence of Δ*paaK1*Δ*paaK2*Δ*cepR*. In the absence of both *cepR*

and PAA-CoA, XR is able to activate certain virulence genes (Fig. 7, bottom panel). We were

unable to detect differences between the levels of PAA-CoA in wild type and PaaK mutants as

the amounts of PAA-CoA were very close to the limit of detection of our method. However, it is

reasonable to assume that wild type can produce PAA-CoA while a PaaK mutant cannot.

Therefore, we propose that, in the absence of PAA-CoA, XR positively regulates the virulence

genes. This alternative regulatory system does not require C8-HSL to activate XR as C8-HSL

was absent in all of the PAA pathway QS mutants. We propose that this alternative pathway may

be relevant in environmental conditions where quorum sensing signals do not accumulate but a

virulent response is necessary.

Fig. 7: Model of the role of the PAA pathway in the regulation of CepIR quorum sensing-

controlled virulence. Our model suggests that PAA-CoA acts as a co-repressor, with an as yet

unidentified transcriptional regulator (XR). In wild type CepIR QS dominates the regulation of

virulence due to the lower concentration of PAA-CoA (top left panel). Whereas in Δ*paaABCDE*

the levels of PAA-CoA are high and the XR:PAA-CoA complex can compete with CepR:C8-

HSL complexes to repress virulence, resulting in an attenuated virulence compared to wild type

(top right panel). CepR mutants in wild type don't have CepR:C8-HSL complexes to regulate

virulence and are therefore attenuated for virulence (bottom panel). However, in a PaaK

background CepR mutants are virulent. The absence of PAA-CoA in these mutants in

combination with the lack of CepR create the perfect conditions for XR to positively regulate

virulence genes.

Normally Δ*cepI* mutants have downregulation of *cepR*, but as we observed in RT-qPCR

of the PAA pathway mutants with Δ*cepI* there was increased *cepR* transcription (Fig. 6A)

suggesting that cepR transcription can be increased even in the absence of AHL molecules.

However, both these mutants are attenuated in pathogenicity so the presence of *cepR* in these

mutants may prevent the action of the alternate transcriptional regulator that bypasses the CepIR

system.

When taken together, the results presented here suggest that *B. cenocepacia* is able to

modulate QS-controlled gene expression in the absence of the CepIR QS system through the

PAA catabolic pathway. *B. cenocepacia* might use precursors of PAA as carbon sources to

support growth in certain environmental conditions. The CepIR QS system has been the most

studied QS system in *B. cenocepacia* due to its role in pathogenicity and its position in the

hierarchy but our results suggest that the regulation of QS in *B. cenocepacia* is still not fully

understood and that an, as yet, unidentified transcriptional regulator may regulate the virulence

in the absence of the CepIR QS system. Efforts to identify this putative transcriptional regulator

are ongoing. While this work contributes to our knowledge of the role of the PAA pathway in the

regulation of QS-related virulence more work is needed to fully understand the complexity that is

PAA degradation during *B. cenocepacia* natural interactions with their diverse hosts.

Materials and Methods

Strains, media preparation and growth conditions

Bacterial strains, listed in Table S1, were grown at 37°C with agitation at 230 rpm in

Lysogeny Broth (LB) medium, unless specified. All cultures were started from isolated colonies

grown 48 hours on LB agar at 37°C, unless specified. Plasmids used are listed in Table S2. For

E. coli strains the following antibiotics and concentrations were used: Ampicillin (100 µg/ml;

200 µg/ml for W14 strain); Chloramphenicol (30µg/ml), Kanamycin (30 µg/ml; 40 µg/ml for

MM290 strain), Trimethoprim (50 µg/ml; 100 µg/ml for W14 strain), Tetracycline (20 µg/ml).

For *B. cenocepacia* strains the following antibiotics and concentrations were used: Trimethoprim

(100 µg/ml), Tetracycline (100 µg/ml), Gentamycin (50 µg/ml). Stock solutions of C8-HSL

(Sigma Aldrich), were suspended in dimethyl sulfoxide (DMSO) and stored at -20°C. For assays

with exogenous addition of C8-HSL cultures were grown in LB supplemented with 50 mM 3-(*N*-

morpholino)propanesulfonic acid (MOPS) to prevent pH-dependent lactonolysis (36). Stock

solutions of L-arabinose (Alfa Aesar, Fisher Scientific) and L-rhamnose (Sigma Aldrich) were

prepared at a concentration of 20% (w/v) in deionized water and filter sterilized. Stock solutions

of phenylalanine and PAA (Sigma Aldrich) were prepared at 100 mM in deionized water, filter

sterilized and stored at room temperature. Stock solutions of PAA-CoA (Sigma Aldrich) were

prepared at 3.5 mM in DMSO and stored at -80°C.

Molecular biology techniques

Genetic manipulation of *B. cenocepacia* K56-2 was performed via tri-parental matings

with *E. coli* MM290/pRK2013 as a helper strain. Plasmids generated were maintained in *E. coli*

SY327λpir or DH5α Z-competent cells depending on their origin of replication. PCRs were

performed with an Eppendorf Mastercycler ep gradient S thermocycler with either HotStar

HiFidelity *Taq* polymerase (Qiagen) or *Taq* DNA polymerase (Qiagen). Conditions were

optimized for each primer set; primers used are listed in Table S3. Restriction enzymes and T4

DNA ligase (New England Biolabs) were used as recommended by the manufacturer. PCR

products and digested DNA were purified by QIAquick Gel or PCR purification kits (Qiagen)

and plasmids were isolated using QIAprep Miniprep kit (Qiagen).

Construction of unmarked deletion mutants in B. cenocepacia

Unmarked deletion mutants were created using the system developed by Flannagan *et al*.

(37). Flanking regions of the target gene were synthesized as a custom gene fragment with *Xba*I

and *Sma*I restriction sites (Integrated DNA Technologies, Inc), see Table 4 for sequences. The

fragment with regions flanking BCAL0404 (*paaK1*) and pGPI-*Sce*I were digested and ligated

together to create pTL17 and pTL18 was created by insertion of the fragment with regions

flanking BCAM1711 (*paaK2*). Each plasmid was introduced into wild-type K56-2 by tri-parental

mating and were integrated into the genome by homologous recombination. Colonies were

picked onto LB medium supplemented with trimethoprim and onto *Burkholderia cepacia*

selective agar (BCSA). Conjugates with a dull phenotype on BCSA were selected. The

integration of the suicide plasmids was confirmed with primers 900/901for pTL17 and primers

902/903 for pTL18. Then, pDAI-*Sce*I, expressing I-*Sce*I endonuclease, was introduced into K56-

2::pTL17 and K56-2::pTL18 by tri-parental mating. The double-stranded break caused by the I-

*Sce*I endonuclease triggered a recombination event that resulted in a reversion of the conjugate to

a wild type allele or a deletion of the fragment between the two flanking regions. Potential

conjugates were tetracycline resistant and trimethoprim sensitive. The deletion event was

confirmed by PCR using primer set 980/981 for *paaK1* and 982/983 for *paaK2*. Positive clones

were cured of the pDAI-*Sce*I plasmid by sequential sub-culturing in LB medium. Loss of the

plasmid was confirmed by the loss of tetracycline resistance using the replica plate method and

PCR with primer set 920/921 for the absence of pDAI-*Sce*I.

Once cured, the mutants were tested for growth defects by washing overnight cultures

twice in PBS, diluting to an OD_{600} of 0.04 in 200 µl of LB in a clear 96-well plate in triplicate.

Plates were incubated at 37 °C in a Biotek Synergy II plate reader with continuous shaking for

24 hours with OD_{600} measured every hour. Deletions of *cepI* and *cepR* were performed in the

wild type and PAA pathway mutant backgrounds using the pDelCepI and pDelCepR plasmids

created by Aubert *et al.* (38). Integration of the plasmids was confirmed using primer set

1082/154 for *cepI* deletions and 1080/154 for *cepR*. pDAI-*Sce*I was introduced and potential

conjugates were selected on appropriate antibiotics and the deletion event was confirmed by

PCR with primer sets 1112/1113 and 1080/1081 for *cepI* and *cepR*, respectively. Positive clones

were cured of the pDAI-*Sce*I and screened for the loss of the plasmid as described above.

Plasmid construction

To create pKF2, *paaK2* (BCAM1711) was amplified from genomic *B. cenocepacia* K56-

2 DNA using primers 1055/1056and restricted with *Xba*I and *Nde*I before ligation into similarly

digested pSCrhaB2 (39). The plasmid construct was verified using primers 1055/848. To create

the *E. coli* reporter system, *cepR* (BCAM1868) was amplified using primers 703/704 and the

amplicon and pBAD30 were restricted with *Eco*RI and *Xba*I and then ligated to form pTL5. To

construct a plasmid expressing the *luxCDABE* operon, the *cepI* promoter region including the

first 15 codons of *cepI* was amplified and an in-frame stop codon was added using primers 712

and 713. Plasmid pBBRlux and the amplicon were digested with *Spe*I and *Bam*HI, ligation then

resulted in pTL7. Plasmids pTL12 and pTL13 were created in the same way with primers

768/769 for the P_{BAD} promoter and regulatory *araC* and primers 770/771 for the *dhfr* promoter

region of pTnMod-STp. To create a plasmid that expressed PaaK and was compatible with the

reporter strain we had to use the pSC101 ori. pTL9 was created by digesting pTnMod-STp with

*Bgl*II to remove the Tn*5* transposase, then the rhamnose inducible promoter and its regulatory

genes were added to create pTL21 using primers 724 and 947. Using primers 725 and 726 the

paaK ligase from *E. coli* W was amplified from pAAD (27) and restriction cloned into pTL21 to

create pTL22.

Growth of B. cenocepacia *strains on various carbon sources*

One ml of culture from a five ml overnight culture in LB with appropriate antibiotic was

washed in PBS twice and the OD_{600} was adjusted to 0.08 in 1 ml of M9 with 25 mM of Glucose

or M9 with 5 mM PAA. 100 µl was aliquoted to each well of a clear 96-well plate with 100 µl of

appropriate media for a starting OD_{600} of 0.04 in triplicate. Plates were incubated at 37°C with

aeration at 230 rpm and the OD_{600} was read after 48 hours of incubation. OD_{600} values were

converted to 1-cm-path-length by prior calibration with a GeneQuant[TM] III 4283, version

4283V1.6.

Quantification of intracellular and extracellular levels of PAA and PAA-CoA by ion pair

reversed phase ultra high performance liquid chromatography tandem mass spectrometry (IP-

RP-UHPLC-MS/MS)

B. cenocepacia cultures were diluted 1:100 into 10 ml of LB broth in a 50 ml Erlenmeyer

flask and incubated at 37°C with agitation at 230 rpm for 6 hours until late exponential phase.

Using the differential method (21, 22), 1 ml of culture was immediately transferred into weighed

tubes containing 5 ml of 60% (v/v) methanol in water held at -40°C in an acetonitrile/dry-ice

bath, vortexed and returned to -40°C. Two ml of culture was drawn into a syringe, equipped with

a 0.45 µm filter, and 1 ml of filtrate was immediately transferred into 5 ml of 60% (v/v)

methanol in water held at -40°C in an acetonitrile/dry-ice bath, vortexed and returned to -40°C.

Tubes were weighed and 500 µl was transferred to a fresh tube and held at -40°C. 5 ml of 75%

(v/v) ethanol was boiled at 95°C for 4 minutes and then poured into the tube with 500 µl of

sample, mixed and placed at 95°C for 3 minutes. Extracted samples were then put back at -40°C

and evaporated in a speedvac (Savant SC110 with refrigerated condensation trap RT100) until

dry. Care was taken to ensure that the samples were not overdried. All measurements were

carried out using IP-RP-UHPLC-MS/MS as described previously (23) with slight modifications.

Briefly, no splitter or make up flow was used and the chromatographic separation was performed

on a narrower column (1x100 mm) with a lower eluant flow (0.075 ml/min).

For the *E. coli* W14 CepR reporter assay strains, cultures were diluted 1:100 in 5 ml of

LB supplemented with 50 mM MOPS and appropriate antibiotics and grown at 30°C, 230 rpm to

$OD_{600} \sim 0.3$. Cultures were provided 0.5 mM PAA and 0.01% rhamnose to induce the expression

of *paaK* to convert PAA to PAA-CoA. Cultures were incubated again until an OD_{600} of 0.6 and

diluted back to OD_{600} 0.3 before incubation for 1h30 minutes to simulate the conditions of the

reporter assay. Samples were immediately prepared using the differential method described

above.

Slow killing assays in Caenorhabditis elegans

Worms were propagated and maintained on nematode growth medium (NGMI) seeded

with a lawn of *E. coli* OP50 at 16°C. Slow killing assays were performed as described previously

(18). Sixty-millimetre Petri plates filled with 10 ml of NGMII were plated with 50 µl of a

stationary-phase culture adjusted to an OD_{600} of 1.7. Plates were incubated overnight at 37°C to

allow formation of a bacterial lawn. 20 to 40 hypochlorite-synchronized *C. elegans* DH26

nematodes at L4 larval stage were inoculated onto each plate and subsequently incubated at 25°C

for the duration of the assay. Worms were scored as live or dead for a period of 7 days.

Exoprotease assays

Cells from an overnight culture were washed with PBS twice and adjusted to an OD_{600} of

1.0. Three microliters of culture were spotted on 2% skim milk agar plates or 2% skim milk agar

supplemented with 2 mM phenylalanine. Plates were incubated at 37°C for 48 hours. Plates were

prepared in duplicate and three biological replicates were performed. Proteolytic activity was

measured as a zone of clearing surrounding the colony. Images were taken at 48 hours and the

area of the zone of clearance was analyzed using ImageJ software (40). A scale was applied to

the images and a threshold analysis was used to determine the zone of clearance. The area of this

zone was then calculated (minus the area of the colony).

CepI enzymatic assay

CepI enzyme activity was determined spectrophotometrically by measuring *holo*-ACP

formation with dichlorophenylindophenol (DCPIP; = 19100 M-1 cm-1) using the assay

described previously (29, 41). Reaction mixtures contained 50 mM 4-(2-hydroxyethyl)-1-

piperazineethanesulfonic acid (HEPES) pH 7.5, 0.005% Nonidet P-40, 0.13 mM DCPIP, 70 µM

C8-ACP, 4 µM CepI. After 10 minutes of pre-incubation the reaction was initiated by addition of

40 µM S-adenosyl methionine. The compounds were assayed in triplicate at a concentration of

200 µM.

Luminescence assays with CepR reporter strain

The reporter assay was performed based on the assay by McInnis and Blackwell (42)

with slight modifications. Briefly, overnight cultures were prepared from glycerol stocks in LB

with appropriate antibiotics and incubated overnight at 30°C with agitation at 230 rpm. The

culture was diluted 1:100 into 5 ml fresh LB medium supplemented with 50 mM MOPS and

appropriate antibiotics. This subculture was incubated at 30°C with shaking at 230 rpm until it

reached an optical density of 0.6. One hundred microliters of culture were plated into wells with

100 µl of appropriate media with C8-HSL, PAA and arabinose as needed, resulting in a starting

OD_{600} of 0.3. For the reporter strain with pTL22 (rha-inducible *paaK*) the culture was grown and

subcultured as described above. This subculture was incubated at 30°C with shaking at 230 rpm

until it reached an optical density of ~0.3. At this time 0.5 mM PAA and 0.01% rhamnose were

added to induce *paaK* expression resulting in the conversion of PAA to PAA-CoA. The cultures

were incubated at 30°C, 230 rpm until an OD_{600} of 0.6. One hundred microliters of culture were

plated into wells with 100 µl of appropriate media with C8-HSL, PAA, and arabinose and

rhamnose as needed, resulting in a starting OD_{600} of 0.3. Arabinose and rhamnose were used at

final concentrations of 0.2% (v/v) and 0.01% (v/v), respectively. C8-HSL was used at a final

concentration of 20 nM, which was the EC_{50} of C8-HSL for the conditions of this assay as

determined using a dose response curve (Supplemental Fig. S1). PAA was tested at several

concentrations (100 µM, 500 µM, 1 mM and 5 mM) and PAA was added at 500 µM in the

presence of 0.01% rhamnose to produce PAA-CoA. White 96-well microplates with clear

bottoms were used and the lids were treated with a TritonX-100/Ethanol solution. Plates were

incubated at 30°C with agitation in a Biotek Synergy 2 plate reader. OD_{600} and relative

luminescence were measured at 30 min intervals for 4 h. Inhibition was measured at the

timepoint with peak signal intensity (2 h for PAA and 3 h for PAA-CoA).

Reverse Transcription quantitative PCR (RT-qPCR)

B. cenocepacia cells were harvested from stationary cultures (OD_{600} ~7). RNA was

purified and DNase treated using the Ribopure bacteria kit (Ambion) with the DNase treatment

extended to 2 h. RNA quality was confirmed by running on a 2% agarose gel. cDNA was

synthesized using iScript Reverse Transcriptase kit (Bio-Rad) and quantified using iQ SYBR

Green mastermix (Bio-Rad) performed on a CFX96 Touch Real-Time PCR Detection System

(Bio-Rad). Primer efficiency was calculated for each primer set and efficiencies between 95%

and 105% were deemed acceptable. Data were analyzed using the comparative C_T method (43,

44). Genes were normalized to the RNA polymerase sigma factor *sigE* (BCAM0918) as a

reference gene (primers 750 and 751) (7, 45, 46).

LC-MS/MS quantification of AHLs

For AHL extraction, five milliliters of *B. cenocepacia* strains were harvested at the same

time as the samples used for RT-qPCR (see above). Samples were extracted and analyzed as

described previously (47) and 5,6,7,8-Tetradeutero-4-hydroxy-2-heptylquinoline (HHQ-d4) was

used as an internal standard at a concentration of 0.2 ppm.

Acknowledgements

This study was supported by a discovery grant from the Natural Sciences and Engineering

Research Council of Canada (NSERC) awarded to S.T. Cardona. The authors would like to

acknowledge the preliminary work of Samuel J. Wolfram, Vincent Henega and Stacey Line. The

authors thank Dr. Miguel Valvano for generously sharing plasmids pDelCepR and pDelCepI.

pBBRlux was a gift from Dr. Aleksandra Sikora. TJL and STC were responsible for the

experimental design and writing the manuscript. TJL and KLF performed the experimental work.

CR detected PAA and PAA-CoA using IP-RP-UHPLC-MS/MS. SB and LRC performed the

CepI enzyme activity assay. MCG and ED performed the LC-MS/MS and data analysis for AHL

quantifiaction. JLS provided knowledge and equipment for the initial detection of PAA by HPLC

and edited the manuscript.

References

1. Bassler BL. 1999. How bacteria talk to each other: regulation of gene expression by

quorum sensing. Curr Opin Microbiol 2:582–587.

2. Fuqua WC, Winans SC, Greenberg EP. 1994. Quorum sensing in bacteria: the LuxR-LuxI

family of cell density-responsive transcriptional regulators. J Bacteriol 176:269–275.

3. Hanzelka BL, Greenberg EP. 1995. Evidence that the N-terminal region of the *Vibrio*

fischeri LuxR protein constitutes an autoinducer-binding domain. J Bacteriol 177:815–817.

4. Winzer K, Williams P. 2001. Quorum sensing and the regulation of virulence gene

expression in pathogenic bacteria. Int J Med Microbiol IJMM 291:131–143.

5. Mahenthiralingam E, Urban TA, Goldberg JB. 2005. The multifarious, multireplicon

Burkholderia cepacia complex. Nat Rev Microbiol 3:144–156.

6. Darling P, Chan M, Cox AD, Sokol PA. 1998. Siderophore production by cystic fibrosis

isolates of *Burkholderia cepacia*. Infect Immun 66:874–877.

7. O'Grady EP, Viteri DF, Malott RJ, Sokol PA. 2009. Reciprocal regulation by the CepIR

and CciIR quorum sensing systems in *Burkholderia cenocepacia*. BMC Genomics 10:441.

8. Malott RJ, Baldwin A, Mahenthiralingam E, Sokol PA. 2005. Characterization of the cciIR

quorum-sensing system in *Burkholderia cenocepacia*. Infect Immun 73:4982–4992.

9. Lewenza S, Sokol PA. 2001. Regulation of ornibactin biosynthesis and *N*-acyl-L-

homoserine lactone production by CepR in *Burkholderia cepacia*. J Bacteriol 183:2212–

2218.

10. Lewenza S, Conway B, Greenberg EP, Sokol PA. 1999. Quorum sensing in *Burkholderia*

cepacia: identification of the LuxRI homologs CepRI. J Bacteriol 181:748–756.

11. Malott RJ, O'Grady EP, Toller J, Inhülsen S, Eberl L, Sokol PA. 2009. A *Burkholderia*

cenocepacia orphan LuxR homolog is involved in quorum-sensing regulation. J Bacteriol

191:2447–2460.

12. Brown SA, Palmer KL, Whiteley M. 2008. Revisiting the host as a growth medium. Nat

Rev Microbiol 6:657–666.

13. Eisenreich W, Dandekar T, Heesemann J, Goebel W. 2010. Carbon metabolism of

intracellular bacterial pathogens and possible links to virulence. Nat Rev Microbiol 8:401–

412.

14. Fuchs TM, Eisenreich W, Heesemann J, Goebel W. 2012. Metabolic adaptation of human

pathogenic and related nonpathogenic bacteria to extra- and intracellular habitats. FEMS

Microbiol Rev 36:435–462.

15. de Lorenzo V. 2014. From the selfish gene to selfish metabolism: revisiting the central

dogma. BioEssays News Rev Mol Cell Dev Biol 36:226–235.

16. Teufel R, Mascaraque V, Ismail W, Voss M, Perera J, Eisenreich W, Haehnel W, Fuchs G.

2010. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. Proc Natl

Acad Sci U S A 107:14390–14395.

17. Teufel R, Friedrich T, Fuchs G. 2012. An oxygenase that forms and deoxygenates toxic

epoxide. Nature 483:359–362.

18.   Pribytkova T, Lightly TJ, Kumar B, Bernier SP, Sorensen JL, Surette MG, Cardona ST.

2014. The attenuated virulence of a *Burkholderia cenocepacia paaABCDE* mutant is due to

inhibition of quorum sensing by release of phenylacetic acid. Mol Microbiol 94:522–536.

19.   Lightly TJ, Phung RR, Sorensen JL, Cardona ST. 2017. Synthetic cystic fibrosis sputum

medium diminishes *Burkholderia cenocepacia* antifungal activity against *Aspergillus*

fumigatus independently of phenylacetic acid production. Can J Microbiol 1–12.

20.   Ijeme AI, Silvia TC. 2011. 3-Hydroxyphenylacetic Acid Induces The *Burkholderia*

cenocepacia Phenylacetic Acid Degradation Pathway. Towards Understanding The

Contribution Of Aromatic Catabolism To Pathogenesis. Front Cell Infect Microbiol 1.

21.   Taymaz-Nikerel H, de Mey M, Ras C, ten Pierick A, Seifar RM, van Dam JC, Heijnen JJ,

van Gulik WM. 2009. Development and application of a differential method for reliable

metabolome analysis in *Escherichia coli*. Anal Biochem 386:9–19.

22.   Van Gulik WM, Canelas AB, Taymaz-Nikerel H, Douma RD, de Jonge LP, Heijnen JJ.

2012. Fast sampling of the cellular metabolome. Methods Mol Biol Clifton NJ 881:279–

306.

23.   Seifar RM, Ras C, Deshmukh AT, Bekers KM, Suarez-Mendez CA, da Cruz ALB, van

Gulik WM, Heijnen JJ. 2013. Quantitative analysis of intracellular coenzymes in

Saccharomyces cerevisiae using ion pair reversed phase ultra high performance liquid

chromatography tandem mass spectrometry. J Chromatogr A 1311:115–120.

24. Sokol PA, Sajjan U, Visser MB, Gingues S, Forstner J, Kooi C. 2003. The CepIR Quorum-

Sensing System Contributes to the Virulence of *Burkholderia cenocepacia* Respiratory

Infections. Microbiology 149:3649–3658.

25. Uehlinger S, Schwager S, Bernier SP, Riedel K, Nguyen DT, Sokol PA, Eberl L. 2009.

Identification of specific and universal virulence factors in *Burkholderia cenocepacia*

strains using multiple infection hosts. Infect Immun 9:4102–4110.

26. Defoirdt T, Brackman G, Coenye T. 2013. Quorum sensing inhibitors: how strong is the

evidence? Trends Microbiol 21:619–624.

27. Ferrandez A, Prieto MA, Garcia JL, Diaz E. 1997. Molecular characterization of PadA, a

phenylacetaldehyde dehydrogenase from *Escherichia coli*. FEBS Lett 406:23–27.

28. Ferrandez A, Minambres B, Garcia B, Olivera ER, Luengo JM, Garcia JL, Diaz E. 1998.

Catabolism of phenylacetic acid in *Escherichia coli*. Characterization of a new aerobic

hybrid pathway. J Biol Chem 273:25974–25986.

29. Scoffone VC, Chiarelli LR, Makarov V, Brackman G, Israyilova A, Azzalin A, Forneris F,

Riabova O, Savina S, Coenye T, Riccardi G, Buroni S. 2016. Discovery of new

diketopiperazines inhibiting *Burkholderia cenocepacia* quorum sensing in vitro and in vivo.

Sci Rep 6:32487.

30. Bhuiyan MS, Ellett F, Murray GL, Kostoulias X, Cerqueira GM, Schulze KE, Mahamad

Maifiah MH, Li J, Creek DJ, Lieschke GJ, Peleg AY. 2016. *Acinetobacter baumannii*

phenylacetic acid metabolism influences infection outcome through a direct effect on

neutrophil chemotaxis. Proc Natl Acad Sci U S A 113:9599–9604.

31. Musthafa KS, Sivamaruthi BS, Pandian SK, Ravi AV. 2012. Quorum sensing inhibition in

Pseudomonas aeruginosa PAO1 by antagonistic compound phenylacetic acid. Curr

Microbiol 65:475–480.

32. Wang J, Dong Y, Zhou T, Liu X, Deng Y, Wang C, Lee J, Zhang LH. 2013. *Pseudomonas*

aeruginosa cytotoxicity is attenuated at high cell density and associated with the

accumulation of phenylacetic acid. PloS One 8:e60187.

33. Niu C, Clemmer KM, Bonomo RA, Rather PN. 2008. Isolation and Characterization of an

Autoinducer Synthase from *Acinetobacter baumannii*. J Bacteriol 190:3386–3392.

34. Latifi A, Winson MK, Foglino M, Bycroft BW, Stewart GS, Lazdunski A, Williams P.

1995. Multiple homologues of LuxR and LuxI control expression of virulence determinants

and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1.

Mol Microbiol 17:333–343.

35. Sobotková L, Grafková J, Kyslík P. 2002. Effect of phenylacetic acid on the growth and

production of penicillin G acylase of recombinant and host strains derived from *Escherichia*

coli W. Enzyme Microb Technol 31:992–999.

36. Yates EA, Philipp B, Buckley C, Atkinson S, Chhabra SR, Sockett RE, Goldner M,

Dessaux Y, Cámara M, Smith H, Williams P. 2002. N-Acylhomoserine Lactones Undergo

Lactonolysis in a pH-, Temperature-, and Acyl Chain Length-Dependent Manner during

Growth of *Yersinia pseudotuberculosis* and *Pseudomonas aeruginosa*. Infect Immun

70:5635–5646.

37.　Flannagan RS, Linn T, Valvano MA. 2008. A system for the construction of targeted

unmarked gene deletions in the genus *Burkholderia*. Environ Microbiol 10:1652–1660.

38.　Aubert DF, O'Grady EP, Hamad MA, Sokol PA, Valvano MA. 2013. The *Burkholderia*

cenocepacia sensor kinase hybrid AtsR is a global regulator modulating quorum-sensing

signalling. Environ Microbiol 15:372–385.

39.　Cardona ST, Valvano MA. 2005. An expression vector containing a rhamnose-inducible

promoter provides tightly regulated gene expression in *Burkholderia cenocepacia*. Plasmid

54:219–228.

40.　Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nat Methods 9:671–675.

41.　Christensen QH, Grove TL, Booker SJ, Greenberg EP. 2013. A high-throughput screen for

quorum-sensing inhibitors that target acyl-homoserine lactone synthases. Proc Natl Acad

Sci U S A 110:13815–13820.

42.　McInnis CE, Blackwell HE. 2011. Thiolactone modulators of quorum sensing revealed

through library design and screening. Bioorg Med Chem 19:4820–4828.

43.　Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods San Diego Calif 25:402–

408.

44.　Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative C(T)

method. Nat Protoc 3:1101–1108.

45.   Peeters E, Sass A, Mahenthiralingam E, Nelis H, Coenye T. 2010. Transcriptional response

of *Burkholderia cenocepacia* J2315 sessile cells to treatments with high doses of hydrogen

peroxide and sodium hypochlorite. BMC Genomics 11:90.

46.   Wong Y-C, Abd El Ghany M, Ghazzali RNM, Yap S-J, Hoh C-C, Pain A, Nathan S. 2018.

Genetic Determinants Associated With *in vivo* Survival of *Burkholderia cenocepacia* in the

Caenorhabditis elegans Model. Front Microbiol 9:1118.

47.   Chapalain A, Groleau M-C, Le Guillouzer S, Miomandre A, Vial L, Milot S, Déziel E.

2017. Interplay between 4-Hydroxy-3-Methyl-2-Alkylquinoline and N-Acyl-Homoserine

Lactone Signaling in a *Burkholderia cepacia* Complex Clinical Strain. Front Microbiol

8:1021.

786
787

Supplemental Material Figure Legends

(see Supplemental Material document for Tables and Figures)

Table S1. Bacterial strains used in this study

Table S2. Plasmids used in this study.

Table S3. Primers used in this study.

Table S4. Gene fragment sequences flanking the regions of target genes.

Fig. S1: 0.02 μM (20 nM) is the effective concentration (EC$_{50}$) of C8-HSL with the CepR reporter

system. The CepR reporter system was tested with varying concentrations of *N*-octanoyl homoserine

lactone (C8-HSL) to determine the concentration required for half maximal signal. The signal was

measured over 4 hours and the time point where the signal was highest (2 hours) was used. Data was

graphed and the EC$_{50}$ was calculated using GraphPad Prism. Error bars are standard deviations of two

replicates. For the conditions of this reporter system the EC$_{50}$ of C8-HSL is 20 nM.

Fig. S2: Growth of *paaK* ligase mutants on M9 with 5 mM PAA was restored by pKF2, a rhamnose-

inducible *paaK* ligase expression vector. *B. cenocepacia* strains were grown on M9 with 25 mM of

glucose or 5 mM of PAA as a sole carbon source. There was no difference of growth of *B. cenocepacia*

K56-2 containing the empty vector (pSCrhaB2) in the presence of 0.01% (v/v) rhamnose on glucose or

PAA as a sole carbon source. When the *paaK* ligase expression vector (pKF2) was induced with

rhamnose in the K56-2*ΔpaaK1ΔpaaK2* and the K56-2*ΔpaaK2ΔpaaK1* strains growth was restored on

PAA. This was not the case for the empty vector controls, indicating that the PAA pathway interruption in

this mutant can be complemented with the introduction of BCAM1711 (*paaK2*) in trans. Error bars

represent standard deviations of three biological replicates.

Fig. S3: PAA does not accumulate intracellularly and PAA-CoA is barely detectable extracellularly.

Cellulare levels of PAA and PAA-CoA were measured with ion pair reversed phase ultra high

performance liquid chromatography tandem mass spectrometry (IP-RP-UHPLC-MS/MS)**.A)**

PAA levels are very low compared to those measured extracellularly. **B)** PAA-CoA was barely detectable

in extracellular samples. Two biological replicates with three technical replicates were performed.

Fig. S4: *C. elegans* slow killing assays show that the *paaK* single deletion mutants have wild type

levels of virulence but the double *paaK* knockout mutants have increased virulence compared to

wild type in *C. elegans* and exoprotease assays. A) *ΔpaaK2ΔpaaK1* had slightly increased virulence

compared to wild type and was determined as significant according to the log-rank test. **B)** The *ΔpaaK1*

and *ΔpaaK2* single deletion mutants have wild type levels virulence in a *C. elegans* model of infection.

This figure is one assay representative of at least three independent experiments. **C)** Proteolytic activity

was measured as the area of the zone of clearance (excluding colonies) on agar containing 2% skim milk

with or without the addition of 2 mM of phenylalanine. The error bars represent the SD of three

independent experiments.

Fig. S5: Transcription of *cepI* and *cepR* and C8-HSL production is not decreased in any of the PAA

pathway mutants. A) RT-qPCR shows that *cepI* and *cepR* gene expression is not decreased compared to

wild type in the PAA pathway mutants. Error bars are standard deviations of three biological replicates.

B) C8-HSL levels were not decreased in any of the mutants, but C8-HSL levels were higher in the

ΔpaaABCDE and *ΔpaaK2ΔpaaK1* mutants. The bar is the mean of three biological replicates. Virulence

is for *C. elegans* slow killing assays and exoprotease assays as seen in Fig. 3.

Fig. S6: Growth of QS mutants in the PAA pathway mutants show no growth defects in LB.

Mutants were grown in LB from a starting OD_{600} of 0.04 over 24 hours and showed no significant growth

defects compared to wild type. Error bars are standard deviation of three biological replicates.

Fig. S7: The virulence of Δ*paaK2*Δ*paaK1*Δ*cepR* is independent of the CepIR QS system. A)

cepI and *cepR* deletions in wild type and Δ*paaABCDE* mutant backgrounds are attenuated for virulence in

C. elegans. The Δ*paaABCDE* mutant is attenuated compared to wild type but is less attenuated than *cepI*

and *cepR* mutants. **B)** As seen with the Δ*paaK1*Δ*paaK2*Δ*cepR* mutant, the *paaK2*Δ*paaK1*Δ*cepR* mutant

displays CepR-independent virulence in *C. elegans*. A *cepI* deletion in the Δ*paaK2*Δ*paaK1* mutant is

attenuated similar to the *cepI* and *cepR* mutants in wild type backgrounds. **C)** The Δ*paaK2*Δ*paaK1*Δ*cepR*

mutant has wild type levels of protease activity with or without phenylalanine whereas the Δ*cepR* mutant

has decreased protease activity. Protease activity was measured as the area of the zone of clearance

(excluding colonies) on agar containing 2% skim milk with or without the addition of 2 mM of

phenylalanine. **D)** *cepR* transcription is restored to wild type levels in Δ*paaABCDE*Δ*cepI* and

Δ*paaK2*Δ*paaK1*Δ*cepI* indicating that PAA increases *cepI* transcription. **E)** *cepI* is downregulated in all of

the Δ*cepR* mutants. **F)** C8-HSL levels were measured and none of the QS mutants had detectable levels of

C8-HSL. Experiments were performed in triplicate. The error bars represent the SD of three independent

experiments.