An hourglass pattern of inter-embryo gene expression variability and of histone regulation in fly embryogenesis

Jialin Liu[1,2]*, Michael Frochaux[2,3], Vincent Gardeux[2,3], Bart Deplancke[2,3], Marc Robinson-Rechavi[1,2]*

1 Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

2 Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland

3 Laboratory of Systems Biology and Genetics, Institute of Bioengineering, School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne (EPFL)

* corresponding authors: jialin.liu@unil.ch, marc.robinson-rechavi@unil.ch

Gene expression is in part a stochastic process [1–3], which is at odds with the need for precise expression levels in biological processes such as embryonic development. To study the tension between precision and stochasticity of gene expression during development, we quantified expression variability of single embryo transcriptomes throughout fly *Drosophila melanogaster* embryogenesis, from 150 samples over eight developmental stages. We found that expression variability follows an hourglass pattern, with lower variability at extended germband, the phylotypic stage. We explain this pattern by stronger histone modification mediated transcriptional noise control at this stage. In addition, we propose that histone modifications contribute to mutational robustness in regulatory elements, and thus to conserved expression levels. These results provide insight into the role of robustness in the phenotypic and genetic patterns of evolutionary conservation in animal development.

Phenotypes can vary even among isogenic individuals in homogenous environments, suggesting that stochastic effects contribute to phenotypic diversity [4,5]. Gene expression variability among genetically identical individuals under uniform conditions, hereafter "variability", is one of the most important stochastic processes in the mapping of genotype to phenotype. It is caused by a combination of molecular noise (stochastic biochemical effects, e.g., transcriptional burst process based transcriptional noise) and other effects (variation in cells and their environment, e.g., distribution of molecules at cell division) [2,3,6]. Precise regulation of gene expression is notably important during development [7], where it controls the exact timing and level of gene expression to make a functional embryo. However, this process inevitably has to deal with stochasticity [8].

This tension between precision and stochasticity in development raises questions, such as how expression variability is controlled or buffered during development. And whether some stages are more robust to gene expression stochasticity. This is especially interesting in relation to the "phylotypic" stage, in mid-embryonic development, and the corresponding "hourglass" model of Evo-Devo [9,10]. The phylotypic stage is characterized by conserved morphology inside a phylum, which extends to the molecular level, notably lower expression divergence between species or strains [11–15]. To answer these questions, we investigated expression variability across fly embryonic development.

We generated 288 single embryo 3' end transcriptomes using BRB-seq [16], at eight

developmental stages covering the whole fly embryogenesis, with 3h intervals (Figure 1A).

After quality control, 239 samples were kept (Figure S1, S2). On average, we obtained over

5 million uniquely mapped reads of protein coding genes per embryo. Based on

multidimensional scaling analysis (MDS), 150 embryos follow the developmental trajectory,

while there is a small cluster of 89 embryos collected at different time points mixed together

(Figure 1B). The samples in this cluster appear to be unfertilized eggs (Methods and Figure

S3). All further analysis was performed only on the 150 fertilized embryos.

Figure 1: Studying expression variability throughout embryogenesis

A. Method outline. We performed single embryo BRB-seq[16] at eight developmental stages,
indicated by different colored dots. The number of samples collected at each stage is
indicated in the colored triangles. Embryo images adapted from [17] and used with
permission from Springer Nature (License Number: 4547630238607) and from the
authors.

B. Multidimensional scaling analysis (MDS) of 239 high quality samples. Different colors
indicate different stages. The samples can be split into two groups: a small cluster in the
top-left delimited by two red lines; and the remaining samples, which are organized
according to embryonic stage. Only the 150 samples which follow embryonic stages
were used for further analysis.

We measured expression variability as Adjusted SD, standard deviation (SD) of expression

between replicates corrected for expression level (Methods and Figures S4-6). This

expression variability follows an hourglass pattern overall, with a global minimum at E3

(Figure 2A), corresponding to the phylotypic stage of fly [14]. There is also a local minimum

at E6. This is consistent with the pattern of transcriptome divergence between fly and

mosquito *Anopheles gambiae*, with the global minimum at E3, and a local minimum at E6

[18]. Our observations are robust to the use of different variability metrics (Figure S7), and to

sampling (bootstrap analysis, Figure S8). Bootstrap results also suggest that the minimum

of variability extends over E3 to E4. The embryo transcriptome is dominated by zygotic

transcripts 2.5h after egg laying[19], so the high variability in E1 and E2 is not directly caused

by maternal transcripts. We didn't find any significant functional enrichment for genes

which follow the hourglass variability pattern. Overall, expression variability is not equally

distributed throughout embryogenesis, and gene expression at the phylotypic stage appears

more robust to stochastic factors than at other stages.

Figure 2: The phylotypic stage (E3) has lower expression variability

The number of individual samples used in each development stage is indicated below each

box. The number of genes analyzed is indicated in the top-left corner of each plot. The lower

and upper intervals indicated by the dashed lines ("whiskers") represent 1.5 times the

interquartile range (IQR), and the box shows the lower and upper intervals of IQR together

with the median. The black dot in each box indicates the mean.

A. Expression variability pattern of all genes which passed quality control. We performed

pairwise Wilcoxon tests between any two stages to test the significance. The multiple test

corrected p-values (Benjamini–Hochberg method) are shown in Table S1; they are all <

10^{-7}.

B. *Expression variability pattern of genes expressed at all stages. We performed pairwise*
Wilcoxon tests between any two stages to test the significance. The multiple test corrected
p-values (Benjamini–Hochberg method) are shown in Table S2; they are all $< 10^{-5}$ except
E2 vs. E4, for which p-values = 0.24.

The variation in expression variability could either be due to changes in the set of active
genes, with genes differing in their intrinsic variability levels, or to genome-wide changes
in the regulation of variability. To test this, we first reproduced our results restricted to the
subset of genes which are expressed at all stages. Under the first explanation, we would
expect to lose the hourglass variability pattern, but the pattern is maintained (Figure 2B).
We performed additional tests: restricting to genes with constant expression level over
development (Figure S9A); restricting to transcription factors (Figure S9B); and contrasting
genes with dispersed or precise promoters (Figure S10), following Schor et al [20]. Dispersed
promoters seem to be more robust to mutations, which might also translate into robustness
to noise. Despite a loss of power with fewer genes, there remains an hourglass pattern of
expression variability in all cases. Interestingly, the precise promoter genes have higher
variability than the dispersed promoter genes except at E3, thus a strongest hourglass pattern.
Overall, these results suggest that the lower variability at E3 is due to genome-wide
regulation mechanisms more than to changes in the gene set.

Histone modifications can regulate transcriptional noise [21–25], notably through the
modulation of transcriptional burst frequency [22–24]. For example, high levels of histone
modifications can increase chromatin accessibility, leading to an increase in transcriptional
burst frequency, which leads to minimizing noise. To check this role of histone modifications,
we analyzed four available euchromatin histone modifications at six developmental stages [26].
For each gene, we calculated the mean modification signal (background-subtracted tag
density) separately for proximal promoters and for gene bodies [23]. Higher modification signal
genes tend to have lower variability for all histone modifications (Figure 3A). This supports a
role in minimizing transcriptional noise, and is consistent with previous studies in yeast and
mammals [22,23].

Figure 3: Histone modification signal and expression variability

Red and blue represent histone modification signals calculated on the proximal promoter (4kb around the transcription start site – TSS) and the gene body, respectively.

A. *Spearman's correlation coefficient between histone modification signal (background-subtracted tag density) and expression variability. Here, for each gene, both its variability and its histone modification signal are the mean value across stages. ***, P < 0.001; **, P < 0.01; *, P < 0.05, NS, P ≥ 0.05.*

B. *Proximal promoter H3K4Me3 signal (Z score relative to intergenic signal) in different stages. Corresponding stages of our expression variability data are indicated below. The lower and upper intervals indicated by the dashed lines ("whiskers") represent 1.5 times the interquartile range (IQR), and the box shows the lower and upper intervals of IQR together with the median. We performed pairwise Wilcoxon tests between any two stages to test the significance. The multiple test corrected p-values (Benjamini–Hochberg method) are shown in Table S3; they are all $<10^{-7}$ except 4-8h vs. 12-16h, for which p-value = 0.68.*

C. *Gene body H3K4Me3 signal (Z score relative to intergenic signal) in different stages. Corresponding stages of our single embryo BRB-seq data are indicated below. The lower and upper intervals indicated by the dashed lines ("whiskers") represent 1.5 times the interquartile range (IQR), and the box shows the lower and upper intervals of IQR together with the median. We performed pairwise Wilcoxon tests between any two stages to test the significance. The multiple test corrected p-values (Benjamini–Hochberg method) are shown in Table S4; they are all $< 10^{-5}$ except 0-4h vs. 20-24h, for which p-value = 0.26; and 8-12h vs. 16-20h, for which p-value = 0.26.*

The gene-level relation between histone modifications and expression variability raises the possibility that the pattern of expression variability across development could be driven by changes in histone modification signal. To compare histone modification signal between stages,

we normalized gene and promoter signal by that on intergenic regions (Methods), which are

not expected to change histone modification signal between stages. All histone marks present

an hourglass-like pattern, with the highest signal at 8-12h (except for H3K4Me1 on gene body,

where it is a local but not global maximum), corresponding to E3 and E4, i.e. the lowest

expression variability, for both promoters and gene body (Figures 3B-C, S11). Moreover, for

all histone marks on gene body, as well as H3K4Me1 on promoters, there is another local

maximum at 16-20h, corresponding to E6. Generally, these results support changes in histone

modification signal over development, with a correspondence between stronger histone

modification signal and lower expression variability.

Several studies have suggested that mechanisms which confer robustness to stochastic

variation can also buffer the effects of genetic variation [7,27,28]. If histone modifications can

buffer the effect of genetic variation on gene expression, we should observe that genes with

higher histone modification signal are less sensitive to mutations in their regulatory regions,

and are thus less conserved. Indeed, genes with higher histone modification signal tend to

have less conserved core promoter sequences [29] (49 bp upstream TSS and 10 bp downstream

from the TSS) between species (phastCons score; Figure 4A). They are also less conserved

within a population (promoter nucleotide diversity π; Figure S12). The phastCons pattern

remains using 200 bp or 400 bp regions, but disappears using 1 kb regions (Figure S13),

indicating a relatively narrow region around the TSS under this balance of selection and

robustness.

Figure 4: Histone modification signal and promoter sequence conservation

The promoter sequence conservation is the mean of the phastCons score over experimentally identified core promoter regions (49 bp upstream TSS to 10 bp downstream of the TSS) [29].

A. *Spearman's correlation coefficient between histone modification signal (background-subtracted tag density) and promoter sequence conservation. Red and blue represent histone modification signals calculated from the proximal promoter (4 kb around the TSS) and gene body respectively. Here, for each gene, the histone modification signal is the mean value across stages. ***, P < 0.001; **, P < 0.01; *, P < 0.05, NS, P ≥ 0.05.*

B. *Variation of promoter sequence conservation for stage specific genes. The number of genes in each development stage is indicated below each box. We performed pairwise Wilcoxon test between any two stages to test the significance. The multiple test corrected p-values (Benjamini–Hochberg method) are shown in Table S5.*

C. *Transcriptome index of promoter phastCons score across development. The grey area indicates 95% confidence interval estimated from bootstrap analysis.*

Since histone modifications appear to buffer genetic variation in gene expression, and since the E3 stage has stronger modification signals, the lower expression divergence in E3 between species [14] might be caused either by stronger purifying selection on mutations in regulatory regions, or by histone modifications buffering the consequences of mutations in these regions. In the first case, we expect genes specifically expressed at E3 to have higher sequence conservation on promoters. In the second case, we expect the opposite pattern, since mutations that are buffered would behave nearly neutrally. To test this, we identified

genes specifically expressed in each stage and compared their promoter sequence
conservation. We found that genes specific of E3 have a relatively weak promoter sequence
conservation (Figure 4B), supporting a stronger buffering mechanism rather than stronger
purifying selection on sequences. The transcriptome indexes of conservation and of diversity
(mean promoter sequence conservation and mean π, respectively, weighted by expression)
extend this observation to the full transcriptome (Figure 4C; Figure S14). These results
support a role of buffering effects on regulatory mutations in the hourglass pattern of
expression divergence in fly embryogenesis.

We have found an uneven distribution of variability, and thus of robustness of the process
of gene expression, across development, which mirrors the hourglass Evo-Devo model [9,10].
Stage E3 is the most robust to stochastic variation on gene expression, with lower expression
variability, and is the phylotypic stage of fly, with conservation between species [14]. Our
results support a role of increased gene and promoter histone modification signal in the lower
variability in the phylotypic stage.

Although mutational robustness can evolve under natural selection theoretically [30], the
conditions are too restrictive to be relevant in practice. Thus, we propose that the mutational
buffering effect of histone modifications is a by-product of selection for minimizing
transcriptional noise. The mutational buffering analyses have two potential caveats. First, our
results were based solely on promoter analysis, and it remains to be seen how much these
observations extend to other regulatory elements. Second, we only considered nucleotide
substitution changes, but not insertion, deletion and turnover. For these large effect
mutations, purifying selection might be more efficient than mutational robustness to keep
conserved expression level. Thus it is possible that both purifying selection and mutational
robustness together shape the lower expression divergence in the fly phylotypic stage.
Overall, our results support an important role for the control of expression variability by
histone modifications in embryonic gene expression and in the evolution of development.

Acknowledgements

We thank Virginie Braman for help with embryo collection and library preparation. We thank Daniel Alpern for help with the BRB-seq technology. We thank Richard Benton, David Garfield and Laurent Keller for critically reading and commenting on the manuscript. We thank Andrea Komljenovic and other members of the Robinson-Rechavi lab for helpful discussions. Part of the computations were performed at the Vital-IT (http://www.vital-it.ch) Center for high-performance computing of the SIB Swiss Institute of Bioinformatics. JL and MRR are supported by Swiss National Science Foundation grants 31003A_153341 / 1 and 31003A_173048. MRR and BD are supported by SystemsX.ch grant AgingX.

Author contributions

JL designed the work with input from MRR, MF and BD. MF led all experiments. JL performed all computational analyses. VG and BD contributed expertise in the BRB-seq experiments. JL and MRR interpreted the results with input from all the other authors. JL wrote the first draft of the paper. JL and MRR finalized the paper with input from all the other authors.

Supplementary figure legends

Figure S1: Relationship between uniquely mapped reads and expressed genes

Each dot represents one sample. The black dots indicate low quality samples with <4500 expressed genes or with <0.3 million uniquely mapped reads. The 239 orange colored samples were retained for downstream analysis ("high quality samples").

Figure S2: Proportion of retained samples in each development stage

The number of retained samples and of total samples in each stage is indicated in the bottom of each bar.

Figure S3: Evidence that the samples from the small cluster are unfertilized eggs

A. Boxplot of Spearman's correlation coefficients (rho) of expression between individual unfertilized eggs and each sample from the small cluster or from the large cluster, showing that the small cluster has an expression profile of unfertilized eggs. The lower and upper intervals indicated by the dashed lines ("whiskers") represent 1.5 times the interquartile range (IQR), and the box shows the lower and upper intervals of IQR together with the median.

B. Expression heat map of meiosis related genes across all samples, showing that their expression decreases over development for the large cluster, but is high in all samples of the small cluster, consistent with unfertilized eggs.

For testing of an alternative explanation of the two clusters as being males and females, see Figure S16.

Figure S4: Relationship between average expression and coefficient of variation at each stage

Pearson's correlation between average expression and coefficient of variation in each development stage is indicated in the top left of each subfigure.

Figure S5: Relationship between average expression and distance to median at each stage

Pearson's correlation between average expression and distance to median in each development stage is indicated in the top left of each subfigure.

Figure S6: Relationship between average expression and adjusted SD at each stage

Pearson's correlation between average expression and adjusted SD in each development stage

is indicated in the top left of each subfigure.

Figure S7: Variation of expression variability across development using alternate

measures of variability

A. Variability measured by adjusted SD; unlike in Figure 2, the variability in E1 was

calculated using all samples from both small and large clusters.

B. Variability measured by coefficient of variation (CV).

C. Variability measured by distance to median (DM).

The legend is the same as for Figure 2. We performed pairwise Wilcoxon test between any

two stages to test the significance. The multiple test corrected p-values (Benjamini–Hochberg

method) are shown in Tables S6, S7 and S8.

Figure S8: Bootstrap analysis of the variability calculation

We performed pairwise Wilcoxon test between any two stages to test the significance. The

multiple test corrected p-values (Benjamini–Hochberg method) are shown in Table S9.

Figure S9: Variation of expression variability across development for different categories

of genes

A. Genes with constant expression level over development.

B. Transcription factor.

The legend is the same as for Figure 2. We performed pairwise Wilcoxon test between any

two stages to test the significance. The multiple test corrected p-values (Benjamini–Hochberg

method) are shown in Tables S10 and S11.

Figure S10: Variation of expression variability across development for dispersed

promoter genes and for precise promoter genes

For each stage, the first and the second box represents dispersed promoter genes and precise

promoter genes respectively. The legend is the same as for Figure 2. We performed pairwise

Wilcoxon test between any two stages to test the significance separately for dispersed promoter

genes and for precise promoter genes. The multiple test corrected p-values (Benjamini–

Hochberg method) are shown in Tables S12 and S13.

Figure S11: Histone modification signal across development

The legend is the same as for Figure 3B and 3C. The median signal value in each development stage is indicated above each box. We performed pairwise Wilcoxon test between any two stages to test the significance. The multiple test corrected *p*-values (Benjamini–Hochberg method) for H3K4Me1, H3K27Ac and H3K9Ac are shown in Tables S14-S19.

Figure S12: Spearman's correlation coefficient between histone modification signal and promoter nucleotide diversity (π).

The legend is the same as for Figure 4A.

Figure S13: Spearman's correlation coefficient between histone modification signal and promoter sequence conservation for different definitions of promoter width

The figure legend is the same as in Figure 4A.

A. Promoter defined as 200 bp around TSS

B. Promoter defined as 400 bp around TSS

C. Promoter defined as 1000 bp around TSS

Figure S14: transcriptome index of π across development.

The legend is the same as for Figure 4C.

Figure S15: Multidimensional scaling analysis for all samples

Different colors indicate different stages. The solid triangles represent high quality samples according to Figure S1; the hollow triangles represent low quality samples which were discarded.

Figure S16: Mapping of X/autosome gene expression ratios to the multidimensional scaling analysis plot

We calculated the ratio of mean expression between genes from the X chromosome and from the autosomes for each sample. Red represents high ratio, blue represents low ratio. For Drosophila, dosage compensation is achieved by increasing expression of X chromosome genes in males. Since the dosage compensation is still incomplete during development, females should have a higher ratio of mean expression between genes from the X chromosome

339 and from the autosomes. Here, we found both high ratio samples and low ratio samples are
340 well mixed in both the cluster and large clusters. Thus, we reject the hypothesis that the two
341 different clusters are due to sex.
342

Figure S17: Relationship between expression variability and protein importance
We used the average variability across all development stages.
A. Spearman's correlation between expression variability and proportion of essential genes.
We split genes into 10 equally sized bins based on expression variability. The Spearman's
correlation is calculated by using the median variability and proportion of essential genes
in the ten bins. The Spearman's correlation coefficient and p-value are indicated in the top-
right.
B. Spearman's correlation between connectivity in a protein-protein interaction network and
expression variability. The coefficient and p-value are indicated in the top-right.
352

353 **Figure S18: Detection of stage specific genes**
354 A. The artificial expression profile.
355 B. The expression of identified stage specific genes. The bold black line represents the
356   median expression, the two gray lines represent 25th and 75th quantiles of expression,
357   respectively.
358

References

1.   Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic Gene Expression
in a Single Cell. *Science*. **297**, 1183–1186 (2002).

2.   Kærn, M., Elston, T. C., Blake, W. J. & Collins, J. J. Stochasticity in gene expression:
from theories to phenotypes. *Nat. Rev. Genet.* **6**, 451–464 (2005).

3.   Raj, A. & van Oudenaarden, A. Nature, nurture, or chance: stochastic gene expression
and its consequences. *Cell* **135**, 216–26 (2008).

4.   Raj, A., Rifkin, S. A., Andersen, E. & van Oudenaarden, A. Variability in gene
expression underlies incomplete penetrance. *Nature* **463**, 913–8 (2010).

5.   Li, X. *et al.* Systems Properties and Spatiotemporal Regulation of Cell Position
Variability during Embryogenesis. *Cell Rep.* **26**, 313-321.e7 (2019).

6.   Eling, N., Morgan, M. D. & Marioni, J. C. Challenges in measuring and understanding
biological noise. doi:10.1038/s41576-019-0130-6

7.   Waddington, C. H. Canalization of development and genetic assimilation of acquired
characters. *Nature* **150**, 563–565 (1942).

8.   Oates, A. C. What's all the noise about developmental stochasticity? *Development*
138, 601–7 (2011).

9.   Raff, R. A. *The shape of life : genes, development, and the evolution of animal form.*
(University of Chicago Press., 1996).

10.   Duboule, D. Temporal colinearity and the phylotypic progression: a basis for the
stability of a vertebrate Bauplan and the evolution of morphologies through
heterochrony. *Development* **1994**, 135–142 (1994).

11.   Hu, H. *et al.* Constrained vertebrate evolution by pleiotropic genes. *Nat. Ecol. Evol.* **1**,
1722–1730 (2017).

12.   Irie, N. & Kuratani, S. Comparative transcriptome analysis reveals vertebrate
phylotypic period during organogenesis. *Nat. Commun.* **2**, 248 (2011).

13.   Levin, M., Hashimshony, T., Wagner, F. & Yanai, I. Developmental milestones
punctuate gene expression in the Caenorhabditis embryo. *Dev. Cell* **22**, 1101–8 (2012).

14.   Kalinka, A. T. *et al.* Gene expression divergence recapitulates the developmental
hourglass model. *Nature* **468**, 811–4 (2010).

15.   Zalts, H. & Yanai, I. Developmental constraints shape the evolution of the nematode
mid-developmental transition. *Nat. Ecol. Evol.* **1**, 0113 (2017).

16.   Alpern, D. *et al.* BRB-seq: ultra-affordable high-throughput transcriptomics enabled

by bulk RNA barcoding and sequencing. *Genome Biol.* **20**, 71 (2019).

17.   Levin, M. *et al.* The mid-developmental transition and the evolution of animal body

plans. *Nature* **531**, 637–41 (2016).

18.   Schep, A. N. & Adryan, B. A Comparative Analysis of Transcription Factor

Expression during Metazoan Embryonic Development. *PLoS One* **8**, e66826 (2013).

19.   Tadros, W. & Lipshitz, H. D. The maternal-to-zygotic transition: a play in two acts.

Development **136**, 3033–3042 (2009).

20.   Schor, I. E. *et al.* Promoter shape varies across populations and affects promoter

evolution and expression noise. *Nat. Genet.* **49**, 550–558 (2017).

21.   Benayoun, B. A. *et al.* H3K4me3 breadth is linked to cell identity and transcriptional

consistency. *Cell* **158**, 673–88 (2014).

22.   Wu, S. *et al.* Independent regulation of gene expression level and noise by histone

modifications. *PLOS Comput. Biol.* **13**, e1005585 (2017).

23.   Nicolas, D., Zoller, B., Suter, D. M. & Naef, F. Modulation of transcriptional burst

frequency by histone acetylation. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7153–7158

(2018).

24.   Weinberger, L. *et al.* Expression Noise and Acetylation Profiles Distinguish HDAC

Functions. *Mol. Cell* **47**, 193–202 (2012).

25.   Faure, A. J., Schmiedel, J. M. & Lehner, B. Systematic Analysis of the Determinants

of Gene Expression Noise in Embryonic Stem Cells. *Cell Syst.* **5**, 471-484.e4 (2017).

26.   Nègre, N. *et al.* A cis-regulatory map of the Drosophila genome. *Nature* **471**, 527–531

(2011).

27.   Lehner, B. Genes Confer Similar Robustness to Environmental, Stochastic, and

Genetic Perturbations in Yeast. *PLoS One* **5**, e9035 (2010).

28.   Meiklejohn, C. D. & Hartl, D. L. A single mode of canalization. *Trends Ecol. Evol.* **17**,

468–473 (2002).

29.   Dreos, R., Ambrosini, G., Périer, R. C. & Bucher, P. The Eukaryotic Promoter

Database: expansion of EPDnew and new promoter analysis tools. *Nucleic Acids Res.*

43, D92–D96 (2015).

30.   Wagner, G. P., Booth, G. & Bagheri-Chaichian, H. A Population Genetic Theory of

Canalization. *Evolution.* **51**, 329 (1997).

31.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

32.   Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Res.* **46**, D754–D761 (2018).

33.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

427   throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

428   34.   Lun, A. T. L., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level

429   analysis of single-cell RNA-seq data. *F1000Research* **5**, 2122 (2016).

430   35.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

431   differential expression analysis of digital gene expression data. *Bioinformatics* **26**,

432   139–40 (2010).

433   36.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear

434   model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).

435   37.   Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray

436   expression data using empirical Bayes methods. *Biostatistics* **8**, 118–127 (2007).

437   38.   Kolodziejczyk, A. A. *et al.* Single Cell RNA-Sequencing of Pluripotent States Unlocks

438   Modular Transcriptional Variation. *Cell Stem Cell* **17**, 471–85 (2015).

439   39.   Avilés-Pagán, E. E. & Orr-Weaver, T. L. Activating embryonic development in

440   Drosophila. *Semin. Cell Dev. Biol* **84**, *100-110* (2018).

441   40.   Raser, J. M. & O'Shea, E. K. Noise in gene expression: origins, consequences, and

442   control. *Science* **309**, 2010–3 (2005).

443   41.   Tung, P.-Y. *et al.* Batch effects and the effective design of single-cell gene expression

444   studies. *Sci. Rep.* **7**, 39921 (2017).

445   42.   Newman, J. R. S. *et al.* Single-cell proteomic analysis of S. cerevisiae reveals the

446   architecture of biological noise. *Nature* **441**, 840–846 (2006).

447   43.   Barroso, G. V., Puzovic, N. & Dutheil, J. Y. The Evolution of Gene-Specific

448   Transcriptional Noise Is Driven by Selection at the Pathway Level. *Genetics* **208**, 173–

449   189 (2018).

450   44.   Lehner, B. Selection to minimise noise in living systems and its implications for the

451   evolution of gene expression. *Mol. Syst. Biol.* **4**, 170 (2008).

452   45.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

453   genomic features. *Bioinformatics* **26**, 841–2 (2010).

454   46.   Liu, J. & Robinson-Rechavi, M. Adaptive Evolution of Animal Proteins over

455   Development: Support for the Darwin Selection Opportunity Hypothesis of Evo-Devo.

456   *Mol. Biol. Evol.* **35**, 2862–2872 (2018).

47.   Alexa, A., Rahnenfuhrer, J. & Lengauer, T. Improved scoring of functional groups

from gene expression data by decorrelating GO graph structure. *Bioinformatics* **22**,

1600–1607 (2006).

48.   Huang, W. *et al*. Natural variation in genome architecture among 205 Drosophila

melanogaster Genetic Reference Panel lines. *Genome Res*. **24**, 1193–208 (2014).

49.   Danecek, P. *et al*. The variant call format and VCFtools. *Bioinformatics* **27**, 2156–8

(2011).

50.   Liu, J. & Robinson-Rechavi, M. Developmental Constraints on Genome Evolution in

Four Bilaterian Model Species. *Genome Biol. Evol.* **10**, 2266–2277 (2018).

51.   Domazet-Loso, T. & Tautz, D. A phylogenetically based transcriptome age index

mirrors ontogenetic divergence patterns. *Nature* **468**, 815–818 (2010).

52.   Carbon, S. *et al*. AmiGO: online access to ontology and annotation data.

Bioinformatics **25**, 288–9 (2009).

53.   Paris, M., Villalta, J. E., Eisen, M. B. & Lott, S. E. Sex Bias and Maternal

Contribution to Gene Expression Divergence in Drosophila Blastoderm Embryos.

PLoS Genet. **11**, e1005592 (2015).

54.   Chen, W.-H., Lu, G., Chen, X., Zhao, X.-M. & Bork, P. OGEE v2: an update of the

online gene essentiality database with special focus on differentially essential genes in

human cancer cell lines. *Nucleic Acids Res*. **45**, D940–D944 (2017).

55.   Siepel, A. *et al*. Evolutionarily conserved elements in vertebrate, insect, worm, and

yeast genomes. *Genome Res*. **15**, 1034–50 (2005).

Methods

Availability of code

Data files and analysis scripts are available on GitHub:

https://github.com/ljljolinq1010/expression-noise-across-fly-embryogenesis.

Availability of data

Expression datasets have been deposited to the Gene Expression Omnibus with accession number GSE128370.

Embryo collection and RNA extraction

Fly lines (w^{1118}) were obtained from the Bloomington stock center and reared at room temperature on a standard fly medium with 12 hours light dark cycle. The fly medium we used is composed of: 6.2 g Agar powder (ACROS N. 400400050), 58.8 g Farigel wheat (Westhove N. FMZH1), 58.8 g yeast (Springaline BA10), 100 mL grape juice; 4.9 mL Propionic acid (Sigma N. P1386), 26.5 mL of Methyl 4-hydroxybenzoate (VWR N. ALFAA14289.0) solution (400 g/L) in 95% ethanol, 1 L Water. 100 to 150 flies were transferred to cages, which were sealed to a grape agar plate (1:1 mixture of 6% agar and grape juice). We used 4 separate cages to collect the embryos. The adults were kept overnight on this plate before being transferred to a new plate supplemented with yeast paste. Synchronization of eggs on this plate lasted for 2 hours before being swapped with a new plate supplemented with yeast paste. We let the adults lay eggs for 30 min before removing the plate and letting the eggs incubate for the desired time. Eggs were harvested using the following protocol. First a 1:1 bleach (Reactolab 99412) 1x PBS mix was poured on the plate and incubated for 2 min. During this incubation, we used a brush to lightly scrape the surface to mobilize the embryos. We then poured the PBS-bleach mixture through a sieve, washed the plate with 1x PBS, and poured the wash on the same sieve. We washed the sieve several time with 1x PBS until the smell of bleach disappeared. Single embryos were then manually transferred to Eppendorf containing 50 μL beads and 350 μL Trizol (lifetechnologies AM9738). The tubes were homogenized in a Precellys 24 Tissue Homogenizer at 6000 rpm for 30 seconds. Samples were then transferred to liquid nitrogen for flash freezing and stored at –80°C. For RNA extraction, tubes were thawed on ice, supplemented with 350 μL of 100% Ethanol before homogenizing again with the same parameters. We then used the Direct-zol™ RNA Miniprep R2056 Kit, with the following modifications: we did not perform DNAse I treatment, we added another 2 min centrifugation

into an empty column after the RNA Wash step, finally elution was performed by adding 8 μL

of RNAse-free water to the column, incubation at room temperature for 2 min and then

centrifugation for 2 min. RNA was transferred to a low-binding 96 well plate and stored at -

80°C.

Bulk RNA Barcoding and sequencing (BRB-seq)

The BRB-seq is a technique for multiplexed RNA-seq [16] which is able to provide high-quality

3' transcriptomic data at a low cost (e.g. 10-fold lower than Illumina Truseq Stranded mRNA-

seq). The data (fastq files) generated from BRB-seq are multiplexed and asymmetrical paired

reads. Read R1 contains a 6 bp sample barcode, while read R2 contains the fragment sequence

to align to the reference genome.

1. Library preparation

RNA quantity was assessed using picogreen (Invitrogen P11496). Samples were then grouped

according to their concentration in 96-well plates and diluted to a final concentration

determined by the lowest sample concentration on the plate. RNA was then used for gene

expression profiling using BRB-seq. In short, the BRB-seq protocol starts with oligo-dT

barcoding, without TSO for the first-strand synthesis (reverse transcription), performed on each

sample separately. Then all samples are pooled together, after which the second-strand is

synthesized using DNA PolII Nick translation. The sequencing library is then prepared using

cDNA augmented by an in-house produced Tn5 transposase preloaded with the same adapters

(Tn5-B/B), and further enriched by limited-cycle PCR with Illumina compatible adapters.

Libraries are then size-selected (200 - 1000 bp), profiled using High Sensitivity NGS Fragment

Analysis Kit (Advanced Analytical, #DNF-474), and measured using Qubit dsDNA HS Assay

Kit (Invitrogen, #Q32851). In total, we generated four libraries. For details of library

information, please check Table S20.

2. Sequencing

Libraries were mixed in equimolar quantities and were then sequenced on an Illumina Hi-Seq

2500 with pair-end protocol (read R2 with 101 bp) at the Lausanne Genomic Technologies

Facility.

RNA-seq analysis

1. Generating expression matrix

The fastq files were first demultiplexed by using the "Demultiplex" tool from BRB-seqTools

suite (available at https://github.com/DeplanckeLab/BRB-seqTools). Then, we trimmed the

polyA sequences of the demultiplexed files by using the "Trim" tool. Next, the STAR aligner

[31] was used to map the trimmed reads to the reference genome of fly *Drosophila melanogaster*

(BDGP6, Ensembl release 91 [32]). Finally, the read count of each gene was obtained with HTSeq

[33].

2. Filtering samples and genes

Low-quality samples need to be filtered out, since they might bias results of downstream

analyses. In order to assess sample quality, we calculated the number of uniquely mapped reads

and of expressed genes for each sample [34]. We removed samples with <0.3 million uniquely

mapped reads or with <4500 expressed genes (Figure S1). We confirmed that these filtered

samples are indeed outliers in a multidimensional scaling analysis (MDS) (Figure S15). Since

lowly expressed genes have larger technical error, to minimize the technical noise, we need

to remove lowly expressed genes as well. We first calculated counts per million (cpm) with

the edgeR package [35] for each gene. Then we removed genes with mean cpm across samples

≤1, as suggested by Lun et al.[34]. Finally, for the remaining genes, we re-transformed their

cpm values to the original count values for the downstream normalization analysis. After

filtering, we obtained an expression count matrix with 239 samples (Figure S2) and 8004

protein coding genes.

3. Normalization and batch effect correction

Because BRB-Seq retains only the 3′ end of the transcript, we performed sample

normalization by using quantile normalization with log transformation in the voom package [36],

but without transcript length normalization. To remove potential batch effects across the

four libraries, we applied the ComBat function in the sva package [37] to the normalized and

log2 transformed expression data. For genes with expression values less than 0 after Combat,

or with original expression values equal to 0, we change its values to 0 after Combat

correction as suggested by Kolodziejczyk et al [38].

Multidimensional scaling analysis (MDS)

A number of factors could be invoked to explain the two groups observed in our

multidimensional scaling analysis (MDS) (Figure 1B), but they should also explain that only

one group is structured according to developmental time. The obvious hypothesis that they

correspond to male and female embryos does not explain that structure, and is also not

supported by examining X/autosome gene expression ratios (Figure S16). An alternative

hypothesis is that samples in the small cluster are unfertilized eggs. If an egg is not fertilized,

after completion of meiosis, development will be arrested [39], but they are visually

indistinguishable. This hypothesis is confirmed by at least two lines of evidence, in addition

to the lack of developmental time structure. First, for expression correlation, all samples in

the small cluster are highly correlated with unfertilized egg, while the correlations in the

other samples gradually decrease with development (Figure S3A). Second, all the samples

from the small cluster are enriched with meiosis related genes (Figure S3B). Thus we

excluded the small cluster for downstream analyses, i.e. we used 150 embryos with an

average of 18 individuals per developmental stage.

589   Metrics of expression variability

Expression variability is generally measured by the coefficient of variation (CV) [40]. However,

a gene's CV is strongly dependent on its RNA abundance (Figure S4). While this is an inherent

property of time-interval counting processes (such as a Poisson process), it makes the

comparison of variability between different conditions difficult [38,41]. Distance to median (DM,

the distance between the squared CV of a gene and its running median) has been proposed as

a variability metric that is independent of expression level [38,41,42]. However, the DM is still

strongly negatively correlated with the mean expression level in our data (Figure S5). To avoid

this dependency, we developed another variability measure, the adjusted standard deviation

(adjusted SD), by calculating the ratio between observed SD and expected SD. Following the

same approach as Barroso et al. [43], we performed polynomial regressions to predict the

expected SD from mean expression. Since the adjusted SD metric works much better than DM

in terms of accounting for the confounding effects of mean expression (Figure S6), we adopted

it as a measure of expression variability in our study. As observed in yeast [42,44], we found that

essential genes and hubs (proteins in the center of protein-protein interaction network) have

lower expression variability than other genes (Figure S17), indicating selection to reduce it.

This observation provides a control that we are indeed measuring biologically relevant

expression variability.

Detailed calculation of expression variability:

1. Adjusted SD.

For each gene, we computed standard deviation (SD) in each stage and over all stages. Then

we fitted a polynomial model to predict the global (across all stages) SD from the global mean

expression. We increased the degrees of the model until there was no more significant

improvement (tested with ANOVA, $p<0.05$ as a significant improvement). Then, based on this

best fitting model, for each gene, we computed its predicted global SD based on its global mean

expression. Finally, the adjusted SD of a gene in one stage is this gene's SD in its corresponding

stage divided by its predicted global SD. This method is derived from Barroso et al. [43], but

computing adjusted SD rather than adjusted variance.

2. Distance to median: the distance between the squared coefficient of variation (CV) of a

gene and its running median.

For each gene, we computed its squared CV in each stage and over all stages. Then, we ordered

genes based on their global (across all stages) mean expression. Next, we defined series of

sliding windows of 50 genes with 25 genes overlap, starting from lowest global mean

expression. Finally, the distance to median of a gene in one stage is the stage specific log10

squared CV minus the median of global log10 squared CV in this gene's corresponding

window. R code was modified from the DM function of the scran package [34].

Bootstrap analysis

For each stage, we randomly sampled the same number of samples. Then, we computed the

adjusted SD based on these random samples. We repeated the first two steps 500 times. Each

time, we only kept the median of the adjusted SD for each stage. Thus in each stage we obtained

500 medians. Finally, we performed a Wilcoxon test to test the significance of the difference

between the bootstrapped values of different stages.

ChIP-Seq data analysis

1. Histone modification signal datasets

The signal data files of four euchromatin histone modification marks (H3K4me1, H3K4me3,

H3K9ac, and H3K27ac) at six developmental stages (0-4h, 4-8h, 8-12h, 12-16h. 16-20h, 20-

24h) were downloaded from modENCODE [26] (NCBI GEO: GSE16013) (March, 2018). The

signal is smoothed, background-subtracted tag density. The signal was precomputed along

the genome in 35-bp windows.

2. Histone modification signal for promoter and gene body

For each gene, as suggested by Nicolas et al. [23], we separately calculated the mean signal

of its proximal promoter (2 kb upstream to 2 kb downstream for transcription start site (TSS))

and of its gene body (TSS to transcription end site (TES)) by using the bedtools "map"

command [45]. The TSS and TES information was retrieved from Ensembl release 91 [32]. For

a gene with several TSS and TES, we use its mean coordinates.

3. Histone modification signal Z score transformation

For each modification mark in each stage, the signal value was transformed into a Z score by

subtracting the mean signal across intergenic regions and dividing by the standard deviation

signal of the intergenic regions. The intergenic region were defined by removing all proximal

promoter regions and gene body regions with the bedtools "subtract" command [45]. Our

assumption is that on average such intergenic regions are not the target of active histone

modification signal, and thus allow to normalize between libraries. Then, for each gene, we

re-calculated the mean signal (Z score) of its proximal promoter (2 kb upstream to 2 kb

downstream for transcription start site (TSS)) and of its gene body (TSS to transcription end

site (TES)) by using the bedtools "map" command [45].

Identification of stage specifically expressed genes

658   Following the same approach as previously [46], we first defined 8 stage specific expressed

659   artificial expression profile (Figure S18A). Then, for each gene, we performed Pearson's

660   correlation between its real expression and this artificial expression. Finally, for each

661   artificial expression profile, we kept genes with top 10% correlation coefficient as the

662   corresponding stage specifically expressed genes (Figure S18B).

Identification of hourglass expression variability genes

665   Similar to the stage specifically expressed gene identification approach, we correlated each

666   gene's variability profile with the median across all genes. Then, we kept genes with the top

667   10% correlation coefficient as the ones following the global hourglass variability profile.

Identification of genes expressed at all stages

670   For each gene, we calculated the average expression across replicates in each stage. Then,

671   we defined the average expression > 1 as expressed.

Identification of genes with constant expression across all stages

674   For each gene, we first preformed one-way analysis of variance (ANOVA) to compare the

675   means of expression in different stage. Then, we calculated the *q-values* for multiple test

676   correction. Finally, the constantly expressed genes were defined as genes with *q-values >*
677   0.05.

Gene ontology (GO) enrichment analysis

680   We performed GO enrichment analysis for hourglass expression variability genes by using
681   the topGO [47] R package with the "elim" algorithm.

Single Nucleotide Polymorphism (SNP) data

684   The SNP data for 205 *D. melanogaster* inbred lines were downloaded from the Drosophila
685   Genetic Reference Panel (DGRP [48]) (December, 2018).

Nucleotide diversity (π) calculation

688   We calculated nucleotide diversity of promoters with vcftools [49].

Transcriptome index analysis

691   A "transcriptome index" [50,51] is a weighted mean of a feature over all genes, where the
692   weights are the expression levels of the genes at each condition (e.g., developmental stage).
693   The transcriptome index of phastCons was calculated as:

$$TPI_s = \frac{\sum_{i=1}^{n} phastCons_i * e_{is}}{\sum_{i}^{n} e_{is}},$$

695   where s is the developmental stage, $phastCons_i$ is the promoter sequence conservation score
696   of gene i, n is the total number of genes, and e_{is} is the expression level (log transformed) of
697   gene i in developmental stage s. For the transcriptome index of nucleotide diversity (π) the
698   same formula is used, replacing $phastCons_i$ by π_i.

Meiosis related genes and transcription factors

701   The Meiosis related genes and transcription factors were downloaded from AmiGO [52] (May,
702   2018).

Individual unfertilized eggs RNA-seq data

705   The normalized and log transformed expression matrix of individual unfertilized eggs was
706   downloaded from NCBI GEO: GSE68062 [53] (May, 2018).

Dispersed and precise promoters

708 The annotation of genes with dispersed or precise promoters was downloaded from Schor et

710 al [20] (June, 2019). Dispersed promoters are often associated with ubiquitously expressed

711 genes, have more dispersed patterns of transcriptional initiation, and do not contain a TATA

712 box. On the contrary, precise promoters are typically associated with restricted tissue-specific

713 expression and with a TATA box, and have a single predominant TSS.

Essential gene annotation and protein connectivity datasets

716 The gene essentiality and protein connectivity datasets were downloaded from OGEE v2 [54]

717 (March, 2018).

PhastCons score

720 The pre-computed sequence conservation score phastCons [55] of fly genome was downloaded

721 from http://hgdownload.soe.ucsc.edu/goldenPath/dm3/phastCons15way/ (February, 2018).

722 Higher value means higher conservation.

Experimentally validated core promoters

Experimentally validated transcription start sites (TSSs) were downloaded from the

Eukaryotic Promoter Database (EPD) [29] (May, 2018). For a gene with several TSSs, we

selected the most representative one (the TSS that has been validated by the largest number

of samples). The core promoter region was defined as 49 bp upstream TSS to 10 bp

downstream of the TSS [29]. We used EPD defined TSSs here because they are more accurate

for defining core promoters, whose function is expected to be related to sequence

conservation. Whereas for histone modification signal for promoter and gene body we used

Ensembl defined TSSs to be consistent with the source of TES information, and precision

was less important in defining broader proximal promoters.