Tissue micro-RNAs associated with colorectal cancer prognosis: a systematic review

Micro-RNAs associated with colorectal cancer prognosis

Igor Lopes dos Santos[1][¶][*], Karlla Greick Batista Dias Penna[1][&], Megmar Aparecida dos Santos Carneiro[2][&], Larisse Silva Dalla Libera[3][&], Jéssica Enocencio Porto Ramos[1][&], Vera Aparecida Saddi[1][¶]

[1] Programa de Mestrado em Ciências Ambientais e Saúde da Pontifícia Universidade Católica de Goiás - Laboratório de Genética e Biodiversidade, Escola de Ciências Médicas, Farmacêuticas e Biomédicas da Pontifícia Universidade Católica de Goiás, Goiânia, Goiás, Brazil

[2] Laboratório de Virologia, Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, Goiânia, Goiás, Brazil

[3] Ciências da Saúde, Universidade Federal de Goiás, Goiânia, Goiás, Brazil

* Corresponding author

E-mail: igorlsantoss@outlook.com

Address: Escola de Ciências Médicas, Farmacêutica e Biomédicas, Pontifícia

Universidade Católica de Goiás, Área IV, Praça Universitária, 1440 - Setor Leste

Universitário, Goiânia - GO, 74605-010, Brazil.

¶ These authors contributed equally to this work

& These authors also contributed equally to this work

The Authors declare that there is no conflict of interest.

This research received no specific grant from any funding agency in the public,

commercial, or not-for-profit sectors.

Abstract

Colorectal cancer (CRC) is a multifactorial disease commonly diagnosed

worldwide, with high mortality rates. Several studies demonstrate important

associations between differential expression of micro-RNAs (miRs) and the prognosis

of CRC. However, only a few systematic reviews emphasize the most relevant miRs

able to contribute to the establishment of new prognostic biomarkers in CRC

patients. The present study aimed to identify differentially expressed tissue miRs

associated with prognostic factors in CRC patients, through a systematic review of

the Literature. Using the PubMed database, Cochrane Library and Web of Science,

studies published in English evaluating miRs differentially expressed in tumor tissue

and significantly associated with the prognostic aspects of CRC were selected. All

the included studies used RT-PCR (Taqman or SYBR Green) for miR expression

analysis and the period of publication was from 2009 to 2018. A total of 115 articles

accomplished the inclusion criteria and were included in the review. The studies investigated the expression of 102 different miRs associated with prognostic aspects in colorectal cancer patients. The most frequent oncogenic miRs investigated were miR-21, miR-181a, miR-182, miR-183, miR-210 and miR-224 and the hyperexpression of these miRs was associated with distant metastasis, lymph node metastasis and worse survival in patients with CRC. The most frequent tumor suppressor miRs were miR-126, miR-199b and miR-22 and the hypoexpression of these miRs was associated with distant metastasis, worse prognosis and a higher risk of disease relapse (worse disease-free survival). Specific tissue miRs are shown to be promising prognostic biomarkers in patients with CRC, given their strong association with the prognostic aspects of these tumors, however, new studies are necessary to establish the sensibility and specificity of the miRs in order to use them in clinical practice.

Key-words: colorectal cancer, micro-RNA, prognosis, systematic review, oncology

Introduction

Colorectal cancer (CRC) is one of the most prevalent neoplasms in the world, being the second type of cancer more frequent in women and the third type more frequent in men. In Brazil, for the 2018-2019 biennium, it is estimated the occurrence of 36.030 new colorectal cancer cases with 17.380 cases in men and 18.980 cases in women[1].

The survival of patients with CRC is directly associated with the pathological

stage (pTNM) of the disease which is determined after microscope analysis of the

sample obtained by biopsy or surgical resection of the tumor [2,3]. The pTNM stage is a

system of stage classification proposed by the American Joint Committee on Cancer

which determines the degree of tumor development according with the T, N and M

categories. The T category informs about tumor growth, depth and stage of adjacent

tissues invasion. The N category informs about lymph nodes metastasis and the M

category describes the quantity of affected anatomical sites by distant metastasis [3].

On one hand, it is estimated that the survival of patients diagnosed at the first

stages of the disease is up to 90%, but on the other, 30 to 45% of the patients with

late diagnosis will present disease recurrence [4,5]. Factors such as low degree of

differentiation and high levels of serum carcinoembryonic antigen (CEA) are also

described as important for the increase of CRC recurrence after tumor resection [2,6].

However, results from recent studies have demonstrated that the analysis of micro-

RNA expression in CRC can be used as a more sensible prognostic biomarker due

to its potential of predicting disease recurrence, survival and therapeutic response [7-

11].

MicroRNAs (miRs) are small non-coding RNA molecules of approximately 21

to 24 nucleotides in length that modulate gene expression by binding with the 3'

untranslated region (3'UTR) of messenger RNAs resulting in its translational

repression or cleavage [12]. In the literature, more than 2000 miRs have been already

described which demonstrate an essential role in the regulation of eukaryotic cells

and in the pathogenesis of several diseases, including cancer [13,14].

Biogenesis of miRs begins with the activity of RNA polymerase enzyme II which transcribes the primary miRs (pri-miRs) recognized afterwards by DGCR8 protein associated with DROSHA enzyme creating a complex which allows the cleavage of the pri-miRs originating the precursor miRs (pre-miRs). The pre-miRs are carried to the cell cytoplasm by Exportin-5 and cleaved by DICER, generating double stranded miRs which will be recognized by Argonaut proteins promoting the release of one strand of the mature miR. Argonaut proteins in association with the mature miR and other proteins will constitute the RNA-Induced Silencing Complex (RISC) whose targets will be specific mRNAs. [12,15,16]. Fig 1 outlines the miRs biogenesis.

Fig 1. **miRNAs biogenesis and proteins involved in the process.**

Recent studies report that miRs regulate cellular growth and apoptosis, therefore their dysregulation could trigger carcinogenesis [17,18]. MiRs are classified as oncogenic or tumoral suppressor when they inhibit the expression of tumoral suppressor genes or oncogenes respectively. Oncogenic miRs are usually upregulated in the neoplastic process while tumor supressor miRs are usually downregulated in cancer [12,18,19]. Consequently, studies have demonstrated by means of analysis of miR expression, that they can be valuable diagnostic and prognostic biomarkers in cancers, including CRC. [20–23].

Despite the recent improvements in the diagnostic methods for CRC detection, the overall survival for CRC patients is only 65% while the disease recurrence after

tumor resection is approximately 45%, emphasizing the importance of describing

new methods in order to guarantee a better prognosis for these patients [4,24].

Several studies investigate the importance of miRs expression in CRC

prognosis [7–9,11,21–23]. However, there is a lack of systematic reviews in the Literature

integrating and identifying the main miRs significantly associated with the prognostic

aspects of CRC.

Given the relationship between the expression of miRs and the prognosis of

CRC, the present study aims to highlight this association by means of a systematic

review of the Literature, in order to identify the main miRs associated with each

prognostic aspect of CRC. This could contribute to develop new prognostic

biomarkers and could help to identify new molecular targets useful in CRC treatment.

Methods

Study design and search strategy

This study represents a systematic review of the Literature. For the elaboration

of this project a reference review in the PubMed database, Cochrane Library and

Web of Science database was performed in order to identify the relevant studies that

evaluated miR expression associated with CRC prognosis. The Literature search was

performed from February of 2009, year of publication of the first and less recent

selected study, to December of 2018, year of publication of the most recent selected

study. The following search strategy was used for the identification and selection of

the articles: miRNA, colorectal cancer, prognosis. The search was performed in

November 2018 and it was limited to English published articles. At the end of the

search, all the results were compared and reviewed for the inclusion of relevant publications about the subject.

Inclusion criteria

The paper selection was based on criteria related to the study classification and design in order to include the most relevant articles in this review. The relevant criteria related to the classification of the studies were: primary and descriptive studies, studies related to CRC patients and studies completely available online. The relevant criteria related to study design were: articles that evaluate CRC prognosis by overall survival and disease free survival; articles that evaluate miR expression in tumoral tissues compared to normal tissues; articles that evaluate miR expression by Real Time Polymerase Chain Reaction (RT-PCR) using Taqman and/or SYBR Green; articles that evaluate miR expression related to tumor growth/size, lymph nodes metastasis, distant metastasis and TNM stage. Publications belonging to case report, systematic reviews and meta-analysis categories were not included.

Data Extration

The data extraction was performed by means of a collection form pre-developed by the authors in which the following data of each publication were collected: name of the first author, year of publication, micro-RNA studied, sample group, origin of cases, type of specimens analyzed, type of methodology used to evaluate the expression of micro-RNAs, target genes and association with pathological and prognostic clinical aspects.

156 This review followed the Preferred Reporting Items for Systematic Reviews
157 and Meta-Analyses (PRISMA) recommendations.

159 **Statistical analysis and quality assessment**

Due to the heterogeneity of the included studies, it was not possible to perform
a weighted analysis of the results extracted from each article. Therefore, the results
were analyzed qualitatively and presented narratively. A meta-analysis and the
assessment of publication bias were not practically applicable for this study.

Problems with the design and execution of individual studies of healthcare
interventions raise questions about the validity of their findings, and therefore, the
quality of the primary studies was assessed by evaluating eight characteristics
extracted from a checklist provided by the STROBE platform: description of the
studying setting, description of study participants, definitions of all variables,
description of the data sources and measurement tools, justification of sample size,
description of statistical methods, description of results and discussion and
interpretation of results. After evaluating the studies, each attribute was pointed in a
range from 0 to 2 points (0 = not described, 1 = poor described, 2 = well described),
with maximum quality score of 16 points in total.

175 **Data analysis**

176   The data were reviewed and synthesized in semi-structured tables of
177 Microsoft® Office Excel version 2013. The results were analyzed by descriptive
178 statistics and qualitatively tabulated.

Results

Articles selection

After searching for the terms: microRNA, colorectal cancer and prognosis in all three data bases, 924 articles were found. By reading the titles, 29 duplicates were removed and 215 abstracts were selected for complete reading. After analyzing the abstracts, 134 articles which evaluate miR expression associated with CRC prognosis were selected for complete reading. After reading, 19 articles that did not achieve the proposed inclusion criteria were excluded. At the end, 115 articles were included in the systematic review. Fig. 2 outlines the study selection process.

Fig 2. Flow diagram outlining the study selection process.

The quality scores of the studies ranged from 14 to 16 with a mean quality rating of 15.7 (S1 File).

Characteristics of the selected studies

In total, 14.894 CRC patients were studied in the 115 included articles. The number of patients varied from 10 to 765 patients per study. The studies were performed in institutes and universities in Greece, South Korea, United States, Czech Republic, Iran, Sweden, Denmark, Japan and China with most of the research performed in the last two countries. The studies evaluated the expression of 102 miRs associated with CRC prognostics aspects. MiR-21 was the most frequent one, being analyzed in six studies [25–30]. MiR-181a [31–33], miR-16 [34–36] and miR-133b [37–39] were individually investigated in three studies. MiR-126 [8,40], miR-133a [41,42], miR-182

[43,44], miR-183 [45,46], miR-199b [47,48], miR-22 [49,50], miR-224 [51,52], miR-24-3p [53,54], miR-7

[55,56], were separately analyzed in two studies. Seven researches studied two miRs

concomitantly [30,34,38,39,57–59], while the others analyzed a single miR. All studies

analyzed miR expression performing RT-PCR by TaqMan and/or SYBR Green

methods in which 64 studies used SYBR Green while 51 studies used Taqman. The

characteristics of the selected studies in this review are available as supplemental

material (S2 Table).

Studies that investigated the prognosis of CRC patients by means of overall

survival and disease-free survival demonstrated that upregulated miRs associated

with worse prognosis were more frequent than downregulated miRs. In terms of

pathologic clinical aspects, downregulated miRs associated with advanced TNM

stage, tumor growth and lymph node metastasis were more frequent than

upregulated miRs. Nonetheless, upregulated miRs were more associated with distant

metastasis. The association of the miRs differentially expressed in CRC and their

respective prognostic aspects are summarized in Table 1.

Table 1. Micro-RNAs diferentially expressed in CRC and associated prognostic aspects.

Prognostic Aspects	Downregulated micro-RNAs	Upregulated micro-RNAs
Advanced TNM stage	miR-302c [67], miR-466 [68], miR-340 [69], miR-196b-5p [70], miR-944 [71], miR-30d [72], miR-22 [50], miR-296 [73], miR-384 [74], miR-199b [48], miR-128 [75], miR-33b [76], miR-155-5p [77], miR-154 [78], miR-149 [79], miR-194 [80], miR-133b [37], miR-15a [34], miR-100 [81], miR-133a [41], miR-335 [82], miR-378a-3p [58], miR-378a-5p [58], miR-218 [83], miR-16 [34]	miR-182 [43,44], miR-1296 [84], miR-888 [64], miR-21 [29], miR-130a [85], miR-7 [55], miR-96 [86], miR-20a-5p [63], miR-517a [87], miR-106b [88], miR-320e [89], miR-592 [90], miR-196a [57], miR-196b [57], miR-720 [91], miR-133a [42], miR-630 [92], miR-181a [31], miR-210 [93], miR-25 [94], miR-191 [95], miR-92a [96], miR-31[59], miR-552 [97]
Tumor Growth	miR-302c [67], miR-466 [68], let-7a-5p [98], miR-132 [99], miR-206 [100], miR-33b [76], miR-154 [78], miR-194 [80], miR-100 [81], miR-378 [101], miR-378a-3p [58], miR-378a-5p [58], miR-148b [102], miR-30c [103]	miR-1296 [84], miR-193a-3p [104], miR-592 [90], miR-720 [91], miR-210 [93], miR-182 [44], miR-429 [105], miR-125b [66], miR-21 [26,28]
Lymph Node Metastasis	miR-6852 [106], miR-302c [67], miR-466 [68], miR-340 [69], miR-944 [71], miR-30d [72], miR-22 [50], miR-296 [73], miR-199b [47], let-7a-5p [98], miR-132 [99], miR-497 [107], miR-206 [100], miR-128 [75], miR-154 [78], miR-149 [79], miR-24-3p [54], miR-15a [34], miR-16 [34,36], miR-193a-5p [108], miR-100 [81], miR-133a [41], miR-205 [109], miR-335 [82], miR-378 [101], miR-218 [83], miR-138 [110], miR-625 [111], miR-212 [112], miR-195 [113], miR-133b [37], miR-30c [103]	miR-1296 [84], miR-200c [114], miR-21 [26,29], miR-130a [85], miR-7 [55], miR-96 [86], miR-1260b [115], miR-517a [87], miR-106b [88], miR-320e [89], miR-196a [57], miR-196b [57], miR-376a [116], miR-720 [91], miR-494 [61], miR-630 [92], miR-181a [31], miR-20a [62], miR-210 [93], miR-182 [43,44], miR-25 [94], miR-191 [95], miR-183 [46], miR-199a-3p [117], miR-429 [105], miR-92a [96], miR-10b [118], miR-155 [30], miR-552 [97]
Distant Metastasis	miR-466 [68], miR-944 [71], miR-30d [72], miR-22 [49,50], miR-296 [73], miR-140-5p [119], miR-199b [48], miR-155-5p [77], miR-99b-5p [120], miR-154 [78], miR-149 [79], miR-96-5p [121], miR-133b [38,39], miR-126 [40], miR-138 [110], miR-625 [111], miR-212 [112]	miR-888 [64], miR-20a-5p [63], miR-517a [87], miR-320e [89], miR-592 [90], miR-720 [91], miR-494 [61], miR-133a [42], miR-210 [38], miR-630 [92], miR-17-5p [60], miR-181a[31], miR-25 [94], miR-191 [95], miR-183 [46], miR-224 [52], miR-199a-3p [117], miR-92a [96], miR-372 [122], miR-185 [39], miR-21 [26,28,30]
Worse Overall Survival	miR-199b [47,48], miR-6852 [106], miR-302c [67], miR-466 [68], miR-650 [123], miR-196b-5p [70], miR-944 [71], miR-30d [72], miR-296 [73], miR-187 [124], let-7a-5p [98], miR-132 [99], miR-206 [100], miR-148a [125], miR-33b [76], miR-99b-5p [120], miR-154 [78], miR-217 [126], miR-149 [79], miR-194 [80], miR-7 [56], miR-24-3p [54], miR-29b [127], miR-15a [34], miR-16 [34,36], miR-204-5p [128], miR-96-5p [121], miR-193a-5p [108], miR-100 [81], miR-133a [41], miR-139-5p [129], miR-124-5p [130], miR-335 [82], miR-378 [101], miR-126 [40], miR-378a-3p [58], miR-378a-5p [58], miR-218 [83], miR-138 [110], miR-625 [111], miR-212 [112], miR-124 [131], miR-22 [49], miR-133b [39], miR-193a-3p [104], miR-195 [113], miR-30c [103]	miR-181a [31–33], miR-182 [43,44], miR-1296 [84], miR-888 [64], miR-28-5p [132], miR-183 [45,46], miR-200c [114], miR-34a [133], miR-15a-5p [134], miR-181c [135], miR-24-3p [53], miR-16 [35], miR-7 [55], miR-96 [86], miR-1260b [115], miR-20a-5p [63], miR-517a [87], miR-503 [136], miR-106b [88], miR-211 [137], miR-320e [89], miR-592 [90], miR-196a [57], miR-196b [57], miR-376a [116], miR-224 [51,52], miR-720 [91], miR-494 [61], miR-223 [138], miR-133a [42], miR-133b [38], miR-210 [38,93], miR-630 [92], miR-17-5p [60], miR-20a [62], miR-25 [94], miR-191 [95], miR-183 [45,46], miR-199a-3p [117], miR-429 [105], miR-92a [96], miR-372 [122], miR-10b [118], miR-185 [39], miR-21 [25–28,30], miR-155 [30], miR-125b [66], miR-552 [97]
Worse Disease-Free Survival	miR-196b-5p [70], miR-944 [71], miR-30d [72], miR-22 [50], miR-296 [73], miR-199b [47], miR-187 [124], let-7a-5p [98], miR-132 [99], miR-29b [127], miR-15a [34], miR-16 [34], miR-34a-5p [65], miR-193a-5p [108], miR-138 [110], miR-126 [8], miR-212 [112], miR-124 [131]	miR-888 [64], miR-28-5p [132], miR-183 [45], miR-200c [114], miR-34a [133], miR-15a-5p [134], miR-181c [135], miR-24-3p [53], miR-16 [35], miR-96 [86], miR-1260b [115], miR-20a-5p [63], miR-517a [87], miR-503 [136], miR-106b [88], miR-320e [89], miR-196a [57], miR-196b [57], miR-224 [51], miR-494 [61], miR-181a [32], miR-21 [25,26,30], miR-155 [30]

Some of the selected studies investigated the target genes of the miRs

associated with the prognostic aspects of CRC. In total, 65 target genes were

analyzed in which the most frequent ones were: PTEN, investigated in six studies

[29,32,33,50,60,61], SMAD4, investigated in three studies [62–64], and TP53 [65,66] investigated

in two studies.

Micro-RNAs and the prognosis of colorectal cancer

Micro-RNAs associated with advanced TNM stage

Twenty-four studies demonstrated the downregulation of twenty-five miRs

associated with advanced TNM stage: miR-302c [67], miR-466 [68], miR-340 [69], miR-

196b-5p [70], miR-944 [71], miR-30d [72], miR-22 [50], miR-296 [73], miR-384 [74], miR-199b [48],

miR-128 [75], miR-33b [76], miR-155-5p [77], miR-154 [78], miR-149 [79], miR-194 [80],

miR-133b [37], miR-15a [34], miR-100 [81], miR-133a [41], miR-335 [82], miR-378a-3p [58], miR-

378a-5p [58], miR-218 [83] and miR-16 [34]. Two studies demonstrated the upregulation of

miR-182 [43,44] associated with advanced TNM stage and others 22 studies pointed 23

different upregulated miRs associated with advanced TNM stage. In detail, miR-1296

[84], miR-888 [64], miR-21 [29], miR-130a [85], miR-7 [55], miR-96 [86], miR-20a-5p [63], miR-517a

[87], miR-106b [88], miR-320e [89], miR-592 [90], miR-196a [57], miR-196b [57], miR-720 [91], miR-

133a [42], miR-630 [92], miR-181a [31], miR-210 [93], miR-25 [94], miR-191 [95], miR-92a [96],

miR-31[59] and miR-552 [97] have been reported.

Micro-RNAs associated with tumor growth

Tumor growth was associated with the downregulation of 14 miRs reported in 13

studies: miR-302c [67], miR-466 [68], let-7a-5p [98], miR-132 [99], miR-206 [100], miR-33b [76],

miR-154 [78], miR-194 [80], miR-100 [81], miR-378 [101], miR-378a-3p [58], miR-378a-5p [58],

miR-148b [102] and miR-30c [103]. Ten studies verified the upregulation of miR-1296 [84],

miR-193a-3p [104], miR-592 [90], miR-720 [91], miR-210 [93], miR-182 [44], miR-429 [105],miR-

125b [66] and miR-21 [26,28] associated with tumor growth as well.

Micro-RNAs associated with the presence of lymph node metastasis

A total of 32 studies demonstrated the association between the downregulation of

32 miRs and the presence of lymph node metastasis such as miR-6852 [106], miR-

302c [67], miR-466 [68], miR-340 [69], miR-944 [71], miR-30d [72], miR-22 [50], miR-296 [73], miR-

199b [47], let-7a-5p [98], miR-132 [99], miR-497 [107], miR-206 [100], miR-128 [75], miR-154 [78],

miR-149 [79], miR-24-3p [54], miR-15a [34], miR-16 [34,36], miR-193a-5p [108], miR-100 [81],

miR-133a [41], miR-205 [109], miR-335 [82], miR-378 [101], miR-218 [83], miR-138 [110], miR-625

[111], miR-212 [112], miR-195 [113], miR-133b [37] and miR-30c [103]. Other 30 studies verified

the upregulation of miR-1296 [84], miR-200c [114], miR-21 [26,29], miR-130a [85], miR-7 [55],

miR-96 [86], miR-1260b [115], miR-517a [87], miR-106b [88], miR-320e [89], miR-196a [57], miR-

196b [57], miR-376a [116], miR-720 [91], miR-494 [61], miR-630 [92], miR-181a [31], miR-20a [62],

miR-210 [93], miR-182 [43,44], miR-25 [94], miR-191 [95], miR-183 [46], miR-199a-3p [117], miR-

429 [105], miR-92a [96], miR-10b [118], miR-155 [30] and miR-552 [97] associated with the

presence of lymph node metastasis.

Micro-RNAs associated with the presence of distant metastasis

The downregulation of miR-466 [68], miR-944 [71], miR-30d [72], miR-22 [49,50], miR-296

[73], miR-140-5p [119], miR-199b [48], miR-155-5p [77], miR-99b-5p [120], miR-154 [78], miR-149

[79], miR-96-5p [121], miR-133b [38,39], miR-126 [40], miR-138 [110], miR-625 [111] and miR-212

[112] was associated with the presence of distant metastasis in 19 studies. The

upregulation of miR-888 [64], miR-20a-5p [63], miR-517a [87], miR-320e [89], miR-592 [90],

miR-720 [91], miR-494 [61], miR-133a [42], miR-210 [38], miR-630 [92], miR-17-5p [60], miR-

181a[31], miR-25 [94], miR-191 [95], miR-183 [46], miR-224 [52], miR-199a-3p [117], miR-92a [96],

miR-372 [122], miR-185 [39] and miR-21 [26,28,30] was associated with the presence of

distant metastasis in 23 studies.

Micro-RNAs associated with worse overall survival

Among the evaluated studies, 49 demonstrated the downregulation of miR-199b

[47,48], miR-6852 [106], miR-302c [67], miR-466 [68], miR-650 [123], miR-196b-5p [70], miR-944

[71], miR-30d [72], miR-296 [73], miR-187 [124], let-7a-5p [98], miR-132 [99], miR-206 [100], miR-

148a [125], miR-33b [76], miR-99b-5p [120], miR-154 [78], miR-217 [126], miR-149 [79], miR-194

[80], miR-7 [56], miR-24-3p [54], miR-29b [127], miR-15a [34], miR-16 [34,36], miR-204-5p [128],

miR-96-5p [121], miR-193a-5p [108], miR-100 [81], miR-133a [41], miR-139-5p [129], miR-124-

5p [130], miR-335 [82], miR-378 [101], miR-126 [40], miR-378a-3p [58], miR-378a-5p [58], miR-

218 [83], miR-138 [110], miR-625 [111], miR-212 [112], miR-124 [131], miR-22 [49], miR-133b [39],

miR-193a-3p [104], miR-195 [113] and miR-30c [103] associated with worse prognosis

(lower overall survival) of CRC patients. Fifty-six studies identified the upregulation of

miR-181a [31–33], miR-182 [43,44], miR-1296 [84], miR-888 [64], miR-28-5p [132], miR-183 [45,46],

miR-200c [114], miR-34a [133], miR-15a-5p [134], miR-181c [135], miR-24-3p [53], miR-16 [35],

miR-7 [55], miR-96 [86], miR-1260b [115], miR-20a-5p [63], miR-517a [87], miR-503 [136], miR-

106b [88], miR-211 [137], miR-320e [89], miR-592 [90], miR-196a [57], miR-196b [57], miR-376a

[116], miR-224 [51,52], miR-720 [91], miR-494 [61], miR-223 [138], miR-133a [42], miR-133b [38],

miR-210 [38,93], miR-630 [92], miR-17-5p [60], miR-20a [62], miR-25 [94], miR-191 [95], miR-183

[45,46], miR-199a-3p [117], miR-429 [105], miR-92a [96], miR-372 [122], miR-10b [118], miR-185 [39],

miR-21 [25–28,30], miR-155 [30], miR-125b [66] and miR-552 [97] was associated with worse

overall survival of CRC patients.

Micro-RNAs associated with worse disease free-survival

In total, 17 studies demonstrated the downregulation of 18 miRs associated with

worse disease free-survival such as miR-196b-5p [70], miR-944 [71], miR-30d [72], miR-22

[50], miR-296 [73], miR-199b [47], miR-187 [124], let-7a-5p [98], miR-132 [99], miR-29b [127], miR-

15a [34], miR-16 [34], miR-34a-5p [65], miR-193a-5p [108], miR-138 [110], miR-126 [8], miR-212

[112] and miR-124 [131]. Twenty-one studies associated the upregulation of miR-888 [64],

miR-28-5p [132], miR-183 [45], miR-200c [114], miR-34a [133], miR-15a-5p [134], miR-181c [135],

miR-24-3p [53], miR-16 [35], miR-96 [86], miR-1260b [115], miR-20a-5p [63], miR-517a [87], miR-

503 [136], miR-106b [88], miR-320e [89], miR-196a [57], miR-196b [57], miR-224 [51], miR-494 [61],

miR-181a [32], miR-21 [25,26,30] and miR-155 [30] with worse disease free-survival.

Target genes

In order to demonstrate the association among miRs and the clinical and prognostic aspects of CRC, 60 studies investigated also the putative target genes for these miRs. Sixty-five target genes were identified and, among these genes PTEN, SMAD4, P53 and KRAS were the most frequent ones. The downregulation of PTEN was associated with the upregulation of miR-181a [32,33], miR-17-5p [60], miR-494 [61], miR-22 [50] and miR-21 [29]. The downregulation of SMAD4 was associated with the upregulation of miR-20a [62], miR-20a-5p [63] and miR-888 [64]. The downregulation of P53 was related to the upregulation of miR-125b [66] and to the downregulation of de miR-34a-5p [65]. The upregulation of KRAS was associated with the downregulation of miR-96-5p [121] and miR-384 [74].

Discussion

Colorectal cancer is one of the most prevalent neoplasms in the world and it is estimated that the five-year overall survival for CRC patients is only 65%[24]. Several factors associated with CRC prognosis have been already studied, however sensible and specific prognostic biomarkers capable of predicting survival and recurrence of the disease are still unknown [139].

Recent reviews have demonstrated strong correlations between the prognostic aspects of breast cancer [140] and leukemia [141] with miR expression, however there are few systematic reviews integrating the main miRs associated with the prognostic aspects of CRC. The present study evaluated 113 articles that investigated the

differential expression of 100 tissue miRs associated with the main clinicopathological

and prognostic aspects of CRC, demonstrating that such molecules may be useful as

valuable prognostic biomarkers in CRC.

MiRNAs expression can be analyzed in different biological samples such as

tumor tissue [46], blood (serum or plasma) [142], feces [143], or urine [144]. In the present study,

articles that evaluated the expression of miRs in tumor tissues were prioritized and

selected. Tumor tissues can be stored in liquid nitrogen or paraffin, allowing greater

stability to the miRs molecules for later analysis in retrospective studies that aim to

evaluate patients' survival. The analysis in the tumor tissues confers greater specificity

to the results since circulating miRs (present in serum or plasma) may come from

different anatomical sites. In addition, tumor samples allow the analysis of miRs

expression in comparison to the adjacent normal tissue of the same patient providing

greater reliability to the results. In addition, studies indicate that factors associated to

the tumor microenvironment influence the expression of miRs making the analysis in

tumors more adequate for this type of study [145,146].

The expression of miRs can be evaluated by different methods such as northern

blotting, immunohistochemistry, real-time PCR (RT-PCR), microarray, and others [147].

Each method has its own particularities and limitations. In the present study, only

articles that evaluated the expression of miRs in the CRC by means of RT-PCR using

TaqMan or SYBR Green were selected, because these techniques often present hight

sensitivity and specificity, besides allowing the comparison of a specific miR in relation

to an endogenous control.

The miRs are classified as oncogenic or tumor suppressors according to their

respective target genes. MiRs that inhibit the expression of tumor suppressor genes are

considered oncogenic and are generally upregulated in neoplasms. The most frequent

oncogenic miRs in this study were miR-21, miR-181a, miR-182, miR-183, miR-210 and

miR-224 and the upregulation of most of these miRs was associated with distant

metastasis, lymph node metastasis and poorer prognosis in patients with CRC.

Conversely, miRs that inhibit the expression of oncogenes are considered tumor

suppressors and are generally downregulated in the carcinogenic processes. The most

frequent tumor suppressor miRs in this study were miR-126, miR-199b and miR-22 and

their downregulation was associated with the presence of distant metastasis, worse

prognosis and a greater risk of disease relapse (worse disease-free survival). In detail,

the downregulation of miR-199b and miR-22 was also associated with more advanced

TNM stage and the presence of lymph node metastasis. Other miRs such as miR-133a,

miR-133b, miR-16, miR-24-3p and miR-7 were also relatively frequent in the present

study, however, investigations analyzing the expression of these miRs have shown

controversial results. The downregulation of these five miRs was associated with a

worse prognosis and with the exception of miR-7, all the other miRs were

downregulated and associated with the presence of lymph node metastasis.

In addition, other 84 miRs were also identified in the present review and their

differential expression was associated with distinct clinicopathological and prognostic

aspects of CRC, however, these miRs were evaluated in a single study. In general, it

was observed that down and upregulated miRs associated with lymph node metastasis

and worse prognosis (overall survival) in CRC were the most frequently studied.

Among the miRs identified, miR-21 was the most frequent one being analyzed in

six studies. In these studies, an upregulation of miR-21 in the tumor compared to the

adjacent normal tissue was found in addition to a strong correlation with more

aggressive CRC characteristics such as tumor size, lymph node metastasis, distant

metastasis and worse prognosis. A systematic review with meta-analysis conducted by

Peng et al. selected studies that analyzed the role of miR-21 as diagnostic biomarker

and prognostic of CRC, and statistically demonstrated that miR-21 upregulation in the

tumor was associated with worse overall survival and worse disease-free survival, as in

the present review [148]. The findings in both studies suggest the potential of miR-21 to

predict survival and relapse of the disease making it promising biomarker for CRC

prognosis.

Another systematic review with meta-analysis conducted by Gao et al. evaluated

the prognostic value of miRs in colorectal cancer as in the present review. However, in

that review, studies that analyzed the expression of miRs in different types of samples

such as serum, plasma and tissue and performing other methods of detection, as

immunohistochemistry, were considered, so they presented miRs not identified in the

present study. Nevertheless, the findings of the review by Gao et al. and the present

review demonstrate similar results regarding the analysis of miRs expression in tumor

tissues associated with the prognostic aspects of CRC, that is, the upregulation of miR-

181a, miR-224, miR-21 and miR-126 downregulation were also present associated with

worse overall survival in CRC patients [21].

The molecular mechanisms by which polyps and adenomas develop to CRC are

diverse. However, the major altered signal pathways responsible for this progression in

CRC are: MAPK / EGFR pathway, TGFβ pathway and via P13K / AKT pathway [149].

Coincidentally, the most frequent target genes for the miRNAs identified in the present

study were also directly associated with these pathways.

The phosphatase and tensin homolog (PTEN) protein is encoded by a tumor

suppressor gene responsible for the inactivation of pathways associated with cell

proliferation. PTEN negatively regulates the PI3K / AKT pathway by hydrolyzing the

Phosphatidylinositol-3,4,5–trisphosphate (PIP3) into Phosphatidylinositol 4,5-

bisphosphate (PIP2) inactivating AKT, a protein that has antiapoptotic action. Similarly,

PTEN inactivates the MAPK pathway by ERK (Extracellular Regulated Kinase)

dephosphorylation, inactivating the transmission of signals through this pathway and

consequently inhibiting cell proliferation. In addition, PTEN activates RAD51, a DNA-

repairing protein, and CENPC (Centromere Protein C), a protein responsible for the

stability of centromere and chromosomal integrity [150]. Situations in which PTEN is

downregulated result in dysregulation in the cell cycle and destabilization of the cellular

DNA and can trigger carcinogenic processes like CRC. In the present study, the

upregulation of miR-181a, miR-17-5p, miR494, miR-22 and miR-21 was correlated with

the downregulation of PTEN gene and worse prognosis of CRC patients.

The transforming growth factor β (TGFβ) is a tumor suppressor cytokinin protein

known to exert important functions in the regulation of apoptosis, differentiation,

migration and cell proliferation. Through the binding of TGFß to its TGFR2 receptor,

numerous proteins of this pathway are phosphorylated and activated, including the

SMAD4 protein. However, studies indicate that mutations or downregulation of SMAD4,

can hyperactivate TGFβ pathway conferring it the potential to promote tumor growth,

invasion and metastasis [149,151]. In the present study, upregulation of miR-20a, miR-20a-

5p and miR-888 was associated with the downregulation of SMAD4 and the presence of

distant metastasis, advanced TNM stage, worse overall survival and worse disease-free

survival in patients with CRC.

The TP53 gene encodes the tumor suppressor protein P53 that plays important

roles in cell cycle control and apoptosis. MiRs belonging to the miR-34 family can

modulate the antiproliferative and pro-apoptotic activity of P53 by inhibiting oncogenes

and cell cycle associated genes. Thus, downregulation of miR-34 or overexpression of

oncogenic miRs that negatively regulate p53 results in a dysregulation of the cell cycle

leading to a carninogenic process [152]. In the present study, the downregulation of P53

was correlated with the upregulation of miR-125b and downregulation of miR-34a-5p,

resulting in a worse prognosis for patients with CRC.

KRAS is a protein belonging to the RAS family of proteins that control the activation

of several pathways related to cell proliferation, such as the MAPK pathway. MiRs

considered to be tumor suppressors may act as regulators of KRAS expression. Thus,

the downregulation of these miRs may increase the activation of KRAS resulting in a

dysregulated cellular proliferation [153]. The present review demonstrated that

upregulation of KRAS was associated with downregulation of miR-96-5p and miR-384,

which in turn was associated with an advanced TNM stage, presence of distant

metastasis and worse overall survival of patients with CRC. Fig 3 outlines the

differential expression of the miRs and their respective target genes.

Fig 3. Upregulation and downregulation of miRs associated with their respective target genes

and cell functions.

By analyzing the results of the present review fully, it was possible to observe the

large number of miRs associated with the different prognostic aspects of CRC, making

them eligible as possible prognostic markers. However, few articles have analyzed the

expression of two or more miRs concomitantly. Thus, the development of studies that

analyze these miRs synergistically, would allow the identification of network

interactions, making them more sensitive and specific in predicting the prognosis of

patients with CRC and assisting in their therapeutic behavior.

Some limitations were observed during the elaboration of the present study. The

large number of studies and miRs investigated made it difficult to conduct a quantitative

and qualitative analysis capable of demonstrating the most sensitive and specific

differentially expressed miRs associated with each clinical-pathological and prognostic

aspect of the CRC. In addition, the study was limited to articles published in English,

making it harder to obtain data from studies published in other languages.

Conclusion

After analyzing the results of the present systematic review, specific tissue miRs

demonstrate to be promising CRC prognostic biomarkers due to their strong correlation

with prognostic aspects of this disease, including advanced TNM stage, presence of

distant metastasis and lymph node metastasis, tumor growth, prediction of worse

prognosis (worse overall survival) and recurrence of the disease (worse disease-free

survival).

The miR-21 was the most frequent studied miR in the present review among the

miRs identified and its upregulation was associated with tumor size, lymph node

metastasis, distant metastasis and worse prognosis making it a promising biomarker of

CRC prognosis. However, new studies are needed to systematize and identify the most

sensitive and specific miRs associated with each prognostic aspect of CRC in order to

choose the most relevant ones and apply them into clinical practice.

In addition, most studies have shown strong correlations between the expression

of miRs and their respective target genes, so the development of functional studies

aiming to elucidate these associations becomes important as a way to help in the

description of new therapeutic targets of this disease.

The vast number of miRs identified in the present study also translates the

complexity and heterogeneity of CRC, emphasizing the importance of new studies that

clarify the regulatory mechanisms of the expression of these miRs in this tumor,

allowing a better understanding of the molecular biology of this disease.

References

1.   INCA. Estimativa 2018-Incidência de câncer no Brasil [Internet]. Instituto Nacional

de Câncer José Alencar Gomes da Silva. 2017. 130 p. Available from:

http://www.inca.gov.br/estimativa/2018/estimativa-2018.pdf

2.   Mayer RJ, Wolpin Brian M. Systemic Treatment of Colorectal Cancer.

Gastroenterology. 2009;134(5):1296–310.

3.   Beth A. Erickson, Alexander B. Olawaire, Adriana Bermudez E c. G and DGM.

American Joint Committee on Cancer Staging Manual. 8th Edition. Cervix Uteri

[Internet]. 2017. 649–659 p. Available from: www.cancerstaging.org

ajcc@facs.org

4.   Wilhelmsen M, Kring T, Jorgensen LN, Madsen MR, Jess P, Bulut O, et al.

Determinants of recurrence after intended curative resection for colorectal cancer.

Scand J Gastroenterol. 2014;49(12):1399–408.

5.   Siegel R, DeSantis C, Virgo K, Stein K, Mariotto A, Smith T, et al. Cancer

treatment and survivorship statistics, 2012. CA Cancer J Clin. 2012;62(4):220–41.

6.   Destri GL, Rubino AS, Latino R, Giannone F, Lanteri R, Scilletta B, et al.

Preoperative Carcinoembryonic Antigen and Independent Prognostic Factor Still

Reliable. Int Surg. 2015;100:617–25.

7.   Pardini B, Rosa F, Naccarati A, Vymetalkova V, Ye Y, Wu X, et al.

Polymorphisms in microRNA genes as predictors of clinical outcomes in

colorectal cancer patients. Carcinogenesis. 2015;36(1):82–6.

8.   Hansen TF, Christensen RDP, Andersen RF, Sørensen FB, Johnsson A,

Jakobsen A. MicroRNA-126 and epidermal growth factor-like domain 7-an

angiogenic couple of importance in metastatic colorectal cancer. Results from the

Nordic ACT trial. Br J Cancer. 2013;109(5):1243–51.

9.   Rasmussen MH, Jensen NF, Tarpgaard LS, Qvortrup C, Rømer MU, Stenvang J,

et al. High expression of microRNA-625-3p is associated with poor response to

first-line oxaliplatin based treatment of metastatic colorectal cancer. Mol Oncol.

2013;7(3):1–10.

10.   Masuda T, Hayashi N, Kuroda Y, Ito S, Educhi H, Mimori K. MicroRNAs as

biomarkers in colorectal cancer. Cancers (Basel). 2017;9(124):1–15.

11.   Toiyama Y, Hur K, Tanaka K, Inoue Y, Kusunoki M, Boland CR, et al. Serum miR-

200c is a novel prognostic and metastasis-predictive biomarker in patients with

colorectal cancer. Ann Surg. 2014;259(4):735–43.

12.   Gregory RI, Shiekhattar R, Gregory RI, Shiekhattar R. MicroRNA Biogenesis and

Cancer MicroRNA Biogenesis and Cancer. 2005;(9):3509–12.

13.   Hammond SM. An overview of microRNAs. Adv Drug Deliv Rev. 2015;87:3–14.

14.   Falzone L, Scola L, Zanghì A, Biondi A, Cataldo A Di, Libra M, et al. Integrated

analysis of colorectal cancer microRNA datasets : identification of microRNAs

associated with tumor development. 2018;10(5):1000–14.

15.   Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, et al. The nuclear RNase III Drosha

initiates microRNA processing. Nature [Internet]. 2003;425(6956):415–9.

Available from: www.nature.com/nature

16.   MacFarlane L-A, R. Murphy P. MicroRNA: Biogenesis, Function and Role in

Cancer. Curr Genomics. 2010;11(7):537–61.

17.   Moutinho C, Esteller M. MicroRNAs and Epigenetics [Internet]. 1st ed. Vol. 135,

Adv in Canc Res. Elsevier Inc.; 2017. 189–220 p. Available from:

http://dx.doi.org/10.1016/bs.acr.2017.06.003

18.   Calin GA, Croce CM. MicroRNA signatures in human cancers. Nat Rev Cancer.

2006;6(11):857–66.

19.   Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor

suppressors. Dev Biol. 2007;302(1):1–12.

20.   Xi Y, Formentini A, Chien M, Weir DB, Russo JJ, Ju J, et al. Prognostic Values of

microRNAs in Colorectal Cancer. Biomark Insights [Internet]. 2007;1(251):113–

21. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2134920/

21.   Gao S, Zhao Z-Y, Wu R, Zhang Y, Zhang Z-Y. Prognostic value of microRNAs in

colorectal cancer□: a meta-analysis. Cancer Manag. 2018;907–29.

22.   Thomas J, Ohtsuka M, Pichler M, Ling H. MicroRNAs: Clinical relevance in

colorectal cancer. Int J Mol Sci. 2015;16(12):28063–76.

23.   Michael MZ, O' Connor SM, van Holst Pellekaan NG, Young GP, James RJ.

Reduced accumulation of specific microRNAs in colorectal neoplasia. Mol Cancer

Res. 2003;1(12):882–91.

24.   Siegel R, Desantis C, Jemal A. Colorectal Cancer Statistics, 2014. CA Cancer J

Clin [Internet]. 2014;64(1):104–17. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24639052

25.   Mima K, Nishihara R, Yang J, Dou R, Masugi Y, Shi Yy, et al. MicroRNA MIR21

549   (miR-21) and PTGS2 expression in colorectal cancer and patient survival. Clin

550   Cancer Res. 2016;22(15):3841–3848.

551   26.   Fukushima Y, Iinuma H, Tsukamoto M, Matsuda K, Hashiguchi Y. Clinical

552   significance of microRNA-21 as a biomarker in each Dukes' stage of colorectal

553   cancer. Oncol Rep. 2015;33(2):573–82.

554   27.   Chen T-H, Wang K-L, Liu C-N, Lee H, Lin C-C, Huang C-C, et al. The prognostic

555   significance of APC gene mutation and miR-21 expression in advanced-stage

556   colorectal cancer. Color Dis. 2013;15(11):1367–74.

557   28.   Toiyama Y, Takahashi M, Hur K, Nagasaka T, Tanaka K, Inoue Y, et al. Serum

558   miR-21 as a diagnostic and prognostic biomarker in colorectal cancer. J Natl

559   Cancer Inst. 2013;105(12):849–59.

560   29.   Wu Y, Song Y, Xiong Y, Wang X, Xu K, Han B, et al. MicroRNA-21 (Mir-21)

561   Promotes Cell Growth and Invasion by Repressing Tumor Suppressor PTEN in

562   Colorectal Cancer. Cell Physiol Biochem. 2017;43(3):945–58.

563   30.   Shibuya H, Iinuma H, Shimada R, Horiuchi A, Watanabe T. Clinicopathological

564   and Prognostic Value of MicroRNA-21 and MicroRNA-155 in Colorectal Cancer.

565   Oncology. 2010;79:313–20.

566   31.   Ji D, Chen Z, Li M, Zhan T, Yao Y, Zhang Z, et al. MicroRNA-181a promotes

567   tumor growth and liver metastasis in colorectal cancer by targeting the tumor

568   suppressor WIF-1. Mol Cancer. 2014;13(1).

569   32.   Nishimura J, Handa R, Yamamoto H, Tanaka F, Shibata K, Mimori K, et al.

570   microRNA-181a is associated with poor prognosis of colorectal cancer. Oncol

571   Rep. 2012;28(6):2221–6.

572   33.   Li Z, Wang H, Xu Z, Sun Y, Han J. Expression and mechanism of microRNA-181a

573   on incidence and survival in late liver metastases of colorectal cancer. Oncol Rep.

574   2016;35(3):1403–8.

575   34.   Xiao G, Tang H, Wei W, Li J, Ji L, Ge J. Aberrant Expression of MicroRNA-15a

576   and MicroRNA-16 Synergistically Associates with Tumor Progression and

577   Prognosis in Patients with Colorectal Cancer. Gastroenterol Res Pract.

578   2014;2014.

579   35.   Diamantopoulos MA, Kontos CK, Kerimis D, Papadopoulos IN, Scorilas A.

580   Upregulated miR-16 expression is an independent indicator of relapse and poor

581   overall survival of colorectal adenocarcinoma patients. Clin Chem Lab Med.

582   2017;55(5):737–47.

583   36.   Qian J, Jiang B, Li M, Chen J, Fang M. Prognostic significance of microRNA-16

584   expression in human colorectal cancer. World J Surg. 2013;37(12):2944–9.

585   37.   Guo Y, Li X, Lin C, Zhang Y, Hu G, Zhou J, et al. MicroRNA-133b inhibits

586   connective tissue growth factor in colorectal cancer and correlates with the clinical

587   stage of the disease. Mol Med Rep. 2015;11(4):2805–12.

588   38.   Ellermeier C, Vang S, Cleveland K, Durand W, Resnick MB, Brodsky AS.

589   Prognostic microRNA expression signature from examination of colorectal primary

and metastatic tumors. Anticancer Res. 2014;34(8):3957–68.

39.   Akçakaya P, Ekelund S, Kolosenko I, Caramuta S, Özata D, Xie H, et al. miR-185

and miR-133b deregulation is associated with overall survival and metastasis in

colorectal cancer. Int J Oncol. 2011;39(2):311–8.

40.   Liu Y, Zhou Y, Feng X, Yang P, Yang J, Ping A, et al. Low Expression of

MicroRNA-126 is Associated with. Wiley Period. 2014;53:358–365.

41.   Wang LL, Du LT, Li J, Liu YM, Qu AL, Yang YM, et al. Decreased expression of

miR-133a correlates with poor prognosis in colorectal cancer patients. World J

Gastroenterol. 2014;20(32):11340–6.

42.   Wan TMH, Lam CSC, Ng L, Chow AKM, Wong SKM, Li HS, et al. The

Clinicopathological Significance of miR-133a in Colorectal Cancer. Dis Markers.

2014;2014.

43.   Rapti SM, Kontos CK, Papadopoulos IN, Scorilas A. Enhanced miR-182

transcription is a predictor of poor overall survival in colorectal adenocarcinoma

patients. Clin Chem Lab Med. 2014;52(8):1217–27.

44.   Liu H, Du L, Wen Z, Yang Y, Li J, Wang L, et al. Up-regulation of miR-182

expression in colorectal cancer tissues and its prognostic value. Int J Colorectal

Dis. 2013;28(5):697–703.

45.   Chen Y, Song W. Wnt/catenin β1/microRNA 183 predicts recurrence and

prognosis of patients with colorectal cancer. Oncol Lett [Internet].

2018;15(4):4451–6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/29541213%0Ahttp://www.pubmedcentral.nih

.gov/articlerender.fcgi?artid=PMC5835957

46. Zhou T, Zhang GJ, Zhou H, Xiao HX, Li Y. Overexpression of microRNA-183 in

human colorectal cancer and its clinical significance. Eur J Gastroenterol Hepatol.

2014;26(2):229–33.

47. Cristóbal I, Caramés C, Rincón R, Manso R, Madoz-Gúrpide J, Torrejón B, et al.

Downregulation of microRNA-199b predicts unfavorable prognosis and emerges

as a novel therapeutic target which contributes to PP2A inhibition in metastatic

colorectal cancer. Oncotarget [Internet]. 2017;8(25):40169–80. Available from:

http://www.oncotarget.com/fulltext/11174

48. Shen ZZ-LZ, Wang B, Jiang KK-WK, Ye CC-XC, Cheng C, Yan Y-CYY-CY, et al.

Downregulation of miR-199b is associated with distant metastasis in colorectal

cancer via activation of SIRT1 and inhibition of CREB/KISS1 signaling.

Oncotarget [Internet]. 2016;7(23):35092–105. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27145368%5Cnhttp://europepmc.org/abstrac

t/med/27145368

49. Zhang G, Xia S, Tian H, Liu Z, Zhou T. Clinical significance of miR-22 expression

in patients with colorectal cancer. Med Oncol. 2012;29(5):3108–12.

50. Xia S-S, Zhang G-J, Liu Z-L, Tian H-P, He Y, Meng C-Y, et al. MicroRNA-22

suppresses the growth, migration and invasion of colorectal cancer cells through

a Sp1 negative feedback loop. Oncotarget [Internet]. 2017;8(22):36266–78.

Available from:

http://www.oncotarget.com/fulltext/16742%0Ahttp://www.ncbi.nlm.nih.gov/pubmed

/28422727%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC54

82653

51. Adamopoulos PG, Kontos CK, Rapti SM, Papadopoulos IN, Scorilas A. MiR-224

overexpression is a strong and independent prognosticator of short-term relapse

and poor overall survival in colorectal adenocarcinoma. Int J Oncol.

2015;46(2):849–59.

52. Liao WT, Li TT, Wang ZG, Wang SY, He MR, Ye YP, et al. microRNA-224

promotes cell proliferation and tumor growth in human colorectal cancer by

repressing PHLPP1 and PHLPP2. Clin Cancer Res. 2013;19(17):4662–72.

53. Kerimis D, Kontos CK, Christodoulou S, Papadopoulos IN, Scorilas A. Elevated

expression of miR-24-3p is a potentially adverse prognostic factor in colorectal

adenocarcinoma. Clin Biochem [Internet]. 2017;50(6):285–92. Available from:

http://dx.doi.org/10.1016/j.clinbiochem.2016.11.034

54. Gao Y, Liu Y, Du L, Li J, Qu A, Zhang X, et al. Down-regulation of miR-24-3p in

colorectal cancer is associated with malignant behavior. Med Oncol. 2015;32(1).

55. NAGANO Y, TOIYAMA Y, OKUGAWA Y, IMAOKA H, FUJIKAWA H, YASUDA H,

et al. MicroRNA-7 Is Associated with Malignant Potential and Poor Prognosis in

Human Colorectal Cancer. Anticancer Res [Internet]. 2016;36(12):6521–6.

Available from: http://ar.iiarjournals.org/content/36/12/6521.abstract

56.   Suto T, Yokobori T, Yajima R, Morita H, Fujii T, Yamaguchi S, et al. MicroRNA-7 expression in colorectal cancer is associated with poor prognosis and regulates cetuximab sensitivity via EGFR regulation. Carcinogenesis. 2015;36(3):338–45.

57.   Ge J, Chen Z, Li R, Lu T, Xiao G. Upregulation of microRNA-196a and microRNA-196b cooperatively correlate with aggressive progression and unfavorable prognosis in patients with colorectal cancer. Cancer Cell Int. 2014;14(1):1–8.

58.   Li H, Dai S, Zhen T, Shi H, Zhang F, Yang Y, et al. Clinical and biological significance of miR-378a-3p and miR-378a-5p in colorectal cancer. Eur J Cancer [Internet]. 2014;50(6):1207–21. Available from: http://dx.doi.org/10.1016/j.ejca.2013.12.010

59.   Wang CJ, Zhou ZG, Wang L, Yang L, Zhou B, Gu J, et al. Clinicopathological significance of microRNA-31, -143 and -145 expression in colorectal cancer. Dis Markers. 2009;26(1):27–34.

60.   Fang L, Li H, Wang L, Hu J, Jin T, Wang J, et al. MicroRNA-17-5p promotes chemotherapeutic drug resistance and tumour metastasis of colorectal cancer by repressing PTEN expression. Oncotarget [Internet]. 2014;5(10):2974–87. Available from: http://www.oncotarget.com/fulltext/1614

61.   Sun HB, Chen X, Ji H, Wu T, Lu HW, Zhang Y, et al. MiR-494 is an independent prognostic factor and promotes cell migration and invasion in colorectal cancer by directly targeting PTEN. Int J Oncol. 2014;45(6):2486–94.

62. Zhang GJ, Li Y, Zhou H, Xiao HX, Zhou T. miR-20a is an independent prognostic factor in colorectal cancer and is involved in cell metastasis. Mol Med Rep. 2014;10(1):283–91.

63. Cheng D, Zhao S, Tang H, Zhang D, Sun H, Yu F, et al. MicroRNA-20a-5p promotes colorectal cancer invasion and metastasis by downregulating Smad4. Oncotarget [Internet]. 2016;7(29):45199–213. Available from: http://www.oncotarget.com/fulltext/9900

64. Gao S-J, Chen L, Lu W, Zhang L, Wang L, Zhu H-H. miR ⎯ 888 functions as an oncogene and predicts poor prognosis in colorectal cancer. Oncol Lett. 2018;15:9101–9.

65. Gao J, Li N, Dong Y, Li S, Xu L, Li X, et al. MiR-34a-5p suppresses colorectal cancer metastasis and predicts recurrence in patients with stage II/III colorectal cancer. Oncogene. 2015;34(31):4142–52.

66. Nishida N, Yokobori T, Mimori K, Sudo T, Tanaka F, Shibata K, et al. MicroRNA miR-125b is a prognostic marker in human colorectal cancer. Int J Oncol. 2011;38(5):1437–43.

67. Ma W, Liu B, Li J, Jiang J, Zhou R, Huang L, et al. MicroRNA-302c represses epithelial–mesenchymal transition and metastasis by targeting transcription factor AP-4 in colorectal cancer. Biomed Pharmacother [Internet]. 2018;105(March):670–6. Available from: https://doi.org/10.1016/j.biopha.2018.06.025

68.   Tong F, Ying Y, Pan H, Zhao W, Li H, Zhan X. Microrna-466 (Mir-466) functions

as a tumor suppressor and prognostic factor in colorectal cancer (CRC). Bosn J

Basic Med Sci. 2018;18(3):252–9.

69.   Zhang L, Xie F, Tang C, Xu W, Ding X, Liang J. miR-340 suppresses tumor

growth and enhances chemosensitivity of colorectal cancer by targeting RLIP76.

Eur Rev Med Pharmacol Sci. 2017;21:2875–86.

70.   Stiegelbauer V, Vychytilova-Faltejskova P, Karbiener M, Pehserl AM, Reicher A,

Resel M, et al. MiR-196b-5p regulates colorectal cancer cell migration and

metastases through interaction with HOXB7 and GALNT5. Clin Cancer Res.

2017;23(17):5255–66.

71.   Wen L, Li Y, Jiang Z, Zhang Y, Yang B, Han F. MiR-944 inhibits cell migration and

invasion by targeting MACC1 in colorectal cancer. Oncol Rep. 2017;37(6):3415–

22.

72.   Yan L, Qiu J, Yao J. Downregulation of microRNA-30d promotes cell proliferation

and invasion by targeting LRH-1 in colorectal carcinoma. Int J Mol Med.

2017;39(6):1371–80.

73.   He Z, Yu L, Luo S, Li M, Li J, Li Q, et al. miR-296 inhibits the metastasis and

epithelial-mesenchymal transition of colorectal cancer by targeting S100A4. BMC

Cancer. 2017;17(1):1–10.

74.   Wang Y-X, Chen Y-R, Liu S-S, Ye Y-P, Jiao H-L, Wang S-Y, et al. MiR-384

inhibits human colorectal cancer metastasis by targeting KRAS and CDC42.

Oncotarget [Internet]. 2016;7(51):84826–38. Available from:

http://www.oncotarget.com/fulltext/12704

75. Wu L, Shi B, Huang K, Fan G. MicroRNA-128 suppresses cell growth and

metastasis in colorectal carcinoma by targeting IRS1. Oncol Rep.

2015;34(5):2797–805.

76. Liao W, Gu C, Huang A, Yao J, Sun R. MicroRNA-33b inhibits tumor cell growth

and is associated with prognosis in colorectal cancer patients. Clin Transl Oncol.

2016;18(5):449–56.

77. Qu YL, Wang HF, Sun ZQ, Tang Y, Han XN, Yu XB, et al. Up-regulated miR-155-

5p promotes cell proliferation, invasion and metastasis in colorectal carcinoma. Int

J Clin Exp Pathol. 2015;8(6):6988–94.

78. Kai Y, Qiang C, Xinxin P, Miaomiao Z, Kuailu L. Decreased miR-154 expression

and its clinical significance in human colorectal cancer. World J Surg Oncol

[Internet]. 2015;13(195):615–9. Available from: http://dx.doi.org/10.1186/s12957-

015-0607-5

79. Xu K, Liu X, Mao X, Xue L, Wang R, Chen L, et al. MicroRNA-149 suppresses

colorectal cancer cell migration and invasion by directly targeting forkhead box

transcription factor FOXM1. Cell Physiol Biochem. 2015;35(2):499–515.

80. Wang B, Shen ZL, Gao ZD, Zhao G, Wang CY, Yang Y, et al. MiR-194,

commonly repressed in colorectal cancer, suppresses tumor growth by regulating

the MAP4K4/c- Jun/MDM2 signaling pathway. Cell Cycle. 2015;14(7):1046–58.

81.   Chen P, Xi Q, Wang Q, Wei P. Downregulation of microRNA-100 correlates with

737   tumor progression and poor prognosis in colorectal cancer. Med Oncol.

738   2014;31(10):1–6.

739   82.   Sun ZF, Zhang Z, Liu Z, Qiu B, Liu K, Dong G. MicroRNA-335 inhibits invasion

740   and metastasis of colorectal cancer by targeting ZEB2. Med Oncol. 2014;31(6).

741   83.   Yu H, Gao G, Jiang L, Guo L, Lin M, Jiao X, et al. Decreased expression of miR-

742   218 is associated with poor prognosis in patients with colorectal cancer. Int J Clin

743   Exp Pathol. 2013;6(12):2904–11.

744   84.   Tao Y, Ma C, Fan Q, Wang Y, Han T, Sun C. MicroRNA-1296 facilitates

745   proliferation, migration and invasion of colorectal cancer cells by targeting SFPQ.

746   J Cancer. 2018;9(13):2317–26.

747   85.   Chen W, Tong K, Yu J. MicroRNA-130a is upregulated in colorectal cancer and

748   promotes cell growth and motility by directly targeting forkhead box F2. Mol Med

749   Rep. 2017;16(4):5241–8.

750   86.   Rapti SM, Kontos CK, Papadopoulos IN, Scorilas A. High miR-96 levels in

751   colorectal adenocarcinoma predict poor prognosis, particularly in patients without

752   distant metastasis at the time of initial diagnosis. Tumor Biol [Internet].

753   2016;37(9):11815–24. Available from: http://dx.doi.org/10.1007/s13277-016-5023-

754   0

755   87.   Ma W, Yu Q, Jiang J, Du X, Huang L, Zhao L, et al. miR-517a is an independent

756   prognostic marker and contributes to cell migration and invasion in human

colorectal cancer. Oncol Lett. 2016;11(1):2583–9.

88.   Zhang GJ, Li JS, Zhou H, Xiao HX, Li Y, Zhou T. MicroRNA-106b promotes

colorectal cancer cell migration and invasion by directly targeting DLC1. J Exp

Clin Cancer Res. 2015;34(1):1–11.

89.   Perez-Carbonell L, Sinicrope FA, Alberts SR, Oberg AL, Balaguer F, Castells A,

et al. MiR-320e is a novel prognostic biomarker in colorectal cancer. Br J Cancer

[Internet]. 2015;113(1):83–90. Available from:

http://dx.doi.org/10.1038/bjc.2015.168

90.   Liu M, Zhi Q, Wang W, Zhang Q, Fang T, Ma Q. Up-regulation of miR-592

correlates with tumor progression and poor prognosis in patients with colorectal

cancer. Biomed Pharmacother [Internet]. 2015;69:214–20. Available from:

http://dx.doi.org/10.1016/j.biopha.2014.12.001

91.   Wang X, Kuang Y, Shen X, Zhou H, Chen Y, Han Y, et al. Evaluation of miR-720

prognostic significance in patients with colorectal cancer. Tumor Biol.

2014;36(2):719–27.

92.   Chu D, Zheng J, Li J, Li Y, Zhang J, Zhao Q, et al. MicroRNA-630 is a prognostic

marker for patients with colorectal cancer. Tumor Biol. 2014;35(10):9787–92.

93.   Qu A, Du L, Yang Y, Liu H, Li J, Wang L, et al. Hypoxia-inducible MiR-210 is an

independent prognostic factor and contributes to metastasis in colorectal cancer.

PLoS One. 2014;9(3).

94. Li X, Yang C, Wang X, Zhang J, Zhang R, Liu R. The expression of miR-25 is

increased in colorectal cancer and is associated with patient prognosis. Med

Oncol. 2014;31(1).

95. Qin S, Zhu Y, Ai F, Li Y, Bai B, Yao W, et al. MicroRNA-191 correlates with poor

prognosis of colorectal carcinoma and plays multiple roles by targeting tissue

inhibitor of metalloprotease 3 S. Neoplasma. 2014;61(1):27–34.

96. Zhou T, Zhang G, Liu Z, Xia S, Tian H. Overexpression of miR-92a correlates with

tumor metastasis and poor prognosis in patients with colorectal cancer. Int J Color

Dis. 2013;28(11):19–24.

97. Wang N, Liu W. Increased expression of miR-552 acts as a potential predictor

biomarker for poor prognosis of colorectal cancer. Eur Rev Med Pharmacol Sci.

2018;22:412–6.

98. Liu TP, Huang CC, Yeh KT, Ke TW, Wei PL, Yang JR, et al. Down-regulation of

let-7a-5p predicts lymph node metastasis and prognosis in colorectal cancer:

Implications for chemotherapy. Surg Oncol [Internet]. 2016;25(4):429–34.

Available from: http://dx.doi.org/10.1016/j.suronc.2016.05.016

99. Mokutani Y, Uemura M, Munakata K, Okuzaki D, Haraguchi N, Takahashi H, et al.

Down-Regulation of microRNA-132 is Associated with Poor Prognosis of

Colorectal Cancer. Ann Surg Oncol. 2016;23:599–608.

100. Sun P, Sun D, Wang X, Liu T, Ma Z, Duan L. miR-206 is an independent

prognostic factor and inhibits tumor invasion and migration in colorectal cancer.

Cancer Biomarkers. 2015;15(4):391–6.

101.   Zhang G, Zhou H, Xiao H, Li Y, Zhou T. MiR-378 is an independent prognostic

factor and inhibits cell growth and invasion in colorectal cancer. BMC Cancer

[Internet]. 2014;14(1):109. Available from:

http://bmccancer.biomedcentral.com/articles/10.1186/1471-2407-14-109

102.   Song Y, Xu Y, Wang Z, Chen Y, Yue Z, Gao P, et al. MicroRNA-148b suppresses

cell growth by targeting cholecystokinin-2 receptor in colorectal cancer. Int J

Cancer. 2012;131(5):1042–51.

103.   Bin L, Zhang M, Lixia L, Aimin Z, Hua Y, Yanhong S, et al. Down-regulation of

miRNA-30c predicts poor prognosis in Colorectal Cancer patients. Revi Rom Med

Labo. 2016;24(4):369–75.

104.   Lin M, Duan B, Hu J, Yu H, Sheng H, Gao H, et al. Decreased expression of miR-

193a-3p is associated with poor prognosis in colorectal cancer. Oncol Lett.

2017;14(1):1061–7.

105.   Li J, Du L, Yang Y, Wang C, Liu H, Wang L, et al. MiR-429 is an independent

prognostic factor in colorectal cancer and exerts its anti-apoptotic function by

targeting SOX2. Cancer Lett [Internet]. 2013;329(1):84–90. Available from:

http://dx.doi.org/10.1016/j.canlet.2012.10.019

106.   Cui BH, Hong X. Mir-6852 serves as a prognostic biomarker in colorectal cancer

and inhibits tumor growth and metastasis by targeting tcf7. Exp Ther Med.

2018;16(2):879–85.

107. Wang L, Jiang C, Li D, Ge X, Shi Z, Li C, et al. MicroRNA-497 inhibits tumor

growth and increases chemosensitivity to 5-fluorouracil treatment by targeting

KSR1. Oncotarget [Internet]. 2016;7(3). Available from:

http://www.oncotarget.com/fulltext/6545

108. Zhang P, Ji DB, Han HB, Shi YF, Du CZ, Gu J. Downregulation of miR-193a-5p

correlates with lymph node metastasis and poor prognosis in colorectal cancer.

World J Gastroenterol. 2014;20(34):12241–8.

109. Orang A V, Safaralizadeh R, Hosseinpour Feizi MA, Somi MH. Diagnostic and

prognostic value of miR-205 in colorectal cancer. Asian Pac J Cancer Prev.

2014;15(9):4033–7.

110. Long L, Huang G, Zhu H, Guo Y, Liu Y, Huo J. Down-regulation of miR-138

promotes colorectal cancer metastasis via directly targeting TWIST2. J Transl

Med. 2013;11(1):1–10.

111. Lou X, Qi X, Zhang Y, Long H, Yang J. Decreased expression of microRNA-625

is associated with tumor metastasis and poor prognosis in patients with colorectal

cancer. J Surg Oncol. 2013;108(4):230–5.

112. Meng X, Wu J, Pan C, Wang H, Ying X, Zhou Y, et al. Genetic and Epigenetic

Down-regulation of MicroRNA-212 Promotes Colorectal Tumor Metastasis via

Dysregulation of MnSOD. Gastroenterology [Internet]. 2013;145(2):426-436.e6.

Available from: http://dx.doi.org/10.1053/j.gastro.2013.04.004

113. Wang X, Wang J, Ma H, Zhang J, Zhou X. Downregulation of miR-195 correlates

with lymph node metastasis and poor prognosis in colorectal cancer. Med Oncol.

2012;29(34):919–27.

114. Roh MS, Lee HW, Jung SB, Kim K, Lee EH, Park M il, et al. Expression of miR-

200c and its clinicopathological significance in patients with colorectal cancer.

Pathol Res Pract [Internet]. 2018;214(3):350–5. Available from:

http://dx.doi.org/10.1016/j.prp.2018.01.005

115. Liu D-R, Guan Q-L, Gao M-T, Jiang L, Kang H-X. miR-1260b is a Potential

Prognostic Biomarker in Colorectal Cancer. Med Sci Monit. 2016;22:2417–23.

116. Mo Z-H, Wu X-D, Li S, Fei B-Y, Zhang B. Expression and Clinical Significance of

MicroRNA-376a in Colorectal Cancer. Asian Pacific J Cancer Prev.

2014;15(21):9265–70.

117. Wan D, He S, Xie B, Xu G, Gu W, Shen C, et al. Aberrant expression of miR-

199a-3p and its clinical significance in colorectal cancers. Med Oncol. 2013;30(1).

118. Nishida N, Yamashita S, Mimori K, Sudo T, Tanaka F, Shibata K, et al.

MicroRNA-10b is a prognostic indicator in colorectal cancer and confers

resistance to the chemotherapeutic agent 5-fluorouracil in colorectal cancer cells.

Ann Surg Oncol. 2012;19(9):3065–71.

119. Yu L, Lu Y, Han X, Zhao W, Li J, Mao J, et al. microRNA -140-5p inhibits

colorectal cancer invasion and metastasis by targeting ADAMTS5 and IGFBP5.

Stem Cell Res Ther [Internet]. 2016;7(1):1–11. Available from:

http://dx.doi.org/10.1186/s13287-016-0438-5

120.   Li W, Chang J, WWang S, Liu X, Junjie P, Huang D, et al. miRNA-99b-5p

suppresses liver metastasis of colorectal cancer by down-regulating mTOR.

Oncotarget [Internet]. 2015;6(27):24448–62. Available from:

http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L

606354568%5Cnhttp://dx.doi.org/10.18632/oncotarget.4423%5Cnhttp://sfx.library

.uu.nl/utrecht?sid=EMBASE&issn=19492553&id=doi:10.18632/oncotarget.4423&

atitle=miRNA-99b-5p+suppresses+l

121.   Ress AL, Stiegelbauer V, Winter E, Schwarzenbacher D, Kiesslich T, Lax S, et al.

MiR-96-5p influences cellular growth and is associated with poor survival in

colorectal cancer patients. Mol Carcinog. 2015;54(11):1442–50.

122.   Yamashita S, Yamamoto H, Mimori K, Nishida N, Takahashi H, Haraguchi N, et

al. MicroRNA-372 is associated with poor prognosis in colorectal cancer.

Oncology. 2012;82(4):205–12.

123.   Zhou C, Cui F, Li J, Wang D, Wei Y, Wu Y, et al. MiR-650 represses high-risk

non-metastatic colorectal cancer progression via inhibition of AKT2/GSK3beta/E-

cadherin pathway. Oncotarget [Internet]. 2017;8(30):49534–47. Available from:

http://www.impactjournals.com/oncotarget/index.php?journal=oncotarget&page=a

rticle&op=download&path%5B%5D=17743&path%5B%5D=56826%0Ahttp://ovids

p.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emex&NEWS=N&AN=61752

8910%0Ahttp://ovidsp.ovid.com/ovidweb.cgi?T=JS&

124.   Wang ZS, Zhong M, Bian YH, Mu YF, Qin SL, Yu MH, et al. MicroRNA-187

inhibits tumor growth and invasion by directly targeting CD276 in colorectal

cancer. Oncotarget [Internet]. 2016;7(28):44266–76. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/27329595%0Ahttps://www.ncbi.nlm.nih.gov

/pmc/articles/PMC5190094/pdf/oncotarget-07-44266.pdf

125. Hibino Y, Sakamoto N, Naito Y, Goto K, Oo HZ, Sentani K, et al. Significance of

miR-148a in colorectal neoplasia: Downregulation of miR-148a contributes to the

carcinogenesis and cell invasion of colorectal cancer. Pathobiology.

2015;82(5):233–41.

126. Wang B, Shen ZL, Jiang KW, Zhao G, Wang C you, Yan Y chao, et al. MicroRNA-

217 functions as a prognosis predictor and inhibits colorectal cancer cell

proliferation and invasion via an AEG-1 dependent mechanism. BMC Cancer

[Internet]. 2015;15(1):1–11. Available from: ???

127. Inoue A, Yamamoto H, Uemura M, Nishimura J, Hata T, Takemasa I, et al.

MicroRNA-29b is a Novel Prognostic Marker in Colorectal Cancer. Ann Surg

Oncol. 2015;22(April):1410–8.

128. Yin Y, Zhang B, Wang W, Fei B, Quan C, Zhang J, et al. MiR-204-5p inhibits

proliferation and invasion and enhances chemotherapeutic sensitivity of colorectal

cancer cells by downregulating RAB22A. Clin Cancer Res. 2014;20(23):6187–

699.

129. Song M, Yin Y, Zhang J, Zhang B, Bian Z, Quan C, et al. MiR-139-5p inhibits

migration and invasion of colorectal cancer by downregulating AMFR and

NOTCH1. Protein Cell. 2014;5(11):851–61.

130. Jinushi T, Shibayama Y, Kinoshita I, Oizumi S, Jinushi M, Aota T, et al. Low

expression levels of microRNA-124-5p correlated with poor prognosis in

colorectal cancer via targeting of SMC4. Cancer Med. 2014;3(6):1544–52.

131. Wang MJ, Li Y, Wang R, Wang C, Yu YY, Yang L, et al. Downregulation of

microRNA-124 is an independent prognostic factor in patients with colorectal

cancer. Int J Colorectal Dis. 2013;28(2):183–9.

132. Tsiakanikas P, Kontos CK, Kerimis D, Papadopoulos IN, Scorilas A. High

microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term

relapse of node-negative patients and poor overall survival of patients with non-

metastatic disease. Clin Chem Lab Med. 2018;56(6):990–1000.

133. Rapti SM, Kontos CK, Christodoulou S, Papadopoulos IN, Scorilas A. miR-34a

overexpression predicts poor prognostic outcome in colorectal adenocarcinoma,

independently of clinicopathological factors with established prognostic value. Clin

Biochem [Internet]. 2017;50(16–17):918–24. Available from:

http://dx.doi.org/10.1016/j.clinbiochem.2017.06.004

134. Kontos CK, Tsiakanikas P, Avgeris M, Papadopoulos IN, Scorilas A. miR-15a-5p,

A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma.

Mol Diagnosis Ther. 2017;21(4):453–64.

135. Yamazaki N, Koga Y, Taniguchi H, Kojima M, Kanemitsu Y, Saito N, et al. High

expression of miR-181c as a predictive marker of recurrence in stage II colorectal

cancer. Oncotarget [Internet]. 2017;8(4):6970–83. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/28036302

136.   Noguchi T, Toiyama Y, Kitajima T, Imaoka H, Hiro J, Saigusa S, et al. MiRNA-503
Promotes Tumor Progression and Is Associated with Early Recurrence and Poor
Prognosis in Human Colorectal Cancer. Oncol. 2016;90(4):221–31.

137.   Sümbül AT, Göğebakan B, Bayram S, Batmacı CY, Öztuzcu S. MicroRNA 211
expression is upregulated and associated with poor prognosis in colorectal
cancer: a case–control study. Tumor Biol. 2015;36(12):9703–9.

138.   Li Z wu, Yang Y mei, Du L tao, Dong Z, Wang L li, Zhang X, et al. Overexpression
of miR-223 correlates with tumor metastasis and poor prognosis in patients with
colorectal cancer. Med Oncol. 2014;31(11):1–5.

139.   Yiu AJ, Yiu CY. Biomarkers in Colorectal Cancer. Anticancer Res. 2016;36:1093–
338.

140.   Soares BA dos S, Penna KGBD, Saddi VA. Prognostic impact of micro-rna
expression in breast cancer: systematic review. Mastology. 2018;23(4):268–75.

141.   Hajizamani S, Shahjahani M, Shahrabi S, Saki N. MicroRNAs as prognostic
biomarker and relapse indicator in leukemia. Clin Transl Oncol. 2017;19(8):951–
60.

142.   Binderup HG, Madsen JS, Helweg Heegaard NH, Houlind K, Andersen RF,
Brasen CL. Quantification of microRNA levels in plasma – Impact of preanalytical
and analytical conditions. PLoS One. 2018;13(7):1DUMMY.

143. Ahmed FE, Ahmed NaC, Gouda MM, Vos PW, Bonnerup C. RT-qPCR for Fecal

Mature MicroRNA Quantification and Validation. Color Cancer Methods Protoc

Methods Mol Biol. 2018;1765:203–15.

144. Fuessel S, Lohse-Fischer A, Van DV, Solomo K, Erdmann K, Wirth MP.

Quantification of MicroRNAs in Urine-Derived Specimens Susanne. Urothelial

Carcinoma Methods Protoc Methods Mol Biol. 2018;1655(16):201–26.

145. Wentz-Hunter KK, Potashkin JA. The Role of miRNAs as Key Regulators in the

Neoplastic Microenvironment. Mol Biol Int. 2011;2011:1–8.

146. Radovich M, Sood AK, Armaiz-Pena GN, Rodriguez-Aguayo C, Zhang W,

Pradeep S, et al. Hypoxia-mediated downregulation of miRNA biogenesis

promotes tumour progression. Nat Commun [Internet]. 2014;5(1):1–13. Available

from: http://dx.doi.org/10.1038/ncomms6202

147. Hennessy E, Driscoll LO. MicroRNA Expression Analysis: Techniques Suitable for

Studies of Intercellular and Extracellular MicroRNA. In: Gene Expression Profiling:

Methods and Protocols, Methods in Molecular Biology. 2011. p. 99–107.

148. Peng Q, Zhang X, Min M, Zou L, Shen P, Yaqun Z. The clinical role of microRNA-

21 as a promising biomarker in the diagnosis and prognosis of colorectal cancer:

a systematic review and meta-analysis. Oncotarget. 2017;8(27):44893–909.

149. Shirafkan N, Mansoori B, Mohammadi A, Shomali N, Ghasbi M, Baradaran B.

MicroRNAs as novel biomarkers for colorectal cancer: New outlooks. Biomed

Pharmacother [Internet]. 2018;97(August 2017):1319–30. Available from:

http://dx.doi.org/10.1016/j.biopha.2017.11.046

150. Chung JH, Eng C. Nuclear-cytoplasmic partitioning of phosphatase and tensin

homologue deleted on chromosome 10 (PTEN) differentially regulates the cell

cycle and apoptosis. Cancer Res. 2005;65(18):8096–100.

151. Zhang B, Halder SK, Kashikar ND, Cho Y, Datta A, David L, et al. Anti-metastatic

Role of Smad4 Signaling in Colorectal Cancer Bixiang. Gastroenterology.

2010;138(3):969–80.

152. Oren M, Bentwich Z, Rosenfeld N, Raver-Shapira N, Moskovits N, Spector Y, et

al. Transcriptional Activation of miR-34a Contributes to p53-Mediated Apoptosis.

Mol Cell. 2007;26(5):731–43.

153. Jones RP, Sutton PA, Evans JP, Clifford R, McAvoy A, Lewis J, et al. Specific

mutations in KRAS codon 12 are associated with worse overall survival in patients

with advanced and recurrent colorectal cancer. Br J Cancer [Internet].

2017;116(7):923–9. Available from: http://dx.doi.org/10.1038/bjc.2017.37