Tissue micro-RNAs associated with colorectal cancer prognosis: a systematic review

Igor Lopes dos Santos[1,¶,*], Karlla Greick Batista Dias Penna[1,&], Megmar Aparecida dos Santos Carneiro[2,&], Larisse Silva Dalla Libera[3,&], Jéssica Enocencio Porto Ramos[1,&], Vera Aparecida Saddi[1,¶]

[1] Programa de Mestrado em Ciências Ambientais e Saúde da Pontifícia Universidade Católica de Goiás - Laboratório de Genética e Biodiversidade, Escola de Ciências Médicas, Farmacêuticas e Biomédicas da Pontifícia Universidade Católica de Goiás, Goiânia, Goiás, Brazil

Escola de Ciências Médicas, Farmacêuticas e Biomédicas, Pontifícia Universidade Católica de Goiás, Goiânia, Goiás, Brazil

[2] Laboratório de Virologia, Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, Goiânia, Goiás, Brazil

[3] Ciências da Saúde, Universidade Federal de Goiás, Goiânia, Goiás, Brazil

* Corresponding author

E-mail: igorlsantoss@outlook.com

¶ These authors contributed equally to this work

22   & These authors also contributed equally to this work

Abstract

Colorectal cancer (CRC) is a multifactorial disease commonly diagnosed

worldwide, with high mortality rates. Several studies demonstrate important

associations between differential expression of micro-RNAs (miRs) and the prognostic

of CRC. However, only a few systematic reviews emphasize the major miRs able to

contribute to the establishment of new prognostic biomarkers in CRC patients. The

present study aimed to identify differentially expressed tissue miRs associated with

prognostic factors in CRC patients, through a systematic review of the literature. Using

the PubMed database, Cochrane Library and Web of Science, studies published in

English evaluating miRs differentially expressed in tumor tissue and significantly

associated with the prognostic aspects of CRC were selected. All the included studies

used RT-PCR (Taqman or SYBR Green) for miR expression analysis and the period

of publication was from 2009 to 2018. A total of 115 articles accomplished the inclusion

criteria and were included in the review. The studies investigated the expression of 102

different miRs associated with prognostic aspects in colorectal cancer patients. The

most frequent oncogenic miRs investigated were miR-21, miR-181a, miR-182, miR-

183, miR-210 and miR-224 and the hyperexpression of these miRs was associated

with distant metastasis, lymph node metastasis and worse survival in patients with

CRC. The most frequent tumor suppressor miRs were miR-126, miR-199b and miR-

22 and the hypoexpression of these miRs was associated with distant metastasis,

worse prognosis and a higher risk of disease relapse (worse disease-free survival). In

general, lymph node metastases and lower overall survival in patients with CRC were

the most frequently investigated prognostic variables associated with the expression

of tumor miRs. Specific tissue miRs are shown to be promising prognostic biomarkers

in patients with CRC, given their strong association with the prognostic aspects of

these tumors, however, new studies are necessary to establish the sensibility and

specificity of the miRs in order to use them in clinical practice.

52   Introduction

Colorectal cancer (CRC) is one of the most prevalent neoplasms in the world,

being the second type of cancer more frequent in women and the third type more

frequent in men. In Brazil, for the 2018-2019 biennium, it's estimated the occurrence

of 36.030 new colorectal cancer cases with 17.380 cases in men and 18.980 cases in

women[1].

The survival of patients with colorectal is directly associated with the

pathological stage (pTNM) of the disease which is determined after microscope

analysis of the sample obtained by biopsy or surgical resection of the tumor [2,3]. The

pTNM stage is a system of stage classification proposed by the American Joint

Committee on Cancer which determines the degree of tumor development according

with the T, N and M categories. The T category informs about tumor growth, depth and

stage of adjacent tissues invasion. The N category informs about lymph nodes

metastasis and the M category describes the quantity of affected anatomical sites by

distant metastasis [3].

It´s estimated that the survival of patients diagnosed at the first stages of the

disease is up to 90%, in the other hand, 30 to 45% of the patients with late diagnosis

will present disease recurrence [4,5]. Factors such as low degree of differentiation and

high levels of serum carcinoembryonic antigen (CEA) are also described as important

for the increase of CRC recurrence after tumor resection [2,6], however, results from

recent studies have demonstrated that the analysis of micro-RNA expression in CRC

can be used as a more sensible prognostic biomarker due to it's potential of predicting

disease recurrence, survival and therapeutic response [7–12].

MicroRNAs (miRs) are small non-coding RNA molecules of approximately 21 to

24 nucleotides in length that modulate gene expression by connecting with the non

translational region 3 (3'UTR) of messenger RNAs resulting in it's translational

repression or cleavage [13]. In the literature, it's been already described more than 2000

miRs which demonstrate an essential role in the regulation of eukaryotic cells and in

the pathogenesis of several diseases, including cancer [14].

Biogenesis of miRs begins with the activity of RNA polymerase enzyme II which

transcribes the primary miRs (pri-miRs) recognized afterwards by DGCR8 protein

associated with DROSHA enzyme creating a complex which allows the cleavage of

the pri-miRs originating the precursor miRs (pre-miRs). The pre-miRs are carried to

the cell cytoplasm by Exportin-5 and cleaved by DICER, originating double strand miRs

which will be recognized by Argonaut proteins promoting the release of one strand of

the mature miR. Argonaut proteins in association with the mature miR and other

proteins will originate the RNA-Induced Silencing Complex (RISC) whose targets will

be specific mRNAs. [13,15,16]. Fig 1 outlines the miRs biogenesis.

Fig 1. **miRNAs biogenesis and proteins involved in the process.**

Recent studies report that the miRs regulate cellular growth and apoptosis,

therefore by silencing or upregulating of specific miRs can trigger the carcinogenic

process. [17,18]. MiRs are classified as oncogenic or tumoral suppressor when they inhibit

the expression of tumoral suppressor genes or oncogenes respectively. Oncogenic

miRs are usually upregulated in the neoplastic process while tumor supressor miRs

are usually downregulated in cancer [13,18,19]. Consequently, studies have demonstrated

by means of analysis of miRs expression, that they can be valuable diagnostic and

prognostic biomarkers in cancers, including the CRC. [7,20–22].

Despite the recent improvements in the diagnostic methods for CRC detection,

the overall survival for CRC patients is only 65% while the disease recurrence after

tumor resection is approximately 45%, emphasizing the importance of describing new

methods in order to guarantee a better prognosis for this patients [4,23].

There are innumerable studies investigating the significance of miRs expression

in CRC prognosis [7–10,12,21,22]. However, there is a lack of systematic reviews in the

literature integrating and identifying the main miRs significantly associated with the

prognostic aspects of CRC.

Given the relationship between the expression of miRs and the prognosis of

CRC, the present study aims to highlight such association by means of a systematic

review of the literature, in order to identify the main miRs associated with each

prognostic aspect of CRC, in order to contribute for the innovation of new prognostic

biomarkers and to help the development of new therapies based on the understanding

of the molecular biology of this disease.

Methods

Study design and search strategy

This study comprehends a systematic review of the literature. For the
elaboration of this project a reference review in the PubMed database, Cochrane
Library and Web of Science database was performed in order to identify the relevant
studies that evaluated miR expression associated with CRC prognosis. The time
period of the search was from February 2009, date from the last found and selected
study, to May 2018, date from the first found and selected study. The following search
strategy was used for the identification and selection of the articles: miRNA, colorectal
cancer, prognosis. The search was performed in November 2018 and it was limited to
English published articles. At the end of the search, all the results were compared and
reviewed for the inclusion of relevant publications about the subject.

Inclusion criteria

As a strategy of including relevant articles for the review, criteria related to study
classification and design were considered. The relevant criteria related to the
classification of the studies were: primary and descriptive studies, studies related to
CRC patients and studies completely available online. The relevant criteria related to
study design were: articles that evaluate CRC prognosis by overall survival and
disease free survival; articles that evaluate miR expression in tumoral tissues

compared to normal tissues; articles that evaluate miR expression by Real Time

Polymerase Chain Reaction (RT-PCR) using Taqman and/or SYBR Green; articles

that evaluate miR expression related to tumor growth/size, lymph nodes metastasis,

distant metastasis and TNM stage. As exclusion criteria, publications belonging to case

report, systematic reviews and meta-analysis categories were not included.

Data Extration

The data extraction was performed by means of a collection form pre-developed

by the authors in which the following data of each publication were collected: name of

the first author, year of publication, micro-RNA studied, sample group, origin of cases,

type of specimens analyzed, type of methodology used to evaluate the expression of

micro-RNAs, target genes and association with pathological and prognostic clinical

aspects.

This review followed the Preferred Reporting Items for Systematic Reviews and

Meta-Analyses (PRISMA) recommendations.

Quality assessment and publication bias

The present study comprehends a systematic review without meta-analysis,

therefore the results extracted from the selected articles were analyzed qualitatively,

thus the assessment of publication bias might not be practically applicable.

Problems with the design and execution of individual studies of healthcare

interventions raise questions about the validity of their findings, and therefore, the

quality of the primary studies were accessed by evaluating eight attributes extracted

from a checklist provided by the STROBE platform: description of the studying setting,

description of study participants, definitions of all variables, description of the data

sources and measurement tools, justification of sample size, description of statistical

methods, description of results and discussion and interpretation of results. After

evaluating the studies, each attribute were pointed in a range from 0 to 2 points (0 =

not described, 1 = poor described, 2 = well described), with maximum quality score of

16 points in total.

Data analysis

171 The data were reviewed and synthesized in semi-structured tables of

172 Microsoft® Office Excel version 2013. The results were analyzed by descriptive

173 statistics and qualitatively tabulated.

Results

Articles selection

After searching for the terms: microRNA, colorectal cancer and prognosis in all

three data bases, 924 articles were found. By reading the titles, 29 duplicates were

removed and 215 abstracts were selected for complete reading. After analyzing the

abstracts, 134 articles which evaluate miR expression associated with CRC prognosis

were selected for complete reading. After reading, 19 articles that did not achieve the

proposed inclusion criteria were excluded. At the end, 115 articles were included in the

systematic review. Fig. 2 outlines the study selection process.

Fig 2. Flow diagram outlining the study selection process.

The quality scores of the studies ranged from 14 to 16 with a mean quality rating of

15.7 (S2 File).

Characteristics of the selected studies

In total, 14.894 CRC patients were studied in the 115 included articles. The

number of patients varied from 10 to 765 patients per study. The studies were

performed in institutes and universities in Greece, South Korea, United States, Czech

Republic, Iran, Sweden, Denmark, Japan and China with most of the research

performed in the last two countries. The studies evaluated the expression of 102 miRs

associated with CRC prognostics aspects. MiR-21 was the most frequent one, being

analyzed in six studies [24–29]. MiR-181a [30–32] , miR-16 [33–35] and miR-133b [36–38] were

individually investigated in three studies. The miRs: miR-126 [9,39], miR-133a [40,41], miR-

182 [42,43], miR-183 [44,45], miR-199b [46,47], miR-22 [48,49], miR-224 [50,51], miR-24-3p [52,53], miR-

7 [54,55], were separately analyzed in two studies. Seven researches studied two miRs

concomitantly [29,33,37,38,56–58], while the others analyzed a single miR. All studies

analyzed miR expression performing RT-PCR by TaqMan and/or SYBR Green

methods in which 64 studies used SYBR Green while 51 studies used Taqman. The

characteristics of the selected studies in this review were summarized in the Table 1.

208 **Tabela 1.** Características dos estudos incluídos na revisão sistemática.

Ref.	Author / Year	Studied micro-RNA	Target Gene	Cases (n)	Cases Origin	Quantification Method	Results
(105)	Cui, 2018	miR-6852	TCF7	CRC (n=44) and normal adjacent tissue (n=44)	Harbin, China	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis and worse overall survival.
(83)	Tao, 2018	miR-1296	SFPQ	CRC (n= 80) and normal adjacent tissue (n= 80)	China and Japan	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis and worse overall survival
(63)	Gao, 2017	miR-888	Smad4	CRC (n= 126) and normal adjacent tissue (n= 126)	Jiangsu, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, distant metastasis, worse overall survival and worse disease free-survival.
(66)	Ma, 2018	miR-302c	AP-4	CRC (n= 90) and adjacent normal tissue (n=90)	Shaanxi, China	RT-qPCR TaqMan e SYBY Green	Downregulation was associated with advanced TNM stage, tumor depth, lymph node metastasis and worse overall survival.
(131)	Tsiakanikas, 2018	miR-28-5p	/	CRC (n= 182) and adjacent normal tissue (n= 86)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease-free survival.
(96)	Wang, 2018	miR-552	/	CRC (n= 183) and adjacent normal tissue (n= 83)	Shijiazhuan, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(44)	Chen, 2018	miR-183	/	CRC (n=42) and adjacent normal tissue (n=42)	Tianjin, China	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease-free survival.
(113)	Roh, 2018	miR-200c	/	CRC (n=109) and adjacent normal tissue (n=109)	Changwon, South Korea	RT-qPCR TaqMan	Upregulation was associated with lymph node metastasis, worse overall survival and worse disease-free survival.
(67)	Tong, 2018	miR-466	Cyclin D1, BAX e MMP-2	CRC (n=100) and adjacent normal tissue (n=100)	Zhejiang, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis, distant metastasis and worse overall survival.

(28)	Wu, 2017	miR-21	PTEN	CRC (n=105) and adjacent normal tissue (n=105)	Sichuan, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage and lymph node metastasis.
(84)	Chen, 2017	miR-130a	Forkhead box F2	CRC (n=53 and adjacent normal tissue (n=53)	Zhejiang, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage and lymph node metastasis.
(105)	Lin, 2017	miR-193a-3p	E-cadherin	CRC (n=90) and adjacent normal tissue (n=90)	Taizhou, China	RT-qPCR SYBR Green	Upregulation was associated with tumor groth and downregulation associated with worse overall survival.
(68)	Zhang, 2017	miR-340	RLIP76	CRC (n=91) and adjacent normal tissue (n=91)	Beijing, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage and lymph node metastasis.
(132)	Rapti, 2017	miR-34a	/	CRC (n=133) and adjacent normal tissue (n=61)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease-free survival.
(122)	Zhou, 2017	miR-650	AKT2/GSK3β/ E-cadherin pathway	CRC patients with better prognosis (n=48) and CRC patients with worse prognosis (n=40)	Shanghai, China	RT-qPCR Taqman	Downregulation was associated with worse overall survival.
(69)	Stiegelbauer, 2017	miR-196b-5p	HOXB7 e GALNT5	Two cohorts: CRC (n=110) and CRC (n=182)	Graz, Austria e Czech Republic	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, worse overall survival and worse disease-free survival.
(70)	Wen, 2017	miR-944	MACC1	CRC (n=86) and adjacent normal tissue (n=86)	Guangzhou, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(71)	Yan, 2017	miR-30d	LRH-1	CRC (n=80) and adjacent normal tissue (n=80)	Xi'an, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(49)	Xia, 2017	miR-22	Sp1 (e PTEN/AKT)	CRC (n=118) and adjacent normal tissue (n=118)	Sichuan, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.

(133)	Kontos, 2017	miR-15a-5p	RECK	CRC (n=182) and adjacent normal tissue (n=86)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease free-survival.
(72)	He, 2017	miR-296	S100A4	CRC (n=90) and adjacent normal tissue (n=90)	Beijing, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(134)	Yamazaki, 2017	miR-181c	/	Two cohorts of CRC patients stage II: CRC (n= 80) and adjacent normal tissue (n=80) and CRC (n=60) and adjacent normal tissue (n=60)	Tokyo, Japan	RT-qPCR TaqMan	Upregulation was associated with worse overall survival and worse disease-free survival.
(52)	kerimis, 2017	miR-24-3p	/	CRC (n=182) and adjacent normal tissue (n=86)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse diasease-free survival.
(34)	Diamantopoulos, 2017	miR-16	/	CRC (n=182) and adjacent normal tissue (n=86)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse diasease-free survival.
(54)	Nagano, 2016	miR-7	/	CRC (n=196) and adjacent normal tissue ((n=14)	Mie, Japan	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(118)	Yu, 2016	miR-140-5p	ADAMTS5, IGFBP5	CRC (n=60) and adjacent normal tissue (n=60)	Dalian, China	RT-qPCR TaqMan	Downregulation was associated iwth tumoral stage and distant metastasis.
(85)	Rapti, 2016	miR-96	/	CRC (n=108) and adjacent normal tissue (n=54)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, lymph node metastasis, worse overall survival and worse disease-free survival.
(73)	Wang, 2016	miR-384	KRAS, CDC42	CRC (n=100) and adjacent normal tissue (n=100)	Guangdong, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage.
(46)	Cristóbal, 2017	miR-199b	PP2A	CRC (n=97)	Madrid, Spain	RT-qPCR TaqMan	Downregulation was associated with lymph node metastasis, worse overall survival and worse disease-free survival.

(114)	Liu , 2016	miR-1260b	/	CRC (n=120) and adjacent normal tissue (n=120)	Gansu, China	RT-qPCR SYBR Green	Upregulation was associated with lymph node metastasis, worse overall survival and worse disease-free survival.
(123)	Wang, 2016	miR-187	CD276	CRC (n=32) and adjacent normal tissue (n=32) CRC (n=80) and adjacent normal tissue (n=80)	Shanghai, China	RT-qPCR SYBR Green	Downregulation was associated with worse overall survival and worse disease free-survival.
(62)	Cheng, 2016	miR-20a-5p	Smad4	CRC (n=544) and adjacent normal tissue (n=544)	Shanghai, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, distant metastasis, worse overall survival and worse disease-free survival.
(97)	Liu, 2016	let-7a-5p	HMGA2	CRC (n=192) and adjacent normal tissue (n=192)	Taipei, Taiwan	RT-qPCR TaqMan	Downregulation was associated with tumor growth, lymph node metastasis, worse overall survival and worse diasease-free survival.
(47)	Shen, 2016	miR-199b	SIRT1; CREB/KISS1	CRC (n=60) and adjacent normal tissue (n=60)	Beijing, China	RT-qPCR SYBR Green	Downregulation associated with advanced TNM stage, distant metastasis and worse overall survival.
(86)	Ma, 2016	miR-517a	FOXJ3	CRC (n=90) and adjacent normal tissue (n=90)	Xi'an, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(135)	Noguchi, 2016	miR-503	CaSR	CRC (n=20), adenomas (n=20) and normal tissue (n=20)	Tsu , Japan	RT-qPCR TaqMan	Upregulation was associated with worse overall survival and worse disease-free survival.
(24)	Mima, 2016	miR-21	/	CRC (n=765)	Boston, MA, USA	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease-free survival.
(98)	Mokutani, 2016	miR-132	ANO1	CRC (n=21 and adjacent normal tissue (n=21)	Osaka, Japan	RT-qPCR TaqMan	Downregulation was associated with tumor growth, lymph node metastasis, worse overall survival and worse disease-free survival.

(32)	Li, 2016	miR-181A	PTEN/AKT	CRC (n=72) and adjacent normal tissue (n=69)	Shandong, China	RT-qPCR SYBR Green	Upregulation in hepatic metastasis patients was associated with worse overall survival.
(102)	Bin, 2016	miR-30c	/	CCR (n=192) and adjacent normal tissue (n=192)	Baoding, China	RT-qPCR SYBR Green	Downregulation was associated with tumor growth, lymph node metastasis and worse overall survival
(106)	Wang, 2016	miR-497	KSR1	CRC (n=62) and adjacent normal tissue (n=62)	Nanjing, China	RT-qPCR TaqMan	Downregulation was associated with lymph node metastasis and advanced CRC stages.
(99)	Sun, 2015	miR-206	/	CRC (n=80) and adjacent normal tissue (n=80)	Jilin, China	RT-qPCR SYBR Green	Downregulation was associated with tumor growth, lymph node metastasis and worse overall survival.
(124)	Hibino, 2015	miR-148a	MMP7	CRC (n=159) and adjacent normal tissue (n=159)	Hiroshima , Japan	RT-qPCR TaqMan	Downregulation was associated with advanced CRC stages and worse overall survival in CRC patients stage III.
(74)	Wu, 2015	miR-128	IRS1	CRC (n=45) and adjacent normal tissue (n=45)	Changchun, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage and lymph node metastasis.
(75)	Liao , 2015	miR-33b	/	CRC (n=60) and adjacent normal tissue (n=60)	Shanghai, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, tumor growth and worse overall survival.
(76)	Qu, 2015	miR-155-5p	/	CRC (n=372) and adjacent normal tissue (n=372)	Xinjiang, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage and distant metastasis.
(119)	Li, 2015	miR-99b-5p	mTOR	CRC (n=56) and adjacent normal tissue (n=56)	Shanghai, China	RT-qPCR TaqMan	Downregulation was associated with distant metastasis and worse overall survival.
(87)	Zhang, 2015	miR-106b	DLC1	CRC (n=93) and adjacent normal tissue (n=93)	Sichuan, People's Republic of China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, lymph node metastasis, worse overall survival and worse disease-free survival.
(136)	Sümbül, 2015	miR-211	/	CRC (n=65), and adjacent normal tissue (n=65) and normal tissue (n=65)	Hatay, Turkey	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival.

(77)	Kai, 2015	miR-154	/	CRC (n=169) and adjacent normal tissue (n=169)	Wenzhou, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis, distant metastasis and worse overall survival.
(88)	Perez-Carbonell, 2015	miR-320e	/	CRC (n=100)	Barcelona, Spain	RT-qPCR TaqMan	Up regulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(125)	Wang, 2015	miR-217	AEG-1	CRC (n=50)	Beijing, P.R. China	RT-qPCR SYBR Green	Downregulation was associated with worse overall survival.
(89)	Liu, 2015	miR-592	/	CRC (n=89) and adjacent normal tissue (n=89)	Shanghai, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, tumor growth, distant metastasis and worse overall survival.
(78)	Xu, 2015	miR-149	FOXM1	CRC (n=78) and adjacent normal tissue (n=20)	Jiangsu, PR China	RT-qPCR TaqMan	Down regulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.
(79)	Wang, 2015	miR-194	MAP4K4/c-Jun/MDM2	CRC (n=50)	Beijing, PR China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, tumor growth and worse overall survival.
(56)	Ge, 2014	miR-196a, miR-196b	/	CRC (n=126) and adjacent normal tissue (n=126)	Hunan, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis, worse overall survival and worse disease-free survival.
(55)	Suto, 2014	miR-7	EGFR	CRC (n=105) and adjacent normal tissue (n=105)	Maebashi, Japan	RT-qPCR TaqMan	Down regulation was associated with worse overall survival.
(53)	Gao, 2015	miR-24-3p	/	CRC (n=95) and adjacent normal tissue (n=95)	Shandong, China	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis and worse overall survival.
(36)	Guo, 2015	miR-133b	CTGF	CRC (n=71)	Hunan, China	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis and advanced TNM stage.

(126)	Inoue, 2015	miR-29b	MCL1 e CDK6	CRC (n=42) and adjacent normal tissue (n=42)	Osaka, Japan	RT-qPCR TaqMan	Downregulation was associated with worse overall survival and worse disease-free survival.
(33)	Xiao, 2014	miR-15a, miR-16	/	CRC (n=126) and adjacent normal tissue (n=126)	Hunan, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, lymph node metastasis, worse overall survival and worse disease-free survival.
(115)	Mo, 2014	miR-376a	/	CRC (n=53) and adjacent normal tissue (n=53)	Changchun, China	RT-qPCR SYBR Green	Upregulation was associated with lymph node metastasis and worse overall survival.
(25)	Fukushima, 2014	miR-21	/	CRC (n=306) and adjacent normal tissue (n=306)	Tokyo, Japan	RT-qPCR TaqMan	Upregulation was associated with tumor growth, lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(50)	Adamopoulos, 2015	miR-224	/	CRC (n=115) and adjacent normal tissue (n=66)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with worse overall survival and worse disease-free survival.
(64)	Gao, 2015	miR-34a-5p	p53	CRC (n=10) and adjacent normal tissue (n=10)	Beijing, China	RT-qPCR TaqMan	Downregulation was associated with worse overall survival.
(127)	Yin, 2014	miR-204-5p	RAB22A	CRC (n=272) and adjacent normal tissue (n=272)	Shanghai, China	RT-qPCR SYBR Green	Downregulation was associated with worse overall survival.
(90)	Wang, 2014	miR-720	/	CRC (n=96) and adjacent normal tissue (n=96)	Suzhou, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis, distant metastasis and worse overall survival.
(60)	Sun, 2014	miR-494	PTEN	CRC (n=247) and adjacent normal tissue (n=247)	Shaanxi, china	RT-qPCR TaqMan	Upregulation was associated with lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(137)	Li, 2014	miR-223	/	CRC (n=62) and adjacent normal tissue (n=62)	Jinan, China	RT-qPCR	Upregulation was associated with worse disease-free survival.
(120)	Ress, 2015	MiR-96-5p	KRAS	CRC (n=80)	Graz, Austria	RT-qPCR SYBR Green	Downregulation was associated with distant metastasis and worse overall survival.

(107)	Zhang, 2014	miR-193a-5p	/	CRC (n=69) and adjacent normal tissue (n=69)	Beijing, China	RT-qPCR TaqMan	Downregulation was associated with lymph node metastasis, worse overall survival and worse disease-free survival.
(80)	Chen, 2014	miR-100	/	CRC (n=138) and adjacent normal tissue (n=138)	Zaozhuang, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis and worse overall survival.
(40)	Wang, 2014	miR-133a	/	CRC (n=169) and adjacent normal tissue (n=169)	Shandong, China	RT-qPCR	Downregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(128)	Song, 2014	miR-139-5p	AMFR and NOTCH1	CRC (n=158) and adjacent normal tissue (n=158)	Shanghai, China	RT-qPCR SYBR Green	Downregulation was associated with worse overall survival.
(41)	Wan, 2014	miR-133a	/	CRC (n=125) and adjacent normal tissue (n=125)	Hong Kong, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, distant metastasis and worse overall survival.
(129)	Jinushi, 2014	miR-124-5p	SMC4	CRC (n=71)	Sapporo, Japan	RT-qPCR SYBR Green e TaqMan	Downregulation was associated with worse overall survival.
(37)	Ellermeier, 2014	miR-133b e miR-210	/	CRC (n=19 and adjacent normal tissue (n=19)	Providence, RI, U.S.A	RT-qPCR TaqMan	Upregulation of both was associated with worse overall survival. Upregulation of miR-210 and downregulation of miR-133b were associated with distant metastasis.
(91)	Chu, 2014	miR-630	/	CRC (n=206) and adjacent normal tissue (n=206)	Xi'na, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.
(108)	Orang, 2014	miR-205	/	CRC (n=40) and adjacent normal tissue (n=40)	Tabriz, Iran	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis.
(59)	Fang, 2014	miR-17-5p	PTEN	CRC (n=295)	Guangzhou, China	RT-qPCR SYBR Green	Upregulation was associated with distant metastasis and worse overall survival.
(81)	Sun, 2014	miR-335	ZEB2	CRC (n=80) and adjacent normal tissue (n=80)	Xi'na, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.

(30)	Ji, 2014	miR-181a	WIF-1	CRC (n=137) and adjacent normal tissue (n=137)	Beijing, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.
(61)	Zhang, 2014	miR-20a	SMAD4	CRC (n=86) and adjacent normal tissue (n=86)	Nanchong, China	RT-qPCR SYBR Green	Upregulation was associated with lymph node metastasis and worse overall survival.
(92)	Qu, 2014	miR-210	VMP1	CRC (n=193) and adjacent normal tissue (n=193)	Shandong, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis and worse overall survival.
(42)	Rapti, 2014	miR-182	/	CRC (n=116) and adjacent normal tissue (n=60)	Athens, Greece	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(100)	Zhang, 2014	miR-378	Vimentin	CRC (n=84) and adjacent normal tissue (n=84)	Sichuan, China	RT-qPCR SYBR Green	Downregulation was associated with tumor growth, lymph node metastasis and worse overall survival.
(39)	Liu, 2014	miR-126	/	CRC (n=92) and adjacent normal tissue (n=92)	Zhanjiang, China	RT-qPCR SYBR Green	Downregulation was associated with distant metastasis and worse overall survival.
(57)	Li, 2014	miR-378a-3p, miR-378a-5p	/	Two cohorts: CRC (n=96) and adjacent normal tissue (n=96); CRC (n=12) and adjacent normal tissue (n=12)	Guangzhou, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, tumor growth and worse overall survival.
(82)	Yu, 2013	miR-218	/	CRC (n=26) and adjacent normal tissue (n=26)	Jiangsu, China	RT-qPCR SYBR Green	Downregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(93)	Li, 2014	miR-25	/	CRC (n=186) and adjacent normal tissue (n=186)	Shaanxi, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.
(94)	Qin, 2014	miR-191	TIMP3	CRC (n=136)	Xinxiang, China	RT-qPCR TaqMan	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.

(109)	Long, 2013	miR-138	TWIST2	CRC (n=187) and adjacent normal tissue (n=187)	Hunan,China	RT-qPCR TaqMan	Downregulation was associated with lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(45)	Zhou, 2014	miR-183	/	CRC (n=94) and adjacent normal tissue (n=94)	Sichuan, China	RT-qPCR SYBR Green	Upregulation was associated with lymph node metastasis, distant metastasis and worse overall survival.
(35)	Qian, 2013	miR-16	/	CRC (n=143) and adjacent normal tissue (n=18)	Jiangsu, China	RT-qPCR TaqMan	Downregulation was associated with advanced TNM stage, lymph node metastasis and worse overall survival.
(9)	Hansen, 2013	miR-126	/	CRC (n=126)	Odense, Denmark/ Lund, Sweden	RT-qPCR SYBR Green	Downregulation was associated with worse overall survival.
(110)	Lou, 2013	miR-625	/	CRC (n=96) and adjacent normal tissue (n=96)	Shanghai, China	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis, distant metastasis and worse overall survival.
(51)	Liao, 2013	miR-224	PHLPP1, PHLPP2	CRC (n=43) and adjacent normal tissue (n=43)	Guangzhou, China	RT-qPCR TaqMan	Upregulation was associated with distant metastasis and worse overall survival.
(26)	Chen, 2013	miR-21	/	CRC (n=195)	Changhua, Taiwan	RT-qPCR TaqMan	Upregulation was associated with worse overall survival.
(27)	Toiyama, 2013	miR-21	/	CRC (n=166) and adjacent normal tissue (n=166)	Japan	RT-qPCR TaqMan	Upregulation was associated with tumor growth, distant metastasis and worse overall survival.
(109)	Meng, 2013	miR-212	MnSOD	CRC (n=30) and adjacent normal tissue (n=30)	Guangzhou, China	RT-qPCR TaqMan	Downregulation was associated with lymph node metastasis, distant metastasis, worse overall survival and worse disease-free survival.
(43)	Liu, 2013	miR-182	/	CRC (n=152) and adjacent normal tissue (n=152)	Jinan, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, tumor growth, lymph node metastasis and worse overall survival.
(116)	Wan, 2013	miR-199a-3p	/	CRC (n=92) and adjacent normal tissue (n=92)	Shanghai, China	RT-qPCR SYBR Green	Upregulation was associated with lymph node metastasis, distant metastasis and worse overall survival.

(104)	Li, 2012	miR-429	SOX2	CRC (n=107) and adjacent normal tissue (n=107)	Shandong, China	RT-qPCR	Upregulation was associated with tumor growth, lymph node metastasis and worse overall survival.
(31)	Nishimura, 2012	miR-181a	PTEN	CRC (n=162)	Beppu, Japan	RT-qPCR TaqMan	Upregulation was associated with worse overall survival and worse disease-free survival.
(130)	Wang, 2013	miR-124	/	CRC (n=96) and adjacent normal tissue (n=96)	Chengdu City, China	RT-qPCR TaqMan	Downregulation was associated with worse overall survival and worse disease-free survival.
(95)	Zhou, 2012	miR-92a	/	CRC (n=82) and adjacent normal tissue (n=82)	Nanchong, China	RT-qPCR SYBR Green	Upregulation was associated with advanced TNM stage, lymph node metastasis, distant metastasis and worse overall survival.
(48)	Zhang, 2012	miR-22	/	CRC (n=86) and adjacent normal tissue (n=86)	Nanchong, China	RT-qPCR SYBR Green	Downregulation was associated with distant metastasis and worse overall survival.
(121)	Yamashita, 2012	miR-372	/	CRC (n=144)	Beppu, Japan	RT-qPCR TaqMan	Upregulation was associated with distant metastasis and worse overall survival.
(117)	Nishida, 2012	miR-10b	BIM	CRC (n=88)	Osaka, Japan	RT-qPCR TaqMan	Upregulation was associated with lymph node metastasis and worse overall survival.
(101)	Song, 2011	miR-148b	CCK2R	CRC (n=96) and adjacent normal tissue (n=96)	Shenyang City, China	RT-qPCR SYBR Green	Downregulation was associated with tumor growth.
(38)	Akçakaya, 2011	miR-185, miR-133b	/	CRC (n=50)	Stockholm, Sweden	RT-qPCR SYBR Green e TaqMan	Upregulation of miR-185 and downregulation of miR-133b were associated with metastasis and worse overall survival.
(29)	Shibuya, 2011	miR-21 e miR-155	PDCD4 and TP53INP1	CRC (n=156) and adjacent normal tissue (n=156)	Tokyo, Japan	RT-qPCR TaqMan	Upregulation of miR-21 was associated with distant metastasis, worse overall survival and worse disease-free survival. Upregulation of miR-155 as associated with lymph node metastasis, worse overall survival and worse disease-free survival.

(65)	Nishida, 2011	miR-125b	p53 e p21	CRC (n=89)	Osaka, Japan	RT-qPCR TaqMan	Upregulation was associated with tumor growth and worse overall survival.
(112)	Wang, 2012	miR-195	/	CRC (n=85) and adjacent normal tissue (n=85)	Nanjing, China	RT-qPCR SYBR Green	Downregulation was associated with lymph node metastasis and worse overall survival.
(58)	Wang, 2009	miR-31, miR-143, miR-145	/	CRC (n=98) and adjacent normal tissue (n=98)	Chengdu, China	RT-qPCR TaqMan	Upregulation of miR-31 was associated with advanced TNM stage.

209 miR: micro-RNA. CRC: colorectal cancer. RT-qPCR: real-time polymerase chain reaction

221   Studies that investigated the prognosis of CRC patients by means of overall

222 survival and disease-free survival demonstrated that upregulated miRs associated with

223 worse prognosis were more frequent than downregulated miRs. In terms of pathologic

224 clinical aspects, downregulated miRs associated with advanced TNM stage, tumor

225 growth and lymph node metastasis were more frequent than upregulated miRs.

226 Nonetheless, upregulated miRs were more associated with distant metastasis. The

227 association of the miRs differentially expressed in CRC and their respective prognostic

228 aspects are summarized in Table 2.

229 **Tabela 2**. Micro-RNAs diferentially expressed in CRC and associated prognostic aspects.

Prognostic Aspects	Downregulated micro-RNAs	Upregulated micro-RNAs
Advanced TNM stage	miR-302c [66], miR-466 [67], miR-340 [68], miR-196b-5p [69], miR-944 [70], miR-30d [71], miR-22 [49], miR-296 [72], miR-384 [73], miR-199b [47], miR-128 [74], miR-33b [75], miR-155-5p [76], miR-154 [77], miR-149 [78], miR-194 [79], miR-133b [36], miR-15a [33], miR-100 [80], miR-133a [40], miR-335 [81], miR-378a-3p [57], miR-378a-5p [57], miR-218 [82], miR-16 [33]	miR-182 [42,43], miR-1296 [83], miR-888 [63], miR-21 [28], miR-130a [84], miR-7 [54], miR-96 [85], miR-20a-5p [62], miR-517a [86], miR-106b [87], miR-320e [88], miR-592 [89], miR-196a [56], miR-196b [56], miR-720 [90], miR-133a [41], miR-630 [91], miR-181a [30], miR-210 [92], miR-25 [93], miR-191 [94], miR-92a [95], miR-31[58], miR-552 [96]
Tumor Growth	miR-302c [66], miR-466 [67], let-7a-5p [97], miR-132 [98], miR-206 [99], miR-33b [75], miR-154 [77], miR-194 [79], miR-100 [80], miR-378 [100], miR-378a-3p [57], miR-378a-5p [57], miR-148b [101], miR-30c [102]	miR-1296 [83], miR-193a-3p [103], miR-592 [89], miR-720 [90], miR-210 [92], miR-182 [43], miR-429 [104], miR-125b [65], miR-21 [25,27]
Lymph Node Metastasis	miR-6852 [105], miR-302c [66], miR-466 [67], miR-340 [68], miR-944 [70], miR-30d [71], miR-22 [49], miR-296 [72], miR-199b [46], let-7a-5p [97], miR-132 [98], miR-497 [106], miR-206 [99], miR-128 [74], miR-154 [77], miR-149 [78], miR-24-3p [53], miR-15a [33], miR-16 [33,35], miR-193a-5p [107], miR-100 [80], miR-133a [40], miR-205 [108], miR-335 [81], miR-378 [100], miR-218 [82], miR-138 [109], miR-625 [110], miR-212 [111], miR-195 [112], miR-133b [36], miR-30c [102]	miR-1296 [83], miR-200c [113], miR-21 [25,28], miR-130a [84], miR-7 [54], miR-96 [85], miR-1260b [114], miR-517a [86], miR-106b [87], miR-320e [88], miR-196a [56], miR-196b [56], miR-376a [115], miR-720 [90], miR-494 [60], miR-630 [91], miR-181a [30], miR-20a [61], miR-210 [92], miR-182 [42,43], miR-25 [93], miR-191 [94], miR-183 [45], miR-199a-3p [116], miR-429 [104], miR-92a [95], miR-10b [117], miR-155 [29], miR-552 [96]
Distant Metastasis	miR-466 [67], miR-944 [70], miR-30d [71], miR-22 [48,49], miR-296 [72], miR-140-5p [118], miR-199b [47], miR-155-5p [76], miR-99b-5p [119], miR-154 [77], miR-149 [78], miR-96-5p [120], miR-133b [37,38], miR-126 [39], miR-138 [109], miR-625 [110], miR-212 [111]	miR-888 [63], miR-20a-5p [62], miR-517a [86], miR-320e [88], miR-592 [89], miR-720 [90], miR-494 [60], miR-133a [41], miR-210 [37], miR-630 [91], miR-17-5p [59], miR-181a[30], miR-25 [93], miR-191 [94], miR-183 [45], miR-224 [51], miR-199a-3p [116], miR-92a [95], miR-372 [121], miR-185 [38], miR-21 [25,27,29]
Worse Overall Survival	miR-199b [46,47], miR-6852 [105], miR-302c [66], miR-466 [67], miR-650 [122], miR-196b-5p [69], miR-944 [70], miR-30d [71], miR-296 [72], miR-187 [123], let-7a-5p [97], miR-132 [98], miR-206 [99], miR-148a [124], miR-33b [75], miR-99b-5p [119], miR-154 [77], miR-217 [125], miR-149 [78], miR-194 [79], miR-7 [55], miR-24-3p [53], miR-29b [126], miR-15a [33], miR-16 [33,35], miR-204-5p [127], miR-96-5p [120], miR-193a-5p [107], miR-100 [80], miR-133a [40], miR-139-5p [128], miR-124-5p [129], miR-335 [81], miR-378 [100], miR-126 [39], miR-378a-3p [57], miR-378a-5p [57], miR-218 [82], miR-138 [109], miR-625 [110], miR-212 [111], miR-124 [130], miR-22 [48], miR-133b [38], miR-193a-3p [103], miR-195 [112], miR-30c [102]	miR-181a [30–32], miR-182 [42,43], miR-1296 [83], miR-888 [63], miR-28-5p [131], miR-183 [44,45], miR-200c [113], miR-34a [132], miR-15a-5p [133], miR-181c [134], miR-24-3p [52], miR-16 [34], miR-7 [54], miR-96 [85], miR-1260b [114], miR-20a-5p [62], miR-517a [86], miR-503 [135], miR-106b [87], miR-211 [136], miR-320e [88], miR-592 [89], miR-196a [56], miR-196b [56], miR-376a [115], miR-224 [50,51], miR-720 [90], miR-494 [60], miR-223 [137], miR-133a [41], miR-133b [37], miR-210 [37,92], miR-630 [91], miR-17-5p [59], miR-20a [61], miR-25 [93], miR-191 [94], miR-183 [44,45], miR-199a-3p [116], miR-429 [104], miR-92a [95], miR-372 [121], miR-10b [117], miR-185 [38], miR-21 [24–27,29], miR-155 [29], miR-125b [65], miR-552 [96]
Worse Disease-Free Survival	miR-196b-5p [69], miR-944 [70], miR-30d [71], miR-22 [49], miR-296 [72], miR-199b [46], miR-187 [123], let-7a-5p [97], miR-132 [98], miR-29b [126], miR-15a [33], miR-16 [33], miR-34a-5p [64], miR-193a-5p [107], miR-138 [109], miR-126 [9], miR-212 [111], miR-124 [130]	miR-888 [63], miR-28-5p [131], miR-183 [44], miR-200c [113], miR-34a [132], miR-15a-5p [133], miR-181c [134], miR-24-3p [52], miR-16 [34], miR-96 [85], miR-1260b [114], miR-20a-5p [62], miR-517a [86], miR-503 [135], miR-106b [87], miR-320e [88], miR-196a [56], miR-196b [56], miR-224 [50], miR-494 [60], miR-181a [31], miR-21 [24,25,29], miR-155 [29]

Some of the selected studies investigated the target genes of the miRs associated with the prognostic aspects of CRC. In total, 65 target genes were analyzed in which the most frequent ones were: PTEN, investigated in six studies [28,31,32,49,59,60], SMAD4, investigated in three studies [61–63], and TP53 [64,65] investigated in two studies.

Micro-RNAs and the prognosis of colorectal cancer

Micro-RNAs associated with advanced TNM stage

Twenty-four studies demonstrated the downregulation of twenty-five miRs associated with advanced TNM stage. Such marks were: miR-302c [66], miR-466 [67], miR-340 [68], miR-196b-5p [69], miR-944 [70], miR-30d [71], miR-22 [49], miR-296 [72], miR-384 [73], miR-199b [47], miR-128 [74], miR-33b [75], miR-155-5p [76], miR-154 [77], miR-149 [78], miR-194 [79], miR-133b [36], miR-15a [33], miR-100 [80], miR-133a [40], miR-335 [81], miR-378a-3p [57], miR-378a-5p [57], miR-218 [82] and miR-16 [33]. In the other hand, two studies demonstrated the upregulation of miR-182 [42,43] associated with advanced TNM stage. Futhermore, others 22 studies pointed 23 different upregulated miRs associated with advanced TNM stage. Those markers were: miR-1296 [83], miR-888 [63], miR-21 [28], miR-130a [84], miR-7 [54], miR-96 [85], miR-20a-5p [62], miR-517a [86], miR-106b [87], miR-320e [88], miR-592 [89], miR-196a [56], miR-196b [56], miR-720 [90], miR-133a [41], miR-630 [91], miR-181a [30], miR-210 [92], miR-25 [93], miR-191 [94], miR-92a [95], miR-31[58] and miR-552 [96].

Micro-RNAs associated with tumor growth

Tumor growth was associated with the downregulation of 14 miRs demonstrated in

13 studies. Those markers were: miR-302c [66], miR-466 [67], let-7a-5p [97], miR-132 [98],

miR-206 [99], miR-33b [75], miR-154 [77], miR-194 [79], miR-100 [80], miR-378 [100], miR-378a-

3p [57], miR-378a-5p [57], miR-148b [101] and miR-30c [102]. However, ten studies verified the

upregulation of miR-1296 [83], miR-193a-3p [103], miR-592 [89], miR-720 [90], miR-210 [92],

miR-182 [43], miR-429 [104], miR-125b [65] and miR-21 [25,27] associated with tumor growth as

well.

Micro-RNAs associated with the presence of lymph node metastasis

A total of 32 studies demonstrated the association between downregulation of 32

miRs and the presence of lymph node metastasis such as miR-6852 [105], miR-302c [66],

miR-466 [67], miR-340 [68], miR-944 [70], miR-30d [71], miR-22 [49], miR-296 [72], miR-199b [46],

let-7a-5p [97], miR-132 [98], miR-497 [106], miR-206 [99], miR-128 [74], miR-154 [77], miR-149 [78],

miR-24-3p [53], miR-15a [33], miR-16 [33,35], miR-193a-5p [107], miR-100 [80], miR-133a [40], miR-

205 [108], miR-335 [81], miR-378 [100], miR-218 [82], miR-138 [109], miR-625 [110], miR-212 [111],

miR-195 [112], miR-133b [36] and miR-30c [102]. In contrast, 30 studies verified the

upregulation of miR-1296 [83], miR-200c [113], miR-21 [25,28], miR-130a [84], miR-7 [54], miR-96

[85], miR-1260b [114], miR-517a [86], miR-106b [87], miR-320e [88], miR-196a [56], miR-196b [56],

miR-376a [115], miR-720 [90], miR-494 [60], miR-630 [91], miR-181a [30], miR-20a [61], miR-210

[92], miR-182 [42,43], miR-25 [93], miR-191 [94], miR-183 [45], miR-199a-3p [116], miR-429 [104],

miR-92a [95], miR-10b [117], miR-155 [29] and miR-552 [96] associated with the presence of

lymph node metastasis.

Micro-RNAs associated with the presence of distant metastasis

Downregulation of miR-466 [67], miR-944 [70], miR-30d [71], miR-22 [48,49], miR-296 [72],

miR-140-5p [118], miR-199b [47], miR-155-5p [76], miR-99b-5p [119], miR-154 [77], miR-149 [78],

miR-96-5p [120], miR-133b [37,38], miR-126 [39], miR-138 [109], miR-625 [110] and miR-212 [111]

was associated with the presence of distant metastasis in 19 studies. In contrast,

downregulation of miR-888 [63], miR-20a-5p [62], miR-517a [86], miR-320e [88], miR-592 [89],

miR-720 [90], miR-494 [60], miR-133a [41], miR-210 [37], miR-630 [91], miR-17-5p [59], miR-

181a[30], miR-25 [93], miR-191 [94], miR-183 [45], miR-224 [51], miR-199a-3p [116], miR-92a [95],

miR-372 [121], miR-185 [38] and miR-21 [25,27,29] was associated with the presence of distant

metastasis in 23 studies.

Micro-RNAs associated with worse overall survival

Among de evaluated studies, 49 demonstrated the downregulation of miR-199b

[46,47], miR-6852 [105], miR-302c [66], miR-466 [67], miR-650 [122], miR-196b-5p [69], miR-944 [70],

miR-30d [71], miR-296 [72], miR-187 [123], let-7a-5p [97], miR-132 [98], miR-206 [99], miR-148a

[124], miR-33b [75], miR-99b-5p [119], miR-154 [77], miR-217 [125], miR-149 [78], miR-194 [79], miR-7

[55], miR-24-3p [53], miR-29b [126], miR-15a [33], miR-16 [33,35], miR-204-5p [127], miR-96-5p [120],

miR-193a-5p [107], miR-100 [80], miR-133a [40], miR-139-5p [128], miR-124-5p [129], miR-335

[81], miR-378 [100], miR-126 [39], miR-378a-3p [57], miR-378a-5p [57], miR-218 [82], miR-138 [109],

miR-625 [110], miR-212 [111], miR-124 [130], miR-22 [48], miR-133b [38], miR-193a-3p [103], miR-

195 [112] and miR-30c [102] associated with worse prognosis (lower overall survival) of

CRC patients. Conversely, 56 studies identified the upregulation of miR-181a [30–32],

miR-182 [42,43], miR-1296 [83], miR-888 [63], miR-28-5p [131], miR-183 [44,45], miR-200c [113],

miR-34a [132], miR-15a-5p [133], miR-181c [134], miR-24-3p [52], miR-16 [34], miR-7 [54], miR-96

[85], miR-1260b [114], miR-20a-5p [62], miR-517a [86], miR-503 [135], miR-106b [87], miR-211 [136],

miR-320e [88], miR-592 [89], miR-196a [56], miR-196b [56], miR-376a [115], miR-224 [50,51], miR-

720 [90], miR-494 [60], miR-223 [137], miR-133a [41], miR-133b [37], miR-210 [37,92], miR-630 [91],

miR-17-5p [59], miR-20a [61], miR-25 [93], miR-191 [94], miR-183 [44,45], miR-199a-3p [116], miR-

429 [104], miR-92a [95], miR-372 [121], miR-10b [117], miR-185 [38], miR-21 [24–27,29], miR-155 [29],

miR-125b [65] and miR-552 [96] was also associated with worse overall survival of CRC

patients.

Micro-RNAs associated with worse disease free-survival

In total, 17 studies demonstrated the downregulation of eighteen miRs associated

with worse disease free-survival such as miR-196b-5p [69], miR-944 [70], miR-30d [71], miR-

22 [49], miR-296 [72], miR-199b [46], miR-187 [123], let-7a-5p [97], miR-132 [98], miR-29b [126], miR-

15a [33], miR-16 [33], miR-34a-5p [64], miR-193a-5p [107], miR-138 [109], miR-126 [9], miR-212

[111] and miR-124 [130]. However, 21 studies associated the upregulation of miR-888 [63],

miR-28-5p [131], miR-183 [44], miR-200c [113], miR-34a [132], miR-15a-5p [133], miR-181c [134],

miR-24-3p [52], miR-16 [34], miR-96 [85], miR-1260b [114], miR-20a-5p [62], miR-517a [86], miR-

503 [135], miR-106b [87], miR-320e [88], miR-196a [56], miR-196b [56], miR-224 [50], miR-494 [60],

miR-181a [31], miR-21 [24,25,29] and miR-155 [29] with worse disease free-survival.

Target genes

In addition to demonstrating the association of miRs and the clinical and prognostic aspects of colorectal cancer, 60 studies also investigated the possible target genes for these miRs. 65 target gens were identified, however PTEN, SMAD4, P53 and KRAS were the most frequent ones. The downregulation of PTEN was associated with de upregulation of miR-181a [31,32], miR-17-5p [59], miR-494 [60], miR-22 [49] and miR-21 [28]. The downregulation of SMAD4 was associated with de upregulation of miR-20a [61], miR-20a-5p [62], miR-888 [63]. The downregulation of P53 was related to the upregulation of miR-125b [65] and to the downregulation of de miR-34a-5p [64]. The upregulation of KRAS was associated with the downregulation of miR-96-5p [120] and miR-384 [73].

Discussion

Colorectal cancer is one of the most prevalent neoplasms in the world and it's estimated that the five-year overall survival for CRC patients is only 65%[23]. Several factors associated with CRC prognosis have been already studied, however sensible and specific prognostic biomarkers capable of predicting survival and recurrence of the disease are still unknown [138].

Recent reviews have demonstrated strong correlations between the prognostic aspects of breast cancer [139] and leukemia [140] with miR expression, yet there are few systematic reviews integrating the main miRs associated with the prognostic aspects of CRC. The present study evaluated 113 articles that investigated the differential expression of 100 tissue miRs associated with the main clinical-pathological and

prognostic aspects of CRC, demonstrating that such molecules may be useful as valuable

prognostic biomarkers in CRC.

Expression of the miRNAs can be analyzed in different biological samples such as

tumor tissue [45], blood (serum or plasma) [141], feces [142], or urine [143]. In the present study,

articles that evaluated the expression of miRs in the tumor tissues were prioritized and

selected. Tumor tissues can be stored in liquid nitrogen or paraffin, allowing greater

stability to the miRs molecules for later analysis in retrospective studies that aim to

evaluate patients' survival. The analysis in the tumor tissues confers greater specificity to

the results since circulating miRs (present in serum or plasma) may come from other

anatomical sites. In addition, tumor samples allow the analysis of miRs expression in

comparison to the adjacent normal tissue of the same patient providing greater reliability

to the results. In addition, studies indicate that factors associated to the tumor

microenvironment influence the expression of miRs making the analysis in tumors more

adequate for this type of study [144,145].

The expression of miRs can be evaluated by different methods such as northern

blotting, immunohistochemistry, real-time PCR (RT-PCR), microarray, and others [146].

Each method has its own particularities and limitations. In the present study, only articles

that evaluated the expression of miRs in the CRC by means of RT-PCR using TaqMan

or SYBR Green were selected, because these techniques often present hight sensitivity

and specificity, besides allowing the comparison of a specific miR in relation to an

endogenous control.

The miRs are classified as oncogenic or tumor suppressors according to their

respective target genes. MiRs that inhibit the expression of tumor suppressor genes are

considered oncogenic and are generally upregulated in neoplasms. The most frequent

oncogenic miRs in this study were miR-21, miR-181a, miR-182, miR-183, miR-210 and

miR-224 and the upregulation of most of these miRs was associated with distant

metastasis, lymph node metastasis and poorer prognosis in patients with CRC.

On the other hand, miRs that inhibit the expression of oncogenes are considered

tumor suppressors and are generally downregulated in the carcinogenic processes. The

most frequent tumor suppressor miRs in this study were miR-126, miR-199b and miR-22

and downregulation was associated with the presence of distant metastasis, worse

prognosis and a greater risk of disease relapse (worse disease-free survival). Particularly,

downregulation of miR-199b and miR-22 was also associated with more advanced TNM

stage and the presence of lymph node metastasis. Other miRs such as miR-133a, miR-

133b, miR-16, miR-24-3p and miR-7 were also relatively frequent in the present study,

however, investigations analyzing the expression of these miRs have shown controversial

results. The downregulation of these five miRs was associated with a worse prognosis

and with the exception of miR-7, all the others also had downregulation associated with

the presence of lymph node metastasis.

In addition, another 84 miRs were also identified in the present review and their

differential expression was associated with distinct clinicopathological and prognostic

aspects of CRC, however, these miRs were evaluated in a single study. In general, it was

observed that down and upregulated miRs associated with lymph node metastasis and

worse prognosis (overall survival) in CRC were the most frequently studied.

Among the miRs identified, miR-21 was the most frequent one being analyzed in

six studies. In these studies, an upregulation of miR-21 in the tumor compared to the

adjacent normal tissue was found in addition to a strong correlation with more aggressive

CRC characteristics such as tumor size, lymph node metastasis, distant metastasis and

worse prognosis. A systematic review with meta-analysis conducted by Peng et al.

selected studies that analyzed the role of miR-21 as diagnostic biomarker and prognostic

of CRC, and statistically demonstrated that miR-21 upregulation in the tumor was

associated with worse overall survival and worse disease-free survival, as in the present

review [147]. The findings in both studies suggest the potential of miR-21 to predict survival

and relapse of the disease making it promising biomarker for CRC prognosis.

Another systematic review with meta-analysis conducted by Gao et al. evaluated

the prognostic value of miRs in colorectal cancer as in the present review. However, in

that review, studies that analyzed the expression of miRs in different types of samples

such as serum, plasma and tissue and performing other methods of detection, as

immunohistochemistry, were considered, so they presented miRs not identified in the

present study. Nevertheless, the findings of the review by Gao et al. and the present

review demonstrate similar results regarding the analysis of miRs expression in tumor

tissues associated with the prognostic aspects of CRC, that is, the upregulation of miR-

181a, miR-224, miR-21 and miR-126 downregulation were also present associated with

worse overall survival in CRC patients [21].

The molecular mechanisms by which polyps and adenomas develop to CRC are

diverse. However, the major altered signal pathways responsible for this progression in

CRC are: MAPK / EGFR pathway, TGFβ pathway and via P13K / AKT pathway [148].

Coincidentally, the most frequent target genes for the miRNAs identified in the present

study were also directly associated with these pathways.

The phosphatase and tensin homolog (PTEN) protein is encoded by a tumor

suppressor gene responsible for the inactivation of pathways associated with cell

proliferation. PTEN negatively regulates the PI3K / AKT pathway by hydrolyzing the

Phosphatidylinositol-3,4,5–trisphosphate (PIP3) into Phosphatidylinositol 4,5-

bisphosphate (PIP2) inactivating AKT, a protein that has antiapoptotic action. Similarly,

PTEN inactivates the MAPK pathway by ERK (Extracellular Regulated Kinase)

dephosphorylation, inactivating the transmission of signals through this pathway and

consequently inhibiting cell proliferation. In addition, PTEN activates RAD51, a DNA-

repairing protein, and CENPC (Centromere Protein C), a protein responsible for the

stability of centromere and chromosomal integrity [149]. Situations in which PTEN is

downregulated result in dysregulation in the cell cycle and destabilization of the cellular

DNA and can trigger carcinogenic processes like CRC. In the present study, the

upregulation of miR-181a, miR-17-5p, miR494, miR-22 and miR-21 was correlated with

the downregulation of PTEN gene and worse prognosis of CRC patients.

The transforming growth factor β (TGFβ) is a tumor suppressor cytokinin protein

known to exert important functions in the regulation of apoptosis, differentiation, migration

and cell proliferation. Through the binding of TGFß to its TGFR2 receptor, numerous

proteins of this pathway are phosphorylated and activated, including the SMAD4 protein.

However, studies indicate that mutations or downreglation of SMAD4, can hyperactivate

TGFβ pathway conferring it the potential to promote tumor growth, invasion and

metastasis [148,150]. In the present study, upregulation of miR-20a, miR-20a-5p and miR-

888 was associated with the downregulation of SMAD4 and the presence of distant

metastasis, advanced TNM stage, worse overall survival and worse disease-free survival

in patients with CRC.

The TP53 gene encodes the tumor suppressor protein P53 that plays important roles

in cell cycle control and apoptosis. MiRs belonging to the miR-34 family can modulate the

antiproliferative and pro-apoptotic activity of P53 by inhibiting oncogenes and cell cycle

associated genes. Thus, downregulation of miR-34 or overexpression of oncogenic miRs

that negatively regulate p53 results in a dysregulation of the cell cycle leading to a

carninogenic process [151]. In the present study, the downregulation of P53 was correlated

with the upregulation of miR-125b and downregulation of miR-34a-5p, resulting in a worse

prognosis for patients with CRC.

KRAS is a protein belonging to the RAS family of proteins that control the activation

of several pathways related to cell proliferation, such as the MAPK pathway. MiRs

considered to be tumor suppressors may act as regulators of KRAS expression. Thus,

the downregulation of these miRs may increase the activation of KRAS resulting in a

dysregulated cellular proliferation [152]. The present review demonstrated that upregulation

of KRAS was associated with downregulation of miR-96-5p and miR-384, which in turn

was associated with an advanced TNM stage, presence of distant metastasis and worse

overall survival of patients with CRC. Fig 3 outlines the differential expression of the miRs

and their respective target genes.

Fig 3. Upregulation and downregulation of miRs associated with their respective target genes

and cell functions.

By analyzing the results of the present review fully, it was possible to observe the large

number of miRs associated with the different prognostic aspects of CRC, making them

eligible as possible prognostic markers. However, few articles have analyzed the

expression of two or more miRs concomitantly. Thus, the development of studies that

analyze these miRs synergistically, would allow the identification of network interactions,

making them more sensitive and specific in predicting the prognosis of patients with CRC

and assisting in their therapeutic behavior.

Some limitations were observed during the elaboration of the present study. The large

number of studies and miRs investigated made it difficult to conduct a quantitative and

qualitative analysis capable of demonstrating the most sensitive and specific differentially

expressed miRs associated with each clinical-pathological and prognostic aspect of the

CRC. In addition, the study was limited to articles published in English, making it harder

to obtain data from studies published in other languages.

471   Conclusion

After analyzing the results of the present systematic review, specific tissue miRs

demonstrate to be promising CRC prognostic biomarkers due to their strong correlation

with prognostic aspects of this disease, including advanced TNM stage, presence of

distant metastasis and lymph node metastasis, tumor growth, prediction of worse

prognosis (worse overall survival) and recurrence of the disease (worse disease-free

survival).

The miR-21 was the most frequent studied miR in the present review among the

miRs identified and its upregulation was associated with tumor size, lymph node

metastasis, distant metastasis and worse prognosis making it a promising biomarker of

CRC prognosis. However, new studies are needed to systematize and identify the most

sensitive and specific miRs associated with each prognostic aspect of CRC in order to

choose the most relevant ones and apply them into clinical practice.

In addition, most studies have shown strong correlations between the expression

of miRs and their respective target genes, so the development of functional studies aiming

to elucidate these associations becomes important as a way to help in the description of

new therapeutic targets of this disease.

The vast number of miRs identified in the present study also translates the

complexity and heterogeneity of CRC, emphasizing the importance of new studies that

clarify the regulatory mechanisms of the expression of these miRs in this tumor, allowing

a better understanding of the molecular biology of this disease.

493   References

1.   INCA. Estimativa 2018-Incidência de câncer no Brasil [Internet]. Instituto Nacional

de Câncer José Alencar Gomes da Silva. 2017. 130 p. Available from:

http://www.inca.gov.br/estimativa/2018/estimativa-2018.pdf

2.   Mayer RJ, Wolpin Brian M. Systemic Treatment of Colorectal Cancer.

Gastroenterology. 2009;134(5):1296–310.

3. Beth A. Erickson, Alexander B. Olawaire, Adriana Bermudez E c. G and DGM.

American Joint Committee on Cancer Staging Manual. 8th Edition. Cervix Uteri

[Internet]. 2017. 649–659 p. Available from: www.cancerstaging.org

ajcc@facs.org

4. Wilhelmsen M, Kring T, Jorgensen LN, Madsen MR, Jess P, Bulut O, et al.

Determinants of recurrence after intended curative resection for colorectal cancer.

Scand J Gastroenterol. 2014;49(12):1399–408.

5. Siegel R, DeSantis C, Virgo K, Stein K, Mariotto A, Smith T, et al. Cancer

treatment and survivorship statistics, 2012. CA Cancer J Clin. 2012;62(4):220–41.

6. Destri GL, Rubino AS, Latino R, Giannone F, Lanteri R, Scilletta B, et al.

Preoperative Carcinoembryonic Antigen and Independent Prognostic Factor Still

Reliable. Int Surg. 2015;100:617–25.

7. Thomas J, Ohtsuka M, Pichler M, Ling H. MicroRNAs: Clinical relevance in

colorectal cancer. Int J Mol Sci. 2015;16(12):28063–76.

8. Pardini B, Rosa F, Naccarati A, Vymetalkova V, Ye Y, Wu X, et al.

Polymorphisms in microRNA genes as predictors of clinical outcomes in

colorectal cancer patients. Carcinogenesis. 2015;36(1):82–6.

9. Hansen TF, Christensen RDP, Andersen RF, Sørensen FB, Johnsson A,

Jakobsen A. MicroRNA-126 and epidermal growth factor-like domain 7-an

angiogenic couple of importance in metastatic colorectal cancer. Results from the

Nordic ACT trial. Br J Cancer. 2013;109(5):1243–51.

10.　Rasmussen MH, Jensen NF, Tarpgaard LS, Qvortrup C, Rømer MU, Stenvang J,

et al. High expression of microRNA-625-3p is associated with poor response to

first-line oxaliplatin based treatment of metastatic colorectal cancer. Mol Oncol.

2013;7(3):1–10.

11.　Masuda T, Hayashi N, Kuroda Y, Ito S, Educhi H, Mimori K. MicroRNAs as

biomarkers in colorectal cancer. Cancers (Basel). 2017;9(124):1–15.

12.　Toiyama Y, Hur K, Tanaka K, Inoue Y, Kusunoki M, Boland CR, et al. Serum miR-

200c is a novel prognostic and metastasis-predictive biomarker in patients with

colorectal cancer. Ann Surg. 2014;259(4):735–43.

13.　Gregory RI, Shiekhattar R, Gregory RI, Shiekhattar R. MicroRNA Biogenesis and

Cancer MicroRNA Biogenesis and Cancer. 2005;(9):3509–12.

14.　Hammond SM. An overview of microRNAs. Adv Drug Deliv Rev. 2015;87:3–14.

15.　Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, et al. The nuclear RNase III Drosha

initiates microRNA processing. Nature [Internet]. 2003;425(6956):415–9.

Available from: www.nature.com/nature

16.　MacFarlane L-A, R. Murphy P. MicroRNA: Biogenesis, Function and Role in

Cancer. Curr Genomics. 2010;11(7):537–61.

17.　Costa E de BO, Pacheco C. Micro-RNAs : Perspectivas atuais da regulação da

expressão gênica em eucariotos. Biosaúde. 2012;14(2):81–93.

18.　Calin GA, Croce CM. MicroRNA signatures in human cancers. Nat Rev Cancer.

2006;6(11):857–66.

19.   Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor

suppressors. Dev Biol. 2007;302(1):1–12.

20.   Xi Y, Formentini A, Chien M, Weir DB, Russo JJ, Ju J, et al. Prognostic Values of

microRNAs in Colorectal Cancer. Biomark Insights [Internet]. 2007;1(251):113–

21. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2134920/

21.   Gao S, Zhao Z-Y, Wu R, Zhang Y, Zhang Z-Y. Prognostic value of microRNAs in

colorectal cancer : a meta-analysis. Cancer Manag. 2018;907–29.

22.   Michael MZ, O' Connor SM, van Holst Pellekaan NG, Young GP, James RJ.

Reduced accumulation of specific microRNAs in colorectal neoplasia. Mol Cancer

Res. 2003;1(12):882–91.

23.   Siegel R, Desantis C, Jemal A. Colorectal Cancer Statistics, 2014. CA Cancer J

Clin [Internet]. 2014;64(1):104–17. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24639052

24.   Mima K, Nishihara R, Yang J, Dou R, Masugi Y, Shi Yy, et al. MicroRNA MIR21

(miR-21) and PTGS2 expression in colorectal cancer and patient survival. Clin

Cancer Res. 2016;22(15):3841–3848.

25.   Fukushima Y, Iinuma H, Tsukamoto M, Matsuda K, Hashiguchi Y. Clinical

significance of microRNA-21 as a biomarker in each Dukes' stage of colorectal

cancer. Oncol Rep. 2015;33(2):573–82.

26. Chen T-H, Wang K-L, Liu C-N, Lee H, Lin C-C, Huang C-C, et al. The prognostic

significance of APC gene mutation and miR-21 expression in advanced-stage

colorectal cancer. Color Dis. 2013;15(11):1367–74.

27. Toiyama Y, Takahashi M, Hur K, Nagasaka T, Tanaka K, Inoue Y, et al. Serum

miR-21 as a diagnostic and prognostic biomarker in colorectal cancer. J Natl

Cancer Inst. 2013;105(12):849–59.

28. Wu Y, Song Y, Xiong Y, Wang X, Xu K, Han B, et al. MicroRNA-21 (Mir-21)

Promotes Cell Growth and Invasion by Repressing Tumor Suppressor PTEN in

Colorectal Cancer. Cell Physiol Biochem. 2017;43(3):945–58.

29. Shibuya H, Iinuma H, Shimada R, Horiuchi A, Watanabe T. Clinicopathological

and Prognostic Value of MicroRNA-21 and MicroRNA-155 in Colorectal Cancer.

Oncology. 2010;79:313–20.

30. Ji D, Chen Z, Li M, Zhan T, Yao Y, Zhang Z, et al. MicroRNA-181a promotes

tumor growth and liver metastasis in colorectal cancer by targeting the tumor

suppressor WIF-1. Mol Cancer. 2014;13(1).

31. Nishimura J, Handa R, Yamamoto H, Tanaka F, Shibata K, Mimori K, et al.

microRNA-181a is associated with poor prognosis of colorectal cancer. Oncol

Rep. 2012;28(6):2221–6.

32. Li Z, Wang H, Xu Z, Sun Y, Han J. Expression and mechanism of microRNA-181a

on incidence and survival in late liver metastases of colorectal cancer. Oncol Rep.

2016;35(3):1403–8.

33.   Xiao G, Tang H, Wei W, Li J, Ji L, Ge J. Aberrant Expression of MicroRNA-15a

and MicroRNA-16 Synergistically Associates with Tumor Progression and

Prognosis in Patients with Colorectal Cancer. Gastroenterol Res Pract.

2014;2014.

34.   Diamantopoulos MA, Kontos CK, Kerimis D, Papadopoulos IN, Scorilas A.

Upregulated miR-16 expression is an independent indicator of relapse and poor

overall survival of colorectal adenocarcinoma patients. Clin Chem Lab Med.

2017;55(5):737–47.

35.   Qian J, Jiang B, Li M, Chen J, Fang M. Prognostic significance of microRNA-16

expression in human colorectal cancer. World J Surg. 2013;37(12):2944–9.

36.   Guo Y, Li X, Lin C, Zhang Y, Hu G, Zhou J, et al. MicroRNA-133b inhibits

connective tissue growth factor in colorectal cancer and correlates with the clinical

stage of the disease. Mol Med Rep. 2015;11(4):2805–12.

37.   Ellermeier C, Vang S, Cleveland K, Durand W, Resnick MB, Brodsky AS.

Prognostic microRNA expression signature from examination of colorectal primary

and metastatic tumors. Anticancer Res. 2014;34(8):3957–68.

38.   Akçakaya P, Ekelund S, Kolosenko I, Caramuta S, Özata D, Xie H, et al. miR-185

and miR-133b deregulation is associated with overall survival and metastasis in

colorectal cancer. Int J Oncol. 2011;39(2):311–8.

39.   Liu Y, Zhou Y, Feng X, Yang P, Yang J, Ping A, et al. Low Expression of

MicroRNA-126 is Associated with. Wiley Period. 2014;53:358–365.

40. Wang LL, Du LT, Li J, Liu YM, Qu AL, Yang YM, et al. Decreased expression of

miR-133a correlates with poor prognosis in colorectal cancer patients. World J

Gastroenterol. 2014;20(32):11340–6.

41. Wan TMH, Lam CSC, Ng L, Chow AKM, Wong SKM, Li HS, et al. The

Clinicopathological Significance of miR-133a in Colorectal Cancer. Dis Markers.

2014;2014.

42. Rapti SM, Kontos CK, Papadopoulos IN, Scorilas A. Enhanced miR-182

transcription is a predictor of poor overall survival in colorectal adenocarcinoma

patients. Clin Chem Lab Med. 2014;52(8):1217–27.

43. Liu H, Du L, Wen Z, Yang Y, Li J, Wang L, et al. Up-regulation of miR-182

expression in colorectal cancer tissues and its prognostic value. Int J Colorectal

Dis. 2013;28(5):697–703.

44. Chen Y, Song W. Wnt/catenin β1/microRNA 183 predicts recurrence and

prognosis of patients with colorectal cancer. Oncol Lett [Internet].

2018;15(4):4451–6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/29541213%0Ahttp://www.pubmedcentral.nih

.gov/articlerender.fcgi?artid=PMC5835957

45. Zhou T, Zhang GJ, Zhou H, Xiao HX, Li Y. Overexpression of microRNA-183 in

human colorectal cancer and its clinical significance. Eur J Gastroenterol Hepatol.

2014;26(2):229–33.

46. Cristóbal I, Caramés C, Rincón R, Manso R, Madoz-Gúrpide J, Torrejón B, et al.

Downregulation of microRNA-199b predicts unfavorable prognosis and emerges

as a novel therapeutic target which contributes to PP2A inhibition in metastatic

colorectal cancer. Oncotarget [Internet]. 2017;8(25):40169–80. Available from:

http://www.oncotarget.com/fulltext/11174

47. Shen ZZ-LZ, Wang B, Jiang KK-WK, Ye CC-XC, Cheng C, Yan Y-CYY-CY, et al.

Downregulation of miR-199b is associated with distant metastasis in colorectal

cancer via activation of SIRT1 and inhibition of CREB/KISS1 signaling.

Oncotarget [Internet]. 2016;7(23):35092–105. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27145368%5Cnhttp://europepmc.org/abstrac

t/med/27145368

48. Zhang G, Xia S, Tian H, Liu Z, Zhou T. Clinical significance of miR-22 expression

in patients with colorectal cancer. Med Oncol. 2012;29(5):3108–12.

49. Xia S-S, Zhang G-J, Liu Z-L, Tian H-P, He Y, Meng C-Y, et al. MicroRNA-22

suppresses the growth, migration and invasion of colorectal cancer cells through

a Sp1 negative feedback loop. Oncotarget [Internet]. 2017;8(22):36266–78.

Available from:

http://www.oncotarget.com/fulltext/16742%0Ahttp://www.ncbi.nlm.nih.gov/pubmed

/28422727%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC54

82653

50. Adamopoulos PG, Kontos CK, Rapti SM, Papadopoulos IN, Scorilas A. MiR-224

overexpression is a strong and independent prognosticator of short-term relapse

and poor overall survival in colorectal adenocarcinoma. Int J Oncol.

2015;46(2):849–59.

51.   Liao WT, Li TT, Wang ZG, Wang SY, He MR, Ye YP, et al. microRNA-224

promotes cell proliferation and tumor growth in human colorectal cancer by

repressing PHLPP1 and PHLPP2. Clin Cancer Res. 2013;19(17):4662–72.

52.   Kerimis D, Kontos CK, Christodoulou S, Papadopoulos IN, Scorilas A. Elevated

expression of miR-24-3p is a potentially adverse prognostic factor in colorectal

adenocarcinoma. Clin Biochem [Internet]. 2017;50(6):285–92. Available from:

http://dx.doi.org/10.1016/j.clinbiochem.2016.11.034

53.   Gao Y, Liu Y, Du L, Li J, Qu A, Zhang X, et al. Down-regulation of miR-24-3p in

colorectal cancer is associated with malignant behavior. Med Oncol. 2015;32(1).

54.   NAGANO Y, TOIYAMA Y, OKUGAWA Y, IMAOKA H, FUJIKAWA H, YASUDA H,

et al. MicroRNA-7 Is Associated with Malignant Potential and Poor Prognosis in

Human Colorectal Cancer. Anticancer Res [Internet]. 2016;36(12):6521–6.

Available from: http://ar.iiarjournals.org/content/36/12/6521.abstract

55.   Suto T, Yokobori T, Yajima R, Morita H, Fujii T, Yamaguchi S, et al. MicroRNA-7

expression in colorectal cancer is associated with poor prognosis and regulates

cetuximab sensitivity via EGFR regulation. Carcinogenesis. 2015;36(3):338–45.

56.   Ge J, Chen Z, Li R, Lu T, Xiao G. Upregulation of microRNA-196a and microRNA-

196b cooperatively correlate with aggressive progression and unfavorable

prognosis in patients with colorectal cancer. Cancer Cell Int. 2014;14(1):1–8.

57.   Li H, Dai S, Zhen T, Shi H, Zhang F, Yang Y, et al. Clinical and biological

significance of miR-378a-3p and miR-378a-5p in colorectal cancer. Eur J Cancer

[Internet]. 2014;50(6):1207–21. Available from:

http://dx.doi.org/10.1016/j.ejca.2013.12.010

58.   Wang CJ, Zhou ZG, Wang L, Yang L, Zhou B, Gu J, et al. Clinicopathological

significance of microRNA-31, -143 and -145 expression in colorectal cancer. Dis

Markers. 2009;26(1):27–34.

59.   Fang L, Li H, Wang L, Hu J, Jin T, Wang J, et al. MicroRNA-17-5p promotes

chemotherapeutic drug resistance and tumour metastasis of colorectal cancer by

repressing PTEN expression. Oncotarget [Internet]. 2014;5(10):2974–87.

Available from: http://www.oncotarget.com/fulltext/1614

60.   Sun HB, Chen X, Ji H, Wu T, Lu HW, Zhang Y, et al. MiR-494 is an independent

prognostic factor and promotes cell migration and invasion in colorectal cancer by

directly targeting PTEN. Int J Oncol. 2014;45(6):2486–94.

61.   Zhang GJ, Li Y, Zhou H, Xiao HX, Zhou T. miR-20a is an independent prognostic

factor in colorectal cancer and is involved in cell metastasis. Mol Med Rep.

2014;10(1):283–91.

62.   Cheng D, Zhao S, Tang H, Zhang D, Sun H, Yu F, et al. MicroRNA-20a-5p

promotes colorectal cancer invasion and metastasis by downregulating Smad4.

Oncotarget [Internet]. 2016;7(29):45199–213. Available from:

http://www.oncotarget.com/fulltext/9900

63. Gao S-J, Chen L, Lu W, Zhang L, Wang L, Zhu H-H. miR‐888 functions as an

oncogene and predicts poor prognosis in colorectal cancer. Oncol Lett.

2018;15:9101–9.

64. Gao J, Li N, Dong Y, Li S, Xu L, Li X, et al. MiR-34a-5p suppresses colorectal

cancer metastasis and predicts recurrence in patients with stage II/III colorectal

cancer. Oncogene. 2015;34(31):4142–52.

65. Nishida N, Yokobori T, Mimori K, Sudo T, Tanaka F, Shibata K, et al. MicroRNA

miR-125b is a prognostic marker in human colorectal cancer. Int J Oncol.

2011;38(5):1437–43.

66. Ma W, Liu B, Li J, Jiang J, Zhou R, Huang L, et al. MicroRNA-302c represses

epithelial–mesenchymal transition and metastasis by targeting transcription factor

AP-4 in colorectal cancer. Biomed Pharmacother [Internet].

2018;105(March):670–6. Available from:

https://doi.org/10.1016/j.biopha.2018.06.025

67. Tong F, Ying Y, Pan H, Zhao W, Li H, Zhan X. Microrna-466 (Mir-466) functions

as a tumor suppressor and prognostic factor in colorectal cancer (CRC). Bosn J

Basic Med Sci. 2018;18(3):252–9.

68. Zhang L, Xie F, Tang C, Xu W, Ding X, Liang J. miR-340 suppresses tumor

growth and enhances chemosensitivity of colorectal cancer by targeting RLIP76.

Eur Rev Med Pharmacol Sci. 2017;21:2875–86.

69. Stiegelbauer V, Vychytilova-Faltejskova P, Karbiener M, Pehserl AM, Reicher A,

Resel M, et al. MiR-196b-5p regulates colorectal cancer cell migration and metastases through interaction with HOXB7 and GALNT5. Clin Cancer Res. 2017;23(17):5255–66.

70. Wen L, Li Y, Jiang Z, Zhang Y, Yang B, Han F. MiR-944 inhibits cell migration and invasion by targeting MACC1 in colorectal cancer. Oncol Rep. 2017;37(6):3415–22.

71. Yan L, Qiu J, Yao J. Downregulation of microRNA-30d promotes cell proliferation and invasion by targeting LRH-1 in colorectal carcinoma. Int J Mol Med. 2017;39(6):1371–80.

72. He Z, Yu L, Luo S, Li M, Li J, Li Q, et al. miR-296 inhibits the metastasis and epithelial-mesenchymal transition of colorectal cancer by targeting S100A4. BMC Cancer. 2017;17(1):1–10.

73. Wang Y-X, Chen Y-R, Liu S-S, Ye Y-P, Jiao H-L, Wang S-Y, et al. MiR-384 inhibits human colorectal cancer metastasis by targeting KRAS and CDC42. Oncotarget [Internet]. 2016;7(51):84826–38. Available from: http://www.oncotarget.com/fulltext/12704

74. Wu L, Shi B, Huang K, Fan G. MicroRNA-128 suppresses cell growth and metastasis in colorectal carcinoma by targeting IRS1. Oncol Rep. 2015;34(5):2797–805.

75. Liao W, Gu C, Huang A, Yao J, Sun R. MicroRNA-33b inhibits tumor cell growth and is associated with prognosis in colorectal cancer patients. Clin Transl Oncol.

2016;18(5):449–56.

76.　Qu YL, Wang HF, Sun ZQ, Tang Y, Han XN, Yu XB, et al. Up-regulated miR-155-

5p promotes cell proliferation, invasion and metastasis in colorectal carcinoma. Int

J Clin Exp Pathol. 2015;8(6):6988–94.

77.　Kai Y, Qiang C, Xinxin P, Miaomiao Z, Kuailu L. Decreased miR-154 expression

and its clinical significance in human colorectal cancer. World J Surg Oncol

[Internet]. 2015;13(195):615–9. Available from: http://dx.doi.org/10.1186/s12957-

015-0607-5

78.　Xu K, Liu X, Mao X, Xue L, Wang R, Chen L, et al. MicroRNA-149 suppresses

colorectal cancer cell migration and invasion by directly targeting forkhead box

transcription factor FOXM1. Cell Physiol Biochem. 2015;35(2):499–515.

79.　Wang B, Shen ZL, Gao ZD, Zhao G, Wang CY, Yang Y, et al. MiR-194,

commonly repressed in colorectal cancer, suppresses tumor growth by regulating

the MAP4K4/c- Jun/MDM2 signaling pathway. Cell Cycle. 2015;14(7):1046–58.

80.　Chen P, Xi Q, Wang Q, Wei P. Downregulation of microRNA-100 correlates with

tumor progression and poor prognosis in colorectal cancer. Med Oncol.

2014;31(10):1–6.

81.　Sun ZF, Zhang Z, Liu Z, Qiu B, Liu K, Dong G. MicroRNA-335 inhibits invasion

and metastasis of colorectal cancer by targeting ZEB2. Med Oncol. 2014;31(6).

82.　Yu H, Gao G, Jiang L, Guo L, Lin M, Jiao X, et al. Decreased expression of miR-

218 is associated with poor prognosis in patients with colorectal cancer. Int J Clin Exp Pathol. 2013;6(12):2904–11.

83. Tao Y, Ma C, Fan Q, Wang Y, Han T, Sun C. MicroRNA-1296 facilitates proliferation, migration and invasion of colorectal cancer cells by targeting SFPQ. J Cancer. 2018;9(13):2317–26.

84. Chen W, Tong K, Yu J. MicroRNA-130a is upregulated in colorectal cancer and promotes cell growth and motility by directly targeting forkhead box F2. Mol Med Rep. 2017;16(4):5241–8.

85. Rapti SM, Kontos CK, Papadopoulos IN, Scorilas A. High miR-96 levels in colorectal adenocarcinoma predict poor prognosis, particularly in patients without distant metastasis at the time of initial diagnosis. Tumor Biol [Internet]. 2016;37(9):11815–24. Available from: http://dx.doi.org/10.1007/s13277-016-5023-0

86. Ma W, Yu Q, Jiang J, Du X, Huang L, Zhao L, et al. miR-517a is an independent prognostic marker and contributes to cell migration and invasion in human colorectal cancer. Oncol Lett. 2016;11(1):2583–9.

87. Zhang GJ, Li JS, Zhou H, Xiao HX, Li Y, Zhou T. MicroRNA-106b promotes colorectal cancer cell migration and invasion by directly targeting DLC1. J Exp Clin Cancer Res. 2015;34(1):1–11.

88. Perez-Carbonell L, Sinicrope FA, Alberts SR, Oberg AL, Balaguer F, Castells A, et al. MiR-320e is a novel prognostic biomarker in colorectal cancer. Br J Cancer

[Internet]. 2015;113(1):83–90. Available from:

http://dx.doi.org/10.1038/bjc.2015.168

89. Liu M, Zhi Q, Wang W, Zhang Q, Fang T, Ma Q. Up-regulation of miR-592

correlates with tumor progression and poor prognosis in patients with colorectal

cancer. Biomed Pharmacother [Internet]. 2015;69:214–20. Available from:

http://dx.doi.org/10.1016/j.biopha.2014.12.001

90. Wang X, Kuang Y, Shen X, Zhou H, Chen Y, Han Y, et al. Evaluation of miR-720

prognostic significance in patients with colorectal cancer. Tumor Biol.

2014;36(2):719–27.

91. Chu D, Zheng J, Li J, Li Y, Zhang J, Zhao Q, et al. MicroRNA-630 is a prognostic

marker for patients with colorectal cancer. Tumor Biol. 2014;35(10):9787–92.

92. Qu A, Du L, Yang Y, Liu H, Li J, Wang L, et al. Hypoxia-inducible MiR-210 is an

independent prognostic factor and contributes to metastasis in colorectal cancer.

PLoS One. 2014;9(3).

93. Li X, Yang C, Wang X, Zhang J, Zhang R, Liu R. The expression of miR-25 is

increased in colorectal cancer and is associated with patient prognosis. Med

Oncol. 2014;31(1).

94. Qin S, Zhu Y, Ai F, Li Y, Bai B, Yao W, et al. MicroRNA-191 correlates with poor

prognosis of colorectal carcinoma and plays multiple roles by targeting tissue

inhibitor of metalloprotease 3 S. Neoplasma. 2014;61(1):27–34.

95.   Zhou T, Zhang G, Liu Z, Xia S, Tian H. Overexpression of miR-92a correlates with

790   tumor metastasis and poor prognosis in patients with colorectal cancer. Int J Color

791   Dis. 2013;28(11):19–24.

792   96.   Wang N, Liu W. Increased expression of miR-552 acts as a potential predictor

793   biomarker for poor prognosis of colorectal cancer. Eur Rev Med Pharmacol Sci.

794   2018;22:412–6.

795   97.   Liu TP, Huang CC, Yeh KT, Ke TW, Wei PL, Yang JR, et al. Down-regulation of

796   let-7a-5p predicts lymph node metastasis and prognosis in colorectal cancer:

797   Implications for chemotherapy. Surg Oncol [Internet]. 2016;25(4):429–34.

798   Available from: http://dx.doi.org/10.1016/j.suronc.2016.05.016

799   98.   Mokutani Y, Uemura M, Munakata K, Okuzaki D, Haraguchi N, Takahashi H, et al.

800   Down-Regulation of microRNA-132 is Associated with Poor Prognosis of

801   Colorectal Cancer. Ann Surg Oncol. 2016;23:599–608.

802   99.   Sun P, Sun D, Wang X, Liu T, Ma Z, Duan L. miR-206 is an independent

803   prognostic factor and inhibits tumor invasion and migration in colorectal cancer.

804   Cancer Biomarkers. 2015;15(4):391–6.

805   100. Zhang G, Zhou H, Xiao H, Li Y, Zhou T. MiR-378 is an independent prognostic

806   factor and inhibits cell growth and invasion in colorectal cancer. BMC Cancer

807   [Internet]. 2014;14(1):109. Available from:

808   http://bmccancer.biomedcentral.com/articles/10.1186/1471-2407-14-109

809   101. Song Y, Xu Y, Wang Z, Chen Y, Yue Z, Gao P, et al. MicroRNA-148b suppresses

cell growth by targeting cholecystokinin-2 receptor in colorectal cancer. Int J Cancer. 2012;131(5):1042–51.

102. Bin L, Zhang M, Lixia L, Aimin Z, Hua Y, Yanhong S, et al. Down-regulation of miRNA-30c predicts poor prognosis in Colorectal Cancer patients. Revi Rom Med Labo. 2016;24(4):369–75.

103. Lin M, Duan B, Hu J, Yu H, Sheng H, Gao H, et al. Decreased expression of miR-193a-3p is associated with poor prognosis in colorectal cancer. Oncol Lett. 2017;14(1):1061–7.

104. Li J, Du L, Yang Y, Wang C, Liu H, Wang L, et al. MiR-429 is an independent prognostic factor in colorectal cancer and exerts its anti-apoptotic function by targeting SOX2. Cancer Lett [Internet]. 2013;329(1):84–90. Available from: http://dx.doi.org/10.1016/j.canlet.2012.10.019

105. Cui BH, Hong X. Mir-6852 serves as a prognostic biomarker in colorectal cancer and inhibits tumor growth and metastasis by targeting tcf7. Exp Ther Med. 2018;16(2):879–85.

106. Wang L, Jiang C, Li D, Ge X, Shi Z, Li C, et al. MicroRNA-497 inhibits tumor growth and increases chemosensitivity to 5-fluorouracil treatment by targeting KSR1. Oncotarget [Internet]. 2016;7(3). Available from: http://www.oncotarget.com/fulltext/6545

107. Zhang P, Ji DB, Han HB, Shi YF, Du CZ, Gu J. Downregulation of miR-193a-5p correlates with lymph node metastasis and poor prognosis in colorectal cancer.

831 World J Gastroenterol. 2014;20(34):12241–8.

108. Orang A V, Safaralizadeh R, Hosseinpour Feizi MA, Somi MH. Diagnostic and prognostic value of miR-205 in colorectal cancer. Asian Pac J Cancer Prev. 2014;15(9):4033–7.

109. Long L, Huang G, Zhu H, Guo Y, Liu Y, Huo J. Down-regulation of miR-138 promotes colorectal cancer metastasis via directly targeting TWIST2. J Transl Med. 2013;11(1):1–10.

110. Lou X, Qi X, Zhang Y, Long H, Yang J. Decreased expression of microRNA-625 is associated with tumor metastasis and poor prognosis in patients with colorectal cancer. J Surg Oncol. 2013;108(4):230–5.

111. Meng X, Wu J, Pan C, Wang H, Ying X, Zhou Y, et al. Genetic and Epigenetic Down-regulation of MicroRNA-212 Promotes Colorectal Tumor Metastasis via Dysregulation of MnSOD. Gastroenterology [Internet]. 2013;145(2):426-436.e6. Available from: http://dx.doi.org/10.1053/j.gastro.2013.04.004

112. Wang X, Wang J, Ma H, Zhang J, Zhou X. Downregulation of miR-195 correlates with lymph node metastasis and poor prognosis in colorectal cancer. Med Oncol. 2012;29(34):919–27.

113. Roh MS, Lee HW, Jung SB, Kim K, Lee EH, Park M il, et al. Expression of miR-200c and its clinicopathological significance in patients with colorectal cancer. Pathol Res Pract [Internet]. 2018;214(3):350–5. Available from: http://dx.doi.org/10.1016/j.prp.2018.01.005

114. Liu D-R, Guan Q-L, Gao M-T, Jiang L, Kang H-X. miR-1260b is a Potential

Prognostic Biomarker in Colorectal Cancer. Med Sci Monit. 2016;22:2417–23.

115. Mo Z-H, Wu X-D, Li S, Fei B-Y, Zhang B. Expression and Clinical Significance of

MicroRNA-376a in Colorectal Cancer. Asian Pacific J Cancer Prev.

2014;15(21):9265–70.

116. Wan D, He S, Xie B, Xu G, Gu W, Shen C, et al. Aberrant expression of miR-

199a-3p and its clinical significance in colorectal cancers. Med Oncol. 2013;30(1).

117. Nishida N, Yamashita S, Mimori K, Sudo T, Tanaka F, Shibata K, et al.

MicroRNA-10b is a prognostic indicator in colorectal cancer and confers

resistance to the chemotherapeutic agent 5-fluorouracil in colorectal cancer cells.

Ann Surg Oncol. 2012;19(9):3065–71.

118. Yu L, Lu Y, Han X, Zhao W, Li J, Mao J, et al. microRNA -140-5p inhibits

colorectal cancer invasion and metastasis by targeting ADAMTS5 and IGFBP5.

Stem Cell Res Ther [Internet]. 2016;7(1):1–11. Available from:

http://dx.doi.org/10.1186/s13287-016-0438-5

119. Li W, Chang J, WWang S, Liu X, Junjie P, Huang D, et al. miRNA-99b-5p

suppresses liver metastasis of colorectal cancer by down-regulating mTOR.

Oncotarget [Internet]. 2015;6(27):24448–62. Available from:

http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L

606354568%5Cnhttp://dx.doi.org/10.18632/oncotarget.4423%5Cnhttp://sfx.library

.uu.nl/utrecht?sid=EMBASE&issn=19492553&id=doi:10.18632/oncotarget.4423&

atitle=miRNA-99b-5p+suppresses+l

120. Ress AL, Stiegelbauer V, Winter E, Schwarzenbacher D, Kiesslich T, Lax S, et al. MiR-96-5p influences cellular growth and is associated with poor survival in colorectal cancer patients. Mol Carcinog. 2015;54(11):1442–50.

121. Yamashita S, Yamamoto H, Mimori K, Nishida N, Takahashi H, Haraguchi N, et al. MicroRNA-372 is associated with poor prognosis in colorectal cancer. Oncology. 2012;82(4):205–12.

122. Zhou C, Cui F, Li J, Wang D, Wei Y, Wu Y, et al. MiR-650 represses high-risk non-metastatic colorectal cancer progression via inhibition of AKT2/GSK3beta/E-cadherin pathway. Oncotarget [Internet]. 2017;8(30):49534–47. Available from: http://www.impactjournals.com/oncotarget/index.php?journal=oncotarget&page=article&op=download&path%5B%5D=17743&path%5B%5D=56826%0Ahttp://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emex&NEWS=N&AN=617528910%0Ahttp://ovidsp.ovid.com/ovidweb.cgi?T=JS&

123. Wang ZS, Zhong M, Bian YH, Mu YF, Qin SL, Yu MH, et al. MicroRNA-187 inhibits tumor growth and invasion by directly targeting CD276 in colorectal cancer. Oncotarget [Internet]. 2016;7(28):44266–76. Available from: https://www.ncbi.nlm.nih.gov/pubmed/27329595%0Ahttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC5190094/pdf/oncotarget-07-44266.pdf

124. Hibino Y, Sakamoto N, Naito Y, Goto K, Oo HZ, Sentani K, et al. Significance of miR-148a in colorectal neoplasia: Downregulation of miR-148a contributes to the

carcinogenesis and cell invasion of colorectal cancer. Pathobiology.

2015;82(5):233–41.

125. Wang B, Shen ZL, Jiang KW, Zhao G, Wang C you, Yan Y chao, et al. MicroRNA-

217 functions as a prognosis predictor and inhibits colorectal cancer cell

proliferation and invasion via an AEG-1 dependent mechanism. BMC Cancer

[Internet]. 2015;15(1):1–11. Available from: ???

126. Inoue A, Yamamoto H, Uemura M, Nishimura J, Hata T, Takemasa I, et al.

MicroRNA-29b is a Novel Prognostic Marker in Colorectal Cancer. Ann Surg

Oncol. 2015;22(April):1410–8.

127. Yin Y, Zhang B, Wang W, Fei B, Quan C, Zhang J, et al. MiR-204-5p inhibits

proliferation and invasion and enhances chemotherapeutic sensitivity of colorectal

cancer cells by downregulating RAB22A. Clin Cancer Res. 2014;20(23):6187–

699.

128. Song M, Yin Y, Zhang J, Zhang B, Bian Z, Quan C, et al. MiR-139-5p inhibits

migration and invasion of colorectal cancer by downregulating AMFR and

NOTCH1. Protein Cell. 2014;5(11):851–61.

129. Jinushi T, Shibayama Y, Kinoshita I, Oizumi S, Jinushi M, Aota T, et al. Low

expression levels of microRNA-124-5p correlated with poor prognosis in

colorectal cancer via targeting of SMC4. Cancer Med. 2014;3(6):1544–52.

130. Wang MJ, Li Y, Wang R, Wang C, Yu YY, Yang L, et al. Downregulation of

microRNA-124 is an independent prognostic factor in patients with colorectal

cancer. Int J Colorectal Dis. 2013;28(2):183–9.

131. Tsiakanikas P, Kontos CK, Kerimis D, Papadopoulos IN, Scorilas A. High

microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term

relapse of node-negative patients and poor overall survival of patients with non-

metastatic disease. Clin Chem Lab Med. 2018;56(6):990–1000.

132. Rapti SM, Kontos CK, Christodoulou S, Papadopoulos IN, Scorilas A. miR-34a

overexpression predicts poor prognostic outcome in colorectal adenocarcinoma,

independently of clinicopathological factors with established prognostic value. Clin

Biochem [Internet]. 2017;50(16–17):918–24. Available from:

http://dx.doi.org/10.1016/j.clinbiochem.2017.06.004

133. Kontos CK, Tsiakanikas P, Avgeris M, Papadopoulos IN, Scorilas A. miR-15a-5p,

A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma.

Mol Diagnosis Ther. 2017;21(4):453–64.

134. Yamazaki N, Koga Y, Taniguchi H, Kojima M, Kanemitsu Y, Saito N, et al. High

expression of miR-181c as a predictive marker of recurrence in stage II colorectal

cancer. Oncotarget [Internet]. 2017;8(4):6970–83. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/28036302

135. Noguchi T, Toiyama Y, Kitajima T, Imaoka H, Hiro J, Saigusa S, et al. MiRNA-503

Promotes Tumor Progression and Is Associated with Early Recurrence and Poor

Prognosis in Human Colorectal Cancer. Oncol. 2016;90(4):221–31.

136. Sümbül AT, Göğebakan B, Bayram S, Batmacı CY, Öztuzcu S. MicroRNA 211

expression is upregulated and associated with poor prognosis in colorectal

cancer: a case–control study. Tumor Biol. 2015;36(12):9703–9.

137. Li Z wu, Yang Y mei, Du L tao, Dong Z, Wang L li, Zhang X, et al. Overexpression

of miR-223 correlates with tumor metastasis and poor prognosis in patients with

colorectal cancer. Med Oncol. 2014;31(11):1–5.

138. Yiu AJ, Yiu CY. Biomarkers in Colorectal Cancer. Anticancer Res. 2016;36:1093–

338.

139. Soares BA dos S, Penna KGBD, Saddi VA. Prognostic impact of micro-rna

expression in breast cancer: systematic review. Mastology. 2018;23(4):268–75.

140. Hajizamani S, Shahjahani M, Shahrabi S, Saki N. MicroRNAs as prognostic

biomarker and relapse indicator in leukemia. Clin Transl Oncol. 2017;19(8):951–

60.

141. Binderup HG, Madsen JS, Helweg Heegaard NH, Houlind K, Andersen RF,

Brasen CL. Quantification of microRNA levels in plasma – Impact of preanalytical

and analytical conditions. PLoS One. 2018;13(7):1DUMMY.

142. Ahmed FE, Ahmed NaC, Gouda MM, Vos PW, Bonnerup C. RT-qPCR for Fecal

Mature MicroRNA Quantification and Validation. Color Cancer Methods Protoc

Methods Mol Biol. 2018;1765:203–15.

143. Fuessel S, Lohse-Fischer A, Van DV, Solomo K, Erdmann K, Wirth MP.

Quantification of MicroRNAs in Urine-Derived Specimens Susanne. Urothelial

Carcinoma Methods Protoc Methods Mol Biol. 2018;1655(16):201–26.

144. Wentz-Hunter KK, Potashkin JA. The Role of miRNAs as Key Regulators in the

Neoplastic Microenvironment. Mol Biol Int. 2011;2011:1–8.

145. Radovich M, Sood AK, Armaiz-Pena GN, Rodriguez-Aguayo C, Zhang W,

Pradeep S, et al. Hypoxia-mediated downregulation of miRNA biogenesis

promotes tumour progression. Nat Commun [Internet]. 2014;5(1):1–13. Available

from: http://dx.doi.org/10.1038/ncomms6202

146. Hennessy E, Driscoll LO. MicroRNA Expression Analysis: Techniques Suitable for

Studies of Intercellular and Extracellular MicroRNA. In: Gene Expression Profiling:

Methods and Protocols, Methods in Molecular Biology. 2011. p. 99–107.

147. Peng Q, Zhang X, Min M, Zou L, Shen P, Yaqun Z. The clinical role of microRNA-

21 as a promising biomarker in the diagnosis and prognosis of colorectal cancer:

a systematic review and meta-analysis. Oncotarget. 2017;8(27):44893–909.

148. Shirafkan N, Mansoori B, Mohammadi A, Shomali N, Ghasbi M, Baradaran B.

MicroRNAs as novel biomarkers for colorectal cancer: New outlooks. Biomed

Pharmacother [Internet]. 2018;97(August 2017):1319–30. Available from:

http://dx.doi.org/10.1016/j.biopha.2017.11.046

149. Chung JH, Eng C. Nuclear-cytoplasmic partitioning of phosphatase and tensin

homologue deleted on chromosome 10 (PTEN) differentially regulates the cell

cycle and apoptosis. Cancer Res. 2005;65(18):8096–100.

150.   Zhang B, Halder SK, Kashikar ND, Cho Y, Datta A, David L, et al. Anti-metastatic

977   Role of Smad4 Signaling in Colorectal Cancer Bixiang. Gastroenterology.

978   2010;138(3):969–80.

979   151.   Oren M, Bentwich Z, Rosenfeld N, Raver-Shapira N, Moskovits N, Spector Y, et

980   al. Transcriptional Activation of miR-34a Contributes to p53-Mediated Apoptosis.

981   Mol Cell. 2007;26(5):731–43.

982   152.   Jones RP, Sutton PA, Evans JP, Clifford R, McAvoy A, Lewis J, et al. Specific

983   mutations in KRAS codon 12 are associated with worse overall survival in patients

984   with advanced and recurrent colorectal cancer. Br J Cancer [Internet].

985   2017;116(7):923–9. Available from: http://dx.doi.org/10.1038/bjc.2017.37

987   # Supporting Information

988   **S1 File. PRISMA Checklist**

989   **S2 File. Quality assessment of the selected studies for the systematic review**

Fig1

Fig2

Fig3