Cell Type-Specific Modulation of Layer 6A Excitatory Microcircuits by Acetylcholine in Rat Barrel Cortex

Danqing Yang[1], Robert Günter[2], Guanxiao Qi[1], Gabriele Radnikow[1] and Dirk Feldmeyer[1,3,4]*

[1]Institute of Neuroscience and Medicine, INM-10, Research Centre Jülich, D-52425 Jülich, Germany

[2]Institute of Neuroscience and Medicine, INM-2, Research Centre Jülich, D-52425 Jülich, Germany

[3]Department of Psychiatry, Psychotherapy and Psychosomatics, RWTH Aachen University, D-52074 Aachen, Germany.

[4]Jülich Aachen Research Alliance, Translational Brain Medicine (JARA Brain), D-52074 Aachen, Germany

*Corresponding author: Dirk Feldmeyer

Institute of Neuroscience and Medicine (INM-10) Research Centre Jülich D-52425 Jülich Germany

Tel.: +49-2461-61-5226

Fax: +49-2461-61-1778

E-mail: d.feldmeyer@fz-juelich.de

Abstract

Acetylcholine (ACh) is known to regulate cortical activity during different behavioral states, e.g.

wakefulness and attention. Here we show a differential expression of muscarinic ACh receptors

(mAChRs) and nicotinic AChRs (nAChRs) in different layer 6A (L6A) pyramidal cell (PC) types

of somatosensory cortex. At low concentrations, ACh induced a persistent hyperpolarization in

corticocortical (CC) but a depolarization in corticothalamic (CT) L6A PCs via M_4 and M_1 mAChRs,

respectively. At ~1 mM ACh depolarized exclusively CT PCs via $\alpha_4\beta_2$ subunit-containing nAChRs

without affecting CC PCs. Miniature EPSC frequency in CC PCs was decreased by ACh but

increased in CT PCs. In synaptic connections with a presynaptic CC PC, glutamate release was

suppressed via M_4 mAChR activation but enhanced by nAChRs via $\alpha_4\beta_2$ nAChRs when the

presynaptic neuron was a CT PC. Thus, in layer 6A the interaction of mAChRs and nAChRs results

in an altered excitability and synaptic release, effectively strengthening corticothalamic output

while weakening corticocortical synaptic signaling.

Keywords: barrel cortex, layer 6, pyramidal cells, acetylcholine, muscarinic receptors, nicotinic

receptors, corticocortical, corticothalamic

Introduction

Acetylcholine (ACh) has been shown to play a major role in memory processing, arousal, attention and sensory signaling[1-7]. It has been demonstrated that the ACh concentration in the cerebrospinal fluid increases during wakefulness and sustained attention[8,9]. In the neocortex release of ACh occurs predominately via afferents originating from cholinergic neurons in the nucleus basalis of Meynert of the basal forebrain[10-12]; their terminals are densely distributed throughout all neocortical layers[13-15]. A classical view is that ACh invariably increases the excitability of excitatory neurons in neocortex 16-20. However, a persistent hyperpolarization in layer 4 (L4) excitatory neurons was found in somatosensory cortex[21,22]. This layer-specific cholinergic modulation may contribute to improving the cortical signal-to-noise ratio[23-25].

Although extensive studies have been conducted on the cholinergic modulation of neocortical excitatory neurons, the action of ACh on the layer 6 (L6) microcircuitry has not been systematically investigated. Two main pyramidal cell (PC) classes exist in cortical layer 6, namely corticothalamic (CT) and corticocortical (CC) PCs. These two neuron types differ in their axonal projection patterns, dendritic morphological features, electrophysiological properties and expression of molecular markers[26-30]. CC PCs have no subcortical target and send intracortical projections mainly within the infra-granular layers[28]; CT PCs, in contrast, have few axons distributed in cortex and send projections directly back to the thalamus thereby contributing to a feedback control of sensory input[31-35]. The question how the function of these two classes of L6 PCs is modulated by ACh has so far not been explored.

Recent optogenetic studies suggest that PCs in L5 and L6 receive direct cholinergic inputs[20,36]. In these neurons, ACh induces a slowly desensitizing inward current in L6 PCs of prefrontal cortex through activation of $\alpha_4\beta_2$ subunit containing synaptic nicotinic acetylcholine receptors (nAChRs)[23,36-39]. However, there are very few studies focusing on the effects of muscarinic acetylcholine receptors (mAChRs) in L6A neurons[29,40]. Here, using single and paired patch-clamp recordings with simultaneous biocytin-filling, we investigated both muscarinic and nicotinic modulation of morphologically identified excitatory neurons and their synaptic connections in layer 6A of rat primary somatosensory barrel cortex. We found that ACh shows a cell type-specific effect on both cellular and synaptic properties in L6A excitatory microcircuits through activation of mAChRs and/or nAChRs. Our results reveal that two functionally and morphologically distinct subpopulations of L6A PCs, CC and CT PCs, are differentially modulated by ACh. We

70 demonstrate that ACh suppresses intracortical synaptic transmission via somatodendritic

71 hyperpolarization and inhibition of presynaptic neurotransmitter release of CC PCs by activating

72 M_4Rs. In contrast, CT PC show a dual cholinergic modulation: These neurons are depolarized via

73 M_1 mAChRs and $\alpha_4\beta_2$ subunit-containing nAChRs while the presynaptic release probability is

74 enhanced by $\alpha_4\beta_2$ nAChRs. In this way, ACh contributes to a facilitation of corticothalamic feedback.

Results

ACh either depolarizes or hyperpolarizes L6A PCs through activation of mAChRs

Whole-cell patch clamp recordings from L6A neurons were performed in acute brain slices of rat barrel cortex with simultaneous biocytin fillings. During recordings, excitatory neurons were distinguished from interneurons by their regular firing pattern with a low maximum firing frequency. Following bath application of 100 µM ACh, one subset of L6A PCs showed a membrane potential hyperpolarization by on average -2.0 ± 1.0 mV (n = 14), whereas another was depolarized by +9.5 ± 6.1 mV (n = 15; **Fig. 1a**). In addition, 1 s current pulses were injected in the recorded neuron to elicit AP firing before and during bath application of ACh. Under the supra-threshold stimulus (100pA above the rheobase current), the firing frequency was decreased by ACh in L6A PCs showing an ACh-induced hyperpolarization but increased in PCs that exhibit a depolarizing ACh response (**Fig. 1a, b**). Notably, both ACh-induced hyperpolarization and depolarization were not transient but persisted until the end of bath application. For L2/3 and L5 PCs it has been reported that the ACh-induced depolarization was preceded by an initial transient hyperpolarization mediated by 'small conductance', Ca^{2+}-activated K^+ channels[21,41-43]. We were able to reproduce this finding under the same recording condition; however, the de- and hyperpolarizing cholinergic response in L6A PCs induced by ACh puff application was always monophasic (**Supplementary Fig. 1**).

To determine which fraction of the membrane potential changes in L6A pyramidal neurons is mediated by mAChRs, 20 µM atropine (ATRO, a general mAChR antagonist) was applied in voltage-clamp mode. Both the ACh-induced outward and inward currents were found to be strongly blocked by ATRO (20 µM) (**Fig. 1 c, d**), suggesting that both ACh response types in L6A excitatory neurons are almost exclusively mediated by mAChRs. We hypothesized that the $G_{i/o}$ protein coupled M_4 mAChR subtype (M_4Rs) mediates the hyperpolarizing effects while the $G_{q/11}$ protein-coupled M_1 mAChR subtype (M_1Rs) is responsible for the depolarization induced by ACh application. To test this, puff application of ACh (100 µM) was performed in the presence and absence of the selective mAChR antagonists in the perfusion solution. In the presence of 1 µM tropicamide (TRO, a selective M_4R antagonist), the ACh-induced hyperpolarization was abolished (**Fig. 1e, g**). Conversely, the ACh-induced depolarization was blocked by 0.5 µM pirenzipine (PIR, a selective M_1R antagonist; **Fig. 1f, h**). These results indicate that the persistent hyperpolarization

and depolarization induced by low concentrations of ACh are mediated exclusively by M_4Rs and

M_1Rs, respectively.

The dose-dependence of the muscarinic effects was investigated by bath application of increasing

concentrations of ACh in the presence of 1 µM mecamylamine (MEC, a general nAChR antagonist)

and 0.5 µM tetrodotoxin (TTX) (0.3 µM to 300 µM; **Fig. 1i, j**). The dose-response curve was

obtained by fitting the data to the Hill equation. For hyperpolarizing L6A PCs, the ACh

concentration for a half-maximum response (EC_{50}) was 6.2 ± 1.3 µM while for depolarizing neurons,

the EC_{50} was 26.7 ± 5.4 µM. Thus, an ACh concentration of 30 µM was adopted for all subsequent

experiments; this concentration resulted in a >50% of the maximum response in both subgroups of

L6A excitatory neurons. In addition, when only 30 µM ACh was used neurons did not respond

with AP firing which was occasionally observed when applying 100 µM ACh.

Cholinergic responses in L6A PCs are cell-type specific

To investigate whether the two different cholinergic response types are specific for a defined

L6A pyramidal cell type, we characterized L6A PCs by their morphological, electrophysiological

and molecular features. Here, a total of 105 excitatory L6A neurons were recorded and

morphologically reconstructed. Previous studies have consistently shown that CC and CT L6A

PCs can be distinguished reliably by their axonal projection patterns[26,27,30,44]. Of all excitatory cells,

74 (70.5%) were identified as putative CT PCs while 31 (29.5%) were putative CC PCs. CC L6A

PCs displayed a dense horizontal axonal projection pattern in infragranular layers spanning

several neighboring barrel columns; CT L6A PCs, on the other hand, showed a sparse columnar

axonal domain with the majority of collaterals projecting directly towards the pia and terminating

predominately in layer 4 **(cf Fig. 2a left and right panels)**. CC PCs have a significantly larger

axonal (15523 ± 5013 µm vs. 5209 ± 1462 µm, P < 0.001) and dendritic length (5921 ± 1346 µm

vs. 5134 ± 1070 µm, P < 0.05) compared to CT PCs. Similar differences were also detected in the

horizontal axonal and dendritic field span (1714 ± 350 µm vs. 358 ± 111 µm, P < 0.001 and 361

± 58 µm vs. 232 ± 28 µm, P < 0.001, respectively, for CC vs. CT L6A PCs). For CC L6A PCs

these values are likely to be strong underestimates (by $\geq 90\%$, cf.[45]) because in acute slice

preparations their long-range axonal collaterals will be severely truncated; however, this does not

prevent an unambiguous cell type identification. In addition, CC L6A PCs have more first order

axon collaterals (p < 0.001) but fewer dendrites (p < 0.001) than CT PCs **(Fig. 2b)**. The features

described above are reflected in the polar plots **(Fig. 2a)**.

In addition, we determined the electrophysiological properties of morphologically identified CC (n

= 11) and CT (n = 9) L6A PCs. Compared to CT PCs, CC PCs showed a significantly lower R_{in} (P

< 0.05), a longer onset time (P < 0.01) for the first action potential (AP) evoked by injecting a

rheobase current and a longer AP half-width (P < 0.05). Trains of spikes were elicited to examine

the firing behavior. The AP adaptation ratio (2^{nd} ISI/10^{th} ISI) of CC PCs was smaller (P < 0.05) than

that of CT cells because they exhibited an initial spike burst. **(Supplementary Fig. 2).** The

differences in passive and active electrophysiological properties found here are in accordance with

previous studies[26,40].

Furthermore, the nuclear transcription factor Fork-head box protein P2 (FoxP2) is co-expressed

with the neurotensin receptor 1 (NtsR1) gene, a molecular marker for CT L6A PCs in mice[29,46]. To

identify the expression of FoxP2 in L6A PCs, we performed whole-cell recordings with

simultaneous filling of biocytin and fluorescent Alexa Fluor® 594 dye (n = 14). Subsequently,

brain slices were processed for FoxP2 immunofluorescence staining. We found that CT L6A PCs

were FoxP2-positive while CC PCs are FoxP2-negative **(Supplementary Fig. 3a,b)**. The tight

correlation between neuronal morphology, electrophysiology and FoxP2 expression demonstrates

the reliability of classification based on axonal projection patterns of CC and CT PCs.

ACh at a concentration of 30 μM was bath-applied to 63 morphological identified L6A neurons.

CC L6A PCs showed a hyperpolarizing response with a mean amplitude of -1.76 ± 4.28 mV (from

-61.9 ± 5.6 mV to -63.6 ± 7.0 mV, P < 0.05, n = 35). In contrast, ACh (30 μM) induced a strong

depolarization with a mean amplitude of +11.4 ± 4.6 mV (from -70.3 ± 5.1 mV to -58.8 ± 7.4 mV,

P < 0.001, n = 14) in CT PCs without exception (**Fig. 2c**). In **Fig. 2d**, ACh-induced membrane

potential changes are plotted against the horizontal axonal field span revealing a strong correlation

between axonal morphology and cholinergic response for the two L6A pyramidal cell types. In

addition, by performing immunostaining we confirmed that M_4Rs were enriched within L6A. We

found that M_4R-positive neurons were FoxP2-negative while virtually no FoxP2-positive neuron

expressed M_4Rs **(Supplementary Fig. 3d)**. This is consistent with our pharmacological result that

only FoxP2-negative CC PCs showed a M_4Rs-mediated hyperpolarization following ACh

application **(Supplementary Fig. 3)**.

CT PCs are selectively activated by high concentrations of ACh via α4β2 nAChRs

As demonstrated above, the depolarizing and hyperpolarizing effects of ACh in L6A PCs can be attributed to the activation of M_1 and M_4 mAChRs, respectively (**Fig. 1c, d**). However, previous studies have shown that ACh excites L6A excitatory neurons by activating nAChRs[23,36,37,39]. In order to investigate the functional role of nAChRs in L6A of rat barrel cortex, we perfused slices continuously with 200 nM ATRO. Under this condition, 30 µM ACh had no effect on both CC and CT PCs ($P > 0.05$ for CC cells, n = 5; $P > 0.05$ for CT cells, n = 5) (**Fig. 3a**); application of 1 mM ACh, however, strongly depolarized CT PCs ($P < 0.001$, n = 12) while CC PCs showed no response ($P > 0.05$, n = 4; **Fig. 3b**). Our results demonstrate that both the muscarinic and nicotinic modulation of L6A PCs is cell-type specific; nAChRs are present solely in CT L6A PCs and activated substantially only by high ACh concentrations. To determine the concentration range in which ACh activates postsynaptic nAChRs, we measured the dose-response curve for ACh in the presence of 200 nM ATRO. A fit of dose-response relationship to the Hill equation gave an EC_{50} of 1.2 ± 0.3 mM (n = 5) for the nicotinic ACh response (**Fig. 3c**), a value more than about two orders of magnitude larger than those of the de- and hyperpolarizing muscarinic response.

It has been reported that the expression of nAChR subtypes in the neocortex exhibits layer-specificity. L6A PCs in prefrontal cortex show a slow inward current to ACh by activating nAChRs containing the α_4 and β_2 subunits[23,36]. To confirm that this nAChR subtype mediates the response, the response of CT L6A PCs to application of 1 mM ACh was recorded in an ATRO-containing perfusion solution. In the presence of DHßE (10 µM), a nicotinic antagonist specific for $\alpha_4\beta_2$ nAChRs, the ACh-dependent depolarization in CT PCs was eliminated, suggesting that CT L6A PCs express postsynaptic $\alpha_4\beta_2$ subunit-containing nAChRs (**Fig. 3d**).

ACh differentially modulate miniature spontaneous activity of CC and CT L6A PCs

In addition to changing the membrane properties and excitability of neurons, ACh is also a powerful modulator of neurotransmitter release. Therefore, we measured the amplitude and frequency of miniature spontaneous activity by performing whole-cell voltage-clamp recordings from L6A PCs. The membrane potential was held at -70 mV and inward miniature EPSCs (mEPSCs) were recorded in the presence of TTX (0.5 µM) and gabazine (10 µM).

We found that ACh differentially modulates miniature spontaneous activity in both L6A CC and CT PCs. The frequency but not the amplitude of mEPSCs in CC L6A PCs was significantly

decreased by application of 30 µM ACh (2.8 ± 0.8 vs. 2.2 ± 1.0 Hz; n = 7, P < 0.01), an effect that

was blocked by the M_4Rs antagonist tropicamide (2.2 ± 1.0 vs. 2.8 ± 1.0 Hz; n = 7, P < 0.01) (**Fig.**

4a-c). This suggests that ACh decreases the neurotransmitter release probability at synapses with

CC L6A PCs via presynaptic M_4Rs. Similarly, when DHßE was co-applied with tropicamide and

ACh, a reduction of mEPSC frequency without a change in mEPSC amplitude was observed

(**Supplementary Fig. 4**). This implies that in addition to M_4Rs, $\alpha_4\beta_2$ nAChRs also play a role in

the cholinergic modulation of excitatory synaptic transmission onto CC PCs.

In contrast to CC L6A PCs, application of 30 µM ACh significantly decreased the inter-event

interval of mEPSCs in CT PCs, reflecting an increase in mEPSC frequency (0.95 ± 0.36 vs. 2.12 ±

0.68 Hz; n = 7, P < 0.01) while the mEPSc amplitude remained unaffected. Because ATRO did not

affect the mEPSC frequency, we argued that the cholinergic effects on spontaneous mEPSCs in

CT L6A PCs were not mediated by mAChRs but exclusively by presynaptic nAChRs. To test this,

10 µM DHßE was co-applied with ACh. In the presence of DHßE, the ACh-induced increase of

mEPSCs frequency in CT L6A PCs was reduced to control level (2.12 ± 0.68 vs. 1.01 ± 0.55; n =

7, P < 0.05; **Fig. 4d-f**). These results suggest that ACh potentiates excitatory synaptic transmission

onto L6A CT PCs exclusively via presynaptic $\alpha_4\beta_2$ subunit containing nAChRs.

ACh induces a reduction of presynaptic neurotransmitter release in CC L6A PCs but an an

increase in CT PCs

To elucidate cholinergic effects on L6A PCs at pre- and postsynaptic sites independently, paired

recordings and simultaneous biocytin fillings of synaptically coupled L6A neurons were performed.

Excitatory neurons were classified as either CT and CC PCs based on the criteria mentioned above.

During recordings, inhibitory interneurons were preliminarily identified by their high frequency

AP firing pattern. After reconstructions, interneurons were further distinguished based on

morphological features such as lack of dendritic spines. 34 excitatory connections were established

by presynaptic CC PCs. We found that ACh suppresses the synaptic efficacy of neuronal

connections established by presynaptic CC PCs regardless of the postsynaptic neuron type **(Fig. 5)**.

The unitary EPSP (uEPSP) amplitude of all synaptic connections with a presynaptic L6A CC PC

were all significantly decreased by ACh (30 µM). For CC-CC connections the uEPSP amplitude

decreased from 0.45 ± 0.32 mV to 0.19 ± 0.14 mV (n = 20 pairs, P < 0.001) and for CC-CT

connections changed from 0.35 ± 0.22 mV to 0.19 ± 0.13 mV (n = 5 pairs, P < 0.05). For CC-

interneuron connections the mean uEPSP was reduced from 0.90 ± 0.90 mV to 0.52 ± 0.58 mV (n

= 9 pairs, P < 0.05) in the presence of ACh. ACh also significantly increased the paired-pulse ratio

(PPR) of CC-CC (1.0 ± 0.4 vs. 1.5 ± 0.7, P < 0.01), CC-CT (1.2 ± 0.7 vs. 2.1 ± 1.6, P < 0.05) and

CC-interneuron (1.0 ± 0.5 vs. 1.2 ± 0.5, P < 0.01) connections. Following ACh application, CC-

CC connections and CC-interneuron connections showed an increase in the CV; at CC-interneuron

connections the failure rate was also significantly increased **(Fig. 5 e; Supplementary Tab. 1)**.

These changes in the EPSP properties suggest that ACh decreases the neurotransmitter release

probability of intra-laminar connections established by a presynaptic L6A CC PC. Other synaptic

properties, like rise time, latency and decay time, were not affected by ACh **(Supplementary Tab.**

1).

Previous studies have shown that ACh may inhibits intracortical excitatory synaptic transmission

at some synaptic connections through activation of presynaptic M_4 mAChRs[21,47,48]. To test whether

the ACh-induced suppression of the efficacy of synaptic connections with a presynaptic CC L6A

PC is mediated by M_4 mAChR activation, 1 µM TRO (a selective antagonist of M_4Rs) was co-

applied with ACh (30 µM) after bath application of ACh alone. The effects of ACh on synaptic

connections established by CC PCs (n = 6 pairs, comprising 2 CC-CC, 1 CC-CT and 3 CC-

interneuron connections) were completely blocked by TRO. The EPSP amplitude decreased from

1.0 ± 0.8 mV to 0.3 ± 0.2 mV during ACh application and fully recovered to 1.0 ± 0.8 mV during

co-application of ACh and TRO. Moreover, TRO also blocked the ACh effects on the CV ($0.8 \pm$

0.2 for control vs. 0.9 ± 0.3 for ACh and TRO; n = 6 pairs, P = 0.75) and failure rate ($27.0 \pm$

17.6 % for control vs. 26.3 ± 19.6 % for ACh and TRO; n = 6 pairs, P = 1.00) **(Fig. 6)**. In

addition to reversing the ACh-induced increase in the PPR, TRO increased the PPR of

connections established by CC PCs. Co-application of ACh and TRO resulted in a smaller PPR

compared to control (1.1 ± 0.3 vs. 1.3 ± 0.4; n = 6 pairs, P < 0.05) **(Fig. 6)**. In order to isolate the

presynaptic effect of ACh on L6A intra-laminar connections established by presynaptic CC PCs,

a CC-CT synaptically coupled pair was recorded. The ACh-induced reduction in synaptic release

probability recovered only after co-application of TRO together with PIR but not when PIR (0.5

µM) was applied alone **(Supplementary Fig. 5)**.

Because of their sparse and narrow axonal domain, L6A CT PCs rarely innervate neurons in their

home layer and their intracortical synaptic connections are remarkably weak and unreliable[44,49,50].

Here we applied ACh (30 µM) to seven synaptic connections established by a presynaptic CT PC

in L6A, including two CT-CT, one CT-CC and four CT-interneuron connections. In all synaptic

connections established by CT PCs ACh significantly enhanced the EPSP amplitude (0.10 ± 0.08

mV vs. 0.15 ± 0.10 mV; n = 7 pairs, $P < 0.05$) and reduced the PPR (3.0 ± 1.7 vs. 0.8 ± 0.6; n = 7

pairs, $P < 0.05$) **(Fig. 7)**. The ACh-mediated reduction in the PPR suggests a presynaptic locus for

synaptic modulation. Because these connections display very small unitary EPSP amplitudes and

frequent failures, the signal-to-noise ratio was often too low to appropriately calculate the CV and

failure rate. No significant differences were detected in rise time, decay time and latency

(Supplementary Tab. 1). Our findings indicate that in contrast to the inhibition of presynaptic

release in L6A CC PCs, ACh enhances the synaptic efficacy of the weak connections established

by a presynaptic CT PC.

To determine the AChR subtype that mediates the increase in synaptic efficacy at these connections,

we tested whether the selective antagonists of the M_1Rs (PIR), homomeric the α_7 subunit-containing

nAChRs (methyllycaconitine, MLA) and the heteromeric $\alpha_4\beta_2$ subunit-containing nAChRs (DHßE)

could block the effect of ACh on synaptic connections with a presynaptic CT PC. While PIR and

MLA had no effect, DHßE blocked the increase of EPSP amplitude and decrease of PPR **(Fig. 8)**.

This indicates that the ACh-induced enhancement of synaptic efficacy is induced by activation of

$\alpha_4\beta_2$ subunit-containing nAChRs in presynaptic CT PCs.

Discussion

We investigated the cholinergic modulation of CT and CC PCs in layer 6A of the barrel cortex. We showed that (i) low concentrations of ACh differentially modulate the L6A microcircuitry by persistently depolarizing CT but hyperpolarizing CC L6A PCs. These effects are monophasic and mediated via M_1 and M_4 mAChRs, respectively; (ii) a nicotinic ACh response was observed exclusively in CT PCs only when a high ACh concentration was applied. In addition, (iii) low concentrations of ACh increases the frequency of miniature EPSCs via presynaptic nAChRs in L6A CT but decreases that of CC PCs via M_4Rs. To better understand the effects of ACh on intralaminar synaptic transmission, recordings were performed from synaptically coupled L6A PC pairs. We found that (iv) in neuronal connections with a presynaptic CC PC the neurotransmitter release probability was reduced via activation of M_4Rs but (v) increased in connections with a presynaptic CT L6A neuron by $\alpha_4\beta_2$ nAChR activation. Our results reveal that two functionally and morphologically distinct subpopulations of L6A PCs are affected differentially by ACh acting on both mAChRs and nAChRs.

Synergistic modulation of L6A PCs by mAChRs and nAChRs

In a number of studies investigating the nAChR response of L6 PCs in different cortical areas only high ACh concentration ($\geq$ 1 mM) have been applied because the ACh affinity of nAChRs is substantially lower than that of mAChRs[23,36,37,39]. Under this condition any mAChR effect is almost entirely masked by the strong nicotinic response so that any involvement of mAChRs has been explicitly ruled out. Here we demonstrate for the first time that mAChRs play crucial roles in both pre- and postsynaptic modulation of L6A PC activity. Both the pre- and postsynaptic effects of mAChRs are already present at low ACh concentrations (1-10 µM) suggesting that neuromodulation via mAChRs is tonically present and mediated by volume transmission[51,52]. On the other hand, only a high concentration of ACh (EC_{50} of ~1 mM) could effectively depolarize CT L6A PCs via nAChRs, but an up-regulation of presynaptic vesicle release via $\alpha_4\beta_2$ subunit-containing nAChRs was already observed in the presence of 30 µM ACh. This $\alpha_4\beta_2$ nAChR-mediated effect was found for spontaneous excitatory synaptic activity as well as for CT-formed monosynaptic connections, suggesting that presynaptic nAChR are expressed on synaptic boutons of both CT L6A axons and other glutamatergic afferents (e.g. from thalamus or other cortical regions). Single channel currents from nAChRs can be activated already by low concentrations of

ACh[53,54]. Because of the electrotonic compact structure of a presynaptic bouton even the opening of
a few nAChR receptor channels may result in a depolarisation that is sufficiently strong to increase
the open probability of presynaptic Ca^{2+} channels and hence the neurotransmitter release.

Furthermore, in some L6 PCs the α_5 nAChR subunit co-assembles with the α_4 and β_2 subunits[23,36].
Nicotinic AChRs containing the α_4, β_2 and α_5 subunits have a higher Ca^{2+} permeability than those
composed of α_4 and β_2 nAChR subunits alone[55]. In the presynaptic terminals, Ca^{2+} entry via $\alpha_4\beta_2\alpha_5$
nAChRs into the presynaptic bouton could enhance neurotransmitter release provided these
channels are located sufficiently close to the release site.

Cholinergic activation by endogenous ACh in neocortical L6

In the cortex, ACh levels change dramatically during different stages of waking and sleep[8,9]. It has
been suggested that high ACh levels serve to enhance the response to sensory stimuli by increasing
the strength of afferent input while low concentration of ACh contributes in the consolidation of
encoded information[56]. Cholinergic signalling has been described to occur via a volume release
mechanism[52], which is slow and unspecific. Volume release of ACh reaches concentrations in a low
micromolar range, spreads widely over neocortical layers and activates predominantly mAChRs.
In addition, cholinergic synapses have been identified particularly in deep layers of neocortex and
less so in superficial cortical layers[20,36,57]. At these cholinergic synapses, ACh reaches a high
concentration in the synaptic cleft, thereby activating postsynaptic nAChRs in L6 PCs[36]. Because
cholinergic synapses in the neocortex are small[58], ACh released into the synaptic cleft may spill
over into the perisynaptic space. Subsequently, the extra-synaptic AChRs on presynaptic boutons
of CT PCs are activated resulting in an increase of release probability (**Fig. 7**). Thus, nAChRs and
mAChRs act on different time scales and at different neurotransmitter concentrations, resulting in
a striking complexity of the cholinergic modulation of neocortical signaling.

The cell type-specific effect of ACh in L6A

ACh has been shown to induce a persistent depolarization of L2/3 and L5 PCs but a
hyperpolarization of excitatory L4 neurons[21,43,59]. Here, we demonstrate that ACh modulates PCs not
only in a layer-specific but also a cell type-specific way that can be attributed to a cell type-
dependent expression of mAChRs (**Fig. 9**). In L6A of barrel cortex, ACh hyperpolarizes CC PCs
but depolarizes CT PCs via activation of M_4Rs and M_1Rs, respectively. The action potential firing

frequency was decreased by ACh in CC PCs but increased in CT PCs, thereby modulating the

excitability and signal propagation in L6A PCs in a cell-specific manner. In addition, CT L6A PCs

but not CC PCs showed a strong $\alpha_4\beta_2$ nAChR-mediated response (**Fig. 9**). This is consistent with

previous findings in L6 of prefrontal cortex that regular spiking neurons have a larger nicotinic

receptor-mediated inward current following ACh application when compared with bursting

neurons[37]. A cell type-specific neuromodulation was also discovered previously in deep layers of

medial prefrontal cortex for neuromodulators such as noradrenaline, dopamine and adenosine[60-64].

By studying miniature spontaneous activity of L6A CC and CT PCs, we found that ACh both

increases the excitatory synaptic release onto CC and CT PCs by activation of $\alpha_4\beta_2$ nAChRs (**Fig.**

9). Because CT PCs express $\alpha_4\beta_2$ nAChRs, an increase in spontaneous activity may result from the

enhanced release probability at CT L6 PC boutons; however, the intracortical axon density of these

PCs is low so that their contribution to the spontaneous mEPSC frequency is minimal. On the other

hand, activation of nAChRs increases thalamocortical transmission onto L3, L4 and L5 neocortical

neurons[47,65,66]. Thus, the increased excitatory transmission onto L6A PCs is probably resulting to a

large degree from a higher release probability at thalamocortical and less so from intracortical

synapses. In addition, CC PCs receive more intra-laminar inputs than CT PCs, which can be

suppressed by ACh via M_4Rs. Therefore, the ACh-induced reduction of mEPSC frequency in CC

PCs could be a combinatorial effect on thalamocortical and intracortical transmission.

It has been proposed that ACh increases the signal-to-noise ratio (SNR) of sensory signaling by

selectively enhancing thalamocortical inputs over intracortical synaptic transmission[47,67,68]. ACh has

been found to suppress the efficacy of excitatory intracortical connections in different layers

including L2/3, L4 and L5[21,47,48]. Here, a differential cholinergic modulation of presynaptic

neurotransmitter release was observed in CC and CT L6A PC types. ACh suppresses synaptic

transmission in excitatory L6A connections with presynaptic CC PCs through activation of M_4Rs

but potentiates connections with a presynaptic CT PCs via presynaptic $\alpha\beta_2$ nAChRs (**Fig. 9**); no α_7

nAChR or M_1R effect on synaptic transmission was observed.

In hippocampus and some subcortical structures such as the ventral tegmental area, glutamatergic

synapses are known to be facilitated by nAChRs located on presynaptic terminals[69,70]. However,

very few studies demonstrate an ACh-mediated enhancement of intracortical excitatory synaptic

transmission. Recently, it has been shown that excitatory synaptic transmission between PCs and

somatostatin-positive interneurons in layer 2 of mouse barrel cortex is increased by ACh via

activating nAChRs[71]. Although an increase of synaptic efficacy was observed in connections with a presynaptic CT PCs, this type of synaptic connections are rare, generally weak and very unreliable[49,50]. Therefore we propose that ACh mainly acts on CT PCs not primarily by increasing intracortical synaptic transmission but rather by facilitating corticothalamocortical feedback; this facilitation will occur already at ACh levels in the low micromolar range.

Acknowledgement

We thank Werner Hucko for excellent technical assistance and Dr. Dr. Karlijn van Aerde for custom-written macros in Igor Pro software. We warmly thank Dr. Chao Ding for helpful discussions and Dr. Vishalini Emmenegger for proofreading the manuscript.

Funding

This work was supported by the Helmholtz Society, the DFG Research Group - BaCoFun (grant no. Fe471/4-2 to D.F.), the European Union's Horizon 2020 Research, Innovation Programme under Grant Agreement No. 785907 (HBP SGA2; to DF) and the China Scholarship Council (to D.Y.).

Author contributions

D.F., G.R. and D.Y. designed the research; D.Y., R.G. and G.Q. performed experiments and data analysis; D.Y., D.F. and G.R. wrote the manuscript. The authors declare no conflicts of interest.

References

Jones, B. E. Activity, modulation and role of basal forebrain cholinergic neurons innervating the cerebral cortex. *Prog Brain Res* **145**, 157-169, doi:10.1016/S0079-6123(03)45011-5 (2004).

Hasselmo, M. E. The role of acetylcholine in learning and memory. *Curr Opin Neurobiol* **16**, 710-715, doi:10.1016/j.conb.2006.09.002 (2006).

Herrero, J. L. *et al.* Acetylcholine contributes through muscarinic receptors to attentional modulation in V1. *Nature* **454**, 1110-1114, doi:nature07141 [pii] 10.1038/nature07141 (2008).

Hasselmo, M. E. & Sarter, M. Modes and models of forebrain cholinergic neuromodulation of cognition. *Neuropsychopharmacology* **36**, 52-73, doi:10.1038/npp.2010.104 (2011).

Thiele, A. Muscarinic signaling in the brain. *Annu Rev Neurosci* **36**, 271-294, doi:10.1146/annurev-neuro-062012-170433 (2013).

Wester, J. C. & Contreras, D. Differential modulation of spontaneous and evoked thalamocortical network activity by acetylcholine level in vitro. *J Neurosci* **33**, 17951-17966, doi:10.1523/JNEUROSCI.1644-13.2013 (2013).

Ma, S., Hangya, B., Leonard, C. S., Wisden, W. & Gundlach, A. L. Dual-transmitter systems regulating arousal, attention, learning and memory. *Neurosci Biobehav Rev* **85**, 21-33, doi:10.1016/j.neubiorev.2017.07.009 (2018).

Himmelheber, A. M., Sarter, M. & Bruno, J. P. Increases in cortical acetylcholine release during sustained attention performance in rats. *Brain Res Cogn Brain Res* **9**, 313-325 (2000).

Teles-Grilo Ruivo, L. M. *et al.* Coordinated Acetylcholine Release in Prefrontal Cortex and Hippocampus Is Associated with Arousal and Reward on Distinct Timescales. *Cell Rep* **18**, 905-917, doi:10.1016/j.celrep.2016.12.085 (2017).

Mesulam, M. M., Mufson, E. J., Wainer, B. H. & Levey, A. I. Central cholinergic pathways in the rat: an overview based on an alternative nomenclature (Ch1-Ch6). *Neuroscience* **10**, 1185-1201 (1983).

Zaborszky, L. *et al.* Neurons in the basal forebrain project to the cortex in a complex topographic organization that reflects corticocortical connectivity patterns: an experimental study based on retrograde tracing and 3D reconstruction. *Cereb Cortex* **25**, 118-137, doi:10.1093/cercor/bht210 (2015).

Paul, S., Jeon, W. K., Bizon, J. L. & Han, J. S. Interaction of basal forebrain cholinergic neurons with the glucocorticoid system in stress regulation and cognitive impairment. *Front Aging Neurosci* **7**, 43, doi:10.3389/fnagi.2015.00043 (2015).

Eckenstein, F. P., Baughman, R. W. & Quinn, J. An anatomical study of cholinergic innervation in rat cerebral cortex. *Neuroscience* **25**, 457-474 (1988).

14 Henny, P. & Jones, B. E. Projections from basal forebrain to prefrontal cortex comprise cholinergic,
GABAergic and glutamatergic inputs to pyramidal cells or interneurons. *Eur J Neurosci* **27**, 654-670,
doi:10.1111/j.1460-9568.2008.06029.x (2008).

15 Kalmbach, A., Hedrick, T. & Waters, J. Selective optogenetic stimulation of cholinergic axons in
neocortex. *J Neurophysiol* **107**, 2008-2019, doi:10.1152/jn.00870.2011 (2012).

16 McCormick, D. A. & Prince, D. A. Two types of muscarinic response to acetylcholine in mammalian
cortical neurons. *Proc Natl Acad Sci U S A* **82**, 6344-6348, doi:10.1073/pnas.82.18.6344 (1985).

17 Mednikova, Y. S., Karnup, S. V. & Loseva, E. V. Cholinergic excitation of dendrites in neocortical
neurons. *Neuroscience* **87**, 783-796 (1998).

18 Desai, N. S. & Walcott, E. C. Synaptic bombardment modulates muscarinic effects in forelimb motor
cortex. *J Neurosci* **26**, 2215-2226, doi:10.1523/JNEUROSCI.4310-05.2006 (2006).

19 Zhang, Z. & Seguela, P. Metabotropic induction of persistent activity in layers II/III of anterior cingulate
cortex. *Cereb Cortex* **20**, 2948-2957, doi:10.1093/cercor/bhq043 (2010).

20 Hedrick, T. & Waters, J. Acetylcholine excites neocortical pyramidal neurons via nicotinic receptors. *J
Neurophysiol* **113**, 2195-2209, doi:10.1152/jn.00716.2014 (2015).

21 Eggermann, E. & Feldmeyer, D. Cholinergic filtering in the recurrent excitatory microcircuit of cortical
layer 4. *Proc Natl Acad Sci U S A* **106**, 11753-11758, doi:10.1073/pnas.0810062106 (2009).

22 Dasgupta, R., Seibt, F. & Beierlein, M. Synaptic Release of Acetylcholine Rapidly Suppresses Cortical
Activity by Recruiting Muscarinic Receptors in Layer 4. *J Neurosci* **38**, 5338-5350,
doi:10.1523/JNEUROSCI.0566-18.2018 (2018).

23 Poorthuis, R. B. *et al.* Layer-specific modulation of the prefrontal cortex by nicotinic acetylcholine
receptors. *Cereb Cortex* **23**, 148-161, doi:10.1093/cercor/bhr390 (2013).

24 Obermayer, J., Verhoog, M. B., Luchicchi, A. & Mansvelder, H. D. Cholinergic Modulation of Cortical
Microcircuits Is Layer-Specific: Evidence from Rodent, Monkey and Human Brain. *Front Neural
Circuits* **11**, 100, doi:10.3389/fncir.2017.00100 (2017).

25 Radnikow, G. & Feldmeyer, D. Layer- and Cell Type-Specific Modulation of Excitatory Neuronal
Activity in the Neocortex. *Front Neuroanat* **12**, 1, doi:10.3389/fnana.2018.00001 (2018).

26 Kumar, P. & Ohana, O. Inter- and intralaminar subcircuits of excitatory and inhibitory neurons in layer
6a of the rat barrel cortex. *J Neurophysiol* **100**, 1909-1922, doi:10.1152/jn.90684.2008 (2008).

27 Zhang, Z. W. & Deschenes, M. Intracortical axonal projections of lamina VI cells of the primary
somatosensory cortex in the rat: a single-cell labeling study. *J Neurosci* **17**, 6365-6379 (1997).

28 Thomson, A. M. Neocortical layer 6, a review. *Front Neuroanat* **4**, 13, doi:10.3389/fnana.2010.00013
(2010).

29 Sundberg, S. C., Lindstrom, S. H., Sanchez, G. M. & Granseth, B. Cre-expressing neurons in visual
cortex of Ntsr1-Cre GN220 mice are corticothalamic and are depolarized by acetylcholine. *J Comp*
Neurol **526**, 120-132, doi:10.1002/cne.24323 (2018).

30 Pichon, F., Nikonenko, I., Kraftsik, R. & Welker, E. Intracortical connectivity of layer VI pyramidal
neurons in the somatosensory cortex of normal and barrelless mice. *Eur J Neurosci* **35**, 855-869,
doi:10.1111/j.1460-9568.2012.08011.x (2012).

31 Beierlein, M. & Connors, B. W. Short-term dynamics of thalamocortical and intracortical synapses onto
layer 6 neurons in neocortex. *J Neurophysiol* **88**, 1924-1932 (2002).

32 Yang, Q. *et al.* Intrinsic properties of and thalamocortical inputs onto identified corticothalamic-VPM
neurons. *Somatosens Mot Res* **31**, 78-93, doi:10.3109/08990220.2013.869495 (2014).

33 Oberlaender, M. *et al.* Cell type-specific three-dimensional structure of thalamocortical circuits in a
column of rat vibrissal cortex. *Cereb Cortex* **22**, 2375-2391, doi:10.1093/cercor/bhr317 (2012).

34 Constantinople, C. M. & Bruno, R. M. Deep cortical layers are activated directly by thalamus. *Science*
340, 1591-1594, doi:10.1126/science.1236425 (2013).

35 Lubke, J. & Feldmeyer, D. Excitatory signal flow and connectivity in a cortical column: focus on barrel
cortex. *Brain Struct Funct* **212**, 3-17, doi:10.1007/s00429-007-0144-2 (2007).

36 Hay, Y. A., Lambolez, B. & Tricoire, L. Nicotinic Transmission onto Layer 6 Cortical Neurons Relies
on Synaptic Activation of Non-alpha7 Receptors. *Cereb Cortex* **26**, 2549-2562,
doi:10.1093/cercor/bhv085 (2016).

37 Kassam, S. M., Herman, P. M., Goodfellow, N. M., Alves, N. C. & Lambe, E. K. Developmental
excitation of corticothalamic neurons by nicotinic acetylcholine receptors. *J Neurosci* **28**, 8756-8764,
doi:10.1523/JNEUROSCI.2645-08.2008 (2008).

38 Alves, N. C., Bailey, C. D., Nashmi, R. & Lambe, E. K. Developmental sex differences in nicotinic
currents of prefrontal layer VI neurons in mice and rats. *PLoS One* **5**, e9261,
doi:10.1371/journal.pone.0009261 (2010).

39 Bailey, C. D., Alves, N. C., Nashmi, R., De Biasi, M. & Lambe, E. K. Nicotinic alpha5 subunits drive
developmental changes in the activation and morphology of prefrontal cortex layer VI neurons. *Biol*
Psychiatry **71**, 120-128, doi:10.1016/j.biopsych.2011.09.011 (2012).

40 Tian, M. K., Bailey, C. D. & Lambe, E. K. Cholinergic excitation in mouse primary vs. associative cortex:
region-specific magnitude and receptor balance. *Eur J Neurosci* **40**, 2608-2618, doi:10.1111/ejn.12622
(2014).

41 Gulledge, A. T. & Stuart, G. J. Cholinergic inhibition of neocortical pyramidal neurons. *J Neurosci* **25**,
10308-10320, doi:10.1523/JNEUROSCI.2697-05.2005 (2005).

42 Gulledge, A. T. & Kawaguchi, Y. Phasic cholinergic signaling in the hippocampus: functional homology
with the neocortex? *Hippocampus* **17**, 327-332, doi:10.1002/hipo.20279 (2007).

43 Dasari, S., Hill, C. & Gulledge, A. T. A unifying hypothesis for M1 muscarinic receptor signalling in
pyramidal neurons. *J. Physiol.* **595**, 1711-1723, doi:10.1113/JP273627 (2017).

44 Mercer, A. *et al.* Excitatory connections made by presynaptic cortico-cortical pyramidal cells in layer 6
of the neocortex. *Cereb Cortex* **15**, 1485-1496, doi:10.1093/cercor/bhi027 (2005).

45 Narayanan, R. T. *et al.* Beyond Columnar Organization: Cell Type- and Target Layer-Specific Principles
of Horizontal Axon Projection Patterns in Rat Vibrissal Cortex. *Cereb. Cortex* **25**, 4450-4468,
doi:10.1093/cercor/bhv053 (2015).

46 Tasic, B. *et al.* Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat Neurosci*
19, 335-346, doi:10.1038/nn.4216 (2016).

47 Gil, Z., Connors, B. W. & Amitai, Y. Differential regulation of neocortical synapses by neuromodulators
and activity. *Neuron* **19**, 679-686 (1997).

48 Levy, R. B., Reyes, A. D. & Aoki, C. Nicotinic and muscarinic reduction of unitary excitatory
postsynaptic potentials in sensory cortex; dual intracellular recording in vitro. *J Neurophysiol* **95**, 2155-
2166, doi:10.1152/jn.00603.2005 (2006).

49 West, D. C., Mercer, A., Kirchhecker, S., Morris, O. T. & Thomson, A. M. Layer 6 cortico-thalamic
pyramidal cells preferentially innervate interneurons and generate facilitating EPSPs. *Cereb Cortex* **16**,
200-211, doi:10.1093/cercor/bhi098 (2006).

50 Crandall, S. R., Patrick, S. L., Cruikshank, S. J. & Connors, B. W. Infrabarrels Are Layer 6 Circuit
Modules in the Barrel Cortex that Link Long-Range Inputs and Outputs. *Cell Rep* **21**, 3065-3078,
doi:10.1016/j.celrep.2017.11.049 (2017).

51 Parikh, V., Kozak, R., Martinez, V. & Sarter, M. Prefrontal acetylcholine release controls cue detection
on multiple timescales. *Neuron* **56**, 141-154, doi:10.1016/j.neuron.2007.08.025 (2007).

52 Sarter, M., Parikh, V. & Howe, W. M. Phasic acetylcholine release and the volume transmission
hypothesis: time to move on. *Nat Rev Neurosci* **10**, 383-390, doi:10.1038/nrn2635 (2009).

53 Colquhoun, D. & Sakmann, B. Fast events in single-channel currents activated by acetylcholine and its
analogues at the frog muscle end-plate. *J Physiol* **369**, 501-557, doi:10.1113/jphysiol.1985.sp015912
(1985).

54 Mazzaferro, S., Bermudez, I. & Sine, S. M. alpha4beta2 Nicotinic acetylcholine receptors: Relationships
between subunit stoichiometry and function at the single chanel level. *J Biol Chem* **292**, 2729-2740,
doi:10.1074/jbc.M116.764183 (2017).

55 Fucile, S. Ca2+ permeability of nicotinic acetylcholine receptors. *Cell Calcium* **35**, 1-8 (2004).

56 Hasselmo, M. E. & McGaughy, J. High acetylcholine levels set circuit dynamics for attention and
encoding and low acetylcholine levels set dynamics for consolidation. *Prog Brain Res* **145**, 207-231,
doi:10.1016/S0079-6123(03)45015-2 (2004).

57 Bennett, C., Arroyo, S., Berns, D. & Hestrin, S. Mechanisms generating dual-component nicotinic
EPSCs in cortical interneurons. *J Neurosci* **32**, 17287-17296, doi:10.1523/JNEUROSCI.3565-12.2012
(2012).

58 Takacs, V. T., Freund, T. F. & Nyiri, G. Neuroligin 2 is expressed in synapses established by cholinergic
cells in the mouse brain. *PLoS One* **8**, e72450, doi:10.1371/journal.pone.0072450 (2013).

59 Gulledge, A. T., Park, S. B., Kawaguchi, Y. & Stuart, G. J. Heterogeneity of phasic cholinergic signaling
in neocortical neurons. *J Neurophysiol* **97**, 2215-2229, doi:10.1152/jn.00493.2006 (2007).

60 Dembrow, N. C., Chitwood, R. A. & Johnston, D. Projection-specific neuromodulation of medial
prefrontal cortex neurons. *The Journal of neuroscience : the official journal of the Society for
Neuroscience* **30**, 16922-16937, doi:10.1523/JNEUROSCI.3644-10.2010 (2010).

61 van Aerde, K. I., Qi, G. & Feldmeyer, D. Cell Type-Specific Effects of Adenosine on Cortical Neurons.
Cerebral cortex, doi:10.1093/cercor/bht274 (2013).

62 Clarkson, R. L., Liptak, A. T., Gee, S. M., Sohal, V. S. & Bender, K. J. D3 Receptors Regulate
Excitability in a Unique Class of Prefrontal Pyramidal Cells. *J Neurosci* **37**, 5846-5860,
doi:10.1523/JNEUROSCI.0310-17.2017 (2017).

63 Anastasiades, P. G., Boada, C. & Carter, A. G. Cell-Type-Specific D1 Dopamine Receptor Modulation
of Projection Neurons and Interneurons in the Prefrontal Cortex. *Cereb Cortex*,
doi:10.1093/cercor/bhy299 (2018).

64 Baker, A. L., O'Toole, R. J. & Gulledge, A. T. Preferential cholinergic excitation of corticopontine
neurons. *J Physiol* **596**, 1659-1679, doi:10.1113/JP275194 (2018).

65 Lambe, E. K., Picciotto, M. R. & Aghajanian, G. K. Nicotine induces glutamate release from
thalamocortical terminals in prefrontal cortex. *Neuropsychopharmacology* **28**, 216-225,
doi:10.1038/sj.npp.1300032 (2003).

66 Kawai, H., Lazar, R. & Metherate, R. Nicotinic control of axon excitability regulates thalamocortical
transmission. *Nat Neurosci* **10**, 1168-1175, doi:10.1038/nn1956 (2007).

67 Hsieh, C. Y., Cruikshank, S. J. & Metherate, R. Differential modulation of auditory thalamocortical and
intracortical synaptic transmission by cholinergic agonist. *Brain Res* **880**, 51-64 (2000).

68 Oldford, E. & Castro-Alamancos, M. A. Input-specific effects of acetylcholine on sensory and
intracortical evoked responses in the "barrel cortex" in vivo. *Neuroscience* **117**, 769-778 (2003).

69 Gray, R., Rajan, A. S., Radcliffe, K. A., Yakehiro, M. & Dani, J. A. Hippocampal synaptic transmission
enhanced by low concentrations of nicotine. *Nature* **383**, 713-716, doi:10.1038/383713a0 (1996).

70 Mansvelder, H. D. & McGehee, D. S. Long-term potentiation of excitatory inputs to brain reward areas
by nicotine. *Neuron* **27**, 349-357 (2000).

71 Urban-Ciecko, J., Jouhanneau, J. S., Myal, S. E., Poulet, J. F. A. & Barth, A. L. Precisely Timed Nicotinic
Activation Drives SST Inhibition in Neocortical Circuits. *Neuron* **97**, 611-625 e615,
doi:10.1016/j.neuron.2018.01.037 (2018).

72 Agmon, A. & Connors, B. W. Thalamocortical responses of mouse somatosensory (barrel) cortex in
vitro. *Neuroscience* **41**, 365-379 (1991).

73 Marx, M., Gunter, R. H., Hucko, W., Radnikow, G. & Feldmeyer, D. Improved biocytin labeling and
neuronal 3D reconstruction. *Nat Protoc* **7**, 394-407, doi:10.1038/nprot.2011.449 (2012).

74 Qi, G. X., Radnikow, G. & Feldmeyer, D. Electrophysiological and Morphological Characterization of
Neuronal Microcircuits in Acute Brain Slices Using Paired Patch-Clamp Recordings. *Jove-J Vis Exp*,
doi:ARTN e52358 10.3791/52358 (2015).

75 Dodt, H. U. & Zieglgansberger, W. Visualizing unstained neurons in living brain slices by infrared DIC-
videomicroscopy. *Brain Res* **537**, 333-336 (1990).

76 Meyer, H. S. *et al.* Number and laminar distribution of neurons in a thalamocortical projection column
of rat vibrissal cortex. *Cereb Cortex* **20**, 2277-2286, doi:10.1093/cercor/bhq067 (2010).

77 Feldmeyer, D., Egger, V., Lubke, J. & Sakmann, B. Reliable synaptic connections between pairs of
excitatory layer 4 neurones within a single 'barrel' of developing rat somatosensory cortex. *J Physiol* **521
Pt 1**, 169-190 (1999).

Materials and Methods

Slice preparation and solutions for electrophysiology

All experiments involving animals were performed in accordance with the EU Directive 2010/63/EU, the German animal welfare act and the guidelines of the Federation of European Laboratory Animal Science Association (FELASA). Wistar rats (Charles River) were maintained on a 12/12-hr light-dark cycle from 7 AM to 7 PM. Rats aged 17-21 postnatal days (P17-21, both sexes) were lightly anaesthetized with a concentration < 0.1% of isoflurane and then decapitated. The brain was quickly removed and transferred into ice-cold artificial cerebrospinal fluid (ACSF) containing a high Mg^{2+}- and a low Ca^{2+}-concentration (4 mM $MgCl_2$ and 1 mM $CaCl_2$) to reduce synaptic activity and bubbled continuously with carbogen (95% O_2 and 5% CO_2). It was then placed on the ramp of a slope of 10° and were cut at an angle of 50° to the midline[72]. Thalamocortical slices were cut at 350 μm thickness using a high vibration frequency and incubated for 30-60 minutes at room temperature (21-24°C) in slicing solution. During whole-cell patch clamp recordings, slices were continuously perfused with a perfusion solution containing (in mM): 125 NaCl, 2.5 KCl, 25 D-glucose, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 2 $CaCl_2$, 1 $MgCl_2$, 3 myo-inositol, 2 sodium pyruvate and 0.4 ascorbic acid, bubbled with carbogen and maintained at a temperature of 30-33°C. Patch pipettes were filled with an internal solution containing (in mM): 135 K gluconate, 4 KCl, 10 HEPES, 10 phosphocreatine, 4 Mg-ATP and 0.3 GTP (pH 7.4, 290-300 mOsm). To stain the patched neurons, biocytin was added at a concentration between 3-5 mg/ml to the pipette solution; a recording time of ~15-30 minutes was necessary for biocytin to diffuse into the dendrites and axons of the recorded cells[73,74]. No biocytin was added to the internal solution of 'searching' pipettes used during searching for synaptic connections.

Cell identification

Slices were placed in the recording chamber under an upright microscope (fitted with 4x plan/ 0.13 numerical aperture and 40x water immersion/0.80 NA objectives; Olympus, Tokyo, Japan) with the pial surface pointing forward. The cortical layers and the barrel field were visualized at 4x magnification; the barrels can be identified in L4 as narrow dark stripes with evenly spaced, light 'hollows' and were visible in 6-8 consecutive slices. L6A neurons were identified in the upper 60% of layer 6 at 40x magnification using infrared differential interference contrast (IR-DIC) microscope[75,76]. Putative PCs and interneurons were differentiated on the basis of their intrinsic

609 action potential firing pattern during recording and after histological processing by their
610 morphological appearance.

Electrophysiological recordings

Whole-cell patch clamp recordings from L6A neurons were performed at 30-33°C for an optimal
oxygenation. Patch pipettes were pulled from thick-wall borosilicate capillaries (outer diameter: 2
mm; inner diameter: 1 mm) to a final resistance of 6-10 MΩ. Recordings were made using an
EPC10 amplifier (HEKA, Lambrecht, Germany), sampled at 10 kHz, and filtered at 2.9 kHz using
the Patch-master software (HEKA). Neurons were selected randomly and excluded from the
analysis when their whole-cell series resistance exceeded 40 MΩ (50 MΩ for neurons from paired-
recodings) or their resting membrane potential was more depolarized than -50 mV immediately
after rupturing the cell membrane. The resting membrane potential of L6A excitatory neurons was
continuously recorded in the current clamp mode to monitor changes in amplitude.

Miniature spontaneous events were recorded in voltage-clamp mode and changes in mEPSC
frequency and amplitude were analyzed. Recordings of L6A excitatory neurons were made in the
presence of tetrodotoxin (TTX, 0.5 µM) and gabazine (10 µM) to inhibit AP firing and inhibitory
postsynaptic currents (IPSCs), respectively. During recordings, the holding potential was set at -
70mV.

Because the connectivity of L6A neurons was low compared to other intra-laminar connections in
rat barrel cortex, we followed the 'searching procedure' described previously after patching a
putative postsynaptic neuron[74,77]. A monosynaptic connection can be found by patching multiple
cells in 'loose cell-attached' mode. When the AP resulted in a unitary excitatory postsynaptic
potential (uEPSP) in the postsynaptic L6A neuron, this presynaptic neuron was repatched with a
new pipette filled with biocytin containing internal solution. APs were elicited by current injection
in the presynaptic neurons and the postsynaptic response were recorded in whole cell (current
clamp) mode, the effects of ACh on unitary EPSPs were then tested.

Drug Application

ACh (1 µM-10 mM) was bath applied via the perfusion system or puff applied through a patch
pipette (tip diameter: 1-2 µm) connected to a PDES-02D device (npi electronic GmbH, Tamm,

Germany). The puff pipette was placed at 10-20 μm from the same recorded neuron and a brief

low pressure was applied for about 1 s. Mecamylamine (10 μM), atropine (200 nM-20 μM),

pirenzepine (0.5 μM), tropicamide (1μM), dihydro-ß-erythroidine (DHßE) (10 μM),

methyllycaconitine (MLA) (10μM), tetrodotoxin (TTX) (0.5 μM) and gabazine (10 μM) were all

bath applied; drugs were purchased from Sigma-Aldrich (Steinheim, Germany) or Tocris (Bristol,

UK).

Histological staining

After single cell or paired recordings, brain slices containing biocytin-filled neurons were

processed as described previously[73]. Slices were fixed at 4°C for at least 12 hours in 100 mM

phosphate buffer (PB, PH 7.4) solution containing 4% paraformaldehyde (PFA) and then incubated

in 0.1% triton X-100 solution containing avidin-biotinylated horseradish peroxidase (Vector ABC

staining kit, Vector Lab. Inc., Burlingame, USA). The reaction was catalyzed using 3,3′-

diaminobenzidine (Sigma-Aldrich, St. Louis, MO, USA) as a chromogen. Slices were again rinsed

with 100 mM PB solution several times, followed by slow dehydration using ethanol and xylene.

After embedding in Eukitt medium (Otto Kindler GmbH, Freiburg, Germany), the dendritic and

axonal structures were clearly visible.

Immunofluorescence staining was performed for the identification of molecular markers expressed

in L6A PCs. During electrophysiological recordings, Alexa Fluor® 594 dye (1:500, Invitrogen,

Darmstadt, Germany) was added to the internal solution for post hoc identification of patched

neurons. After recording, slices (350 μm) were fixed with 4% PFA in 100mM PBS for at least 24

hours at 4°C and then permeabilized in 1% milk power solution containing 0.5% Triton X-100 and

100 mM PBS. Primary and secondary antibodies were diluted in the permeabilization solution (0.5%

Triton X-100 and 100 mM PBS) shortly before experiments. For single cell-FoxP2 staining, slices

were incubated overnight with Goat-anti-FoxP2 primary antibody (1:500, Santa Cruz

Biotechnology, Heidelberg, Germany) at 4°C and then rinsed thoroughly with 100 mM PBS.

Subsequently, slices were treated with Alexa Fluor® secondary antibodies (1:500) for 2-3 hours at

room temperature in the dark. After being rinsed in 100 mM PBS the slices were embedded in

Moviol. The fluorescence images were taken using the Olympus CellSens platform. The position

of the patched neurons were identified by the conjugated Alexa dye, so that the expression of FoxP2

could be tested in biocytin-stained neurons. After acquiring fluorescent images, slices were

incubated in 100 mM PBS overnight and were processed for subsequent morphological analysis.

Co-immunostaining of FoxP2 and M_4Rs (Rabbit-anti-M_4Rs, 1:500, Abbexa, Cambridge, UK) were

performed with 150 μm thin brain slices following the procedure described above.

Morphological reconstructions

3D reconstructions of L6A excitatory and inhibitory neurons or synaptically coupled neuron pairs

labelled with biocytin were made using the NEUROLUCIDA® software (MicroBrightField Inc.,

Williston, VT, USA) and Olympus BX61 microscopy at 1000 X magnification. Slices were

selected to be reconstructed only if the labeling quality was high and the background staining was

low. Barrel borders, demarcation of different layers, pial surface and white matter were delineated

during reconstructions. The cell body, the axonal and dendritic branches were reconstructed

manually under constant visual inspection to detect even small collaterals. Corrections for

shrinkage were performed in all spatial dimensions (factor 1.1 in the x and y axes, factor 2.1 in the

z axes) 73. Analysis of 3D reconstructed neurons was done with NEUROEXPLORER® software

(MicroBrightField Inc., Williston, VT, USA).

The neuronal polarity of reconstruction was calculated with NEUROEXPLORER® software using

cubic spline smoothing. The dendritic and axonal length was averaged for each of the 120 "3°

sectors" around the soma. Data were recalculated, plotted in angular subdivision around the soma

and polar plots were made with Grapher software (GoldenSoftware, Colorado, USA). The radian

depicts degree in angles (°) with 0° towards the pial surface, 90° towards the posterior-median axis,

180° towards the white matter, and 270° towards the anterior-lateral axis.

Data analysis

Custom written macros for Igor Pro 6 (WaveMetrics, Lake Oswego, USA) were used to analyze

the recorded electrophysiological signals. The miniature spontaneous activity was analyzed using

the program SPCN (http://www.spacan.net). A threshold of 5 pA was set manually for detecting

mEPSC events, which is at least 2.5 fold larger than the noise level (< 2pA). No noise filtration

was applied before data analysis.

The synaptic properties were evaluated as described in the previous studies[77]. First, all sweeps were

aligned to their corresponding presynaptic AP peaks and an average sweep was generated as the

mean uEPSP. The EPSP amplitude was calculated as the difference between the mean baseline

amplitude and maximum voltage of the postsynaptic event. The paired pulse ratio was defined as

the second uPSP amplitude divided by the first uPSP amplitude of the mean uPSP elicited by paired

APs with a stimulation frequency of 10 Hz. Failures were defined as events with amplitudes $<1.5\times$

the standard deviation (SD) of the noise within the baseline window and the failure rate refers to

the percentage of failures. The coefficient of variation (CV) was calculated as the SD divided by

the mean uEPSP amplitude.

Statistical tests

For all data, the mean ± SD was given. To assess the differences between two paired groups under

different pharmacological conditions, Wilcoxon signed rank test was performed. The paired

Student's *t* test was used when n = 4 for the paired samples and the Mann-Whitney *U*-test was used

when the sample size was different between two groups. Statistical significance was set at $P < 0.05$,

n indicates the number of neurons or pairs analyzed.

Figures

Fig. 1 Low concentrations of ACh induce either a hyperpolarization or a depolarization of L6A PCs by activating mAChRs.

(a, b) Top, L6A PC either show a hyperpolarizing (a) or a depolarizing (b) response following bath application of 100 μM ACh. Bottom, firing patterns of neurons in response to 1 s depolarising current injection (rheobase + 100 pA) before and during ACh application.

(c, d) Representative voltage-clamp recordings with bath application of ACh showing outward current (I_{out}) (e) or inward current (I_{in}) (f) in L6A PCs. The effects are blocked by 20 μM atropine (ATRO).

(e, f) Representative current-clamp recordings showing that puff application of ACh (100μM) evokes a fast hyperpolarization (HP) (g) or depolarization (DP) (h) in L6A PCs. The specific M_1 mAChR antagonist pirenzipine (PIR, 0.5 μM) or the specific M_4 mAChR antagonist tropicamide (TRO, 1 μM), respectively, were added to the perfusion solution to block ACh-induced membrane potential changes.

(g, h) Summary bar graphs showing the percentage block by general and specific mAChR antagonists. n = 7, p = 0.0006 for I_{out} group, n = 7, p = 0.0022 for I_{in} group, n = 4, p = 0.029 for HP group and n = 6, p = 0.0022 for DP group. Statistical analysis was performed using Mann Whitney U-test . Error bars represent SD.

(i, j) Muscarinic responses of ACh were examined in the presence of 1 μM mecamylamine (MEC) and 0.5 μM TTX. ACh dose-response curves for hyperpolarizing (n = 5) (c) and depolarizing (n = 8) (d) L6A PCs are fitted by the Hill equation. Dashed lines represent half maximal effects. The corresponding EC_{50} is 6.2 ± 1.3 μM for hyperpolarizing PCs and 26.7 ± 5.4 μM for depolarizing PCs. Filled circles represent mean values of different ACh concentrations.

Fig. 2 ACh hyperpolarizes CC L6A PCs but depolarizes CT L6A PCs.

(a) Left, overlay of reconstructions of CC and CT PCs. Reconstructions of PCs were aligned with respect to the barrel center. Right, polar plots of CC and CT PCs. n = 15 for each group. Somatodendrites are shown in red and axons are shown in blue.

(b) Histograms comparing the length, fieldspan and number of first order collaterals of axonal and dendritic structures for the two groups of PCs. n = 21 for CC neurons and n = 54 for CT neurons. Dendritic length: p = 0.0015, dendritic fieldspan: p = 1.4×10^{-10}, number of dendritic main nodes: p = 1.7×10^{-6}; Axonal length: p = 9.6×10^{-8}, dendritic fieldspan: p = 4.8×10^{-11}, number of axoanl main nodes: p = 8.5×10^{-11} for Mann-Whitney *U*-test.

(c) Top, representative current-clamp recordings of a depolarizing CC (orange) and a hyperpolarizing CT pyramidal cell (green) following bath application of 30μM ACh. Bottom, histograms of resting membrane potential (V_m) of L6A CC PCs in control and in the presence of 30 μM ACh (n = 35, p = 0.019 for Wilcoxon signed-rank test) and CT (n = 14, p = 6.1×10^{-5} for Wilcoxon signed-rank test) PCs.

756   **(d)** Plots of the ACh-induced change in V_m vs axonal fieldspan for two subtypes of PCs. Open

757   orange circles, data from individual CC PCs (n = 27); open green circles, data from individual CT

758   PCs (n = 13). Filled orange circle, average data from CC cells; filled green circle, average data

759   from CT cells.

Fig. 3 High concentration of ACh selectively depolarizes CT PCs.

(a) In the presence of 200 nM atropine (ATRO), bath application of low concentration ACh (30 µM, 50s) show no effect on either CC (top) or CT (bottom) PCs. Summary plots show the resting membrane potential (V_m) under control and ACh conditions in L6A CC (n = 5, p = 0.188) and CT (n = 5, p = 0.0625) PCs. NS (not significant) for Wilcoxon signed-rank test. Error bars represent SD.

(b) In the presence of 200 nM ATRO, bath application of 1 mM ACh for 50s has no effect on CC L6A PCs (top) but induces a strong depolarization of CT L6A PCs (bottom). Summary plots showing V_m of CC L6A PCs under control conditions and in the presence of ACh in L6A CC (n = 5, p = 0.156) and CT (n = 12, p = 0.0002) PCs. Statistical analysis was peformed using Wilcoxon signed-rank test. Error bars represent SD.

(c) The dose-response curve of ACh under ATRO application (200 nM) in CT PCs (n = 5) is well fitted by the Hill equation. The dashed line indicates the half maximal effect; the corresponding EC_{50} is 1.2 mM. Filled circles show mean effect of different concentrations while open circles represent individual values. Error bars represent SD.

(d) The depolarization induced by ACh application (in the presence of 200 nM ATRO) is blocked by 10 µM of the specific antagonist of $\alpha_4\beta_2$ subunit-containing nAChRs DHßE in CT L6A PCs. Summary plots showing the amplitude of the depolarization in response to application of 1 mM ACh in the presence of ATRO alone (open bar) and ATRO together with DHßE (n = 7, p = 0.0078 for Wilcoxon signed-rank test). Error bars represent SD.

Fig. 4 ACh differentially modulates miniature spontaneous activity in CC and CT L6A PCs.

(a) Example voltage-clamp recordings of a CC L6A PCs under control (black), bath application of 30 μM ACh (gray) and co-application of ACh and 1 μM TRO (purple). Miniature EPSCs were recorded in the presence of TTX (0.5 μM) and GABAzine (10 μM) at a holding potential of -70 mV.

(b) Cumulative distributions of mEPSCs inter-event interval recorded in CC L6A PCs under control condition, in the presence of ACh alone, and of ACh & TRO. Summary histograms of mEPSC frequency are shown on the right. **P < 0.01 for Wilcoxon signed-rank test. Error bars represent SD.

(c) Cumulative distributions of mEPSCs amplitude recorded in CC L6A PCs under control, ACh and ACh & TRO conditions. Summary histograms of mEPSC amplitude are shown on the right. Control vs. ACh, p = 0.8125; ACh vs. ACh & TRO, p = 0.9375, n = 7 for Wilcoxon signed-rank test. Error bars represent SD.

(d) Example voltage-clamp recordings of a CT L6A PC in control (black), after bath application of 30 μM ACh (gray) and subsequent co-application of ACh and 10 μM DHßE (turquoise). Miniature EPSCs were recorded in the presence of TTX (0.5 μM) and GABAzine (10 μM) at a holding potential of -70 mV.

(e) Cumulative distributions of mEPSCs inter-event interval recorded in CT L6A PCs under control,

ACh and ACh & DHßE conditions. Summary histograms of mEPSC frequency are shown on the

right. **P < 0.01 for Wilcoxon signed-rank test. Error bars represent SD.

(f) Cumulative distributions of mEPSCs amplitude recorded in CT L6A PCs under control, ACh

and ACh & DHßE conditions. Summary histograms of mEPSC amplitude are shown on the right.

Control vs. ACh, p = 0.4258; ACh vs. ACh & DHßE, p = 0.0781, n = 7 for Wilcoxon signed-rank

test. Error bars represent SD.

Fig. 5 ACh-mediated reduction of presynaptic release at CC PC synapses.

(a) Left, Schematic representation of the synaptic connections with a presynaptic CC L6A PC. CC PCs are shown in orange, the CT PC in green and the interneuron in black. Barrel structures indicate layer 4. Right, corresponding firing patterns of pre- and postsynaptic neurons of the same connection type.

(b) Time course of EPSP amplitude changes following bath application of 30μM ACh (gray phases) in a CC-CC, a CC-CT and a CC-interneuron pair.

(c) Overlay of average EPSPs in control (black) and ACh application (gray) phases. Presynaptic APs are shown at the top. Data are recorded from the same pairs as in (b).

(d) The average and SD of several EPSP properties for CC-CC (n = 20), CC-CT (n = 5) and CC-interneuron (n = 9) connections are shown. *P < 0.05, **P < 0.01, ***P < 0.001 for Wilcoxon signed-rank test.

Fig. 6 ACh decreases presynaptic release probability of CC PC via activation of M₄ AChRs.

(a) Left, Overlay of average EPSPs recorded in control, the presence of ACh (30μM), and of ACh & TRO (1 μM) from a representative CC-CC connection. Right, normalising the mean EPSP amplitudes obtained in ACh and ACh & TRO to the first EPSPs amplitude in control reveals changes of PPR. Presynaptic APs are shown at the bottom.

(b) Histograms (n = 6) showing the effect of ACh and TRO blockade of ACh-induced changes on several EPSP properties including EPSP amplitude, PPR, CV and failure rate. Data were recorded from L6A synaptic connections with a presynaptic CC PC. Open circles, individual data points; bars, the average for each condition. Error bars represent SD. *P < 0.05 for Wilcoxon signed-rank test.

Fig. 7 ACh enhances synaptic efficacy of L6A excitatory connections with a presynaptic CT PC.

(a) Left, schematic representation of the synaptic connections with a presynaptic CT PC. Color code as in Fig. 5. Barrel structures indicate layer 4. Right, corresponding firing patterns of pre- and postsynaptic neurons of the same connection type.

(b) Time course of EPSP amplitude change following bath application of 30 μM ACh in a CT-CT, CT-CC, and CT-interneuron pair.

(c) Overlay of average EPSPs in control (black) and ACh application (gray) phases. Presynaptic APs are shown at the top. Data are recorded from the same pairs as in (b).

(d) Summary data (n = 7) of ACh-induced changes in first uEPSP amplitude and paired-pulse ratio for L6A excitatory pairs with a presynaptic CT L6A PC. Bars indicate the average for each condition; Error bars represent SD. *P < 0.05 for Wilcoxon signed-rank test.

Fig. 8 ACh increases presynaptic release probability of CT PC via activation of α₄β₂ nAChRs.

(a) Average EPSPs recorded in control, the presence of ACh (30μM), and of ACh & DHßE (10 μM) from a representative CT-CT connection. Gray phase, bath application of 30μM ACh; Turquoise phase, co-application of ACh and DHßE. The presynaptic APs are shown at the bottom.

(b) Histograms (n = 4) showing the effect of ACh and DHßE blockade of ACh-induced changes in first uEPSP amplitude and paired-pulse ratio. Data were recorded from L6A synaptic connections with a presynaptic CT PC. Open circles, individual data points; bars, the average for each condition. Error bars represent SD. *P < 0.05 for paired Student's t-test.

Fig. 9 Cholinergic actions on muscarinic and nicotinic receptors in L6A PCs.

Schematic summary shows cholinergic modulation of L6A microcircuits in rat barrel cortex. ACh affects membrane excitability and presynaptic release probability of L6A PCs via activating muscarinic and/or nicotinic AChRs. Because of the cell type-specific distribution of AChRs at pre- and postsynaptic sites, CC and CT L6A PCs are affected differentially by ACh. The presynaptic CC and CT L6A PC are shown in orange and green, respectively. The postsynaptic L6A PC and interneuron are shown in gray.