UNCOVERING PHOTOLYASE/CRYPTOCHROME GENES DIVERSITY IN AQUATIC MICROBIOMES EXPOSED TO DIVERSE UV-B REGIMES

Daniel G. Alonso-Reyes[1,2], Maria Eugenia Farias[2] and Virginia Helena Albarracín[1,2,3]*

[1]Laboratorio de Microbiología Ultraestructural y Molecular, Centro Integral de Microscopía Electrónica (CIME, CONICET, UNT) CCT, CONICET, Tucumán, Argentina.

[2]Laboratorio de Investigaciones Microbiológicas de Lagunas Andinas (LIMLA), Planta Piloto de Procesos Industriales y Microbiológicos (PROIMI), CCT, CONICET, Tucumán, Argentina.

[3]Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Tucumán, Argentina.

Running headline: METAGENOMICS OF PHOTOLYASE/CRYPTOCHROMES IN AQUATIC MICROBIOMES

Keywords: UV, PHOTOLYASES, CRYPTOCHROMES, MICROBIOMES, METAGENOMICS

*Corresponding author:

Virginia Helena Albarracín, Centro Integral de Microscopía Electrónica (CIME, CONICET, UNT) Camino de Sirga s/n. Finca El Manantial, Yerba Buena (4107). Tucumán, Argentina.

ARTICLE TYPE

Original Research Article

ABSTRACT

During evolution, microorganisms exposed to high UV-B doses developed a fine-tuned photo-enzymes called "photolyases" to cope with DNA damage by UV-B. These photoreceptors belonging to the Cryptochrome/Photolyase Family (CPF) were well characterized at the genomic and proteomic level in bacteria isolated from a wide range of environments. In this work, we go further towards studying the abundance of CPF on aquatic microbial communities from different geographic regions across the globe. Metagenomics data combined with geo-referenced solar irradiation measurements indicated that the higher the UV-B dose suffered by the microbiome's environment, the higher the abundance of CPF genes and lower the microbial diversity. A connection between CPF abundance and radiation intensity/photoperiod was reported. Likewise, cryptochrome-like genes were found abundant in most exposed microbiomes, indicating a complementary role to standard photolyases. Also, we observed that CPFs are more likely present in dominant taxa of the highly irradiated microbiomes, suggesting an evolutionary force for survival and dominance under extreme solar exposure. Finally, this work reported three novel CPF clades not identified so far, proving the potential of global metagenomic analyses in detecting novel proteins.

1. INTRODUCTION

Solar radiation is an essential factor sustaining complex life on Earth. In the past decades, the drastic reduction of stratospheric ozone produced by increased concentrations of chlorofluorocarbons (CFCs) (Aucamp 2007) and other halogen gases in the upper atmosphere (Russell III et al. 1996) resulted in a rise of biologically harmful UV radiation (UV) at the Earth´s surface together with its detrimental effect on all life forms.

The amount of UV radiation reaching the ground comprises only a small proportion of global radiation, about 6–7 % of UV-A (320–400 nm), less than 1.0 % of UV-B (280–315 nm) and virtually no UV-C (Hu et al. 2008). Biological damage is wavelength-dependent: UV-A produces mainly reactive oxygen intermediates causing indirect damage to DNA, proteins, and lipids. On the other hand, UV-B and UV-C (100 to 280 nm) cause both indirect and direct damage to DNA because of its strong absorption at wavelengths below 320 nm (Mitchell & Karentz 1993). In accordance, numerous studies reported the UV-B effects on plants (Searles et al. 2001, Xiong & Day 2001, Robinson et al. 2005, Ruhland et al. 2005, Jansen et al. 2010, Yan et al. 2012), animals (Robson et al. 2005, Bao et al. 2014), microorganisms (Zaller et al. 2002, Avery et al. 2004, Rinnan et al. 2005, Piccini et al. 2009) and ecosystems (Garcia-Pichel 1994, Karentz 1995, Joux et al. 1999, Ballaré et al. 2011, Häder et al. 2011).

Solar UV radiation can damage aquatic organisms and decrease the productivity of aquatic ecosystems. Beyond the targets for damage by UV radiation (DNA, lipids, protein) that are common for all biological systems, a major site of damage in primary producers is the photosynthetic machinery including photosystem II and the accessory pigments that

funnel light energy to the reaction centers (Häder & Gao 2015). The subsequent damage

will directly reduce primary production. Also, the effects of UV radiation may reduce the

photosynthetic uptake of atmospheric carbon dioxide and affect species diversity,

ecosystem stability, trophic interactions, and global biogeochemical cycles (Häder et al.

2011, Williamson et al. 2019).

During evolution, microorganisms exposed to high UV doses and other DNA

damaging factors have developed specific and highly conserved DNA repair mechanisms.

The most important are photoreactivation, excision repair (NER and BER), mismatch repair

(MMR), and homologous repair (HR). Also, damage tolerance (dimer bypass), SOS

response, checkpoint activation, and programmed cell death or apoptosis efficiently act

against DNA lesions ensuring genomic integrity.

The finest and efficient mechanism to repair DNA damage by UV-B is

photoreactivation, executed by photoreactivating enzymes known as "photolyases," which

target the main products of UV-B, cyclobutane pyrimidine dimers (CPD). Such CPD

lesions bring polymerases to a standstill, eventually leading to cell death. These enzymes

bind tightly to CPDs in the dark and can be activated by different wavelengths, causing the

split of the two C–C bonds in the CPD unit and resulting in the re-formation of the two

separate pyrimidine bases (S Weber 2005). Photolyases, together with the related

cryptochromes (Cry), constitute the Cryptochrome/Photolyase Family (CPF). However,

cryptochromes have no photolyase activity and function as signaling molecules regulating

diverse biological responses such as entrainment of circadian rhythms in plants and animals

(Roenneberg & Merrow 2005, Harmer 2009). It has been argued that CPF had had an early

evolution in the history of life due to the need of the primitive organisms to cope with the

exacerbated solar radiation in the time when no stratospheric ozone was protecting the

Earth (Toh et al. 1997, Portero et al. 2019).

Investigations at metagenome-level about the effects of sunlight on microbial

communities were mostly confined to photoreceptors such as microbial rhodopsins

(Pushkarev & Béjà 2016, Pushkarev et al. 2018) and LOV-domains (Pathak et al. 2012).

Likewise, Singh et al. (2009) reported abundances for several light-related genes in

microbiomes from different environments (Singh et al. 2009). However, abundance and

diversity of CPF on microbiomes in a world-wide scale and light range were not studied so

far, although being these photoreceptors a vital issue in the defense against UV-B. In this

work, we studied aquatic microbiomes exposed to different intensities of UV-B and

photoperiods around the globe through metagenomic analysis of existing published DNA

datasets. Thus, the objective was to characterize the occurrence and diversity of CPF on

microbiomes from aquatic ecological niches, with some of them suffering from extreme

conditions. Our results add a new dimension to the understanding of the short-term effects

of climate change and atmospheric ozone depletion on environmental microbial

communities. Besides, the focus of this work on microbial communities from extreme

environments provides models for early life research.

2. **MATERIALS AND METHODS**

2.1. Metagenome selection and UV-B data processing. The datasets of

metagenomes selected for this work from the NCBI database were obtained and sequenced

by shotgun strategy through Illumina technology except for Lake Diamante metagenome,

which was sequenced with 454 GS FLX Titanium instrument. The accession numbers are

the following: Lake Diamante (DM) (ERR1824222), Socompa stromatolite (SS)

(SRR3341855), Tibetan Plateau sediment (TB) (SRR3322106), Amazon River (AM)

(SRR1790676), Lake Rauer (RA) (SRR6129205), Dewar Creek hot spring (HT)

(SRR5580900), Greenland cryoconite (CR) (SRR5275901), Lake Montjoie   (MT)

(SRR5818193), Olkiluoto Island groundwater (OK) (SRR6976411) and human gut (GU)

(SRR6517782).

An essential aspect of this work was to reanalyze published metagenomic data of

world-wide distributed microbiomes taking into account their overall UV-B exposure (Fig.

1). Monthly mean UV-B glUV datasets (Beckmann et al. 2014) processed using QGIS

(www.qgis.org) were utilized to calculate UV-B exposure of the samples of the selected

metagenomes according to the date/location of sampling available at the "biosample"

linked section of each SRA entry. To compensate for the lack of replicates on most

samples, we performed the analysis by clustering the datasets in three groups according to

UV-exposure regimes: High (UV_{High}), Mid (UV_{Mid}), and low-exposed (UV_{Low}). The

metagenome of the human gut was assumed as a UV-B free environment, acting as a

negative control. Table 1 showed the main characteristics of these environmental datasets

when available. A brief description of the environmental conditions of each sampling site is

summarized below, including citation to previous works describing the corresponding

microbial communities with further detail:

DM samples were taken from red biofilms attached to the bottom of calcareous

stones submerged in Lake Diamante on 15[th] February 2012. This lake is located inside the

Galan Volcano crater placed at 4,589 m above sea level (coordinates: -26.008889, -

67.043333). High pH (9 to 11), high arsenic concentrations (115 to 234 mg l − 1), high

salinity (270 g l − 1, 217 mS cm − 1), high UV radiation (84 W m − 2 of UVA-B at noon),

long day-night temperature range (−20 °C to +20 °C) and low O_2 pressure constitutes a set

of unique extreme conditions that prevails in the lake (Rascovan et al. 2016).

SS belongs to stromatolites found along the southern shore of Socompa Lake (-

24.5361333, -68.20275) at the base of the Socompa Volcano (3,570 masl). These structures

were found around the border of Socompa Lake during the summer when they are partially

exposed depending on the tide and hydrological regime; at the opposite, they are entirely

submerged during winter and spring. The stromatolites are rounded domal structures that

present a clear stratification that appear in vertical sections. The samples were cylinders of

5 cm deep from the top of the stromatolite. The site was exposed to air at the moment of

sampling in February 2011. Extreme environmental conditions of the lake include

hypersalinity, high thermal amplitude with daily temperatures that range from −10 °C to 20

°C in summer and −20 °C to 10 °C in winter, UV solar irradiance that reaches 68 W m−2

19, 32, low O_2 pressure, low nutrient availability and, primarily, high arsenic content (18.5

mg L−1) (Kurth et al. 2017).

TB was collected on 1[st] August 2013 from sediments of the Qiangyong Glacier

Lake (28.51, 90.12) that is located in the southern part of the Tibetan Plateau, in the Indian

monsoon climate region. The temperature, pH, and electric conductivity of surface water

are, on average, 6.89 °C, 8.32, and 136.4 µS cm[-1], respectively. (Chen et al. 2016).

AM water sample was collected on 19[th] May 2011 from Amazon River at Obidos

station (-1.919017, 55.525717) at 33 m of depth by pumping water with a Shurflo

submersible pump through a 297 µm stainless steel screen. The temperature, pH, and O_2

were 28.4 °C, 6.6, and 3.2 m/L respectively on the site of sampling (Satinsky et al. 2015,

Doherty et al. 2017).

RA was sampled on 12 January 2015 from Rauer Lake (Torckler Island; -68.5558

78.1913). Water temperature and approximate depth of water at the sampling site were 9°C

and 10-15cm. The lake was shallow <30cm. There was a crust of salt crystals on the

sediment (which make the lake look frozen from the air). There also appeared to be

stratification in the lake: a top clear layer of ~10cm and a bottom layer of ~5cm. These

layers weren't visible until they were disturbed, and a visible haze produced when these

layers were mixed. (Tschitschko et al. 2018)

CR belongs to a cryoconite sample collected on 29[th] August 2013 from Greenland

ice sheet in the TAS-U-A1 site (65.41, -38.51). pH, altitude, and depth of sampling were

5.53, 580 m, and 20 cm, respectively. Cryoconite was collected from 2–18 holes at each

site, depending on their availability, using sterile 50 ml syringes (Stibal et al. 2015a b,

Hauptmann et al. 2017).

MT microbial sample was collected from Lake Montjoie, in Canada (45.4091, -

72.0994 W) on February 2014, being the lake covered of ice at the time of sampling. The

sample was taken from the epilimnion strata of the lake (0-3 m), and pH in the site of

sampling was 6.41 (Tran et al. 2018).

HT sediment sample was collected from Dewar Creek hot spring (49.9543667, -

116.5155000) near the source of the hot spring on 28[th] September 2012. pH and

temperature of the environment were 7.93 and 66.4 °C (Eloe-Fadrosh et al. 2016).

OK belongs to groundwater collected from a deep subsurface site KR11_0.1

(61.2413, 21.4947) in Olkiluoto Island, on the south-west coast of Finland on 9[th] September

2016. Given the environmental data provided by the authors of the original research

(Bomberg et al. 2016), 60 drill holes were collected from different fractures at different

depths (296-798 mbsl). The groundwater is stratified with a salinity gradient extending

from fresh to brackish water to a depth of 30 m and the highest salinity concentration of

125 g L−1 total dissolved solids (TDS) at 1,000 m depth (Posiva, 2013). Between 100 and

300 m depths, the groundwater originates from ancient (pre-Baltic) seawater and has high

concentrations of SO_2^{-4}. The temperature rises linearly with depth, from ca. 5–6 ∘C at 50 m

to ca. 20 ∘C at 1,000 m depth (Ahokas et al., 2008). The pH of the groundwater is slightly

alkaline throughout the depth profile. The bedrock of Olkiluoto consists mainly of mica

gneiss and pegmatitic granite-type rocks (Kärki and Paulamäki, 2006). The *in-situ*

temperature at 300 m depth in the Olkiluoto bedrock is stable at approximately 10 ∘C and

increases linearly to approximately 16 ∘C at 800 m depth.

GU corresponds to a gut metagenome collected from the faeces of an active

Behcet's disease patient (Ye et al. 2018) was used as negative control with no exposure to

UV.

2.2. Meta-analysis, quality control, and assembly of selected metagenomes.

Quality filtering and merging yielded a range between 0.64 and 13.66 Gb for further

analysis (Table 1). Only TB and AM datasets reported a high percentage of merging with

FLASH (Magoč & Salzberg 2011); however, the low-merged dataset from OK was also

used for downstream steps as it ended with a reasonable size (between the range mentioned

above). MEGAHIT (Liu et al. 2015) was used for assembly since it can deal with large and

complex datasets in a time- and cost-efficient manner. Protein prediction with Prodigal

(Hyatt et al. 2010) over contigs with >999 bp outputted a rate of 1000-1500 proteins / Mb,

which is congruent with the high coding density expected for microbial species.

Adapters were removed from Illumina raw reads using fastq-mcf tool of ea-utils

v1.04.676 (Aronesty 2011). This step was not needed for Lake Diamante metagenome, as it

was sequenced with 454 technologies. The quality filtering and trimming were performed

with the same program using parameters l>50 and q>20. The program kneaddata v0.6.1

(https://bitbucket.org/biobakery/kneaddata/wiki/Home) was used with the --bypass-trim

option to clean human contaminant sequences. Pair end reads were merged with FLASH

v1.2.11 (Magoč & Salzberg 2011) to recover unpaired longer reads. The pair end reads

with a low percentage of merging were left in the paired state for assembly. Assembly of

filtered reads was performed with MEGAHIT v1.1.2 (Liu et al. 2015). Assembled contigs

were annotated with Prodigal v2.6.2 (Hyatt et al. 2010), which outputted translated protein

sequences.

To perform quantitative metagenomics, pair-end reads that remained with a low

percentage of merging were concatenated together by the forward and reverse using a script

available   at   GitHub

(https://github.com/LangilleLab/microbiome_helper/blob/master/concat_paired_end.pl).

2.3. Reference database building. Considering the main functional orthologous of

the CPF group, a reference protein database was built. The two KEGG Orthology number

corresponding to CPF members (K06876, K01669) was linked with the UniProt database,

and then filtered results by EC number, known molecular function or biological process,

uniref 90 clusters, and sequence length >100 subsequently. The single, concatenated, and

merged reads were aligned to the aforementioned databases with PALADIN, which

outputted read counts. The counts were normalized to Reads per Kilobase per Genome, and

the number of genomes computed through MicrobeCensus v1.1.0 (Nayfach & Pollard

2015). The following formula describes the normalization:

$$\frac{\text{Read Counts} \div \text{Number of Genomes}}{\text{Gene Length}}$$

2.4. Estimation of gene abundance. To search and quantify the abundance of the

genes in the different metagenomes, we aligned the preprocessed unpaired reads against the

CPF reference database created before. The protein alignment was performed with

PALADIN v1.3.1 (Westbrook et al. 2017), which outputs a table with alignment counts for

each gene. Alignments were filtered by maximum quality = 60.

Diversity analysis of each metagenome was performed with MetaPhlAn v2.7.7

(Truong et al. 2015) with the **--**bt2_ps parameter set to 'sensitive.' The program MetaPhlAn

profiles the composition of microbial communities from metagenomic shotgun sequencing

data at species-level. It relies on ~1M unique clade-specific marker genes identified from

~17,000 reference genomes (~13,500 bacterial and archaeal, ~3,500 viral, and ~110

eukaryotic). Shannon diversity index was obtained with Vegan 2.5-2 R package (Oksanen

et al. 2018).

2.5. Analysis of the annotated genes. The protein sequences were aligned using

Diamond v0.9.22 software (Buchfink et al. 2014) against the CPF genes of the reference

database built earlier. Alignment parameters were 50% identity, 70% query coverage and e

$< 10^{-5}$. The retrieved files coming from each metagenome were modified at the sequence

headers to hold the name of its respective metagenome. Next, all files were merged in a

single one and filtered by sequence length > 400 residues.

The filtered sequences were used for phylogenetic analysis. The phylogenetic tree

was built with FastME v2.0 (Lefort et al. 2015) using the Jones–Taylor–Thornton rate

matrix (Jones et al. 1992) with 1000 bootstrap replicates. A consensus of the 1000 resulting

trees was selected for further processing and visualization using iTOL (Letunic & Bork

2016). Additionally, a taxonomical identification of the sequences was performed through

BLAST web server (https://blast.ncbi.nlm.nih.gov/Blast.cgi).

3. RESULTS AND DISCUSSION

3.1. UV-B intensity profiles on worldwide microbiomes. The UV-B geo-

referenced data helps us to group microbiomes according to their UV-exposure regimes in

UV_{High}, $UV_{Mid,}$ and UV_{Low} as described in materials and methods. The UV_{High} included

DM, SS, and TB with UV-B intensities of 9677, 9536, and 8885 J/m2/day, respectively. In

contrast, UV_{Low} microbiomes OK, MT, and CR suffered from low UV-B intensities of 77,

719, and 1759 J/m2/day, respectively. In between the UV_{Mid} microbiomes AM, RA, and

HT, received intensity of 5630, 2289, and 2281 J/m2/day, respectively. Important to remark

is that these intensity values were registered taking in account the sampling dates; for

instance, the samples of the UV_{High} group were registered during summer, when the

insolation is maximum on the high altitude environments of the Argentinean Puna and the

Tibetan Plateau (February and August), and UV-B incidence become the highest of the

planet. On the other hand, the UV_{Low} microbiomes were sampled during August,

November, and February, which correspond to summer, autumn and winter seasons in the

northern hemisphere with lower levels of irradiation as compared to the southern

hemisphere. In turn, the UV_{Mid} included the AM microbiome from a tropical environment,

and RA and HT from high-latitudes extreme cold and hot environments, respectively.

The calculated day-length from the UV-datasets for each georeferenced

metagenome (https://www.suncalc.org) revealed that two microbiomes suffered from

peculiar photoperiods at the time of sampling. While the OK microbiome was experiencing

a short photoperiod -only 7.8 hours-, RA was sampled when Antarctica went through an

entire day under the sun. We assumed that the environments were exposed to similar

conditions of indicated solar irradiance for a considerable broad period already before

sampling. In consequence, the diversity of species and genes on genomes are a reflection of

the ecological pressure of radiation on the environment (among other factors) as the

geographic conditions (latitude, altitude, and orography) did not change considerably in

those sampled regions for decades. Although no available, transcriptomic data on these

sites would be an excellent tool to evaluate the reflection of environmental pressure on the

dynamics of the microbial communities. Besides, the intensity of solar irradiation over each

microbiome also depend on their on-site spatial disposition being maximum for samples

taken from soils or shallow water of lakes, springs, oceans, and streams but much lower in

sediments, groundwater or deep water.

It is known that UV-B negatively affects microbial diversity (Ballaré et al. 2011);

nevertheless, this was not assessed so far in a systematic way across different irradiation

regimes on a global scale. The herein results used disperse metagenomic data to assessed

species richness and Shannon diversity index in diverse ecological niches, suggest an

interesting trend (Fig. 2); TB, SS, and DM from the UV_{High} group had the lowest species

number, while the GU microbiome, which received null irradiation, had the maximum

number of species. Moreover, microbiomes exposed to intensities below 4000 $J/m^2/day$,

revealed relatively high species richness (number of species: 11-103), while SS microbiome

with intensities above 8000 J/m2/day only showed 21 species.

Species richness (Fig. 2A) was quite variable for microbiomes whose UV-B

exposure was below a certain threshold (6000 J/m^2/day). In microbiomes receiving higher

UV doses, the number of species decreased, indicating selective pressure. Those species

that do not possess efficient molecular mechanisms to defend themselves from UV-B

become unable to adapt to the new assemblage of species. A similar situation was observed

when applying the Shannon index (Fig. 2B) (which incorporates equitability of the species

abundance in addition to its ability to detect rare species): microbiomes of the UV_{High}

group, and HT (UV_{Mid}) presented the lowest Shannon index values. These results suggest

that those microorganisms with full capacity to defend themselves against UV-B radiation

reach dominance, relegating others less fitted. Nevertheless, this correlation is not

conclusive as many other chemical and physical factors (salinity, nutrients availability,

heavy metals) may be causing selective pressure on microbial diversity.

3.2. Quantitative analysis of CPF. In order to evaluate the abundance of CPF

genes in each microbiome, a database was built using Uniprot sequences linked to KEGG

orthology and then aligned to the query metagenomes. Our results show that the

distribution of CPF genes in metagenomes followed a predicted ecological pattern; CPF

genes were abundant in microbiomes with UV_{High} and UV_{Mid}, while insignificant or null in

UV_{Low} microbiomes -OK and GU respectively. In DM, CPF genes appeared profusely.

Their abundances were 249 e^{-02} RPKG, which were higher than in the rest of the

microbiomes.

The relationship between the abundance of CPF and intensity/time of insolation was

studied (Fig. 3A), showing an upward trend of CPF abundance as UV-B irradiation

increases. However, we have seen that RA, an UV_{Mid} microbiome, completely deviates

from the trend having because of its high abundance. We had previously mentioned that

RA has the most extended photoperiod of the study, so we set out to verify if there is a

connection between the abundance of CPF and the photoperiod. Figure 3B shows that there

is an upward trend of CPF as the photoperiod increases, finding RA quite in line with the

trend. Thus, both factors, the intensity, and the photoperiod could be influencing the

abundance of CPF in the microbial communities

Our study reveals a rising trend of CPF abundance in microbial communities as

their environments receive higher radiation or day duration extends. That is likely due to an

increase in DNA damage caused by UV-B, which are mainly pyrimidine dimers. In such a

situation, populations could increase the expression of CPF by increasing the gene copy

number. Besides, species lacking CPF in their genome would be replaced by species that

contain these genes. In either case, the overall increase in CPF copy abundance would

indicate that the community improves its defense capabilities against UVB using the highly

efficient mechanism of photoreactivation or modulating enzymatic mechanisms triggered

by light sensing by cryptochromes.

3.4. Phylogenetic analysis of CPF genes. The CPF group of proteins comprises

mostly genes of photolyases with different specializations and cryptochromes whose

functions are mostly unknown. The family has been divided into different subgroups

considering phylogeny, chromophore type, specialization, host organism, and structure.

Alignment, editing, and subsequent phylogenetic analysis of 214 CPF sequences found in

the metagenomes configured a robust tree with numerous clades (Fig. 4). The sequences

mostly grouped within well-known subfamilies: the CPD photolyases classes I, II, and III,

DASH cryptochromes or single-strand photolyases (Selby & Sancar 2006), the FeS-BCPs

group that consist of proteins having an iron-sulfur cluster either prokaryotic [6-4]

photolyases or cryptochromes (Graf et al. 2015) and the thermostable CPD photolyases, a

group of unstable position in phylogeny, that have FMN as antenna cofactor and a

thermostable nature (Ueda et al. 2005) (Portero et al. 2019). Also, this work reports three

novel extra clades: i.e., unidentified I, II, and III (UI, UII, and UIII) with 41%, 88%, and

96% of bootstrap support, respectively. The three new clades grouped 24.76% of the whole

sequences in the analysis (Fig. 5), thus offering us a large pool of candidates for new

functions, specializations, or molecular specificities. Interestingly, the clades UI and UII

are mainly composed of sequences of a single microbiome; sequences from the RA

microbiome constitute 85.7% of UI, while 87.5% of UII corresponds to MT sequences. The

UI subfamily is the only clade in DM but dominant in RA; both microbiomes displayed the

highest content of CPF´s genes in this study. Furthermore, UI and UII clades are

taxonomically homogeneous, with their sequences belonging to Halobacteria and

Actinobacteria, respectively. On the other hand, UIII is formed by sequences of different

taxonomic origins, including Proteobacteria, Bacteroidetes, and Verrucomicrobia phylum.

It is clear the widespread presence of cryptochrome-holding subfamilies cry-DASH

and Fes-BCPs among the microbiomes (Fig. 6). The abundance of cry-DASH followed a

pattern in consistence with higher irradiation conditions; the highest number of cry-DASH

sequences are found in SS and TB communities (UVB_{High}), AM (UVB_{Mid}) and RA (the

most prolonged photoperiod). It has been reported in previous works that these

cryptochromes are, in fact, photolyases with an affinity for single-stranded DNA and, in

some cases RNA (Selby & Sancar 2006). Cry-DASH may play a complementary role of

standard photolyases with an affinity for double-stranded DNA, contributing to the global

increase of photoreactivation in these microbiomes. The case for FeS-BCPs group is

intriguing, being present in almost all the communities with similar relative abundance. It

has been suggested that this class complements other photolyases by performing the

function of a [6-4] photolyase (Zhang et al. 2013, Graf et al. 2015), avoiding or decreasing

in this way the use of the inefficient NER system. Our work gives more support to such

affirmation by showing its ubiquitous and homogenous presence in world-wide

microbiomes.

3.5. Relationship between CPF and taxa dominance. The assignment of CPF

genes to a particular taxonomic group was also considered in this work. CPF sequences

were classified by genus and paired with the same information previously obtained for each

microbiome through MetaPhlAn. The CPF sequences grouped with their homolog with a

BLAST identity lower than 70% clustered as an unclassified category. Genus diversity

from the whole microbiome (top bar) and the set of CPF sequences recovered from that

microbiome (lower bar) are shown (Fig. 6). The classification rate of the CPF genes was

generally high (>60%) in DM, SS, TB, AM, RA, and GU but lower in the rest of the

samples. MT had the lowest percentage of sequences (23%) assigned to some genus. This

phenomenon may be a consequence of the poorly referenced diversity of the community

itself (top bar). Despite this, MT had the most substantial diversity of genus (12) with CPF

hits along with SS and RA (11 each).

The most abundant taxa in DM, SS, RA, and HT communities possess CPF genes.

In AM, only *Ralstonia* and *Vibrio* displayed CPF; both genera together represented 35% of

the community. In the case of CR, only *Hymenobacter* and *Agromyces* possessed these

genes, with both groups adding up to 7%. *Pedobacter* was the unique taxon in TB with

CPF genes, and it represented less than 4% of the community. Finally, *Escherichia,* with an

abundance of just 0.44%, was the unique CPF contributor in GU. Neither in MT nor OK

found matches between the MetaPhlAn and BLAST classifications.

These results show that CPF and thus, photoreactivation is usually present in the

most abundant organisms of the UV$_{High}$ microbiomes (Fig. 6). Also, SS and RA show a

high number of taxa in which photoreactivation is present. That suggests that

photoreactivation may act as a successful system for assuring the survival and

predominance of taxa in UV-stressed environments. This fact contrasts with those less

insolated samples, where natural selection doesn't act in favor of photoreactivation

promoting abundance nor diversity of taxa carrying these genes.

4. CONCLUSIONS

In this work, UV-B was considered for the first time as an ecological variable in a

sequenced-based metagenomic study of microbial communities on a worldwide scale. Our

results consistently showed concordance of high UV exposure of a given microbiome with

its low microbial diversity and high CPF abundance.

We also evaluated the CPF diversity in the worldwide microbiomes and reported

three novel CPF clades not identified in previous analyses. CPF was more likely present in

most abundant organisms of the UV$_{High}$ microbiomes, suggesting a significant evolutionary

force for survival and dominance in highly irradiated environments. Also, cryptochrome-

like genes were much more abundant in the most exposed microbiomes indicating a

complementary role to standard photolyases.

Finally, metagenomics proved to be an excellent tool to reveal a clear correspondence between microbiomes, UV-exposure, and UV-B resistance mechanisms – measured in the number of gene copies. Additional methods, such as metatranscriptomics and metaproteomics, should be implemented in order to unveil the molecular dynamics of the CPF upon different light conditions.

5. ACKNOWLEDGEMENTS

VHA and MEF are staff researchers from the National Research Council (CONICET) in Argentina. DA is a recipient of a doctoral fellowship from CONICET. The authors have produced this manuscript despite the delays and shortages of funding execution from National Agencies in Argentina, mainly FONCyT (PICT 2013 2991) and CONICET (PIP 2015 0519, PIO-UNCA y PICT V 3825-2016), and the drastic devaluation of Argentinean currency during the period 2016-2019. This manuscript has been released as a Pre-Print at bioRxiv (Alonso-Reyes et al. 2019).

6. AUTHORS CONTRIBUTIONS STATEMENT

DAR designed the experiments and performed the data analysis. VA planned the research, provided training and analysis tools. DAR and VA wrote the paper. MEF and VA revised the manuscript. MEF performed HAAL sampling expeditions and provided the HAAL´s microbiomes sequences. MEF and VA obtained funding for the original project idea. All authors read and approved this manuscript.

 7. CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding the publication of

this article.

REFERENCES

Alonso-Reyes DG, Farias ME, Albarracín VH (2019) Unveiling The Worldwide UV-

Metaresistome: Metagenomic Analysis of Microbiomes Exposed to Different UV-B

Regimes. bioRxiv:701565.

Aronesty E (2011) Ea-utils: Command-line tools for processing biological sequencing data.

Aucamp PJ (2007) Questions and answers about the effects of the depletion of the ozone

layer on humans and the environment. Photochem Photobiol Sci 6:319–330.

Avery L, C. Thorpe P, Thompson K, Paul N, Grime J, M. West H (2004) Physical

disturbance of an upland grassland influences the impact of elevated UV☐B radiation

on metabolic profiles of below☐ground micro☐organisms.

Ballaré CL, Caldwell MM, Flint SD, Robinson SA, Bornman JF (2011) Effects of solar

ultraviolet radiation on terrestrial ecosystems. Patterns, mechanisms, and interactions

with climate change. Photochem Photobiol Sci 10:226–241.

Bao X, Li Q, Hua J, Zhao T, Liang W (2014) Interactive effects of elevated ozone and UV-

B radiation on soil nematode diversity. Ecotoxicology 23:11–20.

Beckmann M, Václavík T, Manceur AM, Šprtová L, von Wehrden H, Welk E, Cord AF

(2014) GlUV: a global UV-B radiation data set for macroecological studies. Methods

Ecol Evol 5:372–383.

Bomberg M, Lamminmäki T, Itävaara M (2016) Microbial communities and their predicted

metabolic characteristics in deep fracture groundwaters of the crystalline bedrock at

Olkiluoto, Finland . Biogeosciences 13:6031–6047.

Buchfink B, Xie C, Huson DH (2014) Fast and sensitive protein alignment using

DIAMOND. Nat Methods 12:59.

Chen B, Yuan K, Chen X, Yang Y, Zhang T, Wang Y, Luan T, Zou S, Li X (2016)

Metagenomic Analysis Revealing Antibiotic Resistance Genes (ARGs) and Their

Genetic Compartments in the Tibetan Environment. Environ Sci Technol 50:6670–

6679.

Doherty M, Yager PL, Moran MA, Coles VJ, Fortunato CS, Krusche A V, Medeiros PM,

Payet JP, Richey JE, Satinsky BM, Sawakuchi HO, Ward ND, Crump BC (2017)

Bacterial Biogeography across the Amazon River-Ocean Continuum. Front Microbiol

8:882.

Eloe-Fadrosh EA, Paez-Espino D, Jarett J, Dunfield PF, Hedlund BP, Dekas AE, Grasby

SE, Brady AL, Dong H, Briggs BR, Li W-J, Goudeau D, Malmstrom R, Pati A, Pett-

Ridge J, Rubin EM, Woyke T, Kyrpides NC, Ivanova NN (2016) Global metagenomic

survey reveals a new bacterial candidate phylum in geothermal springs. Nat Commun

7:10476.

Garcia-Pichel F (1994) A model for internal self-shading in planktonic organisms and its

implications for the usefulness of ultraviolet sunscreens. Limnol Oceanogr 39:1704–

1717.

Graf D, Wesslowski J, Ma H, Scheerer P, Krauß N, Oberpichler I, Zhang F, Lamparter T

(2015) Key Amino Acids in the Bacterial (6-4) Photolyase PhrB from Agrobacterium

fabrum. PLoS One 10:e0140955.

Häder D-P, Gao K (2015) Interactions of anthropogenic stress factors on marine

phytoplankton . Front Environ Sci 3:14.

Häder D-P, Helbling EW, Williamson CE, Worrest RC (2011) Effects of UV radiation on

aquatic ecosystems and interactions with climate change. Photochem Photobiol Sci

10:242–260.

Harmer SL (2009) The Circadian System in Higher Plants. Annu Rev Plant Biol 60:357–

377.

Hauptmann AL, Sicheritz-Pontén T, Cameron KA, Bælum J, Plichta DR, Dalgaard M,

Stibal M (2017) Contamination of the Arctic reflected in microbial metagenomes from

the Greenland ice sheet. Environ Res Lett 12.

Hu B, Wang Y, Liu G (2008) Influences of the clearness index on UV solar radiation for

two locations in the Tibetan Plateau-Lhasa and Haibei. Adv Atmos Sci 25:885–896.

Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ (2010) Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics 11:119.

Jansen MAK, Martret B Le, Koornneef M (2010) Variations in constitutive and inducible

UV-B tolerance; dissecting photosystem II protection in Arabidopsis thaliana

accessions. Physiol Plant 138:22–34.

Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices

from protein sequences. Comput Appl Biosci 8:275–282.

Joux F, Jeffrey WH, Lebaron P, Mitchell DL (1999) Marine bacterial isolates display

diverse responses to UV-B radiation. Appl Environ Microbiol 65:3820–3827.

Karentz D (1995) Considerations for evaluating ultraviolet radiation-induced genetic

damage relative to Antarctic ozone depletion.

Kurth D, Amadio A, Ordoñez OF, Albarracín VH, Gärtner W, Farías ME (2017) Arsenic

metabolism in high altitude modern stromatolites revealed by metagenomic analysis.

Sci Rep 7:1024.

Lefort V, Desper R, Gascuel O (2015) FastME 2.0: A Comprehensive, Accurate, and Fast

Distance-Based Phylogeny Inference Program: Table 1. Mol Biol Evol 32:2798–2800.

Letunic I, Bork P (2016) Interactive tree of life (iTOL) v3: an online tool for the display

and annotation of phylogenetic and other trees. Nucleic Acids Res 44:W242–W245.

Liu C-M, Li D, Sadakane K, Luo R, Lam T-W (2015) MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics 31:1674–1676.

Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve

genome assemblies. Bioinformatics 27:2957–2963.

Mitchell DL, Karentz D (1993) The Induction and Repair of DNA Photodamage in the

Environment. In: *Environmental UV Photobiology*. Springer US, Boston, MA, p 345–

377

Nayfach S, Pollard KS (2015) Average genome size estimation improves comparative

metagenomics and sheds light on the functional ecology of the human microbiome.

Genome Biol 16:51.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR,

O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H (2018)

Vegan R package version 2.5-2. Community Ecol Packag.

Pathak GP, Losi A, Gärtner W (2012) Metagenome-based screening reveals worldwide

distribution of LOV-domain proteins. Photochem Photobiol 88:107–118.

Piccini C, Conde D, Pernthaler J, Sommaruga R (2009) Alteration of chromophoric

dissolved organic matter by solar UV radiation causes rapid changes in bacterial

community composition. Photochem Photobiol Sci 8:1321–1328.

Portero LR, Alonso-Reyes DG, Zannier F, Vazquez MP, Farías ME, Gärtner W, Albarracín

VH (2019) Photolyases and Cryptochromes in UV-resistant Bacteria from High-

altitude Andean Lakes. Photochem Photobiol 95:315–330.

Pushkarev A, Béjà O (2016) Functional metagenomic screen reveals new and diverse

microbial rhodopsins. ISME J 10:2331–2335.

Pushkarev A, Inoue K, Larom S, Flores-Uribe J, Singh M, Konno M, Tomida S, Ito S,

Nakamura R, Tsunoda SP, Philosof A, Sharon I, Yutin N, Koonin E V, Kandori H,

Béjà O (2018) A distinct abundant group of microbial rhodopsins discovered using

functional metagenomics. Nature 558:595–599.

Rascovan N, Maldonado J, Vazquez MP, Eugenia Farías M (2016) Metagenomic study of

red biofilms from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea.

ISME J 10:299–309.

Rinnan R, Keinänen MM, Kasurinen A, Asikainen J, Kekki TK, Holopainen T, Ro-Poulsen

H, Mikkelsen TN, Michelsen A (2005) Ambient ultraviolet radiation in the Arctic

reduces root biomass and alters microbial community composition but has no effects

on microbial biomass. Glob Chang Biol 11:564–574.

Robinson SA, Turnbull JD, Lovelock CE (2005) Impact of changes in natural ultraviolet

radiation on pigment composition, physiological and morphological characteristics of

the Antarctic moss, Grimmia antarctici. Glob Chang Biol 11:476–489.

Robson TM, Pancotto VA, Scopel AL, Flint SD, Caldwell MM (2005) Solar UV-B

influences microfaunal community composition in a Tierra del Fuego peatland. Soil

Biol Biochem 37:2205–2215.

Roenneberg T, Merrow M (2005) Circadian clocks—the fall and rise of physiology. Nat

Rev Mol Cell Biol 6:965–971.

Ruhland CT, Xiong FS, Clark WD, Day TA (2005) The Influence of Ultraviolet-B

Radiation on Growth, Hydroxycinnamic Acids and Flavonoids of Deschampsia

antarctica during Springtime Ozone Depletion in Antarctica¶†. Photochem Photobiol

81:1086–1093.

Russell III JM, Luo M, Cicerone RJ, Deaver LE (1996) Satellite confirmation of the

dominance of chlorofluorocarbons in the global stratospheric chlorine budget. Nature

379:526–529.

S Weber (2005) Light-driven enzymatic catalysis of DNA repair: a review of recent

biophysical studies on photolyase. Elsevier.

Satinsky BM, Fortunato CS, Doherty M, Smith CB, Sharma S, Ward ND, Krusche A V,

Yager PL, Richey JE, Moran MA, Crump BC (2015) Metagenomic and

metatranscriptomic inventories of the lower Amazon River, May 2011. Microbiome

3:39.

Searles PS, Flint SD, Caldwell MM (2001) A meta-analysis of plant field studies

simulating stratospheric ozone depletion. Oecologia 127:1–10.

Selby CP, Sancar A (2006) A cryptochrome/photolyase class of enzymes with single-

stranded DNA-specific photolyase activity. Proc Natl Acad Sci 103:17696–17700.

Singh AH, Doerks T, Letunic I, Raes J, Bork P (2009) Discovering Functional Novelty in

Metagenomes: Examples from Light-Mediated Processes. J Bacteriol 191:32 LP – 41.

Stibal M, Gözdereliler E, Cameron KA, Box JE, Stevens IT, Gokul JK, Schostag M, Zarsky

JD, Edwards A, Irvine-Fynn TDL, Jacobsen CS (2015a) Microbial abundance in

surface ice on the Greenland Ice Sheet. Front Microbiol 6:225.

Stibal M, Schostag M, Cameron KA, Hansen LH, Chandler DM, Wadham JL, Jacobsen CS

(2015b) Different bulk and active bacterial communities in cryoconite from the margin

and interior of the Greenland ice sheet. Environ Microbiol Rep 7:293–300.

Toh H, Kanai S, Todo T, Kikuno R, Ryo H (1997) Molecular evolution of the photolyase–

blue-light photoreceptor family. J Mol Evol 45:535–548.

Tran P, Ramachandran A, Khawasik O, Beisner BE, Rautio M, Huot Y, Walsh DA (2018)

Microbial life under ice: Metagenome diversity and in situ activity of Verrucomicrobia

in seasonally ice-covered Lakes. Environ Microbiol 20:2568–2584.

Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, Tett A,

Huttenhower C, Segata N (2015) MetaPhlAn2 for enhanced metagenomic taxonomic

profiling. Nat Methods 12:902.

Tschitschko B, Erdmann S, DeMaere MZ, Roux S, Panwar P, Allen MA, Williams TJ,

Brazendale S, Hancock AM, Eloe-Fadrosh EA, Cavicchioli R (2018) Genomic

variation and biogeography of Antarctic haloarchaea. Microbiome 6:113.

Ueda T, Kato A, Kuramitsu S, Terasawa H, Shimada I (2005) Identification and

characterization of a second chromophore of DNA photolyase from Thermus

thermophilus HB27. J Biol Chem 280:36237–36243.

Westbrook A, Ramsdell J, Bergeron RD, Normington L, MacManes MD, Thomas WK,

Schuelke T (2017) PALADIN: protein alignment for functional profiling whole

metagenome shotgun data. Bioinformatics 33:1473–1478.

Williamson CE, Neale PJ, Hylander S, Rose KC, Figueroa FL, Robinson SA, Häder DP,

Wängberg S, Worrest RC (2019) The interactive effects of stratospheric ozone

depletion, UV radiation, and climate change on aquatic ecosystems. Photochem

Photobiol Sci 18:717–746.

Xiong FS, Day TA (2001) Effect of solar ultraviolet-B radiation during springtime ozone

depletion on photosynthesis and biomass production of Antarctic vascular plants. Plant

Physiol 125:738–751.

Yan A, Pan J, An L, Gan Y, Feng H (2012) The response of trichome mutants to enhanced

ultraviolet-B radiation in Arabidopsis thaliana.

Ye Z, Zhang N, Wu C, Zhang X, Wang Q, Huang X, Du L, Cao Q, Tang J, Zhou C, Hou S,

He Y, Xu Q, Xiong X, Kijlstra A, Qin N, Yang P (2018) A metagenomic study of the

gut microbiome in Behcet's disease. Microbiome 6:135.

Zaller JG, Caldwell MM, D. Flint S, Scopel AL, Salo OE, Ballaré CL (2002) Solar UV-B

radiation affects below-ground parameters in a fen ecosystem in Tierra del Fuego,

Argentina: implications of stratospheric ozone depletion. Glob Chang Biol 8:867–871.

Zhang F, Scheerer P, … IO-P of the, 2013 U (2013) Crystal structure of a prokaryotic (6-4)

photolyase with an Fe-S cluster and a 6, 7-dimethyl-8-ribityllumazine antenna

chromophore. Natl Acad Sci.

Tables

Table 1. General traits of the metagenomes used in the current study.

Metagenome	Description	Depth of sampling (cm)	Reference	Size of Raw Metagenomic Data (Gb)	Size after merging and quality filtering (Gb)	N° of assembled Contigs (>999 bp)	Assembly size (bp)	N° of annotated proteins	N° of proteins per Mb
Lake Diamante sediment	Red biofilm from Lake Diamante, located inside the Galan Volcano crater at 4589 m a.s.l.	-	(Rascovan et al. 2016)	0,74	0,64	19537	30047993	47731	1588,492
Socompa stromatolite	Sample from a high-altitude stromatolite in the surroundigns of Lake Socompa	0—5	(Kurth et al. 2017)	11,4	7,57	77652	190723415	219571	1151,254
Tibetan Plateau sediment	Sediment sample collected from Qiangyong Glacier Lake	-	(Chen et al. 2016)	3,4	2,59	42453	80284053	105553	1314,744
Amazon River	Water sample from the Amazon River at a site facing Óbidos city, Brazil.	0—3300	(Satinsky et al. 2015)	9,1	5,7	41430	71904038	108943	1515,117
Lake Rauer	Saline lake microbial sample from Rauer Islands, Antarctica.	10—30	(Tschitschko et al. 2018)	5,1	5,01	64101	177770567	216623	1218,554
Dewar Creek hot spring	Hot spring sediment metagenome from Dewar Creek, Canada.	-	(Eloe-Fadrosh et al. 2016)	9,9	9,47	26840	162485558	169888	1045,558
Greenland cryoconite	Cryoconite sample from Greenland ice sheet	20	(Hauptmann et al. 2017)	8,7	7,77	45376	150415849	173231	1151,68
Lake Montjoie	Freshwater microbial communities from epilimnion strata of Lake Montjoie, Canada.	0—300	(Tran et al. 2018)	14,2	13,66	240051	545748448	693357	1270,47
Olkiluoto Island groundwater	Groundwater microbial communities from the Olkiluoto Island deep subsurface site, Finland	>29600	(Bomberg et al. 2016)	24,9	8,85	74876	159143965	231244	1453,049
Human gut	Metagenome of fecal sample collected from an active Behet's disease patient.	-	(Ye et al. 2018)	10	9,91	47652	205076299	210369	1025,808

Figures

Figure 1. Worldwide metagenomic sampling linked to UV-B georeferenced data obtained

from glUV datasets (https://www.ufz.de/gluv/). The metagenomes were selected from

NCBI using biosample information: DM, Lake Diamante; SS, Socompa stromatolite; TB,

Tibetan Plateau sediment; AM, Amazon River; RA, Lake Rauer; HT, Dewar Creek hot

spring; CR, Greenland cryoconite; MT, Lake Montjoie; OK, Olkiluoto Island groundwater.

Figure 2. Scatterplots showing the impact of UV-B on the microbial diversity in the whole

metagenomic dataset. Two parameters were measured: species richness (A) and the

Shannon index (B). Metagenomes were also colored according the group they belong,

being red for UV_{High}, green for UV_{Mid} and blue for UV_{Low}. Abbreviations: DM, Lake

Diamante; SS, Socompa stromatolite; TB, Tibetan Plateau sediment; AM, Amazon River;

RA, Lake Rauer; HT, Dewar Creek hot spring; CR, Greenland cryoconite; MT, Lake

Montjoie; OK, Olkiluoto Island groundwater; GU, human gut.

Figure 3. Scatterplots showing the impact of UV-B (A) and Day length (B) on the CPF

abundance in the whole metagenomic dataset. Metagenomes were also colored according

the group they belong, being red for UV_{High}, green for UV_{Mid} and blue for UV_{Low}.

Abbreviations: DM, Lake Diamante; SS, Socompa stromatolite; TB, Tibetan Plateau

sediment; AM, Amazon River; RA, Lake Rauer; HT, Dewar Creek hot spring; CR,

Greenland cryoconite; MT, Lake Montjoie; OK, Olkiluoto Island groundwater; GU, human

gut.

Figure 4. Phylogenetic tree built with the CPF sequences from the whole metagenomic

dataset. Each leaf is colored and labeled by sample procedence: DM, Lake Diamante

(purple); SS, Socompa stromatolite (dark green); TB, Tibetan Plateau sediment (brown);

AM, Amazon River (blue); RA, Lake Rauer (grey); HT, Dewar Creek hot spring (orange);

CR, Greenland cryoconite (cyan); MT, Lake Montjoie (red); OK, Olkiluoto Island

groundwater (light green). Abbreviations: CPD I, II and III refers to class I, II and III

cyclobutane pirimidine dimer photolyases; CPD THR, thermostable CPD photolyase;

DASH, DASH cryptochrome; FeS-BCPs, FeS bacterial cryptochromes and photolyases;

UI, II and III refers to unidentified group I, II and III respectively.

Figure 5. Distribution of each subfamily by metagenome. Abbreviations: DM, Lake

Diamante; SS, Socompa stromatolite; TB, Tibetan Plateau sediment; AM, Amazon River;

RA, Lake Rauer; HT, Dewar Creek hot spring; CR, Greenland cryoconite; MT, Lake

Montjoie; OK, Olkiluoto Island groundwater; GU, human gut.

Figure 6. Comparative barplot showing for each metagenome the genus diversity both for

whole microbial community (top bar) and the set of CPF sequences recovered from the

same metagenome (lower bar). The diversity of the microbial community was estimated

with Metaphlan, while the diversity of the set of CPF sequences were computed with Blast

online platform. Abbreviations: DM, Lake Diamante; SS, Socompa stromatolite; TB,

Tibetan Plateau sediment; AM, Amazon River; RA, Lake Rauer; HT, Dewar Creek hot

694   spring; CR, Greenland cryoconite; MT, Lake Montjoie; OK, Olkiluoto Island groundwater;

695   GU, human gut.