Presynaptic Gα$_o$ (GOA-1) signals to depress command neuron excitability and allow

stretch-dependent modulation of egg laying in *Caenorhabditis elegans*

Bhavya Ravi[1,2,3], Jian Zhao[4,5], Sana Chaudhry[2], Mattingly Bartole[1,2], Richard J. Kopchock 3rd [2],

Christian Guijarro[2,6], Lijun Kang[4], and Kevin M. Collins[1,2,7]

[1]Neuroscience Program, University of Miami Miller School of Medicine, Miami, FL USA

[2]Department of Biology, University of Miami, Coral Gables, FL USA

[3]Present address: Department of Neurology, Johns Hopkins University School of Medicine,
Baltimore, MD USA

[4]Department of Neuroscience, Zhejiang University School of Medicine, Hangzhou, Zhejiang,
China

[5]Present address: Department of Molecular Biology, Massachusetts General Hospital, Boston,
MA USA

[6]Present address: ALPS Electric Europe GmbH, Munich, Germany

[7]To whom correspondence should be addressed:

kevin.collins@miami.edu

Department of Biology
University of Miami
1301 Memorial Drive
Coral Gables, FL 33143
Tel: (305) 284-9058
21

Short title: Presynaptic Gα$_o$ inhibits command neuron excitability

54 pages with 7 Figures, 2 Tables, and 7 Videos

Abstract- 205 words, Introduction- 648 words, Discussion- 1483 words.

Abstract

Egg laying in the nematode worm *Caenorhabditis elegans* is a two-state behavior modulated

by internal and external sensory input. We have previously shown that homeostatic feedback

of embryo accumulation in the uterus regulates bursting activity of the serotonergic HSN

command neurons that sustains the egg-laying active state. How sensory feedback of egg

release signals to terminate the egg-laying active state is less understood. We find that $G\alpha_o$, a

conserved Pertussis Toxin-sensitive G protein, signals within HSN to inhibit egg-laying circuit

activity and prevent entry into the active state. $G\alpha_o$ signaling hyperpolarizes HSN, reducing

HSN Ca^{2+} activity and input onto the postsynaptic vulval muscles. Loss of inhibitory $G\alpha_o$

signaling uncouples presynaptic HSN activity from a postsynaptic, stretch-dependent

homeostat, causing precocious entry into the egg-laying active state when only a few eggs are

present in the uterus. Feedback of vulval opening and egg release activates the uv1

neuroendocrine cells which release NLP-7 neuropeptides which signal to inhibit egg laying

through $G\alpha_o$-independent mechanisms in the HSNs and $G\alpha_o$-dependent mechanisms in cells

other than the HSNs. Thus, neuropeptide and inhibitory $G\alpha_o$ signaling maintains a bi-stable

state of electrical excitability that dynamically controls circuit activity in response to both

external and internal sensory input to drive a two-state behavior output.

Introduction

A major goal of neuroscience is to understand how sensory signals control neural circuit activity and changes in animal behavior. Such sensory feedback informs when a behavior should begin, how long it should continue, and when it should end. Extensive evidence has shown that neuromodulators like serotonin and neuropeptides signal through G protein coupled receptors to remodel neural circuit activity and drive behavior state transitions (Jiang et al., 2001; Goulding et al., 2008). Yet, there is no neural circuit in any organism for which we know precisely how each signaling event contributes to sensory modulation of a behavior. Small neural circuits typically found in invertebrates combine anatomical simplicity with genetic and experimental accessibility, allowing for a complete understanding of the molecular and physiological basis for a behavioral output (Marder, 2012).

The *C. elegans* female egg-laying behavior circuit is ideally suited to study how sensory signals modulate circuit functions that underlie decision-making. As shown in Figure 1A, the circuit is comprised of two Hermaphrodite Specific Neurons (HSNs) that synapse onto the egg-laying vulval muscles (White et al., 1986; Shen et al., 2004; Li et al., 2013). During ~2 minute active states, rhythmic HSN Ca^{2+} activity releases serotonin and neuropeptides that signal to promote the excitability of the muscles, driving ejection of ~4-6 eggs in sequence from the uterus into the environment (Waggoner et al., 1998; Shyn et al., 2003; Zhang et al., 2008; Collins et al., 2016; Brewer et al., 2019). External and internal sensory input regulate the onset of egg laying (Horvitz et al., 1982; Trent, 1982; Sawin, 1996; Aprison and Ruvinsky, 2014; Ravi et al., 2018a), and genetic studies have identified neuropeptides, receptors, and two antagonistic heterotrimeric G proteins that signal to regulate HSN activity, neurotransmitter release, and egg laying (Schafer, 2006; Koelle, 2016). $G\alpha_q$ signals through the conserved PLCβ and Trio RhoGEF effector pathways to promote neurotransmitter release and egg laying (Brundage et

al., 1996; Lackner et al., 1999; Miller et al., 1999; Bastiani et al., 2003; McMullan et al., 2006; Williams et al., 2007; McMullan and Nurrish, 2011). Because phorbol ester DAG-mimetics such as PMA rescue synaptic transmission defects of $G\alpha_q$ signaling mutants (Lackner et al., 1999; Williams et al., 2007), DAG production and recruitment of UNC-13 and/or Protein Kinase C effectors is thought to mediate the $G\alpha_q$ dependent modulation of synaptic transmission (Yawo, 1999; Wierda et al., 2007; Lou et al., 2008). $G\alpha_q$ signaling is opposed by $G\alpha_o$ which signals to inhibit neurotransmitter release (Koelle and Horvitz, 1996; Miller et al., 1999; Nurrish et al., 1999), vulval muscle activity (Shyn et al., 2003), and egg laying (Mendel et al., 1995; Segalat et al., 1995). How $G\alpha_o$ signaling antagonizes $G\alpha_q$, Rho, and DAG signaling is not clear. Even though $G\alpha_o$ mediates signaling by numerous neurotransmitters in diverse animals including mammals (Jiang et al., 2001), direct effectors for $G\alpha_o$ have not yet been identified. In *C. elegans*, $G\alpha_o$ mutants resemble animals lacking DAG Kinase; both mutants show increased UNC-13 localization to synapses and hyperactive egg-laying behavior defects, resembling treatment with phorbol esters (Miller et al., 1999; Nurrish et al., 1999; Jose and Koelle, 2005). Because synaptic transmission could be modulated by upstream changes in cell and circuit electrical excitability and/or by downstream effects on synaptic vesicle fusion, what is needed are direct measurements of how discrete changes in G protein and effector signaling affect cell and circuit activity and their consequences on animal behavior.

Here we explore how $G\alpha_o$ inhibits *C. elegans* egg-laying circuit activity and behavior. Our data reveal that $G\alpha_o$ signaling reduces the electrical excitability of the HSN command neurons to promote the inactive state while eggs are produced. Feedback of egg accumulation in the uterus then switches the circuit into active states where rhythmic bursting drives sequential egg-laying events. Thus, modulation by inhibitory $G\alpha_o$ signaling allows for proper

induction of motor behavior circuit upon the alignment of optimal external and internal sensory

conditions.

Results

Reduced inhibitory Gα_o signaling leads to premature egg laying and decreases the

duration of egg-laying inactive states

To better understand how inhibitory Gα_o signaling contributes to egg-laying behavior

state transitions, we used mutants and transgenes to manipulate Gα_o signaling in specific cells

of the egg-laying circuit and analyzed the consequences on cell Ca^{2+} activity and behavior. We

found that Gα_o signaling inhibits the onset of egg laying. We performed a 'time to first egg lay'

assay in wild-type animals and in mutants with excessive or reduced Gα_o signaling. As

previously shown, wild-type animals release their first embryo ~6-7 hours after becoming

adults (McMullen et al., 2012; Ravi et al., 2018a). Animals bearing Gα_o loss-of-function or null

mutations laid their eggs much earlier, 3-4 hours after becoming adults (Figure 1B). *goa-*

1(n1134), a hypomorphic mutant predicted to lack the conserved N-terminal myristoylation and

palmitoylation sequence, and *goa-1(sa734)*, an early stop mutant predicted to be a molecular

null (Segalat et al., 1995; Robatzek and Thomas, 2000), showed similarly precocious onset of egg

laying (Figure 1B). This phenotype was shared in transgenic animals where Gα_o function was

inhibited just in HSNs through the cell-specific expression of Pertussis Toxin (Tanis et al.,

2008). Because the timing of this first egg-laying event requires serotonin along with HSN

function and activity (Ravi et al., 2018a), these results suggest that Gα_o normally signals in HSN

to inhibit neurotransmitter release and thereby delay the first egg-laying active state (Figure

1B). To test the effects of increased $G\alpha_o$ signaling, we analyzed the behavior of *egl-10(md176)*

mutants which lack the major RGS protein that terminates $G\alpha_o$ signaling by promoting $G\alpha_o$

GTP hydrolysis (Koelle and Horvitz, 1996). *egl-10(md176)* mutants showed a strong and

significant delay in the onset of egg laying, laying their first egg ~15 hours after reaching

adulthood (Figure 1B). This delay of egg-laying phenotype was shared in transgenic animals

expressing the constitutively active $G\alpha_o$ (Q205L) mutant specifically in the HSNs (Tanis et al.,

2008) and resembled animals developmentally lacking HSNs (Ravi et al., 2018a), consistent

with $G\alpha_o$ signaling in HSN to inhibit neurotransmitter release that normally drives the onset of

egg laying.

Figure 1. $G\alpha_o$ signaling maintains the inactive egg-laying behavior state.

(**A**) Cartoon of how identified and unidentified neuropeptides released from AWC (Fenk and de
Bono, 2015), BAG (Ringstad and Horvitz, 2008), uv1 (Banerjee et al., 2017), and other sensory
cells bind to G-protein coupled receptors expressed on HSN command neurons (green) which

signal via Gα_o (red) or Gα_q (black) effector pathways to modulate serotonin and NLP-3
neuropeptide release (Tanis et al., 2008; Brewer et al., 2019). The egg-laying vulval muscles
(orange) express receptors for serotonin (Carnell et al., 2005; Dempsey et al., 2005; Hobson et
al., 2006; Fernandez et al., 2020) and possibly NLP-3 which signal to promote vulval muscle
excitability and egg laying (**B**) Scatter plots of the first egg-laying event in wildtype (grey), null
goa-1(sa734) mutants (orange), hypomorphic loss-of-function *goa-1(n1134)* mutants (red), *egl-*
10(md176) null mutants (purple), and transgenic animals expressing Pertussis Toxin (blue) or
GOA-1^{Q205L} in the HSNs (pink). Error bars show 95% confidence intervals for the mean from
≥10 animals. Asterisks indicate p≤0.0001 (One-way ANOVA with Bonferroni correction for
multiple comparisons). (**C**) Representative raster plots showing temporal pattern of egg laying
during three hours in wild-type (black), hypomorphic loss-of-function *goa-1(n1134)* mutant
(red), and *egl-10(md176)* null mutant animals (purple), along with transgenic animals
expressing Pertussis Toxin (blue) or GOA-1^{Q205L} in the HSNs (pink). Vertical lines indicate
single egg-laying events.

To understand how Gα_o signaling controls the normal two-state pattern of egg laying,

we performed long-term recordings of adults as they transitioned into and out of the active

states in which clusters of several eggs are typically laid. We defined intra-cluster intervals as

the time elapsed between consecutive egg-laying events within a single active state (e.g.

intervals < 4 minutes) and inter-cluster intervals as the time elapsed between different active

states (e.g. intervals > 4 minutes), as described (Waggoner et al., 1998; Collins and Koelle,

2013). As expected, wild-type animals displayed a two-state pattern of egg laying with multiple

egg-laying events clustered within brief, ~2-minute active states about every 20-30 minutes

(Figure 1C and Table 1). Animals bearing the *goa-1(n1134)* hypomorphic mutation entered

active states 2-3-fold more frequently, often laying single eggs during active states separated

by only ~12-13 minutes (Figure 1C and Table 1), as previously shown (Waggoner et al., 2000).

We found the duration of inactive states in animals expressing Pertussis Toxin in the HSN

neurons was indistinguishable from the *goa-1(n1134)* mutant, indicating that Gα_o signals in

HSN to reduce the probability of entering the active state (Figure 1C and Table 1). Loss of

inhibitory Gα_o signaling led to active states in which 1-2 embryos were laid almost immediately

after they were positioned in the uterus next to the vulval opening. As a result, successive egg-

laying events were rate-limited by egg production, explaining why the average intra-cluster

intervals were typically double seen in wild-type animals (Figure 1C and Table 1). Conversely,

egl-10(md176) mutant animals or transgenic animals expressing the $G\alpha_o$(Q205L) gain-of-

function mutation in the HSNs showed prolonged inactive periods of 258 and 67 min,

respectively (Figure 1C and Table 1), resembling animals lacking HSNs altogether (Waggoner

et al., 2000). Interestingly, animals without HSNs or with too much inhibitory $G\alpha_o$ signaling still

lay multiple eggs within discrete active states (Figure 1C and Table 1), consistent with our

results showing that a stretch-dependent homeostat can maintain the active state even when

neurotransmitter release from the HSN is inhibited (Collins et al., 2016; Ravi et al., 2018a).

These results show that $G\alpha_o$ signaling does not modulate patterns of egg laying within active

states. Instead, $G\alpha_o$ acts in HSN to depress entry into the egg-laying active state.

Genotype	*Intra*-cluster interval (1/λ₁) Average (min) (95%CI range)	*Inter*-cluster interval (1/λ₂) Average (min) (95%CI range)	Eggs laid per active state Average ± 95%CI	Unlaid eggs per animal Average ± 95%CI
Wild type	0.42 (0.33-0.51)	22.11 (21.19-23.09)	2.5 ± 0.5	15.9 ± 1.5
goa-1(n1134)	1.31 (0.94-1.67)	12.88* (12.31-12.75)	1.2 ± 0.1‡	2.2 ± 0.3
HSN::Pertussis Toxin	1.69 (1.17-2.20)	12.42* (12.09-12.75)	1.1 ± 0.07‡	4.1 ± 0.3#
egl-10(md176)	0.64 (0.49-0.79)	258.6* (231.26-293.42)	2.5 ± 0.2	45.9 ± 3.3
HSN::GOA-1^{Q205L}	0.35 (0.24-0.47)	66.97* (64.14-70.02)	2.2 ± 0.2	36.8 ± 3.8

Table 1. Egg-laying behavior measurements in animals with altered $G\alpha_o$ signaling.

Long-term behavior recordings were used to extract features of egg-laying active and inactive
behavior states for the indicated genotypes, as described (Waggoner et al., 1998). '*' indicates
significant differences compared to wild-type animals (p<0.0001, Kruskal-Wallis test with
Dunn's correction for multiple comparisons), '‡' indicates significant differences compared to

wildtype (p<0.0001, One-way ANOVA with Bonferroni correction), and '#' indicates that this
result was previously reported (Tanis et al., 2008).

$G\alpha_o$ signaling inhibits HSN Ca^{2+} activity to promote the inactive behavior state

Previous work has shown that $G\alpha_o$ signals presynaptically to depress neurotransmitter

release. Whether $G\alpha_o$ signals to inhibit HSN electrical excitability that might be evident in

changes in cell Ca^{2+} activity, or instead signals downstream of Ca^{2+} to regulate steps in UNC-

13-dependent docking, priming, and vesicle fusion, has not been tested directly. To address

how $G\alpha_o$ signals to inhibit HSN neurotransmitter release, we performed ratiometric Ca^{2+}

imaging in our panel of $G\alpha_o$ signaling mutants as they entered spontaneous egg-laying active

states. Freely behaving animals bearing the *goa-1(n1134)* hypomorphic or *goa-1(sa734)* null

mutations showed a clear change in HSN Ca^{2+} activity from two-state burst firing to more tonic

firing (Figure 2A, Videos 1-3). Complete loss of inhibitory $G\alpha_o$ signaling caused a significant

increase in the frequency of HSN Ca^{2+} transients (Figure 2B and 2C). We were surprised that

the *goa-1(n1134)* mutants, which show strongly hyperactive egg-laying behavior

indistinguishable from that of *goa-1(sa734)* null mutants, showed only a modest and

statistically insignificant increase in HSN Ca^{2+} activity compared to wild-type (Figure 2C).

However, our results with *goa-1(n1134)* match previously findings (Shyn et al., 2003). The *goa-*

1(n1134) hypomorphic mutant is expected to have residual $G\alpha_o$ signaling activity in that its

major defect is the absence of a proper membrane anchor sequence (Mumby et al., 1990).

These results show that $G\alpha_o$ signaling depresses HSN presynaptic activity, but that the

hyperactive egg-laying phenotypes observed in *goa-1(n1134)* mutants are separable from an

increase in presynaptic HSN Ca^{2+} activity.

Figure 2. Gα₀ signaling inhibits HSN neuron Ca²⁺ activity and burst firing.

(**A**) Representative GCaMP5:mCherry (ΔR/R) ratio traces showing HSN Ca²⁺ activity in freely behaving wild-type (black), *goa-1(n1134)* loss-of-function mutant (green), *goa-1(sa734)* null mutant (blue), and *egl-10(md176)* null (purple) mutant animals, along with transgenic animals expressing the activated GOA-1(Q205L) in the HSN neurons (pink) during an egg-laying active state. Arrowheads indicate egg-laying events. (**B**) Cumulative distribution plots of instantaneous Ca²⁺ transient peak frequencies (and inter-transient intervals) in wild-type (black open circles), *goa-1(n1134)* (green filled triangles), *goa-1(sa734)* (blue squares), *egl-10(md176)* mutants (purple open squares) along with transgenic animals expressing the activated GOA-1(Q205L) in the HSN neurons (pink open circles). Asterisks indicate $p<0.0001$ (Kruskal-Wallis test with Dunn's correction for multiple comparisons). (**C**) Scatter plots show average Ca²⁺ transient frequency (per min). Error bars indicate 95% confidence intervals for the mean. Asterisk indicates $p<0.0001$; pound (#) indicates $p≤0.0079$; n.s. indicates $p>0.05$ (One-way ANOVA with Bonferroni correction for multiple comparisons).

We next tested how increased inhibitory Gα₀ signaling affects HSN activity. Both *egl-10(md176)* mutants and transgenic animals expressing the activated GOA-1(Q205L) in HSNs showed a significant and dramatic reduction in the frequency of HSN Ca²⁺ transients, with single HSN Ca²⁺ transients occuring several minutes apart (Figure 2A and 2B). The rare egg-laying events seen in animals with increased Gα₀ signaling mostly followed single HSN Ca²⁺ transients, not multi-transient bursts typically seen in wild-type animals (Figure 2A and 2C). In one *egl-10(md176)* animal, we observed an egg-laying event that was not accompanied by an

HSN Ca^{2+} transient. This suggests that elevated $G\alpha_o$ signaling is sufficient to silence the

HSNs, and that, in this case, egg laying becomes HSN-independent. In support of this model,

previous work has shown that *egl-10(md176)* mutants respond to serotonin but are resistant to

the serotonin reuptake inhibitor imipramine (Trent et al., 1983). Alternatively (or additionally)

$G\alpha_o$ signaling may function to depress coordinated activity between the gap-junctioned,

contralateral HSNs, whose Ca^{2+} activity we were unable to observe simultaneously because

our confocal imaging conditions only captures one HSN at a time.

To determine how disruption of inhibitory $G\alpha_o$ signaling in HSN affects its activity, we

recorded HSN Ca^{2+} transients in transgenic animals expressing Pertussis Toxin in the HSNs.

$G\alpha_o$-silenced HSNs showed a dramatic increase in the frequency of HSN Ca^{2+} transients,

leading to a nearly constitutive, tonic Ca^{2+} activity like that observed in *goa-1(sa734)* null

mutants (Figure 3A, 3B, and 3C; Video 4). While control animals showed an average HSN

Ca^{2+} transient frequency of about ~0.4 transients per minute, animals expressing Pertussis

Toxin in HSN showed an average 1.9 transients per minute, a significant increase (Figure 3C).

These results suggest that even under normal growth conditions, unidentified neuromodulators

signal through $G\alpha_o$-coupled receptors on HSN to inhibit Ca^{2+} activity. We hypothesize that the

observed changes in presynaptic UNC-13 localization (Nurrish et al., 1999) and serotonin

biosynthesis (Tanis et al. 2008) when inhibitory $G\alpha_o$ signaling is reduced are consequences,

not causes, of altered presynaptic activity.

Figure 3. Inhibitory Gα$_o$ signaling in HSN is required for two-state Ca^{2+} activity and facilitates modulation by feedback of egg accumulation.

(A) Representative GCaMP5:mCherry (ΔR/R) ratio traces showing HSN Ca^{2+} activity in untreated (fertile) wild-type animals (black open circles), FUDR-sterilized wild-type animals (black filled circles), untreated (fertile) animals expressing Pertussis Toxin (PTX) in the HSN neurons (blue open squares), and in FUDR-sterilized transgenic animals expressing PTX in the HSNs (blue filled squares). Arrowheads indicate egg-laying events. **(B)** Cumulative distribution plots of instantaneous Ca^{2+} transient peak frequencies (and inter-transient intervals) in untreated and FUDR-treated animals. Asterisks indicate p<0.0001 (Kruskal-Wallis test with Dunn's test for multiple comparisons). **(C)** Scatter plots show average Ca^{2+} transient frequency (per min). Error bars indicate 95% confidence intervals for the mean. Asterisks indicate p<0.0001 (One-way ANOVA with Bonferroni's test for multiple comparisons). Data from 10 animals were used for each strain for analysis.

We have previously shown that burst Ca^{2+} activity in the command HSN neurons is

initiated and sustained by a stretch-dependent homeostat. In chemically or genetically

sterilized animals, burst Ca^{2+} activity in HSN is largely eliminated ([Ravi et al., 2018a](#)). As such,

we were surprised to observe high frequency Ca^{2+} transients in animals with reduced Gα$_o$

signaling because these animals typically retain few (1 to 3) eggs in the uterus at steady state,

conditions that normally eliminate HSN burst firing. We hypothesized that the stretch-

dependent homeostat was not required to promote HSN Ca^{2+} activity in Gα$_o$ signaling mutants.

To test this, we chemically sterilized transgenic animals expressing Pertussis Toxin in the

HSNs with Floxuridine (FUDR), a blocker of embryogenesis ([Mitchell et al., 1979](#)), and

recorded HSN Ca^{2+} activity. Wild-type animals treated with FUDR showed a dramatic decrease

in the frequency of HSN Ca^{2+} activity and an elimination of burst firing (Figures 3A-C), as we

have previously shown (Ravi et al., 2018a). Sterilized transgenic animals expressing Pertussis

Toxin in the HSNs showed only a slight reduction in HSN Ca^{2+} frequency (Figure 3A and 3B).

Both fertile and sterile Pertussis Toxin-expressing animals had dramatically and significantly

increased HSN Ca^{2+} transient frequency (~1.9 / min), indicating their HSNs no longer

responded to the retrograde signals of egg accumulation arising from the stretch-homeostat.

One explanation for this result could be that in fertile wild-type animals, feedback of egg

accumulation elevates HSN excitability above a firing threshold that overcomes endogenous

inhibitory Gα_o signaling. Together, these results support a model where Gα_o signals in HSN to

depress cell electrical excitability which allows for the proper two-state pattern of HSN Ca^{2+}

activity that responds to homeostatic feedback of egg accumulation.

Presynaptic Gα_o signaling inhibits postsynaptic vulval muscle activity

To test how changes in inhibitory Gα_o signaling affect the postsynaptic vulval muscles, we

recorded Ca^{2+} activity in the vulval muscles of *goa-1(n1134)* mutant and Pertussis Toxin

expressing transgenic animals. Both wild-type and *goa-1(n1134)* mutants showed increased

vulval muscle Ca^{2+} transients as the animals entered the egg-laying active state (Figure 4A

and 4B). *goa-1(n1134)* mutants also showed elevated muscle activity compared to wild-type

control animals during inactive states when no eggs were laid (Figure 4A and 4B), as

previously shown (Shyn et al., 2003). Surprisingly, Ca^{2+} activity during egg-laying active states

was not significantly different in *goa-1(n1134)* mutant animals (Figure 4A and 4B; compare

Videos 5 and 6), a result consistent with Gα_o signaling primarily to prolong the inactive state.

These results suggest Gα$_o$ signals to inhibit circuit activity during the egg-laying inactive state

and/or that loss of Gα$_o$ elevates circuit activity that prolongs the active state beyond the brief

~2-3 minute window during which eggs are typically laid (Waggoner et al., 1998). Recordings of

vulval muscle Ca^{2+} activity in transgenic animals expressing Pertussis Toxin in the presynaptic

HSNs show a dramatic and significant increase in vulval muscle Ca^{2+} transient frequency

during both active and inactive states (Figure 4C and 4D; Video 7), confirming Gα$_o$ is required

in HSN to inhibit neurotransmitter release.

Figure 4. Gα$_o$ signals in HSN to reduce excitatory modulation of the vulval muscles.

(A) Representative GCaMP5:mCherry (ΔR/R) ratio traces showing vulval muscle Ca^{2+} activity
in wild-type (black) and *goa-1(n1134)* loss-of-function mutant animals (red). Egg-laying events
are indicated by arrowheads, and egg-laying active states are indicated by dashed green lines.
(B) Cumulative distribution plots of instantaneous vulval muscle Ca^{2+} transient peak
frequencies (and inter-transient intervals) in wild-type (black circles) and *goa-1(n1134)* mutant
animals (red squares) in the inactive and active egg-laying states (filled and open,
respectively). **(C)** Representative GCaMP5:mCherry (ΔR/R) ratio traces showing vulval muscle

Ca^{2+} activity in untreated (circles) and FUDR-treated (cross) wild-type animals (black) along
with untreated (filled) or FUDR-treated (open) transgenic animals expressing Pertussis Toxin
in the HSNs (blue squares). Egg-laying events are indicated by arrowheads, and egg-laying
active states are indicated by green dashed lines. **(D)** Cumulative distribution plots of
instantaneous vulval muscle Ca^{2+} transient peak frequencies (and inter-transient intervals) in
wildtype and in transgenic animals expressing Pertussis Toxin in the HSN neurons, with or
without FUDR treatment (black cross or blue square), during the inactive and active egg-laying
states (filled and open, respectively). Asterisks indicate p<0.0001 (Kruskal-Wallis test with
Dunn's test for multiple comparisons).

We have previously shown that egg accumulation promotes vulval muscle excitability

during the active state while sterilization reduces it to that seen during the inactive state

(Collins et al., 2016; Ravi et al., 2018a). Because FUDR-sterilization failed to reduce HSN Ca^{2+}

activity in Pertussis Toxin expressing animals (Figure 3C), we predicted that vulval muscle

Ca^{2+} activity would remain similarly elevated in these animals after sterilization. To our

surprise, FUDR treatment significantly reduced, but did not completely eliminate, the elevated

vulval muscle Ca^{2+} activity in animals expressing Pertussis Toxin in HSN (Figure 4C and 4D).

This result suggests egg accumulation and/or germline activity is still required for full vulval

muscle activity even when HSN Ca^{2+} activity is dramatically increased. However, because

these animals lay eggs almost as soon as they are made, the degree of uterine stretch

necessary to induce the active state must be markedly reduced. This supports a model where

both the stretch-dependent homeostat and Gα_o signaling dynamically interact to regulate egg-

laying behavior states.

Gα_o signaling modulates the HSN resting membrane potential

Reduction of inhibitory Gα_o signaling strongly increased HSN Ca^{2+} activity and burst

firing, prompting us to investigate whether Gα_o signals to modulate HSN electrical excitability.

We recorded the resting membrane potential of the HSN neurons in animals with altered $G\alpha_o$

signaling using the whole-cell patch clamp technique (Figure 5A), as described (Yue et al.,

2018). Hypomorphic *goa-1(n1134)* loss-of-function mutants displayed a trend towards more

depolarized resting potentials (-17.9 mV) compared to HSNs from recorded wild-type animals

(-21.1 mV), but this difference was not statistically significant (Figure 5B). In contrast, the

resting membrane potential of HSNs in *egl-10(md176)* $G\alpha_o$ RGS protein mutant animals with a

global increase in $G\alpha_o$ signaling (Koelle and Horvitz, 1996) was significantly hyperpolarized (-

40.8 mV) compared to wild-type control animals. This hyperpolarization of HSNs in *egl-*

10(md176) mutants explains the reduced frequency of HSN Ca^{2+} transients and their strong

defects in egg-laying behavior. Transgenic animals expressing Pertussis Toxin specifically in

the HSNs had significantly depolarized HSNs (-14.75 mV) compared to the wild-type parental

strain (-21.8 mV). These results show that $G\alpha_o$ signals in the HSNs to promote membrane

polarization, reducing cell electrical excitability, Ca^{2+} activity, and neurotransmitter release.

Inhibition of egg laying by $G\alpha_o$ is not replicated by elevated $\beta\gamma$ expression

Receptor activation of $G\alpha_{i/o}$ heterotrimers releases $\beta\gamma$ subunits which have previously

been shown to bind to activate specific K^+ channels and inhibit Ca^{2+} channels (Reuveny et al.,

1994; Herlitze et al., 1996). RNAi-mediated knockdown of the GPB-1 β subunit in HSN reduces

egg-laying behavior (Esposito et al., 2007). An increase in free $\beta\gamma$ subunits in $G\alpha_o$ mutants

might explain their hyperactive egg-laying behavior phenotypes. To test if $\beta\gamma$ over-expression

in HSN would increase egg laying, we transgenically overexpressed the sole non-RGS $G\beta$

subunit, GPB-1 (Jansen et al., 1999), and the broadly expressed $G\gamma$ subunit, GPC-2 (Yamada et

al., 2009), under the *tph-1* promoter along with GFP. $\beta\gamma$ over-expression in HSN did not cause

any significant differences in steady-state egg accumulation (Figure 5C). The number of eggs

stored *in-utero* in these animals (13.0 ± 1.1 eggs) was comparable to wild-type animals (15.7 ±

1.2 eggs) and less than *egl-10(md176)* mutant animals (44.5 ± 2.3 eggs). These results

suggest that Gα_o signals to inhibit HSN activity in a distinct manner from simple titration or

release of $\beta\gamma$ subunits.

Egg-laying behavior is dysregulated in cAMP and cGMP signaling mutants

Gα_o is in the G$\alpha_{i/o}$ class of G proteins that bind to and inhibit nucleotide cyclases, which

function to reduce cAMP and cGMP levels and their subsequent activation of protein kinases

(Kobayashi et al., 1990; Zhang and Pratt, 1996; Matsubara, 2002; Ghil et al., 2006). The

hyperactive egg-laying behavior phenotypes of animals with reduced Gα_o function could be

explained by a failure to properly terminate cAMP and/or cGMP signaling (Figure 5D, top). To

test this, we analyzed egg-laying behavior in strains bearing mutations that increase Gα_s and

cAMP effector signaling (Reynolds et al., 2005; Schade et al., 2005; Charlie et al., 2006a; Charlie

et al., 2006b). Animals carrying *gsa-1(ce81)* gain-of-function mutations in Gα_s predicted to

increase signaling accumulate fewer eggs compared to wild-type animals (Figure 5D, middle).

Because a reduction in steady-state egg accumulation could be caused by indirect effects of

these mutations on egg production or brood size, we also examined the developmental age of

embryos laid. As previously reported (Bany et al., 2003), loss of inhibitory Gα_o signaling causes

embryos to be laid precociously, before they reach the 8-cell stage (Figure 5D, bottom). Gα_s

gain-of-function mutant animals do not show a corresponding increase in early-stage embryos

that are laid, suggesting the reduction in egg accumulation observed in Gα_s gain-of-function

mutant animals is indirect. In contrast, gain-of-function *acy-1* Adenylate Cyclase mutations or

loss-of-function *pde-4* phosphodiesterase mutations, both predicted to result in increased

cAMP signaling, cause animals to accumulate fewer eggs and lay them at earlier stages

(Figure 5D). Similarly, *kin-2* mutations that disrupt an inhibitory regulatory subunit of Protein

Kinase A showed a modest but significant hyperactive egg-laying phenotype. Together, these

results indicate that $G\alpha_s$, cAMP, and Protein Kinase A signaling promote egg-laying behavior.

Because these mutant phenotypes were not nearly as strong as animals carrying $G\alpha_o$ mutants,

$G\alpha_o$ likely signals to inhibit egg laying via other effectors.

Figure 5. Gα$_o$ depresses HSN resting membrane potential and may inhibit egg-laying behavior in parallel to βγ, cGMP, and cAMP signaling pathways.

(**A**) Cartoon of the glued worm preparation used for patch clamp electrophysiology of HSN. (**B**) Scatter plots show resting membrane potential of wild-type control animals (grey circles), *goa-1(n1134)* loss-of-function mutants (red filled circles), *egl-10(md176)* null mutants (red open circles), and in transgenic animals expressing Pertussis Toxin in HSN (blue filled squares). Error bars indicate 95% confidence intervals for the mean. Asterisks indicate p<0.0001 (One-Way ANOVA with Bonferroni correction for multiple comparisons). N≥7 animals recorded per genotype. (**C**) Scatter plots show average number of eggs retained by wild-type animals (gray filled circles), *egl-10(md176)* null mutants (orange open circles), and in transgenic animals expressing either Gβ (GPB-2) and Gγ (GPC-1) subunits (green filled circles) or nothing (blue filled circles) in HSN from the *tph-1* gene promoter along with GFP. Error bars indicate means with 95% confidence intervals. Asterisk indicates p<0.0001; n.s. indicates p>0.05 (One-Way ANOVA with Bonferroni correction for multiple corrections). (**D**) Top, cartoon of how Gα$_o$ signaling might interact with cAMP and cGMP signaling pathways. Gene names for *C. elegans* orthologs tested here are indicated in parentheses. Middle, scatterplots show average number of eggs retained by wildtype (grey), *egl-10(md176)* (red open circles), *goa-1(n1134)* and *goa-1(sa734)* Gα$_o$ loss of function (red filled circles) mutants, and in animals with altered cAMP effector signaling (green): *gsa-1(ce81)* Gα$_s$ gain-of-function, *acy-1(ce2)* Adenylate Cyclase gain-of-function, *kin-2(ce179)* Protein Kinase A (PKA) inhibitory regulatory subunit loss-of-function, *pde-4(ce268)* Phosphodiesterase (PDE) loss-of-function, *pde-2(qj6)* Phosphodiesterase (PDE) null; altered cGMP effector signaling (pink): *egl-4(ad805)* and *egl-4(mg410)* Protein Kinase G (PKG) gain-of-function (pink filled circles), *egl-4(n478)* and *egl-4(n479)* loss-of-function mutants (pink open circles). Mean totals below ~10 eggs indicate hyperactive egg laying while totals above ~20 eggs indicate egg-laying behavior defects. Bottom, bar graphs indicate percent of embryos laid at early stages of development. Animals laying ≥50% embryos at early stages are considered hyperactive. Error bars indicate ± 95% confidence intervals for the mean proportion. N.D. indicates the stages of eggs laid was not determined because those mutants are egg-laying defective (Egl). Asterisk indicates highlighted significant differences (p≤0.0001; Fisher Exact Test with Bonferroni correction for multiple comparisons).

Extensive work has shown that the cGMP-dependent Protein Kinase G, EGL-4, regulates egg laying in *C. elegans* (Fujiwara et al., 2002; L'Etoile et al., 2002; Raizen et al., 2006; Hao et al., 2011). Mutations which increase EGL-4 activity cause hyperactive egg laying and release of early stage embryos, while elimination of EGL-4 signaling causes egg-laying defects and significant egg accumulation (Figure 5D, middle). To determine whether Gα$_o$ and Protein Kinase G regulate egg laying in a shared pathway, we performed a genetic epistasis experiment. *goa-1(sa734); egl-4(n479)* double null mutants accumulate very few eggs (Figure

5D, middle), resembling the *goa-1(sa734)* null mutant. However, the low brood size of the *goa-1(sa734)* mutant could prevent accurate measurement of these animal's egg-laying defects. To address this, we measured the stage of eggs laid as indicated above. Loss of the EGL-4 Protein Kinase G strongly and significantly suppressed the hyperactive egg-laying behavior of $G\alpha_o$ null mutants (Figure 5D, bottom). *goa-1(sa734)*; *egl-4(n479)* mutants laid 33% of their embryos at early stages compared to 88% for the *goa-1(sa734)* single mutant, an intermediate egg-laying phenotype. These results are consistent with cGMP and/or Protein Kinase G signaling acting downstream of $G\alpha_o$ in the proper regulation of egg-laying behavior.

$G\alpha_o$ signals in the vulval muscles and uv1 neuroendocrine cells to inhibit egg laying

In addition to HSN, GOA-1 is expressed in all neurons of the reproductive circuit, the egg-laying vulval muscles, and the uv1 neuroendocrine cells (Mendel et al., 1995; Segalat et al., 1995; Jose et al., 2007), raising questions as to how $G\alpha_o$ signaling functions in those other cells to regulate egg-laying behavior. Previous work has shown that GOA-1(Q205L) was sufficient to rescue the hyperactive egg-laying behavior of *goa-1(n1134)* mutants when expressed in the HSNs but had little effect when expressed in the VCs or vulval muscles (Tanis et al., 2008). Similarly, transgenic expression of Pertussis Toxin in the VCs or vulval muscles failed to modify egg-laying behavior strongly (Tanis et al., 2008). Previous work failing to identify a function for $G\alpha_o$ in the vulval muscles used a modified *Nde*-box element from the *ceh-24* promoter that drives expression more efficiently in the vm1 muscle cells compared to the vm2 muscles innervated by the HSN and VC neurons (data not shown). We find that expression of Pertussis Toxin in both vm1 and vm2 vulval muscles from a larger region of the *ceh-24* gene promoter (Harfe and Fire, 1998; Ravi et al., 2018a) also failed to cause any significant changes

in the steady-state egg accumulation (Figure 6A). Conversely, expression of the activated

GOA-1(Q205L) mutant in the vulval muscles from the same promoter did cause a modest but

significant egg-laying defect, with animals accumulating 24.1 ± 2.0 eggs compared to

mCherry-expressing control transgenic animals (13.2±0.7 eggs). This egg-laying defect was

significantly weaker than *egl-10(md176)* mutants (46.0 ± 3.3 eggs) or transgenic animals

expressing GOA-1(Q205L) just in the HSNs (36.8 ± 3.8 eggs). We do not believe this modest

egg-laying defect was caused by transgene expression outside of the vulval muscles, as

blocking synaptic transmission via transgenic expression of Tetanus Toxin from the same *ceh-*

24 promoter showed no such egg-laying defect (Figure 6B). Collectively, these results show

that $G\alpha_o$ does not play a significant role in suppressing vulval muscle excitability under state-

state conditions but that activated $G\alpha_o$ can signal in these cells to induce a mild but significant

inhibition of cell activity and egg-laying behavior.

The uv1 cells express the neurotransmitter tyramine along with NLP-7 and FLP-11

neuropeptides which inhibit egg laying (Alkema et al., 2005; Collins et al., 2016; Banerjee et al.,

2017). Based on the function of $G\alpha_o$ signaling in inhibiting neurotransmitter release in neurons,

we would expect that loss of $G\alpha_o$ function in uv1 would enhance their excitability, promoting

release of inhibitory tyramine and neuropeptides, causing a reduction of egg laying.

Surprisingly, previous work has shown that transgenic expression of Pertussis Toxin in uv1

cells increased the frequency of early-stage eggs that are laid, similar to the blocking of

neurotransmitter release by Tetanus Toxin (Jose et al., 2007). A caveat of these experiments is

that the *ocr-2* gene promoter used for transgene expression in uv1 also expresses in the utse

(uterine-seam) associated cells and head sensory neurons (Jose et al., 2007). To test whether

$G\alpha_o$ functions specifically in uv1 to regulate egg laying, we used the *tdc-1* gene promoter

(Alkema et al., 2005) along with the *ocr-2* 3' untranslated region (Jose et al., 2007) to drive

expression more specifically in uv1. Transgenic expression of Pertussis Toxin in uv1 caused a

mild but significant decrease in steady-state egg accumulation (10.9 ± 1.5 eggs) compared to

mCherry-expressing control animals (15.3 ± 1.2 eggs), indicating $G\alpha_o$ signaling facilitates uv1-

dependent inhibition of egg-laying behavior (Figure 6C). We also tested how increased $G\alpha_o$

signaling in uv1 affects egg laying. Transgenic expression of the activated GOA-1(Q205L)

mutant in uv1 cells caused no quantitative differences in egg accumulation (15.5 ± 2.1 eggs)

(Figure 6C). Together, these results show that $G\alpha_o$ has a limited role in regulating egg-laying

behavior in the vulval muscles or uv1 neuroendocrine cells, unlike the strong phenotypes

observed when we genetically manipulate $G\alpha_o$ function in HSN.

Figure 6. Altered $G\alpha_o$ signaling in the vulval muscles and uv1 neuroendocrine cells
causes only modest effects on egg laying.

(A) Scatter plots show average number of eggs retained by wild-type (grey), *goa-1(n1134)* mutants (green), *egl-10(md176)* null mutant animals (orange) along with transgenic animals expressing GOA-1^{Q205L} in the HSNs (red open circles) compared to transgenic animals expressing mCherry only (pink open circles), Pertussis Toxin (blue open circles), or GOA-1^{Q205L} (black open circles) in the vulval muscles from the *ceh-24* gene promoter. **(B)** Scatter plots show average number of eggs retained in transgenic animals expressing mCherry only (gray) or Tetanus Toxin along with mCherry (orange) in the vulval muscles (vm) using the *ceh-24* gene promoter. Error bars indicate means with 95% confidence intervals. **(C)** Scatter plots show average number of eggs retained by wild-type (grey), *goa-1(n1134)* mutant (green), *egl-10(md176)* null mutant (orange) animals, and transgenic animals expressing GOA-1^{Q205L} in the HSNs (red) compared to transgenic animals expressing mCherry only (pink), Pertussis Toxin (blue), or GOA-1^{Q205L} (black open circles) in the uv1 neuroendocrine cells from the *tdc-1* gene promoter. Four or five independent extrachromosomal arrays were generated for each transgene in **(A-C)** and ~10 animals bearing each extrachromosomal array were analyzed. Error bars indicate 95% confidence intervals for the mean. Asterisks indicate p<0.0001; n.s. indicates p>0.05 (One-Way ANOVA with Bonferroni's correction for multiple comparisons or Student's t test).

NLP-7 inhibition of HSN activity and egg-laying requires the EGL-47 receptor and Gα$_o$

Multiple neuropeptides and receptors have been identified to inhibit egg laying by signaling through Gα$_o$-coupled receptors expressed on HSN (Figure 1A). Recent work has identified NLP-7 neuropeptides, synthesized in the VC neurons and uv1 neuroendocrine cells, as potential ligands for EGL-47 receptor signaling through Gα$_o$ (Moresco and Koelle, 2004; Banerjee et al., 2017). Animals overexpressing the NLP-7 neuropeptide are egg-laying defective, accumulating 39.6 ± 3.4 eggs in the uterus (Figure 7A). To test how NLP-7 inhibits egg laying, we crossed NLP-7 over-expressing transgenes into *goa-1* mutant animals and evaluated their egg-laying behavior phenotypes. As shown in Figure 7A, the hypomorphic *goa-1(n1134)* loss-of-function mutant significantly suppressed these egg-laying defects, as previously shown (Banerjee et al., 2017). However, the suppression by *goa-1(n1134)* was incomplete; NLP-7 over-expressing, *goa-1(n1334)* double mutant animals retained more eggs than the *goa-1(n1134)* single mutant or even wild-type control animals (Figure 7A). To confirm whether Gα$_o$ was required for NLP-7 inhibition, we tested the *goa-1(sa734)* null mutant and

found it fully suppressed the egg-laying defects caused by NLP-7 over-expression. To confirm

this epistatic relationship, we measured the stage of embryos laid by these animals. *goa-*

1(sa734) null mutant animals over-expressing NLP-7 laid ~100% of their embryos at early

stages, a level not significantly different *goa-1(sa734)* single mutant animals (Figure 7B).

Together, these results show that NLP-7 neuropeptides cannot inhibit egg-laying behavior in

the absence of $G\alpha_o$ function.

Since the HSNs appear to be the principal sites of inhibitory $G\alpha_o$ signaling, we tested

how NLP-7 over-expression affects HSN Ca^{2+} activity. As expected, over-expression of NLP-7

strongly inhibited HSN Ca^{2+} activity (Figure 7C and 7D), consistent with the strong egg-laying

defects of these animals. To our surprise, loss of $G\alpha_o$ in these animals did now show high

frequency HSN Ca^{2+} transient activity, despite showing the strong hyperactive egg-laying

behavior of *goa-1(sa734)* single mutants (Figure 7C and 7D). This result shows that the

hyperactive egg-laying behavior of $G\alpha_o$ null mutants is unlinked to the increase in HSN Ca^{2+}

activity we observe. Previous work showed NLP-7 inhibition of HSN activity and egg laying

requires the EGL-47 receptor (Banerjee et al., 2017). To show directly whether NLP-7 inhibition

of HSN activity requires the EGL-47 receptor, we recorded HSN Ca^{2+} activity in *egl-47(ok677)*

null mutants. Unexpectedly, animals lacking EGL-47 had significantly fewer HSN Ca^{2+}

transients than wild-type animals whether or not NLP-7 was over-expressed (Figure 7D).

These results show that although NLP-7 signals to silence HSN Ca^{2+} activity, it does not

require EGL-47 or $G\alpha_o$ function to do so.

Figure 7. NLP-7 neuropeptides signal through EGL-47 and Gα₀ to inhibit egg-laying outside of the HSNs

(A) NLP-7 signals through Gα₀ to inhibit egg-laying behavior. Scatter plots show average number of eggs retained by wild-type (gray circles), *goa-1(sa734)* mutants (blue open circles), NLP-7 over-expressing (OE) transgenics in the wild-type (pink triangles or orange squares),

and in NLP-7 over-expressing transgenics in the *goa-1(n1134)* (pink open triangles) and *goa-*
1(sa734) null mutant background (orange open squares). Data in orange squares are from
animals that also carry the *vsIs183* transgene used for HSN Ca^{2+} imaging. Mean totals below
~10 eggs indicate hyperactive egg laying while totals above ~20 eggs indicate egg-laying
defective behavior defects. Error bars indicate 95% confidence intervals for the mean.
Asterisks indicate $p<0.0001$ (One-Way ANOVA with Bonferroni correction for multiple
comparisons). N≥30 animals for each strain. (**B**) Measure of early stage eggs laid by wild-type
(black), *goa-1(sa734)* null mutants (blue), and *goa-1(sa734)* null mutants over-expressing
NLP-7 neuropeptides (orange). Both *goa-1(sa734)* mutant strains also carry the *vsIs183*
transgene used for HSN Ca^{2+} imaging. (**C**) NLP-7 over-expression silences HSN Ca^{2+} activity.
Representative GCaMP5/mCherry ratio traces showing HSN Ca^{2+} activity in wild-type (black),
goa-1(sa734) null mutants, and NLP-7 overexpressing transgenic animals in the wild-type
(pink) and *goa-1(sa734)* null mutant backgrounds (orange). Arrowheads indicate egg-laying
events. (**D**) Scatter plots show HSN Ca^{2+} peaks per minute measurements in wild-type (black),
goa-1(sa734) null mutants, or *egl-47(ok677)* null mutant animals in the absence or presence of
NLP-7 over-expression (OE). Error bars indicate 95% confidence intervals for the mean for
N≥10 animals; n.s. indicates $p>0.05$ while asterisk indicates $p≤0.0001$ (one-way ANOVA with
Bonferroni correction for multiple comparisons). (**E**) NLP-7 inhibition of egg laying is not
exclusive to HSN silencing. Scatter plots show the average number of eggs retained by wild-
type animals, transgenic animals expressing Pertussis Toxin in HSN, and *egl-47(ok677)*
mutant animals in the absence or presence of NLP-7 over-expression (OE) or the absence of
HSNs. Error bars indicate 95% confidence intervals for the mean. Asterisks indicate $p<0.0001$
(One-Way ANOVA with Bonferroni correction for multiple comparisons); N≥30 animals for each
strain. (**F**) Measure of early stage eggs laid by wild-type or transgenic animals expressing
Tetanus Toxin the VC neurons (black outlined boxes) in animals expressing Pertussis Toxin in
the HSNs (green) or in *goa-1(sa734)* animals missing $G\alpha_o$ (blue). Error bars indicate ± 95%
confidence intervals for the mean proportion. Asterisk indicates highlighted significant
differences ($p≤0.0001$) while n.s. indicates $p>0.05$ (Fisher Exact Test with Bonferroni
correction for multiple comparisons).

To confirm whether NLP-7 inhibition of egg laying requires EGL-47 and $G\alpha_o$ function

outside of HSN, we compared egg-laying behavior in NLP-7 transgenic animals where $G\alpha_o$

and synaptic transmission were blocked in defined cells of the egg-laying circuit. As expected

from our Ca^{2+} imaging experiments, NLP-7 over-expressing animals accumulated a similar

number of eggs as *egl-1(n986dm)* animals lacking HSNs (35.7 ± 1.4 eggs vs. 33.4 ±0.9 eggs

Figure 7E). Transgenic expression of Pertussis Toxin in HSNs or loss of the EGL-47 receptor

weakly suppressed egg accumulation in NLP-7 over-expressing animals (Figure 7E). The

suppression of NLP-7 egg-laying defects by Pertussis Toxin expression was specific, as *egl-*

1(n986dm) animals developmentally lacking HSNs showed no comparable decrease in egg

retention and showed a mild but significant increase (Figure 7E), possibly through co-

expression of Pertussis Toxin from the transgene in the NSMs (Tanis et al., 2008). Thus, NLP-7

inhibits egg laying, in part, through an inhibitory G protein like $G\alpha_o$ in HSN and the EGL-47

receptor. Because whole-animal, but not HSN-specific, elimination of $G\alpha_o$ function completely

suppressed NLP-7-induced egg-laying defects (Figure 7A), these data show that NLP-7

signals to inhibit egg laying through $G\alpha_o$-coupled receptors on cells other than HSN. The

cholinergic VC neurons also make extensive synapses onto the vulval muscles, suggesting

they may act alongside the HSNs to promote vulval muscle contractility and egg laying (White

et al., 1986; Waggoner et al., 1998). To test whether enhanced VC neurotransmitter release was

driving the hyperactive egg-laying behavior of $G\alpha_o$ signaling mutants, we transgenically

expressed Tetanus Toxin in the VC neurons using a cell-specific promoter (Bany et al., 2003).

Loss of VC synaptic transmission had no effect on egg-laying behavior by *goa-1(sa734)* null

mutants or animals expressing Pertussis Toxin in HSN (Figure 7F). As previously reported

(Mendel et al., 1995; Segalat et al., 1995), we find that loss of $G\alpha_o$ also bypasses the regulation

by of egg laying by the HSNs, as *goa-1(sa734)*; *egl-1(n986dm)* double mutants

developmentally missing the HSNs still lay their eggs at early stages of development (Figure

7F). Together, these results show that NLP-7 neuropeptides and $G\alpha_o$ signal to inhibit egg-

laying behavior through cellular targets other than the VCs and HSNs.

Discussion

Using a combination of genetic, imaging, physiological, and behavioral approaches, we

found that the conserved G protein, $G\alpha_o$, signals to depress activity in the *C. elegans* egg-

laying circuit, preventing entry into the active behavior state. Neuropeptides, released in

response to aversive external sensory input and feedback of successful egg release, activate

inhibitory receptors and $G\alpha_o$ which signal to reduce cell electrical excitability and

neurotransmitter release. Without inhibitory $G\alpha_o$ signaling, the presynaptic HSN command

neurons remain electrically excited, rendering them resistant to inhibitory sensory input and

feedback from the stretch-dependent homeostat. As a result, animals lacking $G\alpha_o$ enter the

egg-laying active state twice as frequently as wild-type animals and in environments that would

normally be unfavorable for egg laying. Thus, inhibitory $G\alpha_o$ signaling allows for a two-state

pattern of circuit activity that can be properly gated by external and internal sensory input.

Our results inform our understanding of how G protein signaling modulates the stretch-

dependent homeostat that governs egg-laying behavior. Even in 'optimal' environmental

conditions including abundant food, egg laying in wild-type animals typically begins ~6 hours

after the L4-adult molt upon the accumulation of 5-8 eggs in the uterus (Ravi et al., 2018a). Loss

of inhibitory $G\alpha_o$ signaling causes precocious egg laying, with eggs being released soon after

being deposited into the uterus. Animals lacking $G\alpha_o$ function in HSN show tonic Ca^{2+} transient

activity even after chemical sterilization, conditions that normally inhibit HSN and vulval muscle

Ca^{2+} activity in wild-type animals. Genetic perturbations that increase inhibitory $G\alpha_o$ signaling

delay egg laying until ~18 hours after the molt, like animals lacking HSNs altogether. Egg

laying in animals with reduced or eliminated HSN function resumes when the accumulation of

eggs activates the circuit via the stretch-dependent homeostat (Collins et al., 2016; Ravi et al.,

2018a). Wild-type animals acutely exposed to aversive sensory conditions show a rapid

inhibition of HSN activity and egg laying (Sawin, 1996; Zhang et al., 2008; Fenk and de Bono,

2015), but they continue to make and accumulate eggs in the uterus. Egg laying resumes upon

removal of the inhibitory sensory stimulus or a return to food (Sawin, 1996; Dong et al., 2000).

Feedback of egg accumulation and release has long-term consequences on female

reproductive physiology. The cGMP-dependent Protein Kinase G ortholog, EGL-4, regulates

the expression of a novel secreted protein in the uterine epithelium whose levels correlate with

egg-laying rate (Hao et al., 2011). Feedback of egg release regulates *tph-1* gene expression

that increases serotonin biosynthesis in the HSNs, alters sensory responses to male

pheromone, and germline precursor cell proliferation (Aprison and Ruvinsky, 2019b, a).

Because the primary function of all adult *C. elegans* behaviors is survival and reproduction, we

hypothesize that most or all salient sensory signals ultimately converge to modulate egg-laying

circuit activity and animal fecundity.

Our work shows that $G\alpha_o$ also signals in cells other than HSN to inhibit egg-laying circuit

activity and behavior. GOA-1 is likely expressed in all electrically excitable cells including all

cells in the egg-laying circuit (Mendel et al., 1995; Segalat et al., 1995; Jose et al., 2007). Over-

expression of NLP-7 neuropeptides strongly inhibits HSN Ca^{2+} activity and egg-laying

behavior. While complete loss of $G\alpha_o$ suppresses egg-laying defects caused by NLP-7

overexpression, HSN Ca^{2+} activity remains largely inhibited. This suggests NLP-7 signals

through $G\alpha_o$ to inhibit egg laying in cells other than HSN. Consistent with this result, *goa-1* null

mutants remain strongly hyperactive for egg laying even when HSN or VC synaptic

transmission is blocked. Where else does $G\alpha_o$ function to inhibit egg laying? While $G\alpha_o$ is

expressed in the vulval muscles, transgenic expression of Pertussis Toxin or a GTP-locked

form of $G\alpha_o$ causes only modest egg-laying defects. Extensive work has shown that $G\alpha_o$

650 signals to inhibit release of acetylcholine from motor neurons during locomotion (Miller et al.,

651 1999; Nurrish et al., 1999). EM reconstruction that revealed the *C. elegans* synaptic wiring

652 diagram shows that the VA7 and VB6 cholinergic motor neurons that drive body wall muscle

653 contraction for locomotion also synapse onto the vm1 vulval muscles (White et al., 1986).

654 Recent work employing a new fluorescent reporter of acetylcholine shows rhythmic activity in

655 the vm1 muscles (Borden et al., 2020), and this vm1 activity is precisely where we observe

656 weak vulval muscle 'twitch' Ca^{2+} transients (Collins et al., 2016; Brewer et al., 2019). Mutations

657 that increase $G\alpha_s$ and cAMP signaling increase acetylcholine release from motor neurons and

658 promote locomotion (Reynolds et al., 2005; Schade et al., 2005; Charlie et al., 2006a; Charlie et al.,

659 2006b), possibly explaining why these mutations also increase egg laying. We predict that

660 cholinergic neurons like VA7 and VB6, or their command interneurons, express neuropeptide

661 and neurotransmitter receptors that signal through $G\alpha_o$ to depress acetylcholine release and

662 excitation of the vulval muscles. Identifying where such receptors are expressed and how they

663 signal will help explain how each cell in the circuit functions to drive discrete steps in egg

664 laying (Brewer et al., 2019; Fernandez et al., 2020).

665   Our work supports a model where $G\alpha_o$ inhibits synaptic transmission via modulation of

666 presynaptic ion channels. Genetic studies in *C. elegans* have identified the IRK inward

667 rectifying K^+ channels, NCA Na^+ leak channels, and CCA-1 T-type voltage-gated Ca^{2+}

668 channels as potential targets of $G\alpha_o$ signaling (Yeh et al., 2008; Emtage et al., 2012; Topalidou et

669 al., 2017a; Zang et al., 2017). Inward rectifying GIRK K^+ channels are activated by release of $\beta\gamma$

670 subunits (Hille, 1994). Previous work has shown the IRK-1 K^+ channel is expressed in HSN

671 and is required for inhibition of egg laying by the $G\alpha_o$-coupled EGL-6 neuropeptide receptor

672 (Emtage et al., 2012). The egg-laying phenotypes of *irk-1* mutant animals are not as strong as

673 *goa-1* mutants, and over-expression of $\beta\gamma$ subunits in HSN causes little or no effect on egg

laying. These results suggest $G\alpha_o$ signals to inhibit HSN neurotransmitter release via

additional mechanism(s). NALCN Na^+ leak channels are also expressed in HSN, and gain-of-

function mutations increase HSN Ca^{2+} activity and drive hyperactive egg-laying and locomotion

behaviors (Yeh et al., 2008). NALCN channels are genetically downstream of both $G\alpha_q$ and $G\alpha_o$

in the regulation of dopamine signaling and locomotion, suggesting that NALCN channels

could be direct targets for modulation (Lutas et al., 2016; Topalidou et al., 2017b). Physiological

experiments in mammalian neurons support this model. NALCN currents are activated by

neuropeptide signaling through $G\alpha_q$ (Lu et al., 2009) and inhibited by dopamine and GABA

signaling through $G\alpha_{i/o}$ (Philippart and Khaliq, 2018). However, *C. elegans* knockout mutants in

NALCN channel components do not show the strong behavior defects seen in $G\alpha_q$ and $G\alpha_o$

mutants, suggesting other channels act in parallel to NALCN to regulate HSN and circuit

excitability. One candidate are TMC channels that drive a background Na^+ leak conductance

that promotes HSN and vulval muscle cell electrical excitability (Yue et al., 2018). TMC

channels may also regulate cell electrical excitability as mechanosensors (Pan et al., 2018;

Tang et al., 2020). Cl^- channels and transporter proteins also regulate HSN activity and egg-

laying behavior. CLH-3 is a swelling and hyperpolarization-activated, inwardly rectifying

chloride channel that inhibits HSN activity (Branicky et al., 2014). Dynamic expression of Cl^-

transporter proteins in the HSNs as animals mature into egg-laying adults regulates the Cl^-

reversal potential (Tanis et al., 2009; Bellemer et al., 2011; Han et al., 2015). Loss of KCC-2 or

ABTS-1 transporters flips the reduced egg laying behavior of *egl-47(dm)* mutants into

hyperactive egg laying. Modulation of Cl^- transporter expression or activity by G protein

signaling could drive the shifts in cell excitability we observe between the inactive and active

egg-laying behavior states.

The modulation of antagonistic cation and anion channels likely determines the Ca^{2+}

dependent spiking probability of cells in the egg-laying circuit. HSN expresses L-type, P/Q-

type, and T-type voltage-gated Ca^{2+} channels (Mathews et al., 2003; Zang et al., 2017), and

each contributes to a normal serotonin response and egg-laying behavior (Schafer and Kenyon,

1995; Lee et al., 1997; Kwok et al., 2006). The modest but significant changes in HSN resting

membrane potential we observe in $G\alpha_o$ signaling mutants are expected to alter the probability

of eliciting Ca^{2+} spiking activity. Ca^{2+}-dependent action potentials have been found in *C.*

elegans neurons and muscles (Gao and Zhen, 2011; Liu et al., 2011; Liu et al., 2018) and is

regulated by both L-type and T-type Ca^{2+} channels. T-type channels like CCA-1 can contribute

to a 'window current' where the channel can pass current at depolarized potentials that are

insufficient to trigger channel inactivation (Williams et al., 1997; Crunelli et al., 2005; Zang et al.,

2017). Activation of these window currents might allow neurons like HSN to shift from

spontaneous tonic firing to high frequency Ca^{2+} bursting. Future work leveraging the powerful

molecular tools uniquely available in *C. elegans* and the egg-laying circuit along with direct

physiological measurements of membrane potential will allow mechanistic insight into how

neuromodulators like serotonin and neuropeptides signal through effectors to shape patterns

of circuit activity that underlie distinct behavior states.

Materials and Methods

Nematode Culture and Developmental Staging

Caenorhabditis elegans hermaphrodites were maintained at 20°C on Nematode Growth

Medium (NGM) agar plates with *E. coli* OP50 as a source of food as described (Brenner,

1974). For assays involving young adults, animals were age-matched based on the timing of

completion of the L4 larval molt. All assays involving adult animals were performed using age-

matched adult hermaphrodites 20-40 hours past the late L4 stage. Table 2 lists all strains used

in this study and their genotypes.

Plasmid and Strain Construction

Calcium reporter transgenes

HSN Ca^{2+}: HSN Ca^{2+} activity was visualized using LX2004 *vsIs183* [*nlp-3::GCaMP5::nlp-3*

3'UTR + nlp-3::mCherry::nlp-3 3'UTR + lin-15(+)] *lite-1(ce314) lin-15(n765ts) X* strain

expressing GCaMP5G and mCherry from the *nlp-3* promoter as previously described (Collins

et al., 2016). To visualize HSN Ca^{2+} activity in Gα_o signaling mutants, we crossed LX2004

vsIs183 lite-1(ce314) lin-15(n765ts) X males with MT2426 *goa-1(n1134) I*, DG1856 *goa-*

1(sa734) I, or MT8504 *egl-10(md176) V* hermaphrodites, and the fluorescent cross-progeny

were allowed to self, generating MIA210 *goa-1(n1143) I; vsIs183 X lite-1(ce314) lin-15*

(n765ts) X, MIA263 *goa-1(sa734) I; vsIs183 X lite-1(ce314) lin-15 (n765ts) X*, and MIA216 *egl-*

10(md176) V; vsIs183 lite-1(ce314) lin-15(n765ts) X strains, respectively. To visualize how

NLP-7 signals to inhibit HSN Ca^{2+} activity and egg laying, we crossed IZ1236 *ufIs118*

hermaphrodites (Banerjee et al., 2017) with LX2004 males. The fluorescent cross-progeny were

selfed, generating MIA293 *ufIs118; vsIs183 lite-1(ce314) lin-15(n765ts) X.* To test the

requirement for Gα_o signaling in NLP-7 inhibition, we crossed N2 males into MIA263

hermaphrodites, and the *goa-1(sa734)/+* I; *vsIs183 lite-1(ce314) lin-15 (n765ts) X*

heterozygous males obtained were then crossed with MIA293 hermaphrodites to generate

MIA294 *ufIs118; goa-1(sa734) I; vsIs183 lite-1(ce314) lin-15(n765ts) X.* In order to test

whether EGL-47 was a potential receptor for NLP-7, we crossed LX2004 males into RB850

egl-47(ok677) V hermaphrodites. The fluorescent cross-progeny were allowed to self and

homozygous *egl-47(ok677)* mutant animals were identified by duplex PCR genotyping with the

following oligos: CGTCACTTTTCCGTTGCTCTCTCATG,

GTAGCGGGAAAGATGGCAAGAAGTCG, and TCGGTGAAACTCATTGTGCTCATTGTGC,

creating MIA359 *egl-47(ok677) V; vsIs183 lite-1(ce314) lin-15(n765ts) X.* Wild-type N2 males

were then crossed to MIA359 hermaphrodites, and the male *egl-47(ok677)/+ V; vsIs183 lite-*

1(ce314) lin15(n765ts) X cross progeny were then crossed with MIA293 hermaphrodites. The

cross-progeny were selfed and genotyped for the *egl-47(ok677) V* deletion mutant, generating

MIA366 *ufIs118; egl-47(ok677) V; vsIs183 lite-1(ce314) lin-15(n765ts) X.*

During the course of crossing the *vsIs50* transgene expressing the catalytic subunit of

Pertussis Toxin in the HSNs from the *tph-1* promoter (Tanis et al., 2008), we noticed repulsion

between *vsIs183*, suggesting both were linked to the X chromosome. As such, LX850 *vsIs50*

lin-15(n765ts) X males were crossed with LX1832 *lite-1(ce314) lin-15(n765ts) X*

hermaphrodites, the non-Muv progeny were selfed, and homozygous *lite-1(ce314)* non-Muv

animals were kept, generating the strain MIA218 *vsIs50 lite-1(ce314) lin-15(n765ts) X.* MIA218

males were then crossed with LX2007 *vsIs186; lite-1(ce314) lin-15(n765ts) X*; the cross-

progeny were selfed, generating MIA227 *vsIs186; vsIs50 lite-1(ce314) lin-15(n765ts) X.* To test

whether Gα_o acts in HSN to mediate NLP-7 inhibition, we crossed wild-type N2 males into

LX2007 *vsIs186; lite-1(ce314) lin-15 (n765ts) X* hermaphrodites, and the heterozygous

vsIs186/+; lite-1(ce314) lin-15 (n765ts) X cross-progeny were mated with IZ1236 *ufIs118*

hermaphrodites. The fluorescent cross-progeny were selfed, creating MIA358 *ufIs118; vsIs186*

lite-1(ce314) lin-15(n765ts) X. Wild-type N2 males were then crossed into MIA227

hermaphrodites to generate heterozygous *vsIs186/+; vsIs50 lite-1(ce314) lin-15(n765ts) X*

males which were then crossed into MIA358 hermaphrodites. The fluorescent cross-progeny

were selfed, generating MIA360 *ufIs118; vsIs50; vsIs186; lite-1(ce314) lin-15(n765ts) X*.

Heterozygous *vsIs186/+; vsIs50 lite-1(ce314) lin-15(n765ts) X* males were also crossed into

MIA123 *egl-1(n986dm) V; lite-1(ce314) lin-15(n765ts)* X hermaphrodites, the fluorescent cross-

progeny were selfed, and Egl non-Muv animals were selected, generating MIA369 *egl-*

1(n986dm) V; vsIs50; vsIs186 lite-1(ce314) lin-15 (n765ts) X. The presence of the *egl-*

1(n986dm) allele was confirmed with snip-SNP PCR genotyping using oligonucleotides

CTCTGTTCCAGCTCAAATTTCC and GTAGTTGAGGATCTCGCTTCGGC followed by NciI

digestion.

In order to visualize HSN Ca^{2+} activity in transgenic animals expressing a constitutively active

mutant GOA-1^{Q205L} protein that increases Gα_o signaling in the HSN neurons, LX2004 *vsIs183*

lite-1(ce314) lin-15(n765ts) X males were crossed with LX849 *vsIs49; lin-15(n765ts) X*

hermaphrodites. We noted a repulsion between *vsIs183* and *vsIs49* transgenes integrated on

X. As such, we selected a strain MIA277 with trans-heterozygous *vsIs49* and *vsIs183*

transgenes (*lite-1(ce314) vsIs49 X / lite-1(ce314) vsIs183 X*) for Ca^{2+} imaging. The MIA277

strain was maintained by picking phenotypically egg-laying defective adult animals which show

GCaMP5G/mCherry expression.

Vulval Muscle Ca^{2+}: Vulval muscle Ca^{2+} activity was recorded in adult animals using LX1918

vsIs164 [unc-103e::GCaMP5::unc-54 3'UTR + unc-103e::mCherry::unc-54 3'UTR + lin-15(+)]

lite-1(ce314) lin-15(n765ts) X strain as described ([Collins et al., 2016](#)). To visualize vulval

muscle activity in Gα_o signaling mutants, LX1918 males were crossed with MT2426 *goa-*

1(n1134) I and DG1856 *goa-1(sa734) I* hermaphrodites, and the fluorescent cross-progeny

were selfed, generating MIA214 *goa-1(n1134) I; vsIs164 lite-1(ce314) lin-15(n765ts) X* and

MIA295 *goa-1(sa734) I; vsIs164 lite-1(ce314) lin-15(n765ts) X* strains, respectively. To

visualize vulval muscle activity in transgenic animals expressing the catalytic subunit of

Pertussis Toxin in the HSN neurons (Tanis et al., 2008), MIA218 *vsIs50 lite-1(ce314) lin-*

15(n765ts) X males were crossed with LX1919 *vsIs165 [unc-103e::GCaMP5::unc-54 3'UTR +*

unc-103e::mCherry::unc-54 3'UTR + lin-15(+)]; lite-1(ce314) lin-15(n765ts) X hermaphrodites,

and the cross progeny were selfed, generating MIA245 *vsIs50; vsIs165; lite-1(ce314) lin-*

15(n765ts) X. To visualize vulval muscle activity in transgenic animals expressing a

constitutively active mutant GOA-1^{Q205L} protein which increases Gα_o signaling in the HSN

neurons (Tanis et al., 2008), LX849 *vsIs49; lin-15(n765ts) X* males were crossed with LX1919

vsIs165; lite-1(ce314) lin-15(n765ts) X hermaphrodites and the fluorescent cross-progeny were

selfed, generating MIA291 *vsIs165; vsIs50 lite-1(ce314) lin-15(n765ts) X*.

Transgenes used to manipulate Gα_o signaling and synaptic transmission in specific cells of the

egg-laying circuit

HSN neurons: To produce a HSN (and NSM)-specific GPB-1 expressing construct, the *gpb-1*

cDNA fragment was amplified from pDEST-*gpb-1* (Yamada et al., 2009) using the following

oligonucleotides: 5'- GAGGCTAGCGTAGAAAAATGAGCGAACTTGACCAACTTCGA-3' and

5'-GCGGGTACCTCATTAATTCCAGATCTTGAGGAACGAG-3'. The ~1 kb DNA fragment was

digested with NheI/KpnI and ligated into pJT40A (Tanis et al., 2008) to generate pBR30. To

produce an HSN (and NSM)-specific GPC-2 expressing construct, the *gpc-2* cDNA fragment

was amplified from worm genomic DNA using the following forward and reverse

oligonucleotides: 5'-GAGGCTAGCGTAGAAAAATGGATAAATCTGACATGCAACGA-3' and

5'-GCGGGTACCTTAGAGCATGCTGCACTTGCT-3'. The ~250 bp DNA fragment was

digested with NheI/KpnI and ligated into pJT40A to generate pBR31. To co-overexpress the βγ

G protein subunits in the HSN neurons, we injected pBR30 (50 ng/μl), pBR31 (50 ng/μl), and

pJM60 [*ptph-1*::GFP] (80 ng/μl) ([Moresco and Koelle, 2004](#)) into the LX1832 *lite-1(ce314) lin-*

15(n765ts) animals along with pLI5EK (50 ng/μl), generating five independent

extrachromosomal transgenic lines which were used for behavioral assays. One representative

transgenic strain, MIA278 [*keyEx52*; *lite-1(ce314) lin-15(n765ts)*], was kept. To generate a

control strain for comparison in the egg-laying assays, we injected pJM66 [*ptph-1*::empty] (100

ng/μl) ([Tanis et al., 2008](#)) and pJM60 (80 ng/μl) into the LX1832 *lite-1(ce314) lin-15(n765ts)*

animals along with pLI5EK (50 ng/μl) generating five independent extrachromosomal control

transgenes which were used for behavioral assays. One representative transgenic strain,

MIA279 [*keyEx53*; *lite-1(ce314) lin-15(n765ts)*], was kept. To test how loss of HSNs affected

the hyperactive egg-laying behavior of *goa-1(sa734)* null mutants, we mated wild-type N2

males into MIA26 *egl-1(n986dm) V* hermaphrodites, and the heterozygous *egl-1(n986dm)*/+

male cross-progeny were then crossed into DG1856 *goa-1(sa734) I* hermaphrodites. Their

cross-progeny were selfed, and the presence of the *egl-1(n986dm)* and *goa-1(sa734)* were

confirmed by genotyping, creating MIA345 *goa-1(sa734) I*; *egl-1(n986dm) V*.

Vulval muscles: pJT40A *Ptph-1*::Pertussis Toxin ([Tanis et al., 2008](#)) was digested with

NheI/KpnI and ligated into pBR3 (*Pceh-24*::mCherry) to generate pBR20. pBR20 [*Pceh-*

24::Pertussis Toxin] (10 ng/μl) and pBR3 [*Pceh-24*::mCherry] (10 ng/μl) were injected into the

LX1832 *lite-1(ce314) lin-15(n765ts)* animals along with pLI5EK (50 ng/μl) to generate five

independent extrachromosomal transgenes which were used for behavioral assays. One

representative transgenic strain, MIA257 [*keyEx46*; *lite-1(ce314) lin-15(n765ts)*], was kept. To

produce vulval muscle-specific GOA-1(Q205L), the coding sequence of GOA-1(Q205L) was

recovered from pJM70C ([Tanis et al., 2008](#)) after digestion with NheI/SacI and ligated into

pKMC188 (*Punc-103e*::GFP; (Collins and Koelle, 2013)) generating pKMC268 (*Punc-*

103e::goa-1(Q205L)). However, because the *unc-103e* promoter also directs expression in

neurons that might indirectly regulate egg laying, GOA-1(Q205L) coding sequences were

removed from pKMC268 by digesting with NheI/NcoI and ligated into pBR3 to generate

pBR21. pBR21 [*Pceh-24*::GOA-1^{Q205L}] (10 ng/µl) and pBR3 [*Pceh-24*::mCherry] (10 ng/µl)

were injected into the LX1832 *lite-1(ce314) lin-15(n765ts)* animals along with pLI5EK (50 ng/µl)

to generate five independent extrachromosomal transgenes which were used for behavior

assays. One representative transgenic strain, MIA258 [*keyEx47*; *lite-1(ce314) lin-15(n765ts)*],

was kept. To generate control strains for comparison in egg-laying assays, pBR3 [*Pceh-*

24::mCherry] (20 ng/µl) was injected into the LX1832 *lite-1(ce314) lin-15(n765ts)* animals

along with pLI5EK (50 ng/µl) to generate five independent extrachromosomal transgenes

which were used for behavioral assays. One representative control transgenic strain, MIA256

[*keyEx45*; *lite-1(ce314) lin-15(n765ts)*], was kept. To produce a vulval muscle-specific Tetanus

Toxin transgene, Tetanus Toxin coding sequences were amplified from pAJ49 (*Pocr-*

2::Tetanus toxin) (Jose et al., 2007) using the following oligonucleotides: 5'-

GAGGCTAGCGTAGAAAAAATGCCGATCACCATCAACAACTTC-3' and 5'-

GCGCAGGCGGCCGCTCAAGCGGTACGGTTGTACAGGTT-3'. The DNA fragment was

digested with NheI/NotI and ligated into pBR6 to generate pBR27. To block any possible

neurotransmitter release from the vulval muscles, pBR27 (10 ng/µl) and pBR3 (10 ng/µl) was

injected into the LX1832 *lite-1(ce314) lin-15(n765ts)* animals along with pLI5EK (50 ng/µl) to

generate five independent extrachromosomal transgenes which were used for behavior

assays. One representative transgenic strain, MIA262 [*keyEx51*; *lite-1(ce314) lin-15(n765ts)*],

was kept.

uv1 neuroendocrine cells*:* To generate a uv1 cell-specific Pertussis toxin transgene, pBR20

(*Pceh-24*::Pertussis toxin) was digested with NheI/NcoI and the coding sequences of Pertussis

Toxin were then ligated into pAB5 (*Ptdc-1*::mCherry::*ocr-2* 3'UTR) to generate pBR25. pBR25

[*Ptdc-1*::Pertussis Toxin] (10 ng/µl) and pAB5 [*Ptdc-1*::mCherry] (5 ng/µl) were injected into

LX1832 *lite-1(ce314) lin-15(n765ts)* animals along with pLI5EK (50 ng/µl) to generate five

independent extrachromosomal transgenes which were used for behavioral assays. One

representative transgenic strain, MIA260 [*keyEx49*; *lite-1(ce314) lin-15(n765ts)*], was kept. To

generate a uv1 cell-specific GOA-1(Q205L) transgene, pKMC268 (*Punc-103e*::GOA-

1(Q205L)) was digested with NheI/NcoI and the coding sequences of GOA-1(Q205L) were

then ligated into pBR25 to generate pBR26. To increase $G\alpha_o$ signaling in uv1 cells, we injected

pBR26 [*Ptdc-1*::GOA-1^{Q205L}] (10 ng/µl) and pAB5 [*Ptdc-1*::mCherry] (5 ng/µl) into the LX1832

lite-1(ce314) lin-15(n765ts) animals along with pLI5EK (50 ng/µl) to generate five independent

extrachromosomal transgenes which were used for behavioral assays. One transgenic strain

MIA261 [*keyEx50*; *lite-1(ce314) lin-15(n765ts)*] was kept. To generate a control strain for

comparison in our egg-laying assays, pAB5 [*Ptdc-1*::mCherry] (15 ng/µl) was injected into the

LX1832 *lite-1(ce314) lin-15(n765ts)* animals along with pLI5EK (50 ng/µl) to generate five

independent extrachromosomal transgenes which were used for behavioral assays. One

representative transgenic strain, MIA259 [*keyEx48*; *lite-1(ce314) lin-15(n765ts)*], was kept.

VC neurons: A ~1.4 kB DNA fragment encoding Tetanus Toxin was excised from pAJ49 ([Jose

et al., 2007](#)) using AgeI/XhoI and ligated into a similarly digested pKMC145 *(Plin-11::GFP::unc-*

54 3' UTR) ([Collins et al., 2016](#)) to generate pKMC282 *(Plin-11::TeTx::unc-54 3' UTR)*.

pKMC282 (50 ng/µl) was injected along with pL15EK (50 ng/µl; ([Clark et al., 1994](#)) into LX1832

lite-1(ce314) lin-15(n765ts) X generating 4 independent transgenic lines of which one, MIA113

keyEx32 [Plin-11::TeTx::unc-54 3'UTR + lin-15(+)], was used for integration. *keyEx32* was

integrated with UV/TMP generating three independent integrants (*keyIs32*, *keyIs33*, and

keyIs46). Each transgenic line was backcrossed six times to LX1832 generating strains

MIA144, MIA145, and MIA146. The behavioral characterization of these VC-silenced animals

will be described elsewhere in more detail. To eliminate HSNs in these VC-silenced animals,

wild-type N2 males were crossed into MIA26 *egl-1(n986dm) V* hermaphrodites, and the

heterozygous *egl-1(n986dm)*/+ males were then crossed to MIA146 *keyIs46*; *lite-1(ce314) lin-*

15(n765ts) X hermaphrodites, and the cross-progeny were then selfed, generating the

homozygous strain MIA173 *keyIs46*; *egl-1(n986dm) V*; *lite-1(ce314) lin-15(n765ts) X*. To test

whether synaptic transmission from the VCs was required for the hyperactive egg-laying

behavior of $G\alpha_o$ signaling mutants, wild-type N2 males were crossed to MIA227 *vsIs186;*

vsIs50 lite-1(ce314) lin-15(n765ts) X hermaphrodites and the resulting heterozygous cross-

progeny were then crossed MIA144 hermaphrodites, allowed to self, generating MIA370

vsIs186; keyIs33; vsIs50 lite-1(ce314) lin-15(n765ts) X. To test how complete loss of GOA-1

affected egg laying in VC-silenced animals, we crossed wild-type N2 males into DG1856 *goa-*

1(sa734) I hermaphrodites, and the heterozygous *goa-1(sa734)*/+ cross-progeny were then

mated with MIA144 hermaphrodites. The cross-progeny were allowed to self, generating

MIA371 *goa-1(sa734) I; keyIs33; lite-1(ce314) lin-15(n765ts) X.* The presence of *goa-1(sa734)*

was confirmed by Sanger sequencing.

Fluorescence Imaging

Ratiometric Ca^{2+} Imaging: Ratiometric Ca^{2+} recordings were performed on freely behaving

animals mounted between a glass coverslip and chunk of NGM agar, as previously described

(Collins and Koelle, 2013; Li et al., 2013; Collins et al., 2016; Ravi et al., 2018b). Briefly, recordings

were collected on an inverted Leica TCS SP5 confocal microscope using the 8 kHz resonant

scanner at ~20 fps at 256x256 pixel resolution, 12-bit depth and ≥2X digital zoom using a 20x

Apochromat objective (0.7 NA) with the pinhole opened to ~20 µm. GCaMP5G and mCherry

fluorescence was excited using a 488 nm and 561 nm laser lines, respectively. Adult

recordings were performed 24 hours after the late L4 stage. After staging, animals were

allowed to adapt for ~30 min before imaging. During imaging, the stage and focus were

adjusted manually to keep the relevant cell/pre-synapse in view and in focus.

Ratiometric analysis (GCaMP5:mCherry) for all Ca^{2+} recordings was performed after

background subtraction using Volocity 6.3.1 as described (Ravi et al., 2018a). The egg-laying

active state was operationally defined as the period one minute prior to the first egg-laying

event and ending one minute after the last (in the case of a typical active phase where 3-4

eggs are laid in quick succession). However, in cases where two egg-laying events were apart

by >60 s, peaks were considered to be in separate active phases and any transients observed

between were considered to be from an inactive state. In animals where we observed no Ca^{2+}

peaks during the entire recording, the total duration of the recording was considered as one,

long, inter-transient interval. In animals where we observed a single Ca^{2+} transient, the

duration from the start of the recording to the time of the Ca^{2+} transient and the time from the

Ca^{2+} transient to the end of the recording were defined as inter-transient intervals. Animals

overexpressing NLP-7 become highly egg-laying defective, and the embryonic expression of

mCherry from the *nlp-3* promoter in accumulated eggs interfered with the standard image

segmentation protocol based on the mCherry channel. For these animals, unambiguous Ca^{2+}

transient-dependent changes in GCaMP5 fluorescence localized at the HSN cell body were

scored manually, and the frequency of such events were calculated as the number of observed

events per unit recording time, typically 10-20 minutes.

Behavior Assays and Microscopy

Animal sterilization: Animals were sterilized using Floxuridine (FUDR) as described (Mitchell

et al., 1979; Collins et al., 2016; Ravi et al., 2018a). Briefly, 100 µl of 10 mg/ml FUDR was applied

to OP50 seeded NGM plates. Late L4 animals were then staged onto the FUDR plates and the

treated adults were imaged ~24 hours later.

Egg laying assays: Unlaid eggs were quantitated as described (Chase et al., 2004). Staged

adults were obtained by picking late L4 animals and culturing them for 30-40 hr at 20°C. The

percentage of early-stage eggs laid were quantified as described (Koelle and Horvitz, 1996). 30

staged adults were placed on a thin lawn of OP50 bacteria on a nematode growth medium

(NGM) agar plate (Brenner, 1974) and allowed to lay eggs for 30 min. This was repeated with

new sets of staged animals until a total of at least 100 laid eggs were analyzed. Each egg was

examined under a Leica M165FC stereomicroscope and categorized into the following

categories: eggs which have 1 cell, 2 cell, 3-4 cell, 5-8 cell, and embryos with >8 cells. Eggs

with eight cells or fewer were classified as "early stage."

Long-term recording of egg-laying behavior: Egg-laying behavior was recorded at 4-5

frames per second from 24-hour adults after transfer to NGM plates seeded with a thin lawn of

OP50 bacterial food using a Leica M165FC stereomicroscope and camera (Grasshopper 3,

4.1 Megapixel, USB3 CMOS camera, Point Grey Research). N2 wild-type and hyperactive

egg-laying mutant strains (MT2426 and LX850) were recorded for 3 hours, and the egg-laying

defective strains MT8504 and LX849 were recorded for 8-10 hours. Average duration of active

and inactive states from these recordings were calculated as described (Waggoner et al., 1998;

Waggoner et al., 2000).

Electrophysiology

Electrophysiological recordings were carried out on an upright microscope (Olympus BX51WI)

coupled with an EPC-10 amplifier and Patchmaster software (HEKA), as previously described

([Yue et al., 2018](#); [Zou et al., 2018](#)). Briefly, day 2 adult worms were glued on the surface of

Sylgard-coated coverslips using the cyanoacrylate-based glue (Gluture Topical Tissue

Adhesive, Abbott Laboratories). A dorsolateral incision was made using a sharp glass pipette

to expose the cell bodies of HSN neurons for recording. The bath solution contained (in mM)

145 NaCl, 2.5 KCl, 5 $CaCl_2$, 1 $MgCl_2$, and 20 glucose (325–335 mOsm, pH adjusted to 7.3).

The pipette solution contained (in mM) 145 KCl, 5 $MgCl_2$, 5 EGTA, 0.25 $CaCl_2$, 10 HEPES, 10

glucose, 5 Na_2ATP and 0.5 NaGTP (315–325 mOsm, pH adjusted to 7.2). The resting

membrane potentials were tested with 0 pA holding under the Current Clamp model of whole-

cell patch. The high Cl^- content of the intracellular solution used may contribute to the relatively

depolarized membrane potentials we observed ([Liu et al., 2018](#)).

Experimental Design and Statistical Analysis

Sample sizes for behavioral assays followed previous studies ([Chase et al., 2004](#); [Collins and](#)

[Koelle, 2013](#); [Collins et al., 2016](#)). No explicit power analysis was performed before the study.

Statistical analysis was performed using Prism v.6-8 (GraphPad). Ca^{2+} transient peak

amplitudes and inter-transient intervals were pooled from multiple animals (typically ~10

animals per genotype/condition per experiment). No animals or data were excluded. Individual

p values are indicated in each Figure legend, and all tests were corrected for multiple

comparisons (Bonferroni for ANOVA and Fisher exact test; Dunn for Kruskal-Wallis).

Disclosures / Conflict of Interests

978 The authors declare no conflicts of interest.

Acknowledgements

This work was funded by grants from the NIH (R01-NS086932) and NSF (IOS-1844657) to

KMC. RJK 3rd was supported by a University of Miami Maytag Fellowship. We thank Yuichi Iino

for sharing plasmids. Strains used in this study have been provided to the *C. elegans* Genetics

Center, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

We thank Dr. Addys Bode Hernandez and Michael Scheetz for technical assistance. We thank

Drs. Qiang Liu, James Baker, Julia Dallman, Laura Bianchi, Brock Grill, Peter Larsson,

Stephen Roper, along with members of the Collins lab for helpful discussions and feedback on

the manuscript.

Video 1. GCaMP5:mCherry ratio recording of HSN Ca^{2+} activity in a control, wild-type adult

animal during an egg-laying active state. High Ca^{2+} is indicated in red while low calcium is in

blue. Head is at left, tail is at right.

Video 2. GCaMP5:mCherry ratio recording of HSN Ca^{2+} activity in a *goa-1(n1134)* mutant

adult animal during an egg-laying active state. High Ca^{2+} is indicated in red while low calcium

is in blue. Head is at left, tail is at right.

Video 3. GCaMP5:mCherry ratio recording of HSN Ca^{2+} activity in a *goa-1(sa734)* null mutant

adult animal during an egg-laying active state. High Ca^{2+} is indicated in red while low calcium

is in blue. Head is at right, tail is at left.

Video 4. GCaMP5:mCherry ratio recording of HSN Ca^{2+} activity in a transgenic adult animal

expressing Pertussis Toxin in the HSNs from the *tph-1* gene promoter during an egg-laying

active state. High Ca^{2+} is indicated in red while low calcium is in blue. Head is at top, tail is at

bottom.

Video 5. GCaMP5:mCherry ratio recording of vulval muscle Ca^{2+} activity in a control wild-type

adult animal during an egg-laying active state. High Ca^{2+} is indicated in red while low calcium

is in blue. Head is at right, tail is at top.

Video 6. GCaMP5:mCherry ratio recording of vulval muscle Ca^{2+} activity in a *goa-1(n1134)*

mutant adult animal during an egg-laying active state. High Ca^{2+} is indicated in red while low

calcium is in blue. Head is at right, tail is at left.

Video 7. GCaMP5:mCherry ratio recording of vulval muscle Ca^{2+} activity in a transgenic adult

animal expressing Pertussis Toxin in the HSNs from the *tph-1* gene promoter during an egg-

laying active state. High Ca^{2+} is indicated in red while low calcium is in blue. Head is at left, tail

is at right.

Table 2. *Strain names and genotypes for all animals used in this study (behavior assays and*

calcium imaging)

1015

Strain	Feature	Genotype	Reference
N2	Bristol wild-type strain	wild type	(Brenner, 1974)
DA521	EGL-4 (Protein Kinase G) gain-of-function mutant. Hyperactive egg laying, sluggish locomotion	*egl-4(ad805) IV*	(Raizen et al., 2006)
DG1856	GOA-1 (Gα_o) null mutant, hyperactive egg laying	*goa-1(sa734) I*	(Robatzek and Thomas, 2000)
IZ1236	NLP-7 overexpression	*ufls118*	(Banerjee et al., 2017)
KG421	GSA-1 (Gα_s) gain-of-function mutant. Hyperactive locomotion	*gsa-1(ce81) I*	(Schade et al., 2005)
KG518	ACY-1 (Adenylate Cyclase) gain-of-function mutant. Hyperactive locomotion	*acy-1(ce2) III*	(Schade et al., 2005)
KG532	KIN-2 (Protein Kinase A inhibitory regulatory subunit) loss-of-function mutant. Hypersensitive to stimuli	*kin-2(ce179) X*	(Schade et al., 2005)
KG744	PDE-4 (phosphodiesterase) loss-of-function mutant, Hyperactive locomotion	*pde-4(ce268) II*	(Charlie et al., 2006a)
LX849	HSN and NSM activated GOA-1(Q205L), egg laying defective	*vsls49; lin-15(n765ts) X*	(Tanis et al., 2008)
LX850	HSN and NSM Pertussis Toxin, hyperactive egg laying	*vsls50; lin-15(n765ts) X*	(Tanis et al., 2008)
LX1832	Strain for transgene production, blue-light insensitive, multi-vulva at 20°C in the absence of *lin-15(+)* rescue transgene.	*lite-1(ce314) lin-15(n765ts) X*	(Gurel et al., 2012)
LX1918	vulval muscle GCaMP5, mCherry	*vsls164 lite-1(ce314) lin-15(n765ts) X*	(Collins et al., 2016)
LX1919	Vulval muscle GCaMP5, mCherry	*vsls165; lite-1(ce314) lin-15(n765ts) X*	(Collins et al., 2016)
LX2004	HSN GCaMP5, mCherry	*vsls183 lite-1(ce314) lin-15(n765ts) X*	(Collins et al., 2016)
LX2007	HSN GCaMP5, mCherry	*vsls186 lite-1(ce314) lin-15(n765ts) X*	(Collins et al., 2016)
MIA26	no HSNs	*egl-1(n986dm) V*	(Garcia and Collins, 2019)
MIA36	EGL-4 (Protein Kinase G) gain-of-function mutant. Hyperactive egg laying, sluggish locomotion	*egl-4(mg410) IV*	(Hao et al., 2011)
MIA123	no HSNs with lin-15 marked transgenes	*egl-1(n986dm) V; lite-1(ce314) lin-15(n765ts) X*	this study

MIA144	VC Tetanus Toxin	*keyIs33; lite-1(ce314) lin-15(n765ts) X*	this study
MIA210	GOA-1 (Gα_o) reduced-function mutant; HSN GCaMP, mCherry	*goa-1(n1134) I; vsIs183 lite-1(ce314) lin-15(n765ts) X*	this study
MIA214	GOA-1 (Gα_o) reduced-function mutant; vulval muscle GCaMP5, mCherry	*goa-1(n1134) I; vsIs164 lite-1(ce314) lin-15(n765ts) X*	this study
MIA216	Increased Gα_o signaling; HSN GCaMP5, mCherry	*egl-10(md176) V; vsIs183 lite-1(ce314) lin-15(n765ts) X*	this study
MIA218	HSN and NSM Pertussis Toxin in blue-light insensitive, *lin-15* multi-vulva background	*vsIs50 lite-1(ce314) lin-15(n765ts) X*	this study
MIA227	HSN and NSM Pertussis Toxin; HSN GCaMP5, mCherry	*vsIs186; vsIs50 lite-1(ce314) lin-15(n765ts) X*	this study
MIA245	HSN and NSM Pertussis Toxin; vulval muscle GCaMP5, mCherry	*vsIs50 X vsIs165; lite-1(ce314) lin-15(n765ts) X*	this study
MIA256	Vulval muscle mCherry (*ceh-24* promoter)	*keyEx45; lite-1(ce314) lin-15(n765ts) X*	this study
MIA257	Vulval muscle mCherry + Pertussis Toxin (*ceh-24* promoter)	*keyEx46; lite-1(ce314) lin-15(n765ts) X*	this study
MIA258	Vulval muscle mCherry + Activated GOA-1(Q205L) (*ceh-24* promoter)	*keyEx47; lite-1(ce314) lin-15(n765ts) X*	this study
MIA259	uv1 cells mCherry (*tdc-1* promoter, *ocr-2* 3' UTR)	*keyEx48; lite-1(ce314) lin-15(n765ts) X*	this study
MIA260	uv1 cells mCherry + Pertussis Toxin (*tdc-1* promoter, *ocr-2* 3' UTR)	*keyEx49; lite-1(ce314) lin-15(n765ts) X*	this study
MIA261	uv1 cells mCherry + Activated GOA-1(Q205L) (*tdc-1* promoter, *ocr-2* 3' UTR)	*keyEx50; lite-1(ce314) lin-15(n765ts) X*	this study
MIA262	Vulval muscle mCherry + Tetanus toxin (*ceh-24* promoter)	*keyEx51; lite-1(ce314) lin-15(n765ts) X*	this study
MIA263	GOA-1 (Gα_o) null mutant; HSN GCaMP5, mCherry	*goa-1(sa734) I; vsIs183 lite-1(ce314) lin-15(n765ts) X*	this study
MIA277	Increased Gα_o signaling in HSN; HSN GCaMP5, mCherry	*vsIs49/+, +/vsIs183 lite-1(ce314) lin-15(n765ts) X (trans-heterozygote)*	this study
MIA278	HSN/NSM *gpb-1* and *gpc-2* overexpression + GFP (*tph-1* promoter)	*keyEx52; lite-1(ce314) lin-15(n765ts) X*	this study
MIA279	HSN/NSM empty + GFP (*tph-1* promoter)	*keyEx53; lite-1(ce314) lin-15(n765ts) X*	this study
MIA282	PDE-2 (phosphodiesterase) loss-of-function mutant	*pde-2(qj6)*	([Fujiwara et al., 2015](#))

MIA291	Increased Gα$_o$ signaling in HSN; vulval muscle GCaMP5, mCherry	*vsIs49; vsIs165 lite-1(ce314) lin-15(n765ts) X*	this study
MIA293	NLP-7 overexpression; HSN GCaMP5, mCherry	*ufIs118; vsIs183 lite-1(ce314) lin-15(n765ts) X*	(Banerjee et al., 2017) & this study
MIA294	NLP-7 overexpression; *goa-1(Gα$_o$)* null mutant; HSN GCaMP5, mCherry	*ufIs118; goa-1(sa734) I; vsIs183 lite-1(ce314) lin-15(n765ts) X*	(Banerjee et al., 2017) & this study
MIA295	GOA-1 (Gα$_o$) mutant, vulval muscles GCaMP5, mCherry	*goa-1(sa734) I; vsIs164 lite-1(ce314) lin-15(n765ts) X*	this study
MIA331	GOA-1 (Gα$_o$) null mutant; EGL-4 (Protein Kinase G) null mutant	*goa-1(sa734)I; egl-4(n479)IV*	this study
MIA345	GOA-1 (Gα$_o$) null mutant; no HSNs	*goa-1(sa734) I; egl-1(n986dm) V*	this study
MIA359	EGL-47 null mutant; HSN GCaMP5, mCherry	*egl-47(ok677) V; vsIs183 lite-1(ce314) lin-15(n765ts) X*	this study
MIA360	NLP-7 overexpression; HSN and NSM Pertussis Toxin; HSN GCaMP5, mCherry	*ufIs118; vsIs50; vsIs186 lite-1(ce314) lin-15(n765ts) X*	this study
MIA366	NLP-7 overexpression; EGL-47 null mutant; HSN GCaMP5, mCherry	*ufIs118; egl-47(ok677) V; vsIs183 lite-1(ce314) lin-15(n765ts) X*	this study
MIA369	no HSNs; HSN and NSM Pertussis Toxin; HSN GCaMP5, mCherry	*egl-1(n986dm) V; vsIs50; vsIs186 lite-1(ce314) lin-15 (n765ts) X*	this study
MIA370	VC Tetanus Toxin; HSN and NSM Pertussis Toxin; HSN GCaMP5, mCherry	*keyIs33; vsIs186; vsIs50 keyIs33; lite-1(ce314) lin-15(n765ts) X*	this study
MIA371	GOA-1 (Gα$_o$) null mutant; VC Tetanus Toxin	*goa-1(sa734) I; keyIs33; lite-1(ce314) lin-15(n765ts) X*	this study
MT1073	EGL-4 (Protein Kinase G) loss-of-function mutant. Egg-laying defective, roaming locomotion	*egl-4(n478) IV*	(L'Etoile et al., 2002)
MT1074	EGL-4 (Protein Kinase G) null mutant. Egg-laying defective, roaming locomotion	*egl-4(n479) IV*	(L'Etoile et al., 2002)
MT2426	GOA-1 (Gα$_o$) reduced-function mutant, hyperactive egg laying	*goa-1(n1134) I*	(Segalat et al., 1995)
MT8504	Increased Gα$_o$ signaling due to mutation in RGS protein, EGL-10	*egl-10(md176) V*	(Koelle and Horvitz, 1996)
RB850	EGL-47 null mutant	*egl-47(ok677) V*	(Moresco and Koelle, 2004; Consortium, 2012)

1016

References

References

Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR (2005) Tyramine Functions independently of
octopamine in the Caenorhabditis elegans nervous system. Neuron 46:247-260.
Aprison EZ, Ruvinsky I (2014) Balanced trade-offs between alternative strategies shape the
response of C. elegans reproduction to chronic heat stress. PLoS One 9:e105513.
Aprison EZ, Ruvinsky I (2019a) Coordinated Behavioral and Physiological Responses to a Social
Signal Are Regulated by a Shared Neuronal Circuit. Curr Biol 29:4108-4115 e4104.
Aprison EZ, Ruvinsky I (2019b) Dynamic Regulation of Adult-Specific Functions of the Nervous
System by Signaling from the Reproductive System. Curr Biol 29:4116-4123 e4113.
Banerjee N, Bhattacharya R, Gorczyca M, Collins KM, Francis MM (2017) Local neuropeptide
signaling modulates serotonergic transmission to shape the temporal organization of *C.*
elegans egg-laying behavior. PLoS Genet 13:e1006697.
Bany IA, Dong MQ, Koelle MR (2003) Genetic and cellular basis for acetylcholine inhibition of
Caenorhabditis elegans egg-laying behavior. J Neurosci 23:8060-8069.
Bastiani CA, Gharib S, Simon MI, Sternberg PW (2003) Caenorhabditis elegans Galphaq regulates
egg-laying behavior via a PLCbeta-independent and serotonin-dependent signaling
pathway and likely functions both in the nervous system and in muscle. Genetics 165:1805-
1822.
Bellemer A, Hirata T, Romero MF, Koelle MR (2011) Two types of chloride transporters are
required for GABA(A) receptor-mediated inhibition in *C. elegans*. EMBO J 30:1852-1863.
Borden PM et al. (2020) A fast genetically encoded fluorescent sensor for faithful in
vivo acetylcholine detection in mice, fish, worms and flies.
bioRxiv:2020.2002.2007.939504.
Branicky R, Miyazaki H, Strange K, Schafer WR (2014) The voltage-gated anion channels encoded
by clh-3 regulate egg laying in C. elegans by modulating motor neuron excitability. J
Neurosci 34:764-775.
Brenner S (1974) The genetics of *Caenorhabditis elegans*. Genetics 77:71-94.
Brewer JC, Collins KM, Koelle MR, Olsen A (2019) Serotonin and neuropeptides are both released
by the HSN command neuron to initiate C. elegans egg laying. PLoS Genet 15.
Brundage L, Avery L, Katz A, Kim UJ, Mendel JE, Sternberg PW, Simon MI (1996) Mutations in a C.
elegans Gqalpha gene disrupt movement, egg laying, and viability. Neuron 16:999-1009.
Carnell L, Illi J, Hong SW, McIntire SL (2005) The G-protein-coupled serotonin receptor SER-1
regulates egg laying and male mating behaviors in *Caenorhabditis elegans*. J Neurosci
25:10671-10681.
Charlie NK, Schade MA, Thomure AM, Miller KG (2006a) Presynaptic UNC-31 (CAPS) is required to
activate the G alpha(s) pathway of the Caenorhabditis elegans synaptic signaling network.
Genetics 172:943-961.
Charlie NK, Thomure AM, Schade MA, Miller KG (2006b) The Dunce cAMP phosphodiesterase
PDE-4 negatively regulates G alpha(s)-dependent and G alpha(s)-independent cAMP pools
in the Caenorhabditis elegans synaptic signaling network. Genetics 173:111-130.
Chase DL, Pepper JS, Koelle MR (2004) Mechanism of extrasynaptic dopamine signaling in
Caenorhabditis elegans. Nat Neurosci 7:1096-1103.
Clark SG, Lu X, Horvitz HR (1994) The Caenorhabditis elegans locus lin-15, a negative regulator of
a tyrosine kinase signaling pathway, encodes two different proteins. Genetics 137:987.
Collins KM, Koelle MR (2013) Postsynaptic ERG potassium channels limit muscle excitability to
allow distinct egg-laying behavior states in *Caenorhabditis elegans*. J Neurosci 33:761-775.

Collins KM, Bode A, Fernandez RW, Tanis JE, Brewer JC, Creamer MS, Koelle MR (2016) Activity of
1064   the *C. elegans* egg-laying behavior circuit is controlled by competing activation and
1065   feedback inhibition. Elife 5:e21126.
1066   Consortium CeDM (2012) large-scale screening for targeted knockouts in the Caenorhabditis
1067   elegans genome. G3 (Bethesda) 2:1415-1425.
1068   Crunelli V, Toth TI, Cope DW, Blethyn K, Hughes SW (2005) The 'window' T-type calcium current
1069   in brain dynamics of different behavioural states. J Physiol 562:121-129.
1070   Dempsey CM, Mackenzie SM, Gargus A, Blanco G, Sze JY (2005) Serotonin (5HT), fluoxetine,
1071   imipramine and dopamine target distinct 5HT receptor signaling to modulate
1072   *Caenorhabditis elegans* egg-laying behavior. Genetics 169:1425-1436.
1073   Dong MQ, Chase D, Patikoglou GA, Koelle MR (2000) Multiple RGS proteins alter neural G protein
1074   signaling to allow C. elegans to rapidly change behavior when fed. Genes Dev 14:2003-
1075   2014.
1076   Emtage L, Aziz-Zaman S, Padovan-Merhar O, Horvitz HR, Fang-Yen C, Ringstad N (2012) IRK-1
1077   potassium channels mediate peptidergic inhibition of *Caenorhabditis elegans* serotonin
1078   neurons via a G(o) signaling pathway. J Neurosci 32:16285-16295.
1079   Esposito G, Di Schiavi E, Bergamasco C, Bazzicalupo P (2007) Efficient and cell specific knock-
1080   down of gene function in targeted C. elegans neurons. Gene 395:170-176.
1081   Fenk LA, de Bono M (2015) Environmental CO2 inhibits Caenorhabditis elegans egg-laying by
1082   modulating olfactory neurons and evokes widespread changes in neural activity. Proc Natl
1083   Acad Sci U S A 112:E3525-3534.
1084   Fernandez RW, Wei K, Wang EY, Mikalauskaite D, Olson A, Pepper J, Christie N, Kim S, Koelle MR
1085   (2020) Cellular expression and functional roles of all 26 neurotransmitter GPCRs in the
1086   C. elegans egg-laying circuit. bioRxiv:2020.2004.2023.037242.
1087   Fujiwara M, Sengupta P, McIntire SL (2002) Regulation of body size and behavioral state of C.
1088   elegans by sensory perception and the EGL-4 cGMP-dependent protein kinase. Neuron
1089   36:1091-1102.
1090   Fujiwara M, Hino T, Miyamoto R, Inada H, Mori I, Koga M, Miyahara K, Ohshima Y, Ishihara T
1091   (2015) The Importance of cGMP Signaling in Sensory Cilia for Body Size Regulation in
1092   Caenorhabditis elegans. Genetics 201:1497-1510.
1093   Gao S, Zhen M (2011) Action potentials drive body wall muscle contractions in Caenorhabditis
1094   elegans. Proc Natl Acad Sci U S A 108:2557-2562.
1095   Garcia J, Collins KM (2019) The HSN egg-laying command neurons are required for normal
1096   defecation frequency in Caenorhabditis elegans (II). microPublication Biology.
1097   Ghil S, Choi JM, Kim SS, Lee YD, Liao Y, Birnbaumer L, Suh-Kim H (2006) Compartmentalization of
1098   protein kinase A signaling by the heterotrimeric G protein Go. Proc Natl Acad Sci U S A
1099   103:19158-19163.
1100   Goulding EH, Schenk AK, Juneja P, MacKay AW, Wade JM, Tecott LH (2008) A robust automated
1101   system elucidates mouse home cage behavioral structure. Proc Natl Acad Sci U S A
1102   105:20575-20582.
1103   Gurel G, Gustafson MA, Pepper JS, Horvitz HR, Koelle MR (2012) Receptors and other signaling
1104   proteins required for serotonin control of locomotion in *Caenorhabditis elegans*. Genetics
1105   192:1359-1371.
1106   Han B, Bellemer A, Koelle MR (2015) An evolutionarily conserved switch in response to GABA
1107   affects development and behavior of the locomotor circuit of Caenorhabditis elegans.
1108   Genetics 199:1159-1172.
1109   Hao Y, Xu N, Box AC, Schaefer L, Kannan K, Zhang Y, Florens L, Seidel C, Washburn MP, Wiegraebe
1110   W, Mak HY (2011) Nuclear cGMP-dependent kinase regulates gene expression via activity-

dependent recruitment of a conserved histone deacetylase complex. PLoS Genet
7:e1002065.
Harfe BD, Fire A (1998) Muscle and nerve-specific regulation of a novel NK-2 class homeodomain
factor in *Caenorhabditis elegans*. Development 125:421-429.
Herlitze S, Garcia DE, Mackie K, Hille B, Scheuer T, Catterall WA (1996) Modulation of Ca2+
channels by G-protein beta gamma subunits. Nature 380:258-262.
Hille B (1994) Modulation of ion-channel function by G-protein-coupled receptors. Trends
Neurosci 17:531-536.
Hobson RJ, Hapiak VM, Xiao H, Buehrer KL, Komuniecki PR, Komuniecki RW (2006) SER-7, a
Caenorhabditis elegans 5-HT7-like receptor, is essential for the 5-HT stimulation of
pharyngeal pumping and egg laying. Genetics 172:159-169.
Horvitz HR, Chalfie M, Trent C, Sulston JE, Evans PD (1982) Serotonin and octopamine in the
nematode Caenorhabditis elegans. Science 216:1012-1014.
Jansen G, Thijssen KL, Werner P, van der Horst M, Hazendonk E, Plasterk RH (1999) The complete
family of genes encoding G proteins of Caenorhabditis elegans. Nat Genet 21:414-419.
Jiang M, Spicher K, Boulay G, Wang Y, Birnbaumer L (2001) Most central nervous system D2
dopamine receptors are coupled to their effectors by Go. Proc Natl Acad Sci U S A 98:3577-
3582.
Jose AM, Koelle MR (2005) Domains, amino acid residues, and new isoforms of Caenorhabditis
elegans diacylglycerol kinase 1 (DGK-1) important for terminating diacylglycerol signaling
in vivo. J Biol Chem 280:2730-2736.
Jose AM, Bany IA, Chase DL, Koelle MR (2007) A specific subset of transient receptor potential
vanilloid-type channel subunits in Caenorhabditis elegans endocrine cells function as
mixed heteromers to promote neurotransmitter release. Genetics 175:93-105.
Kobayashi I, Shibasaki H, Takahashi K, Tohyama K, Kurachi Y, Ito H, Ui M, Katada T (1990)
Purification and characterization of five different alpha subunits of guanine-nucleotide-
binding proteins in bovine brain membranes. Their physiological properties concerning the
activities of adenylate cyclase and atrial muscarinic K+ channels. Eur J Biochem 191:499-
506.
Koelle MR (2016) Neurotransmitter signaling through heterotrimeric G proteins: insights from
studies in C. elegans. WormBook:1-78.
Koelle MR, Horvitz HR (1996) EGL-10 regulates G protein signaling in the C. elegans nervous
system and shares a conserved domain with many mammalian proteins. Cell 84:115-125.
Kwok TC, Ricker N, Fraser R, Chan AW, Burns A, Stanley EF, McCourt P, Cutler SR, Roy PJ (2006) A
small-molecule screen in C. elegans yields a new calcium channel antagonist. Nature
441:91-95.
L'Etoile ND, Coburn CM, Eastham J, Kistler A, Gallegos G, Bargmann CI (2002) The cyclic GMP-
dependent protein kinase EGL-4 regulates olfactory adaptation in C. elegans. Neuron
36:1079-1089.
Lackner MR, Nurrish SJ, Kaplan JM (1999) Facilitation of synaptic transmission by EGL-30 Gqalpha
and EGL-8 PLCbeta: DAG binding to UNC-13 is required to stimulate acetylcholine release.
Neuron 24:335-346.
Lee RY, Lobel L, Hengartner M, Horvitz HR, Avery L (1997) Mutations in the alpha1 subunit of an
L-type voltage-activated Ca2+ channel cause myotonia in Caenorhabditis elegans. EMBO J
16:6066-6076.
Li P, Collins KM, Koelle MR, Shen K (2013) LIN-12/Notch signaling instructs postsynaptic muscle
arm development by regulating UNC-40/DCC and MADD-2 in *Caenorhabditis elegans*. Elife
2:e00378.

Liu P, Ge Q, Chen B, Salkoff L, Kotlikoff MI, Wang ZW (2011) Genetic dissection of ion currents
1160   underlying all-or-none action potentials in C. elegans body-wall muscle cells. J Physiol
1161   589:101-117.
1162   Liu Q, Kidd PB, Dobosiewicz M, Bargmann CI (2018) C. elegans AWA Olfactory Neurons Fire
1163   Calcium-Mediated All-or-None Action Potentials. Cell 175:57-70 e17.
1164   Lou X, Korogod N, Brose N, Schneggenburger R (2008) Phorbol esters modulate spontaneous and
1165   Ca2+-evoked transmitter release via acting on both Munc13 and protein kinase C. J
1166   Neurosci 28:8257-8267.
1167   Lu B, Su Y, Das S, Wang H, Wang Y, Liu J, Ren D (2009) Peptide neurotransmitters activate a cation
1168   channel complex of NALCN and UNC-80. Nature 457:741-744.
1169   Lutas A, Lahmann C, Soumillon M, Yellen G (2016) The leak channel NALCN controls tonic firing
1170   and glycolytic sensitivity of substantia nigra pars reticulata neurons. Elife 5.
1171   Marder E (2012) Neuromodulation of neuronal circuits: back to the future. Neuron 76:1-11.
1172   Mathews EA, Garcia E, Santi CM, Mullen GP, Thacker C, Moerman DG, Snutch TP (2003) Critical
1173   residues of the Caenorhabditis elegans unc-2 voltage-gated calcium channel that affect
1174   behavioral and physiological properties. J Neurosci 23:6537-6545.
1175   Matsubara H (2002) [Angiotensin II type 2 (AT2) receptor signal and cardiovascular action].
1176   Nihon Yakurigaku Zasshi 119:95-102.
1177   McMullan R, Nurrish SJ (2011) The RHO-1 RhoGTPase modulates fertility and multiple behaviors
1178   in adult C. elegans. PLoS One 6:e17265.
1179   McMullan R, Hiley E, Morrison P, Nurrish SJ (2006) Rho is a presynaptic activator of
1180   neurotransmitter release at pre-existing synapses in C. elegans. Genes Dev 20:65-76.
1181   McMullen PD, Aprison EZ, Winter PB, Amaral LA, Morimoto RI, Ruvinsky I (2012) Macro-level
1182   modeling of the response of C. elegans reproduction to chronic heat stress. PLoS Comput
1183   Biol 8:e1002338.
1184   Mendel JE, Korswagen HC, Liu KS, Hajdu-Cronin YM, Simon MI, Plasterk RH, Sternberg PW (1995)
1185   Participation of the protein Go in multiple aspects of behavior in C. elegans. Science
1186   267:1652-1655.
1187   Miller KG, Emerson MD, Rand JB (1999) Goalpha and diacylglycerol kinase negatively regulate the
1188   Gqalpha pathway in C. elegans. Neuron 24:323-333.
1189   Mitchell DH, Stiles JW, Santelli J, Sanadi DR (1979) Synchronous growth and aging of
1190   Caenorhabditis elegans in the presence of fluorodeoxyuridine. J Gerontol 34:28-36.
1191   Moresco JJ, Koelle MR (2004) Activation of EGL-47, a Galpha(o)-coupled receptor, inhibits function
1192   of hermaphrodite-specific motor neurons to regulate *Caenorhabditis elegans* egg-laying
1193   behavior. J Neurosci 24:8522-8530.
1194   Mumby SM, Heukeroth RO, Gordon JI, Gilman AG (1990) G-protein alpha-subunit expression,
1195   myristoylation, and membrane association in COS cells. Proc Natl Acad Sci U S A 87:728-
1196   732.
1197   Nurrish S, Segalat L, Kaplan JM (1999) Serotonin inhibition of synaptic transmission: Galpha(0)
1198   decreases the abundance of UNC-13 at release sites. Neuron 24:231-242.
1199   Pan B, Akyuz N, Liu XP, Asai Y, Nist-Lund C, Kurima K, Derfler BH, Gyorgy B, Limapichat W,
1200   Walujkar S, Wimalasena LN, Sotomayor M, Corey DP, Holt JR (2018) TMC1 Forms the Pore
1201   of Mechanosensory Transduction Channels in Vertebrate Inner Ear Hair Cells. Neuron
1202   99:736-753 e736.
1203   Philippart F, Khaliq ZM (2018) Gi/o protein-coupled receptors in dopamine neurons inhibit the
1204   sodium leak channel NALCN. Elife 7.

Raizen DM, Cullison KM, Pack AI, Sundaram MV (2006) A novel gain-of-function mutant of the
cyclic GMP-dependent protein kinase egl-4 affects multiple physiological processes in
Caenorhabditis elegans. Genetics 173:177-187.
Ravi B, Garcia J, Collins KM (2018a) Homeostatic feedback modulates the development of two-
state patterned activity in a model serotonin motor circuit in Caenorhabditis elegans. J
Neurosci 38:6283-6298.
Ravi B, Nassar LM, Kopchock RJ, 3rd, Dhakal P, Scheetz M, Collins KM (2018b) Ratiometric calcium
imaging of individual neurons in behaving Caenorhabditis elegans. J Vis Exp.
Reuveny E, Slesinger PA, Inglese J, Morales JM, Iniguez-Lluhi JA, Lefkowitz RJ, Bourne HR, Jan YN,
Jan LY (1994) Activation of the cloned muscarinic potassium channel by G protein beta
gamma subunits. Nature 370:143-146.
Reynolds NK, Schade MA, Miller KG (2005) Convergent, RIC-8-dependent Galpha signaling
pathways in the Caenorhabditis elegans synaptic signaling network. Genetics 169:651-670.
Ringstad N, Horvitz HR (2008) FMRFamide neuropeptides and acetylcholine synergistically inhibit
egg-laying by *C. elegans*. Nat Neurosci 11:1168-1176.
Robatzek M, Thomas JH (2000) Calcium/calmodulin-dependent protein kinase II regulates
Caenorhabditis elegans locomotion in concert with a G(o)/G(q) signaling network. Genetics
156:1069-1082.
Sawin ER (1996) Genetic and Cellular Analysis of Modulated Behaviors in Caenorhabditis elegans.
In, p 316: Massachusetts Institute of Technology, Cambridge, USA.
Schade MA, Reynolds NK, Dollins CM, Miller KG (2005) Mutations that rescue the paralysis of
Caenorhabditis elegans ric-8 (synembryn) mutants activate the G alpha(s) pathway and
define a third major branch of the synaptic signaling network. Genetics 169:631-649.
Schafer WF (2006) Genetics of egg-laying in worms. Annu Rev Genet 40:487-509.
Schafer WR, Kenyon CJ (1995) A calcium-channel homologue required for adaptation to dopamine
and serotonin in Caenorhabditis elegans. Nature 375:73-78.
Segalat L, Elkes DA, Kaplan JM (1995) Modulation of serotonin-controlled behaviors by Go in
Caenorhabditis elegans. Science 267:1648-1651.
Shen K, Fetter RD, Bargmann CI (2004) Synaptic specificity is generated by the synaptic guidepost
protein SYG-2 and its receptor, SYG-1. Cell 116:869-881.
Shyn SI, Kerr R, Schafer WR (2003) Serotonin and Go modulate functional states of neurons and
muscles controlling C. elegans egg-laying behavior. Curr Biol 13:1910-1915.
Tang YQ, Lee SA, Rahman M, Vanapalli SA, Lu H, Schafer WR (2020) Ankyrin Is An Intracellular
Tether for TMC Mechanotransduction Channels. Neuron.
Tanis JE, Moresco JJ, Lindquist RA, Koelle MR (2008) Regulation of serotonin biosynthesis by the G
proteins Galphao and Galphaq controls serotonin signaling in *Caenorhabditis elegans*.
Genetics 178:157-169.
Tanis JE, Bellemer A, Moresco JJ, Forbush B, Koelle MR (2009) The potassium chloride
cotransporter KCC-2 coordinates development of inhibitory neurotransmission and
synapse structure in *Caenorhabditis elegans*. J Neurosci 29:9943-9954.
Topalidou I, Cooper K, Pereira L, Ailion M (2017a) Dopamine negatively modulates the NCA ion
channels in C. elegans. PLoS Genet 13:e1007032.
Topalidou I, Chen PA, Cooper K, Watanabe S, Jorgensen EM, Ailion M (2017b) The NCA-1 and NCA-
2 Ion Channels Function Downstream of Gq and Rho To Regulate Locomotion in
Caenorhabditis elegans. Genetics 206:265-282.
Trent C (1982) Genetic and Behavioral Studies of the Egg-Laying System of *Caenorhabditis elegans*.
Massachusetts Institute of Technology, Cambridge, USA.

Trent C, Tsuing N, Horvitz HR (1983) Egg-laying defective mutants of the nematode Caenorhabditis elegans. Genetics 104:619-647.

Waggoner LE, Zhou GT, Schafer RW, Schafer WR (1998) Control of alternative behavioral states by serotonin in *Caenorhabditis elegans*. Neuron 21:203-214.

Waggoner LE, Hardaker LA, Golik S, Schafer WR (2000) Effect of a neuropeptide gene on behavioral states in Caenorhabditis elegans egg-laying. Genetics 154:1181-1192.

White JG, Southgate E, Thomson JN, Brenner S (1986) The structure of the nervous system of the nematode *Caenorhabditis elegans*. Philos Trans R Soc Lond B Biol Sci 314:1-340.

Wierda KD, Toonen RF, de Wit H, Brussaard AB, Verhage M (2007) Interdependence of PKC-dependent and PKC-independent pathways for presynaptic plasticity. Neuron 54:275-290.

Williams SL, Lutz S, Charlie NK, Vettel C, Ailion M, Coco C, Tesmer JJ, Jorgensen EM, Wieland T, Miller KG (2007) Trio's Rho-specific GEF domain is the missing Galpha q effector in C. elegans. Genes Dev 21:2731-2746.

Williams SR, Toth TI, Turner JP, Hughes SW, Crunelli V (1997) The 'window' component of the low threshold Ca2+ current produces input signal amplification and bistability in cat and rat thalamocortical neurones. J Physiol 505 (Pt 3):689-705.

Yamada K, Hirotsu T, Matsuki M, Kunitomo H, Iino Y (2009) GPC-1, a G protein gamma-subunit, regulates olfactory adaptation in Caenorhabditis elegans. Genetics 181:1347-1357.

Yawo H (1999) Protein kinase C potentiates transmitter release from the chick ciliary presynaptic terminal by increasing the exocytotic fusion probability. J Physiol 515 (Pt 1):169-180.

Yeh E, Ng S, Zhang M, Bouhours M, Wang Y, Wang M, Hung W, Aoyagi K, Melnik-Martinez K, Li M, Liu F, Schafer WR, Zhen M (2008) A putative cation channel, NCA-1, and a novel protein, UNC-80, transmit neuronal activity in C. elegans. PLoS Biol 6:e55.

Yue X, Zhao J, Li X, Fan Y, Duan D, Zhang X, Zou W, Sheng Y, Zhang T, Yang Q, Luo J, Duan S, Xiao R, Kang L (2018) TMC Proteins Modulate Egg Laying and Membrane Excitability through a Background Leak Conductance in C. elegans. Neuron 97:571-585 e575.

Zang KE, Ho E, Ringstad N (2017) Inhibitory peptidergic modulation of *C. elegans* serotonin neurons is gated by T-type calcium channels. Elife 6:e22771.

Zhang J, Pratt RE (1996) The AT2 receptor selectively associates with Gialpha2 and Gialpha3 in the rat fetus. J Biol Chem 271:15026-15033.

Zhang M, Chung SH, Fang-Yen C, Craig C, Kerr RA, Suzuki H, Samuel AD, Mazur E, Schafer WR (2008) A self-regulating feed-forward circuit controlling C. elegans egg-laying behavior. Curr Biol 18:1445-1455.

Zou W, Fu J, Zhang H, Du K, Huang W, Yu J, Li S, Fan Y, Baylis HA, Gao S, Xiao R, Ji W, Kang L, Xu T (2018) Decoding the intensity of sensory input by two glutamate receptors in one C. elegans interneuron. Nat Commun 9:4311.