2 Seed-dispersal networks in tropical forest fragments: area effects, remnant species,

3 and interaction diversity

Carine Emer[1*], Pedro Jordano[2], Marco A. Pizo[3], Milton C. Ribeiro[1], Fernanda R. da Silva[4] and Mauro

Galetti[1,5]

[1]Departamento de Ecologia, Instituto de Biociências, Universidade Estadual Paulista (UNESP), CP

199, 13506-900, Rio Claro SP, Brazil

[2]Integrative Ecology Group, Estación Biológica de Doñana (EBD-CSIC), Consejo Superior de

Investigaciones Científicas, Avenida Americo Vespucio 26, E – 41092, Sevilla, Spain

[3]Departamento de Zoologia, Universidade Estadual Paulista (UNESP), CP 199, 13506-900, Rio Claro

SP, Brazil

[4] Instituto de Biologia, Universidade de Campinas (UNICAMP), R. Monteiro Lobato, 255 - Cidade

Universitária, 13083-862, Campinas SP, Brazil

[5] Department of Biology, University of Miami, Coral Gables, FL, USA.

*corresponding author: c.emer09@gmail.com

24 Submission and Acceptance Dates:

ABSTRACT

Seed dispersal interactions involve key ecological processes in tropical forests that help to maintain

ecosystem functioning. Yet this functionality may be threatened by increasing habitat loss, defaunation

and fragmentation. However, generalist species, and their interactions, can benefit from the habitat

change caused by human disturbance while more specialized interactions mostly disappear. Therefore

changes in the structure of the local, within fragment, networks can be expected. Here we investigated

how the structure of seed-dispersal networks changes along a gradient of increasing habitat

fragmentation. We analysed 16 bird seed-dispersal assemblages from forest fragments of a

biodiversity-rich ecosystem. We found significant species-, interaction- and network-area relationships,

yet the later was determined by the number of species remaining in each community. The number of

frugivorous bird and plant species, their interactions, and the number of links per species decreases as

area is lost in the fragmented landscape. In contrast, network nestedness has a negative relationship

with fragment area, suggesting an increasing generalization of the network structure in the gradient of

fragmentation. Network specialization was not significantly affected by area, indicating that some

network properties may be invariant to disturbance. Still, the local extinction of partner species,

paralleled by a loss of interactions and specialist-specialist bird-plant seed dispersal associations

suggests the functional homogenization of the system as area is lost. Our study provides empirical

evidence for network-area relationships driven by the presence/absence of remnant species and the

interactions they perform.

RESUMO

Interações de dispersão de sementes formam um processo ecológico chave em florestas tropicais onde

colaboram na manutenção do funcionamento do ecossistema. Porém, esta funcionalidade pode estar

ameaçada pelo aumento na perda e fragmentação do habitat. Enquanto espécies generalistas e suas

interações podem se beneficiar da mudança de habitat causada por distúrbios antrópicos, interações

envolvendo espécies mais especialistas são, na maioria, eliminadas. Desta forma, mudanças nas redes

locais, dentro de fragmentos florestais, são esperadas. Neste trabalho nós investigamos como a

estrutura de redes de dispersão de sementes mudam em um gradiente de fragmentação do habitat. Nós

analisamos 16 comunidades de dispersão de sementes espacialmente explícitas e distribuídas em

fragmentos florestais de um ecossistema rico em biodiversidade. Nós encontramos relações

significativas entre a área do fragmento e espécies, interações e estrutura das redes, sendo que o último

foi determinado pelo número de espécies remanescentes em cada comunidade. O número de espécies

de aves frugívoras e plantas e as interações entre eles, bem como o número de links por espécie

diminuíram significativamente conforme a área dos fragmentos é perdida. Por outro lado, o

aninhamento da rede mostrou uma relação negativa com a área do fragmento, sugerindo um aumento

da generalização da estrutura das redes com a fragmentação do habitat. No entanto, o grau de

especialização das redes não foi afetado pela área, indicando que algumas propriedades de rede podem

ser resistentes à perturbação. Sendo assim, a extinção local de espécies parceiras, conjuntamente com a

perda das interações e associações planta-dispersor mais especializadas, sugere uma homogeneização

do sistema conforme a área do fragmento é perdida. Nosso estudo fornece evidências empíricas para as

relações rede-área, sendo estas direcionadas pela presença e/ou ausência das espécies remanescentes

bem como das interações que estas realizam.

Keywords: frugivory; species interactions; human disturbance; defaunation; forest fragmentation;

network-area relationship; Atlantic Forest.

INTRODUCTION

THE ON-GOING, FAST-PACED DEFORESTATION HAS CREATED FRAGMENTED FORESTS POSING INCREASING

challenges to the conservation of species interactions and ecological processes (Janzen 1974, Bruna et

al. 2005, McConkey et al. 2016, Tylianakis & Morris 2017). Even though fragmentation is well

documented to affect species persistence and the connectivity among forest patches (da Silva &

Tabarelli 2000, Rybicki & Hanski 2013, Hagen et al. 2012), the remnant fragments are also proven to

hold an important fraction of biodiversity (Barlow et al. 2007, Sekercioğlu et al. 2007, Bongers et al.

2015). The pool of species and interactions persisting in forest fragments is formed by a non-random

subset of smaller-sized species while the whole spectrum of sizes found at the landscape level can only

occur in large, more pristine areas (Morante-Filho et al. 2015, Emer et al. 2018). Such local

assemblages configure local networks of interacting partner species whose structure may be affected by

the constraints imposed by fragment area (Aizen et al. 2012, Emer et al. 2013, Bomfim et al. 2018)

while resilience to habitat loss may occur in specific systems (e.g., Passmore et al. 2012). A recent

theoretical framework has proposed different mechanisms by which network-area relationships may

emerge, such as the constraints imposed by species-area relationships, the selection towards generalist

species and the dispersal limitation in smaller spatial scales (Galiana et al. 2018). Yet, whether network

structure is robust enough to be maintained across fragmented forests, or whether it changes as habitat

area is lost is still an open question to be tested with empirical data.

91   Ecological networks are the outcomes from associations among interacting species (Jordano et

al. 2003, Lewinshon & Prado 2006). Mutualistic associations between animal and plants are a

fundamental process to maintain forest dynamics (Herrera & Pellmyr 2002, Dennis et al. 2007). Seed

dispersal by birds, in which both interacting species benefit by using resource from one another, are

ubiquitous in tropical forests in which up to 90 percent of the plant species rely on birds to be

effectively dispersed (Howe & Smallwood 1982, Jordano 1995). Such associations between tropical

frugivorous birds and the plants they feed upon can form complex and diverse seed-dispersal networks,

characterized by a large number of species and interactions (Bascompte & Jordano 2007, Schleuning et

al. 2014, Dügger et al. 2018). Disruptions in the functionality of this system can be inferred by the

emergence of ecological patterns depicted from the correct interpretation of the structural properties of

the networks (Bascompte & Jordano, 2007, Memmott 2009, Howe 2016). For instance, changes in the

mean number of interactions (links) per species may indicate a reduction on the number of partners

available to interact with in smaller areas, in which generalist species are favoured (Morante-Filho et

al. 2016, Galiana et al. 2018). Likewise, increases in nestedness, a pervasive pattern in mutualistic

networks that suggests the presence of highly generalist species interacting with both generalist and

specialist ones, may suggest the loss of specialist interactions in disturbed communities (Aizen et al.

2012, Bomfim et al. 2018). Similarly, a reduction of network specialization, estimated by the H2' index

based on Shannon diversity, would suggest the community is losing diversity of interactions among

species, with the prevalence of more generalist species.

Thus, the winner species and their winner interactions able to thrive despite the disturbances

imposed by habitat loss and fragmentation may be the ones defining the configuration of the local

networks (Hobbs et al. 2006, Tabarelli et al. 2012, Vidal et al. 2014). For instance, diet generalist

species are expected to be more resilient to habitat change due to their ability to feed on, and to be

dispersed by, a wider range of interacting partners (Tabarelli et al. 2012, Morante-Filho et al. 2015).

Likewise, generalist small-bodied bird species and plant species typical of secondary forest may form

loose, generalists associations in fragmented landscapes (Emer et al. 2018), which may result in an

increased generalization in seed dispersal networks. Furthermore, habitat fragmentation generates

changes on biotic and abiotic conditions (Murcia 1995, Laurance et al. 2007) such as increases in

temperature and light conditions that may favour the competitive ability of some plant species over the

others (Laurance et al. 2010, Sfair et al. 2016). Associated to the negative effects of habitat reduction

lays defaunation, a pervasive phenomenon characterized by the local or functional extinction of a given

species, generally towards the larger body mass ones and often related to the reduction of habitat area

(Dirzo et al., 2014, Young et al. 2016). Thus, interactions performed by large-bodied species dispersing

large-seeded plants are expected to be absent in smaller areas (Pérez-Mendez et al. 2016, Emer et al.

2018) therefore affecting the local, within-fragment network structure.

126   Here we aim to understand how the reduction of habitat through landscape fragmentation

affects the structure of tropical seed-dispersal networks. We gathered data of avian-seed dispersal

interactions located in 16 fragments of the Atlantic Forest, a highly diverse and threaten tropical biome

(Ribeiro et al. 2009). Our first prediction is that the number of species present in each fragment will

follow a species-area relationship (MacArthur & Wilson 1967, Rybicki & Hanski 2013), i.e., the

number of species reduces according to the loss of habitat area. Secondly, because smaller area

fragments are disturbed habitats inhabited mostly by generalist species (Morante-Filho et al. 2016,

Emer et al. 2018) able to cope with the biotic and abiotic changes promoted by edge-effects associated

to other human-disturbance (Laurance et al. 2007), we expect a positive relationship between fragment

area, the total number of bird-seed dispersal interactions and the mean number of links per species.

Because the number of bird-seed dispersal interactions in forest fragments is selectively reduced

towards generalist species (Morante-Filho et al. 2016, Emer et al. 2018), we expect that the remaining

generalist interactions will lead to changes in network structure towards higher nestedness and lower

specialization, with interaction links associated to a few dominant species.

METHODS

DATASET

We compiled 16 studies of avian plant-frugivore interactions sampled at the community level along the

Atlantic Forest biome, a hotspot of biodiversity highly threaten by increasing human pressure (Ribeiro

et al. 2009, Joly et al. 2014; Fig. 1, Table S1). The study areas varied from 0.66 to 42,000 ha, along a

gradient of disturbance from semi-pristine Biological Reserves and State Parks to secondary forest

fragments and restored private lands. The matrix surrounding the fragments is variable, including sugar

cane fields, crop plantations, secondary forest, and urban areas. Overall the studies included here

originally aimed to record bird-eating-fruit interactions and not necessarily effective seed dispersal;

therefore we carefully checked every dataset and removed any interaction that would not characterize

plant dispersal events, such as seed predation. In the particular case of parrots we excluded them from

the analysis despite some rare evidences pointing to their role in effective seed dispersal of large nuts

(Tella *et al.* 2016); in most cases we could not establish in the original papers whether the frequency of

interactions involving parrots actually implied legitimate dispersal. We updated and standardized plant

and bird species names using the taxize package (Chamberlain & Szocs, 2013) in R (R Development

Core Team 2014).

NETWORK METRICS

For each community we built a correspondent seed-dispersal network as a bipartite graph represented

by a weighted matrix A_{ij} in which the rows represent the i plant species and columns represent the j bird

species. The elements a_{ij} indicate the frequency of interactions between the plant species i and

frugivore bird species j. Changes in network structure were assessed by estimating the following non-

correlated metrics ($r < 0.7$):

(i) the number of bird species (A);

(ii) the number of plant species (P);

(iii) the number of interactions (I);

(iv) the mean number of links per species (L);

(v) weighted nestedness (*wNODF*), which estimates the presence of generalist species interacting with

more specialist ones, based on the NODF index (Almeida-Neto et al. 2008);

(vi) H2' specialization (*H2'*), which estimates how strong species partition their interaction partners. It

measures the deviation of non-specialization in which values close to zero means low specialization of

the interacting species in the community while values close to one indicate high levels of specialization

(Blüthgen et al. 2006).

LANDSCAPE METRICS

Landscape configuration can play an important role in determining species persistence in fragmented

landscapes (Ribeiro et al. 2009) and as such may influence NAR's as well. Therefore, for each studied

community we calculated the fragment area (ha), the functional connectivity, and the average isolation.

We used the same remnant forests map generated by Ribeiro et al. (2009) as input map for the

calculation of the landscape metrics. The maps had 50 m spatial resolution for Albers Equal Area

projection and South America Datum (SAD 69). Because some birds can successfully move among

fragments within a given distance (Pizo et al. 2011; Cornellius et al. 2017; Vélez et al. 2015), and some

interactions functionally persist throughout a fragmented landscape (Emer et al. 2018), we estimated

the functional connectivity as the amount of forest (ha) that any species can have access if it is able to

cross a 500 m matrix (Ribeiro et al., 2009). Average isolation was calculated by randomly choosing

1000 points in a 5 km buffer around the sampling coordinates and extracting the Euclidean mean

distance (in meters) to the nearest forest fragment.

DEFAUNATION

Likewise, for each studied community we estimated the intensity of defaunation relatively to the

regional pool of species expected for the Atlantic Forest biome. To do so, we calculated the difference

between the sum of all bird body masses at the regional landscape scale (considering all frugivore bird

species present in the Atlantic-Frugivory database [Bello et al 2017] as a proxy for the regional pool of

species) and the summed body masses of the birds that were recorded interacting with fruits in each

fragment. Defaunation is based only on the presence/absence of species in a given fragment.

STATISTICAL ANALYSES

To understand whether and how the structure of avian seed-dispersal networks is affected by habitat

change we used Linear Regressions and General Least Square (gls) models in the nlme package

(Pinheiro et al., 2016). Individual models were fit for each network metric therefore considering the

number of plant species, the number of bird species, the number of interactions, the mean number of

interactions per species, nestedness and H2' specialization as response variables. Network metrics were

logit-transformed to solve the issue of being bounded from zero to one, when needed (Warton & Hui,

2011). Because defaunation is negatively correlated with fragment area ($r = -0.89$) and the three

landscape metrics are highly correlated among them (Table S2), we parsimoniously chose area as the

variable representing changes in habitat configuration. Therefore, fragment area was fitted as the

independent variable along with two covariates: (i) sampling intensity, calculated as the root square of

the number of interactions divided by the root square of the network size (sum of plant species and bird

species), to control for differences in sampling effort among studies (following Schleuning et al. 2012),

and (ii) forest type, classified according to the vegetation structure of each community (Ombrofilous

Forest, Semidecidous Forest, Araucaria Forest, and restored forest) to control for possible effects of

biotic and abiotic conditions intrinsic to each forest formation. Our spatially explicit models and model

selection follows the protocols suggested in Zuur et al. (2009). We did not detect spatial correlation

structure in our models after accounting for the geographic coordinates of each study site in the gls

models, according to the Akaike Information Criterion (AICc). Therefore, we fit the full model to

linear regressions with 'Maximum Likelihood estimation' while applying the 'dredge' function (Barton

2013) to select for the best-fixed structure according to AICc. 'Sampling intensity' was used as a fixed

parameter to test for the effects of area and forest type solely. Then, we used the Akaike weight of

evidence (wAICc) to obtain the relative importance of the different models (Burnham and Anderson

2002). Finally, we present the coefficients of each predictor and the variance explained by the optimal

linear model. Lastly, we used two different but complementary null models to test for changes in the

number of links per species, nestedness and H2' specialization independently of network size by fitting

standardized z-scores in the same model structure described above. Both null models constrained

network size and were created using the 'nullmodel' function. First, we used 'method = 1' (Patefield

algorithm) to randomize the distribution of links among species while maintaining the marginal totals

constant, i.e., species have the same total number of interactions in the observed and randomized

matrices, therefore maintaining their biological characteristic of being common or rare in the

community (Dorman et al. 2009). Second, we used 'method = 3' (vaznull algorithm) to randomize the

number of interactions among species without constraining marginal totals but maintaining

connectance constant, i.e., species have the same number of qualitative interactions as in the observed

matrix (Vázquez et al. 2007). Then, the mean and standard deviation of 100 iterations of each null

model were contrasted against the observed values of links per species, nestedness and specialization;

the resultant z-scores were used as the response variable in the statistical models. Network metrics and

null models were estimated in the bipartite package (Dormann et al., 2009). All analyses were run in R

(R Development Core Team, 2014).

RESULTS

Our dataset comprises 281 plant species (plus 54 plants identified and/or morphotyped at the genus or

family level) that interact with 175 frugivore bird species. We recorded a total of 7637 interactions and

2389 unique pairwise links, in which a bird species dispersed the seeds of a plant species in the studied

fragments of the Atlantic Forest.

The mean frequency of interactions per fragment (472.4 ± 363.4) was 16 times greater than the

mean number of bird (32.3 ± 17.9) and plant species (30.7 ± 42.6) recorded in each study site. The

most abundant interactions were those performed by generalist bird species dispersing secondary-forest

plant species, such as *Thraupis sayaca* dispersing seeds of *Cecropia pachystachya* (61 events of seed-

dispersal between three fragments; Fig S1). However, most of the events of seed-dispersal by frugivore

birds were rare given that 64% of the interactions occurred only once across the studied fragments (Fig.

S1).

We found significant species, interaction- and network-area relationships (Fig. 2, Table 1).

First, the number of bird and plant species decreased significantly with fragment area. Second, both the

number of bird seed-dispersal interactions and the mean number of links per species were negatively

affected by the reduction of fragment area. And finally, network nestedness showed a positive and

significant relationship with increasing fragmentation when controlling for sampling intensity while

specialization showed a tendency to decrease as area is lost, even though the variance explained by this

model was quite low. When network size was controlled for by the use of the null models, only

sampling intensity explained changes in the mean number of links per species, network nestedness and

H2' specialization (Table S3).

DISCUSSION

We found significant changes of network structure following the lost of habitat area in the fragments of

the Atlantic Forest. The direction of these changes is driven by the non-random loss of frugivorous bird

and plant species and the interactions they perform. Such changes derived from the lower number of

links per species and the higher network nestedness observed in smaller area fragments, in which bird-

seed dispersal interactions performed by small-sized generalist species are predominant (Emer et al.

2018). Yet, the reduced community size in smaller areas led to the structural changes observed on the

organization of seed-dispersal interactions, depicted by the influence of sampling intensity and the lack

of significant effects when tested against proper null models. Nonetheless, network specialisation

seems to be an invariant property of those communities given that it was not affected by habitat change

or pairwise species losses.

Bird seed-dispersal networks of the Atlantic Forest are losing species and interactions as the

landscape becomes more fragmented. Beyond the potential consequences for network stability (May

1972, Valdovinos 2019) detected by the increasing nestedness and reduced number of links performed

by the remnant species, those local communities are likely to be also losing important ecological

functions. Species- and interaction-richer communities mostly hold higher functional diversity while

the impoverished ones are likely to have gaps on ecosystem functionality (Isbell et al., 2011; Saavedra

et al. 2014, Montoya et al., 2015). The impoverishment of plant species in forest remnants from the

Northeast of the Atlantic Forest has led to a biotic homogenization across the fragmented landscape

(Lobo et al. 2011). Likewise, the impoverishment of interactions found in this study and the persisting

interactions performed by smaller-sized generalist species in the fragmented landscape (Emer et al.

2018) suggests that the Atlantic Forest is losing specific ecological functions, which may trigger

medium to long-term functional homogenization of interactions throughout the landscape (Olden et al.

2004, Laliberté & Tylianakis 2010). The changes on the number of links per species according to

habitat fragmentation may support our *a posteriori* hypothesis of functional homogenization potentially

driven by the persistence of a few, functionally-redundant sub-groups of interacting species that are

able to cope with the unstable conditions found in most forest fragments (Murcia 1995, Harper et al.

2005, Passmore et al. 2012).

The increasing nestedness of the bird-seed dispersal networks indicates that communities in

smaller area fragments predominantly include a core of generalist species that interact with a sub-group

of more specialist ones, while there is a lack, or at least a reduction of, specialist-specialist interactions.

In other words, the reduced species pool on smaller fragments are likely to promote their generalization

in terms of mutualistic partners while using whichever resource is available in these more disturbed

areas. Thus, it may indicate a community-level shift towards more generalist interactions as

fragmentation increases (Hobbs et al. 2006; Kiers et al 2010). Likewise, changes in the behaviour of

the species that persist in smaller fragments, or fluctuations in population dynamics, could also explain

the increasing network generalization (Awade et al. 2017; Cornelius et al. 2017). Furthermore, the

reduced community size in smaller areas apparently leads to some structural changes in the

organization of seed-dispersal interactions, as suggested by the influence of sampling intensity on

network nestedness when tested against proper null models.

In turn, network specialization was not significantly affected by habitat reduction. This suggests

that, despite negative effects of fragmentation on network size, the way remnant species interact in

terms of partners diversity is similar between small assemblages of bird and plant species in smaller

areas, and large assemblages in larger areas. Considering that the minimum value of the H2' index

indicates maximum niche overlap among interacting species (while the opposite indicates maximum

niche divergence; Schleuning et al. 2012), our analysis reveals that niche partitioning of fruit food

resources provided by the plants to the birds, or the seed dispersal service promoted by the birds to the

plants, are not affected by the loss of area in smaller sized fragments. This suggests that network

specialization in terms of niche partitioning is driven by different mechanisms other than habitat area

(Schleuning et al. 20144, Mello et al. 2015). The presence of naturalized alien species in the system,

mainly plants such as *Melia azedarach* and *Psidium guajava,* may numerically compensate the absence

of some native ones, which are more sensitive to habitat change. Yet, their functionality as food

resources for the bird community appears impoverished.

Tropical forest fragments are hyperdynamic systems facing population and community changes

in response to primary effects of habitat loss followed by subsequent edge effects and stochastic events

(Laurance et al. 2002, Laurance et al. 2010). Furthermore, the Atlantic Forest landscape holds very

high levels of beta-diversity and turnover of birds, plants, and their interactions (Morante-Filho et al.

2016, Farah et al. 2017, Emer et al. 2018), which could lead, at least partially, to the observed network-

area relationships. Such community heterogeneities can be further influenced by other anthropogenic

impacts such as hunting and logging, whose frequency and intensity of occurrence are context-

dependent and vary among fragments (Joly et al. 2014, Galetti et al. 2016). Therefore, long-term effects

of habitat fragmentation on network structure may go through different trajectories (Thompson 2005)

but the impoverishment of the system is unlikely to change.

Finally, our findings have important implications for advancing the understanding of network-

area relationships (NARs, *sensu* Galiana et al. 2018). While species-area relationships are one of the

most pervasive rules in ecology (MacArthur & Wilson 1967, Lawton 1999), empirical data

demonstrating whether or not interactions- and network-area follow the same rules are rare (but see

Aizen et al. 2012, Passmore et al. 2012, Emer et al. 2013, Bomfim et al. 2018 for examples on

mutualistic networks). Our results showed that, as expected, bird and plant species as well as their

interactions, were strongly affected by fragment area. Likewise, the number of links per species and

nestedness also changed with area but not network specialization. Yet, when the number of species in

each community was considered using the null models, none of the network properties changed with

area, only with sampling intensity. These findings suggest that area *per se* does affect network

structure, but primarily mediates the presence/absence of remnant species in the fragments, as well as

the interactions they perform. Furthermore, the lack of significant changes in network specialization

independently of local species richness, contrasting with the richness-associated changes of the

previous metrics, may suggest that the stability of complex system depends on the specific parameter

and the system analysed (May 1972, Thébault & Fontaine 2010, Valdovinos et al. 2019). Specialization

of the seed-dispersal interactions could be seen as an invariant property of the networks, at some extent

robust to habitat loss. On the other hand, network size and the consequent number of interactions and

nestedness seem to be more prone to collapse as disturbance increases with landscape fragmentation

and reduced habitat area. Therefore, the loss of species and their interactions following habitat loss may

be a good predictor of the directions of changes in network structure, providing guidance for an overall

theory of NARs.

ACKNOWLEDGMENTS

We thank Carolina Bello to provide the map used in Figure 1 and Carolina Carvalho for statistical

advice. This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo

(BIOTA – FAPESP 2014/01986-0). CE was supported by a postdoctoral fellowship (FAPESP

DATA AVAILABILITY STATEMENT

Upon acceptance, the dataset used in this manuscript will be made available on the main author's

GitHub repository (https://github.com/carineemer/network_area).

LITERATURE CITED

AIZEN, M. A., SABATINO, M. & TYLIANAKIS, J. M. (2012). Specialization and rarity predict non-random

loss of interactions from mutualist networks. *Science* 335: 1486-1489.

ALMEIDA-NETO, M., GUIMARÃES, P., GUIMARÃES, P. R., LOYOLA, R. D. & ULRICH, W. (2008). A

consistent metric for nestedness analysis in ecological systems: reconciling concept and

measurement. *Oikos* 117, 1227-1239.

AWADE, M., CANDIA-GALLARDO, C., CORNELIUS, C. & METZGER, J. P. (2017). High emigration

propensity and low mortality on transfer drives female-biased dispersal of *Pyriglena leucoptera* in

fragmented landscapes. *PLoS ONE* 12: e0170493.

BARLOW, J., GARDNER, T. A., ARAUJO, I. S., AVILA-PIRES, T. C., BONALDO, A. B., COSTA, J. E.,

ESPOSITO, M. C., FERREIRA, L. V., HAWES, J., HERNANDEZ, M. I. M., HOOGMOED, M. S., LEITE, R.

N., LO-MAN-HUNG, N. F., MALCOLM, J. R., MARTINS, M. B., MESTRE, L. A. M., MIRANDA-SANTOS,

R., NUNES-GUTJAHR, A. L., OVERAL, W. L., PARRY, L., PETERS, S. L., RIBEIRO-JUNIOR, M. A., DA

SILVA, M. N. F., & PERES, C. A. (2007). Quantifying the biodiversity value of tropical primary,

secondary, and plantation forests. *Proc. Natl. Acad. Sci. USA* 104:18555-60.

BARTON, K. (2013). MuMIn: multi-model inference. R package version 1.42.1. Available at:

https://cran.r-project.org/web/packages/MuMIn/index.html (accessed 06 Feb 2019).

BASCOMPTE, J. & JORDANO, P. (2007). Plant-animal mutualistic networks: The architecture of

biodiversity. *Ann. R. Ecol. Evol. Syst.* 38, 567-593.

BELLO, C., GALETTI, M., MONTAN, D., PIZO, M. A., MARIGUELA, T. C., CULOT, L., BUFALO, F.,

LABECCA, F., PEDROSA, F., CONSTANTINI, R., EMER, C., SILVA, W. R., SILVA, F. R. DA, OVASKAINEN,

O. & JORDANO, P. (2017). Atlantic-Frugivory: a plant-frugivore interaction dataset for the Atlantic

Forest. *Ecology* 98: 1729.

BLÜTHGEN, N., F. MENZEL & BLÜTHGEN, N. (2006). Measuring specialization in species interaction

networks. *BMC Ecology* 6:9.

BOMFIM, J. DE A., GUIMARÃES JR. , P. R., PERES, C. A., CARVALHO, G. & CAZETTA, E. (2018). Local

extinctions of obligate frugivores and patch size reduction disrupt the structure of seed dispersal

networks. *Ecography* 41: 1899–1909.

BONGERS, F., CHAZDON, R., POORTER, L. & PENA-CLAROS, M. (2015). The potential of secondary

forests. *Science*, 348, 642-643.

BRUNA, E. M., VASCONCELOS, H. L. & HEREDIA, S. (2005). The effect of habitat fragmentation on

communities of mutualists: a test with Amazonian ants and their host plants. *Biolog. Conserv.* 124:

209-216.

BURNHAM, K. P. & ANDERSON, D. R. (2002). Model selection and multimodel inference: A practical

information-theoretic approach. 2ed. Springer-Verlag, New York, USA.

CHAMBERLAIN, S. A. & SZOCS, E. (2013). taxize - taxonomic search and retrieval in R. *F1000Research*,

2:191. Available at: http://f1000research.com/articles/2-191/v2. (accessed 16 Aug 2019).

CORNELIUS, C., AWADE, M., CÂNDIA-GALLARDO, C., SIEVING, K. E. & METZGER, J. P. (2017). Habitat

fragmentation drives inter-population variation in dispersal behavior in a Neotropical rainforest bird.

Perspect. Ecol. Evol., 15, 3–9.

DA SILVA, J. M. C. & TABARELLI, M. (2000). Tree species impoverishment and the future flora of the

Atlantic forest of northeast Brazil. *Nature* 404, 72-74.

DENNIS, A. J., GREEN, R. J., SCHUPP, E. W. & WESTCOTT, D. A. (2007) Seed dispersal: Theory and its

application in a changing world. CABI, Wallingford, UK.

DIRZO, R., YOUNG, H. S., GALETTI, M., CEBALLOS, G., ISAAC, N. J. B. & COLLEN, B. (2014).

Defaunation in the Anthropocene. *Science*, 345, 401-406.

DORMANN, C. F., JOCHEN, F., BLÜTHGEN, N. & GRUBER, B. (2009). Indices, graphs and null models:

analysing bipartite ecological networks. *Open Ecol. J.* 2: 7-24.

DÜGGER, P., BLENDINGER, P. G., BOHNING-GAESE, K., CHAMA, L., CORREIA, M., DEHLING, D. M.,

EMER, C., FARWIG, N., FRICKE, E. D., GALETTI, M., GARCÍA, D., GRASS, I., HELENO, R., JACOMASSA,

F. A. F., MORAES, S., MORAN, C., MUÑOZ, M. C., NEUSCHULZ, E. L., NOWAK, L., PIRATELLI, A.,

PIZO, M. A., QUITIÁN, M., ROGERS, H. S., RUGGERA, R. A., SAAVEDRA, F., SÁNCHEZ, M. S.,

SÁNCHEZ, R., SANTILLÁNS, V., SCHABO, D. G., DA SILVA, F. R., TIMÓTEO, S., TRAVESET, A.,

VOLLSTADT, M. G. R. & SCHLEUNING, M. (2018). Seed-dispersal networks are more specialized in

the Neotropics than in the Afrotropics. *Global Ecol. Biogeog.* DOI 10.1111/geb.12833.

EMER, C., VENTICINQUE, E. M. & FONSECA, C. R. (2013). Effects of dam-induced landscape

fragmentation on Amazonian ant-plant mutualistic networks. *Conserv. Biol.* 27: 763-773.

EMER, C. GALETTI, M., PIZO, M. A., GUIMARÃES JR., P. R., MORAES, S., PIRATELLI, A. & JORDANO. P.

(2018). Seed-dispersal interactions in fragmented landscapes – a metanetwork approach. *Ecol. Lett.*

21: 484-493.

FARAH, F. T., DE MUYLAERT, R. L., RIBEIRO, M. C., RIBEIRO, J. W., MANGUEIRA, J. R. DE S. A., Souza,

V. C. & Rodrigues, R. R. (2017). Integrating plant richness in forest patches can rescue overall

biodiversity in human-modified landscapes. *For. Ecol. Manag.* 397: 78–88.

GALIANA, N., LURGI, M., CLARAMUNT-LÓPEZ, B., FORTIN, M-J., LEROUX, S., CAZELLES, K., GRAVEL,

D. & MONTOYA, J. M. (2018). The spatial scaling of species interaction networks. *Nat. Ecol. Evol.* 2:

782-790.

GALETTI, M., BROCARDO, C. R., BEGOTI, R. A., HORTENCI, L., ROCHA-MENDES, F., BERBARDO, C. S. S.,

BUENO, R. S., NOBRE, R., BOVENDORP, R. S., MARQUES, R. M., MEIRELLES, F., GOBBO, S. K., BECA,

G., SCHMAEDECKE, G. & SIQUEIRA, T. (2016). Defaunation and biomass collapse of mammals in the

largest Atlantic Forest remnant. *Anim. Conserv.* 20: 270-281.

HAGEN, M., KISSLING, W. D., RASMUSSEN, C., DE AGUIAR, M. A. M., BROWN, L. E., CARSTENSEN, D.

W., ALVES-DOS-SANTOS, I., DUPONT, Y. L., EDWARDS, F. K., GENINI, J., GUIMARAES JR., P. R.,

JENKINS, G. B., JORDANO, P., KAISER-BUNBURY, C. N., LEDGER, M. E., MAIA, K. P., MARQUITTI, F.

M. D., MCLAUGHLIN, O., MORELLATO, L. P. C., O'GORMAN, E. J., TROJELSGAARD, K., TYLIANAKIS,

J. M., VIDAL, M. M., WOODWARD, G. & OLESEN, J. M. (2012). Biodiversity, species interactions and

ecological networks in a fragmented world. *Adv. Ecol. Res.* 46: 89-210.

HARPER, K. A., BROSOFSKE, K. D., SAUNDERS, S. C. & ROBERTS, D. A. R. (2005). Edge influence on

forest structure and composition in fragmented landscapes. *Conserv. Biol.* 19: 768-782.

HERRERA, C. M. & PELLMYR, O. (2002). Plant-animal interactions: An evolutionary approach.

Blackwell Publishing, Malden, USA.

HOBBS, R. J., ARICO, S., ARONSON, J., BARON, J. S., BRIDGEWATER, P., CRAMER, V. A., EPSTEIN, P. R.,

EWEL, J. J., KLINK, C. A., LUGO, A. E., NORTON, D., OJIMA, D., RICHARDSON, D. M., SANDERSON, E.

W., VALLADARES, F., VILA, M., ZAMORA, R. & ZOBEL, M. (2006). Novel ecosystems: theoretical and

management aspects of the new ecological world order. *Glob. Ecol. Biogeog.* 15: 1-7.

HOWE, H. F. & SMALLWOOD, J. (1982) Ecology of seed dispersal. *Annu. Rev. Ecol. Syst.* 13: 201-228.

HOWE, H. F. (2016). Making dispersal syndromes and networks useful in tropical conservation and

restoration. *Glob. Ecol. Biogeog*. 6: 152-178.

ISBELL, F., CALCAGNO, V., HECTOR, A., CONNOLLY, J., HARPOLE, W. S., REICH, P. B., SCHERER-

LORENZEN, M., SCHMID, B., TILMAN, D., VAN RUIJVEN, J., WEIGELT, A., WILSEY, B. J., ZAVALETA,

E. S. & LOREAU, M. (2011). High plant diversity is needed to maintain ecosystem services. *Nature*

477: 199-U96.

JANZEN, D. H. (1974). Deflowering of Central-America. *Nat. Hist.* 83: 48-53.

JOLY, C. A., METZGER, J. P. & TABARELLI, M. (2014). Experiences from the Brazilian Atlantic Forest:

ecological findings and conservation initiatives. *New Phyt.* 204: 459-473.

JORDANO, P. (1995). Angiosperm fleshy-fruits and seed dispersers - a comparative-analysis of

adaptation and constraints in plant-animal interactions. *Am. Nat.* 145: 163-191.

JORDANO, P., BASCOMPTE, J. & OLESEN, J. M. (2003). Invariant properties in coevolutionary networks

of plant-animal interactions. *Ecol. Lett.* 6: 69-81.

KIERS, E. T., PALMER, T. M., IVES, A. R., BRUNO, J. F. & BRONSTEIN, J. L. (2010). Mutualisms in a

changing world: an evolutionary perspective. *Ecol. Lett.* 13: 1459-1474.

LALIBERTÉ, E. & TYLIANAKIS, J. M. (2010). Deforestation homogenizes tropical parasitoid-host

networks. *Ecology* 91: 1740-1747.

LAURANCE, W. F. (2002). Hyperdynamism in fragmented habitats. *J. Veg. Sci.* 13: 595-602.

LAURANCE, W. F., NASCIMENTO, H. E. M., LAURANCE, S. G., ANDRADE, A., EWERS, R. M., HARMS, K.

E., LUIZAO, R. C. C. & RIBEIRO, J. E. (2007). Habitat fragmentation, variable edge effects, and the

landscape-divergence hypothesis. *Plos One* 2: e1017.

LAURANCE, W. F., CAMARGO, J. L. C., LUIZÃO, R. C. C., LAURANCE, S. G., PIMM, S. L., BRUNA, E. M.,

STOUFFER, P. C., WILLIAMSON, G. B., BENÍTEZ-MALVIDO, J., VASCONCELOS, H. L., VAN HOUTAN, K.

S., ZARTMAN, C. E., BOUYLE, S. A., Didham, R. K., Andrade, A. & Lovejoy, T. E. (2010). The fate

of Amazonian forest fragments: A 32-year investigation. *Biol. Conserv.* 144: 56-57.

LAWTON, J. H. (1999). Are there general laws in ecology? *Oikos* 84: 177–192.

LEWINSOHN, T. M. & PRADO, P. I. (2006). Structure in plant-animal interaction assemblages. *Oikos* 113:

174-184.

LOBO, D., LEÃO, T., MELO, F. P. L., SANTOS, A. M. M. & TABARELLI, M. (2011). Forest fragmentation

drives Atlantic Forest of northeastern Brazil to biotic homogenization. *Divers. Distrib.* 17: 287-296.

MACARTHUR, R. H. & WILSON, E. O. (1967). The theory of island biogeography. Princeton Univ.

Press: Princeton.

MAY, R. M. (1972). Will a large complex system be stable? *Nature* 238: 413-414.

MCCONKEY, K. R. & O'FARRILL, G. (2016). Loss of seed dispersal before the loss of seed dispersers.

Biol. Conserv. 201: 38-49.

MELLO, M. A. R., RODRIGUES, F. A., COSTA, L. DA F., KISSLING, W. D., SEKERCIOĞLU, Ç. H.,

MARQUITTI, F. M. D. & KALKO, E. K. V. (2015). Keystone species in seed dispersal networks are

mainly determined by dietary specialization. *Oikos* 124: 1031-1039.

MEMMOTT, J. (2009). Food webs: a ladder for picking strawberries or a practical tool for practical

problems? *Philos. Trans. R. Soc. Lond. B Biol. Sci*. 364: 1693-1699.

MONTOYA, D., YALLOP, M. L. & MEMMOTT, J. (2015). Functional group diversity increases with

modularity in complex food webs. *Nature Comm.* 6: 7379-7379.

MORANTE-FILHO, J. C., ARROYO-RODRIGUEZ, V. & FARIA, D. (2016). Patterns and predictors of beta-

diversity in the fragmented Brazilian Atlantic forest: a multiscale analysis of forest specialist and

generalist birds. *J. Anim. Ecol.* 85: 240-250.

MORANTE-FILHO, J. C., FARIA, D., MARIANO-NETO, E. & RHODES, J. (2015). Birds in Anthropogenic

Landscapes: The Responses of Ecological Groups to Forest Loss in the Brazilian Atlantic Forest.

Plos One 10: e0128923.

MURCIA, C. (1995). Edge effects in fragmented forests: implications for conservation. *Trends Ecol.*

Evol. 10: 58-62.

OLDEN, J. D., POFF, N. L., DOUGLAS, M. R., DOUGLAS, M. E. & FAUSCH, K. D. (2004). Ecological and

evolutionary consequences of biotic homogenization. *Trends Ecol. Evol.* 19: 18-24.

PASSMORE, H. A., BRUNA, E. M., HEREDIA, S. M. & VASCONCELOS, H. L. (2012). Resilient networks of

ant-plant mutualists in Amazonian forest fragments. *Plos One* 7: e39108.

PÉREZ-MENDEZ, N., JORDANO, P., GARCIA, C. & VALIDO, A. (2016). The signatures of Anthropocene

defaunation: cascading effects of the seed dispersal collapse. *Sci. Rep.* 6: 24820.

PINHEIRO, J., BATES, D. & DEBROY, S. (2016). nlme: Linear and Nonlinear Mixed Effects Models.

Barton, K. (2013). R package version 3.1-137. Available at: https://cran.r-

project.org/web/packages/nlme/index.html (accessed 16 Aug 2019).

PIZO, M. A. & DOS SANTOS, B. T. P. (2011). Frugivory, post-feeding flights of frugivorous birds and the

movement of seeds in a Brazilian fragmented landscape. *Biotropica* 43: 335-342

R DEVELOPMENT CORE TEAM (2014). R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. Available at: http://www.R-project.org

(accessed 5 April 2018).

RIBEIRO, M. C., METZGER, J. P., MARTENSEN, A. C., PONZONI, F. J. & HIROTA, M. M. (2009). The

Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed?

Implications for conservation. *Biol. Conserv.* 142: 1141-1153.

RYBICKI, J. & HANSKI, I. (2013). Species–area relationships and extinctions caused by habitat loss and

fragmentation. *Ecol. Lett.* 16: 27–38.

SAAVEDRA, F., HENSEN, I., BECK, S. G., BOEHNING-GAESE, K., LIPPOK, D., TOEPFER, T. & SCHLEUNING,

M. (2014). Functional importance of avian seed dispersers changes in response to human-induced

forest edges in tropical seed-dispersal networks. *Oecologia* 176: 837-848.

SCHLEUNING, M., INGMANN, L., STRAUSS, R., FRITZ, S. A., DALSGAARD, B., DEHLING, D. M., PLEIN, M.,

SAAVEDRA, F., SANDEL, B., SVENNING, J. C., BOHNING-GAESE, K. & DORMANN, C. F. (2014).

Ecological, historical and evolutionary determinants of modularity in weighted seed-dispersal

networks. *Ecol. Lett.* 17: 454-463.

SCHLEUNING, M., FRUND, J., KLEIN, A. M., ABRAHAMCZYK, S., ALARCON, R., ALBRECHT, M.,

ANDERSSON, G. K. S., BAZARIAN, S., BOHNING-GAESE, K., BOMMARCO, R., DALSGAARD, B.,

DEHLING, D. M., GOTLIEB, A., HAGEN, M., HICKLER, T., HOLZSCHUH, A., KAISER-BUNBURY, C. N.,

KREFT, H., MORRIS, R. J., SANDEL, B., SUTHERLAND, W. J., SVENNING, J. C., TSCHARNTKE, T.,

WATTS, S., WEINER, C. N., WERNER, M., WILLIAMS, N. M., WINQVIST, C., DORMANN, C. F. &

BLÜTHGEN, N. (2012). Specialization of mutualistic interaction networks decreases toward tropical

latitudes. *Curr. Biol.* 22: 1925-1931.

SEKERCIOĞLU, C. H., LOARIE, S. R., BRENES, F. O., EHRLICH, P. R. & DAILY, G. C. (2007). Persistence

of forest birds in the Costa Rican agricultural countryside. *Conserv. Biol.* 21: 482-494.

SFAIR, J. C., ARROYO-RODRIGUEZ, V., SANTOS, B. A. & TABARELLI, M. (2016). Taxonomic and

functional divergence of tree assemblages in a fragmented tropical forest. *Ecol. Appl.* 26: 1816-

1826.

TABARELLI, M., PERES, C. A. & MELLO, F. P. L. (2012). The 'few winners and many losers' paradigm

revisited: Emerging prospects for tropical forest biodiversity. *Biol. Conserv.* 155: 136-140.

TELLA, J. L., DENES, F. V., ZULIAN, V., PRESTES, N. P., MARTINEZ, J., BLANCO, G. & HIRALDO, F.

(2016). Endangered plant-parrot mutualisms: seed tolerance to predation makes parrots pervasive

dispersers of the Parana pine. *Sci. Rep.* 6: 31709.

THÉBAULT, E. & FONTAINE, C. (2010). Stability of ecological communities and the architecture of

mutualistic and trophic networks. *Science* 329: 853-856.

THOMPSON, J. (2005). The Geographic Mosaic of Coevolution. University of Chicago Press, Chicago,

IL.

TYLIANAKIS, J. M. & MORRIS, R. M. (2017). Ecological networks across environmental gradients. *Ann.

Rev. Ecol. Evol. Syst.* 48: 25-48.

VALDOVINOS, F. S. (2019) Mutualistic Networks: Moving closer to a predictive theory. *Ecol. Lett.* doi:

10.1111/ele.13279.

VÁZQUEZ, D. P., C. J. MELIAN, N. M. WILLIAMS, N. BLÜTHGEN, B. R. KRASNOV & R. POULIN. (2007).

Species abundance and asymmetric interaction strength in ecological networks. *Oikos* 116: 1120-

1127.

VÉLEZ, M. C. D., W. R. SILVA, M. A. PIZO & L. GALETTO. (2015). Movement patterns of frugivorous

birds promoting functional connectivity among chaco serrano woodland fragments in Argentina.

Biotropica 47:475-483

VIDAL, M. M., HASUI, E., PIZO, M. A., TAMASHIRO, J. Y., SILVA, W. R. & GUIMARÃES JR., P. R. (2014).

Frugivores at higher risk of extinction are the key elements of a mutualistic network. *Ecology* 95:

3440-3447.

WARTON, D. I. & HUI, F. K. C. (2011). The arcsine is asinine: the analysis of proportions in ecology.

Ecology 92: 3-10.

ZUUR, A. F., IENO, E. N., WALKER, N. J., SAVELIEV, A. A. & SMITH, G. M. (2009). Mixed Effects

Models and Extensions in Ecology with R. Springer, New York, USA.

YOUNG, H. S., MCCAULEY, D. J., GALETTI, M. & DIRZO, R. (2016). Patterns, causes, and consequences

of Anthropocene defaunation. *Annu. Rev. Ecol. Syst.* 47: 333-358.

TABLES

TABLE 1. Results of the Linear Regression Models after AICc model selection testing the effects

of fragment area on the structure of avian seed-dispersal networks. Area corresponds to the

logarithmic scale of hectares per fragment. Intensity corresponds to the average number of

interactions per species and was used as a fixed parameter to control for differences in sampling

effort among studies. Only models with delta AICc < 2 were selected as plausible explanations

for the observed patterns (see Table S1 for full model selection). wAICc gives an estimate of the

probability of that given model to be the best choice under the AICc criteria. r^2 gives an

estimation of the variance explained by the optimal model while the coefficient t corresponds to

the importance of that parameter within the model ($t > 2$ indicates the coefficient is significant

with > 95% confidence). '+' indicates that the next presented variable is co-varying to explain the

changes on the corresponding network metric in the corresponding model.

TABLE 1.

Network metric	Best model	*w*AICc	Est	SE	*t*	*r*^{2}
Number of bird species	Area +	0.75	0.08	0.03	2.39	0.29
	Intensity		0.17	0.14	1.28	
Number of plant species	Area +	0.84	0.14	0.05	2.66	0.28
	Intensity		-0.23	0.22	-1.05	
Number of interactions	Area +	0.96	0.12	0.03	3.45	0.72
	Intensity		0.70	0.14	4.98	
Number of links	Area +	0.86	0.15	0.05	3.25	0.41
	Intensity		0.19	0.19	1.03	
Nestedness	Area	0.72	-1.13	0.47	-2.43	0.50
	Intensity		6.74	1.89	3.56	
H2' specialization	Intensity	0.68	0.08	0.07	1.02	0.01
	Area +	0.31	-0.02	0.02	-1.35	0.06
	Intensity		0.09	0.07	1.18	

576
577

FIGURE LEGENDS

FIGURE 1. The structure of the 16 avian seed-dispersal networks and their location in the

Atlantic Forest, SE Brazil. In each network, blue dots represent plant species, orange dots bird

species, and the grey lines are the seed-dispersal interactions among them; the width of the grey

lines correspond to the frequency of interactions in each area. Communities are numbered

according to a decreasing order of fragment area (see Table S1 for details of each site): 1- Parque

Estadual (PE) Intervales; 2- PE Carlos Botelho; 3- PE Ilha do Cardoso; 4- PE Poço das Antas; 5-

PE Ilha Anchieta; 6-Minas Gerais fragment; 7- Estação Ecológica de Aracuri; 8- Mata Santa

Genebra; 9- Itatiba fragment; 10- Rio Claro fragment; 11- restored area 15 years old; 12- restored

area 25 years old; 13- restored area 57 years old; 14- São Paulo fragment; 15- restored area 8

years old; 16- Rio de Janeiro fragment.

FIGURE 2. The effects of fragment area on the structure of seed-dispersal networks of the

Atlantic Forest. Each point corresponds to a single forest fragment. The gradient of colours

correspond to the intensity of defaunation ('def') in each fragment, calculated as the difference

between the total body mass of the frugivore birds expected for the Atlantic Forest (based on

Bello et al. 2017) and the total body mass of bird frugivores recorded in each network.

Defaunation and fragment area are highly correlated (r = - 0.89).

SUPPLEMENTARY INFORMATION

Content

Table S1-S4.

Fig. S1