Seed-dispersal networks lose complexity in tropical forest fragments

Carine Emer[1*], Pedro Jordano[2], Marco A. Pizo[3], Milton C. Ribeiro[1], Fernanda R. da

Silva[4] and Mauro Galetti[1]

[1]Departamento de Ecologia, Instituto de Biociências, Universidade Estadual Paulista

(UNESP), CP 199, 13506-900, Rio Claro SP, Brazil

[2]Integrative Ecology Group, Estación Biológica de Doñana (EBD-CSIC), Consejo

Superior de Investigaciones Científicas, Avenida Americo Vespucio 26, E – 41092,

Sevilla, Spain

[3]Departamento de Zoologia, Universidade Estadual Paulista (UNESP), CP 199,

13506-900, Rio Claro SP, Brazil

[4] Instituto de Biologia, Universidade de Campinas (UNICAMP), R. Monteiro Lobato,

255 - Cidade Universitária, 13083-862, Campinas SP, Brazil

*corresponding author: c.emer09@gmail.com

ABSTRACT

Seed dispersal interactions are a key ecological process in tropical forests that help to

maintain ecosystem functioning. Yet this functionality may be threatened by

increasing habitat loss, defaunation and fragmentation. However, generalist species,

and their interactions, can benefit from the habitat change caused by human

disturbance while more specialized interactions mostly disappear. Therefore changes

in the structure of the local, within fragment networks can be expected. Here we

investigated how the structure of seed-dispersal networks changes in a gradient of

increasing habitat fragmentation. We analysed 16 bird seed-dispersal assemblages

located in forest fragments of a biodiversity-rich ecosystem. We found significant

species-, interaction- and network-area relationships. The number of frugivore bird

and plant species, and their interactions, decreases as area is lost in the fragmented

landscape. Contrarily, network nestedness has a negative relationship with fragment

area, suggesting an increasing generalization of network structure with habitat

fragmentation. Modularity and interaction evenness were not significantly affected by

area though. Therefore, the local extinction of the partner species, paralleled by a loss

of interactions and more specialized seed dispersal associations, suggests the lost of

community complexity linked to a functional homogenization of the system as area is

lost. Our study provides empirical evidence for network-area relationships, yet the

lack of significant changes in modularity and interaction evenness point towards the

persistence of functional groups throughout fragmented landscapes.

RESUMO

Interações de dispersão de sementes formam um processo ecológico chave em

florestas tropicais onde colaboram na manutenção do funcionamento do ecossistema.

Porém, esta funcionalidade pode estar ameaçada pelo aumento na perda e

fragmentação do habitat. Enquanto espécies generalistas e suas interações podem se

beneficiar da mudança de habitat causada por distúrbios antrópicos, interações

envolvendo espécies mais especialistas são, na maioria, eliminadas. Desta forma,

mudanças nas redes locais, dentro de fragmentos florestais, são esperadas. Neste

trabalho nós investigamos como a estrutura de redes de dispersão de sementes mudam

em um gradiente de fragmentação do habitat. Nós analisamos 16 comunidades de

dispersão de sementes espacialmente explícitas e distribuídas em fragmentos florestais

de um ecossistema rico em biodiversidade. Nós encontramos relações significativas

entre área do fragmento e espécies, interações e estrutura das redes. O número de

espécies de aves frugívoras, plantas e as interações entre eles diminui conforme a área

dos fragmentos é perdida. Por outro lado, o aninhamento da rede mostrou uma relação

negativa com a área do fragmento, sugerindo um aumento da generalização da

estrutura das redes com a fragmentação do habitat. Modularidade e equitabilidade das

interações não foram afetadas pela área. Sendo assim, a extinção local de espécies

parceiras, conjuntamente com a perda das interações e associações planta-dispersor

mais especializadas, sugere uma perda de complexidade na comunidade ligada à

homogeneização do sistema conforme a área do fragmento é perdida. Nosso estudo

fornece evidências empíricas para as relações rede-área; por outro lado, a falta de

mudanças significativas na modularidade e equitabilidade das interações aponta para a

persistência de grupos funcionais em paisagens fragmentadas.

Keywords: avifauna; frugivory; species interactions; human disturbance; defaunation;

forest fragmentation; network-area relationship; Atlantic Forest.

INTRODUCTION

THE ON-GOING, FAST-PACED DEFORESTATION HAS CREATED FRAGMENTED FORESTS

POSING INCREASING challenges to the conservation of species interactions and

ecological processes in human-modified landscapes (Janzen 1974, Bruna et al.

2005, McConkey et al. 2016, Tylianakis & Morris 2017). Even though

fragmentation is well documented to affect species persistence, gene flow and the

connectivity among forest patches (da Silva & Tabarelli 2000, Rybicki & Hanski

2013, Hagen et al. 2012, Carvalho *et al.* 2016), the remnant fragments are also

proven to hold an important fraction of biodiversity (Barlow *et al.* 2007, Bongers

et al. 2015, Emer et al. 2018). The pool of species and interactions persisting in

forest fragments is formed by a non-random subset of smaller-sized species while

the whole spectrum of sizes found at the landscape level can only occur in large,

more pristine areas (Aizen et al. 2012, Tylianakis & Morris 2017, Emer et al.

2018). Such local assemblages configure local networks of interacting partner

species whose structure may be affected by the constraints imposed by fragment

area (Thompson & Townsend 2005, Aizen et al. 2008, Bomfim et al. 2018) while

resilience to habitat loss may occur in specific systems (e.g., Passmore et al.

2012). A recent theoretical framework has proposed different mechanisms by

which network-area relationships may emerge, such as the constraints imposed by

species-area relationships, the selection towards generalist species and the

dispersal limitation in smaller spatial scales (Galiana et al. 2018). Yet, whether

network structure is robust enough to be maintained across fragmented forests, or

whether it changes as habitat area is lost is still an open question to be tested with

empirical data.

Ecological networks are the outcomes from the associations between

interacting species. Mutualistic associations between animal and plants are a

fundamental process to maintain forest dynamics (Dennis et al. 2007, Herrera &

Pellmyr 2002). Seed dispersal by birds, in which both interacting species benefit

by using resource from one another, are ubiquitous in tropical forests in which up

to 90 percent of the plant species rely on birds to be effectively dispersed (Howe

& Smallwood 1982, Jordano 1995). Such associations between tropical frugivore

birds and the plants they feed on can form complex and diverse seed-dispersal

networks, characterized by a large number of species and interactions (Bascompte

& Jordano 2007, Schleuning et al. 2014, Dügger et al. 2018). Disruptions in the

functionality of this system can be inferred by the emergence of ecological

patterns depicted from the correct interpretation of the structural properties of the

networks (Bascompte & Jordano, 2007, Memmott, 2009, Howe 2016). For

instance, changes in the distribution of links between species can be inferred by

estimating interaction evenness, which indicates the presence of dominant and/or

abundant species in the community that concentrates most of the interactions

(Dormann et al. 2009). Likewise, increases in nestedness, a pervasive pattern on

mutualistic networks that indicates the presence of highly generalist species

interacting with both generalist and specialist ones, may suggest the loss of

specialist species in the community (Almeida-Neto et al. 2008). Similarly,

decreases in network modularity, i.e, the occurrence of subgroups of species that

interact more strongly among themselves than with species outside their group

(Olesen et al. 2007) may point towards the lost of interaction specialization or a

shift to generalist species associations (Fortuna et al. 2010, Emer et al. 2013).

125   Thus, the winner species and their winner interactions able to thrive

despite the disturbances imposed by habitat loss and fragmentation may be the

ones defining the configuration of the local network structures (Hobbs et al. 2006,

Tabarelli et al. 2012, Vidal et al. 2014). For instance, diet generalist species are

expected to be more resilient to habitat change due to their ability to feed on, and

to be dispersed by, a wider range of interacting partners (Tabarelli et al. 2012,

Morante-Filho et al. 2015). Likewise, generalist small-bodied bird species and

plant species typical of secondary forest may form loose, generalized

associations in fragmented landscapes (Emer et al. 2018), which may result in

an increased generalization in seed dispersal networks. Furthermore, habitat

fragmentation generates changes on biotic and abiotic conditions (Murcia 1995,

Laurance et al. 2007) such as increases in temperature and light conditions that

may favour the competitive ability of some plant species over the others (Laurance

et al. 2010, Sfair et al. 2016). Associated to the negative effects of habitat

reduction lays defaunation, a pervasive phenomenon characterized by the local or

functional extinction of a given species, generally towards the larger body mass

species and often related to the reduction of habitat area (Dirzo et al., 2014, Young

et al. 2016). Thus, interactions performed by large-bodied species dispersing

large-seeded plants are expected to be absent in smaller area (Pérez-Mendez et al.

2016, Emer et al. 2018) therefore affecting the local, within-fragment network

structure.

Here we aim to understand how the reduction of habitat through landscape

fragmentation affects the structure of tropical seed-dispersal networks. We

gathered data of avian-seed dispersal interactions located in 16 fragments of the

Atlantic Forest, a highly diverse and threaten tropical biome (Ribeiro et al. 2009).

Our first prediction is that the number of species present in each fragment will

follow a species-area relationship (MacArthur & Wilson 1967, Rybicki & Hanski

2013), i.e., the number of species reduces according to the loss of habitat area.

Secondly, because smaller area fragments are disturbed habitats inhabited mostly

by generalist species (Morante-Filho et al. 2016, Emer et al. 2018) able to cope

with the biotic and abiotic changes promoted by edge-effects associated to other

human-disturbance (Laurance et al. 2007), we expect a positive relationship

between fragment area and bird-seed dispersal interactions. Because the number of

bird-seed dispersal interactions in forest fragments is selectively reduced towards

generalist species (Morante-Filho et al. 2016, Emer et al. 2018), we expect that the

remaining generalist interactions will lead to changes in network structure towards

higher nestedness and lower modularity, with the concentration of interaction links

within a few dominant species, therefore decreasing interaction evenness.

METHODS

DATASET

We compiled 16 studies of avian plant-frugivore interactions sampled at the

community level along the Atlantic Forest biome, a hotspot of biodiversity highly

threaten by increasing human pressure (Ribeiro et al. 2009, Joly et al. 2014; Fig. 1,

Table S1). The study areas varied from 0.66 to 42,000 ha, in a gradient of

disturbance from semi-pristine Biological Reserves and State Parks to secondary

forest fragments and restored private lands. The matrix surrounding the fragments

is variable, including sugar cane fields, crop plantations, secondary forest, and

urban areas. Overall the studies included here originally aimed to record bird-

eating-fruit interactions and not necessarily effective seed dispersal; therefore we

carefully checked every dataset and removed any interaction that would not

characterize plant dispersal events, such as seed predation. In the particular case of

parrots we excluded them from the analysis despite some rare evidences pointing

to their role in effective seed dispersal of large nuts (Tella *et al.* 2016); in most

cases we could not establish in the original papers whether the frequency of

interactions involving parrots actually implied legitimate dispersal. We updated

and standardized plant and bird species names using the taxize package

(Chamberlain & Szocs, 2013) in R (R Development Core Team 2014).

NETWORK METRICS

For each community we built a correspondent seed-dispersal network as a bipartite

graph represented by a weighted matrix A_{ij} in which the rows represent the i plant

species and columns represent the j bird species. The elements a_{ij} indicate the

frequency of interactions between the plant species i and frugivore bird species j.

Changes in network structure were assessed by estimating:

(i) the number of bird species (A);

(ii) the number of plant species (P);

(iii) the number of interactions (I);

(iv) interaction evenness, which is based on Shannon diversity index and

calculates how equally distributed the interactions are among the species within

the network. Values close to 1 indicates that species performed an equal number of

interactions; values close to 0 indicates that the interactions are mostly

concentrated in a few species, and can be indicative of the presence of highly

generalist and/or dominant species in the system;

(v) weighted nestedness (*wNODF*), which estimates the presence of highly

generalist species interacting with less specialist ones, based on the NODF index

(Almeida-Neto et al., 2008);

(vi) modularity (M_Q), which identifies groups of species that interact more

strongly within than among groups. We used the DIRTLPAwb + algorithm

(Beckett 2016) to obtain the maximum likelihood of modularity following Girvan

& Newman (2002).

LANDSCAPE METRICS

Landscape configuration can play an important role in determining species

persistence in fragmented landscapes (Ribeiro et al. 2009) and as such may

influence NAR's as well. Therefore, for each studied community we calculated the

fragment area (ha), the functional connectivity, and the average isolation. We used

the same remnant forests map generated by Ribeiro et al. (2009) as input map for

the calculation of the landscape metrics. The maps had 50 m spatial resolution for

Albers Equal Area projection and South America Datum (SAD 69). Because some

birds can successfully move among fragments within a given distance (Pizo et al.

2011; Cornellius et al. 2015; Vélez et al. 2015), and some interactions

functionally persist throughout a fragmented landscape (Emer et al. 2018), we

estimated the functional connectivity as the amount of forest (ha) that any species

can have access if it is able to cross a 500 m matrix (Ribeiro et al., 2009; Jorge et

al., 2013). Average isolation was calculated by randomly choosing 1000 points in

a 5 km buffer around the sampling coordinates and extracting the Euclidean mean

distance (in meters) to the nearest forest fragment.

DEFAUNATION

Likewise, for each studied community we estimated the intensity of defaunation

relatively to the regional pool of species expected for the Atlantic Forest biome.

To do so, we calculated the difference between the sum of all bird body masses at

the regional landscape scale (considering all frugivore bird species present in the

Atlantic-Frugivory database [Bello et al 2017] as a proxy for the regional pool of

species) and the summed body masses of the birds that occur in each fragment.

Defaunation is based only on the presence/absence of species in a given fragment

STATISTICAL ANALYSES

To understand whether and how the structure of avian seed-dispersal networks is

affected by habitat fragmentation we used Linear Regressions and General Least

Square (gls) models in the nlme package (Pinheiro et al., 2016). Individual models

were fit for each network metric therefore considering the number of plant species,

the number of bird species, the number of interactions, evenness, nestedness and

modularity as response variables. Network metrics were logit-transformed to solve

the issue of being bounded from zero to one, when needed (Warton & Hui, 2011).

Because defaunation is negatively correlated with fragment area (r = - 0.89) and

the three landscape metrics are highly correlated among them (Table S2), we

parsimoniously chose area as the variable representing changes in habitat

configuration. Therefore, fragment area was fitted as the independent variable

along with two covariates: (i) sampling intensity, calculated as the root square of

the number of interactions divided by the root square of the network size (sum of

plant species and bird species), to control for differences in sampling effort among

studies (following Schleuning et al. 2012), and (ii) forest type, classified according

to the vegetation structure of each community (Ombrofilous Forest, Semidecidous

Forest, Araucaria Forest, and restored forest) to control for possible effects of

biotic and abiotic conditions intrinsic to each forest formation. Our spatially

explicit models and model selection follows the protocols suggested in Zuur et al.

(2009). We did not detect spatial correlation structure in our models after

accounting for the geographic coordinates of each study site in the gls models,

according to the Akaike Information Criterion (AICc). Therefore, we fit the full

model to linear regressions with 'Maximum Likelihood estimation' while applying

the 'dredge' function (Barton 2013) to select for the best fixed structure according

to AICc. 'Sampling intensity' was used as a fixed parameter to test for the effects

of area and forest type solely. Then, we used the Akaike weight of evidence

(wAICc) to obtain the relative importance of the different models (Burnham and

Anderson 2002). Finally, we present the coefficients of each predictor and the

variance explained by the optimal linear model. Last, because nestedness and

modularity can be dependent of network size, we also tested for changes in those

metrics by fitting standardized z-scores in the same model structure described

above. We started by creating null models that randomize the interactions among

species while maintaining connectance constant (Vázquez et al. 2009). Then, the

mean and standard deviation of 100 iterations were contrasted against the

observed values and the resultant z-scores were used as the response variable in

the statistical models. Network metrics and null models were estimated in the

bipartite package (Dormann et al., 2009). All analyses were run in R (R

Development Core Team, 2014).

RESULTS

Our dataset comprises 281 plant species (plus 54 plants identified and/or

morphotyped at the genus or family level) that interact with 175 frugivore bird

species. We recorded a total of 7637 interactions and 2389 unique pairwise links,

in which a bird species dispersed the seeds of a plant species in the studied

fragments of the Atlantic forest.

279   The mean frequency of interactions per fragment (472.4 ± 363.4) was 16

times greater than the mean number of bird (32.3 ± 17.9) and plant species (30.7 ±

42.6) recorded in each study site. The most abundant interactions were those

performed by generalist bird species dispersing secondary-forest plant species,

such as *Thraupis sayaca* dispersing seeds of *Cecropia pachystachya* (61 events of

seed-dispersal between three fragments; Fig S1). However, most of the events of

seed-dispersal by frugivore birds were rare given that 64% of the interactions

occurred only once across the studied fragments Fig. S1).

We found significant interaction- and network-area relationships (Fig. 2,

Table 1). The number of bird seed-dispersal interactions was negatively affected

by the reduction of fragment area while network structure became more nested as

habitat is reduced, when controlling for sampling intensity. Likewise, the species-

area relationship was also significant such as the number of bird and plant species

decreases as fragment area reduces. Modularity and interaction evenness did not

show any relationship with the reduction of fragment area though. Furthermore,

when network size was controlled for by the use of null models, sampling

intensity was the only significant predictor of nestedness while modularity remain

not affected by any of the tested predictors (Table S3).

DISCUSSION

We found significant changes on network structure following the lost of habitat

area in the fragments of the Atlantic Forest. The direction of these changes is

driven by the non-random loss of frugivore bird and plant species and the

interactions they perform, which led to an increasing generalisation of the system.

Such generalization is derived from higher network nestedness in smaller area

fragments, in which bird-seed dispersal interactions performed by small-sized

generalist species are predominant (Emer et al. 2018).

Even though nestedness and modularity are not necessarily opposite

metrics (Lewinsohn & Prado 2006; Fortuna et al. 2010) here only the first one was

significantly affected by fragment area. The increasing nestedness of the structure

of the bird-seed dispersal networks in the Atlantic Forest indicates that the

communities in smaller area fragments are connected by a distinct core of

generalist species, mostly performing generalist interactions, while there is a lack,

or at least a reduction of, specialist-specialist interactions. In other words, the

reduced pool of species in smaller fragments are likely to be using whichever

resource is available, indicating a community-level shift towards more generalist

interactions as fragmentation increases (Hobbs et al. 2006; Kiers et al., 2010).

Likewise, changes in behaviour of the species that persist in smaller fragments, or

fluctuations in population dynamics, could also lead to increasing network

generalization (Awade et al. 2017; Cornelius et al. 2017). Furthermore, the

reduced community size in smaller areas seems to be leading to some structural

changes on the organization of seed-dispersal interactions, as suggested by the

influence of sampling intensity on network nestedness when null models were

used for.

In turn, modularity, which was not significantly affected by habitat

reduction, indicates that even though network size was reduced by the loss of

species and interactions, the establishment of subgroups within the community

that interact stronger among themselves is driven by different mechanisms other

than habitat area (Schleuning et al. 2014, Mello et al. 2015). Likewise, the

distribution of interaction links among species did not change with habitat loss, as

demonstrated by the lack of significance on interaction evenness tested against

fragment area. The latest may be a result of the presence of naturalized alien

species in the system, mainly plants such as *Melia azedarach* and *Psidium*

guajava. Alien species may numerically compensate the absence of the native

ones, yet their functionality as food resource for the bird community may be

impoverished.

Species and interaction richer communities are likely to hold higher

functional diversity while the impoverished ones are likely to have gaps on

ecosystem functionality (Isbell et al., 2011; Saavedra et al. 2014, Montoya et al.,

2015), as indicated here by the reduction of network size in the gradient of

increasing fragmentation. The impoverishment of plant species in forest remnants

from the Northeast of the Atlantic Forest has led to a biotic homogenization across

the fragmented landscape (Lobo et al. 2011). Likewise, the impoverishment of

interactions found in this study and the persisting interactions performed by

smaller-sized generalist species (Emer et al. 2018) suggests that the Atlantic Forest

fragments are losing some specific functions, which may trigger a medium to

long-term functional homogenization of interactions throughout the landscape

(Olden et al. 2004, Laliberté & Tylianakis 2010) despite the current high beta-

diversity (Emer et al. 2018). The lack of significant changes on modularity and

interaction evenness according to fragment area may support our *a posteriori*

hypothesis of functional homogenization because their apparent 'robustness' to

habitat loss may be driven by the persistence of a few sub-groups among

fragments that are able to cope with the unstable conditions found in most forest

fragments (Murcia 1995, Harper et al., 2005, Passmore et al. 2012).

355   Tropical forest fragments are hyperdynamic systems facing population and

community changes in response to primary effects of habitat loss followed by

subsequent edge effects and stochastic events (Laurance et al. 2002, Laurance et

al. 2010). Furthermore, the Atlantic Forest landscape holds very high levels of

beta-diversity and turnover of birds, plants, and their interactions (Morante-Filho

et al. 2016, Farah et al. 2017, Emer et al. 2018), which could lead, at least

partially, to the observed network-area relationships. Such community

heterogeneities can be further influenced by other anthropogenic impacts such as

hunting and logging, which frequency and intensity of occurrence are context-

dependent and vary among fragments (Galetti et al. 2016). Therefore, long-term

effects of habitat fragmentation on network structure may go through different

trajectories (Thompson 2005) but the impoverishment of the system is unlikely to

change.

368   Finally, our findings have important implications for advancing the

understanding of network-area relationships (NARs, *sensu* Galiana et al. 2018).

While species-area relationships are one of the most pervasive rules in ecology

(MacArthur & Wilson 1967, Lawton 1999), empirical data demonstrating whether

or not interactions- and network-area follow the same rules are rare (but see Aizen

et al. 2012, Passmore et al. 2012, Emer et al. 2013, Bomfim et al. 2018 for

examples on mutualistic networks). We found that the spatial scale in which

ecological networks are found does influence some parameters of network

structure, but not all, at least in our study system with plant-frugivore interactions.

Furthermore, the persistence of some metrics and the changes in others may

suggest that the stability of complex system depends on the parameter analysed

(May 1972). Modularity and interaction evenness of the seed-dispersal networks

could be seen as robust to habitat loss, while network size and nestedness seem to

be not (Thébault & Fontaine 2010). Therefore, the loss of complexity and gain of

generalisation of the networks following habitat loss may provide further evidence

for an overall theory of NARs.

ACKNOWLEDGMENTS

We thank Carolina Bello to provide the map used in Figure 1 and Carolina

Carvalho for statistical advice. This work was supported by the Fundação de

Amparo à Pesquisa do Estado de São Paulo (BIOTA – FAPESP 2014/01986-0).

CE was supported by a postdoctoral fellowship (FAPESP 2015/15172-7). MG

received a fellowship from the National Council for Scientific and Technological

Development.

DATA AVAILABILITY STATEMENT

Upon acceptance, the dataset used in this manuscript will be made available on the

main author's GitHub (https://github.com/carineemer/network_area).

LITERATURE CITED

Aizen, M. A., Sabatino, M. & Tylianakis, J. M. (2012). Specialization and rarity

399   predict non-random loss of interactions from mutualist networks. *Science*

400   335: 1486-1489.

401   Almeida-Neto, M., Guimarães, P., Guimarães, P.R., Loyola, R.D. & Ulrich, W.

402   (2008). A consistent metric for nestedness analysis in ecological systems:

403   reconciling concept and measurement. *Oikos*, 117, 1227-1239.

404   Awade, M., Candia-Gallardo, C., Cornelius, C. & Metzger, J. P. (2017). High

405   emigration propensity and low mortality on transfer drives female-biased

406   dispersal of *Pyriglena leucoptera* in fragmented landscapes. *PlosOne*, 12:

407   e0170493.

408   Barlow, J., Gardner, T.A., Araujo, I.S., Avila-Pires, T.C., Bonaldo, A.B., Costa,

409   J.E., Esposito, M.C., Ferreira, L.V., Hawes, J., Hernandez, M.I.M.,

410   Hoogmoed, M.S., Leite, R.N., Lo-Man-Hung, N.F., Malcolm, J.R.,

411   Martins, M.B., Mestre, L.A.M., Miranda-Santos, R., Nunes-Gutjahr, A.L.,

412   Overal, W.L., Parry, L., Peters, S.L., Ribeiro-Junior, M.A., da Silva,

413   M.N.F., & Peres, C.A. (2007). Quantifying the biodiversity value of

414   tropical primary, secondary, and plantation forests. *Proc. Natl. Acad. Sci.*

415   *USA* 104:18555-60.

416   Barton, K. (2013). MuMIn: multi-model inference. R package version 1.42.1.

417   Available at: https://cran.r-project.org/web/packages/MuMIn/index.html

418   (accessed 06 Feb 2019).

419   Bascompte, J. & Jordano, P. (2007). Plant-animal mutualistic networks: The

420   architecture of biodiversity. *Ann. R. Ecol. Evol. Syst.* 38, 567-593.

Bruna, E. M., Vasconcelos, H. L. & Heredia, S. (2005). The effect of habitat

fragmentation on communities of mutualists: a test with Amazonian

ants and their host plants. *Biolog. Conserv.* 124: 209-216

Bello, C., Galetti, M., Montan, D., Pizo, M. A., Mariguela, T. C., Culot, L.,

Bufalo, F., Labecca, F., Pedrosa, F., Constantini, R., Emer, C., Silva, W.

R., Silva, F. R. da, Ovaskainen, O. & Jordano, P. (2017). Atlantic-

Frugivory: a plant-frugivore interaction dataset for the Atlantic Forest.

Ecology 98: 1729.

Bomfim, J. de A., Guimarães Jr. , P. R., Peres, C. A., Carvalho, G. & Cazetta, E.
(2018). Local extinctions of obligate frugivores and patch size reduction
disrupt the structure of seed dispersal networks. *Ecography* 41: 1899–
1909.
Bongers, F., Chazdon, R., Poorter, L. & Pena-Claros, M. (2015). The potential of

secondary forests. *Science*, 348, 642-643.

Burnham, K. P. & Anderson, D. R. (2002). Model selection and multimodel

inference: A practical information-theoretic approach. 2ed. Springer-

Verlag, New York, USA.

Carvalho, C.S., Galetti, M., Colevatti, R.G. & Jordano, P. (2016). Defaunation

leads to microevolutionary changes in a tropical palm. *Sci. Rep.* 6: 31957.

Chamberlain, S.A. & Szocs, E. (2013). *taxize - taxonomic search and retrieval in*

R. F1000Research.

Cornelius, C., Awade, M., C^andia-Gallardo, C., Sieving, K.E. & Metzger, J.P.

(2017). Habitat fragmentation drives inter-population variation in dispersal

behavior in a Neotropical rainforest bird. Perspect. Ecol. Evol., 15, 3–9.

da Silva, J.M.C. & Tabarelli, M. (2000). Tree species impoverishment and the

future flora of the Atlantic forest of northeast Brazil. *Nature* 404, 72-74.

Dennis, A. J., Green, R. J., Schupp, E. W. & Westcott, D. A. (2007) Seed

dispersal: Theory and its application in a changing world. CABI,

Wallingford, UK.

Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaac, N.J.B. & Collen, B.

(2014). Defaunation in the Anthropocene. *Science*, 345, 401-406.

Dormann, C.F., Jochen, F., Blüthgen, N. & Gruber, B. (2009). Indices, graphs and

null models: analyzing bipartite ecological networks. *Open Ecol. J.* 2: 7-

24.

Dügger, P., Blendinger, P.G., Bohning-Gaese, K., Chama, L., Correia, M.,

Dehling, D.M., Emer, C., Farwig, N., Fricke, E.D., Galetti, M., García, D.,

Grass, I., Heleno, R., Jacomassa, F.A.F., Moraes, S., Moran, C., Muñoz,

M.C., Neuschulz, E.L., Nowak, L., Piratelli, A., Pizo, M.A., Quitián, M.,

Rogers, H.S., Ruggera, R.A., Saavedra, F., Sánchez, M.S., Sánchez, R.,

Santilláns, V., Schabo, D.G., da Silva, F. R., Timóteo, S., Traveset, A.,

Vollstadt, M.G.R. & Schleuning, M. (2018). Seed-dispersal networks are

more specialized in the Neotropics than in the Afrotropics. *Global Ecol.*

Biogeog. DOI 10.1111/geb.12833.

Emer, C., Venticinque, E.M. & Fonseca, C.R. (2013). Effects of Dam-Induced

Landscape Fragmentation on Amazonian Ant-Plant Mutualistic Networks.

Conserv. Biol. 27: 763-773.

Emer, C. Galetti, M., Pizo, M.A., Guimarães, P.R., Moraes, S., Piratelli, A. &

Jordano. P. (2018). Seed-dispersal interactions in fragmented landscapes –

a metanetwork approach. *Ecol. Lett.* 21: 484-493.

Farah, F.T., de Muylaert, R.L., Ribeiro, M.C., Ribeiro, J.W. & de Mangueira,

J.R.S.A., et al. (2017). Integrating plant richness in forest patches can

rescue overall biodiversity in human-modified landscapes. *For. Ecol.*

Manag. 397: 78–88.

Fortuna, M.A., Stouffer, D.B., Olesen, J.M., Jordano, P., Mouillot, D., Krasnov,

B.R., Poulin, R. & Bascompte, J. (2010). Nestedness versus modularity in

ecological networks: two sides of the same coin? *J. Anim. Ecol.* 79: 811-

817.

Galiana, N., Lurgi, M., Claramunt-López, B., Fortin, M-J., Leroux, S., Cazelles,

K., Gravel, D. & Montoya, J.M. (2018). The spatial scaling of species

interaction networks. *Nat. Ecol. Evol.* 2: 782-790.

Hagen, M., Kissling, W.D., Rasmussen, C., De Aguiar, M.A.M., Brown, L.E.,

Carstensen, D.W., Alves-Dos-Santos, I., Dupont, Y.L., Edwards, F.K.,

Genini, J., Guimaraes, P.R., Jenkins, G.B., Jordano, P., Kaiser-Bunbury,

C.N., Ledger, M.E., Maia, K.P., Marquitti, F.M.D., McLaughlin, O.,

Morellato, L.P.C., O'Gorman, E.J., Trojelsgaard, K., Tylianakis, J.M.,

Vidal, M.M., Woodward, G. & Olesen, J.M. (2012). Biodiversity, species

interactions and ecological networks in a fragmented world. *Adv. Ecol.*

Res. 46: 89-210.

Harper, K.A., Brosofske, K.D., Saunders, S.C. & Roberts, D.A.R. (2005). Edge

influence on forest structure and composition in fragmented landscapes.

Conserv. Biol. 19: 768-782.

Herrera, C. M. & Pellmyr, O. (2002). Plant Animal Interactions: An Evolutionary
493   Approach. Blackwell Publishing, Malden, USA.
494   Hobbs, R.J., Arico, S., Aronson, J., Baron, J.S., Bridgewater, P., Cramer, V.A.,
495   Epstein, P.R., Ewel, J.J., Klink, C.A., Lugo, A.E., Norton, D., Ojima, D.,
496   Richardson, D.M., Sanderson, E.W., Valladares, F., Vila, M., Zamora, R.
497   & Zobel, M. (2006). Novel ecosystems: theoretical and management
498   aspects of the new ecological world order. *Glob. Ecol. Biogeog.* 15: 1-7.
499   Howe, H.F. & Smallwood, J. (1982) Ecology of seed dispersal. *Annu. Rev. Ecol.*
500   *Syst.* 13: 201-228.
501   Howe, H.F. (2016). Making dispersal syndromes and networks useful in tropical
502   conservation and restoration. *Glob. Ecol. Biogeog.* 6: 152-178.
503   Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W.S., Reich, P.B.,
504   Scherer-Lorenzen, M., Schmid, B., Tilman, D., van Ruijven, J., Weigelt,
505   A., Wilsey, B.J., Zavaleta, E.S. & Loreau, M. (2011). High plant diversity
506   is needed to maintain ecosystem services. *Nature* 477: 199-U96.
507   Janzen, D.H. (1974). Deflowering of Central-America. *Nat. Hist.* 83: 48-53.
508   Joly, C.A., Metzger, J.P. & Tabarelli, M. (2014). Experiences from the Brazilian
509   Atlantic Forest: ecological findings and conservation initiatives. *New Phyt.*
510   204: 459-473.
511   Jordano, P. (1995). Angiosperm fleshy-fruits and seed dispersers - a comparative-
512   analysis of adaptation and constraints in plant-animal interactions. *Am.*
513   *Nat.* 145: 163-191.
514   Jordano, P., Bascompte, J. & Olesen, J.M. (2003). Invariant properties in
515   coevolutionary networks of plant-animal interactions. *Ecol. Lett.* 6: 69-81.
Kiers, E.T., Palmer, T.M., Ives, A.R., Bruno, J.F. & Bronstein, J.L. (2010).

Mutualisms in a changing world: an evolutionary perspective. *Ecol. Lett.*

13: 1459-1474.

Laliberté, E. & Tylianakis, J.M. (2010). Deforestation homogenizes tropical

parasitoid-host networks. *Ecology* 91: 1740-1747.

Laurance, W.F. (2002). Hyperdynamism in fragmented habitats. *J. Veg. Sci.* 13:

595-602.

Laurance, W.F., Nascimento, H.E.M., Laurance, S.G., Andrade, A., Ewers, R.M.,

Harms, K.E., Luizao, R.C.C. & Ribeiro, J.E. (2007). Habitat

fragmentation, variable edge effects, and the landscape-divergence

hypothesis. *Plos One* 2: e1017.

Laurance, W. F., Camargo, J. L. C., Luizão, R. C. C., Laurance, S. G., Pimm, S.

L., Bruna, E. M., Stouffer, P. C., Williamson, G. B., Benítez-Malvido, J.,

Vasconcelos, H. L., van Houtan, K. S., Zartman, C. E., Bouyle, S. A.,

Didham, R. K., Andrade, A. & Lovejoy, T. E. (2010). The fate of

Amazonian forest fragments: A 32-year investigation. *Biol. Conserv.* 144:

56-57.

Lawton, J. H. (1999). Are there general laws in ecology? *Oikos* 84: 177–192.

Lewinsohn, T. M. & Prado, P. I. (2006). Structure in plant-animal interaction

assemblages. *Oikos* 113: 174-184.

Lobo, D., Leão, T., Melo, F. P. L., Santos, A. M. M. & Tabarelli, M. (2011).

Forest fragmentation drives Atlantic Forest of northeastern Brazil to biotic

homogenization. *Divers. Distrib.* 17: 287-296.

MacArthur, R. H. & Wilson, E. O. (1967). The Theory of Island Biogeography.

Princeton Univ. Press: Princeton.

May, R. M. (1972). Will a large complex system be stable? *Nature* 238: 413-414.

McConkey, K. R. & O'Farrill, G. (2016). Loss of seed dispersal before the loss of

seed dispersers. *Biol. Conserv.* 201: 38-49.

Mello, M. A. R., Rodrigues, F. A., Costa, L. da F., Kissling, W. D., Sekercioğlu,

Ç. H., Marquitti, F. M. D. & Kalko, E. K. V. (2015). Keystone species in

seed dispersal networks are mainly determined by dietary specialization.

Oikos 124: 1031-1039.

Memmott, J. (2009). Food webs: a ladder for picking strawberries or a practical

tool for practical problems? *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364:

1693-1699.

Montoya, D., Yallop, M. L. & Memmott, J. (2015). Functional group diversity

increases with modularity in complex food webs. *Nature Comm.* 6: 7379-

7379.

Morante, J. C., Arroyo-Rodriguez, V. & Faria, D. (2016). Patterns and predictors

of beta-diversity in the fragmented Brazilian Atlantic forest: a multiscale

analysis of forest specialist and generalist birds. *J. Anim. Ecol.* 85: 240-

250.

Morante, J. C., Faria, D., Mariano-Neto, E. & Rhodes, J. (2015). Birds in

Anthropogenic Landscapes: The Responses of Ecological Groups to Forest

Loss in the Brazilian Atlantic Forest. *Plos One* 10: e0128923.

Murcia, C. (1995). Edge effects in fragmented forests: implications for

conservation. *Trends Ecol. Evol.* 10: 58-62.

Olden, J. D., Poff, N. L., Douglas, M. R., Douglas, M. E. & Fausch, K. D. (2004).

564   Ecological and evolutionary consequences of biotic homogenization.

565   *Trends Ecol. Evol.* 19: 18-24.

566   Passmore, H. A., Bruna, E. M., Heredia, S. M. & Vasconcelos, H. L. (2012).

567   Resilient networks of ant-plant mutualists in Amazonian forest fragments.

568   *Plos One* 7: e39108.

569   Perez-Mendez, N., Jordano, P., Garcia, C. & Valido, A. (2016). The signatures of

570   Anthropocene defaunation: cascading effects of the seed dispersal collapse.

571   *Sci. Rep.* 6: 24820.

572   Pinheiro, J., Bates, D. & DebRoy, S. (2016). *nlme: Linear and Nonlinear Mixed*

573   *Effects Models*.

574   Pizo, M. A. e B. T. P. dos Santos. 2011. Frugivory, post-feeding flights of

575   frugivorous birds and the movement of seeds in a Brazilian fragmented

576   landscape. Biotropica 43: 335-342

577   R Core Team (2014). R: a language and environment for statistical computing. R

578   Foundation for Statistical Computing, Vienna, Austria. Available at:

579   http://www.R-project.org (accessed 5 April 2018).

580   Ribeiro, M. C., Metzger, J. P., Martensen, A. C., Ponzoni, F. J. & Hirota, M. M.

581   (2009). The Brazilian Atlantic Forest: How much is left, and how is the

582   remaining forest distributed? Implications for conservation. *Biol. Conserv.*

583   142: 1141-1153.

584   Rybicki, J. & Hanski, I. (2013). Species–area relationships and extinctions caused

585   by habitat loss and fragmentation. *Ecol. Lett.* 16: 27–38.

Saavedra, F., Hensen, I., Beck, S. G., Boehning-Gaese, K., Lippok, D., Toepfer, T.

& Schleuning, M. (2014). Functional importance of avian seed dispersers

changes in response to human-induced forest edges in tropical seed-

dispersal networks. *Oecologia* 176: 837-848.

Schleuning, M., Ingmann, L., Strauss, R., Fritz, S. A., Dalsgaard, B., Dehling, D.

M., Plein, M., Saavedra, F., Sandel, B., Svenning, J. C., Bohning-Gaese,

K. & Dormann, C. F. (2014). Ecological, historical and evolutionary

determinants of modularity in weighted seed-dispersal networks. *Ecol.*

Lett. 17: 454-463.

Schleuning, M., Frund, J., Klein, A. M., Abrahamczyk, S., Alarcon, R., Albrecht,

M., Andersson, G. K. S., Bazarian, S., Bohning-Gaese, K., Bommarco, R.,

Dalsgaard, B., Dehling, D. M., Gotlieb, A., Hagen, M., Hickler, T.,

Holzschuh, A., Kaiser-Bunbury, C. N., Kreft, H., Morris, R. J., Sandel, B.,

Sutherland, W. J., Svenning, J. C., Tscharntke, T., Watts, S., Weiner, C.

N., Werner, M., Williams, N. M., Winqvist, C., Dormann, C. F. &

Bluthgen, N. (2012). Specialization of Mutualistic Interaction Networks

Decreases toward Tropical Latitudes. *Curr. Biol.* 22: 1925-1931.

Sekercioglu, C. H., Loarie, S. R., Brenes, F. O., Ehrlich, P. R. & Daily, G. C.

(2007). Persistence of forest birds in the Costa Rican agricultural

countryside. *Conserv. Biol.* 21: 482-494.

Sfair, J. C., Arroyo-Rodriguez, V., Santos, B. A. & Tabarelli, M. (2016).

Taxonomic and functional divergence of tree assemblages in a fragmented

tropical forest. *Ecol. Appl.* 26: 1816-1826.

Tabarelli, M., Peres, C. A. & Mello, F. P. L. (2012). The 'few winners and many

610   losers' paradigm revisited: Emerging prospects for tropical forest

611   biodiversity. *Biol. Conserv.* 155: 136-140.

612   Tella, J. L., Denes, F. V., Zulian, V., Prestes, N. P., Martinez, J., Blanco, G. &

613   Hiraldo, F. (2016). Endangered plant-parrot mutualisms: seed tolerance to

614   predation makes parrots pervasive dispersers of the Parana pine. *Sci. Rep.*

615   6: 31709.

616   Thébault, E. & Fontaine, C. (2010). Stability of ecological communities and the

617   architecture of mutualistic and trophic networks. *Science* 329: 853-856.

618   Thompson, J. (2005). *The Geographic Mosaic of Coevolution*. University of

619   Chicago Press, Chicago, IL.

620   Thompson, R. M. & Townsend, C. R. (2005). Food-web topology varies with

621   spatial scale in a patchy environment. *Ecology* 86: 1916-1925.

622   Tylianakis, J. M. & Morris, R. M. (2017). Ecological networks across

623   environmental gradients. *Ann. Rev. Ecol. Evol. Syst.* 48: 25-48.

624   Vélez, M. C. D., W. R. Silva, M. A. Pizo e L. Galetto. 2015. Movement patterns

625   of frugivorous birds promoting functional connectivity among chaco

626   serrano woodland fragments in Argentina. Biotropica 47:475-483

627   Vidal, M. M., Hasui, E., Pizo, M. A., Tamashiro, J. Y., Silva, W. R. & Guimarães,

628   P. R. Jr. (2014). Frugivores at higher risk of extinction are the key

629   elements of a mutualistic network. *Ecology* 95: 3440-3447.

630   Warton, D. I. & Hui, F. K. C. (2011). The arcsine is asinine: the analysis of

631   proportions in ecology. *Ecology* 92: 3-10.

Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A. & Smith, G. M. (2009).

Mixed Effects Models and Extensions in Ecology with R. Springer, New

York.

Young, H. S., McCauley, D. J., Galetti, M. & Dirzo, R. (2016). Patterns, causes,

and consequences of Anthropocene defaunation. *Annu. Rev. Ecol. Syst.* 47:

333-358.

TABLES

TABLE 1. Results of the Linear Regression Models after AICc model selection testing the effects

of fragment area on the structure of avian seed-dispersal networks. Area corresponded to the

logarithmic scale of hectares per fragment. Intensity corresponds to the average number of

interactions per species and was used to control for differences in sampling effort among studies.

Only models with delta AICc < 2 were selected as plausible explanations for the observed

patterns (see Table S1 for full model selection). *w*AICc gives an estimate of the probability of

that given model to be the best choice under the AICc criteria. r^2 gives an estimation of the

variance explained by the optimal model while the coefficient *t* corresponds to the importance of

that parameter within the model (*t* > 2 indicates the coefficient is significant with > 95%

confidence).

TABLE 1.

Network metric	Best model	*w*AICc	Est	SE	*t*	*r²*
Number of bird species	Area +	0.75	0.08	0.03	2.39	0.29
	Intensity		0.17	0.14	1.28	
Number of plant species	Area +	0.84	0.14	0.05	2.66	0.28
	Intensity		-0.23	0.22	-1.05	
Number of interactions	Area +	0.96	0.12	0.03	3.45	0.72
	Intensity		0.70	0.14	4.98	
Evenness	Intensity	0.57	0.01	0.03	0.20	-0.07
	Area +	0.40	0.01	0.01	1.62	0.04
	Intensity		<0.01	0.03	0.03	
Nestedness	Area +	0.72	-1.13	0.47	-2.43	0.50
	Intensity		6.74	1.89	3.56	
Modularity	Intensity	0.84	0.03	0.03	1.03	0.01

655
656

FIGURE LEGENDS

FIGURE 1. The structure of the 16 avian seed-dispersal networks and their location in the

Atlantic Forest. In each network, blue dots represent plant species, orange dots bird species, and

the grey lines are the seed-dispersal interactions among then; the width of the grey lines

correspond to the frequency of interactions in each area. Communities are numbered according to

a decreasing order of fragment area (see Table S1 for details of each site): 1- Parque Estadual

(PE) Intervales; 2- PE Carlos Botelho; 3- PE Ilha do Cardoso; 4- PE Poço das Antas; 5- PE Ilha

Anchieta; 6-Minas Gerais fragment; 7- Estação Ecológica de Aracuri; 8- Mata Santa Genebra; 9-

Itatiba fragment; 10- Rio Claro fragment; 11- restored area 15 years old; 12- restored area 25

years old; 13- restored area 57 years old; 14- São Paulo fragment; 15- restored area 8 years old;

16- Rio de Janeiro fragment.

FIGURE 2. The effects of fragment area on the structure of seed-dispersal networks of the

Atlantic Forest. Each point corresponds to a single forest fragment. The gradient of colours

correspond to the intensity of defaunation in each fragment, calculated as the difference from the

total body mass of the frugivore birds expected for the Atlantic Forest (based on Bello et al.

2017) and the total body mass of bird frugivores recorded in each network. Defaunation and

fragment area are highly correlated (r = - 0.89).

SUPPLEMENTARY INFORMATION

Content

Table S1-S4.

Fig. S1