Potential for transmission of *Elizabethkingia anophelis* by *Aedes albopictus* and the role of

microbial interactions in Zika virus competence

Onyango MG[1], Payne AF[1], Stout J[1], Dieme C[1], Kuo L[1], Kramer LD[1, 2], Ciota AT[1, 2*]

[1] The Arbovirus Laboratory, Wadsworth Center, New York State Department of Health, 5668

State Farm Road, Slingerlands, 12159, New York

[2] School of Public Health, State University of New York Albany, 1400 Washington Avenue,

Albany, 12222, New York

Email addresses:

MGO: Maria.Onyango@health.ny.gov

PA: anne.payne@health.ny.gov

SJ: Jessica.Stout@health.ny.gov

DC: Constentin.Dieme@health.ny.gov

KL: lili.kuo@health.ny.gov

KL: laura.kramer@health.ny.gov

CA: alexander.ciota@health.ny.gov

[*]Corresponding author

Abstract

Elizabethkingia anophelis has been the cause of four outbreaks with significant morbidity and

mortality. Its transmission routes remain unknown and no point source of infection has been

identified. Here we show that *E. anophelis* can be found in the saliva of *Aedes* mosquitoes,

suggesting the novel possibility of vector-borne transmission of this bacterium. We additionally

characterized diverse microbial communities in *Aedes* midguts, salivary glands and saliva. To

the best of our knowledge, this represents the first description of the microbiome of *Aedes*

saliva. Further, we demonstrate that increased abundance of *E. anophelis* is associated with

decreased susceptibility and replication of Zika virus (ZIKV) in the midgut of *Aedes* mosquitoes,

suggesting a novel transmission barrier for arboviruses transmitted by *Aedes* mosquitoes.

Together, these results demonstrate the complex relationships between the mosquito, the

midgut microbial community and arboviruses and offer insights into the epidemiology and

control of emerging bacterial and viral pathogens.

Keywords: *Elizabethkingia anophelis*, Zika virus and *Aedes* mosquitoes

Author Summary

Elizabethkingia anophelis has in the recent past caused outbreaks different parts of the world

resulting both in morbidity and mortality. Until now, to the best of our knowledge, no study has

been able to demonstrate that this bacterium can be transmitted by mosquitoes. We have

demonstrated for the first time that *Elizabethkingia anophelis* is present in the saliva of both

infected and non-infected *Aedes* mosquitoes. Further, we have shown that it confers an

inhibitory effect on Zika virus establishment in the midguts of *Aedes* mosquitoes. Together,

these results potentially display the potential for vector borne transmission of *E. anophelis* as

46   well as a novel transmission barrier of ZIKV. Lastly, we have for the first time characterized

47   salivary microbes of *Aedes* mosquitoes necessitating the investigation of the impact of salivary

48   microbes in severity of disease in vertebrate hosts.

49
50
51   **Introduction**
52
Elizabethkingia is a gram-negative rod-like, aerobic, non-fermenting, non-motile and non-spore

forming bacteria widely distributed in natural environments, including soil and fresh water as

well as hospital environments [1]. It was originally called Flavobacterium or belonged to CDC

group IIa but later reclassified as Chryseobacterium in 1994 [2]. On the basis of shared

phenotypic and phylogenetic characteristics, *Chryseobacterium meningosepticum* and *C.*

miricola were re-classified to a new genus, *Elizabethkingia* [3]. The most clinically important

species of the genus *Elizabethkingia* include *E. anophelis, E. meningoseptica and E. miricola* [4].

E. anophelis, originally isolated from the midgut of *Anopheles gambiae* [5] has been the cause

of moderate to large-scale outbreaks resulting in morbidity and mortality [6–10] as well as

clinical cases across different geographical settings [1,4,11,12] . *E. anophelis* is associated with

clinically significant infections and high mortality [4] relative to *E. meningoseptica* [9].

The genus *Elizabethkingia* is prevalent in *Aedes* [13–15] and *Anopheles* [16–18] species of

mosquitoes and has been detected in both field-caught and laboratory-reared mosquitoes in

Africa, Europe and North America [15,18,19]. The microbe can be transmitted among

mosquitoes vertically, transstadially and horizontally [13,17]. In humans, evidence for perinatal

vertical transmission of *E. anophelis* has been identified and associated with neonatal

meningitis [11]. While some infections are thought to be nosocomial and mechanical

ventilation has been identified as a risk factor for transmission in hospital settings [1],

transmission routes have not been identified in a number of recent individual cases or clusters.

The transition of *Elizabethkingia* from a common environmental bacterial species to a human

pathogen of clinical significance, as well as the previous identification of *Elizabethkingia* in a

number of mosquito species, prompted us to assess the abundance and tropism of

Elizabethkingia bacteria in *Aedes* mosquitoes. We identified *Elizabethkingia* in midguts, salivary

glands and saliva of *Ae. albopictus*. Sequencing of *Elizabethkingia* identified in the salivary gland

demonstrated 100% homology to *E. anophelis*.

In order to assess the relative abundance of *Elizabethkingia* within microbial communities in

these tissues, we additionally performed 16S *rRNA* sequencing. Further, to better define

interactions with arboviruses vectored by *Aedes* mosquitoes, we assessed the correlation

between *Elizabethkingia* and other microbes with Zika virus (ZIKV) in the midgut of *Ae.*

albopictus. We measured a negative correlation between *E. anophelis* and ZIKV in the midgut

and in mosquito cell culture. These data together demonstrate the complex interactions

between microbial communities and arboviruses that could have important implications for the

emergence and transmission of both viral and bacterial pathogens.

Results

Microbial profiles in *Aedes albopictus* are tissue-specific

Illumina MiSeq 16S *rRNA* sequencing resulted in 2 369 009 reads. The number of reads for

individual samples ranged from 11 925 - 105 039 with a mean of 67 334. Both the positive

spike-in (PhiX) and negative controls (water and non-template control) had limited number of

sequences reads, ranging from 1620 – 2823, and hence were discarded in downstream analysis.

In order to control for alpha and beta diversity biases, each sample was rarefied to a total

frequency of 30 000 reads, hence two samples were discarded from downstream analysis. A

total of 154 OTUs were identified in this study, 59% were Bacteroidetes, 40% Proteobacteria

and 1% Cyanobacteria. In order to visualize the distribution of bacterial genera across individual

tissues, an additional cut-off read frequency of 7 000 was applied (Figure 1).

The microbial profile was diverse across tissues (PERMANOVA, F-value=0.9998, R^2= 0.3111; P

<0.001; Figure 2). Midguts were dominated by Bacteroidetes with a higher relative abundance

identified in ZIKV infected midguts relative to the unexposed midguts. Salivary glands, on the

other hand, were dominated by Proteobacteria. Surprisingly, bacterial taxa from mosquito

saliva was much more diverse with a range of Bacteroidetes, Proteobacteria and Cyanobacteria

(Chao1 [ANOVA], P < 0.001, F-value= 20.58). This analysis clearly demonstrates the dominance

of specific genera in individual tissues. For instance, ZIKV infected midguts and salivary glands

were overwhelmingly dominated by *Elizabethkingia*. Uninfected midguts were dominated by

either *Elizabethkingia* or *Wolbachia*, yet salivary glands from uninfected individuals were

dominated by Enterobacteriaceae (Figure 1).

108
109   ***Aedes* saliva contains diverse microbiota including *Elizabethkingia anophelis***
110
The core microbes in the saliva included: *Pseudomonas, Cytophagaceae, Chitinophagaceae,*

Betaproteobacteria, Gammaproteobacteria, Comamonadaceae, Chryseobacterium,

Flavisolibacter, Bradyrhizobiaceae, MLE1_12 and *Sphingomonas*.

The most abundant taxa present in the saliva (*Elizabethkingia, Chitinophagaceae,*

Enterobacteriaceae, Wolbachia, Pseudomonas and *Betaproteobacteria*) were also present in

the salivary glands and midgut samples (Figure 1).

We observed presence of taxa of public health importance in the saliva of *Ae. albopictus*

exposed to infectious and non-infectious blood meals (Figure 1). These include *Elizabethkingia*,

Enterobacteriaceae and *Pseudomonas*.

We employed a quality control measure to assess potential sources of saliva contamination. A

cotton pad obtained from the container housing mosquitoes exposed to an infectious blood

meal, a non-infectious blood meal, a sterile cotton pad, filter sterilized 10% sucrose solution

used to feed mosquitoes, and filter sterilized mosquito diluent used to collect mosquito saliva,

were incubated overnight in LB media. The control was sterile LB media. Bacterial growth was

observed in the culture tube that had the cotton pad used to feed both mosquitoes exposed to

the infectious blood meal as well as the non-infectious blood meal. No growth occurred in the

culture tube with sterile cotton pad, sterile 10% sucrose solution, the mosquito diluent or the

negative control. Introduction of bacteria during feeding by mosquitoes cannot be ruled out for

taxa identified exclusively in saliva.

Elizabethkingia was present in both ZIKV -infected and uninfected midguts, salivary glands and

saliva. On the other hand, *Enterobacteriaceae* and *Pseudomonas* were present in the saliva and

salivary glands, but not midguts, of *Ae. albopictus* exposed to both ZIKV (infectious) and non-

infectious blood meals (Figure 1; Figure 3).

Relationship between *Elizabethkingia* and ZIKV in *Ae. albopictus*

The 16S *rRNA* amplicons of saliva, salivary glands and midgut samples were sequenced on the

Illumina MiSeq platform. *Elizabethkingia* was identified in both saliva (10 out of 18) and

salivary gland (6 out of 6) samples of *Ae. albopictus* exposed to non-infectious and infectious

blood meals (Figure 1). *Elizabethkingia* sequences were also present in the midguts of *Ae.*

albopictus exposed to infectious or non-infectious blood meal (10 out of 10) (Figure 1).

Furthermore, a higher proportion of *Elizabethkingia* sequences were identified in saliva

collected following non-infectious blood meals relative to the saliva collected following ZIKV

infection (T- test; Mann Whitney test, P = 0.0256; Figure 3; Figure 4A). Conversely, while

Elizabethkingia was identified in both the ZIKV-infected and uninfected *Aedes* salivary glands,

ZIKV-infected salivary glands had a higher proportion of *Elizabethkingia* reads relative to their

uninfected counterparts (F test, P< 0.0001; Figure 4A). Overall, the ZIKV - infected *Ae.*

albopictus midguts had a higher relative amount of *Elizabethkingia* compared to other tissues

(ANOVA; Kruskal-Wallis test, P = 0.0027; Figure 4A).

Although exposure of *Ae. albopictus* midguts to non-infectious blood meal resulted in an

increase in the amount of *Elizabethkingia* in the midgut, Real-Time (RT) qPCR assay

demonstrated a higher load of *Elizabethkingia* in the midguts exposed to ZIKV infectious blood

meals as compared to the *Ae. albopictus* midguts exposed to non-infectious blood meals (T-

test, Mann Whitney test, P = 0.0051; Figure 4B).

To further investigate the relationship between *Elizabethkingia* and ZIKV measured in *Ae.*

albopictus midguts, *Elizabethkingia* bacterial loads were quantified together with ZIKV viral

loads both *in vivo* (ZIKV-infected midguts, (N = 22) and *in vitro* (U4.4 cells co-infected with

Elizabethkingia and ZIKV; three biological replicates, 1 biological replicate, n = 4 wells) using RT-

qPCR.

Despite the fact that feeding on a ZIKV infectious blood meal was associated with higher levels

of *Elizabethkingia* (Figure 4B), fold change in expression of the *Elizabethkingia rpoB* gene from

ZIKV-infected *Ae. albopictus* (N = 22) was negatively correlated to ZIKV absolute quantities

(Spearman r = -0.3985; P < 0.05; Figure 5A). ZIKV viral loads ranged from 3 - 11 $\log_{10}$ ZIKV RNA

copies/ml, while fold change of *Elizabethkingia* ranged from -1.8 - 2.2 $\log_{10}$ CFU/ml.

To clarify the potential for direct interaction between *Elizabethkingia* and ZIKV, we performed

an *in vitro* assay on the *Ae. albopictus* derived U4.4 cell line. Cells were co-infected with

Elizabethkingia and ZIKV or infected with ZIKV only, and viral loads were compared.

At 2 dpi, we measured high amounts of ZIKV in the control, ZIKV/*Elizabethkingia* OD 1.0 (8.5

Logs CFU/ ml) and ZIKV/ *Elizabethkingia* OD0.5 (8.0 Logs CFU/ml) co-infected cells while the

ZIKV/ *Elizabethkingia* OD0.2 (7.7 Logs CFU/ml) co-infected cells had significantly lower ZIKV

titers (ANOVA; Kruskal-Wallis test, P = 0.0070). On the other hand, at 4 dpi, we measured a

significant reduction in ZIKV titers in all ZIKV/*Elizabethkingia* co-infected U4.4 cells relative to

the control (ANOVA, Kruskal-Wallis test, P = 0.0079; Figure 5B).

Phylogenetic analysis of *Elizabethkingia*

To taxonomically classify the *Elizabethkingia* strain that we identified we sequenced the

Elizabethkingia rpoB gene amplified from the cDNA of the *Ae. albopictus* salivary gland. The

phylogenetic analysis demonstrated that the *Elizabethkingia anophelis* strain identified in our

study has a 100% sequence identity to *Elizabethkingia anophelis* which was originally isolated

from the midgut of *Anopheles gambiae* [20]. This strain has a 99% identity to *Elizabethkingia*

endophytica, originally isolated from sweet corn [21]. It was closely related to the

Elizabethkingia genomosp. 4 [22], *Elizabethkingia miricola,* isolated from condensation water in

space station [23] and *Elizabethkingia bruuniana*[24]. *E. anophelis* has a 92% identity to

Elizabethkingia meningosepticum, originally isolated from human cerebrospinal fluid [25]

(Figure 6).

Discussion

This study has demonstrated *Elizabethkingia* in the saliva of *Ae. albopictus* mosquitoes. Further,

phylogenetic analysis has identified the strain of *Elizabethkingia* isolated from an *Ae. albopictus*

salivary gland to have 100% sequence identity to *Elizabethkingia anophelis*, first isolated from

the midgut of the mosquito *Anopheles gambiae*[26]. The *Elizabethkingia* genus was previously

found to be the dominant bacterial genera in the salivary glands of both *An. gambiae* and *An. coluzii*

[27]. To the best of our knowledge, this is the first study to demonstrate *E. anophelis* in the

saliva of *Aedes* mosquitoes. Our findings bear clinical significance as *E. anophelis* has been

identified as a cause of sepsis in adults and children, as well as a cause of neonatal meningitis

[9]. By 2016, *E. anophelis* had caused an outbreak in Central African Republic [6], Singapore [7]

and two outbreaks in the Midwest, United States that resulted in morbidity and mortality of ~

30% [8–10]. The Midwest outbreak occurred primarily in community settings [8] and despite

extensive investigations, no point source of infection has been identified in either the Midwest

or Singapore outbreaks [9,10]. While there is currently no evidence of mosquito-borne

transmission with any of these outbreaks, our data demonstrate the potential for involvement

of mosquitoes in the maintenance of *E. anophelis* and other bacterial pathogens.

The identification of *Elizabethkingia* in the saliva of *Ae. albopictus* infected with ZIKV

necessitates the investigation of the impact of *Elizabethkingia* on transmission efficiency of

ZIKV and other arboviruses, and additionally on how co-transmission of bacteria could influence

the early events, progression and severity of viral infections in the host. Our results present a

new dimension into the vector-host interface dynamics. Several molecules present in mosquito

saliva have been shown to suppress the host inflammatory or immune responses [28–33] thus

enhancing dissemination [34]. Specific salivary proteins such as biogenic amine-binding D7

protein [35], saliva serine protease CLIPA3 [36], salivary factor LTRIN [37] and saliva protein

AgBR1[38] modulate vertebrate infection. However, a gap in knowledge exists on the

contributory role of mosquito salivary microbes in potentiating pathogenicity or transmission

efficiency in vertebrate hosts. Salivary microbes may provide crucial targets for the control of

disease-causing pathogen transmitted by mosquitoes.

In addition to *Elizabethkingia*, we observed a proliferation of other taxa associated with human

disease throughout *Aedes* mosquitoes following exposure to a blood meal. This included

bacteria belonging to the family Enterobacteriaceae, which have been shown to be

opportunistic pathogens [39–41]. Enterobacteriaceae are gram negative bacteria that includes

Escherichia coli, Klebsiella, Salmonella, Shigella and *Yersinia pestis. Enterobacter* spp., which we

identified as among the primary bacteria associated with *Ae. albopictus*. These bacteria are

known to produce molecules with hemolytic activity [42,43] and the increase in abundance of

Enterobacteriaceae in the saliva of *Ae. albopictus* may therefore be an evolutionary mechanism

that exploits blood digestion to ensure their horizontal transfer as well as infection of new

hosts.

Wolbachia was identified in the saliva, salivary gland and midguts of *Ae. albopictus* in this study.

Wolbachia is an intracellular endosymbiont primarily transmitted vertically through the female

germline[44,45]. It can also undergo transfer between somatic and germ line cells. Horizontal

transfer independent of cell to cell contact utilizing host cell phagocytic and clathrin/dynamin-

dependent endocytic machinery is possible [46]. Our results provide further proof of the

possibility of horizontal transfer of *Wolbachia* through saliva. *Wolbachia* has been identified in

the phloem vessels and in some novel "reservoir" spherules along the phloem after *Wolbachia*-

infected whiteflies fed on cotton leaves [47]. These findings could have important implications

for the exploitation of *Wolbachia* as a biological control of tropical diseases.

A significant reduction in *Ae. albopictus* midgut microbial diversity resulting from ZIKV

infection was evident, yet the effect on individual genera was highly variable. Available

evidence demonstrates that commensal microbiota regulate and are in turn regulated by

invading viruses [48–60]. Viral infection often causes disturbance of the commensal host

microbiota, resulting in dysbiosis and subsequently affecting viral infectivity of the host

[50,51,56,61–65]. The dysbiosis of the commensal microbiota due to virus infection is mainly

due to modulation of the immune responses by the virus resulting in immune suppression

[66,67]. Our results are consistent with the findings of a study on the response of the total gut

microbiota diversity of *A. japonicus* and *A. triseriatus* to infection with La Crosse virus, which

revealed higher midgut bacterial richness among the midguts obtained from newly emerged

individuals and lower richness among midguts exposed to viral infection or a non-infectious

blood meals [50]. In addition, our study corroborates previous findings with West Nile virus

infection on the bacterial diversity of *Culex pipiens,* which demonstrated that lower diversity

was associated with virus exposure[51].

It is also notable that *Elizabethkingia* and *Wolbachia* appeared to have an exclusive presence in

the midgut of the *Ae. albopictus* mosquitoes. It is possible that there is competitive exclusion of

Wolbachia in the midgut by *Elizabethkingia*. Studies have shown that *Ae. albopictus* and *Cx.*

quinquefasciatus naturally infected by both *Asaia* and *Wolbachia* have a co-exclusion

relationship[51]. In species uninfected by *Wolbachia* such as *An. gambiae, An. stephensi* and

Ae. aegypti, Asaia is capable of colonizing reproductive organs and salivary glands [68]. The

predominance of *E. meningosepticum* in the *An. stephensi* midgut was shown to be due to its

antimicrobial properties suppressing the growth of other midgut bacteria [69].

We demonstrated a significant negative correlation between ZIKV and *Elizabethkingia* as well

as a reduction in the ZIKV titer in U4.4 cells co-infected with *Elizabethkingia* and ZIKV. Similarly,

it has been demonstrated that *E. meningoseptica* metabolites obtained from the midgut of

Anopheles stephensi exhibited antiparasitic effect on *P. falciparum* parasites *in vitro* and

demonstrated *in vitro* toxicity on gametocytes[69]. We also measured a significant increase in

Elizabethkingia bacterial load in the midgut infected with ZIKV relative to those exposed to a

non-infectious blood meal. These findings corroborate the results of a previous study aimed at

understanding the effect of ZIKV on the dynamic bacterial community harbored by *Ae. aegypti*

[70], which demonstrated that the Flavobacteriacae family was the most abundant bacterial

taxa. This was particularly the case among the ZIKV-exposed blood fed individuals, yet

Flavobacteriacae decreased in abundance among ZIKV-exposed gravid individuals [70]. The

interactions between bacteria and viruses are clearly complex. The mechanisms of interaction

mostly benefit virus infiltration into cells directly by virus binding to a bacterial cell or viral

utilization of a bacterial product [71]. Indirect mechanisms include: virus-induced increase of

bacterial receptor concentrations; virus damage to underlying epithelial cells; virus

displacement of commensal bacteria and virus suppression of the host immune system[71]. In

the direct interactions, the commensal bacteria are considered capable of outcompeting viral

pathogens and limiting tissue accessibility by competing for receptor binding sites reducing the

binding of pathogenic bacteria or viral attachment. This may potentially explain the negative

correlation that we observed in *Ae. albopictus* midguts. Our *In vitro* results which demonstrate

decreased viral loads with *Elizabethkingia* co-infection, particularly at day 4 post-infection,

corroborate previous studies [70] and clearly demonstrate the capacity for this bacterium to

independently suppress viral replication and/or persistence. Further studies delineating the

mechanism that *Elizabethkingia* utilizes to inhibit viral fitness in *Ae. albopictus* are needed, yet

these data suggest the identification of a novel transmission barrier that could ultimately be

exploited in the design of novel control measures.

Overall, to the best of our knowledge, this is the first study to demonstrate and characterize the

microbiome of the saliva of mosquitoes. These data may have significant implications on the

pathogen - host interface and our understanding of the impact of these microbes on the

severity of disease in the vertebrate host. We have also demonstrated that the salivary gland

and saliva of *Aedes* mosquitoes exposed to infectious or non-infectious bloodmeals contain the

bacterial pathogen *Elizabethkingia anophelis*. Finally, our study has demonstrated both by *in*

vivo and *in vitro* studies that *Elizabethkingia* and ZIKV are negatively correlated in the mosquito

midgut. These findings together have implications for our understanding of the interaction of

bacterial and viral agents in mosquitoes that could ultimately improve our understanding of

disease epidemiology and inform novel control measures.

291
292
293
294

Materials and methods

Virus

This study utilized frozen virus sample. The ZIKV strain was HND (2016-19563, GenBank

accession no. KX906952).

Insect sampling and DNA preparation.

 Ae. albopictus were collected from Suffolk County, NYS in 2015 (kindly provided by I. Rochlin)

and colonized at the New York State Department of Health (NYSDOH) Arbovirus Laboratory.

F15 Ae. albopictus were vacuum-hatched and maintained at 27°C under standard rearing

conditions. At eight days post mating, female adults were orally exposed to blood meals. The

blood meal consisted of either 1:1 dilution of defibrinated sheep blood plus 2.5% sucrose;

sodium bicarbonate (to adjust pH to 8.0) and virus, or a non-infectious blood meal containing a

final concentration of 2.5% sucrose solution. Infectious blood meals contained 8.3 $\log_{10}$ PFU/ml

ZIKV HND [72]. The female mosquitoes were exposed to blood meals in a 37°C pre-heated

Hemotek membrane feeding system (Discovery Workshops, Acrington, UK) with a porcine

sausage casing membrane. After an hour, the mosquitoes were anaesthetized with CO_2 and

immobilized on a pre-chilled tray connected to 100% CO_2. Engorged females were separated

and placed in three separate 0.6 L cardboard cartons (30 individuals per carton). In addition, 1

ml of each blood meal was transferred to a 1.5 ml Safe Seal microtube (Eppendorf, Hamburg

Germany) and stored at -80°C. Blood fed females were maintained on 10% sucrose solution

provided ad libitum.

At 14 dpi, the female mosquitoes were immobilized using triethylamine (Sigma Aldrich, St.

Louis, MO, USA). To examine for ZIKV dissemination in the mosquito, the legs were removed

from the mosquitoes before dissecting the gut and salivary gland from every individual. Saliva

was collected under sterile conditions by inserting the proboscis of the female mosquitoes into

a capillary tube containing ~20 µl fetal bovine serum plus 50% sucrose 1:1 for 30 minutes and

subsequently ejecting the mixture into 125 µl Mosquito Diluent (MD; 20% heat-inactivated fetal

bovine serum in Dulbecco phosphate-buffered saline plus 50 µg/ml penicillin/streptomycin, 50

µg/ml gentamicin and 2 µg/ml Fungizone (Sigma Aldrich, St. Louis, MO, USA). The midgut was

dissected from individual mosquitoes and rinsed twice in sterile PBS before transfer to sterile

microcentrifuge tubes and storage at -80°C until tested. Mosquito carcass and legs were then

placed in individual tubes containing 500 µl MD and a bead. All samples were held at -80°C until

assayed. A ZIKV-specific quantitative PCR assay that targets the NS1 region was utilized to

obtain viral titer [73] from the legs, carcass, saliva, salivary glands and midguts as described by

[72].

As a precautionary measure to prevent contamination of saliva, all the reagents used during the

rearing and blood feeding of mosquitoes were filter sterilized and cotton pads were autoclaved.

To test for potential contamination during rearing, we incubated overnight (in LB media) a

piece of cotton pad obtained from the gallon housing mosquitoes exposed to an infectious

blood meal, mosquitoes exposed to a non-infectious blood meal, a sterile cotton pad not

exposed to mosquitoes, filter sterilized 10% sucrose solution that was used to feed the

mosquitoes, filter sterilized mosquito diluent, and sterile LB Media used as a control.

Sequencing and analysis of microbiome of *Aedes* mosquito saliva, salivary glands and

midguts.

To determine the microbial profile in the *Ae. albopictus* saliva, salivary glands and midguts of

each individual mosquito, midgut, legs, carcass and saliva samples were screened for virus. The

individuals that were successfully infected and or transmitting the virus were identified. The

cDNA was prepared from the RNA samples using iScript cDNA kit (Biorad, Hercules, California).

To amplify starting material for downstream analysis, the saliva, salivary gland and midgut

samples were individually whole transcriptome amplified according to the REPLI-g WTA single

cell kit protocol (Hilden, Germany). The resulting amplified cDNA was diluted 1:100. Thereafter,

16S *rRNA* amplicon were amplified using the 16S primer set and conditions in Table 1.

The barcoded high-throughput amplicon sequencing of the bacterial 16S V3-V4 hypervariable

rRNA was done at Wadsworth Center Applied Genomics Core (WCAGC). PCR reactions were

carried out in a total volume of 50 μl, 5μM of 16S *rRNA* V3-V4 hypervariable region primers

consisting of an Illumina barcode (Table 1) [51], 2 μl WTA cDNA, 8 μl deionized filter sterilized

water and 36 μl AccuStart II PCR supermix (Quanta biosciences, Beverly, USA). To ensure that

any potential contamination either at the PCR stage or at the sequencing step would be

identified, two negative controls: water and non-template control were included at the PCR

step and sequenced as well as a positive spike included during the sequencing run. A fragment

size of ~460 bp of each sample was submitted to the WCAGC for sequencing together with the

negative controls. Automated cluster generation and paired-end sequencing (250-bp reads)

was performed on the Illumina MiSeq 500 cycle.

A total of 37 samples [6 infected midguts (INF MG); 4 non-infected midguts (NINF MG); 12

infected saliva (INF SAL); 6 non-infected saliva (NINF SAL); 3 infected salivary glands (INF SALG);

3 non- infected salivary gland (NINF SALG); 1 negative control (NC); 1 non-template control

(NTC) and 1 positive spike (PC) were included in this study.

Analysis of the data was carried out on QIITA (https://qiita.ucsd.edu/) and MicrobiomeAnalyst

[74]. In summary, to convert the de-multiplexed FASTQ files to the format used by QIITA, the

command split libraries FASTQ was used. The clustering of the sequences into operational

taxonomic units (OTUs), was done by the utilization of a closed-reference OTU picking based on

the GreenGenes 16S reference database which matches the sequences based on a 97%

sequence identity threshold before assigning a taxonomy. A BIOM-formatted OTU table was

subsequently generated, to reduce the potential of alpha and beta diversity biases, the data

was rarefied to a depth of 30 000 reads per sample. Finally, to visualize the taxonomic profiles

of each sample, taxa bar plots were generated using the rarefied data artifact. Following the

generation of the relative quantity of each taxon, we generated a taxa bar plot based on taxa

with sequences ≥ 4 000.

The reads have been deposited in NCBI GenBank Short Read Archives with accession numbers:

PRJNA608507, PRJNA608505, PRJNA608503, PRJNA608501, PRJNA608500, PRJNA608499,

PRJNA608472, PRJNA608496, PRJNA608495, PRJNA608493, PRJNA608492, PRJNA608491,

PRJNA608489, PRJNA608483, PRJNA608471, PRJNA608482, PRJNA608470, PRJNA608456,

PRJNA608450, PRJNA608481, PRJNA608449, PRJNA608448, PRJNA608480, PRJNA608479,

PRJNA608477, PRJNA608437, PRJNA608434, PRJNA608427, PRJNA608428, PRJNA608429,

PRJNA608375, PRJNA608426, PRJNA608424, PRJNA608416, PRJNA608418, PRJNA608422,

PRJNA608436 and PRJNA608419.

The alpha diversity profiling (the number of species of taxa in a single sample) was calculated

using the Chao1 diversity measure and significance testing was done using the ANOVA method

in Microbiome Analyst. Chao1 index estimates the richness of taxa in a sample by extrapolating

out the number of rare organisms that may have been omitted due to under sampling. The

index assumes that the number of observations for a taxon has a Poisson distribution and

hence corrects for variance [75].

Beta diversity, a comparison of microbial community composition between two samples or two

conditions, was analyzed by utilizing the Bray-Curtis dissimilarity statistic [76]. The distance

matrix generated after comparing each sample to the other was visualized for dissimilarity

between samples using the Principle Coordinate Analysis ordination method.

Clustering and Biomarker analysis

Patterns of relative abundance of taxa species in response to different experimental factors was

analyzed in Microbiome Analyst [74]. The distance measure was Euclidean [77] and the

clustering algorithm utilized was Ward, an agglomerative hierarchical clustering procedure [78].

The Euclidean distance, a measure of the true straight-line distance between two points in

Euclidean space, examines the root of square differences between coordinates of a pair of

objects.

To provide an estimate of the most important taxa for classification of data resulting from

different experimental factors, biomarker analysis was carried out using the random forest

algorithm within Microbiome Analyst [74]. The default setting of the number of trees to grow

and number of predictors to try was applied (500 and 7, respectively) with the randomness

setting left on. Random forest algorithm [79] is a supervised classification algorithm of trees

created by using bootstrap samples while training data and random feature selection in tree

induction. It is an ensemble of unpruned classification or regression trees trained with the

bagging method [80].

Quantification of *Elizabethkingia* and Zika virus

Both *in vitro* and *in vivo* analyses to quantify the *Elizabethkingia* loads were performed using

the primer pairs *rpoB* gene specific to *Elizabethkingia* using the PCR conditions described in

Table 1. The ribosomal protein *S7* gene was utilized to normalize transcript levels (Table 1).

Absolute quantity of ZIKV in individual midguts was obtained by isolating RNA using Trizol

(ThermoFisher scientific, Massachusetts, USA). A ZIKV-specific quantitative PCR assay that

targets the NS1 region was used for detection [73] and serially diluted ZIKV of known quantities

was used to generate the standard curve and extrapolate viral load.

Phylogenetic analysis of the *Elizabethkingia* isolated from *Ae. albopictus* salivary gland.

In order to validate the identity of *Elizabethkingia* taxa identified in the *Ae. albopictus* salivary

gland and saliva samples, *rpoB* gene (GenBank Accession number: MT096047) was amplified

from the salivary gland cDNA using the *rpoB* primer pairs and PCR conditions as described in

Table 1. The 558 bp PCR amplicon was excised from Agarose gel and cleaned using QIAquick gel

extraction kit (Qiagen, UK). The amplicons were Sanger sequenced at the WCAGC.

The sequences were manually trimmed using Bioedit v7.1.9 and aligned using MUSCLE [81,82].

Upon trimming the primer region and the low-quality regions of the sequence, the FASTA

sequence of the respective genes were blasted against the NCBI database. A total of 9 gene

sequences published with the following GenBank accession numbers: CP023010.2, KY587659.1,

CP016377.1, KY587657.1, CP035811.1, CP035809.1, CP011059.1, CP039929.1 and MN327643.1

were added to the phylogenetic analysis.

To construct the *Elizabethkingia rpoB* phylogenetic tree, Molecular Evolutionary Genetic

Analysis (MEGA) software [83] was utilized. The phylogenetic tree was constructed using the

Maximum Likelihood statistical method [84] with nucleotide substitution type.

The test of phylogeny was by bootstrap method [85] with 1000 bootstrap replications. The

nucleotide substitution model was the Tamura 3-parameter model [86]. The rates among sites

were uniform rates while the Heuristic method was the Nearest-Neighbor-Interchange.

***In vitro* assay of *Elizabethkingia* and ZIKV co-infection**

In order to study the growth kinetics of ZIKV co-infected with *Elizabethkingia*, U4.4 cells (*Ae.*

albopictus), were co-infected with *Elizabethkingia* (kindly provided by Steven Blaire, The

Connecticut Agricultural Experiment Station) and ZIKV.

Elizabethkingia bacterial strain was streaked on Luria-Bertani (LB) plates and incubated in a 10%

CO_2 incubator at 28°C overnight. A single colony was picked using a sterile plastic loop and used

to inoculate 500 ml of sterile LB broth. The inoculated broth media was incubated overnight at

28°C in an orbital shaker shaking at 225 rpm. The *Elizabethkingia* overnight culture was

subsequently diluted to OD 1.0 (8.5 $\log_{10}$ CFU/ml), OD 0.5 (8.0 $\log_{10}$ CFU/ml) and OD 0.2 (7.7

$\log_{10}$ CFU/ml).

Individual wells were seeded with 3 ml of 2 X 10^5 U4.4 cells/ml suspended in M&M medium

without FBS and incubated at 28°C for 1 h before adding FBS to each well at a final

concentration of 20%. The cells were then incubated for 5 days in a 10% CO_2 incubator at 28°C.

After 5 days, supernatant was removed carefully, and the plates were infected with 100 µl of

Elizabethkingia at OD 1.0 (8.5 $\log_{10}$ CFU/ml), OD 0.5 (8.0 $\log_{10}$ CFU/ml) and OD 0.2 (7.7 $\log_{10}$

CFU/ml) for three biological replicates. The negative control plates were infected with 100 µl of

8.3 $\log_{10}$ PFU/ml. After 1 h of incubation at 28°C, 3 ml of M&M media containing 20% fetal

bovine serum (FBS) were added to each well and incubated at 28°C.

To determine whether the U4.4 cells were successfully infected with *Elizabethkingia,* a single

well from each replicate of the different dilutions of *Elizabethkingia* infection was harvested at

2 dpi.

The cells were spun at 1000 rpm and washed three times with sterile phosphate buffered saline

(PBS). After the third wash, 100 µl of the washed cells were used to inoculate 3 ml of LB media

and incubated overnight at 28°C in an orbital shaker at 225 rpm.

The experimental wells were infected with 100 µl of 8.3 $\log_{10}$ PFU/ml ZIKV after a 2 day

incubation with *Elizabethkingia*. Cells which were co-infected with *Elizabethkingia* and ZIKV, as

well as the ZIKV only control, were harvested at both 2 dpi and 4 dpi. RNA was extracted from

the cells using Trizol (ThermoFisher scientific, Massachusetts, USA). The ZIKV quantities were

measured as previously described.

Statistical analysis

Statistical analysis was performed with GraphPad Prism version 5.0. Differences in

Elizabethkingia quantities were assessed using two-tailed T test. Viral loads were compared

using an ANOVA test. Finally, the *Elizabethkingia* /ZIKV correlation analysis was carried out

using Pearson tests.

Acknowledgements

This publication was supported by the Cooperative Agreement Number U01CK000509 funded

by the Center for Disease Control and Prevention. Its content is solely the responsibility of the

authors and do not necessarily represent the official views of the Center for Disease Control

and Prevention or the Department of Health and Human Services. These studies were also

partially funded by the National Institutes of Health award R21AI131683. We would like to

extend our gratitude to Illia Rochlin of Suffolk County Health Department for providing us with

Ae. albopictus mosquitoes and the Arbovirus laboratory insectary crew for mosquito

maintenance. We additionally thank the Wadsworth Center Applied Genomics core for

sequencing and the Wadsworth Center Tissue and Media facility for providing cells and media.

References

1.   Choi MH, Kim M, Jeong SJ, Choi JY, Lee I, Yong T, et al. Risk Factors for Elizabethkingia
Acquisition and Clinical Characteristics of Patients , South Korea. Emerg Infect Dis.
2019;25: 42–51.

2.   Vandamme P, Bernardet J-F, Segers P, Kersters K, Holmes B. New Perspectives in the
Classification of the Flavobacteria⬚: Description of Chryseobacterium gen. nov .,
Bergeyella gen. nov ., and Empedobacter nom. rev. Int J Syst Bacteriol. 1994;44: 827–
831.

3.   Kim KK, Kim MK, Lim JH, Park HY, Lee S. Transfer of Chryseobacterium meningosepticum
and Chryseobacterium miricola to Elizabethkingia gen . nov . as Elizabethkingia

meningoseptica. Int J Syst Evol Microbiol. 2005;55: 1287–1293. doi:10.1099/ijs.0.63541-0

4.   Lau SKP, Chow WN, Foo CH, Curreem SOT, Lo GCS, Teng JLL, et al. Elizabethkingia

anophelis bacteremia is associated with clinically significant infections and high

mortality. Sci Rep. 2016;6: 8–17. doi:10.1038/srep26045

5.   Janda JM, Lopez DL. Mini review: New pathogen profiles: Elizabethkingia anophelis.

Diagn Microbiol Infect Dis. 2017;88: 201–205. doi:10.1016/j.diagmicrobio.2017.03.007

6.   Bobossi-Serengbe G, Gody J, Beyam N, Bercion R. First documented case of

Chryseobacterium meningosepticum meningitis in Central African Republic. Med Trop.

2006;66: 182–184.

7.   Teo J, Tan SY-Y, Tay M, Ding Y, Kjelleberg S, Givskov M, et al. First case of E anophelis

outbreak in an intensive-care unit. Lancet. 2013;382: 855–856. doi:10.1016/S0140-

6736(13)61858-9

8.   Perrin A, Larsonneur E, Nicholson AC, Edwards DJ, Gundlach KM, Whitney AM, et al.

Evolutionary dynamics and genomic features of the Elizabethkingia anophelis 2015-2016

Wisconsin outbreak strain. Nat Commun. 2017;8: 1–12. doi:10.1038/ncomms15483

9.   Janda JM, Lopez DL. Mini review: New pathogen profiles: Elizabethkingia anophelis.

Diagn Microbiol Infect Dis. 2017;88: 201–205. doi:10.1016/j.diagmicrobio.2017.03.007

10.   Navon L, Clegg W, Morgan J, Austin C, McQuiston J, Blaney D, et al. Notes from the field:

Investigation of Elizabethkingia anophelis cluster - Illinois, 2014-2016. 2016.

11.   Lau SKP, Wu AKL, Teng JLL, Tse H, Curreem SOT, Tsui SKW, et al. Evidence for

Elizabethkingia Anophelis transmission from mother to infant, Hong Kong. Emerg Infect

Dis. 2015;21: 232–241. doi:10.3201/eid2102.140623

12.   Reed TAN, Watson G, Kheng C, Tan P, Roberts T, Ling CL, et al. Elizabethkingia anophelis

Infection in infants, Cambodia, 2012-2018. Emerg Infect Dis. 2020;26: 320–322.

13.   Chen S, Bagdasarian M, Walker ED. Elizabethkingia anophelis: Molecular manipulation

and interactions with mosquito hosts. Appl Environ Microbiol. 2015;81: 2233–2243.

doi:10.1128/AEM.03733-14

14.   Perrin A, Larsonneur E, Nicholson AC, Edwards DJ, Gundlach KM, Whitney AM, et al.

Evolutionary dynamics and genomic features of the Elizabethkingia anophelis 2015 to

524   2016 Winconsin outbreak strain. Nat Commun. 2017;8: 1–12.

525   doi:10.1038/ncomms15483

526 15. Terenius O, Lindh JM, Eriksson-Gonzales K, Bussière L, Laugen AT, Bergquist H, et al.

527   Midgut bacterial dynamics in Aedes aegypti. FEMS Microbiol Ecol. 2012;80: 556–565.

528   doi:10.1111/j.1574-6941.2012.01317.x

529 16. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, Chandra R, et al. Exploring

530   Anopheles gut bacteria for Plasmodium blocking activity. Environ Microbiol. 2015;16:

531   2980–2994. doi:10.1111/1462-2920.12381.Exploring

532 17. Akhouayri IG, Habtewold T, Christophides GK. Melanotic Pathology and Vertical

533   Transmission of the Gut Commensal Elizabethkingia meningoseptica in the Major Malaria

534   Vector Anopheles gambiae. PLoS One. 2013;8: 1–12. doi:10.1371/journal.pone.0077619

535 18. Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, et al. Midgut microbiota

536   of the malaria mosquito vector Anopheles gambiae and interactions with Plasmodium

537   falciparum infection. PLoS Pathog. 2012;8: 1–12. doi:10.1371/journal.ppat.1002742

538 19. Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J. Dynamic gut microbiome across life history

539   of the malaria mosquito anopheles gambiae in Kenya. PLoS One. 2011;6: 1–9.

540   doi:10.1371/journal.pone.0024767

541 20. Kämpfer P, Matthews H, Glaeser SP, Martin K, Lodders N, Faye I. Elizabethkingia

542   anophelis sp. nov., isolated from the midgut of the mosquito Anopheles gambiae. Int J

543   Syst Evol Microbiol. 2011;61: 2670–2675. doi:10.1099/ijs.0.026393-0

544 21. Kämpfer P, Busse HJ, McInroy JA, Glaeser SP. Elizabethkingia endophytica sp. nov.,

545   isolated from Zea mays and emended description of Elizabethkingia anophelis Kämpfer

546   et al. 2011. Int J Syst Evol Microbiol. 2015;65: 2187–2193. doi:10.1099/ijs.0.000236

547 22. Holmes B, Steigerwalt AG, Nicholson AC. DNA-DNA hybridization study of strains of

548   Chryseobacterium, Elizabethkingia and Empedobacter and of other usually indole-

549   producing non-fermenters of CDC groups IIc, IIe, IIh and IIi, mostly from human clinical

550   sources, and proposals of Chryseobacterium be. Int J Syst Evol Microbiol. 2013;63: 4639–

551   4662. doi:10.1099/ijs.0.054353-0

552 23. Li Y, Kawamura Y, Fujiwara N, Naka T, Hongsheng L, Huang X, et al. Chryseobacterium

miricola sp. nov., A novel species isolated from condensation water of space station Mir.

Syst Appl Microbiol. 2003;26: 523–528.

24. Nicholson AC, Gulvik CA, Whitney AM, Humrighouse BW, Graziano J, Emery B, et al.

Revisiting the taxonomy of the genus Elizabethkingia using whole-genome sequencing,

optical mapping, and MALDI-TOF, along with proposal of three novel Elizabethkingia

species: Elizabethkingia bruuniana sp. nov., Elizabethkingia ursingii sp. nov., and Eliz.

Antonie van Leeuwenhoek, Int J Gen Mol Microbiol. 2018;111: 55–72.

doi:10.1007/s10482-017-0926-3

25. King E. Studies on a group of previously unclassified bacteria associated with meningitis

in infants. Am J Clin Pathol. 1959;31: 241–247.

26. Ka P, Matthews H, Glaeser SP, Martin K, Lodders N, Faye I. Elizabethkingia anophelis sp .

nov ., isolated from the midgut of the mosquito Anopheles gambiae. 2019; 2670–2675.

doi:10.1099/ijs.0.026393-0

27. Mancini M V, Damiani C, Accoti A, Tallarita M, Nunzi E, Cappelli A, et al. Estimating

bacteria diversity in different organs of nine species of mosquito by next generation

sequencing. BMC Microbiol. 2018;18: 1–10.

28. Lombardo F, Capurro ML. Salivary gland-specific gene expression in the malaria vector

Anopheles gambiae. 1999.

29. Arca B, Lombardo F, Lanfrancotti A, Spanos L, Veneri M, Louis C, et al. A cluster of four

D7-related genes is expressed in the salivary glands of the African malaria vector

Anopheles gambiae. Insect Mol Biol. 2002;11: 47–55.

30. Kerlin R., Hughes S. Enzymes in saliva from four parasitic arthropods. Med Vet Entomol.

1992;6: 121–126.

31. Mellink JJ, Vos BJ. Primary Lymph Nodes Response to Mosquito Bites. Zeitschrift fur

Parasitenkd Parasitol Res. 1977;51: 187–188.

32. Ribeiro J. Characterization of a vasodilator from the salivary glands of the yellow fever

mosquito Aedes aegypti. J exp Biol. 1992;165: 61–71.

33. Schneider B., Soong L, Zeidner N., Higgs S. Aedes aegypti Salivary Gland Extracts

Modulate Anti-Viral and TH1/TH2 Cytokine Responses to Sindbis Virus Infection. Viral

Immunol. 2004;17: 565–573.

34.   Titus RG, Bishop J V, Mejia JS. The immunomodulatory factors of arthropod saliva and

the potential for these factors to serve as vaccine targets to prevent pathogen

transmission. 2006; 131–141. doi:10.1111/j.1365-3024.2006.00807.x

35.   Conway MJ, Londono-renteria B, Troupin A, Watson AM, Klimstra WB, Fikrig E, et al.

Aedes aegypti D7 Saliva Protein Inhibits Dengue Virus Infection. PLoS Negl Trop Dis.

2016; 1–19. doi:10.1371/journal.pntd.0004941

36.   Conway MJ, Watson AM, Colpitts TM, Dragovic SM, Li Z, Wang P, et al. Mosquito Saliva

Serine Protease Enhances Dissemination of Dengue. J Virol. 2014;88: 164–175.

doi:10.1128/JVI.02235-13

37.   Jin L, Guo X, Shen C, Hao X, Sun P, Li P, et al. Salivary factor LTRIN from Aedes aegypti

facilitates the transmission of Zika virus by interfering with the lymphotoxin-β receptor.

Nat Immunol. 2018;19: 342–353. doi:10.1038/s41590-018-0063-9

38.   Uraki R, Hastings AK, Marin-lopez A, Sumida T, Takahashi T, Grover JR, et al. Aedes

aegypti AgBR1 antibodies modulate early Zika virus infection of mice. Nat Microbiol.

doi:10.1038/s41564-019-0385-x

39.   Rubinstien E, Klevjer-Anderson P, Smith C, Drouin M, Patterson J. Enterobacter taylorae,

a New Opportunistic Pathogen: Report of Four Cases. J Clin Microbiol. 1993;31: 249–254.

40.   Hurrell E, Kucerova E, Loughlin M, Caubilla-barron J, Hilton A, Armstrong R, et al.

Neonatal enteral feeding tubes as loci for colonisation by members of the

Enterobacteriaceae. BMC Infect Dis. 2009;9: 1–9. doi:10.1186/1471-2334-9-146

41.   Guentzel M. Escherichia, Klebsiella, Enterobacter, Serratia, Citrobacter and Proteus. 4th

ed. In: Samuel Barron, editor. Medical Microbiology. 4th ed. Galveston, Texas: University

of Texas Medical Branch at Galveston; 1996.

42.   Gaio ADO, Gusmão DS, Santos A V, Berbert-molina MA, Pimenta PFP, Lemos FJA.

Contribution of midgut bacteria to blood digestion and egg production in Aedes aegypti (

diptera: culicidae) (L .). 2011; 1–10.

43.   Simi S, Carbonell G, Falcon R, Gatti M, Joazeiro P, Darini A, et al. A low molecular weight

enterotoxic hemolysin from clinical Enterobacter cloacae. Can J Microbiol. 2003;49: 479–

611  482.

612 44. Serbus LR, Sullivan W. A Cellular Basis for Wolbachia Recruitment to the Host Germline.

613   PLoS Pathog. 2007;3: 1930–1937. doi:10.1371/journal.ppat.0030190

614 45. Werren JH, Baldo L, Clark ME. Wolbachia⬚: master manipulators of invertebrate biology.

615   Nat Rev Microbiol. 2008;6: 741–751. doi:10.1038/nrmicro1969

616 46. White PM, Pietri JE, Debec A, Russell S, Patel B, Sullivan W. Mechanisms of Horizontal

617   Cell-to-cell Transfer of Wolbachia spp. in Drosophila melanongaster. Appl Environ

618   Microbiol. 2017;83: 1–12.

619 47. Li S, Ahmed MZ, Lv N, Shi P, Wang X, Ji-Lei H, et al. Plant – mediated horizontal

620   transmission of Wolbachia between whiteflies. ISME J. 2017;11: 1019–1028.

621   doi:10.1038/ismej.2016.164

622 48. Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, Moreira LA, et al. Harnessing mosquito-

623   Wolbachia symbiosis for vector and disease control. Acta Trop. 2014;132: S150–S163.

624   doi:10.1016/j.actatropica.2013.11.004

625 49. Dickson LB, Jiolle D, Minard G, Moltini-Conclois I, Volant S, Ghozlane A, et al. Carryover

626   effects of larval exposure to different environmental bacteria drive adult trait variation in

627   a mosquito vector. Sci Adv. 2017;3: 1–15. doi:10.1126/sciadv.1700585

628 50. Muturi EJ, J BJ, Rooney AP, K. HA. Midgut fungal and bacterial microbiota of Aedes

629   triseriatus and Aedes japonicus shift in response to La Crosse virus infection. Mol Ecol.

630   2016;25: 4075–4090. doi:10.1111/mec.13741

631 51. Zink SD, van Slyke GA, Palumbo MJ, Kramer LD, Ciota AT. Exposure to west nile virus

632   increases bacterial diversity and immune gene expression in culex pipiens. Viruses.

633   2015;7: 5619–5631. doi:10.3390/v7102886

634 52. Zouache K, Michelland RJ, Failloux AB, Grundmann GL, Mavingui P. Chikungunya virus

635   impacts the diversity of symbiotic bacteria in mosquito vector. Mol Ecol. 2012;21: 2297–

636   2309. doi:10.1111/j.1365-294X.2012.05526.x

637 53. Ramirez JL, Short SM, Bahia AC, Saraiva RG, Dong Y, Kang S, et al. Chromobacterium

638   Csp_P Reduces Malaria and Dengue Infection in Vector Mosquitoes and Has

639   Entomopathogenic and In Vitro Anti-pathogen Activities. PLoS Pathog. 2014;10.

doi:10.1371/journal.ppat.1004398

54.   Apte-Deshpande A, Paingankar M, Gokhale MD, Deobagkar DN. Serratia odorifera a

midgut inhabitant of aedes aegypti mosquito enhances its susceptibility to dengue-2

virus. PLoS One. 2012;7. doi:10.1371/journal.pone.0040401

55.   Weiss B, Aksoy S. Microbiome influences on insect host vector competence. Trends

Parasitol. 2011;27: 514–522. doi:10.1016/j.pt.2011.05.001

56.   Li N, Ma W, Pang M, Fan Q, Hua J. The Commensal Microbiota and Viral Infection : A

Comprehensive Review. Front Immunol. 2019;10: 1–16. doi:10.3389/fimmu.2019.01551

57.   Karst SM. The influence of commensal bacteria on infection with enteric viruses. Nat Rev

Microbiol. 2016;14: 197–204. doi:10.1038/nrmicro.2015.25.The

58.   Berger AK, Mainou BA. Interactions between Enteric Bacteria and Eukaryotic Viruses

Impact the Outcome of Infection. Viruses. 2018;10: 4–6. doi:10.3390/v10010019

59.   Pfeiffer JK, Virgin HW. Transkingdom control of viral infection and immunity in the

mammalian intestine. Science (80-). 2017;351: 1–12.

doi:10.1126/science.aad5872.Transkingdom

60.   Abt MC, Osborne LC, Monticelli LA, Doering TA, Alenghat T, Sonnenberg GF, et al.

Commensal Bacteria Calibrate the Activation Threshold of Innate Antiviral Immunity. Cell.

2012;37: 158–170. doi:10.1016/j.immuni.2012.04.011

61.   Kumar A, Vlasova AN, Deblais L, Huang H, Wijeratne A, Kandasamy S, et al. Impact of

nutrition and rotavirus infection on the infant gut microbiota in a humanized pig model.

BMC Gastroenterol. 2018;18: 1–17.

62.   Jang J, Kim S, Kwon M, Lee J, Yu D, Song R, et al. Rotavirus-mediated alteration of gut

microbiota and its correlation with physiological characteristics in neonatal calves. J

Microbiol. 2019;57: 113–121. doi:10.1007/s12275-019-8549-1

63.   Ma X, Wang Q, Li H, Xu C, Cui N, Zhao X. 16S rRNA genes Illumina sequencing revealed

differential cecal microbiome in specific pathogen free chickens infected with diff erent

subgroup of avian leukosis viruses. Vet Microbiol. 2017;207: 195–204.

doi:10.1016/j.vetmic.2017.05.016

64.   Zhao N, Li M, Luo J, Wang S, Liu S, Wang S, et al. Impacts of canine distemper virus

669  infection on the giant panda population from the perspective of gut microbiota. Sci Rep.

670  2017; 1–10. doi:10.1038/srep39954

671 65. Ding ZF, Cao MJ, Zhu XS, Xu GH, Wang RL. Changes in the gut microbiome of the Chinese

672  mitten crab (Eriocheir sinensis) in response to White spot syndrome virus (WSSV)

673  infection. J Fish Dis. 2017;40: 1561–1571. doi:10.1111/jfd.12624

674 66. Mayuzumi H, Okamoto Y. Interleukin- 17A is required to suppress invasion of Salmonella

675  enterica serovar Typhimurium to enteric mucosa. 2010; 377–385. doi:10.1111/j.1365-

676  2567.2010.03310.x

677 67. Deriu E, Boxx GM, He X, Pan C, Benavidez SD, Cen L, et al. Influenza Virus Affects

678  Intestinal Microbiota and Secondary Salmonella Infection in the Gut through Type I

679  Interferons. PLoS Pathog. 2016; 1–26. doi:10.1371/journal.ppat.1005572

680 68. Rossi P, Ricci I, Cappelli A, Damiani C, Ulissi U, Mancini MV, et al. Mutual exclusion of

681  Asaia and Wolbachia in the reproductive organs of mosquito vectors. Parasites and

682  Vectors. 2015;8: 1–10. doi:10.1186/s13071-015-0888-0

683 69. Ngwa C, Glockner V, Abdelmohsen U, Scheuermayer M, Fischer R, Hentschel U, et al. 16S

684  rRNA gene-based identification of Elizabethkingia meningoseptica

685  (Flavobacteriales:Flavobacteriaceae) as a dominant midgut bacterium of the Asian

686  malaria vector Anopheles stephensi (Diptera: Culicidae) with antimicrobial activities. J

687  Med Entomol. 2013;50: 404–14.

688 70. Villegas LEM, Campolina TB, Barnabe NR, Orfano AS, Chaves BA, Norris DE, et al. Zika

689  virus infection modulates the bacterial diversity associated with Aedes aegypti as

690  revealed by metagenomic analysis. PLoS One. 2018;13: e0190352.

691  doi:10.1371/journal.pone.0190352

692 71. Almand E, Moore M, Jaykus L. Virus-Bacteria Interactions : An Emerging Topic in Human

693  Infection. Viruses. 2017;9: 1–10. doi:10.3390/v9030058

694 72. Ciota A, Bialosuknia S, Zink S, Brecher M, Ehrbar D, Morrissette M, et al. Effects of Zika

695  Virus Strain and Aedes Mosquito Species on Vector Competence. Emerg Infect Dis.

696  2017;23: 1110–1117.

697 73. Lanciotti RS, Calisher CH, Gubler DJ, Chang GJ, Vorndam a V, Vorndamt a V. Rapid

698   detection and typing of dengue viruses from clinical samples by using reverse

699   transcriptase-polymerase chain reaction. J Clin Microbiol. 1992;30: 545–51.

700   74.   Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. MicrobiomeAnalyst: A web-based

701   tool for comprehensive statistical, visual and meta-analysis of microbiome data. Nucleic

702   Acids Res. 2017;45: W180–W188. doi:10.1093/nar/gkx295

703   75.   Chao A. Nonparametric Estimation the Number of Classes in a Population. Ann Math

704   Stat. 1984;20: 265–270. doi:10.1214/aoms/1177729949

705   76.   Bray RJ, Curtis JT. An Ordination of the Upland Forest Communities of Southern

706   Wisconsin. Ecol Monogr. 1957;27: 325–349.

707   77.   Motzkin T. The euclidean algorithm. Bulletin of the American Society. 1949. pp. 1142–

708   1146.

709   78.   Ward J, Hook M. Application of an hierarchical grouping procedure to a problem of

710   grouping profiles. Educ Psychol Meas. 1963;XXIII.

711   79.   Svetnik V, Liaw A, Tong C, Christopher Culberson J, Sheridan RP, Feuston BP. Random

712   Forest: A Classification and Regression Tool for Compound Classification and QSAR

713   Modeling. J Chem Inf Comput Sci. 2003;43: 1947–1958. doi:10.1021/ci034160g

714   80.   Liaw A, Wiener M. Classification and Regression by randomForest. R news. 2002;2: 18–

715   22. doi:10.1023/A:1010933404324

716   81.   Hall T. BioEdit: An important software for molecular biology. GERF Bull Biosci. 2011;2:

717   60–61.

718   82.   Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high

719   throughput. Nucleic Acids Res. 2004;32: 1792–7. doi:10.1093/nar/gkh340

720   83.   Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular

721   evolutionary genetics analysis using maximum likelihood, evolutionary distance, and

722   maximum parsimony methods. Mol Biol Evol. 2011;28: 2731–9.

723   doi:10.1093/molbev/msr121

724   84.   Felsenstein J. Evolutionary Trees from DNA Sequences⍰: A Maximum Likelihood

725   Approach. J Mol Evol. 1981;17: 368–376.

726   85.   Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap.

727   Evolution (N Y). 1985;39: 783–791.

728   86.   Tamura K, Nei M. Estimation of the Number of Nucleotide Substitutions in the Control

729   Region of Mitochondrial DNA in Humans and Chimpanzees. Mol Biol Evol. 1993;10: 512–

730   526.

Author Contributions

MGO: Involved in the study design, conducted the experiments, analyzed the results and

drafted the paper

PA: Trained me in the cell culture, was involved in the antibiotic treatment of the Aedes

mosquitoes and assisted in mosquito manipulation exercises

SJ: Involved in the mosquito rearing, assisted in mosquito manipulation exercises

DC: Assisted in mosquito manipulation exercises

KL: Generated and propagated the ZIKV infectious clone utilized in this study

KLD: Involved in the study design, participated in the writing of the paper

CAT: Involved in the study design, participated in the writing of the paper

Competing Interests

No competing interests

Materials and correspondence

Correspondence and material requests should be addressed to CAT

Tables and figures

List of tables

Table 1. List of Primers used in this study

Primer name	Primer sequence	Annealing temperature
Ribosomal protein S7 _F	GAG ATC GAG TTC AAC AGC AAG A	
Ribosomal protein S7 _R	GAG AAC TTC TTC TCC AGC TCA C	55°C
*Elizabethkingia*_rpoB_F	CTC CGG AAG GAC CAA ACA TTG	
*Elizabethkingia*_rpoB_R	CAA CCG TCC AGT CAG ATC C	55°C
*Elizabethkingia*_qrpoB_F	GCT AGG ACA CAT GGT TGA TGA	
*Elizabethkingia*_qrpoB_R	ACC ACC GAA CTG AGC TTT AC	55°C
16S Microbiome_F	TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CCT ACG GGN GGC WGC AG	
16S Microbiome_R	GTC TCG TGG GCT CGG AGA TCT GTA TAA GAG ACA GGA CTA CHV GGG TAT CTA ATC C	55°C

Figures

Figure 1. Relative abundance of bacterial taxa in individual *Aedes* saliva, salivary gland and midgut samples. Illumina MiSeq 16S rRNA sequencing was used to identify bacterial taxa and distributions were visualized using a cut-off of 7000 reads. Each stacked bar plot illustrates the taxonomic distribution of bacteria identified in individual *Aedes* saliva, salivary gland or midgut samples. Samples were taken at 7 days post-feeding. Blood fed individuals were further separated into NINF (fed non-infectious blood meal [BM]) and INF (fed Zika virus-positive BM and acquired an infection). SAL_INF represents saliva samples obtained from individuals fed on INF BM and successfully transmitting the virus; SAL_NINF represent saliva samples obtained from individuals exposed to NINF BM; SALG_INF represent individual salivary gland samples exposed to INF BM and infected; SALG_NINF represent salivary glands from individuals fed on NINF BM; GUT_INF represent midgut samples exposed to INF BM and infected, and GUT_NINF represent midgut samples from individuals fed on NINF BM.

Figure 2. Tissue-specific diversity of microbes in *Ae. albopictus*. Principal component analyses were completed to assess the genetic relationships among and between bacterial taxa identified in different tissue types. Despite a number of shared individual bacterial genera, significant separation by tissue type was identified [PERMANOVA; F-value:6.9998; R-squared: 0.3111; P <0.001].

Figure 3. Association of bacterial taxa with both tissue and Zika virus infection status in *Ae. albopictus* mosquitoes. The Random forest classifier was used to determine the association of differentially abundant microbial genera with both tissue-type and Zika virus infection status of *Aedes* mosquitoes. Mean Decrease Accuracy is reported for each of the taxa. This measure is obtained by removing the relationship of a taxa and measuring increase in error. The taxa with highest mean decrease in accuracy is considered to have the highest association with the state. The highest mean decrease in accuracy was identified for *Elizabethkingia*. In addition, *Elizabethkingia* and *Enterobacteriaceae*, taxa associated with human disease, were found to be present in the midguts, saliva and salivary glands of *Aedes* mosquitoes exposed to both infectious (INF) and noninfectious (NINF) blood meals (BM).

Figure 4. Relative *Elizabethkingia* levels in *Ae. albopictus.* A. **Tissue tropism.** Relative

quantification of *Elizabethkingia* in the midgut, salivary gland and saliva of *Ae. albopictus*. Levels

were tissue-specific and dependent on Zika virus (ZIKV) exposure. GUT_INF- infected gut;

SAL_INF - infected saliva; SALG_INF - infected salivary gland; GUT_NINF – gut samples of *Ae.*

albopictus exposed to NINF BM; SAL_NINF – saliva samples from *Ae. albopictus* exposed to NINF

BM; SALG_NINF – salivary gland samples of *Ae. albopictus* exposed to NINF BM. **B. Relative**

levels of *Elizabethkingia* in Zika virus infected and uninfected *Ae. albopictus* midguts. Infection

of *Ae. albopictus* midguts by ZIKV (INF) was associated with higher *Elizabethkingia rpoB*

transcript levels (T-test; Mann Whitney test P = 0.0051) relative to midguts following non-

infectious blood meals (NINF BM) at 7 days post-feeding.

Figure 5. Relationship between *Elizabethkingia* and Zika virus proliferation. A. *In vivo*

comparison in *Ae. albopictus* midguts. *Aedes albopictus* were infected with a bloodmeal

containing 8.3 $\log_{10}$ PFU/ml ZIKV and of both expression of *Elizabethkingia rpoB* gene and ZIKV

levels were assessed at 7 days post infection. The RT qPCR results were normalized using S7

gene. The X axis represents the log fold change of *Elizabethkingia*, while Y axis represents $\log_{10}$

ZIKV PFU/midgut. A negative correlation between ZIKV and *Elizabethkingia* was demonstrated

(Spearman r = -0.3985; P < 0.05). **B. *In-vitro* assay of *Elizabethkingia* and Zika virus in U4.4 cells**

U4.4 cells were infected with 8.3 $\log_{10}$ PFU/ml ZIKV following 2 days exposure to *Elizabethkingia*

(EBK, experimental) or media alone (control). Experimental wells were previously exposed to

either OD 1.0 (8.5 $\log_{10}$ CFU/ml EBK), OD 0.5 (8.0 $\log_{10}$ CFU/ml EBK) or OD 0.2 (7.7 $\log_{10}$ CFU/ml

EBK). At both 2 and 4 days post infection (dpi) ZIKV levels were significantly higher in the ZIKV

only controls relative to the EBK co-infected cells (ZIKV + *Elizabethkingia*; ANOVA; Kruskal-Wallis test, **P < 0.01).

Figure 6. **Phylogenetic analysis of *Elizabethkingia rpoB* gene.** Phylogenetic affiliation of *Elizabethkingia* taxa from the salivary gland of *Ae. albopictus* based on a partial sequence of the *rpoB* gene (GenBank Accession number: MT096047). Phylogenetic dendrograms were constructed in MEGA. Bootstrap proportions are shown on branches. *Chryseobacterium* was used as an outgroup.

Figure 1.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6.