Zika virus infection and temperature significantly impact *Aedes albopictus* gut microbiota

Onyango MG[1], Bialosuknia S[1,2], Payne A[1], Mathias N[1], Kuo L[1], Kramer LD[1,2], Ciota AT[1, 2§]

[1] New York State Department of Health, Griffin Laboratory, 5668 State Farm Road, Slingerlands,

12159, New York

[2] School of Public Health, State University of New York Albany, 1400 Washington Avenue,

Albany, 12222, New York

Email addresses:

MGO: Maria.Onyango@health.ny.gov

BS: Sean.Bialosuknia@health.ny.gov

PA: anne.payne@health.ny.gov

MN: Nicholas.Mathias@health.ny.gov

KL: lili.kuo@health.ny.gov

CA: alexander.ciota@health.ny.gov

KL: laura.kramer@health.ny.gov

[§]Corresponding author

Abstract

Background: Vector-borne pathogens must survive and replicate in the hostile environment in

the insect gut before successful transmission. The insect gut bacteria can have a modulatory

effect on their vectorial capacity. The relationship between infection, temperature and the

mosquito gut microbiome are not well characterized.

Methodology/ Principal findings: This study aimed to delineate the effect of Zika virus (ZIKV)

infection and temperature variation on the microbial profile of *Aedes albopictus* mosquitoes.

Ae. albopictus were reared at diurnal temperature of day 28°C/ night 24°C and day 30°C/ night

26°C. Mosquitoes were given infectious blood meals with 8.3 $\log_{10}$ PFU/ml ZIKV and 16S *rRNA*

sequencing was performed on guts at 7 days post-infectious bloodmeal exposure.

Blood meal acquisition and ZIKV exposure significantly reduced the gut taxa diversity while

increase in temperature significantly altered the bacterial composition. A number of individual

bacterial genera, most notably *Elizabethkingia* and *Neoasaia*, were found to be associated with

blood meal digestion, ZIKV exposure and susceptibility to infection. To the best of our

knowledge, we identified *Neoasaia* for the first time in the midgut of an arthropod vector.

The findings of this study demonstrate that both temperature and ZIKV infection modulate the

midgut microbial profile of *Ae. albopictus* mosquitoes. This may impact arbovirus transmission

dynamics in future climate change scenarios and inform novel control strategies.

Key words: Zika virus, *Aedes albopictus*, microbiome, *Elizabethkingia*, *Neoasaia*

Introduction

The microbiota of mosquitoes has been shown to interfere with mosquito pathogen infection and influence mosquito fitness, hence affecting their vectorial capacity (Romoli & Gendrin, 2018). The importance of the microbiota in the modulation of vector competence due to complex interactions between the pathogen, the mosquito and its bacterial microbiota has become an area of interest and target for development of novel biological control tools (Guégan et al., 2018). Abundant communities of diverse microorganisms reside in insects with a major concentration in the intestinal sections (Dillon & Dillon, 2004).

The major bottleneck of pathogen transmission by the mosquito is pathogen invasion of the midgut epithelium (Romoli & Gendrin, 2018) and the gut microbiota may modulate pathogen susceptibility and replication by activation of its basal immunity (Vallet-Gely et al., 2009). Studies investigating the interaction of gut microbial taxa and arboviruses of medical importance have mainly been limited to *Wolbachia* (Asad et al., 2018; Lindsey et al., 2019; Mohanty et al., 2018; Moretti et al., 2018). *Wolbachia* has subsequently been utilised for large scale control of dengue virus (DENV) and Zika virus (ZIKV) (Callaway, 2016; Nguyen et al., 2015). Despite the effectiveness of *Wolbachia*, it is important to further explore other microbes that could potentially be used as novel biological markers and tools. Our study provides insights into the interactions of ZIKV with the resident gut microbiota of *Ae. albopictus*, considered as a secondary vector of DENV and ZIKV.

In addition, the relationship between increases in temperature and gut microbiota of mosquitoes is not well-characterized. This information is important to understand the potential impact of future climate change on mosquito-pathogen interactions. This study sought to delineate the impact of ZIKV infection and temperature shifts on the gut microbiota of *Ae.*

albopictus. The identification of symbionts that are associated with altered vector competence

or altered temperatures can potentially act as targets for informing novel vector control

measures.

Materials and Methods

Mosquito sampling and DNA preparation

Ae. albopictus (ALB LI) were collected from Suffolk County, Long island, NY in 2015 (kindly

provided by I. Rochlin) (Figure 1). Field-collected mosquitoes were colonized at the New York

State Department of Health (NYSDOH) Arbovirus Laboratory while being maintained at 27°C

under standard rearing conditions.

Mosquitoes were hatched and reared under two thermal conditions; 14 hours light at 30°C: 10

hours dark at 26°C (M temperature regimen) and 14 hours light at 28°C: 10 hours dark at 24°C

(L temperature regimen). The L temperature regime modeled current average temperatures

during peak transmission in New York State and the M temperature regime modeled a

predicted 2°C increase, as has been suggested will occur with climate change (Kjellstrom et al.,

2018).

To obtain mosquito midgut samples for microbiome sequencing of ZIKV - infected guts, the F15

ALB LI eggs were vacuum-hatched. Before hatching, the water was incubated for 3 h at the two

different day time temperatures. The immature stages of the mosquitoes were reared at the

two diurnal temperature regimes. At eight days post mating, the female adults were orally

exposed to blood meals containing 8.3 $\log_{10}$ PFU/ml ZIKV. Subsets of the same population of

mosquitoes were either not offered blood meals (UNFED) or offered non-infectious blood

meals (UNEXP).

A total of 15 pools (three mosquitoes/pool) of *Ae. albopictus* female mosquito midguts

collected 7 days post infectious (dpi) blood meal were stored at -80°C until total RNA and

genomic DNA extraction were carried out using the Trizol protocol (ThermoFisher scientific,

Massachusetts, USA). A ZIKV-specific quantitative PCR was utilised to screen the guts for ZIKV

infection as described by Ciota A.T et al., 2017. The ZIKV strain utilised in this study was HND

(2016-19563, GenBank accession no. KX906952). In addition to ZIKV exposed midguts (EXP),

midguts from UNF and UNEXP midguts were collected at equivalent timepoints for comparison.

Sequencing of Microbiome of *Aedes* mosquito guts and Analysis

To determine the gut microbial profile of ZIKV-exposed individuals reared at the two

temperature regimes, barcoded high-throughput amplicon sequencing of the bacterial 16S V3-

V4 hypervariable *rRNA* was done. PCR reactions were carried out in a total volume of 50 µl,

5µM of 16S *rRNA* V3-V4 hypervariable region primers (Table 1), 10µl genomic DNA and 36 µl

AccuStart II PCR supermix (Quanta biosciences, Beverly, USA). A fragment size of ~460 bp of

each sample was submitted to the Wadsworth Center Applied Genomics Core for sequencing.

Automated cluster generation and paired-end sequencing (250-bp reads) was performed on the

Illumina MiSeq system.

Analysis of the data was carried out on QIITA (https://qiita.ucsd.edu/) and MicrobiomeAnalyst

(Dhariwal et al., 2017). In summary, to convert the de-multiplexed FASTQ files to the format

used by QIITA, the command split libraries FASTQ was used, the samples were then trimmed to

100 bp size to allow for meta-analysis. The clustering of the sequences into operational

taxonomic units (OTUs), was done by the utilization of a closed-reference OTU picking based on

the GreenGenes 16S reference database which matches the sequences based on a 97%

sequence identity threshold before assigning a taxonomy. A BIOM-formatted OTU table was

subsequently generated, to reduce the potential of alpha and beta diversity biases, the OTU

data was rarefied to a sample depth of 44000 OTUs per sample. Finally, to visualize the

taxonomic profiles of each sample, taxa bar plots were generated using the rarefied data

artifact.

The alpha diversity profiling (the number of species of taxa in a single sample) was calculated

using the Chao1 diversity measure and significance testing was done using the T-test/ANOVA

method in Microbiome Analyst. Chao1 index estimates the richness of taxa in a sample by

extrapolating out the number of rare organisms that may have been omitted due to under

sampling. The index assumes that the number of observations for a taxon has a Poisson

distribution and hence corrects for variance (Chao, 1984). Shannon diversity index was also

utilised to calculate the alpha diversity (Shannon, 1964). This diversity index accounts for both

abundance and evenness of the species present. The proportion of species i relative to the total

number of species (p_i) is calculated, this is then multiplied by the natural logarithm of the

proportion (lnp_i).

Beta diversity, a comparison of microbial community composition between two samples, was

analyzed by utilizing the Bray-Curtis dissimilarity statistic (Bray & Curtis, 1957). The distance

matrix generated after comparing each sample to the other was visualized for dissimilarity

between samples using the Principle Coordinate Analysis ordination method.

Clustering and Biomarker analysis

Patterns of relative abundance of taxa species in response to different experimental factors was

analyzed in Microbiome Analyst (Dhariwal et al., 2017). The distance measure was Euclidean

(Motzkin, 1949) and the clustering algorithm utilised was Ward, an agglomerative hierarchical

clustering procedure (Ward & Hook, 1963). The Euclidean distance, a measure of the true

straight-line distance between two points in Euclidean space, examines the root of square

differences between coordinates of a pair of objects.

To provide an estimate of the most important taxa for classification of data resulting from

different experimental factors, biomarker analysis was carried out using the random forest

algorithm within Microbiome Analyst (Dhariwal et al., 2017). The default setting of the number

of trees to grow and number of predictors to try was applied (500 and 7 respectively) with the

randomness setting left on. Random forest algorithm (Svetnik et al., 2003) is a supervised

classification algorithm of trees created by using bootstrap samples while training data and

random feature selection in tree induction. It is an ensemble of unpruned classification or

regression trees trained with the bagging method (Liaw & Wiener, 2002).

Phylogenetic analysis of the *Neoasaia* and *Elizabethkingia rpoB* genes

The *Neoasaia rpoB* (502 bp) and *Elizabethkingia rpoB* (588 bp) genes were sequenced from PCR

products excised from agarose gel and cleaned using QIAquick gel extraction kit (Qiagen, UK) at

the Wadsworth Center Applied Genomics Core. The sequences were manually edited using

Bioedit v7.1.9 and aligned using MUSCLE (Edgar, 2004; Hall, 2011). Upon trimming the primer

region and the low-quality regions of the sequence, the FASTA sequence of the respective

genes were blasted against the NCBI database. A total of 11 *rpoB* gene sequences published

with the following GenBank accession numbers: KY587657.1; FJ999737.1; KY587659.1;

KY587658.1; FJ999739.1; FN391773.1; FN391805.1; KF537505.1; FN391750.1; FN391807.1,

HG329623.1 and NC_00913.3 were added to the phylogenetic analysis.

Bayesian analysis was performed on the sequences using MR. BAYES v3.1.2 (Ronquist et al.,

2012). The analysis was run for 1000000 generations using the general time reversible (GTR)

substitution model, sampling for convergence every 5 000 generations.

Results

Gut taxa microbial profile of *Aedes* mosquitoes

Next generation sequencing resulted in a total of 7419965 reads with a maximum read length

of 301 bp and minimum read length of 280 bp. After trimming the reads to 100 bp to allow

efficient comparison, a total of 6062 OTUs were derived from a closed reference OTU picking

with a frequency of 6036630 across the 34 individuals ranging between 44255 and 583632. The

OTU were rarefied to 44000 to allow downstream analysis.

The OTU generated in this study belonged to a total of 35 phyla; Bacteroidetes (0.58),

Proteobacteria (0.36), Actinobacteria (0.03), Firmicutes (0.02), Cyanobacteria (0.01) and OD1

(0.004) were the dominant taxa while the rest of the phyla had a frequency of < 0.004 (Figure

1). To generate a taxa bar plot to visualize the distribution of bacterial genera across individual

guts in each group, a cut-off taxa frequency of 150 was applied and a taxa bar plot generated

(Figure 2).

Blood feeding alters *Ae. albopictus* gut microbiota diversity and composition

Blood feeding significantly altered the gut taxa abundance of *Ae. albopictus* regardless of

grouping ([PERMANOVA] F-value: 15.686, R-squared, 0.18095, P<0.001; Figure 2) and the

overall alpha diversity decreased independent of ZIKV exposure or infection (chao1 T-

test/ANOVA P(0.006) [T-test] statistic -3.1216; Figure 3A). Although less apparent at the higher

temperature regime (M), in all groups the decrease in diversity was driven primarily by an

increase the relative abundance of the genus *Elizabethkingia* (Figure 2). The Random forest

classifier, used to determine the association of differentially abundant taxa in the microbial

community with certain states, confirmed that low relative abundance of *Elizabethkingia* was

strongly associated with UNFED individuals (Figure 4). Numerous taxa decreased or were

eliminated as a result of blood feeding. Most notable were *Acidovorax, Bdellovirio* and

Pseudomonas (Figures 2 and 4).

ZIKV exposure alters *Ae. albopictus* gut microbiota diversity and composition.

Comparisons of pools of midguts from ZIKV exposed individuals (FED EXP) to those fed non-

infectious blood meals (FED UNEXP) demonstrates that ZIKV exposure independently decreases

alpha diversity at 7 days post-feeding (chao1 T-test/ANOVA P (0.005) [T-test] statistic -3.1084;

Figure 3B). These differences in diversity at the group level were also visible at the individual

sample level (Figure 2).   Taxa for which relative abundance was decreased as a result of ZIKV

exposure included *Delftia, Chryseobacterium, Pseudomonas, Moraxellaceae,* and *Acidovorax*

(Figures 2 and 4).

There was no significant difference in taxa richness within and between pools that

disseminated the virus and those that were uninfected following exposure (EXP UNINF v. EXP

DIS; Alpha diversity Chao1 Index P[7.574e-05(ANOVA) F-value: 10.379]; Beta diversity

[PERMANOVA] F-value: 2.7121; R-squared: 0.14892; P [<0.041]; Figure 3C). Despite this, some

individual taxa were associated with infection status.   Although blood feeding alone increased

the relative amount of *Elizabethkingia*, the highest abundance of this genus was associated

with exposed uninfected midguts (EXP UNINF; Figure 2, Figure 4A). This was particularly the

case with individuals reared at the lower temperature regime (EXP UNINF[L]; Figure 2).

Elizabethkingia is only taxa associated with 'resistance' to infection. Pools of guts with

individuals that disseminated the virus demonstrated higher relative amounts of *Neoasaia* and

additonal taxa belonging to the order Rhodospirillales, family Acetobacteraceae (*Roseomonas*,

and *Swaminathania*). Other taxa associated with infection and dissemination included *E. coli*,

Salmonella and *Tanticharoenia* (Figures 2 and 4; Supplementary Figure 1).

Higher temperature enriches the gut microbial community of *Ae. albopictus*

Increased temperatures altered the microbial composition and increased overall microbial

diversity in the absence of ZIKV exposure ([Alpha diversity, Shannon index P (0.022377); T-test

statistic: -2.4172]; [Beta diversity [PERMANOVA] F-value: 3.1477; R-squared: 0.089557; P

(<0.045)]; Figure 3E). The higher diversity is primarily due to a lower relative abundance of

Elizabethkingia at the higher (M) temperature regime (Figure 2), which is also the primary

driver of the measured difference in beta diversity (Figure 3F).   The fact that this difference in

alpha diversity among temperature regimes in the UNEXP mosquitoes is significant only when

using the Shannon, rather than Chao1 index, is due to the increased likelihood of rare taxa in

the M temperature regime (Figure 3D/E). Regardless of index, this difference is lost following

ZIKV exposure (Figure 3D/E), where diversity is significantly lower at both L and M temperature

regimes relative to UNEXP mosquitoes. Despite this lack of temperature-dependent variability

in alpha diversity among ZIKV EXP mosquitoes, the relative abundance of *Elizabethkingia*

remains lower at higher temperatures (Figure 2).

***rpoB* gene sequencing of bacterial strains isolated from *Aedes* mosquitoes**

Given the relative abundance of *Elizabethkingia* and *Neoasaia*, together with their association

with phenotypes of interest (resistance to infection and dissemination, respectively) these taxa

were chosen for further sequencing.

To the best of our knowledge, this is the initial reporting of detection of *Neoasaia* bacterial

strain from an insect. The only other strain present in the literature was derived from a plant

(*Neoasaia chiangmaiensis*)(Yukphan et al., 2005). The *Neoasaia* strain detected in mosquitoes

in this study clustered with this previously identified strain (Figure 5A). The *Elizabethkingia*

strain isolated from the gut of the *Ae. albopictus* in this study clustered with pathogenic strain

of medical importance. Unexpectedly, it did not share a clade with the *Elizabethkingia*

anophelis, which was previously isolated from Anopheline mosquitoes (Kämpfer et al., 2011)

(Figure 5B).

Discussion

This study has demonstrated a significant reduction in *Ae. albopictus* gut microbial diversity

resulting from blood feeding which is further exacerbated by ZIKV exposure and modified at

different temperature regimes. These data further identified individual gut microbes in *Aedes*

mosquitoes that potentially play a modulating role in ZIKV susceptibility and competence.

Recent studies have confirmed the importance of microbiota in the modulation of vector

competence (Apte-Deshpande et al., 2012; Bourtzis et al., 2014; Dickson et al., 2017; Ramirez et

al., 2014; Weiss & Aksoy, 2011). Microbiota can influence competence through inhibition of

the pathogen by secreted enzymes or toxins, or by the induction of the host immune system

targeting the invading pathogenic microbe (Weiss & Aksoy, 2011).

With the aim of delineating possible taxa modulating ZIKV infection of the mosquito midguts,

we characterized the midgut microbial profile before and after feeding on non-infectious and

infectious blood meals and further compared profiles based on infection and dissemination

status. In addition, we measured the effect of temperature on the midgut microbial community

in an effort to understand how climate change might ultimately alter microbial taxa that

influence vectorial capacity for ZIKV and other arboviruses.

Our data support previous studies demonstrating that bacterial composition of the mosquito

midgut was significantly altered by blood meal acquisition and digestion. This is likely brought

about by an oxidative shift in the midgut after blood digestion (Champion & Xu, 2017; Zouache

et al., 2012). While blood feeding significantly decreased overall microbial diversity, substantial

increases in the relative abundance of *Elizabethkingia* were measured, clearly indicating that

the gut environment is more amenable to the propagation of this species following blood meal

digestion. This alteration in microbial composition could be beneficial to *Ae. albopictus.*

Specifically, *Elizabethkingia anophelis* has been shown to produce OxyR regulon and

antioxidants that in turn potentially protect the mosquitoes from the oxidative stress brought

about by blood digestion. The bacteria also displays hemolytic activity and hence may assist in

erythrocyte digestion in the mosquito midgut (Kukutla et al., 2014).

Our data additionally demonstrated that ZIKV exposure significantly altered microbial

composition independent of the effect of blood feeding. Specifically, ZIKV exposure was

associated with elimination of a number of taxa that were present in the mosquito midgut

before and after non-infectious blood meals. This resulted in a few select taxa dominating the

midgut microbial community of the ZIKV exposed midguts. While *Elizabethkingia* was the

dominant genus in all ZIKV exposed guts, the relative abundance of this microbe was higher in

guts that did not acquire infections.

Elizabethkingia species have been previously identified with *Anopheles* mosquitoes, *Aedes*

mosquitoes and most recently, *Culicoides* (Mee et al., 2017, Boissière et al., 2012; Kukutla et al.,

2014; Terenius et al., 2012; Wang et al., 2011). The association of *Elizabethkingia* abundance

with ZIKV resistance suggests a potential antiviral effect. While further studies are needed to

define this relationship, one potential mechanism is the interaction of *Elizabethkingia* with the

Reactive Oxygen Species (ROS) pathway, which is documented to influence *Aedes* competence

for arboviruses (Li et al., 2015). The genus *Elizabethkingia* (formerly Chryseobacterium or

Flavobacterium), a gram-negative bacilli, is additionally a source of public health concern (Janda

& Lopez, 2017). Three species of this genus have been shown to cause human infections of

medical importance: *E. meningoseptica*, *E. anopheles* and *E. miricola* (Janda & Lopez, 2017; Lau

et al., 2016). Its existence in a similar clade as these medically important strains (*E. miricola* and

E. meningoseptica), as shown by our *rpoB* phylogenetic analysis, demonstrates that further

investigation into its pathogenicity in humans is warranted.

To the best of our knowledge, our study is the initial report of an isolation of *Neoasaia spp* from

an arthropod vector, with a single previous isolation of *Neoasaia* reported from a red ginger

flower (Yukphan et al., 2005). While the mechanistic basis for a potential interaction is unclear,

we found *Neoasaia* positively associated with the development of disseminated ZIKV infections,

underscoring the need for further transcriptomic or metabolomic studies. *Neoasaia spp*, an

acetic acid bacteria, belongs to the family Acetobacteraceae which were initially recovered

from bees (Bruce White, 1921). The acetic acid bacteria have been found to colonize insects

that mainly rely on sugar-based diets such as nectars, fruit sugars or phloem (Russell et al.,

2017). The closely related *Asaia spp* has also been found in association with the pathogen-

transmitting mosquitoes *An. Stephensis, An. Maculipennnis, An. gambiae* and *Ae. aegypti*

(Crotti et al., 2009; Damiani et al., 2010; Favia et al., 2007).

In addition to these taxa, data suggested other bacterial species are associated with ZIKV

competence. For instance, *Escherichia coli* was found to have a higher relative abundance in

midguts of individuals with disseminated infections. *E. coli* has been shown to induce hypoxia in

the midgut of mosquitoes through cytochrome bd oxidase (Coon et al., 2017) and it is feasible

that this could indirectly influence ZIKV competence. It is also notable that the *Ae. albopictus*

midgut samples had very little detectable *Wolbachia*, however whole-body samples were found

to have very high amounts of *Wolbachia* (data not shown). This is consistent with the majority

of *Wolbachia* being in the germline cells. It is possible that there is competitive exclusion of

Wolbachia in the midgut by another microbe. Studies have shown that *Ae. albopictus* and *Cx.*

quinquefasciatus naturally infected by both *Asaia* and *Wolbachia* have a co-exclusion type of a

relationship. In species not colonized by *Wolbachia* such as *An. gambiae, An. stephensi* and *Ae.*

aegypti, Asaia is capable of colonizing reproductive organs and salivary gland (Rossi et al.,

2015).

It is well-documented that temperature can significantly alter vectorial capacity of *Aedes*

mosquitoes for arboviruses including ZIKV (Ali et al., 2017;Paz & Semenza, 2016; Onyango et al.,

under review), yet the relationship between altered vectorial capacity, temperature and

microbial communities is not well documented. Temperature is an important component in

shaping the microbial communities of organisms (Fuhrman et al., 2008; Lokmer & Wegner,

2015) and our data clearly demonstrated a complex relationship between *Ae. albopictus* midgut

microbial diversity and temperature. This complexity is partially due to the confounding effects

of blood meal digestion and virus exposure, but also clearly a result of temperature

differentially affecting unique bacterial taxa. For example, higher temperatures are associated

with a decrease in the relative abundance of *Elizabethkingia* and an increase in the relative

abundance of *Neoasaia*. Given the association of these taxa with ZIKV resistance and

dissemination, respectively, these data suggest that modifications to microbial communities at

higher temperatures could be a contributing factor to increased transmissibility of ZIKV and

other arboviruses with climate change. This also demonstrates that individual bacterial species

that may potentially act as biomarkers for infection and/or be utilized as novel vector control

tools could be differentially affected by climate change. These relationships should therefore be

considered when assessing the potential effectiveness of such strategies.

Acknowledgements

All sequencing was completed at the Wadsworth Center Applied Genomics Core and media was

provided by the Wadsworth Center Media and Tissue Culture facilities. This publication was

supported by the Cooperative Agreement Number U01CK000509 funded by the Centers for

Disease Control and Prevention. Its contents are solely the responsibility of the authors and do

not necessarily represent the official views of the Centers for Disease Control and Prevention or

the Department of Health and Human Services. We would like to extend our gratitude to Illia

Rochlin of Suffolk County Health Department for providing us with *Ae. albopictus* mosquitoes.

References

Ali, S., Gugliemini, O., Harber, S., Harrison, A., Houle, L., Ivory, J., … Mordecai, E. A. (2017).
Environmental and Social Change Drive the Explosive Emergence of Zika Virus in the
Americas. *PLoS Neglected Tropical Diseases*, 11(2).

https://doi.org/10.1371/journal.pntd.0005135

Apte-Deshpande, A., Paingankar, M., Gokhale, M. D., & Deobagkar, D. N. (2012). Serratia
odorifera a midgut inhabitant of aedes aegypti mosquito enhances its susceptibility to
dengue-2 virus. *PLoS ONE*, *7*(7). https://doi.org/10.1371/journal.pone.0040401

Asad, S., Hussain, M., Hugo, L., Osei-Amo, S., Zhang, G., Watterson, D., & Asgari, S. (2018).
Suppression of the pelo protein by Wolbachia and its effect on dengue virus in Aedes
aegypti. *PLoS Neglected Tropical Diseases*, *12*(4), 1–20.
https://doi.org/10.1371/journal.pntd.0006405

Bourtzis, K., Dobson, S. L., Xi, Z., Rasgon, J. L., Calvitti, M., Moreira, L. A., … Gilles, J. R. L. (2014).
Harnessing mosquito-Wolbachia symbiosis for vector and disease control. *Acta Tropica*,
132(1), S150–S163. https://doi.org/10.1016/j.actatropica.2013.11.004

Bray, R. J., & Curtis, J. T. (1957). An Ordination of the Upland Forest Communities of Southern
Wisconsin. *Ecological Monographs*, *27*(4), 325–349.

Bruce White, P. (1921). The Normal Bacterial Flora of the Bee. *The Journal of Pathology*, *24*(1).

Callaway, E. (2016). Infected mosquitoes fight Zika. *Nature*, *539*(7627), 17–18.
https://doi.org/10.1038/nature.2016.20878

Champion, C. J., & Xu, J. (2017). The impact of metagenomic interplay on the mosquito redox
homeostasis. *Free Radical Biology and Medicine*, *105*(August 2016), 79–85.
https://doi.org/10.1016/j.freeradbiomed.2016.11.031

Chao, A. (1984). Nonparametric Estimation the Number of Classes in a Population. *The Annals*
of Mathematical Statistics, *20*(4), 265–270. https://doi.org/10.1214/aoms/1177729949

Ciota, A., Bialosuknia, S., Zink, S., Brecher, M., Ehrbar, D., Morrissette, M., & Kramer, L. D.
(2017). Effects of Zika Virus Strain and Aedes Mosquito Species on Vector Competence.
Emerging Infectious Diseases, *23*(7), 1110–1117.

Coon, K. L., Valzania, L., McKinney, D. A., Vogel, K. J., Brown, M. R., & Strand, M. R. (2017).
Bacteria-mediated hypoxia functions as a signal for mosquito development. *Proceedings of*
the National Academy of Sciences, *114*(27), E5362–E5369.
https://doi.org/10.1073/pnas.1702983114

Crotti, E., Damiani, C., Pajoro, M., Gonella, E., Rizzi, A., Ricci, I., … Daffonchio, D. (2009). Asaia, a
versatile acetic acid bacterial symbiont, capable of cross-colonizing insects of
phylogenetically distant genera and orders. *Environmental Microbiology*, *11*(12), 3252–
3264. https://doi.org/10.1111/j.1462-2920.2009.02048.x

Damiani, C., Ricci, I., Crotti, E., Rossi, P., Rizzi, A., Scuppa, P., … Favia, G. (2010). Mosquito-
Bacteria Symbiosis: The Case of Anopheles gambiae and Asaia. *Microbial Ecology*, *60*(3),
644–654. https://doi.org/10.1007/s00248-010-9704-8

Dhariwal, A., Chong, J., Habib, S., King, I. L., Agellon, L. B., & Xia, J. (2017). MicrobiomeAnalyst: A
web-based tool for comprehensive statistical, visual and meta-analysis of microbiome
data. *Nucleic Acids Research*, *45*(W1), W180–W188. https://doi.org/10.1093/nar/gkx295

Dickson, L. B., Jiolle, D., Minard, G., Moltini-Conclois, I., Volant, S., Ghozlane, A., … Lambrechts,
L. (2017). Carryover effects of larval exposure to different environmental bacteria drive
adult trait variation in a mosquito vector. *Science Advances*, *3*(8), 1–15.
https://doi.org/10.1126/sciadv.1700585

Dillon, R. J., & Dillon, V. M. (2004). The Gut Bacteria of Insects: Nonpathogenic Interactions.
Annual Review of Entomology, *49*, 71–92.

Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Research*, *32*(5), 1792–1797.
https://doi.org/10.1093/nar/gkh340

Engla, N. E. W., & Journal, N. D. (2010). New engla nd journal, 1061–1063.

Favia, G., Ricci, I., Damiani, C., Raddadi, N., Crotti, E., Marzorati, M., … Daffonchio, D. (2007).
Bacteria of the genus Asaia stably associate with Anopheles stephensi, an Asian malarial
mosquito vector. *Proceedings of the National Academy of Sciences*, *104*(21), 9047–9051.
https://doi.org/10.1073/pnas.0610451104

Fuhrman, J. A., Steele, J. A., Hewson, I., Schwalbach, M. S., Brown, M. V., Green, J. L., & Brown,
J. H. (2008). A latitudinal diversity gradient in planktonic marine bacteria. *Proceedings of*
the National Academy of Sciences, *105*(22), 7774–7778.
https://doi.org/10.1073/pnas.0803070105

Guégan, M., Zouache, K., Démichel, C., Minard, G., Tran Van, V., Potier, P., … Valiente Moro, C.
(2018). The mosquito holobiont: fresh insight into mosquito-microbiota interactions.
Microbiome, *6*(1), 49. https://doi.org/10.1186/s40168-018-0435-2

Hall, T. (2011). BioEdit: An important software for molecular biology. *GERF Bulletin of*
Biosciences, *2*(1), 60–61.

Janda, J. M., & Lopez, D. L. (2017a). Mini review: New pathogen profiles: Elizabethkingia
anophelis. *Diagnostic Microbiology and Infectious Disease*, *88*(2), 201–205.
https://doi.org/10.1016/j.diagmicrobio.2017.03.007

Janda, J. M., & Lopez, D. L. (2017b). Mini review: New pathogen profiles: Elizabethkingia
anophelis. *Diagnostic Microbiology and Infectious Disease*, *88*(2), 201–205.
https://doi.org/10.1016/j.diagmicrobio.2017.03.007

Kämpfer, P., Matthews, H., Glaeser, S. P., Martin, K., Lodders, N., & Faye, I. (2011).
Elizabethkingia anophelis sp. nov., isolated from the midgut of the mosquito Anopheles
gambiae. *International Journal of Systematic and Evolutionary Microbiology*, *61*(11), 2670–
2675. https://doi.org/10.1099/ijs.0.026393-0

Kjellstrom, E., Nikulin, G., Strandberg, G., Christensen, O., Jacob, D., Keuler, K., … Vautard, R.
(2018). European climate change at global mean temperature increases of 1.5 and 2C
above pre-industrial conditions as simulated by the EURO-CORDEX regional climate
models. *Earth System Dynamics*, *9*, 459–478. https://doi.org/ethz-b-000265761

Kukutla, P., Lindberg, B. G., Pei, D., Rayl, M., Wanqin, Y., Steritz, M., … Xu, J. (2014). Insights
from the genome annotation of Elizabethkingia anophelis from the malaria vector
anopheles gambiae. *PLoS ONE*, *9*(5). https://doi.org/10.1371/journal.pone.0097715

Lau, S. K. P., Chow, W. N., Foo, C. H., Curreem, S. O. T., Lo, G. C. S., Teng, J. L. L., … Woo, P. C. Y.
(2016). Elizabethkingia anophelis bacteremia is associated with clinically significant
infections and high mortality. *Scientific Reports*, *6*(March), 8–17.
https://doi.org/10.1038/srep26045

Li, Y., Liu, Y., Chew, S. C., Tay, M., Margarette, M., Salido, S., … Yang, L. (2015). Complete
Genome Sequence and Transcriptomic Analysis of the Novel Pathogen Elizabethkingia
anophelis in Response to Oxidative Stress, *7*(6), 1676–1685.
https://doi.org/10.1093/gbe/evv101

Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News*,
2(December), 18–22. https://doi.org/10.1023/A:1010933404324

Lindsey, A. R. I., Bhattacharya, T., Newton, I. L. G., & Hardy, R. W. (2018). Conflict in the
intracellular lives of endosymbionts and viruses: A mechanistic look at Wolbachia-

434  mediated pathogen-blocking. *Viruses*, *10*(4). https://doi.org/10.3390/v10040141

435 Lokmer, A., & Mathias Wegner, K. (2015). Hemolymph microbiome of Pacific oysters in

436  response to temperature, temperature stress and infection. *ISME Journal*, *9*(3), 670–682.

437  https://doi.org/10.1038/ismej.2014.160

438 McLean, B. J., Dainty, K. R., Flores, H. A., & O'Neill, S. L. (2019). Differential suppression of

439  persistent insect specific viruses in trans-infected wMel and wMelPop-CLA Aedes-derived

440  mosquito lines. *Virology*, *527*(August 2018), 141–145.

441  https://doi.org/10.1016/j.virol.2018.11.012

442 Mee, P. T., Lynch, S. ., Walker, P. J., Melville, L., & Duchemin, J.-B. (2017). Detection of

443  Elizabethkingia spp. in Culicoides Biting Midges, Australia. *Emerging Infectious Diseases*,

444  *23*(8), 1409–1410. https://doi.org/10.1093/tropej/49.5.308

445 Mohanty, I., Rath, A., Swain, S. P., Pradhan, N., & Hazra, R. K. (2018). Wolbachia Population in

446  Vectors and Non-vectors: A Sustainable Approach Towards Dengue Control. *Current*

447  *Microbiology*, *0*(0), 0. https://doi.org/10.1007/s00284-018-1596-8

448 Moretti, R., Yen, P. S., Houé, V., Lampazzi, E., Desiderio, A., Failloux, A. B., & Calvitti, M. (2018).

449  Combining Wolbachia-induced sterility and virus protection to fight Aedes albopictus-

450  borne viruses. *PLoS Neglected Tropical Diseases*, *12*(7), 1–20.

451  https://doi.org/10.1371/journal.pntd.0006626

452 Motzkin, T. (1949). The euclidean algorithm. In *Bulletin of the American Society* (pp. 1142–

453  1146).

454 Nguyen, T. H., Nguyen, H. Le, Nguyen, T. Y., Vu, S. N., Tran, N. D., Le, T. N., … Hoffmann, A. A.

455  (2015). Field evaluation of the establishment potential of wmelpop Wolbachia in Australia

456  and Vietnam for dengue control. *Parasites and Vectors*, *8*(1), 1–14.

457  https://doi.org/10.1186/s13071-015-1174-x

458 Paz, S., & Semenza, J. C. (2016). Correspondence El Niño and climate. *The Lancet*, *387*(10020),

459  745. https://doi.org/10.1016/S0140-6736(16)00256-7

460 Ramirez, J. L., Short, S. M., Bahia, A. C., Saraiva, R. G., Dong, Y., Kang, S., … Dimopoulos, G.

461  (2014). Chromobacterium Csp_P Reduces Malaria and Dengue Infection in Vector

462  Mosquitoes and Has Entomopathogenic and In Vitro Anti-pathogen Activities. *PLoS*

463  *Pathogens*, *10*(10). https://doi.org/10.1371/journal.ppat.1004398

464 Romoli, O., & Gendrin, M. (2018). The tripartite interactions between the mosquito, its

465  microbiota and Plasmodium. *Parasites and Vectors*, *11*(1), 1–8.

466  https://doi.org/10.1186/s13071-018-2784-x

467 Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., … Huelsenbeck,

468  J. P. (2012). Mrbayes 3.2: Efficient bayesian phylogenetic inference and model choice

469  across a large model space. *Systematic Biology*, *61*(3), 539–542.

470  https://doi.org/10.1093/sysbio/sys029

471 Rossi, P., Ricci, I., Cappelli, A., Damiani, C., Ulissi, U., Mancini, M. V., … Favia, G. (2015). Mutual

472  exclusion of Asaia and Wolbachia in the reproductive organs of mosquito vectors.

473  *Parasites and Vectors*, *8*(1), 1–10. https://doi.org/10.1186/s13071-015-0888-0

474 Russell, J. A., Sanders, J. G., & Moreau, C. S. (2017). Hotspots for symbiosis: Function, evolution,

475  and specificity of ant-microbe associations from trunk to tips of the ant phylogeny

476  (Hymenoptera: Formicidae). *Myrmecological News*, *24*(21), 43–69.

477  https://doi.org/10.1128/AEM.01336-10

478 Shannon, C. E. (1964). A mathematical theory of communication. *ACM SIGMOBILE Mobile*

Computing and Communications Review, *5*(1), 3. https://doi.org/10.1145/584091.584093

Svetnik, V., Liaw, A., Tong, C., Christopher Culberson, J., Sheridan, R. P., & Feuston, B. P. (2003).
Random Forest: A Classification and Regression Tool for Compound Classification and
QSAR Modeling. *Journal of Chemical Information and Computer Sciences*, *43*(6), 1947–
1958. https://doi.org/10.1021/ci034160g

Vallet-Gely, I., Novikov, A., Augusto, L., Liehl, P., Bolbach, G., Pechy-Tarr, M., … Lemaitre, B.
(2009). Implication of the Mosquito Midgut Microbiota in the Defense against Malaria
Parasites. *Applied and Environmental Microbiology*, *76*(3), 910–921.
https://doi.org/10.1371/journal.ppat.1000423

Ward, J., & Hook, M. (1963). Application of an hierarchical grouping procedure to a problem of
grouping profiles. *Educational and Psychological Measurement*, *XXIII*(1).

Weiss, B., & Aksoy, S. (2011). Microbiome influences on insect host vector competence. *Trends*
in Parasitology, *27*(11), 514–522. https://doi.org/10.1016/j.pt.2011.05.001

Yukphan, P., Malimas, T., Potacharoen, W., Tanasupawat, S., Tanticharoen, M., & Yamada, Y.
(2005). Neoasaia chiangmaiensis gen. nov., sp. nov., a novel osmotolerant acetic acid
bacterium in the alpha-Proteobacteria. *J Gen Appl Microbiol*, *51*(5), 301–311.
https://doi.org/10.2323/jgam.51.301

Zouache, K., Michelland, R. J., Failloux, A. B., Grundmann, G. L., & Mavingui, P. (2012).
Chikungunya virus impacts the diversity of symbiotic bacteria in mosquito vector.
Molecular Ecology, *21*(9), 2297–2309. https://doi.org/10.1111/j.1365-294X.2012.05526.x

Author Contributions

MGO: Involved in the study design, conducted the experiments, analyzed the results and
drafted the paper
BS: Involved in the mosquito rearing, assisted in mosquito manipulation exercises
PA: Assisted in mosquito manipulation exercises
MN: Involved in the mosquito rearing, assisted in mosquito manipulation exercises
KL: Generated and propagated the ZIKV infectious clone utilised in this study
KLD: Involved in the study design, participated in the writing of the paper
CAT: Involved in the study design, participated in the writing of the paper

Figures

Figure 1. Proportions of bacterial phyla identified in *Ae. albopictus* midguts.

Data indicates proportions identified by 16s sequencing for all pools of midguts combined

(n=34). Pools consisted of 3 individual midguts. All samples were acquired at 7 days post-

feeding or at equivalent age for unfed individuals.

Figure 2. Relative abundance of bacterial genera in individual pools of *Ae. albopictus* midguts.

Each stacked bar plot illustrates the taxa distribution at the genus level for pools of 3 *Ae. albopictus* midguts. Samples were taken at 7 days post-feeding or equivalent age in unfed (UNF) individuals. Blood fed individuals were further separated into UNEXP (fed non-infectious blood meal), EXP UNINF (fed ZIKV-positive blood meal but did not acquire infection) or EXP DIS (fed ZIKV positive blood meal and developed a disseminated infection). Mosquitoes were hatched and reared under two thermal conditions; 14 hours light at 30°C: 10 hours dark at 26°C (M temperature regimen) or 14 hours light at 28°C: 10 hours dark at 24°C (L temperature regimen).

Figure 3. Blood feeding, ZIKV exposure and temperature have a significant effect on microbial diversity in *Ae. albopictus* midguts.

Distribution of alpha diversity within groups as measured by Chao1 or Shannon Index measures. The differences in diversity between groups was measured by Bray Curtis diversity

A. The effect of blood feeding on *Ae. albopictus* midgut microbial diversity. Comparisons of pools of midguts from unfed individuals (UNF) to those fed non-infectious blood meals demonstrates that blood meal acquisition independently decreases alpha diversity at 7 days post-feeding (chao1 T-test/ANOVA P(0.006) [T-test] statistic -3.1216). **B.** The effect of Zika virus exposure on *Ae. albopictus* midgut microbial diversity. Comparisons of pools of midguts from ZIKV exposed individuals (FED EXP) to those fed non-infectious blood meals demonstrates that ZIKV exposure independently decreases alpha diversity at 7 days post-feeding (chao1 T-test/ANOVA P(0.005) [T-test] statistic -3.1084). **C.** The effect of ZIKV infection on *Ae. albopictus* midgut microbial diversity. Comparisons of pools of midguts from ZIKV exposed individuals that were not infected (EXP UNINF) to those that were infected and developed disseminated infections (EXP DIS) demonstrates that ZIKV infection and dissemination do not independently decreases alpha diversity at 7 days post-feeding (chao1 T-test/ANOVA P(0.005) [T-test] statistic -3.1084). **D-F.** The effect of temperature on *Ae. albopictus* midgut microbial diversity. Although comparison of microbial diversity for individuals held at L and M temperature regimes demonstrates no statistical difference when using the Chao1 index (**D**), a significant temperature effect on alpha diversity is measured with the Shannon index (**E**; Shannon P(0.0065); [ANOVA] F-value: 5.4042) such that higher temperature are associated with decreased microbial diversity for ZIKV exposed individuals (EXP) but increased microbial diversity for individuals fed non-infectious blood meals (UNEXP). A significant effect of temperature on beta diversity was additionally measured (**F**; [PERMANOVA] F-value: 3.7463; R-squared: 0.34861; P[=0.018]).

Figure 4. *Ae. albopictus* **midgut microbial signatures associated with blood feeding, ZIKV exposure and ZIKV infection.**

The Random forest classifier was used to determine the association of differentially abundant taxa in the microbial community with certain states. Mean Decrease Accuracy is reported for each of the taxa, which is obtained by removing the relationship of a taxa and measuring increase in error. The taxa with highest mean decrease in accuracy is considered to have the highest association with the state. **A.** Taxa were listed in order of the strength of their level of association with states including UNFED (no blood meal), UNEXP (blood fed non-infectious blood), EXP UNINF (infectious blood meal but no infection established) and EXP INF (ZIKV positive following infectious blood meal). **B.** Taxa were listed in order of the strength of their level of association with states including EXP DIS (fed ZIKV positive blood meal and developed a disseminated infection), EXP UNINF (fed ZIKV-positive blood meal but did not acquire infection) and UNEXP (fed non-infectious blood meal).

Figure 5. Phylogenetic analysis of *Neoasaia* and *Elizabethkingia rpoB* gene

Phylogenetic affiliation of the *Neoasaia* (**A**) and *Elizabethkingia* (**B**) taxa from the midgut of

Aedes mosquitoes (shown in red) based on a partial sequence of the *rpoB* gene. Phylogenetic

dendrograms were constructed in BAYES. Bootstrap proportions are shown on branches.

Escherichia coli was used as an outgroup.

Supplementary Figure 1. A heat map showing the relative abundance of taxa at OTU level and

their distribution across groups.