Rotavirus re-programs multiple IFN receptors and restricts their intestinal antiviral and inflammatory functions

Adrish Sen[1,2]**, Nima D. Namsa**[1, 2, 3]**, Ningguo Feng**[1,2]**, Harry B. Greenberg**[1,2]

The Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, California, USA[1]; VA Palo Alto Health Care System, Palo Alto, California, USA[2], Department of Molecular Biology and Biotechnology, Tezpur University (A Central University), Napaam 784 028, Assam, India[3]

ABSTRACT

Rotaviruses (RV) cause acute severe diarrhea in the absence of substantial intestinal inflammation. They are also highly infectious in their homologous host species. The efficient replication capacity of RV in the small bowel is substantially linked to its ability to inhibit different types of interferons (IFNs). Here, we find that during RV infection *in vitro*, both virus-infected and uninfected bystander cells resist STAT1 phosphorylation and IRF7 induction in response to exogenous IFN. Functionally, cellular transcription in response to exogenous stimulation with IFN, but not intracellular dsRNA, was inhibited by RV. Further, IFNAR1 stimulation during RV infection significantly repressed a set of virus-induced transcripts. Regulation of IFN signaling *in vivo* was studied in suckling mice using the highly infectious homologous murine EW RV strain. Kinetic studies indicated that whereas sustained EW RV replication and IFN induction occurred in the small intestine, IFN-stimulated transcripts significantly decreased over time. In addition, LPS-mediated intestinal damage, driven by STAT1-induced inflammation, was prevented in EW RV-infected mice. Remarkably, ectopic stimulation of either IFNAR1 or IFNGR1 in murine RV-infected mice eliminated several intestinal antiviral and inflammatory transcriptional responses to RV. In contrast to homologous murine RV, infection with a STAT1-sensitive heterologous simian RV strain induced multiple IFN-stimulated transcripts, inflammatory cytokines, and intestinal expression of STAT1-pY701. Finally, RV strain-specific STAT1 regulation in the gut plays a prominent role in the activation of multiple intestinal caspases. On the other hand, the simian RRV strain, but not murine EW RV, uniquely triggers the cleavage of both extrinsic and intrinsic caspases (-8, -9, and -3) in a STAT1-mediated manner. Collectively, these findings

reveal efficient re-programing of multiple IFN receptors in the gut towards a negative

feedback mode of signaling, accompanied by suppression of IFN-mediated antiviral,

apoptotic, and inflammatory functions, during natural RV intestinal infection.

<u>INTRODUCTION</u>

Rotaviruses are non-enveloped segmented dsRNA viruses that cause severe

dehydrating diarrhea in many mammals and an estimated 200,000 human infant deaths

annually. Acute RV infection in a homologous host (i.e. the host species in which a

specific RV strain replicates to high titer and spreads from individual to individual) is an

extremely efficient process; severe diarrhea can be initiated by exposure to a very small

infectious inoculum[1]. The host innate interferon (IFN) response is a multi-pronged line of

defense against many viral pathogens as its induction can activate hundreds of different

antiviral genes and trigger inflammatory signals – thus eliminating infected cells, limiting

the spread of virus infection to uninfected bystander cells, and shaping both the

magnitude and quality of subsequent adaptive immune responses[2].

All three major IFN types (α/β, γ, and λ) are induced during homologous RV

infection of the small bowel[3,4], but their antiviral effects on viral replication are modest at

best. Although relatively poorly understood, the inflammatory effects of IFNs are also

likely impaired during RV infection, which is associated with only mild intestinal

inflammation compared to other acute diarrheal pathogens[5-7]. Previously, in a suckling

mouse model of RV disease, we demonstrated that the IFNAR1, IFNGR1, and IFNLR1

receptors, as well as the downstream convergent transcription factor STAT1, play an

important role in restricting heterologous RV (RV not typically isolated from the

symptomatically infected host species) replication[1,3,4,8,9]. Depletion of IFNRs in suckling

mice also results in an acute biliary inflammatory disease (often lethal) following

infection with a heterologous (rhesus RRV) but not a homologous (murine EW RV)

strain[8,10,11]. How homologous RV efficiently restricts the ability of different IFNs to

execute their antiviral and inflammatory programs is not well understood but important

both for more effective strategies to combat RV-associated mortalities and to better

understand the varying pathogenic potentials of naturally circulating RV strains.

MATERIALS AND METHODS

Viruses and reagents. The tissue culture-adapted simian RRV and porcine SB1-A RV

strains were propagated in African green monkey kidney cells (MA104) as previously

described[12]. Virus titers were determined by plaque assays in MA104 cells. HT-29 cells,

a colon cancer derived cell line with a variety of small bowel features[13], were plated in 6-

well or 24-well cluster plates and completely confluent monolayers were infected 2-3d

later as described[14]. The non-tissue culture-adapted murine EW RV strain was prepared

as an intestinal homogenate from RV-infected Sv129 suckling mice. The diarrheal dose

(DD_{50}) was determined as described[9]. Human intestinal epithelial HT29 and embryonic

kidney HEK293 cells were purchased from the American Type Culture Collection

(ATCC) and maintained in Advanced D-MEM Ham's F12 (Cellgro) or D-MEM medium

(D-MEM, Cellgro) containing 10% fetal calf serum (Invitrogen) supplemented with non-

essential amino acids, glutamine, penicillin, and streptomycin, respectively. Purified

human IFN-β (PBL) was used for stimulation of cells. Purified carrier-free universal type

I IFN A/D (PBL), murine IFN-γ (PBL), LPS (Sigma) were used in animal studies. High-

molecular weight LyoVec poly(I:C) (Invivogen) was used at 2.5 µg/mL.

Mouse infection and tissue harvest.   All mice were housed in the VA Palo Alto

Health Care System (VAPAHCS) Veterinary Medicine Unit and all experimentation

followed VAPAHCS Institutional Animal Care and Use Committee approved protocols.

3-5 day old Sv129 pups were infected via oral gavage with 10^4 DD_{50} of murine EW RV,

4 x 10^6 PFU of the simian RRV, or mock infected as described[9]. At the indicated time-

points, the small intestines were removed. Sections were collected in Trizol for RNA

purification or in 2X Laemmli buffer for protein analysis by immunoblotting, as

described[4] and stored at -80 °C, or fixed in 4% paraformaldehyde for processing and

subsequent analyses as described[15].

Mouse stimulation and Luminex assays: To ectopically stimulate IFNAR1 and

IFNGR1, suckling mice infected with 10^4 DD_{50} of murine EW RV (or mock-infected) for

12 hours were intraperitoneally administered purified murine IFN-γ (1μg IFN-γ per pup)

or universal type I IFN A/D (1μg per pup) and sacrificed 12 h later for analysis of

transcripts by qRT-PCR as described[4]. For endotoxin stimulation, 3-5 day old 129sv

mice were orally inoculated by gavage with 10^4 DD_{50} of murine RV-EW. At 3 dpi, mice

were intraperitoneally injected with PBS or purified LPS (Sigma, 10 mg/kg body weight)

and sacrificed 6 h later for analysis of paraformaldehyde-fixed and paraffin embedded

small bowel tissue sections by hematoxylin and eosin staining. For measurement of

intestinal cytokines, mouse small intestines were harvested and weighed, and pooled

tissues (from 3 pups) were homogenized in ice-cold PBS containing a cocktail of

protease and phosphatase inhibitors (Sigma) using a handheld micropestle (Thermo

Fisher). Lysates were clarified by two sequential rounds of centrifugation at 10,000X G

for 10 min, and analyzed for cytokine levels at the Human Immune Monitoring Center at

Stanford University. Mouse 38-plex kits were purchased from eBiosciences/Affymetrix

and used according to the manufacturer's recommendations. Plates were read using a

Luminex 200 instrument with a lower bound of 50 beads per sample per cytokine.

Custom assay Control beads by Radix Biosolutions are added to all wells. Murine IFN-β

was measured using a commercial ELISA kit as per the manufacturer's instructions

(PBL).

Western blot analysis. Preparation of cell lysates and immunoblotting *in vitro* was

performed as described[12]. For mouse samples, intestinal tissue pieces were lysed in

radioimmunoprecipitation assay buffer (RIPA) supplemented with protease and

phosphatase inhibitors (Roche). Equal volumes of 2X Laemmli buffer (Sigma-Aldrich)

were added to each sample. Viscosity was reduced by passing repeatedly through a

22G syringe and samples were boiled for 5 minutes prior to analysis by SDS-PAGE and

immunoblotting. Blots were probed with primary antibodies directed towards the

following antigens: STAT1-pY701, cleaved caspase 8, caspase 9, cleaved caspase 3,

(Cell Signaling Technologies), RV VP6 (Santa Cruz Biotechnologies), and B-Actin

(Sigma). The blots were subsequently incubated with either anti-rabbit or anti-mouse

secondary antibodies conjugated to horseradish peroxidase (HRP, Amersham),

incubated with ECL detection reagent, and exposed to film (GE Healthcare). All

densitometry quantifications of protein blots were performed using ImageJ software

analysis and normalized to actin protein bands.

Transcriptional analysis. For *in vivo* analysis of transcripts, mice were sacrificed and

sections of the small intestine were collected on ice in Trizol (Life Technologies). For *in*

vitro studies, cells were first washed in PBS and then lysed in Trizol. Total RNA was

extracted following the manufacturer's instructions and subjected to DNAse digestion

before use in qRT-PCR. Synthesis of cDNA and subsequent microfluidics PCR on the

Fluidigm platform was done as described earlier[16] using commercially available taqman

assays (Thermo-Fisher) and RV NSP5-specific assays described earlier[3,4,9]. Serial 10-

fold dilutions of mouse or human reference RNA (Agilent) and blank controls were run in

duplicate for each PCR run.

Flow cytometry analysis. Staining and flow cytometry analysis of HT-29 cells was

performed as described[14]. Cells were plated in 24-well cluster plates and infected with

RV. Infected cells were harvested at 6 hpi or 16 hpi and transfected cells at 48 after

transfection for flow cytometry analysis. Cells were washed in PBS and fixed at room

temperature for 10 min using 1.6% (v/v) methanol-free paraformaldehyde (Electron

Microscopy Services). Cells were washed in FACS staining buffer (PBS containing

0.5% w/v bovine serum albumin) and permeabilized in cold methanol at 4 °C for 10min.

Cells were washed in FACS staining buffer, stained using antibodies directed to either

STAT1-pY701 or IRF7 (from BD Biosciences), or to the RV VP6 antigen[14] at 4 °C for 30

min, and washed prior to analysis by flow cytometry on a LSRII instrument (BD

Biosciences). The flow data was analyzed using FlowJo software.

Statistical analysis. Analysis of variance (ANOVA) was carried out by Kruskal-Wallis

method using Dunn correction for multiple comparisons. Protein bands from SDS-

PAGE blots were analyzed using Holm-Sidak t-test. P values <0.05 were considered

significant as indicated in the figure legends. All error bars represent the standard error

of the mean.

RESULTS

Rotavirus restricts IFN-directed STAT1 activation and IRF7 expression. Prior

studies using several RV strains identified two distinct RV strategies to regulate IFN

induction that are exemplified by either porcine (degrades β-TRCP) or simian (degrades

IRFs) RV strains[17]. In contrast to this distinction, we reported that both porcine SB1-A

and simian RRV strains similarly mediated the lysosomal degradation of endogenous

IFNAR1, IFNGR1, and IFNLR1 subunits in RV infected (RV+) cells[15] and prevented

IFN-directed STAT1 phosphorylation at the Y701 epitope[14], which is a critical

convergent signal transduction event downstream of the different IFNRs. Surprisingly, in

other *in vitro* studies, SB1-A was also found to inhibit IFN-directed STAT1-Y701

phosphorylation in RV- bystander cells, which expressed normal levels of IFN

receptors[14]. Here, we extended our studies by examining whether IFN signaling is also

inhibited in RV- bystander cells by the simian RV strain RRV. HT-29 cells were infected

with RRV at a multiplicity of infection resulting in approximately 35% RV infected VP6+

cells (Fig. 1A, column 1) and at 12 hpi cells were stimulated for 2h with increasing

doses of purified universal type I IFN (IFN-A/D) to induce STAT1-Y701 phosphorylation

(Fig. 1A, columns 2-6). Analysis of RV VP6 and STAT1-pY701 expression by FACS

demonstrated that mock-infected controls underwent a dose-dependent increase in

STAT1-pY701 expression in response to ectopic IFN (Fig. 1A, row 1 and Fig. 1B). In

contrast, both RRV VP6+ (RV+) and RRV VP6- (bystander) cell populations effectively

resisted IFN-stimulated STAT1 phosphorylation, even at the highest concentration of

IFN-A/D tested (Fig. 1A, row 2 and Fig. 1B). These findings demonstrate that the IFN

signaling pathway is inhibited in uninfected bystander cells *in vitro* by the simian RV

(RRV), similar to our earlier findings using a porcine RV[14]. Thus, although uninfected

RV bystander cells express normal levels of IFNRs[15] during infection, their ability to

transduce IFN-directed signals is significantly impaired.

To extend these observations, we examined the effect of simian RRV strain on

levels of endogenous IRF7, an IFN-stimulated protein that is also targeted for

degradation by the RRV-encoded NSP1 protein[18]. Analysis of RV VP6 and IRF7 by

FACS was performed in HT-29 cells as described above (Fig. 1C). In mock controls,

IRF7 was constitutively expressed by ~33% cells even in the absence of IFN

stimulation. Following stimulation with either a low (312.5 IU/ml) or high (5,000 IU/ml)

dose of IFN-A/D, a similar increase in IRF7 expression occurred (~60% cells were

IRF7+ post-IFN stimulation at either dosage) (Fig. 1C, row 1 and Fig. 1D), indicating

that relatively low IFN concentrations are sufficient to maximally induce IRF7 expression

in HT-29 cells. Following RV infection, RRV VP6- cells exhibited elevated basal IRF7

expression in the absence of any exogenous IFN stimulation (i.e. ~43% of all cells,

almost entirely deriving from the VP6- gate, were IRF7+), which could be possibly

triggered by low levels of IFN induced earlier during RV infection[12,19]. However, IFN

stimulation failed to induce any further increase in IRF7 expression in RRV VP6- cells

(Fig. 1C, row 2 and Fig. 1D). In contrast, RV+ HT-29 cells expressed substantially lower

levels of IRF7 (only ~4% VP6+ cells were IRF7+). Notably, no detectable increase in

IRF7 expression occurred in infected cells following IFN-A/D stimulation at either dose.

Overall, in these cell monolayers the percentage of HT-29 cells expressing IRF7 after

stimulation with 5,000 IU/ml IFN was lower during RRV infection (~39% of all cells) than

mock infection (~61% of all cells). By qRT-PCR analysis, we confirmed that although

RRV infection induced IRF7 transcription compared to mock controls, it prevented any

further exogenous IFN-stimulated IRF7 transcription (Fig. 1E). These initial data

demonstrate that RV-infected cells are highly resistant to IFN-mediated induction of

IRF7 or STAT1-pY701. In addition, the findings indicate that although RRV induces

IRF7 in bystander cells during infection, it effectively prevents further IRF7 induction by

saturating doses of IFN stimulus, which is in general agreement with the observed

blockage of STAT1 phosphorylation in RV bystander cells (Fig. 1A). Since optimal IFN-

directed STAT1-pY701 and IRF7 induction are important for IFNR positive feedforward

transcriptional amplification[20], it seemed likely that RV infection might perturb this

process in both infected and bystander cells.

Rotavirus perturbs interferon-stimulated feedforward cellular transcriptional

response. To gain further insight into the regulation of the IFN signaling response by

RV, we next examined the ability of ectopic IFN to upregulate antiviral and inflammatory

transcription following RRV infection of HT-29 cells. Since several ISGs and

inflammatory transcripts upregulated during IFN signaling are also induced by

intracellular dsRNA[21], primarily through NF-kB and IRF3 signaling pathways, we

compared cellular transcriptional responses to either exogenously applied IFN-β or

intracellular long dsRNA introduced by transfection. Following RV infection for 6h, cells

were stimulated with either purified IFN-β or with liposomally-complexed long dsRNA for

an additional 6h and then analyzed by qRT-PCR. The results (Fig. 2A) demonstrate that

mock-infected HT-29 cells underwent significant upregulation of several transcripts in

response to either IFN and/or dsRNA stimulation (*IFIT3, IFI6, CCL5, TNF, HERC5* were

induced by both types of stimuli; *SERPING, CXCL10, IFI16, TNFAIP3* by IFN-A/D only;

IL29, CCL3, IFNG by dsRNA only). In RV-infected samples, significant transcriptional

upregulation of *CCL5, TNF, IL29, CCL3*, and *IRF8* occurred only in response to

intracellular dsRNA. In addition, *RSAD2, IFIT3, IFI6, CXCL10, IFI16* and *TNFAIP3*

transcripts were also weakly, but not significantly, elevated in response to dsRNA during

RV infection. In contrast, we failed to detect any significant upregulation of measured

transcripts in response to IFN-β with the sole exception of *SERPING*, whose induction

was weaker in RV-infected samples than in mock-infected controls (Fig. 2A). The

stimulus-specific RV effect on cellular transcription was unlikely due to differences in the

level of RV replication, since it was unaltered by either IFN-β or dsRNA stimulation (Fig.

2B).

Interestingly, ectopic stimulation of IFNAR1 during RRV infection also

significantly repressed the expression of a set of transcripts from their levels in the

unstimulated RRV samples (*IFNG, IRF8, HERC5, IL27, IL10*, and *IL10RA*) (Fig. 2A,

bottom row). Although RV infection completely prevented (or repressed) the IFN-

directed upregulation of transcripts, by FACS analysis we found that only ~30% of the

cells in the monolayer were RRV VP6+ (Fig. 2C). Hence, these results suggested that

RV inhibits IFN-directed cellular transcription in both RV-infected and in uninfected

bystander cells.

Rotavirus infection also restricts intestinal IFN amplification responses *in vivo*.

The transcriptional amplification of IFNs (e.g. *IL28*, *IFNG*), antiviral genes (e,g, *MX2*),

and inflammatory cytokines (e.g. *CXCL10*) occurs via STAT1 activation as part of the

IFN feedforward signaling circuit[22]. We next examined the kinetics of selected IFN

stimulated transcripts in the small intestine of suckling mice infected with a highly

infectious homologous murine EW RV strain[1] for either 12h or 72h (Fig. 3A-B). This

analysis revealed that *IFNA4* (a primarily induced IFN gene, i.e. prior to the IFN

amplification phase[20]) and *RV NSP3* transcripts continued to significantly increase

between 12-72 hpi. In contrast, a significant decrease in RV-induced intestinal type II

and III IFNs (*IL28* and *IFNG), RSAD2, IFIT2,* and *CXCL10* transcription occurred in EW

RV-infected mice between 12-72 hpi (Fig. 3A). RV-induced transcription of *MX2*, a

specific marker of the intestinal epithelial response to IFN stimulation[23], also significantly

declined from 12 to 72 hpi. These results indicate that during murine EW RV infection of

suckling mice, despite sustained viral replication and IFN induction, several intestinal

IFN amplification responses are down-regulated by 72 hpi. To further examine this

finding, we directly tested whether intestinal damage caused by endotoxin, which

mediates inflammation in a STAT1-dependent manner[24], is also down-regulated by prior

infection with EW RV. Suckling mice orally infected with EW RV (or mock) for 72h were

systemically administered endotoxin and resulting intestinal damage assessed 6h later.

In mock-infected pups, parenteral administration of endotoxin led to severe intestinal

necrosis and tissue destruction (Fig. 3C). In contrast, suckling mice that were infected

with EW RV prior to endotoxin challenge exhibited minimal small intestinal damage (Fig.

3C). These results further indicate that RV infection in the small bowel can suppress

intestinal IFN-associated amplification responses.

Rotavirus reprograms intestinal IFN type I and II receptors towards negative

feedback signaling. Our data *in vitro* (Fig. 2A) indicated that ectopic IFN-stimulated

transcription was suppressed during RV infection. Therefore, we next examined

intestinal transcriptional responses to ectopically administered type I and type II IFNs in

suckling mice infected with murine RV. Following EW RV infection for 12h, pups were

parenterally administered either purified IFN-A/D or IFN-γ (or PBS), and small intestines

were collected after an additional 12h for transcriptional qRT-PCR analysis (Fig. 4A). In

the absence of any ectopic IFN stimulation, RV infection transcriptionally induced

several intestinal IFNs, ISGs, and inflammatory genes, as expected from our prior

findings (Fig. 3A). Remarkably, ectopic stimulation of IFNAR1 or IFNGR1 in the murine

RV infected pups resulted in either a significant suppression (*IL28, IFIH1*) or the

complete elimination (*IFNB1, IFNA5, IFIT1, MX2, TNFSF10, CXCL10, STAT1*) of

several innate transcriptional responses to the RV infection. A similar IFN-dependent

effect was not observed for *PDCD1*, *PCNA*, *GAPDH*, or *IKBKB* transcripts, or for *ARG2*

and *IL12B,* whose expression during RV infection significantly increased following

administration of either type of IFN (Fig. 4A). To ascertain whether the suppressive

effect of IFN stimulation on RV associated transcription could be explained by the

exogenous IFN mediated inhibition of EW RV replication, we measured intestinal RV

RNA by qRT-PCR using two different EW RV-specific PCR assays (Fig. 4B). These

results demonstrated that EW RV replication was not significantly altered following

either IFN-A/D or IFN-γ stimulation under these conditions (Fig. 4B). These results

reveal that during RV infection, exogenous stimulation of either type I and II IFNRs in

the small intestine has the potential to efficiently suppress both antiviral and

inflammatory transcriptional responses.

Intestinal IFN signaling is regulated differently during infection with homologous

versus heterologous RV*s*. The finding that murine EW RV impairs IFN signaling in

suckling mice is consistent with studies in mice[8] and human enteroids[25] that

demonstrate RV replication in the homologous host species is IFN-resistant. In contrast,

we previously observed that replication of the heterologous simian RRV strain in

suckling mice is highly sensitive to different types of IFN and the presence of STAT1 (by

2-4 logs)[3,4,8,9]. We thus compared intestinal IFN-stimulated antiviral genes and

cytokines in mice infected with the heterologous simian RRV, homologous murine EW,

or mock-infected. As shown in Figure 5A, at 12 hpi the intestinal replication of

heterologous RRV was substantially and significantly lower than the homologous EW

RV. However, RRV infection triggered significantly higher transcription levels of the

ISG20, IFI203, and PELI1 genes as compared to EW RV (Fig. 5A). In addition,

heterologous RV infection also resulted in significantly elevated intestinal protein levels

of IFN-β and the IFN-stimulated pro-inflammatory cytokines IP10 and RANTES when

compared to homologous RV (Fig. 5B). The expression of type III IFN (IL28) was also

elevated in RRV-infected pups compared to either the EW RV or mock groups.

Analysis of intestines collected from EW and RRV infected mice at 1 dpi by

immunoblotting showed that compared to either mock or EW infection, RRV induces

significant phosphorylation of STAT1 at Y701 (Fig. 5C), which is a convergent signaling

response to different types of IFNs in the intestinal epithelium[15]. Collectively, these

findings are consistent with the conclusion that, in contrast to homologous murine RV,

heterologous simian RV strongly activates IFN-dependent STAT1 and pro-inflammatory

responses in the small bowel.

STAT1 activation status regulates the cleavage of multiple intestinal caspases

during homologous vs. heterologous RV infection *in vivo*. Different IFNs, as well as

several IFN-stimulated cytokines (such as IP10 and RANTES), induce caspase-

regulated cell death to potentiate inflammation[26,27]. Differences between EW and RRV

strains in their ability to regulate pSTAT1 and cytokine expression prompted us to

examine the status of intestinal caspases following infection of suckling mice with these

two RV strains. Using immunoblotting analysis, the cleavage status of the initiator

caspases 8 and 9, and the executioner caspase 3, were examined in the small

intestines of WT and STAT1$^{-/-}$ mice infected with either EW or RRV RV (Fig. 6A). WT

mice infected with RRV RV for 24 h undergo substantially higher levels of cleavage of

caspases 8, 9, and 3 when compared to their mock-infected or EW RV counterparts

(Fig. 6A-B), indicating that activation of both extrinsic and intrinsic intestinal apoptotic

pathways is induced by heterologous RRV more efficiently than the homologous murine

EW strain.

Of note, the enhanced cleavage of caspases during RRV infection was

dependent on the intestinal IFN response. Specifically, we found that differences in

caspase 8, 9, and 3 cleavage between mock and RRV infected WT pups were

abrogated following infection in STAT1 knockout mice (Fig. 6A-B). The cleavage of

caspases during EW RV infection was also STAT1-mediated. However, EW RV-

infected STAT1$^{-/-}$ mice demonstrated cleaved caspases 8, 9, and 3 at levels much lower

than in mock-infected controls or RRV infected mice (Fig. 6A and 6B), indicating that the

basal cleavage efficiency of multiple caspases (i.e. occurring in the absence of

infection) was impaired by homologous EW but not heterologous RRV RV.

Impaired caspase 8 cleavage can spontaneously induce acute intestinal

inflammation through necroptosis, resulting in the secretion of the pro-inflammatory

factor IL1α[28]. Therefore, we examined IL1α expression in the small intestine during EW

and RRV infection (Fig. 6C). While the intestinal level of IL1α was no different between

mock- and EW RV-infected suckling mice at 1 dpi, it was significantly elevated after

infection with RRV RV[29] (Fig. 6C). A similar difference was also observed for IL6,

another common marker of intestinal inflammation[30] (Fig. 6C). Thus, although murine

EW RV inhibits caspase 8 cleavage when compared to simian RRV, it avoids inducing

the pro-inflammatory cytokines IL1α or IL6 by a yet unknown mechanism. Collectively,

these data identify an unrecognized role for STAT1 in mediating the cleavage of

multiple intestinal caspases during RV infection. The findings also indicate that the

ability of homologous RV to regulate IFN and STAT1 signaling *in vivo* (Figs. 4 and 5)

likely dampens IFN regulated cell death and inflammation.

<u>DISCUSSION</u>

Although they have been isolated from many different mammalian species, RVs are

generally highly infectious only in their homologous host species. Robust RV replication

occurs predominantly in the small intestinal mature villous tip epithelium and causes

severe diarrhea, but only mild inflammation[6,7]. We and others demonstrated that both

RV intestinal replication and accompanying inflammation are likely to be substantially

regulated by host interferon (IFN) responses. Specifically, intestinal replication of a

heterologous simian RRV strain can be significantly enhanced in suckling mice lacking

IFNAR1, IFNGR1, and/or IFNLR1 receptors, or the convergent downstream

transcription factor STAT1[3]. In the absence of STAT1, infection of suckling mice with the

normally restricted RRV strain causes severe inflammation (mostly in the biliary tract),

systemic spread, and mortality[8]. In contrast, replication of a homologous murine EW RV

strain is quite resistant to IFNs[8], and mice infected with EW RV are subsequently

protected from lethal endotoxemia[15], which is STAT1-mediated[24]. Paradoxically, EW RV

robustly induces all three major IFN types[3,4], raising the possibility that subsequent

IFNR-directed antiviral and inflammatory signaling pathways are effectively blocked

during homologous infection.

How homologous RV successfully inhibits the ability of intestinal IFNs to mediate

antiviral and inflammatory effects is not yet well understood and was therefore

examined in this study. We found that between 12-72 hpi, mice infected with murine EW

RV exhibit sustained viral replication and increased transcription of the primary (i.e.

induced prior to IFN amplification) murine type I IFN gene (IFNA4, Fig. 3A-B)[20].

However, transcriptional induction of the IFNL2/3, MX2, IFNG, RSAD2, IFIT2, and

CXCL10 (Fig. 3A) genes in response to ongoing EW RV replication was considerably

diminished by 72 hpi. Together, these data suggest that although ongoing EW RV

replication triggers intestinal IFN induction signaling pathways, it progressively restricts

intestinal IFN amplification responses. This conclusion is further supported by our

finding that suckling mice infected with EW RV for 72 h are protected from severe small

intestinal injury that normally occurs following systemic LPS exposure (Fig. 3B). This

injury was reported to be driven primarily by STAT1 signaling in the mouse model of

endotoxemia[24]. Using an alternate approach, we established that suckling mice infected

with the STAT1-sensitive[3] simian RRV strain exhibit significantly higher levels of

intestinal ISG transcriptional induction (Fig. 5A), IFN-β protein expression (Fig. 5B), and

STAT1-pY701 expression (Fig. 5C) compared to the homologous murine EW RV strain.

In addition, expression of the IFN-stimulated pro-inflammatory cytokines IP10[31,32] and

RANTES[32-34] are also higher in the intestines of heterologous simian RRV-infected mice

(Fig. 5B). Collectively, these data support the conclusion that infection with homologous

RV *in vivo* results in an inhibition of both IFN-mediated antiviral and inflammatory

functions in the small intestine.

Caspase cleavage is a key parameter for determining both the magnitude and

nature of inflammatory responses driven by IFNs and other cytokines[35,36]. Exposure to

different IFNs can either induce caspase 8-mediated apoptosis[37] or alternately, trigger

inflammatory cell death (necroptosis) under circumstances when caspase 8 cleavage is

impaired[38]. Data presented here (Fig. 6) demonstrate that murine EW RV infected mice

express reduced cleavage of intestinal caspase 8, 9, and 3 when compared to RRV RV;

cleavage levels being similar to the mock-infected control pups (Fig. 6A-B). In contrast,

infection with RRV triggered the cleavage of all three caspases in the small intestine,

indicating that both extrinsic and intrinsic apoptosis pathways are induced by the

heterologous RRV despite its restricted replication capacity. Interestingly, activation of

different caspases by RRV was contingent on the presence of STAT1, as differences in

cleaved caspase 8, 9, and 3 expression between RRV and mock animals were

eliminated in a STAT1 deficient background (Fig. 6A-B). The genetic KO of STAT1 in

suckling mice infected with EW RV resulted in cleaved caspase levels lower than those

in uninfected mock controls (Fig 6B). This was particularly evident for cleaved caspase

8 and 3 expression in the EW RV infected STAT1$^{-/-}$ mice that were at or below the limit

of detection (Fig. 6A and B). A likely explanation for these findings is that during the

course of infection, EW RV regulation of IFN feedforward signaling (Fig. 4) and STAT1

phosphorylation (Fig. 5C) inhibits the efficiency of intestinal caspase 8 and 3 cleavage.

Interestingly,   inhibition of caspase 8 cleavage has been reported to induce

spontaneous intestinal inflammation, primarily via necroptosis[28]. However, expression of

IL-1α – a member of the IL-1 family that is released from necroptotic cells[39], and of the

pro-inflammatory cytokine IL-6, were not detected in EW RV (or mock infected control)

pup intestines, but were significantly induced by RRV RV (Fig. 6C). Future studies will

be required to understand how homologous RV inhibits the cleavage of intestinal

caspases yet avoids causing intestinal inflammation. These findings nevertheless reveal

a STAT1-dependent intestinal caspase activation pathway that is likely regulated by

homologous RV to control inflammation.

Findings in this and earlier reports[3,4,8,9,15] indicate that mechanisms by which RV

regulates STAT1 signaling are likely critical determinants of rotaviral pathogenicity *in*

vivo. We found earlier that several RVs (porcine, bovine, and simian RV strains)

mediate the lysosomal degradation of multiple IFN receptor subunits (i.e. IFNAR1,

IFNGR1, and IFNLR1) in RV VP6+ (i.e. infected) cells, whereas RV VP6- (i.e.

bystander) cells expressed normal levels of the IFN receptors[15]. Remarkably, a porcine

RV strain (SB1-A) was shown to also efficiently inhibit IFN-directed STAT1-Y701

phosphorylation in uninfected (i.e. VP6-) bystander cells[14], indicating the existence of an

alternative STAT1 inhibitory mechanism that operates beyond the virus-infected cell. In

this study, we extended these findings and found that the simian RRV strain can also

block IFN-β mediated STAT1-pY701 induction in both VP6-positive and -negative HT-29

cell populations (Fig. 1A-B). In addition, RRV infection prevented IFNβ-directed

induction of the IRF7 protein in both infected and bystander cells (Fig. 1C). Together

with our previous observation that both simian and porcine RV strains can mediate

IFNR degradation[15], these findings suggest that RV strategies to inhibit STAT1 signaling

in infected and bystander cells are conserved across multiple strains that differ in the

manner in which they regulate the IFN induction phase[17].

Findings of this study also revealed a potential mechanism by which RV confers

resistance to ectopic IFN stimulation to uninfected bystander cells[14]. Specifically, under

infection conditions *in vitro* where only ~30% HT-29 cells were RRV VP6+, ectopic

stimulation with IFN-I failed to evoke significant cellular transcriptional responses (Fig.

2). This inhibitory effect of RRV specifically targeted IFN signaling, as several of these

measured transcripts could be efficiently induced during RRV infection by intracellular

dsRNA stimulation (Fig. 2A). In addition, ectopic IFNAR1 stimulation during RRV

infection also resulted in a significant decline of certain innate immune transcripts

induced in response to RV (Fig. 2A). These collective findings not only demonstrate that

RV infection can block IFNR-directed feedforward transcription but also suggest that RV

represses transcription following ectopic IFN stimulation, which is characteristic of an

IFNR-directed negative feedback transcription mode[2].

The interesting possibility of negative feedback transcription effects is also

supported from our studies in suckling mice which we previously showed that

exogenous IFN I or II induced STAT1-Y701 in the small intestine with rapid kinetics (i.e.

30 min) when administered intra-peritoneally[15]. Remarkably, here we find that ectopic

stimulation of either IFNR early during EW RV infection (i.e. at 12 hpi) substantially

repressed several measured intestinal antiviral transcriptional responses (Fig. 4A). In

particular, intestinal induction of both type I (*IFNB1* and *IFNA5*) and type III (*IL28*) IFN

transcripts by EW RV was significantly dampened when IFNAR1 or IFNGR1 were

ectopically stimulated. In addition, RV-induced MX2 transcription, which is reported to

be primarily IFNLR1-driven in the intestinal epithelial compartment[23], was eliminated

following stimulation with either types I or II IFN.

Ectopic stimulation of IFNGR1 and IFNAR1 has been reported to elicit negative

feedback signaling leading to IFN-resistance in the context of cancer and chronic viral

infections[2,40]. The ability of homologous RV to elicit IFNAR1- and IFNGR1-stimulated

feedback signaling suggests a potential mechanism underlying the ability of EW RV-

infected mice to resist lethal endotoxemia[15] and the accompanying intestinal damage

(Fig. 3C). It will be interesting to identify the RV-mediated effectors involved in the broad

suppression of multiple intestinal IFNR-dependent antiviral and inflammatory functions.

As both type I and III IFNs were found to exert a profound antiviral activity against

heterologous RV[3], one possibility is that the endogenous induction of these IFNs during

homologous EW RV infection[3,4,19] serves to dampen intestinal antiviral, apoptotic, and

inflammatory transcription and promote efficient viral replication.

ACKNOWLEDGEMENTS

This work was supported by grants 1RO1 AI021362 (HBG), R56 AI125249 (HBG), and

1RO1 AI1125249 (HBG) from the National Institutes of Health, and 1IO 1BX000158-

01A1 from the Veterans Affairs (HBG).

FIGURE LEGENDS

Figure 1. Rotavirus inhibits IFN-stimulated cellular responses *in vitro*. (A-B) HT-29

cells were infected with RRV (or mock infected) at a MOI of 1. At 10 hpi cells were

stimulated with purified IFN-β at the indicated concentrations for 2 hours and analyzed

by flow cytometry. (C-D) cells were infected with RRV and analyzed by FACS as in (A).

(E) Cells infected in triplicate as in (C) and at 12 hpi were stimulated with PBS (black

bars) or 500 IU/ml of purified IFN-β (blue bars) for 6h before analysis by qRT-PCR for

IRF7 and HPRT transcript levels. (*p<0.02, *ns* not significant). Data shown are

representative of two or more independent experiments.

Figure 2. Rotavirus infection inhibits IFN-stimulated feedforward cellular

transcription. (A-B) HT-29 cells grown in triplicate wells were infected with RRV strain

at a MOI of 1 (or mock-infected) for 6h and then stimulated with 2µg of [>200 bp long

dsRNA:liposome] complexes, or 500 IU/ml of purified IFN-β. At 12 hpi cells were lysed

for RNA purification and then analyzed by qRT-PCR for expression of indicated host (A)

and RV (B) transcripts. (*ns,* not significant, *p<0.02, **p<0.002, ***p<0.0002;

significance is shown relative to untreated samples in mock and RV infected groups).

(C) Cells were infected as in (A), and the percentage of cells expressing RV VP6

antigen before lysis was determined by FACS. Box indicates gates used to estimate

RV-infected cells in the monolayer). Data shown is representative of two experiments.

Figure 3. Effects of homologous RV infection on intestinal IFN amplification

responses. (A-B) Suckling mice were infected with murine EW RV (or were mock

infected, N=3-8 mice per group) for the times indicated, and small intestines collected to

analyze the expression of transcripts shown. Data shown is fold change in EW RV

versus mock infected mice. (C) Suckling mice were mock- or EW RV-infected and 3 d

later were parenterally administered purified endotoxin (15µg/pup) for 6 h before

harvesting small intestines for histological examination by hematoxylin and eosin

staining. Data shown is representative of 2 pups per group.

Figure 4. Rotavirus reprograms intestinal IFNAR1 and IFNGR1 towards a negative

feedback mode of transcription. Schematic of the experimental approach used is

shown at the top. Mice infected with EW RV (or mock) for 12 h were administered

purified universal IFN-A/D (IFN-I), murine IFNγ (IFN-II), or PBS by the i.p. route. Twelve

hours later mice were sacrificed, and small intestinal RNA analyzed by qRT-PCR for

different host (A) or RV gene (B) transcripts. (*p<0.02, **p<0.002, ***p<0.0002;

significance in each panel is relative to PBS-treated EW RV infected mice, unless

indicated otherwise). Data is measured from 3-5 pups per group.

Figure 5. Differences between homologous and heterologous RV strains on IFN

stimulated genes, inflammatory cytokine expression, and STAT1-Y701 activation

in vivo. (A) Suckling mice (N= 3-5 pups per group) were infected with homologous

murine EW or heterologous simian RRV RV strains (or mock infected). At 12 hpi mice

were sacrificed for subsequent analysis of small intestinal transcripts by qRT-PCR. Data

shown is fold relative to mock-infected mice. (B) Mice infected as in (A) were sacrificed

and small intestinal biopsies were analyzed for the expression of the proteins shown

either by ELISA (IFN-β) or Luminex-based cytokine assays. (C) Mice infected as in (A)

were sacrificed and small intestines analyzed for immunoblotting analysis.

Densitometric quantitation of pSTAT1 bands is shown in the bar diagram at the right.

(*p<0.02, **p<0.002, ***p<0.0002).

Figure 6. Homologous rotavirus inhibits the STAT1-mediated cleavage of multiple

intestinal caspases. (A-B) Either wild type (WT) or STAT1-deficient mice were infected

with EW or RRV RV strains (or mock infected). Twenty-four hours later, mice were

sacrificed, and small intestinal lysates were analyzed by immunoblotting (A).

Densitometric estimation of bands is shown in (B). (C) Mice infected as in (A) were

sacrificed and intestines analyzed for the expression of cytokines by luminex-based

assays. (*$p<0.02$, **$p<0.002$, ***$p<0.0002$)

REFERENCES

Broome, R. L., Vo, P. T., Ward, R. L., Clark, H. F. & Greenberg, H. B. Murine rotavirus genes encoding outer capsid proteins VP4 and VP7 are not major determinants of host range restriction and virulence. *J Virol* **67**, 2448-2455 (1993).

Ivashkiv, L. B. & Donlin, L. T. Regulation of type I interferon responses. *Nat Rev Immunol* **14**, 36-49, doi:10.1038/nri3581 (2014).

Lin, J. D. *et al.* Distinct Roles of Type I and Type III Interferons in Intestinal Immunity to Homologous and Heterologous Rotavirus Infections. *PLoS Pathog* **12**, e1005600, doi:10.1371/journal.ppat.1005600 (2016).

Sen, A. *et al.* Innate immune response to homologous rotavirus infection in the small intestinal villous epithelium at single-cell resolution. *Proc Natl Acad Sci U S A* **109**, 20667-20672, doi:10.1073/pnas.1212188109 (2012).

Greenberg, H. B. & Estes, M. K. Rotaviruses: from pathogenesis to vaccination. *Gastroenterology* **136**, 1939-1951, doi:10.1053/j.gastro.2009.02.076 (2009).

Lundgren, O. & Svensson, L. Pathogenesis of rotavirus diarrhea. *Microbes Infect* **3**, 1145-1156 (2001).

Morris, A. P. & Estes, M. K. Microbes and microbial toxins: paradigms for microbial-mucosal interactions. VIII. Pathological consequences of rotavirus infection and its enterotoxin. *Am J Physiol Gastrointest Liver Physiol* **281**, G303-310, doi:10.1152/ajpgi.2001.281.2.G303 (2001).

Feng, N. *et al.* Role of interferon in homologous and heterologous rotavirus infection in the intestines and extraintestinal organs of suckling mice. *J Virol* **82**, 7578-7590, doi:10.1128/JVI.00391-08 (2008).

Feng, N., Yasukawa, L. L., Sen, A. & Greenberg, H. B. Permissive replication of homologous murine rotavirus in the mouse intestine is primarily regulated by VP4 and NSP1. *J Virol* **87**, 8307-8316, doi:10.1128/JVI.00619-13 (2013).

10   Riepenhoff-Talty, M. *et al.* Group A rotaviruses produce extrahepatic biliary obstruction
in orally inoculated newborn mice. *Pediatr Res* **33**, 394-399, doi:10.1203/00006450-
199304000-00016 (1993).

11   Allen, S. R. *et al.* Effect of rotavirus strain on the murine model of biliary atresia. *J Virol*
81, 1671-1679, doi:10.1128/JVI.02094-06 (2007).

12   Sen, A., Feng, N., Ettayebi, K., Hardy, M. E. & Greenberg, H. B. IRF3 inhibition by
rotavirus NSP1 is host cell and virus strain dependent but independent of NSP1
proteasomal degradation. *J Virol* **83**, 10322-10335, doi:10.1128/JVI.01186-09 (2009).

13   Adachi, A. *et al.* Productive, persistent infection of human colorectal cell lines with
human immunodeficiency virus. *J Virol* **61**, 209-213 (1987).

14   Sen, A., Rott, L., Phan, N., Mukherjee, G. & Greenberg, H. B. Rotavirus NSP1 protein
inhibits interferon-mediated STAT1 activation. *J Virol* **88**, 41-53, doi:10.1128/JVI.01501-
13 (2014).

15   Sen, A., Sharma, A. & Greenberg, H. B. Rotavirus degrades multiple type interferon
receptors to inhibit IFN signaling and protects against mortality from endotoxin in
suckling mice. *J Virol*, doi:10.1128/JVI.01394-17 (2017).

16   Sen, A. *et al.* Innate immune response to homologous rotavirus infection in the small
intestinal villous epithelium at single-cell resolution. *Proc Natl Acad Sci U S A* **109**,
20667-20672, doi:10.1073/pnas.1212188109 (2012).

17   Arnold, M. M. & Patton, J. T. Diversity of interferon antagonist activities mediated by
NSP1 proteins of different rotavirus strains. *J Virol* **85**, 1970-1979,
doi:10.1128/JVI.01801-10 (2011).

18   Arnold, M. M., Barro, M. & Patton, J. T. Rotavirus NSP1 mediates degradation of
interferon regulatory factors through targeting of the dimerization domain. *J Virol* **87**,
9813-9821, doi:10.1128/JVI.01146-13 (2013).

19   Sen, A., Pruijssers, A. J., Dermody, T. S., Garcia-Sastre, A. & Greenberg, H. B. The early
interferon response to rotavirus is regulated by PKR and depends on MAVS/IPS-1, RIG-I,
MDA-5, and IRF3. *J Virol* **85**, 3717-3732, doi:10.1128/JVI.02634-10 (2011).

20   Marie, I., Durbin, J. E. & Levy, D. E. Differential viral induction of distinct interferon-alpha
genes by positive feedback through interferon regulatory factor-7. *EMBO J* **17**, 6660-
6669, doi:10.1093/emboj/17.22.6660 (1998).

21   Dalet, A., Gatti, E. & Pierre, P. Integration of PKR-dependent translation inhibition with
innate immunity is required for a coordinated anti-viral response. *FEBS Lett* **589**, 1539-
1545, doi:10.1016/j.febslet.2015.05.006 (2015).

22   Crotta, S. *et al.* Type I and type III interferons drive redundant amplification loops to
induce a transcriptional signature in influenza-infected airway epithelia. *PLoS Pathog* **9**,
e1003773, doi:10.1371/journal.ppat.1003773 (2013).

23   Pott, J. *et al.* IFN-lambda determines the intestinal epithelial antiviral host defense. *Proc*
Natl Acad Sci U S A **108**, 7944-7949, doi:10.1073/pnas.1100552108 (2011).

24   Karaghiosoff, M. *et al.* Central role for type I interferons and Tyk2 in lipopolysaccharide-
induced endotoxin shock. *Nat Immunol* **4**, 471-477, doi:10.1038/ni910 (2003).

25   Saxena, K. *et al.* A paradox of transcriptional and functional innate interferon responses
of human intestinal enteroids to enteric virus infection. *Proc Natl Acad Sci U S A* **114**,
E570-E579, doi:10.1073/pnas.1615422114 (2017).

26 Apelbaum, A., Yarden, G., Warszawski, S., Harari, D. & Schreiber, G. Type I interferons
induce apoptosis by balancing cFLIP and caspase-8 independent of death ligands. *Mol*
Cell Biol **33**, 800-814, doi:10.1128/MCB.01430-12 (2013).
27 Chawla-Sarkar, M., Leaman, D. W. & Borden, E. C. Preferential induction of apoptosis by
interferon (IFN)-beta compared with IFN-alpha2: correlation with TRAIL/Apo2L
induction in melanoma cell lines. *Clin Cancer Res* **7**, 1821-1831 (2001).
28 Gunther, C. *et al.* Caspase-8 regulates TNF-alpha-induced epithelial necroptosis and
terminal ileitis. *Nature* **477**, 335-339, doi:10.1038/nature10400 (2011).
29 Gabay, C. Interleukin-6 and chronic inflammation. *Arthritis Res Ther* **8 Suppl 2**, S3,
doi:10.1186/ar1917 (2006).
30 Atreya, R. & Neurath, M. F. Involvement of IL-6 in the pathogenesis of inflammatory
bowel disease and colon cancer. *Clin Rev Allergy Immunol* **28**, 187-196,
doi:10.1385/CRIAI:28:3:187 (2005).
31 Majumder, S. *et al.* p48/STAT-1alpha-containing complexes play a predominant role in
induction of IFN-gamma-inducible protein, 10 kDa (IP-10) by IFN-gamma alone or in
synergy with TNF-alpha. *J Immunol* **161**, 4736-4744 (1998).
32 Wen, X. *et al.* Predominant interferon-gamma-mediated expression of CXCL9, CXCL10,
and CCL5 proteins in the brain during chronic infection with Toxoplasma gondii in
BALB/c mice resistant to development of toxoplasmic encephalitis. *J Interferon Cytokine*
Res **30**, 653-660, doi:10.1089/jir.2009.0119 (2010).
33 Kim, M. O., Suh, H. S., Brosnan, C. F. & Lee, S. C. Regulation of RANTES/CCL5 expression
in human astrocytes by interleukin-1 and interferon-beta. *J Neurochem* **90**, 297-308,
doi:10.1111/j.1471-4159.2004.02487.x (2004).
34 Sung, P. S. *et al.* CXCL10 is produced in hepatitis A virus-infected cells in an IRF3-
dependent but IFN-independent manner. *Sci Rep* **7**, 6387, doi:10.1038/s41598-017-
06784-x (2017).
35 Feltham, R., Vince, J. E. & Lawlor, K. E. Caspase-8: not so silently deadly. *Clin Transl*
Immunology **6**, e124, doi:10.1038/cti.2016.83 (2017).
36 Wurstle, M. L., Laussmann, M. A. & Rehm, M. The central role of initiator caspase-9 in
apoptosis signal transduction and the regulation of its activation and activity on the
apoptosome. *Exp Cell Res* **318**, 1213-1220, doi:10.1016/j.yexcr.2012.02.013 (2012).
37 Balachandran, S. *et al.* Alpha/beta interferons potentiate virus-induced apoptosis
through activation of the FADD/Caspase-8 death signaling pathway. *J Virol* **74**, 1513-
1523 (2000).
38 Thapa, R. J. *et al.* Interferon-induced RIP1/RIP3-mediated necrosis requires PKR and is
licensed by FADD and caspases. *Proc Natl Acad Sci U S A* **110**, E3109-3118,
doi:10.1073/pnas.1301218110 (2013).
39 England, H., Summersgill, H. R., Edye, M. E., Rothwell, N. J. & Brough, D. Release of
interleukin-1alpha or interleukin-1beta depends on mechanism of cell death. *J Biol*
Chem **289**, 15942-15950, doi:10.1074/jbc.M114.557561 (2014).
40 Kim, H. S. *et al.* STAT1 deficiency redirects IFN signalling toward suppression of TLR
response through a feedback activation of STAT3. *Sci Rep* **5**, 13414,
doi:10.1038/srep13414 (2015).
637

Figure 1

Figure 2

Figure 3

Figure 4

EW RV   **-/+ IFN**

qRT-PCR^4B analysis

12h   12h

4A

4B

IFNB1

IFIT1

MX2

TNFSF10

ARG2

CXCL10

STAT1

IFNA5

IL28-2

IL12B

PDCD1

PCNA

GAPDH

IFIH1

IKBKB

Gene11-2

ETDG11F1

Figure 5

5A

5B

5C

Figure 6

6A

WT mice   STAT1$^{-/-}$ mice

Mock   EW   RRV   Mock   EW   RRV

Mouse #: 1 2 3 1 2 3 1 2 3   1 2 3 1 2 3 1 2 3

α-Cl-Casp8
α-Casp9
α-Cl-Casp3
Actin
RV VP6

6B

WT mice   STAT1$^{-/-}$ mice

Relative Band Intensity

Cl-Casp8
Cl-Casp9
Cl-Casp3

Mock   EW   RRV   Mock   EW   RRV

6C

IL1α   IL6

● Mock
● EW
● RRV

concentration (pg/ml)