Enhanced leaky sex expression is an adaptive plastic response to pollen limitation in the dioecious plant *Mercurialis annua*

Guillaume Cossard[1] and John R. Pannell[2]

Department of Ecology and Evolution

Biophore Building

University of Lausanne

CH-1015 Lausanne

Switzerland

[1] Current address: Algal Genetics Group

Integrative Biology of Marine Organisms Department (UMR8227)

Station Biologique CNRS UPMC

Place Georges Teissier CS 90074

29688 Roscoff cedex

France

[2] author for correspondence: john.pannell@unil.ch

Running title: Adaptive plasticity in sex expression

Keywords: Dioecy, inconstancy, sex allocation, plasticity, androdioecy, reproductive assurance, hermaphroditism

Author contributions: Conceived the study: GC and JRP; Data collection and analysis: GC; Wrote the manuscript: GC and JRP.

Acknowledgements: We thank J. Gerchen and T. Martignier for comments on the manuscript. The research was funded by the Swiss National Science Foundation.

Abstract

Phenotypic plasticity is all but ubiquitous across the tree of life, but clear demonstrations that it is adaptive are rare. In dioecious plants, males and females frequently show 'leaky' sex expression, with individuals occasionally producing flowers of the opposite sex. Here, we ask whether leaky sex expression in the wind-pollinated dioecious herb *Mercurialis annua* is plastically responsive to its mating context, and whether such plasticity is adaptive. We compared experimental populations of females growing either with or without males. Females growing in the absence of males were leakier in their sex expression than controls with males, producing more than twice as many male flowers, and they enjoy much greater fitness. We discuss differences in the degree of plasticity between ploidal races of annua *Mercurialis* in terms of likely differences in the reliability of the signal plants may perceive for the presence of males in their populations. Our results provide a striking instance of adaptive plasticity in the reproductive behavior of plants and draw attention to possible constraints on plasticity when the environmental signals individuals receive are unreliable.

Introduction

The expression of an individual's phenotype will commonly depend on its environmental context (Schlichting and Pigliucci 1998; Pigliucci 2001). Such phenotypic plasticity is measurable as a genotype by environment (or G x E) interaction, or as a 'norm of reaction', and can have several causes (Van Kleunen and Fischer 2005; Josephs 2018). First, it may be a passive response to different growth conditions, such as variation in resource availability that limits the size of individuals or their capacity to invest in particular structures (Sultan 1987; Dorn et al. 2000; Van Kleunen and Fischer 2005). Such passive responses, which are likely unavoidable consequences of the physiological limits set in different environments, probably account for a much of the phenotypic variation observed among individuals with similar genotypes. Second, phenotypic plasticity may be the result of developmental programs that are altered by exposure to unaccustomed stress in novel environments, such as excessive heat, toxicity or physical damage (Van Kleunen and Fischer 2005). Third, phenotypic plasticity may be an adaptive expression of genotypes habitually exposed to heterogeneous environments (Gomulkiewicz and Kirkpatrick 1992). In this case, plastic genotypes have greater fitness, averaged over the environments they may encounter, than non-plastic 'canalized' genotypes that always express the same phenotype (Releya 2002; Van Kleunen and Fischer 2005). Alternatively, plasticity may be viewed as adaptive if it results in phenotypes that confer local fitness optima across each of the environments concerned (Conover and Schultz 1995; Ghalambor et al. 2007). Either way, whether an observed norm of reaction can properly be interpreted as adapted (in the sense of it being an adaptation; (Dobzhansky 1969; Sultan 1987)) ultimately depends on whether it reflects an evolved response to past selection over essentially the same domain of environmental variation as that observed (Gotthard and Nylin 1995). Being adapted is thus be more difficult to demonstrate than simply whether a response is adaptive.

Although phenotypic plasticity is all but ubiquitous, plants are particularly plastic organisms (Bradshaw 1965; Sultan 1987; Josephs 2018). This is likely due both to their sessile nature, which means that they cannot actively choose their habitat and must make the best of the conditions to which they are exposed, as well as to their modularity, which allows individuals to modify the phenotype of their modules (e.g., branches, inflorescences or flowers) as they are produced in response to environmental variation over time and space. Nevertheless, evidence that plastic variation in plants is adaptive remains surprisingly thin, with few clear demonstrations that the different phenotypes expressed in different environments actually

improve fitness (Van Kleunen and Fischer 2005; Hendry 2015; Wagner and Mitchell-Olds 2018). Several studies have demonstrated that plant responses to shading are adaptive, with fitness benefits to individuals that produce longer internodes and achieve greater height when overtopped by or growing beside potential competitors (e.g., Dudley and Schmitt 1995, 1996). The plastic induction of increased defense in response to herbivory has also been shown to increase individuals' fitness (e.g., Karban et al. 1997; Agrawal 1998; Agrawal 1999). Further, Baythavong (Baythavong and Stanton 2010; 2011) showed that variation in a number of morphological and phenological traits was adaptive in environments with small-scale variation in soil chemistry, and Kenney et al. (2014) showed that plasticity in water-use efficiency (WUE) conferred fitness benefits under dry conditions – though Nicotra and Davidson (2010) reported inconsistent findings in their review of plasticity in WUE among studies.

Plant reproductive traits may also be phenotypically plastic. Examples include: variation in reproductive effort or reproductive allocation, which is sensitive to resource availability and competition (Weiner 2004); sex allocation, e.g., in terms of the relative numbers of male versus female flowers produced by monoecious individuals, which varies with plant size and resource status (Dorken and Barrett 2003; Paquin and Aarssen 2004, Pannell, 1997 #873); and floral longevity and floral display size, with plants adjusting their attractiveness to pollinators in response to the relatedness of their neighbors (Torices et al. 2018) or as a function of relative pollinator abundance and visitation rates (Harder and Johnson 2005). This latter example is particularly interesting in the context of our study here, because it indicates the extent to which plants may alter their reproductive allocation decisions specifically in response to plant mating opportunities. Specifically, Harder and Johnson (2005) found that floral display in the hermaphroditic orchid *Satyrium longicauda* was enhanced when pollinator visitation rates were low, increasing the possibility for later pollen receipt or geitonogamous self-pollination, with likely fitness benefits. Similarly, Lopez et al. (2003) found that in the monoecious plant *Begonia gracilis*, individuals whose female flowers were pollen-limited produced more male flowers than those whose female flowers enjoyed experimentally augmented pollen deposition, suggesting that plants can respond to the operational sex ratio of the population (though without demonstrating a clear effect on fitness in natural populations). In homosporous ferns, gametophytes are more likely to develop as males when females or hermaphrodites are locally abundant, a switch mediated by interplant chemical

signaling (Banks 1997) and interpreted as an adaptive response to mating opportunities (Banks 1997).

Many angiosperms with separate sexes also show variation in sex expression. Specifically, the males and females of dioecious plants commonly display inconstant or 'leaky' sex expression, with the occasional production of a few flowers of the opposite sex (e.g., Baker 1967; Lloyd 1972; Lloyd and Bawa 1984; Diggle 1991; Korpelainen 1998; Delph 2003; Venkatasamy et al. 2007). Such leaky sex expression, which is more common in males than females ((Delph and Wolf 2005; Ehlers and Bataillon 2007), though see (Cossard and Pannell 2019)), has probably been important in facilitating evolutionary transitions from dioecy to hermaphroditism under conditions of mate limitation (Ehlers and Bataillon 2007; Crossman and Charlesworth 2014; Käfer et al. 2017). It is plausible that leaky dioecy may be adaptive by assuring reproductive success under pollen- or mate-limited conditions (Sultan 1987; Korpelainen 1998; Ghalambor et al. 2007), e.g., in colonising species, whose population densities (and consequently whose mate availability) may vary greatly; colonising species are thought to be particularly likely to benefit from plastic strategies in general (Baker 1965; Stebbins 1965; Sultan and Spencer 2002). However, to our knowledge the adaptiveness of leaky dioecy has not been demonstrated for any dioecious plant population. Indeed, leaky sex expression can be elicited by external stimuli such as temperature, drought, simulated herbivory or exogenous hormone application (Kuhn 1939; Westergaard 1958; Korpelainen 1998; Delph and Wolf 2005; Golenberg and West 2013), none of which suggest an obvious adaptive function. Importantly, there appears to be no empirical support to date for the possibility that leaky sex expression might be prompted by pollen or mate limitation, which would be more plausibly adaptive.

Here, we asked whether leaky sex expression in the dioecious, wind-pollinated annua herb *Mercurialis annua* is plastic, and whether it responds to variation in mate availability. *M. annua* is a particularly good model for addressing these questions. First, the dioecious populations of *M. annua* are known to have expanded their range recently from the eastern Mediterranean Basin into western Europe (Obbard et al. 2006; Gonzalez-Martinez et al. 2017), during which populations establishing at the colonizing front are likely to have been exposed to mate-limited conditions. Second, the species has a metapopulation structure and dynamic, with frequent population turnover and substantial fluctuations in population size from generation to generation (Eppley and Pannell 2007a; Dorken et al. 2017), conditions under which a leaky expression of the opposite sex would likely confer fitness benefits on an on-going basis, even

after a range expansion ended (Golenberg and West 2013). Third, previous work has shown that females of *M. annua* quickly become pollen-limited at low density (Hesse and Pannell 2011). Finally, inbreeding depression in western European populations is low (Eppley and Pannell 2009), perhaps as a result of the range expansion (Pujol et al. 2009; Gonzalez-Martinez et al. 2017), so that selfing by leaky individuals under pollen limitation would seem to be particularly likely to be beneficial (Wolf and Takebayashi 2004; Pannell 2008). Taken together, these features suggest that leaky sex expression in *M. annua*, if it occurs, would be adaptive (see Discussion). Our experiment involved growing females of *M. annua* in populations with or without males . We hypothesized that, under a scenario of adaptive leaky sex expression, females growing without males would be more likely to produce male flowers, and would produce more of them.

Material and methods

We established six experimental populations of dioecious *Mercurialis annua* in separate common gardens on the campus of the University of Lausanne and in gardens around the city. Three 'control' populations were established at a 1:1 sex ratio (90 males and 90 females), and three populations 'female-only' population comprised only females (180 females). Plants were established for the experiment from seeds pooled from 35 demes of a metapopulation in north-western Spain (Labouche and Pannell 2016), and seedlings were first raised for six weeks in a glasshouse before being transplanted into their experimental plots. After seven weeks of growth, we recorded the number and mass of male flowers and seeds for 35 to 50 females per population, and calculated the male and female reproductive efforts (MRE and FRE) by dividing the total male-flower mass and seed mass, respectively, by the total above-ground biomass.

We analyzed the expression of leaky sex expression by females in terms of (1) the proportion of leaky females in the population and (2) their average MRE. To calculate proportions, we defined a leaky female as one with an MRE greater than the 95 percentile MRE across the control populations. By this definition, an average 5% of females were identified as leaky across the three control populations. We chose the 95% cut-off for our definition because it can be easy to miss one or two small male flowers on a large plant, and for coherence with other work on leaky sex expression in the lab (Cossard and Pannell 2019). An analysis based on an absolute measure of leakiness (including all females with any male-flower production at all) yielded qualitatively similar results.

We compared the proportion of inconstant individuals between female-only and control populations using a generalized linear model with a binomial error distribution, and with population replicates included as a random factor. MRE and FRE were compared using permutation t-tests, with 100,000 bootstraps and two degrees of freedom (i.e., with population replicate as the unit of observation).

Results

There was a tendency for leaky females to be more frequent in female-only than control populations, but the difference was not significant (Figure 1a; P = 0.16). Nor could we detect any significant difference in the FRE between the two treatments (P = 0.35), although variance of FRE was greater in male-deprived populations (F-test P = 7.06 x10^{-7}; variance = 1.19 x10^{-3} and 4.83 x10^{-4} for female-only and control populations, respectively; Figure 1c). In stark contrast, the MRE of females in female-only populations was 2.33 times that of females in control replicates (P = 4.71 x10^{-3}; Figure 1b).

Discussion

Our results provide evidence for a plastic component to the expression of sex inconstancy in *M. annua*, with a slight tendency for more leaky females in populations lacking males and, significantly, females that on average produced more than twice as many male flowers in the absence of males than in their presence. Plastic responses to pollen limitation in plants have been recorded previously, in the context of the increased production of cleistogamous flowers and corresponding selfing rates in *Viola praemorsa* (Jones et al. 2013) and *Collomia grandiflora* (Albert et al. 2011). However, although leaky sex expression is a common feature of the reproduction of dioecious plants (Lloyd and Bawa 1984; Korpelainen 1998; Ehlers and Bataillon 2007; Cossard and Pannell 2019), our study provides demonstrates that such leakiness may be plastic and sensitive to mating opportunities. It is too early to speculate on the frequency of adaptive plasticity in leaky sex expression in plants more generally, but given that mate availability likely varies substantially in natural populations, it would seem to be a trait that dioecious populations should often evolve.

M. annua is a ruderal species that occupies disturbed habitat, and previous work has shown that its populations are subject to substantial fluctuations in population size and population turnover, with local extinctions and colonization by seed dispersal being key features of the

species' ecology (Pannell 1997; Eppley and Pannell 2007b; Dorken et al. 2017). In sparse populations of *M. annua*, female reproduction may be strongly pollen-limited (Hesse and Pannell 2011), and a capacity to produce male flowers and to self-fertilize under these conditions is likely to be adaptive, especially as inbreeding depression is low (Eppley and Pannell 2009). It is thus plausible that plasticity in leaky sex expression has evolved and/or has been maintained under conditions of fluctuating population size and density in metapopulations. *M. annua* came to occupy its broad range in western Europe via a recent range expansion from the eastern Mediterranean Basin (Obbard et al. 2006; Gonzalez-Martinez et al. 2017), and it is plausible that the repeated demographic bottlenecks that occurred during the range expansion also favored the maintenance of plasticity in leaky sex expression.

Ultimately, plasticity in sex expression in dioecious *M. annua* should be maintained as long as its benefits outweigh its costs. Potential costs of plasticity, which might constrain its evolution (reviewed in Van Kleunen and Fischer 2005), include that of acquiring accurate information about the state of the environment, and that of maintaining the machinery required to sense and respond to environmental cues. Despite considerable effort, however, it has been difficult to find any evidence for such costs (Auld et al. 2010). Rather, it seems more likely that the evolution of plasticity is held in check by so-called 'limits' associated with its deployment, such as the penalty paid by individuals that respond incorrectly to an unreliable signal, or the disadvantage of having to wait until a signal is perceived before expression of an appropriate phenotype (Auld et al. 2010; Murren et al. 2015). We do not know which of these limits might apply to plasticity in leaky sex expression, but both seem plausible. In short-lived species like *M. annua*, or in species with a short reproductive season, there might be strong disadvantages associated with delaying leaky sex expression until mating prospects have been perceived. Yet such delays seem inevitable if, as seems likely, the signal to which plants are responding is the actual deposition of pollen on their stigmas, or physiological signals arising from seed-filling or fruit-set that follow pollination.

Regional variation in the plasticity of sex expression in *M. annua* hints at possible limits to its adaptive expression in terms of signal reliability. Although leaky sex expression is plastic in dioecious *M. annua*, as this study shows, polyploid populations of the species in the Iberian Peninsula comprise females that always express a male function, i.e., they are canalized in their sex expression as monoecious individuals. The reason for this difference is not

immediately clear, because the two lineages are both colonizers of very similar disturbed habit (Buggs and Pannell 2007) and might both benefit from the plasticity we have reported here. However, whereas the absence of males from a population in the dioecious diploid lineage would be reliably perceived by females in terms of an absence of pollen on their stigmas, or low rates of fertilization of their ovules, monoecious 'females' of the polyploid lineage would receive pollen from other monoecious individuals even when males are absent, which occurs frequently (Pannell et al. 2014). Thus, although polyploid monoecious individuals would likely benefit from suppressing the production of male flowers when males are present, because males are so much better at dispersing pollen and outcross siring (Eppley and Pannell 2007a; Santos del Blanco et al. 2018), this strategy might be inaccessible to them for want of a suitably reliable signal.

In conclusion, our study has demonstrated that leaky sex expression in *M. annua* has a substantial plastic component, and our discussion above suggests that this strategy is likely adaptive. Consistent with this adaptive interpretation, Cossard et al. (2019) have recently shown that females of *M. annua* deprived of their mates, as in the female-only lines here, are indeed under strong positive selection for enhanced leaky sex expression, and respond to this selection by evolving enhanced leakiness. Our conjecture that plastic leakiness in sex expression in *M. annua* is adaptive would thus seem to be well-founded. Leaky sex expression in dioecious populations is almost certainly a necessary requirement for evolutionary transitions to occur from dioecy to hermaphroditism (Ehlers and Bataillon 2007; Crossman and Charlesworth 2014; Käfer et al. 2017). While we might expect such transitions to be slowed if leaky expression is strongly plastic, because plasticity diminished a trait's heritability (Falconer and Mackay 1996), it is tempting to speculate that environmentally induced induced leakiness, as found in diploid *M. annua*, might actually enhance the potential for the evolution of canalized monoecy or hermaphroditism via genetic assimilation (Waddington 1942, 1953, 1959; Pigliucci and Murren 2003; Price et al. 2003; West-Eberhard 2003; Ghalambor et al. 2007; Schlichting 2008).

References

Agrawal A. A. 1998. Induced responses to herbivory and increased plant performance. Science 279: 1201–1202.

Agrawal A. A. 1999. Induced responses to herbivory in wild radish: Effects on several herbivores and plant fitness. Ecology 80: 1713-1723.

Albert L. P., L. G. Campbell, and K. D. Whitney. 2011. Beyond simple reproductive assurance: cleistogamy allows adaptive plastic responses to pollen limitation. Int. J. Pl. Sc. 172: 862-869.

Auld J. R., A. A. Agrawal, and R. A. Relyea. 2010. Re-evaluating the costs and limits of adaptive phenotypic plasticity. Proceedings of the Royal Society B-Biological Sciences 277: 503-511.

Baker H. G. 1965. Characteristics and modes of origin of weeds. Pages 141-172 *in* H. G. Baker, and G. L. Stebbins, editors. The Genetics of Colonising Species. Academic Press, London.

Baker H. G. 1967. Support for Baker's Law - as a rule. Evolution 21: 853-856.

Banks J. A. 1997. Sex determination in the fern Ceratopteris. Trends Pl. Sci. 2: 175-180.

Baythavong B. S. 2011. Linking the spatial scale of environmental variation and the evolution of phenotypic plasticity: selection favors adaptive plasticity in fine-grained environments. Amer. Nat. 178: 75-87.

Baythavong B. S., and M. L. Stanton. 2010. Characterizing selection on phenotypic plasticity in response to natural environmental heterogeneity. Evolution 64: 2904-2920.

Bradshaw A. D. 1965. Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics 13: 115–155.

Buggs R. J. A., and J. R. Pannell. 2007. Ecological differentiation and diploid superiority across a moving ploidy contact zone. Evolution 61: 125-140.

Conover D. O., and E. T. Schultz. 1995. Phenotypic similarity and the evolutionary significance of countergradient variation. Trends 10: 248–252.

Cossard G., and J. R. Pannell. 2019. A functional decomposition of sex inconstancy in the dioecious, colonizing plant *Mercurialis annua*. Amer. J. Bot. 106: 722-732.

Crossman A., and D. Charlesworth. 2014. Breakdown of dioecy: models where males acquire cosexual functions. Evolution 68: 426-440.

Delph L. F. 2003. Sexual dimorphism in gender plasticity and its consequences for breeding system evolution. Evol. Devel. 5: 34-39.

Delph L. F., and D. E. Wolf. 2005. Evolutionary consequences of gender plasticity in genetically dimorphic breeding systems. New Phytol. 166: 119-128.

Diggle P. K. 1991. Labile sex expression in andromonoecious *Solanum hirtum*: floral development and sex determination. Amer. J. Bot. 78: 377–393.

Dobzhansky T. 1969. On Cartesian and Darwinian aspects of biology. Pages 165–178 *in* S. Morgenbesser, P. Suppes, and M. White, editors. Philosophy, Science, and Method. St. Martins Press, New York.

Dorken M. E., and S. C. H. Barrett. 2003. Gender plasticity in *Sagittaria sagittifolia* (Alismataceae), a monoecious aquatic species. Pl. Syst. Evol. 237: 99-106.

Dorken M. E., R. P. Freckleton, and J. R. Pannell. 2017. Small-scale and regional spatial dynamics of an annual plant with contrasting sexual systems. J. Ecol. 105: 1044-1057.

Dorn L. A., E. H. Pyle, and J. Schmitt. 2000. Plasticity to light cues and resources in Arabidopsis thaliana: testing for adaptive value and costs

. Evolution 54: 1982 –1994.

Dudley S. A., and J. Schmitt. 1995. Genetic differentiation in morphological responses to simulated foliage shade between populations of Impatiens capensis from open and woodland sites. Func. Ecol. 9: 655-666.

Dudley S. A., and J. Schmitt. 1996. Testing the adaptive plasticity hypothesis: Density-dependent selection on manipulated stem length in Impatiens capensis. Amer. Nat. 147: 445-465.

Ehlers B. K., and T. Bataillon. 2007. 'Inconstant males' and the maintenance of labile sex expression in subdioecious plants. New Phytol. 174: 194-211.

Eppley S. M., and J. R. Pannell. 2007a. Density-dependent self-fertilization and male versus hermaphrodite siring success in an androdioecious plant. Evolution 61: 2349-2359.

Eppley S. M., and J. R. Pannell. 2007b. Sexual systems and measures of occupancy and abundance in an annual plant: testing the metapopulation model. Amer. Nat. 169: 20-28.

Eppley S. M., and J. R. Pannell. 2009. Inbreeding depression in dioecious populations of the plant *Mercurialis annua*: comparisons between outcrossed progeny and the progeny of self-fertilized feminized males. Heredity 102: 600-608.

Falconer D. S., and T. F. C. Mackay. 1996. Introduction to Quantitative Genetics. 4 edition. Longman, Essex, England.

Ghalambor C. K., J. K. McKay, S. P. Carroll, and D. N. Reznick. 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. Func. Ecol. 21: 394-407.

Golenberg E. M., and N. W. West. 2013. Hormonal interactions and gene regulation can link monoecy and environmental plasticity to the evolution of dioecy in plants. Amer. J. Bot. 100: 1022-1037.

Gomulkiewicz R., and M. Kirkpatrick. 1992. Quantitative genetics and the evolution of reaction norms. Evolution 46: 390-411.

Gonzalez-Martinez S. C., K. Ridout, and J. R. Pannell. 2017. Range expansion compromises

adaptive evolution in an outcrossing plant. Curr. Biol. 27: 2544-2549.

Gotthard K., and S. Nylin. 1995. Adaptive plasticity and plasticity as an adaptation: a selective review of plasticity in animal morphology and life-history. Oikos 74: 3-17.

Harder L. D., and S. D. Johnson. 2005. Adaptive plasticity of floral display size in animal-pollinated plants. Proceedings of the Royal Society B-Biological Sciences 272: 2651-2657.

Hendry A. P. 2015. Key questions on the role of phenotypic plasticity in eco-evolutionary dynamics. J. Hered. 107: 1-17.

Hesse E., and J. R. Pannell. 2011. Density-dependent pollen limitation and reproductive assurance in a wind-pollinated herb with contrasting sexual systems J. Ecol. 99: 1531–1539.

Jones N. T., B. C. Husband, and A. S. MacDougall. 2013. Reproductive system of a mixed-mating plant responds to climate perturbation by increased selfing. Proceedings of the Royal Society B-Biological Sciences 280: 6.

Josephs E. B. 2018. Determining the evolutionary forces shaping G x E. New Phytol. 219: 31-36.

Käfer J., G. A. B. Marais, and J. R. Pannell. 2017. On the rarity of dioecy in flowering plants. Mol. Ecol. 26: 1225-1241.

Karban R., A. A. Agrawal, and M. Mangel. 1997. The benefits of induced defenses against herbivores. Ecology 78: 1351-1355.

Kenney A. M., J. K. McKay, J. H. Richards, and T. E. Juenger. 2014. Direct and indirect selection on flowering time, water-use efficiency (WUE, delta C-13), and WUE plasticity to drought in *Arabidopsis thaliana*. Ecol. Evol. 4: 4505-4521.

Korpelainen H. 1998. Labile sex expression in plants. Biological Reviews of the Cambridge Philosophical Society 73: 157-180.

Kuhn E. 1939. Selbstbestäubungen subdiöcischer Blütenpflanzen, ein neuer Beweis für die genetische Theorie der Geschlechtsbestimmung. Planta 30: 457-470.

Labouche A. M., and J. R. Pannell. 2016. A test of the size-constraint hypothesis for a limit to sexual dimorphism in plants. Oecologia 105: 197–208.

Lloyd D. G. 1972. Breeding systems in Cotula L. (Compositae, Anthemedae). I. The array of monoclinous and diclinous systems. New Phytol. 71: 1181-1194.

Lloyd D. G., and K. S. Bawa. 1984. Modification of the gender of seed plants in varying conditions. Evol. Biol. 17: 255-338.

Lopez S., and C. A. Dominguez. 2003. Sex choice in plants: facultative adjustment of the sex

ratio in the perennial herb *Begonia gracilis*. J. Evol. Biol. 16: 1177-1185.

Murren C. J., J. R. Auld, H. Callahan*, et al.* 2015. Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. Heredity 115: 293-301.

Nicotra A. B., and A. Davidson. 2010. Adaptive phenotypic plasticity and plant water use. Functional Plant Biology 37: 117-127.

Obbard D. J., S. A. Harris, and J. R. Pannell. 2006. Sexual systems and population genetic structure in an annual plant: testing the metapopulation model. Amer. Nat. 167: 354-366.

Pannell J. 1997. Variation in sex ratios and sex allocation in androdioecious *Mercurialis annua*. J. Ecol. 85: 57-69.

Pannell J. R. 2008. Consequences of inbreeding depression due to sex-linked loci for the maintenance of males and outcrossing in branchiopod crustaceans. Genetical Research 90: 73-84.

Pannell J. R., S. M. Eppley, M. E. Dorken, and R. Berjano. 2014. Regional variation in sex ratios and sex allocation in androdioecious *Mercurialis annua*. J. Evol. Biol. 27: 1467–1477.

Paquin V., and L. W. Aarssen. 2004. Allometric gender allocation in *Ambrosia artemisiifolia* (Asteraceae) has adaptive plasticity. Amer. J. Bot. 91: 430-438.

Pigliucci M. 2001. Phenotypic Plasticity: Beyond Nature and Nurture. The John Hopkins Univeristy Press, Baltimore.

Pigliucci M., and C. J. Murren. 2003. Genetic assimilation and a possible evolutionary paradox: can macroevolution sometimes be so fast as to pass us by? Evolution 57: 1455–1464.

Price T. D., A. Qvarnstrom, and D. E. Irwin. 2003. The role of phenotypic plasticity in driving genetic evolution. Proceedings of the Royal Society of London, Series B 270: 1433–1440.

Pujol B., S. R. Zhou, J. Sahchez-Vilas, and J. R. Pannell. 2009. Reduced inbreeding depression after species range expansion. Proceeding of the National Academy of Sciences of the United States of America 106: 15379-15383.

Releya R. A. 2002. Costs of phenotypic plasticity. Amer. Nat. 159.

Santos del Blanco L., E. Tudor, and J. R. Pannell. 2018. Low siring success of females with an acquired male function illustrates the legacy of sexual dimorphism in constraining the breakdown of dioecy. Ecol. Lett. 22: 486-497.

Schlichting C. D. 2008. Hidden reaction norms, cryptic genetic variation, and evolvability. Annals of the New York Academy of Sciences 1133: 187-203.

Schlichting C. D., and M. Pigliucci. 1998. Phenotypic Evolution: A Reaction Norm Perspective.

Sinauer, Sunderland, Mass.

Stebbins G. L. 1965. Colonizing species of the native California flora. Pages 173-191 *in* H. G. Baker, and G. L. Stebbins, editors. The Genetics of Colonising Species. Academic Press, London.

Sultan S. E. 1987. Evolutionary implications of phenotypic plasticity in plants. Evol. Biol. 21: 127-178.

Sultan S. E., and H. G. Spencer. 2002. Metapopulation structure favors plasticity over local adaptation. Amer. Nat. 160: 271–283.

Torices R., J. M. Gomez, and J. R. Pannell. 2018. Kin discrimination allows plants to modify investment towards pollinator attraction. Nature Communications 9: 6.

Van Kleunen M., and M. Fischer. 2005. Constraints on the evolution of adaptive phenotypic plasticity in plants. New Phytol. 165: 49–60.

Venkatasamy S., G. Khittoo, and S. Keeley. 2007. Leaky dioecy in *Diospyros* (Ebenaceae) endemic to the Island of Mauritius. Plant Ecology 189: 139-146.

Waddington C. H. 1942. Canalization of development and the inheritance of acquired characters. Nature 150: 563-565.

Waddington C. H. 1953. Genetic assimilation of an acquired character. Evolution 7: 118-126.

Waddington C. H. 1959. Canalization of develoment and genetic assimilation of acquired characters. Nature 183: 1654-1655.

Wagner M. R., and T. Mitchell-Olds. 2018. Plasticity of plant defense and its evolutionary implications in wild populations of *Boechera stricta*. Evolution 72: 1034-1049.

Weiner J. 2004. Allocation, plasticity and allometry in plants. Perspect Plant Ecol 6: 207-215.

West-Eberhard M. J. 2003. Developmental Plasticity and Evolution. Oxford University Press, New York.

Westergaard M. 1958. The mechanism of sex determination in dioecious plants. Advances in Genetics 9: 217-281.

Wolf D. E., and N. Takebayashi. 2004. Pollen limitation and the evolution of androdioecy from dioecy. Amer. Nat. 163: 122-137.

Figure 1. (a) Proportion of females with leaky sex expression, (b) male-reproductive effort (MRE) of females, and (c) female reproductive effort (FRE) of females in male-only (blue) and control populations (red). Means and standard errors are shown. ****: $0.001 < P < 0.01$.