Whole-chromosome hitchhiking driven by a male-killing endosymbiont

Simon H. Martin*[1,2], Kumar Saurabh Singh[3], Ian J. Gordon[4], Kennedy Saitoti Omufwoko[5,6],
Steve Collins[7], Ian A. Warren[2], Hannah Munby[2], Oskar Brattström[2], Walther Traut[8], Dino J.
Martins[5,6], David A. S. Smith[9], Chris D. Jiggins[2], Chris Bass[3], Richard H. ffrench-Constant[3]

[1]Institute of Evolutionary Biology, University of Edinburgh, Edinburgh EH9 3FL, United

Kingdom

[2]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, United Kingdom

[3]Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, TR10

9EZ, United Kingdom

[4]BirdLife International Kigali Office, Box 2527, Kigali Post Office, Kigali, Rwanda

[5]Mpala Research Centre, P O Box 555, Nanyuki, 10400 Kenya.

[6]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544,

USA

[7]African Butterfly Research Institute, Box 14308-0800, Nairobi, Kenya

[8]Institut für Biologie, Zentrum für medizinische Struktur- und Zellbiologie, Universität Lübeck,

Ratzeburger Allee 160, 23538 Lübeck, Germany

[9]Natural History Museum, Eton College, Windsor SL4 6DW, UK

*Correspondence to: simon.martin@ed.ac.uk

Abstract

Neo-sex chromosomes are found in many taxa, but the forces driving their emergence and

spread are poorly understood. The female-specific neo-W chromosome of the African monarch

(or queen) butterfly *Danaus chrysippus* presents an intriguing case study because it is restricted

to a single 'contact zone' population, involves a putative colour patterning supergene, and co-

occurs with infection by the the male-killing endosymbiont *Spiroplasma*. We investigated the

origin and evolution of this system using whole genome sequencing. We first identify the 'BC

supergene', a broad region of suppressed recombination across nearly half a chromosome, which

links two colour patterning loci. Association analysis suggests that the genes *yellow* and *arrow* in

this region control the forewing colour pattern differences between *D. chrysippus* subspecies. We

then show that the same chromosome has recently formed a neo-W that has spread through the

contact zone within ~2200 years. We also assembled the genome of the male-killing

Spiroplasma, and find that it shows perfect genealogical congruence with the neo-W, suggesting

that the neo-W has hitchhiked to high frequency as the male killer has spread through the

population. The complete absence of female crossing-over in the Lepidoptera causes whole-

chromosome hitchhiking of a single neo-W haplotype, carrying a single allele of the BC

supergene, and dragging multiple non-synonymous mutations to high frequency. This has created

a population of infected females that all carry the same recessive colour patterning allele, making

the phenotypes of each successive generation highly dependent on uninfected male immigrants.

Our findings show how hitchhiking can occur between the unlinked genomes of host and

endosymbiont, with dramatic consequences.

Introduction

Structural changes to the genome play an important role in evolution by altering the extent of recombination among loci. This is best studied in the context of chromosomal inversions that cause localised recombination suppression, and can be favoured by selection if they help to maintain clusters of co-adapted alleles (or 'supergenes') in the face of genetic mixing [1–4]. A greater extent of recombination suppression occurs in the formation of heteromorphic sex chromosomes, which can link sex-specific alleles similarly to supergenes [5]. However, suppressed recombination can also have costs. In particular, male-specific Y and female-specific W chromosomes can be entirely devoid of recombination, making them vulnerable to genetic hitchhiking and the accumulation of deleterious mutations through 'Muller's ratchet', which may explain their deterioration over time [6–8]. These contrasting benefits and costs of recombination suppression are of particular interest in the evolution of neo-sex chromosomes, which can form through fusion of autosomes to existing sex chromosomes. There is accumulating evidence that neo-sex chromosomes are common in animals [9–15], but the processes underlying their emergence, spread and subsequent evolution have not been widely studied. In particular, there are few studied examples of recently-formed neo-sex chromosomes that are not yet fixed in a species.

The African monarch (or queen) butterfly *Danaus chrysippus*, provides a unique test case for the causes and consequences of changes in genome architecture and recombination suppression. Like its American cousin (*D. plexippus*), it feeds on milkweeds and has bright colour patterns that warn predators of its distastefulness. However, within Africa *D. chrysippus* is divided into four subspecies with distinct colour patterns and largely distinct ranges (Fig. 1A). Predator learning should favour the maintenance of a single monomorphic warning in any single area. For this reason, researchers have long been puzzled by the large polymorphic contact zone in East and Central Africa where all four *D. chrysippus* subspecies meet and interbreed [16–18] (Fig. 1A). Crosses have shown that colour pattern differences between the subspecies are controlled by Mendelian autosomal loci, including the tightly linked 'B' and 'C' loci (putatively a 'BC supergene' [19]) that define three common forewing patterns [20,21] (Fig. 1A). However, crosses with females from the contact zone revealed that the BC chromosome has become sex linked, forming a neo-W that is unique to this population [19,22]. Since female meiosis is achiasmatic (it lacks crossing-over) in the Lepidoptera, the formation of a neo-W would

71   instantaneously cause perfect linkage, not just of the B and C loci, but of an entire non-

72   recombining chromosome, along with other maternally-inherited DNA.

73   What is particularly striking is that the presence of the neo-W coincides with infection by a

maternally-inherited 'male killer' endosymbiont related to *Spiroplasma ixodetis,* which kills male

offspring and leads to highly female-biased sex-ratios where infection is common [22–24]. The

combination of neo-W and male killing is expected to dramatically alter the inheritance and

evolution of the BC chromosome [22,25]: Infected females typically give rise to all-female

broods who should always inherit the same colour patterning allele on their neo-W, along with

the male-killer, while the other maternal allele is systematically eliminated in the dead sons (Fig.

1B), forming a genetic sink for all colour pattern alleles not on the neo-W. It has been suggested

that the restriction of male killing to females with the neo-W, and only in the region in which

hybridisation occurs between subspecies, may not be a coincidence [19,22,25–27]. However, the

genomic underpinnings of this system – the genetic controllers of colour pattern, the source and

spread of the neo-W, and its relationship with the male killer – have until now remained a

mystery. We generated a reference genome for *D. chrysippus* and used whole genome

sequencing of population samples to uncover the interconnected evolution of the BC supergene,

neo-W and *Spiroplasma.* Our findings reveal a recent whole-chromosome selective sweep

caused by hitchhiking between the host and endosymbiont genomes.

Fig 1. Geography and genetics of colour pattern. (**A**) Approximate ranges of the four subspecies of *D. chrysippus*, with the contact zone outlined. Sampling locations for each of the subspecies and the contact zone are indicated. Cartoon chromosomes show the genotypes of each subspecies at the A (white hindwing patch), B (brown background colour) and C (forewing tip) colour patterning loci, based on previous crosses [20]. Note the linkage of B and C, putatively forming a 'BC supergene' [19]. Two examples of heterozygotes that can be found in the contact zone are shown. Note that *Cc* heterozygotes can exhibit the *transiens* phenotype with white markings on the forewing with ~50% penetrance. (**B**) Model showing how fusion of the BC autosome to the W chromosome has produced a neo-W (purple) in contact zone females (top left), while males have two autosomal copies of the BC chromosome (top right). Daughters inherit the neo-W, while sons inherit the other BC chromosome haplotype from their mother. The latter allele is then lost due to male killing by *Spiroplasma*.

Results and Discussion

Identification of the BC supergene

We assembled a high quality draft genome for *D. chrysippus,* with a total length of 322 Megabases (Mb), a scaffold N50 length of 0.63 Mb, and a BUSCO [28] completeness score of 94% (S1 Table-S8 Table). We then further scaffolded the genome into a pseudo-chromosomal assembly based on homology with the *Heliconius melpomene* genome [29–31] accounting for known fusions that differentiate these species [9,30,32] (Fig. S1). We also re-sequenced 42 individuals representing monomorphic populations of each of the four subspecies and a polymorphic population from a known male-killing hotspot near Nairobi, in the contact zone (Fig 1A, S9 Table).

To identify the putative BC supergene, we scanned for genomic regions showing high

differentiation between the subspecies and an association with colour pattern. Genetic

differentiation (F_{ST}) and excessive divergence (d_{XY}) is largely restricted to a handful of broad

peaks, with a background level of approximately zero (Fig. 2A, S2 Fig, S3 Fig). This low

background level implies a nearly panmictic population across the continent. The effective

population size appears to be very large, as average genome-wide diversity at putatively neutral

4-fold degenerate 3^{rd} codon positions is 0.042, which is among the highest values reported for

animals [33,34]. The islands of differentiation that stand out from this background imply

selection for local adaptation maintaining particular differences between the subspecies, similar

to patterns seen between geographic races of *Heliconius* butterflies [35]. However, here the

peaks of differentiation are broad, covering several Mb, implying some mechanism of

recombination suppression such as inversions that differentiate the subspecies.

The inclusion of the polymorphic contact-zone samples, and the fact that three of the

subspecies each carry a unique colour pattern allele (Fig 1A), allowed us to identify particular

differentiated regions associated with the three major colour pattern traits. A ~3 Mb region on

chromosome 4 is associated with the white hindwing patch (A locus) and a ~5 Mb region on

chromosome 15 (hereafter chr15) is associated with both background orange/brown (B locus)

and the forewing black tip (C locus) (Fig 2A, S2 Fig). Below, we refer to this region on chr15,

which spans over 200 protein-coding genes, as the BC supergene [19], although we note that

additional associated SNPs on chromosome 22 suggest that background wing melanism may also

be influenced by other loci.

Clustering analysis based on genetic distances reveals three clearly distinct alleles at the

BC supergene (Fig 2C). This further supports the hypothesis of recombination suppression,

although a number of individuals show mosaic ancestry consistent with occasional

recombination (S4 Fig). The three main alleles correspond to the three common forewing

phenotypes, so we term these $BC^{chrysippus}$ (orange background with black forewing tip, formerly

bbcc), $BC^{dorippus}$ (orange without black tip, formerly *bbCC)*, and $BC^{orientis}$ (brown background with

black forewing tip, formerly *BBcc*) (Fig 2C). Fifteen of the twenty contact zone individuals are

heterozygous, carrying two distinct BC alleles, and a few carry putative recombinant alleles, as

do some of the southern African *orientis* individuals (S4 Fig). As shown previously, $BC^{dorippus}$

(which includes the dominant *C* allele) and *BC^orientis^* (which includes the dominant *B* allele) are

both dominant over the recessive *BC^chrysippus^* (Fig. 2C, S4 Fig).

Fig 2. Identification of the BC supergene on chromosome 15. (**A**) Pairwise genetic
differentiation (F_{ST}), plotted in 100 kb sliding windows with a step size of 20 kb across all
chromosomes. Three subspecies pairs with sample sizes ≥ 6 are shown (see legend in panel B),
revealing strong overlap in patterns of differentiation. Locations of SNPs most strongly
associated with the A, B and C loci (Wald test, 99.99% quantile) are plotted above in grey, brown
and black, respectively. See S2 Fig for a more detailed plot. (**B**) Expanded plot of F_{ST} across a 10
Mb portion of chromosome 15 (chr15). Note that the first ~6 Mb of the chromosome is not
shown due to complex structural variation (see main text). Scaffolds are indicated below the plot
in alternating shades. The locations of our most likely candidate genes for B (*yellow*) and C
(*arrow*) are indicated. (**C**) Allelic clustering on chr15 in six representative individuals: three
homozygotes and three heterozygotes (see S4 Fig for all individuals, and see panel B for
chromosome positions). Coloured blocks indicate 20 kb windows in which sequence haplotypes
could be assigned to one of three clusters based on pairwise genetic distances (see Methods for
details). Windows in grey indicate insufficient relative divergence to be assigned to a cluster and
white indicates missing data. Allele names reflect the subspecies from which they come. In
heterozygotes, the name of the dominant allele is in bold.

Although it can be challenging to identify particular functional mutations in regions of

suppressed recombination, the presence of some recombinant individuals allowed us to narrow

down candidate regions for the B and C loci. A cluster of SNPs most strongly associated with

background colour (B locus) is found just upstream of the gene *yellow*, and a phylogenetic

network for a 30 kb region around *yellow* groups individuals nearly perfectly by phenotype,

although some individuals classed as heterozygous were intermingled with homozygotes (S5

Fig). Differential expression of Yellow in *Drosophila* is associated with different levels of

melanism [36] and *yellow* knockouts in other butterflies show reduced melanin pigmentation

[37], making this a compelling candidate for the background colour polymorphism. The

strongest associations with forewing tip (C locus) occur at the gene *arrow*, and a phylogenetic

network for a 100 kb region around this gene similarly clusters individuals by phenotype (S5

Fig). In *Drosophila*, Arrow is essential for Wnt signalling in wing development [38]. Wnt

signalling is known to underlie variation in colour pattern in *Heliconius* butterflies [39] and

knock-out mutants for the Wnt ligand gene *WntA* in *D. plexippus* show a loss of pigmentation

[40]. This makes *arrow* a promising candidate for the C locus gene. While these genes represent

our best candidates, numerous strongly-associated SNPs occurred closer to other genes in this

region (S10 Table). Future studies will aim to narrow down the associated regions and

performing functional validation.

Irrespective of their precise mode of action, the patterns of association imply that the B and

C loci are ~1.6 Mb apart (S5 Fig), and would therefore be fairly loosely linked under normal

recombination. This physical distance translates to around 7.6 cM, assuming crossover rates

similar to those in *Heliconius* [31,41], whereas the estimated recombination distance between B

and C based on crosses is 1.9 cM [42]. Theory predicts that recombination suppression can be

favoured if it maintains linkage disequilibrium (LD) between co-adapted alleles in the face of

gene flow [1–4]. Our study is one of only a few cases in which it can be shown that alleles at

distinct loci that each influence a component of a complex trait are maintained in LD by

suppressed recombination [43,44].

It is likely that chromosomal rearrangements contribute to recombination suppression at

the BC supergene. Although our short-read data do not allow us to test directly for inversions,

they do reveal dramatic variation in sequencing coverage over the proximal end of the

chromosome. Comparison of coverage among individuals suggests a large (~5 Mb) polymorphic

insertion in this region that tends to occur in individuals carrying the $BC^{dorippus}$ allele (S6 Fig).

Synteny comparison with *H. melpomene* reveals that this insertion involves an expansion in copy

number of a region of several hundred kb. Comparison of copy numbers for two of the genes in

the expansion with several other species confirms that it is derived in *D. chrysippus* (S7 Fig).

The expansion appears to occur just a few kb from the coding region of *arrow* (S6 Fig), and is

also perfectly associated with the presence of the dominant *dorippus* phenotype (absence of

black forewing tip) (S7 Fig). It is possible that it has a causal effect on the phenotype by

influencing the expression of *arrow*, but it might also be simply linked to the causative mutation.

Either way, we suggest that this large structural change, which increases the length of the

chromosome by nearly a third, contributes to recombination suppression between the $BC^{dorippus}$

allele and other supergene alleles by interfering with chromosome pairing in heterozygotes.

A neo-W chromosome traps a single haplotype of chromosome 15 in contact zone females

Previous crossing experiments indicated that the BC chromosome has become sex-linked

in contact zone females [22]. To confirm this hypothesis using genetic tools, we created a 'cured

line' by treating a female from an all-female brood with tetracycline to eliminate *Spiroplasma*

and allow the survival of male offspring [23]. A cross using this female confirms perfect sex-

linkage of forewing phenotype (n=22, chi-square test p=0.00002; S8 Fig). We then used PCR

assays on a subsequent sibling cross from the cured line to confirm that maternal alleles for

chr15 segregate with sex (n=22, p<0.00003) whereas paternal alleles segregate randomly (n=22,

p=0.36; S8 Fig). These results exactly match the model (Fig. 1B) in which the BC supergene has

become linked to the W chromosome in females, but continues to segregate as an autosome in

males.

Although we were unable to definitively identify any scaffolds from the ancestral W

chromosome, which is likely to be highly repetitive, we can test whether chr15 shows the

expected hallmarks of a young neo-W, hypothesised to have formed through fusion to the

ancestral W [22]. Due to the complete absence of recombination in females, we expect that a

single fused haplotype of chr15 would be spreading in the population. Any unique mutations

specific to this haplotype should therefore occur at high frequency in females and be absent in

males. We scanned for such high-frequency female-specific mutations, and found them to be

abundant across the entire length of chr15 and nearly absent throughout the rest of the genome

(Fig 3A). At the individual level, we can clearly identify 15 females (14 collected in the contact

zone and the single 'cured line' female) that consistently share these high-frequency mutations

(S9 Fig). Genetic distance among these females in the colinear region of chr15 (outside the BC

supergene) is reduced, indicating that they all share a similar haplotype of the fused chromosome

(Fig. 3B).

Fig 3. Recent sweep of a young neo-W. (**A**) The number of high-frequency female-specific
mutations (> 20% in females and absent in males) in 100 kb sliding windows (20 kb step size).
(**B**) Distance-based phylogenetic network for the distal colinear region of chr15 outside of the
BC supergene reveals that most contact zone females carry the conserved neo-W haplotype.
Cartoons show how the colinear region of chr15 is outside of the BC supergene but would still
capture reduced divergence among individuals carrying a shared non-recombining neo-W. (**C**)
Boxplot comparing the density of heterozygous sites in 100 kb windows in the colinear region of
chr15 between wild type individuals from the contact zone (CZ) and those carrying the neo-W.
Cartoon chromosomes above the plot match those shown in panel B. A relative lack of elevated
heterozygosity in the neo-W lineage indicates a lack of divergence of the fused neo-W haplotype,
consistent with the fusion being recent. (**D**) Boxplot of nucleotide diversity (π) within each
population for the same colinear region of chr15. On the far right, π is shown for the haploid neo-
W haplotype specifically, based on partial sequences isolated from this haplotype (see Methods
and S10 Fig for details). The near absence of genetic diversity implies a very rapid spread of the
neo-W through the population.

The neo-W formed recently and spread rapidly

Genetic variation accumulated in the neo-W lineage since its formation can tell us about its

age. Sequence divergence between the neo-W and autosomal copies of chr15 (inferred from the

density of heterozygous sites in the colinear region of chr15 in females carrying the neo-W) is

not significantly different from that between the autosomal copies in 'wild type' individuals that

lack the fusion (Fig. 3C, Wilcoxon signed rank test, p=0.36, n=48 100 kb windows). This implies

that insufficient time has passed since the fusion event for significant accumulation of new

mutations. The limited divergence of the neo-W haplotype from the autosomal copy of chr15 in

each female makes it challenging to isolate. Nonetheless, by using diagnostic mutations that are

unique to, and fixed in the neo-W lineage, we were able to identify sequencing reads from the

shared haplotype and reconstruct a partial neo-W sequence for each female (S10 Fig). A dated

genealogy based on these sequences places the root of the neo-W lineage at ~2200 years (26,400

generations) ago (posterior mean = 2201, std. dev. = 318).

The neo-W is present in all but one of the contact zone females, implying a rapid spread

since its formation. This process is similar to a selective sweep of a beneficial mutation, except

that complete recombination suppression in females means that the sweep affects the entire

chromosome equally. Unlike a conventional sweep, it is not expected to eliminate genetic

diversity from the population as these females will also carry an autosomal copy of chr15

inherited from their father (Fig 1B). Indeed, we see a 20% reduction in overall nucleotide

diversity (π) on chr15 in females of the neo-W lineage (Fig 3D). However, when we consider

only the neo-W haplotype in each of these females we see a nearly complete absence of genetic

variation, with a π of 0.00007, two orders of magnitude lower than for autosomal copies of chr15

(0.0228) (Fig. 3D). These results further support a very recent and rapid spread of the neo-W.

The neo-W haplotype carries the recessive $BC^{chrysippus}$ allele at the BC supergene (S4 Fig).

However, previous work [22] shows that at the the focal sampling site in the contact zone, most

males are immigrants homozygous for the dominant $BC^{dorippus}$ allele, and the vast majority of

females (84%) are heterozygous $BC^{dorippus}/BC^{chrysippus}$, as expected if most inherit $BC^{dorippus}$ from

their father and $BC^{chrysippus}$ (on the neo-W) from their mother. The dominant *dorippus* phenotype is

therefore by far the most abundant in this population. Since aposematic colouration should be

under positive frequency dependent selection, it is highly unlikely that the spread of the neo-W

can be explained by selection on colour pattern, highlighting the question of what else might

have driven its spread.

Hitchhiking between the neo-W and *Spiroplama*

We hypothesised that the neo-W has spread as a result of co-inheritance with the male-

killing *Spiroplasma*, which is itself spreading through the population as a selfish element.

Experiments have suggested that infected female offspring have increased survival relative to

those from uninfected broods, possibly due to reduced competition for resources, although other

factors such as improved immunity have not been tested [45]. For this to drive the spread of the

neo-W, there would also need to be strict vertical inheritance of *Spiroplasma* down the female

line, such that it is always co-inherited with the neo-W.

We identified nine scaffolds making up the 1.75 Mb *Spiroplasma* genome in our *D.*

chrysippus assembly (S11 Fig). Infected individuals are clearly identifiable by mapping

resequencing reads to the *Spiroplasma* scaffolds (S11 Fig), and this was confirmed by PCR. As

predicted, all females in the neo-W lineage are infected (with the exception of the cured line

female, in which *Spiroplasma* had been eliminated). Moreover, all infected females fall into the

same mitochondrial clade (Fig 4A), consistent with matrilineal inheritance. To confirm that the

Spiroplasma is strictly vertically inherited and always associated with a single female lineage,

we used PCR assays for *Spiroplasma* and mitochondrial haplotype and expanded our sample size

to 158 individuals, including samples used in previous studies going back two decades

[19,23] (S12 Table, S12 Fig). This confirms the perfect association: 100% of infected individuals

(n=42) carry the same mitochondrial haplotype, and this haplotype is otherwise rare, occurring in

8% of uninfected individuals (n=116) (S12 Fig).

Like the neo-W, the *Spiroplasma* genomes carry limited variation among individuals ($\pi =$

0.0005), consistent with a single and recent outbreak of the endosymbiont. Although the lack of

variation makes it challenging to infer genealogies, our inferred maximum likelihood

genealogies for the neo-W and *Spiroplasma* are strikingly congruent (Fig 4B). The low bootstrap

support for multiple nodes is unsurprising given that these sequences descend from a recent

common ancestor, such that most nodes will be defined by only a few informative sites. This

does not weaken the support for congruence however, as the probability of two incorrectly

inferred topologies matching by chance is infinitesimally small. In a permutation test for

congruence between the two distance matrices [46], the observed level of congruence exceeds all

100,000 random permutations. There is therefore strong support for co-inheritance of the neo-W

and *Spiroplasma* [47].

Fig 4. Matrilineal inheritance causes coupling between neo-W and *Spiroplasma*. (A)
Maximum likelihood phylogeny for the whole mitochondrial genome. Individuals are coloured
according to population of origin (see Fig 1A), and those carrying the neo-W ('W') and
Spiroplasma ('S') are indicated (including one cured individual in which *Spiroplasma* was
eliminated). Females are indicated by circles and males by squares. (**B**) Maximum likelihood
phylogenies for the neo-W haplotype and *Spiroplasma* genome isolated from infected females.
Corresponding clades are shaded to indicate congruence. Note that two samples are excluded in
panel B: the cured sample, which lacked *Spiroplasma* due to tetracycline treatment, and one
infected female found to lack the neo-W. Whether the latter represents an ancestral state or
secondary loss requires further investigation. In all trees, nodes supported by more than 70 of
100 bootstrap replicates are indicated by circles.

The combined spread of three physically unlinked DNA molecules – the mitochondrial

genome, neo-W and *Spiroplasma* genome – constitutes a form of genetic hitchhiking, but is

facilitated by their strict matrilineal inheritance rather than physical linkage. We cannot entirely

rule out the possibility that the neo-W is contributing to this spread, or even driving it entirely,

through direct selection or meiotic drive. In theory this is testable by examining broods that carry

the neo-W but lack *Spiroplasma*. We raised 11 such broods in our cured line, and Smith

[48] reported 10 natural broods that showed sex-linked colour pattern and no male killing.

Across these 21 broods, totalling 528 adult offspring, 51% were female. This is far from

significantly different from the null expectation of 50% (binomial test p=0.7). However, we note

that in order to detect meiotic drive causing a 1% female bias with good power would require a

far larger samples size of >15,000. Given the evidence from other work showing that male

killers can spread rapidly as selfish elements [49], and the endosymbionts can convey other

fitness benefits [50], this remains the most parsimonious explanation in this case. Selfish

elements have been shown to drive hitchhiking of the mitochondrial genome or a portion of a

chromosome through a population and even across species boundaries [51–53]. Our findings

show how an entire chromosome can be captured in the same way. Hitchhiking may therefore be

of general importance in driving the spread of neo-sex chromosomes.

In *D. chrysippus*, it is currently unclear whether the neo-W or male-killer emerged first. It

is also unclear whether their co-occurrence in a single ancestor was simply a coincidence or

instead reflects some functional connection, such as the suggestion that the neo-W might confer

susceptibility to the male killer [22]. It is important to note that this is not the first time a neo-sex

chromosome has formed in this taxon. A fusion of chromosome 21 to the ancestral Z

chromosome occurred in an ancestor of all *Danaus* species, producing a neo-Z [9,32,54]. It is

speculated that a complementary fusion of chromosome 21 to the ancestral W also occurred

[9,54], but this is difficult to conclusively verify due to degradation of the W chromosome over

longer timescales. If this hypothesis of an ancient neo-W is correct, then the neo-W we describe

(W-chr15) might in fact be better described as a neo-neo-W (W-chr21-chr15). It is possible that

the spread of the original W-chr21 was also driven by hitchhiking with a selfish endosymbiont.

Genetic and phenotypic consequences of recombination suppression

Sex chromosome evolution in many other taxa involves the progressive spread of

recombination suppression outward from the a sex-determining locus [55]. By contrast, the

absence of crossing over in female meiosis means that a lepidopteran neo-W experiences

complete and immediate recombination suppression over its entire length. Butterfly W

chromosomes are therefore thought to be highly degenerated and repetitive, and to our

knowledge none have been successfully assembled to date. The young age of the *D. chrysippus*

neo-W therefore provides a rare opportunity to study the early evolutionary consequences of

recombination suppression across an entire chromosome. Two related processes could shape its

evolution: hitchhiking of pre-existing deleterious mutations that were initially rare in the

population [6], and accumulation of novel deleterious mutations due to reduced purging through

recombination and selection (i.e. Muller's Ratchet) [7].

As a proxy for the 'genetic load' of deleterious mutations in the population, we considered

P_n/P_s, the normalised ratio of non-synonymous to synonymous polymorphisms. Due to purifying

selection, non-synonymous polymorphisms are typically rare, and where they do occur the

mutant allele typically occurs at low frequency in the population [56]. When considering all

polymorphisms in the neo-W lineage, P_n/P_s for chr15 (excluding the BC supergene, to avoid

bias) is very slightly (~5%) higher than for other autosomes (Fig S13). Of 1000 bootstrap

replicates, 916 reproduced this bias, corresponding to a p-value of 0.084. However, when we

partition polymorphisms by allele frequency, we see that chr15 carries a large excess of non-

synonymous polymorphisms in the highest frequency class (i.e. minor allele at 50%), with a

P_n/P_s ratio >3 times larger than on other autosomes (S13 Fig). This holds across all 1000

bootstrap replicates (i.e. p<0.001). A change in the frequency distribution of non-synonymous

variants, without a significant change in their abundance, is best explained by hitchhiking of pre-

existing mildly-deleterious alleles that were initially rare in the population but were inadvertently

carried to high frequency along with the neo-W haplotype, and are therefore now found in all

females in this lineage. In fact, P_n/P_s for high-frequency polymorphisms on chr15 is higher than

that for singletons on autosomes (p=0.044), suggesting that accumulation of additional mildly-

deleterious alleles on the neo-W might have occurred early during its spread through the

population.

At the phenotypic level, perhaps counter-intuitively, the spread of a single supergene allele

on the neo-W has not caused homogenization of warning pattern among contact zone females,

and might in fact have the opposite effect. In locations where the neo-W and *Spiroplasma* are

nearly fixed, such as our sampling site near Nairobi, the high incidence of male killing implies

that the population is strongly shaped by immigrant males. Since the $BC^{chrysippus}$ allele on the neo-

W is universally recessive, daughters will tend to match the phenotype of their immigrant father.

However, because the neo-W is always transmitted to daughters, the paternal chr15 copy will be

lost to male killing after one generation, creating a genetic sink for immigrant male genes

[22] (S14 Fig). This combination of processes results in a female population that is highly

sensitive to the source of immigrants, which is known to fluctuate seasonally with monsoon

winds [16,57] (S14 Fig). This model leads to the testable prediction that seasonal fluctuations in

female phenotypes should be most dramatic where male killing is most abundant.

Future evolutionary trajectories

The future of the neo-W and *Spiroplasma* outbreak is uncertain. A lack of males could lead

to local extinctions [27], but extinction of the entire infected lineage is unlikely given the high

dispersal ability and seasonal influxes of males in the contact zone. Indeed, it is notable that

Spiroplasma infection has only been recorded within the contact zone population (with the

exception of a single South African female reported here, S12 Table), especially given theory

showing that male killers should spread very rapidly across the geographical range of a

panmictic population if they provide even a very weak selective advantage [49]. Future work

will investigate whether its spread might be curtailed by environmental factors, for example if

oviposition behaviour or host plant availability only leads to sibling competition (and consequent

benefits for all-female broods) under certain conditions [45]. An alternative and non-mutually-

exclusive hypothesis is that dispersal rates of infected females are strongly reduced. In other

systems, sex-ratio distortion has driven adaptive responses by the host, including changes to the

mating system [58] and the evolution of resistance to male-killing [59,60]. The absence of

evidence for these phenomena in *D. chrysippus* might simply reflect the recency of the male

killing outbreak. Eventually, we also expect the non-recombining neo-W to begin to degenerate

through further hitchhiking, gene loss and the spread of repetitive elements [8,55]. This young

system provides a rare opportunity to study how these phenomena unfold through time and

space.

Methods

Reference genome sequencing, assembly and annotation

Detailed methods for generation of the *D. chrysippus* reference genome are provided in S1

Text. Briefly, a draft assembly was generated using SPAdes [61] from a combination of paired-

end and mate-pair libraries of various insert sizes. Scaffolding and resolution of haplotypes was

performed using Redundans [62] and Haplomerger2 [63]. The assembly was annotated using a

combination of de-novo gene predictors yielding 16,654 protein coding genes. Mitochondrial

genomes were assembled using NOVOplasty [64].

Although we currently lack linkage information for further scaffolding, we generated a
pseudo-chromosomal assembly based on homology with the highly contiguous *Heliconius*
melpomene genome [30,31,65], adjusted for known karyotypic differences [9,30–32,54].
Although these genomes are diverged by approximately 90 million years, this homology-based
approach has been shown previously to be successful for reconstruct chromosomes in a
fragmented *D. plexippus* genome [9]. In total, 282 Mb (87% of the genome) could be confidently
assigned to chromosomes (S1 Fig).

Scaffolds representing the *Spiroplasma* genome were identified based on read depth of re-
mapped reads (S11 Fig) and homology to other available *Spirolasma* genomes. Annotation was
performed using the RAST server pipeline [66,67].

Population sample resequencing and genotyping

This study made use of 42 newly sequenced *D. chrysippus* individuals, as well as
previously sequenced individuals of the sister species, *D. petilia* (n=1), and next closest outgroup
D. gilippus (n=2) [68] (S9 Table). Details of DNA extraction, sequencing and genotyping are
provided in S1 Text. Briefly, DNA was extracted from thorax tissue and sequenced (paired-end,
150 bp) to a mean depth of coverage 20x or greater. Reads were mapped to the *D. chrysippus*
reference assembly using Stampy [69] v1.0.31 and genotyping was performed using GATK
version 3 [70,71]. Genotype calls were required to have an individual read depth ≥ 8, and
heterozygous and alternate allele calls were further required to have an individual genotype
quality (GQ) ≥ 20 for downstream analyses.

Genomic differentiation and associations with wing pattern

We used the fixation index (F_{ST}) to examine genetic differentiation across the genome
among the three subspecies for which we had six or more individuals sequenced. F_{ST} was
computed using the script popgenWindows.py (github.com/simonhmartin/genomics_general
release 0.2) with a sliding windows of 100 kb, stepping in increments of 20 kb. Windows with
fewer than 20,000 genotyped sites after filtering (see above) were ignored.

To identify SNPs associated with the three Mendelian colour pattern traits (i.e. the A B and
C loci) (Fig 1A), we used PLINK v1.9 [72] with the '--assoc' option, and provided quantitative
phenotypes of 0, 1 or 0.5 for assumed heterozygotes, which causes PLINK to use the Wald test
for quantitative traits. In addition to the quality and depth filters above, SNPs used for this

analysis were required to have genotypes for at least 40 individuals, a minor allele count of at

least 2, and to be heterozygous in no more than 75% of individuals. SNPs were also thinned to a

minimum distance of 100 bp.

To examine relationships among diploid individuals in specific regions of interest, we

constructed phylogenetic networks using the Neighbor-Net [73] algorithm, implemented in

SplitsTree [74]. Pairwise distances used for input were computed using the script distMat.py

(github.com/simonhmartin/genomics_general release 0.2).

Haplotype cluster assignment

To assign haplotypes to clusters in the BC supergene region, we first phased genotypes

using SHAPEIT2 [75,76] using SNPs filtered as for association analysis above, except with a

minor allele count of at least 4, and no thinning. Default parameters were used for phasing except

that the effective population size was set to 3×10^6. To minimise phasing switch errors, we

analysed each 20 kb window separately. Cluster assignment for both haplotypes from each

individual was based on average genetic distance to all haplotypes from each of three reference

groups: *D. c. dorippus*, *D. c. orientis*, or *D. c. alcippus* (which is representative of *chrysippus* as

they share the same alleles at the BC supergene). A haplotype was assigned to one of the three

groups if its average genetic distance to members of that group was less than 80% the average

distance to the other two groups, otherwise it was left as unassigned. Genetic distances were

computed using the script popgenWindows.py (github.com/simonhmartin/genomics_general

release 0.2).

Identification of neo-W specific sequencing reads

To identify females carrying the neo-W chromosome, we visualised the distribution of

female-specific derived mutations that occur at high-frequency. Allele frequencies were

computed using the script freq.py (github.com/simonhmartin/genomics_general release 0.2). Due

to the absence of female meiotic crossing over in Lepidoptera, all females carrying the neo-W

fusion should share a conserved chromosomal haplotype for the entire fused chromosome. To

isolate this shared fused haplotype from the autosomal copy, we first identified diagnostic

mutations as those that are present in a single copy in each member of the 'neo-W lineage' and

absent from all other individuals and outgroups. We then isolated the sequencing read pairs from

each of these females that carry the derived mutation (S10 Fig). This resulted in a patchy

alignment file, with a stack of read pairs over each diagnostic mutation. Based on these aligned

reads, we genotyped each individuals as described above, except here setting the ploidy level to

1, and requiring a minimum read depth of 3.

Diversity and divergence of the neo-W

The lack of recombination across the neo-W makes it possible to gain insights into its age.

Over time, mutations will arise that differentiate the neo-W from the recombining autosomal

copies of the chromosome. We estimated this divergence based on average heterozygosity in

females carrying the neo-W, and compared it to heterozygosity from contact-zone individuals

not carrying the neo-W. Heterozygosity was computed using the Python script

popgenWindows.py (github.com/simonhmartin/genomics_general release 0.2) focusing only on

the colinear portion of the chromosome (i.e. the distal portion from 11 Mb onwards), which is

outside of the BC supergene. Heterozygosity was computed in 100 kb windows, and windows

were discarded if they contained fewer than 20,000 sites genotyped in at least two individuals

from each population.

A recent spread of the neo-W through the population should also be detectable in the form

of strong conservation of the neo-W haplotype in all females that carry it (i.e. reduced genetic

diversity). We therefore computed nucleotide diversity (π) in 100 kb windows as above.

Reported values of π and heterozygosity represent the mean +- standard deviation across 100 kb

windows.

Genealogical analyses

We produced maximum likelihood trees for the mitochondrial genome, neo-W and

Spiroplasma genome, using PhyML v3 [77] with the GTR substitution model. Given the small

number of SNPs in both the neo-W and *Spiroplasma* genome, regions with inconsistent coverage

across individuals were excluded manually. Only sites with no missing genotypes were included.

We estimated the root node age for the neo-W using BEAST2 [78,79] version 2.5.1 with a

fixed clock model and an exponential population growth prior. For all other priors we used the

defaults as defined by BEAUti v2.5.1. We assumed a mutation rate of 2.9×10^{-9} per generation

based on a direct estimate for *Heliconius* butterflies [80] and 12 generations per year [81].

BEAST2 was run for 500,000,000 iterations, sampling every 50,000 generations, and we used

Tracer [82] version 1.7.1 to check for convergence of posterior distributions and compute the

root age after discarding a burn-in of 10%.

We tested for congruence between the neo-W and *Spiroplasma* trees using PACo [46],

which assesses the goodness-of-fit between host and parasite distance matrices, with 100,000

permutations. Distance matrices were computed using the script distMat.py

(github.com/simonhmartin/genomics_general release 0.2).

Analysis of synonymous and non-synonymous polymorphism

We computed P_n/P_s as as the ratio of non-synonymous polymorphisms per non-

synonymous site to synonymous polymorphisms per synonymous site. Synonymous and non-

synonymous sites were defined conservatively as 4-fold degenerate and 0-fold degenerate codon

positions, respectively, with the requirement that the other two codon positions are invariant

across the entire dataset. Only sites genotyped in all 15 females in the neo-W lineage were

considered, and counts were stratified by minor allele frequency using the script sfs.py

(github.com/simonhmartin/genomics_general release 0.1).

Butterfly rearing and molecular diagnostics

To generate a stock line that is cured of *Spiroplasma* infection, we treated caterpillars from

all-female broods with Tetracycline, following Jiggins et al. [23]. A 'cured line' was initiated

from a single treated female that had the heterozygous *Cc* '*transiens*' phenotype (Fig 1A). This

female was crossed to a wild male (homozygous *cc*) to test for sex linkage of phenotype. The

cured line was maintained through sibling crosses for six generations and the persistence of

males indicated that *Spiroplasma* had been eliminated.

We then applied a molecular test for sex linkage of chr15 using the F5 brood from the

cured line. We designed two separate PCR diagnostics based on SNPs segregating on chr15 to

distinguish between the two chromosomes of the male and the female parents (S11 Table). PCR

was performed using the Phusion HF Master Mix and HF Buffer (New England Biosciences).

To screen for *Spiroplasma* infection, we designed a PCR assay targeting the

glycerophosphoryl diester phosphodiesterase (GDP) gene (S11 Table). PCR was performed as

above. We confirmed the the sensitivity of this diagnostic by analyzing individuals of known

infection status based on whole genome sequencing (12 infected and 11 uninfected).

To investigate whether *Spiroplasma* infection was always associated with a single

mitochondrial haplotype, we designed a PCR RFLP for the Cytochrome Oxidase Subunit I (COI)

that differentiates the infected 'K' lineage from uninfected (S10 Table). PCR was performed as

above. A subset of products were verified by Sanger sequencing after purification using the
QIAquick PCR Purification Kit (Qiagen, USA).

Ethics Statement

Butterfly collection was performed under permit where relevant:
NACOSTI/P15/3290/3607, NACOSTI/P15/2403/3602 (National Commission for Science and
Technology, Kenya), MINEDUC/S&T/459/2017 (Ministry of Education, Rwanda),
EMDEP006/17 (Environmental Management Division, St Helena Government); and always with
permission of the land owner and/or local authorities. We also worked with local researchers
wherever possible, including authors DJM, KSS & IJG, and with the Lepidopterists Society of
Africa.

Data Availability

Raw genomic data and assemblies are available from GenBank (accession numbers to be
provided upon acceptance), all processed data files underlying all figures are available from the
Dryad Repository (accession number available upon acceptance). Scripts used for data analysis
are available from https://github.com/simonhmartin/genomics_general.

References

1.   Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation.
Genetics. 2006;173: 419–34. doi:10.1534/genetics.105.047985

2.   Bürger R, Akerman A. The effects of linkage and gene flow on local adaptation: A two-
locus continent–island model. Theor Popul Biol. Academic Press; 2011;80: 272–288.
doi:10.1016/J.TPB.2011.07.002

3.   Guerrero RF, Rousset F, Kirkpatrick M. Coalescent patterns for chromosomal inversions
in divergent populations. Philos Trans R Soc B Biol Sci. 2012;367: 430–438.
doi:10.1098/rstb.2011.0246

4.   Charlesworth D, Charlesworth B. Selection on recombination in clines. Genetics. 1979;91:
575–580.

5.   Charlesworth D. The status of supergenes in the 21st century: Recombination suppression
in Batesian mimicry and sex chromosomes and other complex adaptations. Evol Appl.
John Wiley & Sons, Ltd (10.1111); 2016;9: 74–90. doi:10.1111/eva.12291

6.   Rice WR. Genetic hitchhiking and the evolution of reduced genetic activity of the Y sex
chromosome. Genetics. 1987;116: 161–167.

7.   Charlesworth B. Model for evolution of Y chromosomes and dosage compensation. Proc
Natl Acad Sci U S A. 1978;75: 5618–22.

8.   Bachtrog D, Charlesworth B. The temporal dynamics of processes underlying Y
chromosome degeneration. Genetics. Genetics; 2008;179: 1513–25.
doi:10.1534/genetics.107.084012

9.   Mongue AJ, Nguyen P, Voleníková A, Walters JR. Neo-sex chromosomes in the monarch
butterfly, Danaus plexippus. G3. 2017;7: 3281–3294. doi:10.1534/g3.117.300187

10.   Bracewell RR, Bentz BJ, Sullivan BT, Good JM. Rapid neo-sex chromosome evolution
and incipient speciation in a major forest pest. Nat Commun. Springer US; 2017;8.
doi:10.1038/s41467-017-01761-4

11.   Pala I, Naurin S, Stervander M, Hasselquist D, Bensch S, Hansson B. Evidence of a neo-
sex chromosome in birds. Heredity. Nature Publishing Group; 2012;108: 264–272.
doi:10.1038/hdy.2011.70

12.   Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, et al. A role for a neo-sex
chromosome in stickleback speciation. Nature. Nature Publishing Group; 2009;461:
1079–1083. doi:10.1038/nature08441

13.   Nguyen P, Sykorova M, Sichova J, Kuta V, Dalikova M, Capkova Frydrychova R, et al.
Neo-sex chromosomes and adaptive potential in tortricid pests. Proc Natl Acad Sci.
2013;110: 6931–6936. doi:10.1073/pnas.1220372110

14.   Carabajal Paladino LZ, Provazníková I, Berger M, Bass C, Aratchige NS, López SN, et al.
Sex Chromosome Turnover in Moths of the Diverse Superfamily Gelechioidea. Barluenga
M, editor. Genome Biol Evol. Narnia; 2019;11: 1307–1319. doi:10.1093/gbe/evz075

15.   Steinemann M, Steinemann S. Enigma of Y chromosome degeneration: neo-Y and neo-X
chromosomes of Drosophila miranda a model for sex chromosome evolution. Genetica.
1998;102–103: 409–20.

16.   Smith DAS, Owen DF, Gordon IJ, Lowis NK. The butterfly Danaus chrysippus (L .) in
East Afiica : polymorphism and morph-ratio clines within a complex , extensive and
dynamic hybrid zone. Zool J Linn Soc. 1997;120: 51–78.

17.   Smith DAS. Heterosis, epistasis and linkage disequilibrium in a wild population of the
polymorphic butterfly Danaus chrysippus (L .). Zool J Linn Soc. 1980;69: 87–109.

18.   Smith D a S, Owen DF, Gordon IJ, Owiny AM. Polymorphism and evolution in the
butterfly Danaus chrysippus (L.) (Lepidoptera: Danainae). Heredity. 1993;71: 242–251.
doi:10.1038/hdy.1993.132

19. Smith DAS, Gordon IJ, Allen JA. Reinforcement in hybrids among once isolated
semispecies of Danaus chrysippus (L.) and evidence for sex chromosome evolution. Ecol
Entomol. 2010;35: 77–89. doi:10.1111/j.1365-2311.2009.01143.x

20. Smith DAS. Genetics of Some Polymorphic Forms of the African Butterfly Danaus
chrysippus L. (Lepidoptera: Danaidae). Insect Syst Evol. Brill; 1975;6: 134–144.
doi:10.1163/187631275X00235

21. Clarke CA, Sheppard PM, Smith AG. genetics of fore and hindwing colour in crosses
between Danaus chrysippus from Australia and from Sierra Leone (Danaidae). Lepid Soc
J. 1973;

22. Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, et al. A neo-W
chromosome in a tropical butterfly links colour pattern, male-killing, and speciation. Proc
R Soc B Biol Sci. 2016;283: 20160821. doi:10.1098/rspb.2016.0821

23. Jiggins FM, Hurst GD, Jiggins CD, v d Schulenburg JH, Majerus ME. The butterfly
Danaus chrysippus is infected by a male-killing Spiroplasma bacterium. Parasitology.
2000;120 (Pt 5: 439–46. doi:10.1017/S0031182099005867

24. Herren JK, Gordon I, Holland PWH, Smith D. The butterfly Danaus chrysippus
(Lepidoptera: Nymphalidae) in Kenya is variably infected with respect to genotype and
body size by a maternally transmitted male-killing endosymbiont (Spiroplasma). Int J
Trop Insect Sci. 2007;27: 62. doi:10.1017/S1742758407818327

25. Gordon IJ, Ireri P, Smith DAS. Hologenomic speciation: Synergy between a male-killing
bacterium and sex-linkage creates a "magic trait" in a butterfly hybrid zone. Biol J Linn
Soc. 2014;111: 92–109. doi:10.1111/bij.12185

26. Lushai G, Allen JA, Goulson D, Maclean N, Smith DAS. The butterfly Danaus chrysippus
(L.) in East Africa comprises polyphyletic, sympatric lineages that are, despite behavioural
isolation, driven to hybridization by female-biased sex ratios. Biol J Linn Soc. 2005;86:
117–131. doi:10.1111/j.1095-8312.2005.00526.x

27. Idris E, Saeed M. Hassan S. Biased sex ratios and aposematic polymorphism in African
butterflies: A hypothesis. Ideas Ecol Evol. 2013;6: 5–16. doi:10.4033/iee.2013.6.2.n

28. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. Oxford University Press; 2015;31: 3210–3212.
doi:10.1093/bioinformatics/btv351

29. Heliconius Genome Consortium T. Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. Nature. 2012;487: 94–8. doi:10.1038/nature11041

30. Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. Major
Improvements to the Heliconius melpomene Genome Assembly Used to Confirm 10
Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. G3. 2016;6: 695–
708. doi:10.1534/g3.115.023655

31. Davey JW, Barker SL, Rastas PM, Pinharanda A, Martin SH, Durbin R, et al. No evidence
for maintenance of a sympatric Heliconius species barrier by chromosomal inversions.
Evol Lett. 2017;1: 138–154. doi:10.1002/evl3.12

32. Ahola V, Lehtonen R, Somervuo P, Salmela L, Koskinen P, Rastas P, et al. The Glanville
fritillary genome retains an ancient karyotype and reveals selective chromosomal fusions
in Lepidoptera. Nat Commun. 2014;5: 1–9. doi:10.1038/ncomms5737

33. Leffler EM, Bullaughey K, Matute DR, Meyer WK, Ségurel L, Venkat A, et al. Revisiting
an old riddle: what determines genetic diversity levels within species? PLoS Biol.
2012;10: e1001388. doi:10.1371/journal.pbio.1001388

34. Mackintosh A, Laetsch DR, Hayward A, Charlesworth B, Waterfall M, Vila R, et al. The
determinants of genetic diversity in butterflies. Nat Commun. Nature Publishing Group;
2019;10: 3466. doi:10.1038/s41467-019-11308-4

35. Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, Simpson F, et al.
Genome-wide evidence for speciation with gene flow in Heliconius butterflies. Genome
Res. 2013;23: 1817–1828. doi:10.1101/gr.159426.113

36. Wittkopp PJ, Vaccaro K, Carroll SB. Evolution of yellow Gene Regulation and
Pigmentation in Drosophila. Curr Biol. Cell Press; 2002;12: 1547–1556.
doi:10.1016/S0960-9822(02)01113-2

37. Zhang L, Martin A, Perry MW, van der Burg KRL, Matsuoka Y, Monteiro A, et al.
Genetic Basis of Melanin Pigmentation in Butterfly Wings. Genetics. Genetics; 2017;205:
1537–1550. doi:10.1534/genetics.116.196451

38. Rives AF, Rochlin KM, Wehrli M, Schwartz SL, DiNardo S. Endocytic trafficking of
Wingless and its receptors, Arrow and DFrizzled-2, in the Drosophila wing. Dev Biol.
Academic Press; 2006;293: 268–283. doi:10.1016/j.ydbio.2006.02.006

39. Martin A, Papa R, Nadeau NJ, Hill RI, Counterman BA, Halder G. Diversification of
complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. Proc
Natl Acad Sci U S A. 2012;109: 12632–12637. doi:doi.org/10.1073/pnas.1204800109

40. Mazo-Vargas A, Concha C, Livraghi L, Massardo D, Wallbank RWR, Zhang L, et al.
Macroevolutionary shifts of WntA function potentiate butterfly
wing-pattern diversity. Proc Natl Acad Sci. 2017;114: 10701 LP – 10706.

41.　Van Belleghem SM, Rastas P, Papanicolaou A, Martin SH, Arias CF, Supple MA, et al.
Complex modular architecture around a simple toolkit of wing pattern genes. Nat Ecol
Evol. 2017;1: 0052. doi:10.1038/s41559-016-0052

42.　Smith DAS. Evidence for autosomal meiotic drive in the butterfly Danaus chrysippus L.
Heredity. 1976;36: 139–142.

43.　Coughlan JM, Willis JH. Dissecting the role of a large chromosomal inversion in life
history divergence throughout the *Mimulus guttatus* species complex. Mol Ecol. John
Wiley & Sons, Ltd (10.1111); 2019;28: 1343–1357. doi:10.1111/mec.14804

44.　Lee CR, Wang B, Mojica JP, Mandáková T, Prasad KVSK, Goicoechea JL, et al. Young
inversion with multiple linked QTLs under selection in a hybrid zone. Nat Ecol Evol.
2017;1. doi:10.1038/s41559-017-0119

45.　Gordon IJ, Ireri P, Smith DAS. Preference for isolated host plants facilitates invasion of
Danaus chrysippus (Linnaeus, 1758) (Lepidoptera: Nymphalidae) by a bacterial male-
killer Spiroplasma. Austral Entomol. 2015;54: 210–216. doi:10.1111/aen.12113

46.　Balbuena JA, Míguez-Lozano R, Blasco-Costa I. PACo: A Novel Procrustes Application
to Cophylogenetic Analysis. Moreau CS, editor. PLoS One. Public Library of Science;
2013;8: e61048. doi:10.1371/journal.pone.0061048

47.　Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, et al.
Population Genomics of the Wolbachia Endosymbiont in Drosophila melanogaster. PLoS
Genet. 2012;8. doi:10.1371/journal.pgen.1003129

48.　Smith DAS. African Queens and Their Kin: A Darwinian Odyssey. Taunton, UK:
Brambleby Books; 2014.

49.　Duplouy A, O'Neill SL. Rapid spread of male-Killing Wolbachia in the butterfly
Hypolimnas bolina. Evol Biol - Concepts, Mol Morphol Evol. 2010;23: 209–227.
doi:10.1007/978-3-642-12340-5_13

50.　Hurst GDD, Hutchence KJ. Host defence: Getting by with a little help from our friends.
Curr Biol. Elsevier Ltd; 2010;20: R806–R808. doi:10.1016/j.cub.2010.07.038

51.　Jiggins FM. Male-killing Wolbachia and mitochondrial DNA: selective sweeps, hybrid
introgression and parasite population dynamics. Genetics. 2003;164: 5–12.

52.　Hurst GDD, Jiggins FM. Problems with mitochondrial DNA as a marker in population,
phylogeographic and phylogenetic studies: The effects of inherited symbionts. Proc R Soc
B Biol Sci. 2005;272: 1525–1534. doi:10.1098/rspb.2005.3056

53.　Palopoli MF, Wu CI. Rapid evolution of a coadapted gene complex: Evidence from the
segregation Distorter (SD) system of meiotic drive in Drosophila melanogaster. Genetics.
1996;143: 1675–1688.

54.   Traut W, Ahola V, Smith DAS, Gordon IJ, Ffrench-Constant RH. Karyotypes versus
699   Genomes: The Nymphalid Butterflies Melitaea cinxia, Danaus plexippus, and D.
700   chrysippus. Cytogenet Genome Res. 2018;153: 46–53. doi:10.1159/000484032

701   55.   Wright AE, Dean R, Zimmer F, Mank JE. How to make a sex chromosome. Nature
702   Communications. Nature Publishing Group; 2016. p. 12087. doi:10.1038/ncomms12087

703   56.   Fay JC, Wyckoff GJ, Wu C-I. Positive and Negative Selection on the Human Genome.
704   Genetics. 2001;158: 1227–1234.

705   57.   Smith DAS, Owen DF. Colour genes as markers for migratory activity: The butterfly
706   Danaus chrysippus in Africa. Oikos. 1997;78: 127–135.

707   58.   Jiggins FM, Hurst GDD, Majerus MEN. Sex-ratio-distorting Wolbachia causes sex-role
708   reversal in its butterfly host. Proc R Soc B Biol Sci. 2000;267: 69–73.
709   doi:10.1098/rspb.2000.0968

710   59.   Hornett EA, Charlat S, Duplouy AMR, Davies N, Roderick GK, Wedell N, et al.
711   Evolution of male-killer suppression in a natural population. PLoS Biol. 2006;4: 1643–
712   1648. doi:10.1371/journal.pbio.0040283

713   60.   Wilfert L, Jiggins FM. The dynamics of reciprocal selective sweeps of host resistance and
714   a parasite counter-adaptation in Drosophila. Evolution. Wiley/Blackwell (10.1111);
715   2013;67: 761–773. doi:10.1111/j.1558-5646.2012.01832.x

716   61.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A
717   New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J
718   Comput Biol. Mary Ann Liebert, Inc. 140 Huguenot Street, 3rd Floor New Rochelle, NY
719   10801 USA ; 2012;19: 455–477. doi:10.1089/cmb.2012.0021

720   62.   Pryszcz LP, Gabaldón T. Redundans: an assembly pipeline for highly heterozygous
721   genomes. Nucleic Acids Res. Oxford University Press; 2016;44: e113–e113.
722   doi:10.1093/nar/gkw294

723   63.   Huang S, Kang M, Xu A. HaploMerger2: rebuilding both haploid sub-assemblies from
724   high-heterozygosity diploid genome assembly. Berger B, editor. Bioinformatics. Oxford
725   University Press; 2017;33: 2577–2579. doi:10.1093/bioinformatics/btx220

726   64.   Dierckxsens N, Mardulyn P, Smits G. NOVOPlasty: de novo assembly of organelle
727   genomes from whole genome data. Nucleic Acids Res. Oxford University Press; 2016;45:
728   gkw955. doi:10.1093/nar/gkw955

729   65.   Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.
730   Butterfly genome reveals promiscuous exchange of mimicry adaptations among species.
731   Nature. Nature Publishing Group; 2012;487: 94–8. doi:10.1038/nature11041

66.   Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, et al. The SEED and the
Rapid Annotation of microbial genomes using Subsystems Technology (RAST). Nucleic
Acids Res. Oxford University Press; 2014;42: D206–D214. doi:10.1093/nar/gkt1226

67.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server:
Rapid Annotations using Subsystems Technology. BMC Genomics. BioMed Central;
2008;9: 75. doi:10.1186/1471-2164-9-75

68.   Zhan S, Zhang W, Niitepõld K, Hsu J, Haeger JF, Zalucki MP, et al. The genetics of
monarch butterfly migration and warning colouration. Nature. 2014;514: 317–21.
doi:10.1038/nature13812

69.   Lunter G, Goodson M. Stampy: a statistical algorithm for sensitive and fast mapping of
Illumina sequence reads. Genome Res. 2011;21: 936–9. doi:10.1101/gr.111120.110

70.   DePristo M a, Banks E, Poplin R, Garimella K V, Maguire JR, Hartl C, et al. A framework
for variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet. 2011;43: 491–8. doi:10.1038/ng.806

71.   Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A,
et al. From fastQ data to high-confidence variant calls: The genome analysis toolkit best
practices pipeline. Curr Protoc Bioinforma. Hoboken, NJ, USA: John Wiley & Sons, Inc.;
2013;43: 11.10.1-11.10.33. doi:10.1002/0471250953.bi1110s43

72.   Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. Gigascience. Narnia; 2015;4:
7. doi:10.1186/s13742-015-0047-8

73.   Bryant D, Moulton V. Neighbor-Net: An Agglomerative Method for the Construction of
Phylogenetic Networks. Mol Biol Evol. 2003;21: 255–265. doi:10.1093/molbev/msh018

74.   Huson DH, Bryant D. Application of Phylogenetic Networks in Evolutionary Studies. Mol
Biol Evol. 2006;23: 254–267. doi:10.1093/molbev/msj030

75.   Delaneau O, Zagury J-F, Marchini J. Improved whole-chromosome phasing for disease
and population genetic studies. Nat Methods. 2013;10: 5–6. doi:10.1038/nmeth.2307

76.   Delaneau O, Marchini J, Zagury JF. A linear complexity phasing method for thousands of
genomes. Nat Methods. 2012;9: 179–181. doi:10.1038/nmeth.1785

77.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst Biol. 2010;59: 307–21. doi:10.1093/sysbio/syq010

78.   Bouckaert RR. DensiTree: making sense of sets of phylogenetic trees. Bioinformatics.
2010;26: 1372–3. doi:10.1093/bioinformatics/btq110

79.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: A
Software Platform for Bayesian Evolutionary Analysis. Prlic A, editor. PLoS Comput
Biol. Public Library of Science; 2014;10: e1003537. doi:10.1371/journal.pcbi.1003537

80.   Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, et al.
Estimation of the Spontaneous Mutation Rate in Heliconius melpomene. Mol Biol Evol.
2015;32: 239–243. doi:10.1093/molbev/msu302

81.   Owen DF, Chanter DO. Population biology of tropical African butterflies. Sex ratio and
genetic variation in Acraea encedon. J Zool. 1969;157: 345–374. doi:10.1111/j.1469-
7998.1969.tb01707.x

82.   Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in
Bayesian Phylogenetics Using Tracer 1.7. Susko E, editor. Syst Biol. 2018;67: 901–904.
doi:10.1093/sysbio/syy032

Acknowledgements

We are grateful to Godfrey Amoni Etelej, Laura Hebberecht-Lopez and Glennis Julian for
support with butterfly rearing. We thank Jenny York, Frank Jiggins and Deborah Charlesworth
helpful comments. This work was funded by European Research Council (ERC) European Union
Horizon 2020 research and innovation programme grant 646625 (CB), ERC grant 339873 (CDJ,
SHM), National Geographic Society Research Grant WW-138R-17 (IJG), and a Royal Society
University Research Fellowship (SHM). The funders had no role in study design, data collection
and analysis, decision to publish, or preparation of the manuscript.