Assessing genomic diversity and signatures of selection in Original Braunvieh cattle using whole-genome sequencing data

Meenu Bhati[1, *], Naveen Kumar Kadri[1], Danang Crysnanto[1], Hubert Pausch[1]

[1]Animal Genomics, ETH Zürich, Zürich, Switzerland

Authors and email addresses:

Meenu Bhati: meenu.bhati@usys.ethz.ch

Naveen Kumar Kadri: naveen.kadri@usys.ethz.ch

Danang Crysnanto: danang.crysnanto@usys.ethz.ch

Hubert Pausch: hubert.pausch@usys.ethz.ch

[*]Corresponding author

Abstract

Background: Local cattle breeds represent an important source of genetic variation because they might carry alleles that enable them to adapt to harsh environments and scarce food conditions. One such local Swiss cattle breed is Original Braunvieh (OB) which is used for beef and milk production in alpine areas. Using whole-genome sequencing data of 49 key ancestors, we characterize genomic diversity, genomic inbreeding, and signatures of selection in Swiss OB cattle at nucleotide resolution.

Results: We annotated 15.7 million SNPs and 1.5 million Indels including 10,738 and 2,763 missense deleterious and high impact variants, respectively, that were discovered in 49 OB key ancestors. Six Mendelian trait-associated variants that were previously detected in breeds other than OB, segregated in the sequenced key ancestors including variants causal for recessive xanthinuria and albinism. The average nucleotide diversity (1.6×10^{-3}) was higher in OB than many mainstream European cattle breeds. Accordingly, the average genomic inbreeding ($F_{ROH}=0.14$) was relatively low in OB cattle. However, genomic inbreeding increased in recent years due to a higher number of long (> 1 Mb) runs of homozygosity in cattle born in more recent generations. Using two complementary approaches, i.e., composite likelihood ratio test and integrated haplotype score, we identified 95 and 162 genomic regions encompassing 136 and 157 protein-coding genes, respectively, that showed evidence ($p < 0.005$) of past and ongoing selection. These selection signals were enriched for quantitative trait loci related to beef traits including meat quality, feed efficiency and body weight and pathways related to blood coagulation, nervous and sensory stimulus.

Conclusions: We provide a comprehensive overview of genomic variation segregating in Swiss OB cattle. Using whole-genome sequencing data, we observe higher genomic diversity and less inbreeding in OB than many European mainstream cattle breeds. Footprints of selection were detected in genomic regions that are possibly relevant for meat

quality and adaptation to harsh environments. Considering the low population size and increasing rates of genomic inbreeding in the past generations, the implementation and adoption of genomic mating programs seems warranted to maintain genetic diversity in the Swiss OB cattle population.

Introduction

Following the domestication of cattle, both natural and artificial selection led to the formation of breeds with distinct phenotypic characteristics including morphological, physiological and adaptability traits [1]. An increasing demand for animal-based food products required intensified selection of cattle with high milk and beef production ability in the past decades. The selection of animals from specialized beef and dairy breeds has resulted in a sharp decline in population size of many local breeds which are primarily used for beef and milk production under extensive conditions [2, 3]. Local breeds might carry alleles that enable them to adapt to harsh environments and scarce food conditions. Thus, characterizing and maintaining genetic diversity in local breeds is important to produce beef and milk under changing environmental conditions [4, 5].

With the advancement of high-throughput genotyping technology, millions of cattle have been genotyped using low, medium or high-density SNP arrays [6, 7]. The availability of dense genotype data is a prerequisite to characterize genomic diversity and detect signatures of selection within and between cattle breeds [8–14]. Moreover, genome-wide association testing between dense array-called genotypes and a broad spectrum of phenotypes revealed thousands of quantitative trait loci (QTL) for complex and monogenic traits [15]. Most of the widely used bovine SNP microarrays were designed in a way that they yield reliable genotypes at genetic markers that are polymorphic in mainstream breads [16]. Ascertainment bias is inherent in the resulting genotype data because rare, breed-specific, and less-accessible genetic markers are underrepresented in most microarrays [17]. This limitation causes observed allele frequency distributions to deviate from expectations which can distort population genetics estimates [18–20]. With an increasing number of individuals being whole-genome sequenced, it has become possible to discover sequence variant genotypes at population scale [21]. While sequence variant

genotypes might be biased toward the reference allele, this reference bias is less of a concern when the sequencing coverage is high [22].

Recently, we sequenced the genomes of 49 key ancestors of the Swiss Original Braunvieh (OB) cattle breed to an average fold-coverage of 12.75 per animal and discovered more than 17 million variants using a genome graph-based sequence variant genotyping approach [23]. OB cattle have been bred in Switzerland for centuries. However, in the early 1960s, cattle breeders began inseminating OB cows with semen from US Brown Swiss sires to increase milk yield, reduce calving difficulties and to improve mammary gland morphology of the Swiss OB population [24]. The extensive cross-breeding of OB cows with Brown Swiss sires decreased the number of female OB calves entering the herdbook to less than 2,000 by the mid of the 1990s (Additional file 1). Since then, the Swiss OB population is slowly but steadily growing due to governmental subsidies. In spite of the small population size, pedigree- and microarray-derived estimates of population genetic parameters indicate that genetic diversity is higher in OB than many mainstream breeds likely due to the use of many sires in natural mating [24, 25].

In the present study, we analyze whole-genome sequence variant genotypes of 49 key ancestors of the Swiss OB cattle population to assess genomic diversity and detect signatures of past or ongoing selection in the breed. Moreover, we provide a detailed overview of genomic inbreeding in the population using runs of homozygosity.

Material and Methods

Sequence variant genotyping

We considered genotypes at 17,303,689 biallelic variants (15,722,811 SNPs and 1,580,878 Indels) that were discovered and genotyped previously [23] in the autosomes of 49 key ancestors of the OB population using a genome graph-based sequence variant genotyping

approach [26]. In brief, 49 OB cattle were sequenced at between 6 and 38-fold genome coverage using either Illumina HiSeq 2500 (30 animals) or Illumina HiSeq 4000 (19 animals) instruments. The sequencing reads were filtered and subsequently aligned to the UMD3.1 assembly of the bovine genome [27] using the mem-algorithm of the Burrows Wheeler Aligner (BWA) software package [28]. Single nucleotide and short insertion and deletion polymorphisms were discovered and genotyped using the Graphtyper software [26]. Following recommended filtration criteria (see Crysnanto et al. [23] for more details), 15,722,811 SNPs and 1,580,878 Indels were retained for subsequent analyses. Beagle [29] phasing and imputation was applied to improve the primary genotype calls from Graphtyper and infer missing genotypes. Unless stated otherwise, imputed genotypes were considered for all subsequent analyses, because Beagle imputation considerably improved the primary genotype calls particularly in samples that have been sequenced at low coverage [23].

Variant annotation and evaluation

Functional consequences of 15,722,811 SNPs and 1,580,878 Indels were predicted according to the Ensembl (release 91) annotation of the bovine genome assembly UMD3.1 using the Variant Effect Predictor tool (VEP v.91.3) [30] with default parameter settings. The impacts of amino acid substitutions on protein function were predicted using the SIFT (version 5.2.2) [31] algorithm that has been implemented in the VEP tool. Variants with SIFT scores less than 0.05 were considered to be likely deleterious to protein function. In order to assess if known Mendelian trait-associated variants segregate among 49 sequenced OB cattle, we downloaded genomic coordinates of 155 trait-associated variants that are curated in the Online Mendelian Inheritance in Animals (OMIA) database [32, 33].

Population genetic analysis

We estimated average pairwise nucleotide diversity using VCFtools v0.1.15 (in windows of 10 kb) [34].

Detection of runs of homozygosity

Runs of homozygosity (ROH) were identified using a hidden Markov model (HMM)-based approach implemented in the BCFtools/RoH software [35, 36]. The recombination rate was assumed to be constant along the genome at 10^{-8} per base pair (1 cM/Mb). For the HMM-based detection of ROH, we considered Phred-scaled likelihoods (PL) and allele frequencies of 15,722,811 filtered SNPs before Beagle imputation. Because samples that are sequenced at low coverage are enriched for ROH [37], we considered only 33 samples with average sequencing coverage greater than 10-fold for the detection of ROH (Additional file 2). We only considered ROH longer than 50 kb because they were less likely to contain false-positives (Phred-score > 67 in our data, Additional file 2). Genomic inbreeding (F_{ROH}) was calculated for each animal as $F_{ROH}=\sum L_{ROH}/L_{GENOME}$, where $\sum L_{ROH}$ is the length of all ROH longer than 50 kb and L_{GENOME} is the length of the genome covered by SNPs [38], i.e., 2,512,054,768 bp in our data.

Further, ROH were classified into short (50 - 100 kb), medium (0.1 - 2 Mb) and long ROH (>2 Mb) reflecting ancient, historical, and recent relatedness, respectively [39]. The contribution of each ROH category to F_{ROH} was calculated for each animal. Average genomic inbreeding was compared between animals born before and after 1989 using the two samples t-test.

Detection of signatures of selection

To avoid potential bias arising from extended relationships among the sequenced animals, we did not consider nine sons from sire-son pairs for the detection of signatures of selection. For the remaining 40 cattle, we considered genotypes at 9,051,833 SNPs for which the ancestral allele provided by Rocha et al. [40] was detected in at least two species other than cattle and where it agreed with either the reference or alternate allele in our data. Haplotypes were phased using the Eagle2 [41] software with default parameter settings and assuming a constant recombination rate along the chromosome.

Integrated haplotype score (iHS)

To identify signatures of ongoing selection, integrated haplotype scores (iHS) were calculated for 8,903,614 variants with minor allele frequency (MAF) greater than 0.01 using the R package rehh v.2 [42]. Subsequently, |iHS| values were averaged for non-overlapping windows of 40 kb. Windows with either less than 10 SNPs or no |iHS| values were removed. To test if variants with similar |iHS| properties were pooled in 40 kb windows, we followed the approach of Granka et al. [43], i.e., we randomly selected the same number of SNPs that were pooled in 40 kb windows and calculated the mean variance of |iHS| in the true and permuted 40 kb windows for each chromosome. This procedure was repeated for 10,000 randomly selected 40 kb windows. The variance of |iHS| in the non-overlapping 40 kb windows (0.24) was significantly ($p < 0.01$) less than in windows of randomly selected SNPs (0.37) indicating that SNPs that were grouped in 40 kb windows had |iHS| values that were more similar than random SNPs.

Composite likelihood ratio (CLR)

Composite likelihood ratio (CLR) tests were carried out to identify alleles that are either close to fixation or already reached fixation due to past selection. Following the recommendation of Huber et al. [44], we removed 118,124 SNPs from the data which were fixed for the ancestral alleles because such sites are not informative for CLR tests. Using a pre-computed empirical allele frequency spectrum that was based on the remaining 8,933,709 SNPs for which ancestral and derived alleles were assigned (see above), we calculated CLR statistics in non-overlapping 40 kb windows using SweepFinder2 [45, 46]. A window size of 40 kb was chosen to allow comparison and alignment between |iHS| and CLR values.

Empirical p values were calculated for CLR and |iHS| windows [47] and the top 0.5% of windows were considered as candidate signatures of selection. Adjacent top 0.5% windows were merged using BEDTools v2.27.1 [48]. For each merged candidate signature of selection, the lowest p value among the merged windows was retained.

Characterization of signatures of selection

Genes within candidate signatures of selection were determined based on the Ensembl (release 91) annotation of the UMD3.1 assembly of the bovine genome. Gene-set enrichment analysis of genes within candidate signatures of selection was performed using PANTHER v.14.1 [49].

To determine the overlap between QTL and candidate signatures of selection, we downloaded genomic coordinates for 122,893 known QTL from the Animal QTL Database [15, 50]. We classified 85,722 unique QTL that were located on the 29 autosomes into six trait categories, i.e., exterior, health, milk, meat and carcass, production and reproduction (Additional file 3). QTL with identical genomic coordinates in associated trait categories were considered as one QTL. We used the intersect module of BEDTools v2.27.1 [48] to identify QTL that overlapped with candidate regions from CLR and |iHS| for each of the six trait categories separately. To test if QTL were enriched in candidate signatures of selection, we used a permutation test with 10,000 permutations. In each permutation, we randomly sampled the same number of regions of the same size as the candidate signatures of selection from CLR and |iHS| for each chromosome separately, and overlapped them with QTL of the respective trait categories as detailed using BEDTools (see above). The number of QTL that overlapped permuted regions was used as the empirical null distribution to calculate p values. P values less than 0.05 were considered as indicators for a significant enrichment of QTL in candidate signatures of selection.

Results

Overview of genomic diversity in OB cattle

We annotated 15,722,811 biallelic SNPs and 1,580,878 Indels that were discovered in 49 OB cattle [23]. Of these, 546,419 (3.5%) SNPs and 307,847 (19.5%) Indels were novel, i.e., they were not among the 102,091,847 polymorphic sites of the latest version (150) of the bovine dbSNP database. Functional annotation of the polymorphic sites revealed that the

vast majority of SNPs was located in either intergenic (73.8%) or intronic regions (25.2%). Only 1% of SNPs (160,707) were located in the exonic regions (Table 1). In protein-coding sequences, we detected 58,387, 47,249 and 1,264 synonymous, missense, and high impact SNPs, respectively. According to the SIFT scoring, 10,738 (22.7%) missense SNPs were classified as likely deleterious to protein function (SIFT score < 0.05). Among the high impact variants, we detected 580, 33, 106, 273 and 272 stop gain, stop lost, start lost, splice donor and splice acceptor variants, respectively. Deleterious and high impact variants were more frequent in the low than high allele frequency classes (Additional file 4).

Table 1 Number of SNPs and Indels in sequence ontology classes annotated using the VEP software

Sequence ontology class	SNP	Indel
splice_acceptor_variant	272	84
splice_donor_variant	273	71
stop_gained	580	16
frameshift_variant	0	1324
stop_lost	33	0
start_lost	106	4
inframe_insertion	0	290
inframe_deletion	0	440
missense_variant	47429	0
protein_altering_variant	0	12
splice_region_variant	9553	1059
stop_retained_variant	45	2
synonymous_variant	58387	
coding_sequence_variant	166	125
mature_miRNA_variant	83	23
5_prime_UTR_variant	6744	600

3_prime_UTR_variant	30716	4074
non_coding_transcript_exon_variant	6296	434
intron_variant	3960673	422764
non_coding_transcript_variant	24	25
upstream_gene_variant	526000	56715
downstream_gene_variant	454753	51672
intergenic_variant	10620678	1041144
Total	15722811	1580878

The majority of 1,580,878 Indels was in either intergenic (72.7%) or intronic (26.7%) regions. Only 2,213 (0.14%) Indels affected coding sequences. Among these, 1,499 were classified as high impact variants including 1,324, 16, 4, 71 and 84 frameshift, stop gain, start lost, splice donor and splice acceptor variants, respectively. Similar to previous studies in cattle [21, 51], coding regions were enriched for Indels with lengths in multiples of three indicating that they are less likely to be deleterious to protein function than frameshift variants (Additional File 5).

OMIA variants segregating in the OB population

We obtained genomic coordinates of 155 variants that are associated with Mendelian traits in cattle from the OMIA database to analyze if they segregate among the 49 OB cattle. It turned out that six OMIA variants were also detected in the 49 OB cattle including two variants in the *MOCOS* and *SLC45A2* genes that are associated with severe recessive disorders (Additional file 6). Two OB key ancestor bulls born in 1967 and 1974 (ENA SRA sample accession numbers SAMEA4827662 and SAMEA4827664) were heterozygous carriers of a single base pair deletion (BTA24:g.21222030delC) in the *MOCOS* gene (OMIA 001819-9913) that causes xanthinuria in the homozygous state in Tyrolean grey cattle [52]. Another two OB key ancestor bulls (sire and son; ENA SRA sample accession numbers SAMEA4827659 and SAMEA4827645) that were born in 1967 and 1973 were heterozygous

carriers of two missense variants in *SLC45A2* (BTA20:g.39829806G>A and BTA20:g.39864148C>T) that are associated with oculocutaneous albinism (OMIA 001821-9913) in Braunvieh cattle [53].

Population diversity

Nucleotide diversity (π) measures the degree of polymorphism in the population. The average genome wide nucleotide diversity within the OB breed was 0.001637/bp.

Runs of homozygosity and genomic inbreeding

Runs of homozygosity (ROH) were detected in autosomes of 33 OB animals with average sequencing depth greater than 10-fold. We found an average number of 2,044 (±79) ROH per individual with an average length of 179 kb (±17.6 kb). The length of the ROH ranged from 50 kb (minimum size considered, see methods) to 5,025,959 bp. On average, 14.58% (between 12 and 18.8%) of the genome (excluding sex chromsome) was in ROH (Additional file 7). Average genomic inbreeding for the 29 chromosomes ranged from 11.5% (BTA29) to 18.6% (BTA26) (Figure 1a).

Fig.1 ROH statistics in 33 OB cattle with average sequencing depth greater than 10-fold (a) Average genomic inbreeding and corresponding standard error for the 29 autosomes. (b) Average genomic inbreeding (F_{ROH}) calculated from short (50-100 kb), medium (0.1 - 2 Mb) and long (>2 Mb) ROH. (c) Average number of short, medium and long ROH.

In order to study the demography of the OB population [54–57], we calculated the contributions of short (50-100 kb), medium (0.1-2 Mb) and long (>2 Mb) ROH to the total genomic inbreeding (Additional file 7). The medium-sized ROH were the most frequent class (50.46%), and contributed most (75.01%) to the total genomic inbreeding. While short ROH occurred almost as frequent (49.17%) as medium-sized ROH, they contributed only 19.52% to total genomic inbreeding (Figure 1b&c; Additional file 7). Long ROH were rarely (0.36%) observed among the OB key ancestors and contributed little (5.47%) to total genomic inbreeding. The number of long ROH was positively correlated (r=0.77) with genomic inbreeding.

Genomic inbreeding was significantly (p = 0.0002) higher in 20 animals born between 1990 and 2012 than in 13 animals born between 1965 and 1989 (0.16 vs. 0.14) (Additional file 8). The high genomic inbreeding in animals born in more recent generations was mainly due to more long (>2 Mb; P=0.00004) and medium-sized ROH (0.1-1 Mb; p=0.001) (Figure 2).

Fig.2 Cumulative genomic inbreeding (%) in animals born between 1965-1989 (blue lines) and 1990-2012 (red lines) from ROH sorted on length and binned in windows of 10 kb. Thin dashed lines represent individuals and thick solid lines represent the average cumulative genomic inbreeding of the two groups of animals.

Signatures of selection

We identified candidate signatures of selection using two complementary methods, i.e., the composite likelihood ratio test (CLR) and the integrated haplotype score (iHS). The CLR test detects so called 'hard sweeps', i.e., genomic regions encompassing beneficial adaptive alleles that recently reached fixation [58]. The iHS detects 'soft sweeps', i.e., genomic regions where selection for beneficial alleles is still ongoing [59, 60]. The CLR was calculated for non-overlapping 40 kb windows. iHS values were calculated for each SNP and

subsequently averaged in 62,297 non-overlapping 40 kb windows. Windows with top 0.5% of CLR and iHS statistics were considered as candidate signatures of selection and adjacent windows were merged into regions (Figure 3a & 3b). We detected 95 and 162 candidate regions of signatures of selection (p < 0.005) using CLR and iHS, respectively, encompassing 12.56 Mb and 12.48 Mb (Additional file 9; Additional file 10). These candidate signatures of selection were not evenly distributed over the genome (Figure 3c). Functional annotation revealed that 136 and 157 protein-coding genes overlapped with 50 and 86 candidate regions from CLR and iHS analyses, respectively. All other candidate signatures of selection were located in intergenic regions. Closer inspection of the top selection regions of both analyses revealed that 16 CLR candidate regions overlapped with 25 iHS candidate regions on chromosomes 5, 7, 11, 14, 15, 17 and 26 (Figure 3c) encompassing 36 coding genes.

Fig.3 - Genome wide distribution of top 0.5% signatures of selection from CLR (a) and iHS (b) analyses and their overlap (c). Each point represents a non-overlapping window of 40 kb along the autosomes.

Top candidate signatures of selection

On chromosome 11, we identified 12 and 36 candidate regions using CLR and iHS analyses, respectively. The top CLR candidate region was located between 65.9 Mb and 68.5 Mb (Figure 4a) and it encompassed 24 protein-coding genes. The same region was also in ROH in 77% of the 33 sequenced animals. The top CLR windows were located between 67.5 and 68.2 Mb and they had CLR values higher than 5,000 and encompassed 5 genes (Figure 4a & 4e). The SNP density in these windows was low and their allele frequency distribution was highly skewed indicating the presence of a hard sweep (Figure 4c). The top iHS candidate region was located between 68.4 and 69.2 Mb (|iHS|-value of 4.5) encompassing 7 genes (Figure 4b & 4f). The allele frequencies of the SNPs within the top iHS region are approaching fixation showing ongoing selection possibly due to hitchhiking with the previous large sweep (Figure 4d).

Fig.4 Detailed view of a candidate selection region on chromosome 11 in OB that was detected using in CLR (a) and iHS (b). Each point represents a non-overlapping window of 40 kb. The dotted horizontal lines indicate the cutoff values (top 0.5%) for CLR (210) and iHS (2.13) statistics. The allele frequencies of the derived (red) or alternate alleles (black) (c and d) and genes (e and f) in top selection windows of CLR (67.5-68.4 Mb) and top selection windows of iHS (68.5-69.2 Mb). Gene names in green colour are present on the forward strand and black colour on the reverse strand of DNA.

Another striking CLR signal (CLR value 2,031) was detected on chromosome 6 between 38.5 and 39.4 Mb. This genomic region encompasses the *DCAF16, FAM184B, LAP3, LCORL, MED28* and *NCAPG* genes, and the window with the largest CLR value overlapped the *NCAPG* gene (Figure 5a & 5c). This signature of selection coincides with a QTL that is associated with stature, feed efficiency and fetal growth [61–63]. Most SNPs detected within this region were fixed for the alternate allele in the OB cattle of the present study. For instance, all 49 sequenced OB cattle were homozygous for the Chr6:38777311 G-allele which results in a likely deleterious (SIFT score 0.01) amino acid substitution (p.I442M) in the *NCAPG* gene that is associated with increased pre- and postnatal growth and calving difficulties [61].

Fig.5 Top CLR candidate region on chromosome 6 (a). Each point represents a non-overlapping window of 40 kb. The frequencies of the derived (red) or alternate alleles (black) (c) and genes (e) annotated between 38.5 and 39.4 Mb. Gene names in green colour are present on the forward strand and black colour on the reverse strand of DNA.

GO enrichment analysis

Genes within candidate signatures of selection from CLR and iHS analyses were enriched (after correcting for multiple testing) in the panther pathway (P00011) related to "Blood coagulation". Genes from CLR were also enriched in the pathway "P53 pathway feedback loops 1" (Additional file 11). Although we did not find any enrichment of GO-slim biological processes after correcting for multiple testing, 21 GO-slim biological processes including cellular catabolic processes, oxygen transport and different splicing pathways were nominally enriched for genes within CLR candidate signatures of selection and 14 GO-slim

biological processes including nervous system, sensory perception (olfactory receptors) and multicellular processes were nominally enriched for genes within iHS candidate signatures of selection (Additional file 11).

QTL enrichment analysis

We investigated the overlap between the candidate selection regions and QTL using QTL information curated at the Animal QTL Database (Animal QTLdb). We found that 74.7% and 83.9% of CLR and iHS candidate signatures of selection, respectively, were overlapping at least one QTL (Additional file 12). We tested for enrichment of these signatures of selection in QTL for six trait classes, i.e. exterior, health, milk, meat, production, and reproduction using permutation. It turned out that QTL associated with meat quality (P_{CLR} = 0.0004, P_{iHS} = 0.0003) and production traits (P_{CLR} = 0.0027, P_{iHS} = 0.0039) were significantly enriched in both CLR and iHS candidate signatures of selection. We did not detect any enrichment of QTL associated with milk, reproduction, health, and exterior traits neither in CLR nor in iHS candidate signatures of selection.

Discussion

We discovered 107,291 variants in coding sequences of 49 sequenced OB cattle, i.e., 15,558 more coding variants than a previous study in 43 Fleckvieh animals [64]. The larger number of coding variants detected is likely due to a larger sample size, higher sequencing coverage and a more informative gene annotation considered in our study. In agreement with previous studies in cattle [21], missense deleterious and high impact variants occurred predominantly at low allele frequency likely indicating that variants which disrupt physiological protein functions are removed from the population through purifying selection [65]. However, deleterious variants may reach high frequency in livestock populations due to, e.g., the frequent use of individual carrier animals in artificial insemination [66], hitchhiking with favorable alleles under artificial selection [67, 68], or demography effects such as population bottlenecks [69]. Because we predicted functional consequences of

missense variants using computational inference, they have to be treated with caution in the absence of experimental validation [70]. High impact variants that segregated among the 49 sequenced OB key ancestors were also listed as Mendelian trait-associated variants in the OMIA database including frameshift and missense variants in *MOCOS* and *SLC45A2*, respectively, associated with recessive xanthinuria [52] and oculocutaneous albinism [53]. To the best of our knowledge, calves neither with xanthinuria nor oculocutaneous albinism have been reported in the Swiss OB cattle population. The absence of affected calves is likely due to the low frequencies of the deleterious alleles and avoidance of matings between closely related heterozygous carriers. Among 49 sequenced cattle, we detected respectively only two bulls that carried the disease-associated alleles in *MOCOS* and *SLC45A2* in the heterozygous state. However, the frequent use of individual carrier bulls in artificial insemination might result in many diseased animals within a short time even when the frequency of the deleterious allele is low in the population [71]. Because we sequenced key ancestors that were born decades ago, we cannot preclude that these disease-associated alleles were lost during the recent population bottleneck in OB (Additional file1). A frameshift variant in *SLC2A2* (NM_001103222:c.771_778delTTGAAAAGinsCATC, rs379675307, OMIA 000366-9913) causes a recessive disorder in cattle that is similar to human Fanconi-Bickel syndrome [72–74]. Recently, the frameshift variant causing the disease was detected in the homozygous state in an OB calf with retarded growth due to liver and kidney disease [75]. We did not detect the disease-associated allele in our study likely because it is located on a rare haplotype that does not segregate in the 49 sequenced cattle. Most of the sequenced animals of the present study were selected for sequencing using the key ancestor approach, i.e., their genes contributed significantly to the current population [23, 76]. More sophisticated methods to select animals for sequencing might prioritize rarer haplotypes, thus increasing the likelihood to detect rarer deleterious alleles when the sequencing budget is constrained [77–79].

Breed diversity and genomic inbreeding

Original Braunvieh is a local cattle breed with approximately 10,000 cows registered in the breeding population and 4,500 calves entering the herd book every year (Additional file 1). In spite of the small population size, the nucleotide diversity ($\pi = 1.6 \times 10^{-3}$) is higher in OB than many European taurine cattle breeds with considerably more breeding animals including Holstein, Jersey and Fleckvieh (~$1.2\text{-}1.4 \times 10^{-3}$) [80, 81]. However, nucleotide diversity is lower in OB than African ($2.0\text{-}4.0 \times 10^{-3}$), New Danish Red ($1.7 \times 10^{-3}$) and Yakutian cattle breeds (1.7×10^{-3}) [81–83]. The average genomic inbreeding estimated from whole-genome sequencing data was 0.14 in OB. This is lower than estimates in Holstein (0.18), Jersey (0.24), Old Danish Red (0.23) and Belgian Blue (0.3) cattle [84, 85]. However, genomic inbreeding is slightly higher in OB than New Red Danish cattle (0.11), an admixed breed that contains genes from old Danish and other red breeds [84]. The relatively high genomic diversity in OB cattle is assumed to be the result of many different sires contributing to the gene pool due to frequent use of natural mating [25]. Our whole-genome sequencing based estimate of genomic inbreeding (0.14) is substantially higher than previous estimates obtained using either 50K SNP microarray data ($F_{ROH}= 0.029$, [25]) or pedigree records ($F_{PED}=0.02$, [24]) for the same population. Genotype data obtained using SNP microarrays with medium density (e.g., BovineSNP50) can only capture long ROH (>1 Mb) due to lower SNP density (~1 SNP per 50 kb) and underestimates the genomic inbreeding, particularly when short and medium-sized ROH contribute substantially to genomic inbreeding as it is the case in our study. In our data, short and medium-sized ROH accounted for 80.48% of total inbreeding. Most short and medium-sized ROH are not reliably detectable with SNP microarrays used in Signer-Hasler et al. [25], resulting in a massive underestimation of genomic inbreeding. Our estimate of the genomic inbreeding using WGS variants also includes short and medium-sized ROH that were previously missed using SNP array data, thus representing a realistic estimate of total genomic inbreeding.

Our findings show that medium-sized ROH which occurred due to historical inbreeding contribute most to the genomic inbreeding of the current OB population. The minor

contribution of long ROH to the genomic inbreeding indicates that recent inbreeding is relatively low in OB possibly due to use of many sires in natural matings. Our results based on ROH inferred from whole-genome sequencing variants corroborate that genomic inbreeding is lower in OB than most mainstream breeds [25]. However, comparing the number and size distribution of ROH across studies is difficult and subject to bias because misplaced genomic segments might break ROH into multiple small- and medium-sized ROHs and different ROH-detection approaches yield results that are not readily comparable [29, 34, 56, 86, 87]. Genomic inbreeding is increasing in the OB population in recent years mainly due to an increase in occurrence of long ROH. The recent population bottleneck in the OB population (Additional file 1) might promote matings between closely related animals that caused inbreeding to increase in recent generations. In this regard, genome-based mating strategies seem to be warranted to achieve sufficient genetic gain while maintaining genetic diversity and avoiding matings between carriers of disease-associated alleles [88, 89].

Signatures of selection

Using whole-genome sequencing data, we were able to identify more signatures of selection compared to SNP array data [25, 90], even though we used only 9 million SNP for which we could readily assign ancestral and derived alleles [40]. Using two complementary approaches, we found several new and known candidate regions that seem to be targets of recent or ongoing selection in OB. Many signatures of selection were located in non-coding regions corroborating that selection frequently acts on regulatory sites [91]. However, it is possible that an improved annotation of the bovine genome might place these regions in yet to be annotated coding regions. We applied methods to detect signatures of selection that depend on frequency changes of alleles (CLR) and haplotypes (iHS). The detected signatures of selection may be confounded by other evolutionary forces including genetic drift and background selection [19, 92, 93]. By considering only the top 0.5% of selection

signatures, we likely have reduced the number of false-positive signatures at the cost of missing some true signatures of selection.

We detected candidate selection regions in OB cattle that harbor genes associated with stature or milk production (*NCPAG, LCORL, LAP3*), feed efficiency or lipid metabolism (*R3HDM1, AOX1*), and unknown functions (*SLC25A33, TMEM201*) that were reported to be targets of selection in taurine and indicine cattle breeds [9, 91, 94, 95]. The presence of signatures of selection that are common in several breeds indicates that selection at these regions has happened either before the breeds diverged or independently after the formation of breeds [91, 95]. A number of genes that are targets of selection in various cattle breeds are associated with coat colour (*MC1R*, *KIT*), milk production (*DGAT1*, *ABCG2*, *GHR*) or stature (*PLAG1*) [47, 81, 95, 96]. These genes were not detected within the top 0.5% CLR and iHS windows in OB cattle possibly either due to absence of associated genomic variation in our data or because these genes are not under selection in OB cattle. While some cattle breeds including Holstein and Fleckvieh are selected for particular coat colour patterns [9, 47], the coat colour of OB cattle is uniform and animals with variation in coat colour are rarely observed [97]. Moreover, due to the use of OB cattle for both milk and beef production under extensive conditions, the milk production-associated variants that are under selection in many dairy breeds seem to be less important in OB cattle [94].

Because some of the genes that were identified as targets of selection in specialized breeds contain variants that explain a large proportion of the phenotypic variation for economically important traits, we investigated if such variants segregate in the sequenced OB cattle although they were not detected in the signature of selection analyses. A number of variants in high linkage disequilibrium stimulate the expression of *PLAG1*, thus increasing pre- and postnatal growth in cattle [62, 63, 98]. The OB key ancestors sequenced in our study are fixed for the stature-decreasing alleles at six variants whereas other candidate causal variants are either fixed for the stature-increasing allele or segregate at low allele frequency

indicating that a recombinant haplotype might segregate in Swiss OB cattle that could allow for further fine-mapping of that region (Additional File 13). Among known mutations affecting milk production traits, a mutation in *ABCG2* (p.Y581S, rs43702337 at 38,027,010 bp) did not segregate in our population of Swiss OB cattle [99] which agrees with a previous study in Brown Swiss cattle [100]. A variant (BTA14, g.1802265_1802266GC>AA, p.A232K) of the *DGAT1* gene is associated with milk production traits in various cattle breeds [101, 102]. The milk fat-enhancing and milk yield-lowering lysine-allele segregates in OB cattle at low frequency (0.03). A missense variant (BTA20, g.31909478A>T, p.Y279F) in the *GHR* gene is associated with milk protein percentage [103]. The protein fat percentage-lowering T-allele segregates at low frequency (0.06) in OB.

We observed a striking signature of selection on chromosome 11 that has previously been detected in a number of cattle breeds including Swiss Fleckvieh, Simmental, Eringer and Evolèner using microarray-called genotypes [25, 90]. Our results in OB cattle indicate that this region harbors a rapid sweep which seems to act on alleles with selection advantage [59]. While large sweeps are easy to detect using dense sequencing data, pinpointing causal alleles underpinning such regions remains challenging. Most of the variants in such regions are either fixed or segregate at very low frequency [91] which we also observed for the signature of selection on chromosome 11. In our study, the signature of selection on chromosome 11 encompassed millions of base pairs (between 66 and 72 Mb) and many genes, rendering the identification of underpinning genes and variants a difficult task. The windows with the highest CLR and |iHS| values did not encompass *PROKR1*, which was previously suggested to be the target of selection at this region due to its association with fertility [25, 90]. However, closer inspection of the sequence variants detected in our study revealed that a stop-gained variant in *PROKR1* (g.66998234C>A, rs476744845, p.Y293*) segregates at high frequency in the 49 sequenced OB key ancestors. Yet, it remains to be elucidated, if the presence of a high-impact variant in immediate proximity to a massive selective sweep is causal or just due to hitchhiking. The window with the largest |iHS| value

on chromosome 11 was right next to the *CAPN13* gene which is associated with meat tenderness and was also suggested as a potential target of selection by Signer-Hasler et al. [25].

Genes within the top 0.5% CLR and iHS windows were enriched in pathways related to reactive oxygen species, metabolic process, blood coagulation and nervous system, indicating that the identified regions under selection might harbor genomic variants that confer adaptive advantage to harsh environments. Moreover, QTL associated with meat quality and production traits including feed efficiency and body weight were enriched in selection signatures possibly indicating that OB cattle harbor variants that enabled them to adapt to particular feed conditions. Combining results from selection signature and association analyses might reveal phenotypic characteristics associated with genomic regions that showed evidence of past or ongoing selection [47], thus providing additional hints why particular genomic regions are under selection in OB cattle.

Conclusions

We provide a comprehensive overview of genomic variation segregating in the Swiss OB cattle population using whole-genome sequencing data of 49 key ancestor bulls. In spite of the small population size, genetic diversity is higher and genomic inbreeding is lower in OB than many other mainstream cattle breeds. However, genomic inbreeding is increasing in recent generations mainly due to large ROH which should be considered in future management of this breed. Finally, this study highlights regions that show evidence of past and ongoing selection in OB which are enriched for QTL related to meat quality and production traits and pathways related to blood coagulation, cellular metabolic process, and nervous system.

List of abbreviations

CLR – Composite likelihood ration; iHS – Integrated haplotype score; OB – Original

Braunvieh; OMIA – Online Mendelian Inheritance in Animals; QTL – Quantitative trait loci;

ROH – runs of homozygosity; SNP – Single nucleotide polymorphism

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Raw sequencing read data of all animals are available from the European Nucleotide

Archive (ENA) (http://www.ebi.ac.uk/ena) under primary accession PRJEB28191.

Funding

This study was financially supported by the Federal Office for Agriculture (FOAG), Bern. The

funding body was not involved in the design of the study and collection, analysis, and

interpretation of data and in writing the manuscript.

Author's contributions

Analysis of whole-genome sequencing data: MB NKK CD HP; Conceived and designed the

experiments: MB NKK HP; Wrote the paper: MB HP; Read and approved the final version of

the manuscript: all authors

Acknowledgements

We thank Braunvieh Schweiz for providing pedigree and genotype data of Original Braunvieh cattle.

References

1. Food and Agriculture Organization. The Second Report on the State of the World's Animal Genetic Resources for Food and Agriculture. Rome: FAO Commission on Genetic Resources for Food and Agriculture Assessments. 2015. www.fao.org/publications.

2. Marker-assisted conservation of European cattle breeds: an evaluation. Anim Genet. 2006;37:475–81.

3. Medugorac I, Medugorac A, Russ I, Veit-Kensch CE, Taberlet P, Luntz B, et al. Genetic diversity of European cattle breeds highlights the conservation value of traditional unselected breeds with high effective population size. Mol Ecol. 2009;18:3394–410.

4. Boettcher PJ, Hoffmann I, Baumung R, Drucker AG, McManus C, Berg P, et al. Genetic resources and genomics for adaptation of livestock to climate change. Front Genet. 2015;5:461.

5. Biscarini F, Nicolazzi E, Alessandra S, Boettcher P, Gandini G. Challenges and opportunities in genetic improvement of local livestock breeds. Front Genet. 2015;6:33.

6. García-Ruiz A, Cole JB, VanRaden PM, Wiggans GR, Ruiz-López FJ, Van Tassell CP. Changes in genetic selection differentials and generation intervals in US Holstein dairy cattle as a result of genomic selection. Proc Natl Acad Sci U S A. 2016;113:E3995-4004. doi:10.1073/pnas.1519061113.

7. Georges M, Charlier C, Hayes B. Harnessing genomic information for livestock improvement. Nat Rev Genet. 2019;20:135–56. doi:10.1038/s41576-018-0082-2.

8. Sørensen a C, Sørensen MK, Berg P. Inbreeding in Danish dairy cattle breeds. J Dairy Sci. 2005;88:1865–72. doi:10.3168/jds.S0022-0302(05)72861-7.

9. Gutiérrez-Gil B, Arranz JJ, Wiener P. An interpretive review of selective sweep studies in

Bos taurus cattle populations: Identification of unique and shared selection signals across breeds. Front Genet. 2015;6:167.

10. Randhawa IAS, Khatkar MS, Thomson PC, Raadsma HW. Composite Selection Signals for Complex Traits Exemplified Through Bovine Stature Using Multibreed Cohorts of European and African Bos taurus. G3 (Bethesda). 2015;5:1391–401.

11. Utsunomiya YT, Pérez O'Brien AM, Sonstegard TS, Van Tassell CP, do Carmo AS, Mészáros G, et al. Detecting Loci under Recent Positive Selection in Dairy and Beef Cattle by Combining Different Genome-Wide Scan Methods. PLoS One. 2013;8:1–11.

12. Gurgul A, Szmatoła T, Ropka-Molik K, Jasielczuk I, Pawlina K, Semik E, et al. Identification of genome-wide selection signatures in the Limousin beef cattle breed. J Anim Breed Genet. 2016;133:264–76.

13. Browett S, McHugo G, Richardson IW, Magee DA, Park SDE, Fahey AG, et al. Genomic characterisation of the indigenous Irish Kerry cattle breed. Front Genet. 2018;9:51.

14. Burren A, Neuditschko M, Signer-Hasler H, Frischknecht M, Reber I, Menzi F, et al. Genetic diversity analyses reveal first insights into breed-specific selection signatures within Swiss goat breeds. Anim Genet. 2016;47:727–39.

15. Hu ZL, Park CA, Reecy JM. Building a livestock genetic and genomic information knowledgebase through integrative developments of Animal QTLdb and CorrDB. Nucleic Acids Res. 2019;47:D701–10.

16. Matukumalli LK, Lawley CT, Schnabel RD, Taylor JF, Allan MF, Heaton MP, et al. Development and Characterization of a High Density SNP Genotyping Assay for Cattle. PLoS One. 2009;4.

17. Rowan TN, Hoff JL, Crum TE, Taylor JF, Schnabel RD, Decker JE. A Multi-Breed Reference Panel and Additional Rare Variation Maximizes Imputation Accuracy in Cattle. bioRxiv. 2019;:517144. doi:10.1101/517144.

18. Albrechtsen A, Nielsen FC, Nielsen R. Ascertainment biases in SNP chips affect measures of population divergence. Mol Biol Evol. 2010;27:2534–47.

19. Biswas S, Akey JM. Genomic insights into positive selection. Trends Genet.

2006;22:437–46.

20. Lachance J, Tishkoff SA. SNP ascertainment bias in population genetic analyses: why it is important, and how to correct it. Bioessays. 2013;35:780–6.

21. Daetwyler HD, Capitan A, Pausch H, Stothard P, van Binsbergen R, Brøndum RF, et al. Whole-genome sequencing of 234 bulls facilitates mapping of monogenic and complex traits in cattle. Nat Genet. 2014;46:858–65. doi:10.1038/ng.3034.

22. Günther T, Nettelblad C. The presence and impact of reference bias on population genomic studies of prehistoric human populations. bioRxiv. 2018;:487983. doi:10.1101/487983.

23. Crysnanto D, Wurmser C, Pausch H. Accurate sequence variant genotyping in cattle using variation-aware genome graphs. Genet Sel Evol. 2019;51:21. doi:10.1101/460345.

24. Hagger C. Estimates of genetic diversity in the brown cattle population of Switzerland obtained from pedigree information. J Anim Breed Genet. 2005;122:405–13.

25. Signer-Hasler H, Burren A, Neuditschko M, Frischknecht M, Garrick D, Stricker C, et al. Population structure and genomic inbreeding in nine Swiss dairy cattle populations. Genet Sel Evol. 2017;49:1–13.

26. Eggertsson HP, Jonsson H, Kristmundsdottir S, Hjartarson E, Kehr B, Masson G, et al. Graphtyper enables population-scale genotyping using pangenome graphs. Nat Genet. 2017;49:1654–60. doi:10.1038/ng.3964.

27. Zimin A V, Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D. A whole-genome assembly of the domestic cow, Bos taurus. Genome Biol. 2009;10. doi:10.1186/gb-2009-10-4-r42.

28. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:13033997v2 [q-bioGN]. 2013;00:1–3. http://arxiv.org/abs/1303.3997.

29. Browning SR, Browning BL. Rapid and Accurate Haplotype Phasing and Missing-Data Inference for Whole-Genome Association Studies By Use of Localized Haplotype Clustering. Am J Hum Genet. 2007;81:1084–97. doi:10.1086/521987.

30. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl

Variant Effect Predictor. Genome Biol. 2016;17:042374. doi:10.1186/s13059-016-0974-4.

31. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. SIFT missense predictions for genomes. Nat Protoc. 2016;11:1–9. doi:10.1038/nprot.2015.123.

32. Online Mendelian Inheritance in Animals database (OMIA). https://omia.org/home. Accessed 10 Oct 2018.

33. Nicholas FW, Hobbs M. Mutation discovery for Mendelian traits in non-laboratory animals: A review of achievements up to 2012. Anim Genet. 2014;45:157–70.

34. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. Bioinformatics. 2011;27:2156–8.

35. Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. BCFtools/RoH: A hidden Markov model approach for detecting autozygosity from next-generation sequencing data. Bioinformatics. 2016;32:1749–51.

36. Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M, et al. Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. Science (80-). 2015;348:242–5.

37. van der Valk T, Díez-del-Molino D, Marques-Bonet T, Guschanski K, Dalén L. Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in Eastern Gorillas. Curr Biol. 2019;29:165–70.

38. McQuillan R, Leutenegger AL, Abdel-Rahman R, Franklin CS, Pericic M, Barac-Lauc L, et al. Runs of Homozygosity in European Populations. Am J Hum Genet. 2008;83:359–72.

39. Kirin M, McQuillan R, Franklin CS, Campbell H, McKeigue PM, Wilson JF. Genomic runs of homozygosity record population history and consanguinity. PLoS One. 2010;5:e13996. doi:10.1371/journal.pone.0013996.

40. Rocha D, Billerey C, Samson F, Boichard D, Boussaha M. Identification of the putative ancestral allele of bovine single-nucleotide polymorphisms. J Anim Breed Genet. 2014;131:1–4. doi:10.1111/jbg.12095.

41. Loh PR, Danecek P, Palamara PF, Fuchsberger C, Reshef YA, Finucane HK. Reference-based phasing using the haplotype reference consortium panel. Nat Genet.

2016;48. doi:10.1038/ng.3679.

42. Gautier M, Klassmann A, Vitalis R. rehh 2.0: a reimplementation of the R package rehh to detect positive selection from haplotype structure. Mol Ecol Resour. 2017;17:78–90.

43. Granka JM, Henn BM, Gignoux CR, Kidd JM, Bustamante CD, Feldman MW. Limited evidence for classic selective sweeps in African populations. Genetics. 2012;192:1049–64.

44. Huber CD, DeGiorgio M, Hellmann I, Nielsen R. Detecting recent selective sweeps while controlling for mutation rate and background selection. Mol Ecol. 2015;25:142–56.

45. Nielsen R. Molecular Signatures of Natural Selection. Annu Rev Genet. 2005;39:197–218. doi:10.1146/annurev.genet.39.073003.112420.

46. Degiorgio M, Huber CD, Hubisz MJ, Hellmann I, Nielsen R. SweepFinder2: Increased sensitivity, robustness and flexibility. Bioinformatics. 2016;32:1895–7.

47. Qanbari S, Pausch H, Jansen S, Somel M, Strom TM, Fries R, et al. Classic Selective Sweeps Revealed by Massive Sequencing in Cattle. PLoS Genet. 2014;10. doi:10.1371/journal.pgen.1004148.

48. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. 2010;26:841–2.

49. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: More genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res. 2019;47:D419–26.

50. The Animal Quantitative Trait Loci (QTL) Database. https://www.animalgenome.org/QTLdb/cattle. Accessed 8 Mar 2019.

51. Das A, Panitz F, Gregersen VR, Bendixen C, Holm LE. Deep sequencing of Danish Holstein dairy cattle for variant detection and insight into potential loss-of-function variants in protein coding genes. BMC Genomics. 2015;16:1–12. doi:10.1186/s12864-015-2249-y.

52. Murgiano L, Jagannathan V, Piffer C, Diez-Prieto I, Bolcato M, Gentile A, et al. A frameshift mutation in MOCOS is associated with familial renal syndrome (xanthinuria) in Tyrolean Grey cattle. BMC Vet Res. 2016;12:1–10. doi:10.1186/s12917-016-0904-4.

53. Rothammer S, Kunz E, Seichter D, Krebs S, Wassertheurer M, Fries R, et al. Detection

of two non-synonymous SNPs in SLC45A2 on BTA20 as candidate causal mutations for oculocutaneous albinism in Braunvieh cattle. Genet Sel Evol. 2017;49:1–8.

54. Pemberton TJ, Absher D, Feldman MW, Myers RM, Rosenberg NA, Li JZ. Genomic patterns of homozygosity in worldwide human populations. Am J Hum Genet. 2012;91:275–92.

55. Purfield DC, Berry DP, McParland S, Bradley DG. Runs of homozygosity and population history in cattle. BMC Genet. 2012;13.

56. Bosse M, Megens H, Madsen O, Paudel Y, Frantz LAF, Schook LB, et al. Regions of Homozygosity in the Porcine Genome□: Consequence of Demography and the Recombination Landscape. PLoS Genet. 2012;8.

57. Ceballos FC, Joshi PK, Clark DW, Ramsay M, Wilson JF. Runs of homozygosity: windows into population history and trait architecture. Nat Rev Genet. 2018;19:220–34. doi:10.1038/nrg.2017.109.

58. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. Genomic scans for selective sweeps using SNP data. Genome Res. 2005;15:1566–75.

59. Sabeti PC, Schaffner SF, Fry B, Lohmueller J, Varilly P, Shamovsky O, et al. Positive natural selection in the human lineage. Science. 2006;312:1614–20. doi:10.1126/science.1124309.

60. Voight BF, Kudaravalli S, Wen X, Pritchard JK. A Map of Recent Positive Selection in the Human Genome. PLOS Biol. 2006;4. doi:10.1371/journal.pbio.0040072.

61. Eberlein A, Takasuga A, Setoguchi K, Pfuhl R, Flisikowski K, Fries R, et al. Dissection of genetic factors modulating fetal growth in cattle indicates a substantial role of the non-SMC condensin I complex, subunit G (NCAPG) gene. Genetics. 2009;183:951–64.

62. Pausch H, Flisikowski K, Jung S, Emmerling R, Edel C, Götz KU, et al. Genome-wide association study identifies two major loci affecting calving ease and growth-related traits in cattle. Genetics. 2011;187:289–97.

63. Bouwman AC, Daetwyler HD, Chamberlain AJ, Ponce CH, Sargolzaei M, Schenkel FS, et al. Meta-analysis of genome-wide association studies for cattle stature identifies common

genes that regulate body size in mammals. Nat Genet. 2018;50:362–7.

64. Jansen S, Aigner B, Pausch H, Wysocki M, Eck S, Benet-Pagès A. Assessment of the genomic variation in a cattle population by re-sequencing of key animals at low to medium coverage. BMC Genomics. 2013;14. doi:10.1186/1471-2164-14-446.

65. Miosge LA, Field MA, Sontani Y, Cho V, Johnson S, Palkova A, et al. Comparison of predicted and actual consequences of missense mutations. Proc Natl Acad Sci. 2015;112:E5189–98. doi:10.1073/pnas.1511585112.

66. Pausch H, Kölle S, Wurmser C, Schwarzenbacher H, Emmerling R, Jansen S, et al. A Nonsense Mutation in TMEM95 Encoding a Nondescript Transmembrane Protein Causes Idiopathic Male Subfertility in Cattle. PLoS Genet. 2014;10.

67. Kadri NK, Sahana G, Charlier C, Iso-Touru T, Guldbrandtsen B, Karim L, et al. A 660-Kb Deletion with Antagonistic Effects on Fertility and Milk Production Segregates at High Frequency in Nordic Red Cattle: Additional Evidence for the Common Occurrence of Balancing Selection in Livestock. PLoS Genet. 2014;10.

68. Pausch H, Ammermüller S, Wurmser C, Hamann H, Tetens J, Drögemüller C, et al. A nonsense mutation in the COL7A1 gene causes epidermolysis bullosa in Vorderwald cattle. BMC Genet. 2016;17:1–9. doi:10.1186/s12863-016-0458-2.

69. Bosse M, Megens HJ, Derks MFL, de Cara ÁMR, Groenen MAM. Deleterious alleles in the context of domestication, inbreeding, and selection. Evol Appl. 2019;12:6–17.

70. Dong C, Wei P, Jian X, Gibbs R, Boerwinkle E, Wang K, et al. Comparison and integration of deleteriousness prediction methods for nonsynonymous SNVs in whole exome sequencing studies. Hum Mol Genet. 2015;24:2125–37.

71. Schwarzenbacher H, Wurmser C, Flisikowski K, Misurova L, Jung S, Langenmayer MC, et al. A frameshift mutation in GON4L is associated with proportionate dwarfism in Fleckvieh cattle. Genet Sel Evol. 2016;48:1–10.

72. Santer R, Schneppenheim R, Dombrowski A, Gotze H, Steinmann B. Mutations in GLUT2, the gene for the liver-type glucose transporter, in patients with Fanconi-Bickel syndrome. Nat Genet. 1997;17:324–6.

73. Pausch H, Schwarzenbacher H, Burgstaller J, Flisikowski K, Wurmser C, Jansen S, et al. Homozygous haplotype deficiency reveals deleterious mutations compromising reproductive and rearing success in cattle. BMC Genomics. 2015;16:1–13. doi:10.1186/s12864-015-1483-7.

74. Burgstaller J, Url A, Pausch H, Schwarzenbacher H, Egerbacher M, Wittek T. Clinical and biochemical signs in Fleckvieh cattle with genetically confirmed Fanconi-Bickel syndrome (cattle homozygous for Fleckvieh haplotype 2). Berl Munch Tierarztl Wochenschr. 2016;129:132–7.

75. Joller S, Stettler M, Locher I, Dettwiler M, Seefried F, Meylan M, et al. Fanconi-Bickel-Syndrom□: eine bislang unerkannte Erbkrankheit beim Braunvieh. Schweizer Arch Tierheilkd. 2018;160:179–84.

76. Goddard ME, Hayes BJ. Genomic selection based on dense genotypes inferred from sparse genotypes. Proc 18th Conf Assoc Adv Anim Breed Genet. 2009;18:26–9.

77. Gusev A, Shah MJ, Kenny EE, Ramachandran A, Lowe JK, Salit J, et al. Low-pass genome-wide sequencing and variant inference using identity-by-descent in an isolated human population. Genetics. 2012;190:679–89.

78. Bickhart DM, Hutchison JL, Null DJ, VanRaden PM, Cole JB. Reducing animal sequencing redundancy by preferentially selecting animals with low-frequency haplotypes. J Dairy Sci. 2016;99:5526–34. doi:10.3168/jds.2015-10347.

79. Ros-Freixedes R, Gonen S, Gorjanc G, Hickey JM. A method for allocating low-coverage sequencing resources by targeting haplotypes rather than individuals. Genet Sel Evol. 2017;49:1–17.

80. Charlier C, Li W, Harland C, Littlejohn M, Coppieters W, Creagh F, et al. NGS-based reverse genetic screen for common embryonic lethal mutations compromising fertility in livestock. Genome Res. 2016;26:1333–41.

81. Mei C, Wang H, Liao Q, Wang L, Cheng G, Wang H, et al. Genetic architecture and selection of Chinese cattle revealed by whole genome resequencing. Mol Biol Evol. 2018;35:688–99.

82. Kim J, Hanotte O, Mwai OA, Dessie T, Salim B, Diallo B, et al. The genome landscape of indigenous African cattle. Genome Biol. 2017;18:1–14.

83. Weldenegodguad M, Popov R, Pokharel K, Ammosov I, Ming Y, Ivanova Z, et al. Whole-Genome Sequencing of Three Native Cattle Breeds Originating From the Northernmost Cattle Farming Regions. Front Genet. 2018;9:728.

84. Zhang Q, Guldbrandtsen B, Bosse M, Lund MS, Sahana G. Runs of homozygosity and distribution of functional variants in the cattle genome. BMC Genomics. 2015;16:1–16. doi:10.1186/s12864-015-1715-x.

85. Solé M, Gori AS, Faux P, Bertrand A, Farnir F, Gautier M, et al. Age-based partitioning of individual genomic inbreeding levels in Belgian Blue cattle. Genet Sel Evol. 2017;49:1–18. doi:10.1186/s12711-017-0370-x.

86. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet. 2007;81:559–75. doi:10.1086/519795.

87. Bertrand AR, Kadri NK, Flori L, Gautier M, Druet T. RZooRoH: an R package to characterize individual genomic autozygosity and identify homozygous-by-descent segments. Methods Ecol Evol. 2019;2019. doi:10.1111/2041-210X.13167.

88. Wellmann R. Optimum contribution selection for animal breeding and conservation: The R package optiSel. BMC Bioinformatics. 2019;20:1–13.

89. Cole JB. A simple strategy for managing many recessive disorders in a dairy cattle breeding program. Genet Sel Evol. 2015;47:1–13.

90. Rothammer S, Seichter D, Förster M, Medugorac I. A genome-wide scan for signatures of differential artificial selection in ten cattle breeds. BMC Genomics. 2013;14.

91. Boitard S, Boussaha M, Capitan A, Rocha D, Servin B. Uncovering adaptation from sequence data: Lessons from genome resequencing of four cattle breeds. Genetics. 2016;203:433–50.

92. Akey JM. Constructing genomic maps of positive selection in humans: Where do we go from here? Genome Res. 2009;:711–22.

93. Charlesworth B. The role of background selection in shaping patterns of molecular evolution and variation: Evidence from variability on the Drosophila X chromosome. Genetics. 2012;191:233–46.

94. Zhao F, McParland S, Kearney F, Du L, Berry DP. Detection of selection signatures in dairy and beef cattle using high-density genomic information. Genet Sel Evol. 2015;47:1–12. doi:10.1186/s12711-015-0127-3.

95. Xu L, Bickhart DM, Cole JB, Schroeder SG, Song J, Van Tassell CP, et al. Genomic signatures reveal new evidences for selection of important traits in domestic cattle. Mol Biol Evol. 2015;32:711–25.

96. Randhawa IAS, Khatkar MS, Thomson PC, Raadsma HW. Composite Selection Signals for Complex Traits Exemplified Through Bovine Stature Using Multibreed Cohorts of European and African *Bos taurus*. G3: Genes|Genomes|Genetics. 2015;5:1391–401. doi:10.1534/g3.115.017772.

97. Durkin K, Coppieters W, Drögüller C, Ahariz N, Cambisano N, Druet T, et al. Serial translocation by means of circular intermediates underlies colour sidedness in cattle. Nature. 2012;482:81–4.

98. Karim L, Takeda H, Lin L, Druet T, Arias JAC, Baurain D, et al. Variants modulating the expression of a chromosome domain encompassing PLAG1 influence bovine stature. Nat Genet. 2011;43:405–15.

99. Cohen-Zinder M, Seroussi E, Larkin DM, Loor JJ, Everts-Van Der Wind A, Lee JH, et al. Identification of a missense mutation in the bovine ABCG2 gene with a major effect on the QTL on chromosome 6 affecting milk yield and composition in Holstein cattle. Genome Res. 2005;15:936–44.

100. Pausch H, Emmerling R, Gredler-Grandl B, Fries R, Daetwyler HD, Goddard ME. Meta-analysis of sequence-based association studies across three cattle breeds reveals 25 QTL for fat and protein percentages in milk at nucleotide resolution. BMC Genomics. 2017;18:1–11.

101. Grisart B, Coppieters W, Farnir F, Karim L, Ford C, Berzi P, et al. Positional candidate

cloning of a QTL in dairy cattle: Identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. Genome Res. 2002;12:222–31.

102. Winter A, Kramer W, Werner FAO, Kollers S, Kata S, Durstewitz G, et al. Association of a lysine-232/alanine polymorphism in a bovine gene encoding acyl-CoA:diacylglycerol acyltransferase (DGAT1) with variation at a quantitative trait locus for milk fat content. Proc Natl Acad Sci. 2002;99:9300–5. doi:10.1073/pnas.142293799.

103. Blott S, Kim JJ, Moisio S, Schmidt-Küntzel A, Cornet A, Berzi P. Molecular dissection of a quantitative trait locus: a phenylalanine-to-tyrosine substitution in the transmembrane domain of the bovine growth hormone receptor is associated with a major effect on milk yield and composition. Genetics. 2003;163:253–66.

Supplementary Information

Additional file 1 File S1

File format: .docx

Title: Original Braunvieh herdbook population

Description: Number of female calves entering the OB herdbook between 1980 and 2016.

Additional file 2 File S2

File format: .pdf

Title: Runs of homozygosity in 49 OB cattle

Description: a) Total genome fraction in ROH in 49 cattle with high (<10x) and low (<10x) coverage (b) Phred confidence score for ROH in 33 cattle sequenced at average sequencing depth higher than 10-fold. Red dots indicate mean confidence scores for ROH.

Additional file 3 File S3

File format: .xlsx

Title: Bovine QTL information downloaded from the AnimalQTL database.

Description: Number of QTL for each trait. Traits are grouped in six trait categories.

Additional file 4 File S4

File format: .pdf

Title: Allele frequency distribution in different functional annotations

Description: Allele frequency of SNPs with different consequences according to VEP prediction, i.e., high impact, deleterious (missense SNP with sift score < 0.05). tolerated (missense SNPs with sift score > 0.05) and synonymous SNPs.

Additional file 5 File S5

File format: .pdf

Title: Distribution of length of Indels

Description: a) Number of Indels (x1000) with size less than 12 bp detected according to the number of affected bases. b) number of Indels detected in coding sequences.

Additional file 6 File S6

File format: .xlsx

Title: OMIA variants detected in OB

Description: Six OMIA variants detected in 49 sequenced OB animals with their respective frequency and information.

Additional file 7 File S7

File format: .xlsx

Title: ROH statistics of each animal (high coverage)

Description: Number, length, average length and genomic fraction of all ROH in each animal. Also categorized in short, medium and long ROH category.

Additional file 8 File S8

File format: .pdf

Title: Genomic inbreeding in OB cattle stratified by birth year

Description: Genomic inbreeding in two groups of animals born either between 1960 and 1989 or between 1990 and 2012.

Additional file 9 File S9

File format: .xlsx

Title: Candidate selection signatures detected using CLR

Description: Genomic coordinates, CLR values, p values and encompassed genes for 95 candidate selection signatures.

Additional file 10 File S10

File format: .xlsx

Title: Candidate selection signatures detected using iHS

Description: Genomic coordinates, |iHS| values, p values and encompassed genes for 162 candidate selection signatures.

Additional file 11 File S11

File format: .xlsx

Title: Summary of Gene Ontology enrichment analysis

Description: PANTHER and GO-Slim pathways enriched using genes encompassed in signatures of selection from CLR and iHS analyses.

Additional file 12 File S12

File format: .xlsx

Title: Overlap between QTL and signatures of selection

Description: QTL (from all 6 categories) that overlapped with CLR and iHS selection signatures.

Additional file 13 File S13

File format: .xlsx

Title: Frequency of candidate causal variants for a stature QTL on BTA14

Description: Genomic coordinates and allele frequencies of 14 variants nearby bovine *PLAG1* that were reported as candidate causal variants for a stature QTL in cattle by Karim et al. [95].